Title

Nuclear Argonaute HRDE-1 Sustains Chromatin-Independent Transcriptional Silencing

and Epigenetic Inheritance

Authors

Arantxa M. L. Rojas[1], Almira Chervova[1,2], Loan Bourdon[1], Jagannath Jayaraj[1,3], Ana Karina

Morao[1], Olivier Feudjio[1], Sofija Illic[1], Sara Capurso[1], Piergiuseppe Quarato[1], Julie Ovieve[1],

Germano Cecere[1#]

Affiliations

[1]Mechanisms of Epigenetic Inheritance, Department of Developmental and Stem Cell Biology,

Institut Pasteur, CNRS UMR3738, Paris

[2]Bioinformatics and Biostatistics Hub, Department of Computational Biology, Institut Pasteur,

USR 3756, CNRS, Paris

[3]Université Paris Cité, BioSPC, Paris, France

[#]Corresponding author: germano.cecere@pasteur.fr

Abstract

Epigenetic inheritance of transcriptional silencing is typically attributed to chromatin-based mechanisms in which Argonaute–small RNA complexes recruit histone-modifying enzymes. Here we show that, in *Caenorhabditis elegans*, the nuclear Argonaute HRDE-1 sustains germline repression independently of canonical heterochromatin marks. Inducible HRDE-1 depletion combined with germline nuclear sorting, CUT&Tag, and nascent transcription analysis revealed endogenous genes that become transcriptionally activated in the absence of HRDE-1 despite retention of H3K9me3 and H3K23me3. HRDE-1 further promotes polyUG-dependent amplification of antisense 22G-RNAs in perinuclear condensates, redefining nuclear Argonautes as active drivers of small RNA biogenesis rather than passive acceptors. Loss of HRDE-1 triggers progressive erosion of 22G-RNAs and prevents rapid re-establishment of silencing, underscoring the necessity of nuclear–cytoplasmic coupling for epigenetic memory. Our findings challenge the long-standing assumption that heterochromatin is required for transcriptional repression and redefine the mechanism by which nuclear Argonautes enforce epigenetic inheritance.

Introduction

Epigenetic inheritance of transcriptional silencing is widely attributed to chromatin-based repression[1] . In many systems, nuclear Argonaute proteins bound to small RNAs are thought to enforce this repression by recruiting histone-modifying enzymes, leading to deposition of repressive marks that can be stably transmitted across generations[2] . This model has shaped the prevailing view that heritable gene silencing requires canonical heterochromatin marks, yet whether such modifications are an obligate requirement for Argonaute-mediated silencing remains unresolved.

In *Caenorhabditis elegans*, small RNAs play a central role in transgenerational epigenetic inheritance. Exogenous double-stranded RNAs (dsRNAs) or endogenous PIWI-interacting RNAs (piRNAs) initiate the RNA-dependent RNA Polymerase (RdRP) transcription of antisense 22G-RNAs, which are loaded into Worm-specific Argonautes (WAGOs) to maintain silencing across generations[3] . The nuclear WAGO HRDE-1 (Heritable RNAi Deficient-1) is essential for this process[4] . Loss of HRDE-1 disrupts both dsRNA- and piRNA-triggered inheritance[4–6] . HRDE-1 activity has been associated with increased H3K9me3 and H3K23me3 at target loci[4,7–9], reinforcing the prevailing view that these histone modifications are required to maintain heritable RNA-mediated silencing.

Recent work has challenged this model. The inheritance of RNA interference (RNAi) initiated by double-stranded RNAs (dsRNAs) can persist in the absence of nuclear factors, relying instead on perinuclear condensates that propagate silencing via polyUGylated RNA templates and small RNA amplification[10,11] . Cytoplasmic factors such as ZNFX-1 and WAGOs can maintain silencing post-transcriptionally[10,11] , and in some cases silencing can continue even without the histone methyltransferases that deposit H3K9 and H3K23[12] . These findings raise the question of whether canonical heterochromatin marks are truly essential for HRDE-

1–mediated repression, or whether alternative heritable signals can substitute for chromatin

modifications.

To address this, we combined inducible germline-specific depletion of HRDE-1 with

nuclear sorting, CUT&Tag profiling of repressive histone modifications[13] , and nascent

transcription analysis[14] . This approach enabled direct comparison of transcriptional activity

and chromatin state at HRDE-1 target loci with germline resolution. We show that HRDE-1

maintains transcriptional silencing of a defined set of endogenous genes without requiring

canonical heterochromatin marks. Strikingly, some targets became transcriptionally active

while retaining these repressive modifications, demonstrating that HRDE-1 can enforce

transcriptional repression independently of chromatin state. Furthermore, we find that HRDE-

1 couples nuclear repression to the transgenerational maintenance of antisense 22G-RNAs

through amplification in perinuclear condensates. Together, these findings redefine the

mechanism of nuclear Argonaute–mediated silencing and highlight small RNAs themselves as

the primary heritable signal for a subset of germline targets.

Results

HRDE-1 is required in every generation to maintain germline gene silencing.

Most functional studies of HRDE-1 have relied on transgenic silencing reporters, which,

while informative, may not capture the full range of endogenous targets. To systematically

define HRDE-1–dependent repression of endogenous genes, we combined germline-specific

auxin-inducible degradation (AID)[15] with genome-wide transcriptional profiling. Depletion of

HRDE-1 using this system (**Fig. 1a** and **Extended Data Fig. 1a,b**) efficiently reduced protein

levels (**Extended Data Fig. 1b**) and recapitulated the heritable RNAi defect of the *hrde-1*

mutant in a GFP sensor assay (**Extended Data Fig. 1c**). We depleted HRDE-1 in one

generation from the first to the last larval stage (G0) and maintained worms for four successive

generations in auxin (G4) (**Fig. 1a** and **Extended Data Fig. 1d**). At each generation, synchronized L4-stage animals were collected using a COPAS Biosorter, as previously described[6] , to ensure precise developmental staging across samples. Whole-worm RNA-seq revealed that HRDE-1 depletion caused de-silencing of endogenous protein-coding and pseudogenes in both G0 and G4 animals, while very few genes were downregulated (**Fig. 1b**), indicating that we are primarily detecting genes directly repressed by HRDE-1. The total number of misregulated genes was lower than in the *hrde-1* mutant[16], consistent with the degron approach not fully depleting HRDE-1 protein. Importantly, the number of de-silenced genes did not increase between G0 and G4 (**Extended Data Fig. 1f and Fig. 1c**), indicating that HRDE-1 is required in each generation to maintain germline silencing, rather than having a transgenerational effect.

HRDE-1 acts downstream of RNAi- or piRNA-triggered silencing to mediate transcriptional repression. To compare its targets with those of PRG-1 (the *C. elegans* PIWI Argonaute), we examined available RNA-seq to compare expression changes in *hrde-1 (tm1200)* and *prg-1 (n4357)* mutants[16] for the 69 HRDE-1–silenced genes (**Extended Data Fig.1f**). Most of these genes were upregulated in *hrde-1* but not in *prg-1* mutants (**Fig. 1c,d**). RT-qPCR confirmed that PRG-1 loss had little effect on these targets (**Fig. 1e**). In contrast, mutation of NRDE-2, a nuclear silencing cofactor[4,17] , caused de-silencing to the same extent as *hrde-1* (**Fig. 1e**), confirming that these genes are bona fide nuclear pathway targets. The fact that these loci remain repressed by HRDE-1 in the absence of PRG-1 suggests they are distinct from the spermatogenic targets identified in our previous study[6], which require piRNA trigger. Consistent with this, RNA single-molecule fluorescence *in situ* hybridization (smFISH)[18] for one of the most strongly upregulated genes, F07B7.1, showed robust expression throughout the germline of *hrde-1* mutants but not in wild-type gonads (**Fig. 1f** and **Extended Data Fig. 1g**). HRDE-1 depletion from L1 to L4 recapitulated this pattern (**Fig. 1g** and **Extended Data**

Fig. 1h), demonstrating that repression requires HRDE-1 during germline development. This broad expression pattern contrasts with piRNA-regulated spermatogenic genes, which are only transiently expressed in pachytene nuclei[6], and supports the conclusion that these loci represent a distinct class of constitutively silenced, HRDE-1–dependent targets. Therefore, this defined gene set provides a tractable paradigm to dissect HRDE-1's mechanisms in small RNA–mediated epigenetic inheritance.

HRDE-1 depletion in G0 causes transcriptional de-silencing in germline nuclei.

To confirm that these changes were germline-specific, we used a nuclear GFP reporter to sort germline nuclei from HRDE-1–depleted worms (**Fig. 2a** and **Extended Data Fig. 2a-c**). We performed nuclear RNA-seq and GRO-seq from these sorted nuclei, obtaining similar number of de-silenced genes and very few downregulated, as we observe for the whole worm RNA-seq (**Fig. 1**). Moreover, the high level of de-silencing detected with both approaches strongly supports that these are nuclear targets of HRDE-1 (**Fig. 2b**), and that HRDE-1 maintains its transcriptional silencing every generation. We compared all the genes (n=364) with significant changes (FDR ≤ 0.05) either in RNA-seq or GRO-seq from germline nuclei and found that most of the genes present similar changes in both, total and nascent RNA levels (**Fig. 2e**). Altogether, these results show that depleting HRDE-1 from L1 to L4 is sufficient to transcriptionally de-silence its targets.

HRDE-1 maintains silencing independently of canonical heterochromatin marks.

Silencing by nuclear Argonautes is generally associated with deposition of repressive histone marks[2,4,7–9]. To determine whether the transcriptional de-repression observed upon HRDE-1 depletion results from loss of such marks, we developed a germline-specific CUT&Tag approach by combining GFP-based germline nuclear sorting with histone modification profiling, enabling high-resolution analysis of chromatin changes in the germline (**Fig. 3a, Extended Data Fig. 2 and 3,** and **Methods**). This design overcomes two major

technical limitations of previous studies: (1) whole-worm profiling dilutes germline signals

with abundant somatic H3K9me3, masking germline-specific changes; and (2) the widely used

anti-H3K9me3 antibody (*Abcam 8898*) in *C. elegans* that was recently shown to cross-react

with other repressive marks (H3K27me3, H4K20me3)[19], potentially confounding

interpretation. We instead identified and validated an antibody with high specificity for

H3K9me3 in *C. elegans* (*Active Motif 39161*) (**Extended Data Fig. 2a-c**). As a critical test, we

depleted SET-25, the sole H3K9me3 methyltransferase in *C. elegans*[20], from the germline

between L1 and L4, which completely erased H3K9me3 peaks in sorted germline nuclei

(**Extended Data Fig. 2d,e**), confirming antibody specificity and the sensitivity of our

approach.

We next examined H3K9me3 at genes with changes by nuclear RNA-seq on the same

sorted germline nuclei upon HRDE-1 or SET-25 depletion (**Fig. 2b**, **Fig. 3a-c** and **Extended**

Data Fig. 4c-e). Of 140 upregulated genes, 80 and 75 overlapped with H3K9me3 peaks, with

72 showing no significant reduction in signal upon HRDE-1 depletion (**Fig. 3b,c**). Only three

genes exhibited both a loss of H3K9me3 and transcriptional upregulation (**Fig. 3c**). Moreover,

most of the downregulated genes upon HRDE-1 depletion have no changes in H3K9me3

(**Extended Data Fig. 4c-e**). Importantly, depleting SET-25 did not de-repress HRDE-1 target

genes (**Fig. 3d and Extended Data Fig. 4a,b**), indicating that H3K9me3 loss alone is

insufficient for transcriptional de-repression.

H3K23me3 is another heterochromatin mark implicated in HRDE-1–mediated

silencing[12] . HRDE-1 depletion caused more widespread changes for this mark, with 288 peaks

increased and 672 peaks reduced (**Fig. 3e** and **Extended Data Fig. 4f**). Among the 140

transcriptionally upregulated genes, 117 overlapped H3K23me3 peaks (**Fig. 3e**); of these, 56

showed reduced H3K23me3, three showed increases, and the majority exhibited no significant

change (**Fig. 3e,f**). Notably, the widespread loss of H3K23me3 did not correspond to broad

transcriptional upregulation (**Fig. 3e**), and increased deposition of this mark did not result in

transcriptional repression (**Extended Data Fig. 4g,h**), indicating that H3K23me3 changes are

largely uncoupled from gene expression in this context.

It has also been proposed that heritable transcriptional silencing could involve removal

of the active promoter mark H3K4me3, in addition to recruitment of repressive histone

modifiers[21] . To test this possibility, we profiled H3K4me3 in HRDE-1–depleted germline

nuclei. Only a modest number of H3K4me3 peaks were gained genome-wide (38 total), with

just 11 of 140 derepressed genes showing increased H3K4me3 (**Extended Data Fig. 4i–k**).

Moreover, the majority of upregulated genes did not overlap any H3K4me3 peak, indicating

that the observed transcriptional derepression cannot be explained by H3K4me3 accumulation.

Together, these results demonstrate that HRDE-1 maintains germline transcriptional gene

silencing independently of these histone modification marks, and suggest that deposition of

repressive chromatin modifications is a downstream consequence of transcriptional silencing

rather than its primary cause. Importantly, several HRDE-1 targets were robustly de-repressed

while still retaining wild-type levels of H3K9me3 and/or H3K23me3, demonstrating that

transcription can proceed even in the presence of canonical repressive histone modifications.

HRDE-1 promotes the transgenerational maintenance of 22G-RNAs.

Having established that HRDE-1 sustains transcriptional silencing without canonical

heterochromatin marks, we next investigated whether its function is instead linked to

maintaining the small RNA populations required for transgenerational silencing. Loss of

HRDE-1 has previously been associated with large-scale changes in small RNA levels,

particularly antisense 22G-RNAs[16]. Artificial HRDE-1 tethering can also trigger *de novo*

production of heritable small RNAs[22], suggesting a direct role in promoting their biogenesis.

We sequenced small RNAs from G0, G1, G2, and G4 HRDE-1–depleted L4 worms (**Fig.**

4a,b and **Extended Data Fig. 5a**). Across generations, we predominantly observed a reduction

in antisense 22G-RNAs, with the extent of dysregulation increasing gradually (**Fig. 4b**). The progressive nature of this effect suggests that it cannot be explained solely by reduced stability of existing small RNAs, but rather reflects a transgenerational requirement for HRDE-1 in promoting 22G-RNA biogenesis.

Although 22G-RNAs were also reduced in G0 animals, none of the 69 genes consistently de-silenced upon HRDE-1 depletion (**Extended Data Fig. 1f** and) showed decreased 22G-RNAs (**Fig. 4c** and **Extended Data Fig. 5b,c**). Thus, in the early generation, transcriptional de-repression can occur without detectable changes in small RNA levels, and conversely, small RNA changes do not necessarily correspond to transcriptional changes (**Fig. 4c** and **Extended Data Fig. 5b,c**). This also suggests that the 22G-RNAs available for loading into other WAGOs in G0 HRDE-1-depleted animals is insufficient to compensate for the lack of HRDE-1 and maintain repression post-transcriptionally, as previously observed[10] .

The number of genes losing 22G-RNAs increased markedly across generations (**Fig. 4a,b** and **Extended Data Fig. 5b**), in contrast to the number of de-silenced genes, which remained stable (**Fig. 1a,b**). By G4, 22 of the 69 de-silenced genes had significantly reduced antisense 22G-RNAs in both replicate experiments (**Extended Data Fig. 5c**). These genes showed no change in 22G-RNAs in G0 but a strong reduction in G4, with a gradual decline across generations (**Fig. 4d**), as also shown for F15D4.5 as an example (**Fig. 4e**). These observations indicate that HRDE-1 has two separable roles: direct transcriptional silencing and maintenance of 22G-RNA populations across generations.

Mutator foci, but not Z granules, are required for HRDE-1–dependent 22G-RNA biogenesis and transcriptional repression.

To dissect the cytoplasmic requirements for HRDE-1–dependent silencing, we began by testing whether its targets require polyUGylation, a key step in 22G-RNA biogenesis that follows endonucleolytic cleavage of mRNA targets and is catalyzed by the polymerase RDE-

3^{23}. For several HRDE-1 targets with transgenerational 22G-RNA loss, polyUGylation in *hrde-*

1 mutants was as impaired as in *rde-3* mutants (**Fig. 5a**), indicating a strong defect in this step

of small RNA amplification. This contrasts with previous findings[6] showing that

polyUGylation of spermatogenic genes triggered by piRNAs is independent of HRDE-1 (e.g.,

D2096.1 and F55A6.3), suggesting that most HRDE-1 targets identified here do not require

piRNAs to initiate 22G-RNA biogenesis but do require HRDE-1 to sustain polyUGylation and

22G-RNA accumulation.

We next tested the involvement of the NYN domain-containing endonucleases RDE-8,

NYN-1, and NYN-2 acting upstream of polyUGylation[23,24]. PolyUGylation impairment in

hrde-1 mutants mirrored that observed in *rde-8* mutants or *nyn-1;nyn-2* double mutants (**Fig.**

5b). Consistent with this, expression levels of HRDE-1 target genes in *rde-8;nyn-1;nyn-2* triple

mutants ("endo") were comparable to *hrde-1* mutants (**Fig. 5c**).

Given this dependence on core cytoplasmic biogenesis factors, we asked whether HRDE-

1–dependent 22G-RNA production requires specific perinuclear condensates. Loss of MUT-

16, an essential scaffold for Mutator foci that house RDE-3, RDE-8, and other 22G-RNA

amplification factors[25] , phenocopied *hrde-1* mutants, causing similar de-silencing of HRDE-1

targets (**Fig. 5c**). These results indicate that Mutator foci are essential for both the cytoplasmic

biogenesis and the transcriptional repression of HRDE-1 targets, providing the key interface

between nuclear Argonaute activity and small RNA amplification.

Given the established role of Z granules in cytoplasmic post-transcriptional

silencing[10,26] , we next asked whether they contribute to the maintenance of endogenous

HRDE-1–mediated transcriptional repression. Z granules are perinuclear condensates

nucleated by the helicase ZNFX-1[26]. ZNFX-1 cooperates with cytoplasmic Argonautes to

sustain RNAi inheritance and to maintain silencing in the absence of nuclear HRDE-1, acting

through polyUGylated RNA templates and small RNA amplification[10,11]. We therefore tested

whether ZNFX-1 is similarly required for HRDE-1–regulated endogenous targets. Publicly available small RNA-seq data[10] together with our polyUGylation assays revealed that *znfx-1* mutants have a substantial reduction in 22G-RNAs and impaired polyUGylation for several HRDE-1-regulated genes (**Extended Data Fig. 6a–c**), similar to *hrde-1* mutants. Notably, however, RT-qPCR, and smFISH all showed only mild or no transcriptional derepression of these targets in *znfx-1* mutants compared to wild type (**Fig. 5c,d** and **Extended Figure 6d**). Moreover, combined loss of HRDE-1 and ZNFX-1 did not enhance derepression in either G0 or G1 (**Fig. 5e,f and Extended Fig. 6e**).

Thus, although Z granules influence the abundance of small RNAs for HRDE-1 targets, they are not required to maintain transcriptional repression. Unlike RNAi inheritance, Z granules do not provide a compensatory cytoplasmic pathway for these HRDE-1 targets, which relies selectively on Mutator foci to couple transcriptional repression with heritable small RNA amplification.

Silencing recovery mediated by HRDE-1 requires 22G-RNAs.

By applying inducible depletion of HRDE-1, we uncovered the dynamics of silencing maintenance across generations. We next investigated how silencing is re-established once HRDE-1 expression is restored. Worms depleted of HRDE-1 for four generations (G4) were removed from auxin treatment and propagated to the first (R1), fourth (R4), and sixth (R6) generations before collection at the L4 stage (**Fig. 6a** and **Extended Data Fig. 1a,d,e**). Notably, whereas depleting HRDE-1 during germline development (G0) was sufficient to de-silence these genes (**Fig. 1** and **Fig. 6b,c**), re-expressing HRDE-1 during the same developmental period failed to restore repression (**Fig. 6b,c**). RNA-seq analysis revealed that ~50% of the 69 protein-coding and pseudogenes de-silenced in G4 remained strongly upregulated in R1 (**Fig. 6b**). Instead, silencing was gradually re-established over several generations, with some genes remaining de-silenced even at R6 (**Fig. 6b,c**).

Since most targets showed a pronounced loss of antisense 22G-RNAs by G4, we next

assessed 22G-RNA levels during recovery. In R1, failure to restore silencing correlated with a

complete lack of 22G-RNA recovery, indicating that HRDE-1 requires these small RNAs to

act in transcriptional repression (**Fig. 6b,c**). In later generations, 22G-RNAs progressively

returned, paralleling transcriptional repression, consistent with the idea that their restoration,

like their maintenance, is transgenerationally promoted by HRDE-1.

To determine whether recovery depends on the same cytoplasmic biogenesis steps

required for maintenance, we monitored polyUGylated RNA intermediates for two HRDE-1

targets with distinct recovery kinetics, F15D4.5 and C38D9.2 (**Fig. 6c,d**), alongside a piRNA

target (*bath-13*) whose 22G-RNA accumulation are HRDE-1–independent[6] (**Extended Data**

Fig. 7a). The polyUGylation of *bath-13* remained unchanged across all generations, whereas

F15D4.5 and C38D9.2 showed a strong G4 reduction resembling *hrde-1* mutants and no

improvement in R1 (**Fig. 6c,d** and **Extended Data Fig. 7a**). PolyUG signal increased by R4

and R6, although for C38D9.2 the pattern remained weaker and more discrete than in controls,

matching its slower transcriptional recovery. These results suggest that gene-specific

differences in HRDE-1 efficiency at promoting polyUGylation underlie variable silencing

restoration rates, echoing earlier observations[12].

Finally, to directly test whether recovery failure in R1 is caused by small RNA depletion,

we repeated the experiment with shorter HRDE-1 depletion (G0) (**Fig. 6e**). Since 22G-RNA

levels are unaffected in G0 for de-silenced genes, we predicted that restoring HRDE-1 in the

next generation would enable immediate repression. Indeed, small RNA sequencing in R1 and

R2 after G0 depletion confirmed wild-type 22G-RNA levels (**Fig. 6e**), and RT-qPCR showed

that repression was re-established in R1 (**Fig. 6f** and **Extended Data Fig. 7b**). This

demonstrates that silencing recovery by HRDE-1 strictly depends on the presence of target-

matching 22G-RNAs and that their biogenesis, once lost, must be rebuilt over multiple

generations.

Discussion

By combining inducible germline depletion of HRDE-1 with a method we developed to

isolate and profile sorted germline nuclei (see methods), we captured both immediate and

transgenerational consequences of nuclear Argonaute loss with temporal resolution that

conventional mutants cannot provide. A key technical advance was the application of germline

CUT&Tag with a validated H3K9me3 antibody, which overcomes two limitations of previous

studies: masking of germline signals in whole-worm assays and the use of non-specific

antibodies[19]. This approach enabled direct, high-resolution interrogation of the chromatin state

at HRDE-1 targets.

This approach allowed us to identify a set of endogenous genes whose repression depends

exclusively on HRDE-1 and to dissect two distinct modes of silencing: (1) maintaining

transcriptional repression independently of canonical heterochromatin marks, and (2)

promoting the transgenerational maintenance of 22G-RNAs through polyUG-dependent

amplification in Mutator foci.

Chromatin independence of HRDE-1–mediated silencing

Our data reveal that HRDE-1 maintains transcriptional silencing without reliance on

canonical heterochromatin. Most HRDE-1–derepressed targets retain wild-type levels of

H3K9me3 and/or H3K23me3, and loss of either or both modifications has no impact on their

repression. This challenges the prevailing model, based largely on genetic evidence[12,27], that

nuclear Argonautes sustain silencing by recruiting histone methyltransferases. Instead, our

results support a model in which repressive marks are deposited downstream of HRDE-1–

mediated repression. We also considered whether derepression could be explained by changes

in the active promoter mark H3K4me3, which has been proposed to contribute to heritable silencing[21]. However, H3K4me3 gains were modest and affected only a minority of derepressed genes, indicating that promoter activation in this context is not driven by canonical active chromatin marks.

Consistent with this model, no direct interaction between HRDE-1 and histone methyltransferases has been detected[28]. In contrast, proteomic analyses have identified RNA polymerase II subunits among HRDE-1–associated factors[28], raising the possibility that HRDE-1 influences transcription directly. Our GRO-seq data are consistent with this view: HRDE-1 loss leads to increased nascent transcription, ruling out a model based solely on nuclear transcript degradation and suggesting impaired Pol II recruitment or initiation. Notably, this differs from the mechanism described for the somatic nuclear Argonaute NRDE-3, which inhibits Pol II during elongation[29]. Together, these findings indicate a distinct mode of repression by HRDE-1 in the germline and revise the current model of nuclear Argonaute function.

A distinct, piRNA-independent class of HRDE-1 targets.

The genes derepressed upon HRDE-1 loss differ from previously characterized spermatogenic targets, which are silenced anew each generation downstream of piRNAs and PRG-1[6]. In contrast, the genes we identify are constitutively repressed in the germline and remain silenced in every generation under normal conditions. We propose that their silencing may have originated from ancestral piRNA targeting but no longer requires ongoing piRNA input, leaving HRDE-1 as the sole factor responsible for their repression. Consistent with this, antisense 22G-RNAs for these genes are unaffected or even increased in *prg-1* mutants, reminiscent of the hyper-amplification of small RNAs triggered by RNAi in this background, which can perpetuate silencing indefinitely[30]. This distinct group of HRDE-1-dependent loci

348 therefore represents a paradigm of Argonaute-mediated repression that is self-sustaining and

349 independent of piRNA input.

Coupling nuclear silencing to cytoplasmic 22G-RNA amplification.

Our results show that HRDE-1 sustains transgenerational silencing not through stable

chromatin imprints, but by maintaining antisense 22G-RNA populations via polyUG-

dependent amplification in Mutator foci. This continual replenishment is essential, since

repression is rapidly lost in its absence. Thus, nuclear target recognition by HRDE-1 is coupled

to cytoplasmic small RNA biogenesis, ensuring that small RNAs themselves, rather than

heterochromatin, serve as the heritable memory.

Mechanistically, HRDE-1–dependent 22G-RNA biogenesis requires RDE-8/NYN-1/2–

mediated cleavage, RDE-3–mediated polyUGylation, and intact Mutator foci (MUT-16), but

not Z granules. This argues against the idea that HRDE-1 promotes small RNA amplification

in the nucleus[10], instead supporting a strictly cytoplasmic process. Although the Z granule

helicase ZNFX-1 has been implicated in nuclear-independent RNAi inheritance[11] and, in some

contexts, shown to act redundantly with HRDE-1[10], we find it is not required here: *znfx-1*

mutants display reduced 22G-RNAs at HRDE-1 targets but no transcriptional de-repression,

and combined loss of HRDE-1 and ZNFX-1 does not enhance defects. These results challenge

the view that cytoplasmic Argonautes can broadly compensate for nuclear silencing, and

instead re-establish HRDE-1 as central to transgenerational inheritance.

Small RNAs as the heritable memory of repression.

Our depletion–recovery experiments revealed that HRDE-1 requires pre-existing small

RNAs to re-establish silencing. After four generations without HRDE-1, repression was

restored only gradually across multiple generations, in parallel with the slow re-accumulation

of 22G-RNAs. By contrast, short-term depletion (one generation) left small RNA levels intact

and allowed immediate recovery. Recovery dynamics also varied between targets: for example,

the paralogs *F15D4.5* and *C38D9.2* displayed distinct resilience, likely reflecting differences

in polyUGylation efficiency and sequence complementarity. These findings demonstrate that

heritable silencing depends not on a persistent chromatin state, but on the continuous

availability of small RNAs to guide HRDE-1.

377 Broader implications for chromatin and transgenerational inheritance.

Our findings establish HRDE-1 as a central integrator of nuclear transcriptional

repression and cytoplasmic small RNA amplification, operating independently of canonical

heterochromatin marks. The observation that transcription can be reactivated despite the

continued presence of H3K9me3 and H3K23me3 parallels rare cases in *S. pombe* [31], where

transcriptional changes occur without loss of repressive chromatin, and in *Drosophila*[32], where

specialized transcriptional machinery drives piRNA cluster expression within a

heterochromatic environment. By extending this paradigm to the germline of *C. elegans*, our

GRO-seq results suggest that a nuclear Argonaute can repress transcription at the level of Pol

II recruitment, indicating that chromatin marks are not always the decisive signal for

transcriptional control.

More broadly, these results redefine how nuclear Argonautes sustain transgenerational

gene silencing: by coupling nuclear recognition of targets with continual small RNA

replenishment in cytoplasmic condensates, ensuring that small RNAs — rather than chromatin

imprints — serve as the heritable memory. By showing that transcriptional derepression occurs

independently of both repressive (H3K9me3, H3K23me3) and active (H3K4me3) histone

marks, our work compels a re-evaluation of the long-standing assumption that chromatin is the

central medium of epigenetic inheritance, and highlights nuclear Argonautes as active drivers

of small RNA-based epigenetic memory across generations.

396
397
398
399

Figures and Extended Data Figures.

Fig. 1: HRDE-1 is required in every generation to maintain germline gene silencing. a, Schematic of worm cultivation for transcriptome profiling. HRDE-1 was depleted in the germline using an auxin-inducible degron from L1 to L4 (G0) or continuously for four generations (G4), followed by total RNA-seq from whole worms. **b,** MA-plot showing $\log_2$ fold change versus mean expression in G0 and G4 samples for protein-coding and pseudogenes. Upregulated genes are shown in orange ($\log_2 FC \geq 1$, FDR ≤ 0.05); downregulated genes in blue ($\log_2 FC \leq -1$, FDR ≤ 0.05). n = 2 biological replicates. **c,** Box plot of $\log_2$ fold change (FC) in RNA-seq for the 68 upregulated genes in G0 and G4 (**see Extended Data Fig. 1f**), and their changes in *hrde-1* and *prg-1* mutants (Barucci et al., 2020). **d,** Genome browser views of representative targets, showing RNA-seq for HRDE-1-depleted (control, blue; HRDE-1 depletion, red) and *hrde-1* or *prg-1* mutants (WT, grey; *hrde-1*, green; *prg-1*, pink). TE = mapped transposable elements. Data are averaged from two biological replicates. **e**, RT–qPCR validation of selected HRDE-1–regulated genes in *hrde-1*, *prg-1*, and *nrde-2* mutants at the L4 stage, normalized to *act-3*. Bars show mean $\log_2 FC$; dots represent individual replicates (n = 3). **f**, **g**, RNA FISH for F07B7.1 (magenta) in L4 stage *hrde-1* mutants (**f**) or HRDE-1–depleted G0 worms (**g**) compared to controls. DAPI (cyan) marks nuclei; gonads are outlined with dashed white lines. Details of the two areas outlined in yellow are shown below for each gonad. Corresponding single-channel images are in **Extended Data Fig. 1g,h**.

Fig. 2: De-silencing in G0 HRDE-1-depleted germline nuclei is transcriptional. a, Schematic of germline nuclei isolation after HRDE-1 depletion for RNA-seq and GRO-seq analysis. **b, c,** MA-plot showing $\log_2$ fold change versus mean expression in G0 for protein-coding and pseudogenes in RNA-seq (**b**) and GRO-seq experiments (**c**). Upregulated genes are shown in orange ($\log_2 FC \geq 1$, FDR ≤ 0.05); downregulated genes in blue ($\log_2 FC \leq -1$, FDR ≤ 0.05). n = 4 biological replicates **d,** Genome browser views of representative targets, showing

RNA-seq and GRO-seq for G0 HRDE-1-depleted (control, blue; HRDE-1 depletion, red)

germline nuclei, RNA-seq from G0 whole worm (as in **Fig. 1d**), and RNA-seq and GRO-seq

from whole worms *hrde-1* mutants are included for comparison (WT, grey; *hrde-1*, green)

(Data from Barucci et al., 2020 and Cornes et al., 2022). TE = mapped transposable elements.

Data are averaged from two or four biological replicates. **e,** Heatmap summarizing significant

expression changes (FDR $\leq$ 0.05) for HRDE-1–silenced genes in G0 HRDE-1–depleted

germline nuclei in RNA-seq or GRO-seq.

Fig. 3: HRDE-1 maintains silencing independently of canonical heterochromatin marks.

a, Schematic of germline nuclei isolation after HRDE-1 depletion for histone modification

profiling by CUT&Tag. **b**, MA-plot of H3K9me3 CUT&Tag signal in G0 germline nuclei after

HRDE-1 depletion. Genes upregulated in G0 HRDE-1–depleted nuclei (from **Fig. 2b**) that

intersect H3K9me3 peaks are shown in orange; significantly reduced H3K9me3 peaks are in

dark grey. n = 4 biological replicates. **c**, Box plot of $\log_2$ fold change (FC) for upregulated genes

from G0 HRDE-1–depleted germline nuclei (**Fig. 2b**) intersecting with H3K9me3 peaks.

Categories: ns (no significant change but H3K9me3 peak present), no peak, or DOWN

(significantly reduced H3K9me3). **d**, RT–qPCR validation of selected genes in HRDE-1–

depleted (green) or SET-25–depleted (pink) L4 germlines, normalized to *act-3*. Bars show

mean $\log_2$FC; dots indicate biological replicates. Bars below indicate $\log_2$FC of H3K9me3 peak

signal for each gene; "na" = no detected peak. **e**, MA-plot of H3K23me3 CUT&Tag signal in

G0 germline nuclei after HRDE-1 depletion, with color coding as in (**b**). n = 4 biological

replicates. **f**, Box plot of $\log_2$FC for upregulated genes from G0 HRDE-1–depleted germline

nuclei (**Fig. 2b**) intersecting with H3K23me3 peaks. Categories as in (**c**).

Fig. 4: HRDE-1 is required for transgenerational maintenance of 22G-RNAs. a,

Experimental design for small RNA-seq across generations following germline HRDE-1

depletion. HRDE-1 was depleted during development (G0) or continuously for one (G1), two

(G2), or four (G4) generations. **b**, MA-plot of antisense 22G-RNAs mapping to protein-coding

and pseudogenes in HRDE-1–depleted versus control worms. Orange = significantly increased

22G-RNAs ($\log_2$FC $\geq$ 1, FDR $\leq$ 0.05), blue = significantly decreased 22G-RNAs ($\log_2$FC $\leq$ –

1, FDR $\leq$ 0.05). n = 2 biological replicates. **c**, Genome browser view of F15D4.5, a

representative HRDE-1 target. mRNA expression is similarly de-silenced in G0 and G4 (top;

RNA-seq, from **Fig. 1b**), whereas 22G-RNAs are reduced only in G4 (bottom). Control = blue,

HRDE-1 depletion = red. TE = mapped transposable elements. **d**, Box plot of $\log_2$ fold change

(FC) at G0, G1, G2, and G4 in antisense 22G-RNAs for 21 HRDE-1-silenced genes that lose

small RNAs in G4 (**see Extended Data Fig. 5c**). **f**, Genome browser view showing gradual,

generation-dependent reduction of antisense 22G-RNAs at F15D4.5. Control = blue, HRDE-1

depletion = red. TE = mapped transposable elements.

Fig. 5: Mutator foci, but not Z granules, are required for HRDE-1–dependent 22G-RNA

biogenesis and transcriptional repression. a,b, PolyUGylation assay for HRDE-1–silenced

genes that lose 22G-RNAs transgenerationally in different genetic backgrounds. The gene *gsa-*

1 is included as a control. **c**, RT–qPCR analysis of HRDE-1–silenced genes in *hrde-1*, *rde-*

8;nyn-1;nyn-2 triple mutants ("endo"), *mut-16* and *znfx-1* mutants at the L4 stage. Expression

values are normalized to *act-3*. Bars represent mean $\log_2$ fold change (FC), dots indicate

biological replicates (n = 3). **d**, RNA FISH for F07B7.1 (magenta) in L4-stage wild-type, *hrde-*

1, *znfx-1*, and *hrde-1;znfx-1* mutants. DAPI (cyan) marks nuclei; gonads are outlined with

dashed white lines. e, RT–qPCR analysis of selected HRDE-1-silenced genes as in HRDE-1-

depleted G0 and G1 worms at the L4 stage in control and *znfx-1* mutant backgrounds,

normalized to *act-3*. Bars show mean $\log_2$FC; dots represent biological replicates (n = 5). In

Extended Data Fig. 6d, the expression changes in *znfx-1* and wild-type worms grown in

control or auxin are shown.

Fig. 6: HRDE-1 silencing maintenance requires 22G-RNA biogenesis. a, Experimental design to assess transgenerational maintenance and recovery of HRDE-1–dependent silencing. **b,** Heatmap summarizing expression and 22G-RNA changes for the 69 HRDE-1–silenced genes (from **Fig. 1b**) in G0 and G4 HRDE-1–depleted worms (red) and in R1, R4, and R6 generations after HRDE-1 re-expression (purple). **c,** Genome browser views of representative genes: F15D4.5 and C38D9.2 (HRDE-1 targets) showing RNA-seq and 22-RNAs changes. Control samples are in blue, HRDE-1 depletion in red, and recovery in purple. TE = mapped transposable elements. **d,** PolyUGylation assay for F15D4.5 and C38D9.2 across control (blue), G0 and G4 depletion (red), and R1, R4, and R6 recovery generations (purple). *gsa-1* serves as a control. **e,** Top: Experimental design testing silencing recovery after one (G0) or four (G4) generations of depletion. Bottom: Genome browser snapshots showing 22G-RNA levels for F15D4.5 and C38D9.2 in control (blue), depletion (red, G0), and recovery (purple, R1 and R2). **f,** RT–qPCR analysis of selected genes in G0, R1, and R2 worms at the L4 stage, normalized to *act-3*. Bars show mean log_2FC; dots represent biological replicates (n = 3–5). Bottom: corresponding average fold change of 22G-RNAs for each gene across generations.

Extended Data Fig. 1: Validation of the auxin-inducible degron system and experimental setup for HRDE-1 depletion and recovery. a, Schematic of the auxin-inducible degron (AID) strategy for HRDE-1 depletion in the germline. **b,** Western blot of AID::3xFLAG::HRDE-1 in wild-type (untagged HRDE-1) and AID-tagged strains with or without germline TIR1, in the presence or absence of auxin. Anti-FLAG antibody was used for detection; GAPDH served as a loading control. **c,** RNAi inheritance assay in AID::3xFLAG::HRDE-1; GFP::piRNA sensor worms injected with dsRNA against GFP and developed on ethanol (control) or auxin (HRDE-1 depleted). Silencing was monitored in the F1 generation. **d,** Stage distribution of L4 larvae sorted with COPAS used for RNA-seq and small RNA-seq in **Fig. 1a,b**, **Fig. 6a-d**, and **Extended Data Fig. 4b,c**. **e,** Western blot of L4 larvae at G0 and G4 (HRDE-1 depleted, red)

and R1 (recovery, purple), compared to control (blue). Two biological replicates used for sequencing are shown (replicates 3 and 4). **f,** Venn diagram showing the intersection between G0 and G4 upregulated genes (FDR <= 0.05 and log2FC >=1) from RNA-seq of HRDE-1 depleted worms (**Fig. 1b**). **g,h,** Corresponding single-channel images of RNA FISH shown in **Fig. 1 f,g.**

Extended Data Fig. 2: Set up for germline nuclei isolation from HRDE-1-depleted worms. **a,** Schematics showing the preparation of HRDE-1-depleted germline nuclei for downstream approaches (Corresponding to **Figs. 2 and 3**). **b,** Western blot of L4 lysates corresponding to samples used in **Fig. 2** to test HRDE-1 depletion in each replicate (n = 4 biological replicates). **c,** Record examples of nuclei sorting using FACS, P4 is the sorting gate for GFP-positive nuclei.

Extended Data Fig. 3: Setup for H3K9me3 analysis using CUT and Tag in sorted germline nuclei. a, Genome browser views of a representative region in Chromosome V, showing CUT and Tag results using four different antibodies to detect H3K9me3 in wild-type and *met-2;set-25* L4 worms. **b,** Genome browser views of the same representative region as in **a**, showing CUT and Tag results using the Active Motif antibody to detect H3K9me3, with addition of synthetic nucleosomes (spike-in, Epicypher) in wild-type and *met-2;set-25* sorted L4 worms. Coverage normalized by Counts per million or using spike-in scaling factors is shown. **c,** Antibody specificity table for H3K9me3 antibody corresponding to the experiment shown in **b**. **d,** Schematics showing the preparation of SET-25-germline-depleted germline nuclei for CUT and Tag to assess changes in H3K9me3. On the right, MA plot showing H3K9me3 changes upon SET-25 depletion (G0). **g,** Genome browser views of the same representative region as in **a**, showing CUT and Tag results using the Active Motif antibody to detect H3K9me3 in germline nuclei depleted for SET-25 (G0) and control from the experiment shown in **d**. The coverage was normalized using a spike-in-derived scaling factor.

Extended Data Fig. 4: HRDE-1 maintains gene silencing independently of canonical histone modifications. a, MA-plot of H3K9me3 CUT&Tag signal in germline nuclei upon SET-25 depletion with orange highlighting genes upregulated in HRDE-1–depleted nuclei (**Fig. 2b**) that intersect H3K9me3 peaks. b, Same as in **a**, but with blue highlighting genes downregulated in HRDE-1–depleted nuclei. **c**, MA-plot of H3K9me3 signal upon HRDE-1 depletion (G0). **d**, Same as (**c**), highlighting genes downregulated in HRDE-1–depleted nuclei (**Fig. 2b**) that intersect H3K9me3 peaks. **e**, Box plot of $\log_2$ fold change (FC) for downregulated genes from HRDE-1–depleted germline nuclei grouped by H3K9me3 status: ns (no significant change, peak present), no peak, or DOWN (significantly reduced peak). **f**, MA-plot of H3K23me3 signal upon HRDE-1 depletion (G0). **g**, Same as **f**, with blue highlighting downregulated genes intersecting H3K23me3 peaks. **h**, Box plot of $\log_2$FC for downregulated genes from HRDE-1–depleted germline nuclei grouped by H3K23me3 status. **i**, MA-plot of H3K4me3 signal upon HRDE-1 depletion (G0). **j**, Same as **i**, with orange highlighting upregulated genes intersecting H3K4me3 peaks. **k**, Same as **i**, with blue highlighting downregulated genes intersecting H3K4me3 peaks.

Extended Data Fig. 5: Transgenerational dynamics of 22G-RNAs upon HRDE-1 depletion. a, Schematic of worm cultivation for small RNA analysis and Western blot confirming HRDE-1 depletion in sorted L4 larvae used for library preparation, corresponding to samples used in **Fig. 4b**. **b**, MA-plot of antisense 22G-RNA levels in G0 and G4 worms. Orange = upregulated 22G-RNAs ($\log_2$FC $\geq$ 1, FDR $\leq$ 0.05), blue = downregulated 22G-RNAs ($\log_2$FC $\leq$ -1, FDR $\leq$ 0.05). **c,** Venn diagram showing overlap between the 69 HRDE-1–silenced genes (**Fig. 1b**) and those with significant 22G-RNA changes in G0 and G4 in **b**.

Extended Data Fig. 6: The granule component, ZNFX-1, is required for small RNA maintenance but not silencing of HRDE-1-silenced genes. a, Box plot of $\log_2$ fold change (FC) of 22G-RNAs in *hrde-1*, *znfx-1*, and *znfx-1;hrde-1* compared to wild-type for the 69 genes

upregulated in G0 and G4 HRDE-1-depleted L4 larvae (**Fig. 1b**). Data from Ouyang et al.,

2022. **b,** Genome browser views of representative targets, F07B7.1 and F15D4.5, showing

22G-RNAs for *hrde-1*, *znfx-1*, and *znfx-1;hrde-1* (WT, grey; *hrde-1*, green; *znfx-1*, yellow;

double mutant, lime). TE = mapped transposable elements. Data are averaged from two

biological replicates (Ouyang et al., 2022). **c,** PolyUGylation assay for HRDE-1–silenced genes

that lose 22G-RNAs transgenerationally in wildtype (wt), *hrde-1* and *znfx-1*, backgrounds. The

housekeeping gene *gsa-1* is included as a control. **d,** RT–qPCR analysis of HRDE-1–silenced

genes from **Fig. 5e** in *znfx-1* (yellow) and wild-type (green) exposed to auxin or control at the

L4 stage. Expression values are normalized to *act-3*. Bars represent mean $\log_2$ fold change

(FC), dots indicate biological replicates (n = 3-5). **e,** RT–qPCR analysis of selected genes as in

Fig. 5f in HRDE-1-depleted G0 and G1 worms at the L4 stage in control and *znfx-1* mutant

backgrounds, normalized to *act-3*. Bars show mean $\log_2$FC; dots represent biological replicates

(n = 4–5).

Extended Data Fig. 7: HRDE-1 silencing maintenance requires 22G-RNA biogenesis. a,

PolyUGylation assay for *bath-13* across control (blue), G0 and G4 depletion (red), and R1, R4,

and R6 recovery generations (purple). *gsa-1* serves as control. **b,** RT–qPCR analysis of selected

genes as in **Fig. 6f** in G0, R1, and R2 worms at the L4 stage, normalized to *act-3*. Bars show

mean $\log_2$FC; dots represent biological replicates (n = 3–5). Bottom: corresponding average

fold change of 22G-RNAs for each gene across generations.

Methods

<u>Strains and worm culture</u>

Worms were grown at 20ºC in NGM plates and fed on OP50 using standars methods[33] . For

staged L4 larvae, worms were synchronized by bleaching. Briefly, gravid adults were incubated

5 min at 60 RPM in a 0.2 N NaOH and 1.5% NaOCl solution, washed three times, and

incubated overnight for hatching with M9 buffer. 40,000 starved L1 larvae were placed on 150

mm NGM plates with 1mL of fresh 100X OP50 or NGM supplemented with Ethanol (0.2%

v/v) or Auxin (indole-3-acetic-acid, 500 µM in Ethanol) for Auxin-inducible degradation

experiments. For transgenerational experiments, L4 larvae were collected at each generation

(around 44 hph), while one plate was kept until 72 hph for bleaching gravid adults to obtain

the next generation.

<u>Generation of CRISPR–Cas9 lines</u>

Cas9-guide RNA (gRNA) ribonucleoprotein complexes were microinjected into the

hermaphrodite syncytial gonad[34]. AID CRISPR-Cas9 tags were generated using double-

stranded DNA repair templates (30bp homology arms on each side) amplified by PCR using

an AID::3xFlag gBlock (IDT) sequence. Silent mutations were included where necessary in the

repair templates to prevent Cas9 cleavage. Mix concentrations were adapted from a previous

study[35]. In brief, 10 µl mixes typically contained the following final concentrations: 0.1µg/µL

Cas9-NLS protein (TrueCut V2, Invitrogen), 100 ng/µl in vitro transcribed target-gene gRNA,

80ng/µl of target-gene single-stranded DNA (ssDNA) repair template or 300 ng/µl target-gene

double-stranded DNA (dsDNA) repair template, and 80 ng/µL pRF4 (roller marker). Cas9 and

the target-gene gRNA were pre-incubated for 10-15 minutes at 37°C before adding the other

components to the mixture. dsDNA repair templates were subjected to a melting/annealing step

before addition to the final mix.

<u>Worm population sorting</u>

Worms were prepared and collected as described before[6]. Briefly, we obtain large populations

of precisely developmentally staged worms for genome-wide approaches, using a COPAS

Biosorter (Union Biometrica). For obtaining synchronous L4 population, among 40,000 to

80,000 larvae were harvested at 44 hph. To establish the sorting gate, for each sample, ~ 100

worms were collected and inspected under a 40X objective brightfield microscope. The number

of worms on each substage of the L4 development window was scored according to the vulva

shape[36] . Larvae were sorted using a gate enriched for L4.4-4.5 larvae. All conditions of the

same biological replicate were collected on the same day. 1,000 worm aliquots were harvested

and spun down for 0.5 min at 0.7 g to remove sorting liquid, and flash-frozen in dry ice.

<u>RNA fluorescent in situ hybridization (RNA FISH)</u>

RNA FISH was performed as described before[6]. For F07B7.1, probes were designed in the

exon regions that do not overlap with annotated transposable element sequences. We used

unlabeled primary probes and Cy5 fluorescently labeled secondary detector oligonucleotides.

Worms were harvested at 44 hph, incubated in fixative solution for 40 min at room temperature,

and stored in 70% Ethanol. Fixated larvae were washed once in wash buffer (10% formamide,

2× SSC buffer (Ambion)) and hybridized with the corresponding FLAP-containing probes in

100 µL hybridization buffer (10% dextran sulfate, 2 mM vanadyl-ribonucleoside complex,

0.02% RNAse-free BSA, 50 µg E. coli tRNA, 2× SSC, 10% formamide) at 30 ˚C overnight.

Hybridized embryos were washed twice with wash buffer, with DAPI in the second wash for

nuclei staining, and once in 2× SSC buffer before imaging. Right before imaging, embryos

were resuspended in 100 µL antifade buffer (0.4% glucose, 10 µM Tris-HCl pH 8, 2× SSC)

with 1 µL catalase (Sigma-Aldrich) and 1 µL glucose oxidase (3.7 mg/mL, Sigma-Aldrich).

Imaging was performed in an Andor BC43 spinning disk confocal microscope, and image

processing was done in FIJI[37] .

<u>Germline nuclei isolation.</u>

260,000 starved L1 larvae were plated in NGM-ethanol (control) or NGM-auxin plates

(HRDE-1 or SET-25 germline-depletion) and grown for 44 h for L4 larvae harvesting. Worms

were washed out from plates and rinsed three times with 1X M9 buffer by centrifuging for 30

seconds at 0.7 g. After the last wash, M9 was removed as much as possible, and the worms

were resuspended in 1.5 mL of cold Nuclear Extraction Buffer (NEB)[14] with 1X EDTA-free

Protease Inhibitors (Thermo Scientific, 78425). Worms were lysed in a Steel-Dounce diameter XX by applying 65-70 strokes. Worm lysis was monitored by inspection under a stereoscope (Zeiss), and the presence of GFP-positive nuclei was determined by mixing 2 µl of lysate with 2 µl of VECTASHIELD® Antifade Mounting Medium with DAPI (Vector Laboratories) and inspection under a Zeiss Axio Imager M2 microscope equipped with a Princeton Instrument PIXIS 926 1024 camera, using Plan-Apochromat 20x/1.40 objectives. Lysate was split into ~300 µl aliquots in six tubes, and 1 mL of NEB was added. The lysate was cleared by 4-minute centrifugation at 100 g and 4°C. Approximately 800-1,000 µl of supernatant were transferred to a new tube and centrifuged for 4 minutes at 1,000 g and 4 °C to pellet the nuclei. Supernatant was discarded, and the pellet was resuspended in ~ 100 µl of the remaining NEB buffer and pooled into two tubes. Nuclei were pelleted by 4-minute centrifugation at 1,000 g and 4°C, resuspended in NEB without glycerol and 1X EDTA-free Protease Inhibitors, and kept on ice until sorting.

Germline nuclei sorting.

Nuclei suspension was filtered and diluted 1:3-1:5 in cold NEB without glycerol and 1X EDTA-free Protease Inhibitors (Thermo Scientific™, 78425). Nuclei were analyzed and sorted using an Aria FACS III system (BD Biosciences). Aliquots of 500,000 GFP-positive nuclei were collected in 1.5 mL tubes with 100 µl of NEB without glycerol and 1X EDTA-free Protease Inhibitors and kept on ice. After sorting, four to six aliquots were centrifuged for 4 minutes at 1,100 g and 4°C, the supernatant was discarded, leaving ~50-100 µl to not perturb the nuclei pellet (not visible). Nuclei were gently resuspended avoiding the formation of bubbles and pooled into one tube. Nuclei were pellet by centrifuging for 4 minutes at 1,100 g and 4°C, the supernatant was discarded, and nuclei were resuspended in 100 µl of NEB with glycerol and 1X EDTA-free Protease Inhibitors per initial 500,000 nuclei and stored at -20°C until further processing.

For RNA extraction, nuclei were resuspended in 50 µl per 500,000 initial nuclei, and 1 mL of

TriZol was added before freezing.

For GRO-seq, nuclei were resuspended in 30 µL of GRO-seq Nuclear freezing buffer[14] and

nuclear run-on was performed immediately, as described below.

<u>Western Blot</u>

Western blots were performed to monitor AID:Flag:HRDE-1 levels and its depletion upon

auxin exposure; control worms were collected in parallel. For sorted larvae, 1,000 worms were

resuspended in 50 µl of 4X NuPage LDS and 1X NuPage Reducing agent (Thermo

Scientific™). For larvae used for nuclei preparation, 5 to 10 µl of packed worms were taken

before extraction and mixed with 100 µl of 4X LDS and 1X Reducing agent. Worms were lysed

by 10 minutes incubation at 95ºC. Then centrifuged for 10 minutes at 20ºC and 12000 g, ~30-

50% of sample was used for SDS-PAGE electrophoresis in 4-12% NuPage polyacrylamide

gels. Gels were run 50 min at 180V in 1X MOPS buffer (Thermo Scientific™), and transfer to

Nylon membrane by wet transfer in 1X NuPage wet transfer with 10% Methanol at 4ºC for 2.5

hours at 30 V. Transfers were monitored with Ponceau staining, and unstained before blocking

with 5% blocking reagent for 30 min at room temperature. After the membranes were rinsed

twice with 1X PBS and 0.1% Tween 20, all incubations were performed in the same buffer.

Primary antibody incubation was performed overnight at 4 °C with agitation. After, membranes

were washed four times before secondary antibody incubation for 2 hours at room temperature

and agitation. Antibody dilutions used: mouse anti-Flag 1:5000 (Sigma, F1804), mouse anti-

GADPH (Abcam, AB125247) 1:2000 and the secondary antibodies used were goat anti-mouse

Alexa Fluor 488 (Invitrogen, A11001) and goat anti-rabbit Alexa Fluor 568 (Invitrogen

A11011) antibodies at a dilution of 1:10000.

<u>RNA extraction.</u>

1 mL of TriZol reagent (Invitrogen) was used following the manufacturer's instructions. When

extracting RNA from whole worms, 1000-2000 L4 sorted larvae or 10 µl of L4 (44 hph) packed

worms were mixed with TriZol, and subjected to four to five cycles of freeze-thaw in dry ice

and 2 minutes of agitation at 25ºC and 2000 rpm in a Thermomixer (Eppendorf). Then, 100 µl

of 1-Bromo-3-chloropropane (Sigma, B9673) was added and vortexed. After 15 minutes of

incubation at room temperature, samples were centrifuged for 15 minutes at 12000 g and 4ºC,

and the aqueous phase was transferred to a new tube. 2 µl of glycoblue (Invitrogen, AM9515)

and 1 volume of Isopropanol (Sigma) were added and vortexed to mix. Samples were incubated

at room temperature for 15 min and RNA was precipitated by centrifuging 15 min at 12000 g

and 4ºC. Pellet was washed twice in 75% Ethanol and centrifuging 5 minutes at 7500 g and

4ºC, pellet was resuspended in nuclease-free water. Concentration and quality were estimated

using Nanodrop (Thermo Scientific™).

RT-qPCR.

Reverse transcription was performed using M-MLV reverse transcriptase (Invitrogen), and

qPCR was performed using Applied Biosystems PowerUp SYBR Green PCR Master mix and

using QuantStudio 3 Real-Time PCR System (Applied Biosystems).

Total RNA library preparation.

Total RNA was extracted as detailed above. RNA was treated with 1 unit of Turbo DNAse

(Invitrogen, AM2238) for 30 min at 37ºC and immediately purified using 1.8X ArgentCourt

RNA purification beads and 3X Isopropanol. This step was skipped for RNA extracted from

isolated germline nuclei. RNA integrity number was assessed with Agilent Tapestation 2200

and used to determine the fragmentation time during library preparation. 100 ng of total RNA

from whole worm or RNA from 500,000 germline nuclei were subjected to ribosomal RNA

depletion using DNA probed and DNAse H as described before[16]. DNA probes were degraded

by Turbo DNAse treatment and bead purification as explained before. Strand-specific libraries

were prepared with NEBNext® Ultra™ II Directional RNA Library Prep Kit (E7770).

<u>Small RNA library preparation.</u>

Libraries were prepared as described before[16]. Briefly, around 5 µg of total RNA was mixed

1:1 volume with 2X Novex® TBE-Urea Sample Buffer, denatured for 3 min at 95ºC, and kept

on ice until gel loading. RNAs were size separated in 15% NuPage 7% Urea polyacrylamide

gels (Invitrogen) in 1X TBE buffer for 45 min at 180V. The RNA cut and extracted from gel

was precipitated with 2 µL of Glycoblue (Invitrogen, AM9515) and 3 volumes of 100%

Ethanol. After incubation on ice, the samples were centrifuged and washed once with 75%

ethanol. Isolated small RNAs were treated with 5' Polyphosphatase, ligated with 3' and 5' end

adaptors, and amplified by PCR using Illumina indexed primers. PCRs were purified with

ArgentCourt DNA beads and analyzed using Tapestation. Libraries were pooled and size

selected using Pippinprep. Libraries were purified again using ArgentCourt beads and

quantified using Qubit to estimate molarity, based on a library size of 148 bp. An equimolar

proportion of each library was mixed for sequencing.

<u>GRO-seq library preparation.</u>

GRO-seq was performed as previously described[14] . Briefly, nuclear Run-On reactions were

performed by incorporating 1 mM Bio-11UTP, followed by RNA extraction with Trizol and

RNA fragmentation. Biotinylated nascent RNA was then purified using DynabeadTM

MyOneTM Streptavidin C1 (Cat#65001). To prepare libraries, purified RNAs were first treated

with T4 Polynucleotide Kinase (New England Biolabs) to repair the 5'-OH ends. This was

followed by 3' and 5' adaptor ligation. Adaptor-ligated RNAs were then reverse transcribed

using SuperScript IV Reverse Transcriptase (Thermo Fisher Scientific), following the

manufacturer's protocol with slight modifications: the reaction was incubated for 1 hour at

50°C and 10 minutes at 80°C. cDNA was PCR amplified using specific primers and the

NEBNext Ultra II Q5 Master Mix 2x (New England Biolabs) for 15 cycles. Libraries were analyzed using the Agilent 2200 TapeStation System with high sensitivity D1000 screentapes and quantified with the Qubit Fluorometer High Sensitivity dsDNA assay kit (Thermo Fisher Scientific, Q32851). The multiplexed libraries were then sequenced on the Illumina NextSeq 2000 platform.

CUT&Tag library preparation

Cut&Tag was performed following the EpiCypher protocol with few modifications. Briefly. 100 μL of the nuclei resuspension generated as described above was used for binding to Concanavalin A beads (Epicypher 21-1401). Bead-bound nuclei were resuspended in 100 μL of Antibody150 buffer (20 mM HEPES pH 7.5, 150 mM NaCl, 0.01% digitonin, 2 mM EDTA, protease inhibitors, 0.5 mM spermidine), and 1 μg of primary antibody was added. Reactions were incubated overnight in a thermocycler shaker set up at 4°C with shaking cycles at 2000 rpm, 15 seconds ON 45 seconds OFF. 1 μg of secondary antibody was added, followed by tagmentation with Tn5 (Epicypher 15-1117). Resulting libraries were amplified with i5 and i7 primers using 13 to 15 PCR cycles and purified using AMPure beads. Libraries were analyzed using the Agilent 2200 TapeStation System with high-sensitivity D1000 screentapes and quantified with the Qubit Fluorometer High Sensitivity dsDNA assay kit (Thermo Fisher Scientific, Q32851). Four biological replicates were performed for each experiment.

Library sequencing.

All library preparations were compatible for Illumina sequencing using NextSeq2000 sequencers and conventional or XLEAP-SBS technology, P1 or P2 100 cycles NextSeq2000 reagents.

Sequencing data analysis

All the sequencing data were demultiplexed with Illumina bcl2fastq converter (version v2.17.1.14), and quality control was performed with fastQC (version v0.11.5).

RNA-seq analysis

HISAT2 version 2.0.4[38] was used for mapping RNA-seq reads aligned to the *C. elegans* genome

sequence (ce11, *C. elegans* Sequencing Consortium WBcel235). After alignment, reads

mapping to annotated protein-coding genes were counted using featureCounts (version

2.0.1)[39]. Counted reads for protein-coding genes were used for differential expression analysis

using the R/Bioconductor package DESeq2 (version 1.26.0)[40].

Small RNA-seq analysis

The analysis of small RNA sequencing was performed as previously described[16].

GRO-seq analysis

Analysis of the sequencing data was performed as previously described[43] . The scripts and

workflows are available at https://gitlab.pasteur.fr/bli/bioinfo_utils.

CUT and Tag

Reads were aligned with Bowtie2[42] to the WBcel235 genome, with the following options: "--

local --very-sensitive-local --dovetail --soft-clipped-unmapped-tlen". Reads were additionally

deduplicated using samtools and filtered to keep those with a single discovered alignment.

Peaks were called using MACS2 with the following parameters: "callpeak -q 0.05 -g ce --keep-

dup all". Peaks were further filtered to keep those with MACS2 FDR < 0.01 in merged reads

from all conditions, and FDR < 0.05 in at least two individual replicates. Quantifications for

the selected set of peaks were done for each dataset using Subread featureCounts and analyzed

using a DESeq2 generalized linear model.

<u>Statistics and reproducibility</u>

All the experiments shown in this study were performed independently at least twice, and the

exact number of biological replicates is indicated on each figure legend.

<u>Data and code availability</u>

All the sequencing data are available upon request.

Acknowledgments

We would like to thank all the members of the Cecere laboratory for the helpful discussions on

the manuscript and suggestions on the project. We thank Dr. Florian Muller, head of the

Photonic BioImaging platform, and Dr. Gaëlle Letort, research engineer in the Department of

Developmental and Stem Cell Biology, for their assistance on RNA FISH imaging

interpretation. We thank the Caenorhabditis Genetics Center (CGC) for *C. elegans* strains,

supported by the National Institutes of Health - Office of Research Infrastructure Programs

(P40 OD010440). We thank Carolyn Phillips for sharing rde-8 and nyn-1:nyn-2 strains. We

thank the Flow Cytometry, Photonic BioImaging, and Biomics facilities at Institut Pasteur. This

project has received funding from the Institut Pasteur, the CNRS, and the European Union's

Horizon 2020 research and innovation program (grant agreement no. 101002999). A.M.L.R.

received funding from the European Union under the HORIZON-MSCA-2022-PF-01 program

no. 101109836 — MOMENTS and A.K.M. received funding from LabEx Revive fellowship

ANR-10-LBX-73.

Author contributions

A.M.L.R and G.C. identified and developed the core questions of the work and supervised the

whole project, contributed to the data analysis, and wrote the manuscript. A.K.M. developed

the germline nuclei isolation protocol, and together with J.J. and A.M.L.R., adapted it for

downstream genomic approaches (total RNA-seq, CUT&Tag, and GRO-seq). P.Q. and

A.M.L.R. performed antibody screenings for the CUT&Tag protocol. A.C. developed pipelines

for small RNA-seq, RNA-seq, and CUT&Tag analysis, and assisted A.M.L.R. in downstream

data examination. A.M.L.R. and J.J. performed CUT&Tag experiments. A.M.L.R. performed

RNA, small RNA, GRO-seq, and RNA FISH experiments. L.B. performed together with O.F.

the RNAi inheritance assay. A.M.L.R. and S.I. performed the polyUGylation assays. A.M.L.R.

and S.C. performed RT-qPCR of AID and mutant strains. L.B. assisted A.M.L.R. with the

COPAS operation for staged-worm sorting. L.B. and J.O. generated all the strains and crosses

used in this project and helped with worm growing and maintenance.

Competing interests

804 The authors declare no competing interests.

Materials & Correspondence

Correspondence and material requests should be addressed to Germano Cecere:

germano.cecere@pasteur.fr

References

1. Heard, E. & Martienssen, R. A. Transgenerational Epigenetic Inheritance: Myths and
Mechanisms. *Cell* **157**, 95–109 (2014).

2. Skvortsova, K., Iovino, N. & Bogdanović, O. Functions and mechanisms of epigenetic
inheritance in animals. *Nat Rev Mol Cell Bio* **19**, 774–790 (2018).

3. Cecere, G. Small RNAs in epigenetic inheritance: from mechanisms to trait transmission.
Febs Lett **595**, 2953–2977 (2021).

4. Buckley, B. A. *et al.* A nuclear Argonaute promotes multigenerational epigenetic
inheritance and germline immortality. *Nature* **489**, 447–451 (2012).

5. de Albuquerque, B. F. M., Placentino, M. & Ketting, R. F. Maternal piRNAs Are Essential
for Germline Development following De Novo Establishment of Endo-siRNAs in
Caenorhabditis elegans. *Dev Cell* **34**, 448–456 (2015).

6. Cornes, E. *et al.* piRNAs initiate transcriptional silencing of spermatogenic genes during C.
elegans germline development. *Dev Cell* **57**, 180-196.e7 (2022).

7. Schwartz-Orbach, L. *et al.* Caenorhabditis elegans nuclear RNAi factor SET-32 deposits
the transgenerational histone modification, H3K23me3. *Elife* **9**, e54309 (2020).

8. Gu, S. G. *et al.* Amplification of siRNA in Caenorhabditis elegans generates a
transgenerational sequence-targeted histone H3 lysine 9 methylation footprint. *Nat Genet* **44**,
157–164 (2012).

9. Kalinava, N., Ni, J. Z., Peterman, K., Chen, E. & Gu, S. G. Decoupling the downstream
effects of germline nuclear RNAi reveals that H3K9me3 is dispensable for heritable RNAi
and the maintenance of endogenous siRNA-mediated transcriptional silencing in
Caenorhabditis elegans. *Epigenetics Chromatin* **10**, 6 (2017).

10. Ouyang, J. P. T., Zhang, W. L. & Seydoux, G. The conserved helicase ZNFX-1
memorializes silenced RNAs in perinuclear condensates. *Nat. Cell Biol.* **24**, 1129–1140
(2022).

11. Rieger, I. *et al.* Nucleus-independent transgenerational small RNA inheritance in
Caenorhabditis elegans. *Sci. Adv.* **9**, eadj8618 (2023).

12. Kalinava, N. *et al.* C. elegans Heterochromatin Factor SET-32 Plays an Essential Role in
Transgenerational Establishment of Nuclear RNAi-Mediated Epigenetic Silencing. *Cell Rep.*
25, 2273-2284.e3 (2018).

13. Kaya-Okur, H. S. *et al.* CUT&Tag for efficient epigenomic profiling of small samples
and single cells. *Nat. Commun.* **10**, 1930 (2019).

14. Quarato, P. & Cecere, G. Global Run-On sequencing to measure nascent transcription in
C. elegans. *Star Protoc* **2**, 100991 (2021).

15. Zhang, L., Ward, J. D., Cheng, Z. & Dernburg, A. F. The auxin-inducible degradation
(AID) system enables versatile conditional protein depletion in C. elegans. *Dev. (Camb.,*
Engl.) **142**, 4374–4384 (2015).

16. Barucci, G. *et al.* Small-RNA-mediated transgenerational silencing of histone genes
impairs fertility in piRNA mutants. *Nat. Cell Biol.* **22**, 235–245 (2020).

17. Guang, S. *et al.* An Argonaute Transports siRNAs from the Cytoplasm to the Nucleus.
Science **321**, 537–541 (2008).

18. Tsanov, N. *et al.* smiFISH and FISH-quant – a flexible single RNA detection approach
with super-resolution capability. *Nucleic Acids Res.* **44**, e165–e165 (2016).

19. Chen, S. & Phillips, C. M. HRDE-2 drives small RNA specificity for the nuclear
Argonaute protein HRDE-1. *Nat. Commun.* **15**, 957 (2024).

20. Towbin, B. D. *et al.* Step-Wise Methylation of Histone H3K9 Positions Heterochromatin
at the Nuclear Periphery. *Cell* **150**, 934–947 (2012).

21. Greer, E. L. *et al.* A Histone Methylation Network Regulates Transgenerational
Epigenetic Memory in C. elegans. *Cell Rep.* **7**, 113–126 (2014).

22. Ding, Y.-H., Ochoa, H. J., Ishidate, T., Shirayama, M. & Mello, C. C. The nuclear
Argonaute HRDE-1 directs target gene re-localization and shuttles to nuage to promote small
RNA-mediated inherited silencing. *Cell Rep.* **42**, 112408 (2023).

23. Shukla, A. *et al.* poly(UG)-tailed RNAs in genome protection and epigenetic inheritance.
Nature **582**, 283–288 (2020).

24. Tsai, H.-Y. *et al.* A Ribonuclease Coordinates siRNA Amplification and mRNA
Cleavage during RNAi. *Cell* **160**, 407–419 (2015).

25. Phillips, C. M., Montgomery, T. A., Breen, P. C. & Ruvkun, G. MUT-16 promotes
formation of perinuclear Mutator foci required for RNA silencing in the C. elegans germline.
Gene Dev **26**, 1433–1444 (2012).

26. Wan, G. *et al.* Spatiotemporal regulation of liquid-like condensates in epigenetic
inheritance. *Nature* **557**, 679–683 (2018).

27. Ashe, A. *et al.* piRNAs Can Trigger a Multigenerational Epigenetic Memory in the
Germline of C. elegans. *Cell* **150**, 88–99 (2012).

28. Akay, A. *et al.* The Helicase Aquarius/EMB-4 Is Required to Overcome Intronic Barriers
to Allow Nuclear RNAi Pathways to Heritably Silence Transcription. *Dev Cell* **42**, 241-
255.e6 (2017).

29. Guang, S. *et al.* Small regulatory RNAs inhibit RNA polymerase II during the elongation
phase of transcription. *Nature* **465**, 1097–1101 (2010).

30. Shukla, A., Perales, R. & Kennedy, S. piRNAs coordinate poly(UG) tailing to prevent
aberrant and perpetual gene silencing. *Curr Biol* (2021) doi:10.1016/j.cub.2021.07.076.

31. Duempelmann, L. *et al.* Inheritance of a Phenotypically Neutral Epimutation Evokes
Gene Silencing in Later Generations. *Mol Cell* **74**, 534-541.e4 (2019).

32. Andersen, P. R., Tirian, L., Vunjak, M. & Brennecke, J. A heterochromatin-dependent
transcription machinery drives piRNA expression. *Nature* **549**, 54–59 (2017).

33. Stiernagle, T. Maintenance of C. elegans. *WormBook : online Rev. C elegans Biol.* 1–11
(2006) doi:10.1895/wormbook.1.101.1.

34. Paix, A., Folkmann, A., Rasoloson, D. & Seydoux, G. High Efficiency, Homology-
Directed Genome Editing in Caenorhabditis elegans Using CRISPR-Cas9 Ribonucleoprotein
Complexes. *Genetics* **201**, 47–54 (2015).

35. Dokshin, G. A., Ghanta, K. S., Piscopo, K. M. & Mello, C. C. Robust Genome Editing
with Short Single-Stranded and Long, Partially Single-Stranded DNA Donors in
Caenorhabditis elegans. *Genetics* **210**, 781–787 (2018).

36. Mok, D. Z. L., Sternberg, P. W. & Inoue, T. Morphologically defined sub-stages of C.
elegans vulval development in the fourth larval stage. *BMC Dev. Biol.* **15**, 26 (2015).

37. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat.*
Methods **9**, 676–682 (2012).

38. Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory
requirements. *Nat. Methods* **12**, 357–360 (2015).

39. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: An efficient general-purpose program
for assigning sequence reads to genomic features. *arXiv* (2013)
doi:10.48550/arxiv.1305.3347.

40. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

41. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing
and microarray studies. *Nucleic Acids Res.* **43**, e47–e47 (2015).

42. Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: precision weights unlock linear
model analysis tools for RNA-seq read counts. *Genome Biol.* **15**, R29–R29 (2014).

43. Quarato, P. *et al.* Germline inherited small RNAs facilitate the clearance of untranslated
maternal mRNAs in C. elegans embryos. *Nat Commun* **12**, 1441 (2021).

Figures

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Extended Data Figures.

Ext. Data Figure 1

Ext. Data Figure 2

Ext. Data Figure 3

Ext. Data Figure 4

Ext. Data Figure 5

Ext. Data Figure 6

Ext. Data Figure 7