Title: Brawn before bite in endemic Asian mammals after the end-Cretaceous extinction

Authors:

Z. Jack Tseng[1,2,5]*, Qian Li[3]**, Suyin Ting[2,4]

Affiliations:

[1]Department of Integrative Biology and Museum of Paleontology, University of California, Berkeley; California 94720, U.S.A.

[2]Department of Vertebrate Paleontology, Natural History Museum of Los Angeles County; California 90007, U.S.A.

[3]Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences; Beijing 100044, China

[4]Museum of Natural Science, Louisiana State University; Baton Rouge, Louisiana 70803, U.S.A.

[5]Lead Contact

*Correspondence: zjt@berkeley.edu

**Correspondence: liqian@ivpp.ac.cn

Summary: The first 10 million years (Myr) following the Cretaceous-Paleogene (K-Pg) mass extinction marked a period of global greenhouse conditions and dramatic rise of placental mammals. Because ~80% of known terrestrial sections capturing post-K-Pg mammal recovery come from North America, a substantial knowledge gap exists in the tempo and mode of recovery in Asia, where only 3% of sites are located and most contain species found nowhere else. We show that isolated Paleocene placental assemblages from China (1) reached high tooth size disparity early in the Paleocene, (2) tracked regional and global environmental changes in their dental shape later in the Paleocene, and (3) achieved maximum dental shape-performance integration near the end of the first 10 Myr post-K-Pg. This 'brawn before bite' transformation, coupled with

prolonged dental shape versus performance variability, favors a scenario whereby many

living orders of placental mammals were borne out of phenotypically and functionally

plastic ancestral assemblages, including those in tropical south China, during the

Paleocene.

Main Text:

The Cretaceous-Paleogene (K-Pg) mass extinctions accelerated the formation of

modern-day global biota. In particular, the more than 6,000 species of living placental

mammals trace their origins to the diversification of major orders around or after the K-

Pg boundary [1–3]. Different hypotheses (for example: long-fuse, explosive, and short-

fuse) about the timing of their adaptive radiation have been variably supported by

analyses of molecules, fossils, and model-based approaches [1,4–6]. Regardless of the

diversification scenarios favored by the competing explanations, the Paleocene epoch

(66-56 Myr ago) has been highlighted as the most critical time interval for establishing

phylogenetic and rate parameters for the placental adaptive radiation [5,7]. However,

substantial asymmetry exists in the quantity and quality of terrestrial fossil records that

document the first 10 Myr of the 'Age of Mammals', with ~80% of known terrestrial K-Pg

boundary sections occurring in North America [8]. Earliest Paleocene mammals are not

known from Europe [9,10], rendering detailed reconstructions of post-K-Pg recovery in that

continent difficult. The post-K-Pg fossil assemblages from Asia have hitherto not been

considered in analyses of post-K-Pg recovery dynamics [11].

The North American fossil record suggests that post-K-Pg mammal taxonomic

diversity recovery was relatively rapid at higher latitudes; most occurred within the first

~0.3 to 1 Myr of the Paleocene epoch [12–17]. This initial placental mammal diversification

was driven by archaic groups (i.e., stem lineages and those with no living relatives),

followed by the first appearance of many modern orders during two peak hyperthermal

events in the past 66 Myr, at the end-Paleocene and early Eocene climatic maxima,

respectively [18,19]. Distinct from these taxonomic recovery patterns, high selectivity of

mammalian ecomorphological extinction across the K-Pg boundary indicates a primary

productivity filter at the start of the Cenozoic Era (66 Myr ago to the present),

disproportionately impacting large herbivores and their predators [20]. Additionally, the

first 10 Myr of mammalian brain evolution after the K-Pg was marked by niche

partitioning along a size gradient; endocranial traits reflecting more complex sensory

processing did not appear until the Eocene (56 Myr ago). This phenomenon has been

termed the 'brawn before brains' hypothesis [21]. A similar pattern in initial size-driven

diversification followed by expansion of ecomorphological disparity is also observed in
mammal jaws, indicating a general dynamic across phenotypic systems [22]. Whether this
mode of evolution, whereby size disparity increase precedes other ecometric traits,
should be understood as a global phenomenon during the post-K-Pg placental radiation
remains untested, in large part because no such analyses have centered on non-North
American continental fossil mammal records.

We investigated dental topography-performance shifts and timing of
ecomorphological diversification by developing and leveraging the largest dataset to
date of Paleocene Asian mammal assemblages. Our analyses focused on three of the
most fossiliferous and biogeographically isolated Paleocene sedimentary sequences in
paleotropical Asia: The Nanxiong, Qianshan, and Chijiang Basins in present-day south
China [23–27] (Fig. S1). We generated a new phenotypic dataset of 200 Asian Paleocene
mammal teeth using high-resolution microcomputed tomography and laser scanning,
capturing 37 species endemic to low-latitude east Asia and which are brought to bear on
K-Pg recovery dynamics for the first time (Data S1-S2). We used dentitions as
ecological indicators [28] and examined temporal shifts in tooth crown complexity,
curvature, and height and their association with tooth deformation resistance using
topographic and simulation analyses. Our dataset spans the Paleocene, the first 10 Myr
of the Cenozoic, enabling us to test the hypothesis that dental topography and tooth
puncturing and shearing performance linkages showed delayed niche expansion
relative to body size disparity increase during this initial period of post-K-Pg mammal
recovery in Asia.

Dental traits paralleled Paleocene global and regional environmental conditions

We found that dental topographic trait variability in Paleocene mammals in south
China tracked global and regional climatic changes despite stasis in high-level
taxonomic composition over the course of the first 10 Myr after the K-Pg transition (Fig.
1). Dental height and sharpness variability were low in the beginning and end of the
time interval, with a spike in the middle Paleocene corresponding to a short-lived
negative excursion in global temperature. In contrast, elevated low-level taxonomic
turnover of genera and species within the Paleocene indicates cladogenetic shifts,

rather than anagenetic adaptation, underpin this dental topographic evolution (Data

S1)[11]. These findings suggest that in addition to its impact on crown group mammals,

the K-Pg extinctions and subsequent climatic fluctuations also filtered out archaic

placental mammals with lower speciation rates in favor of rapidly speciating taxa [4].

By contrast, there is no support for significant shifts in dental topographic trait

mean values from the early to the middle Paleocene for the majority of analytical

iterations of clade and dental data partitions (Figs 1A, S4; Table 1; Data S7). However,

in most analyses we observed a significant shift in at least one dental topographic

metric from the middle to the late Paleocene (Table 1; Data S7). The larger-bodied

Chinese endemic pantodont (an ungulate-like archaic placental group; "CEP") mammals

tend to increase dental complexity (OPCR, orientation patch count rotated) and

curvature (DNE, Dirichlet normal energy) whereas smaller, non-pantodonts (Chinese

arctostylopids and anagalids; rodent-like archaic placental groups) in the dataset

exhibited no significant dental traits shifts from the middle to late Paleocene (Fig. S4;

Data S7). The transformation by CEPs in complexity and curvature indices relates to the

capacity of teeth to resist wear and coincides with temperature and aridity increase

towards the end-Paleocene thermal maximum event (Figs. 1-2). Multiple geological and

geochemical proxies suggest that paleoclimate in and around the Nanxiong Basin K-Pg

section in south China reflects a latitudinally much broader global tropical zone during

the Paleocene [29] relative to present-day Earth, as well as rapid shifts between more

versus less humid intervals during the first 10 Myr of the Cenozoic [30]. Results from

feldspar-quartz (F:Q) ratios, clay mineral composition analysis, diffuse reflectance

spectroscopy (DRS), stable carbon isotope analysis, and total organic carbon (TOC)

analysis all support this regional paleoclimate profile [30,31]. Local climate reconstructions

for the latest Cretaceous indicate relatively warm and dry intervals, followed by warm

and humid climates during the earlier Paleocene, then a return to less humid but still

warm conditions in the later Paleocene [31]. Chinese endemic pantodont dentitions

tracked shifts in this paleoenvironmental progression.

The overall increase in dental complexity and curvature also coincided with an

increase in drought-tolerant flora in south China (Fig. 2) and specifically with

paleoenvironmental reconstructions in the Nanxiong Basin [32]. Palynological evidence

suggests that in addition to a predominance of broad-leaved, deciduous plants mixed

with a smaller percentage of conifers than in fossil localities further north [33], south China

also recorded drought resistant taxa such as the maidenhair fern *Pterisisporites* in the

early Paleocene Shanghu formation [34] and again in late Paleocene samples [35].

Expanded comparisons across time and space suggest fossil pollen taxa that are

indicators of seasonal aridity increased from 20.3% of late Paleocene pollen samples to

34.3% of early Eocene pollen samples [35]. These shifts appeared to have stabilized by

the end of the Eocene, when taxonomically modern floras became established in Asia

[36]. In addition to tracking Paleocene temperature trends (Fig. 1), Asian mammal dental

topography (OPCR and DNE in particular) also mimicked trends in the marine realm,

where global planktonic foraminifera records demonstrate a similar pattern in species

richness curve, as well as with global $\delta^{13}C$ [37,38] and south Pacific CO_2 levels [39]. These

broader associations offer strong indications that endemic Paleocene mammals in south

China composed a dynamic assemblage evolving in step with regional and global

environmental changes during the first 10 Myr after the K-Pg mass extinctions.

We detected a shift and expansion of mammal dental morphospace in the late

Paleocene (Fig. 2). An overall shift towards increased dental topographic trait

magnitudes in late Paleocene samples is driven mainly by CEPs (Table 1), despite the

fact that they constitute a minority of the overall data (41% of teeth) as well as late

Paleocene partition (21% of teeth). Additionally, dental metric disparity is significantly

higher in the late Paleocene partition than in the two preceding time bins (Table 1). This

pattern is driven both by increased dental curvature and complexity (DNE & OPCR) in

larger-bodied CEPs and crown height (RFI) in smaller-bodied non-pantodonts, in

addition to differences in disparity among the clades (Data S7; bootstrapped variance

for DNE, OPCR, and RFI are at least twice as large in non-pantodonts compared to

CEPs). This suggests that an expansion of dental disparity in the late Paleocene

occurred across the size spectrum of endemic Asian mammals. Over the same time

interval examined, body size disparity appeared to have reached maximum saturation

early on, in the early Paleocene (Fig. 1B, S6A, Table 1, Data S5), indicating that

substantial increases in the disparity of dental complexity, curvature, and height lagged

behind maximum size disparity during the Paleocene. This marked dental evolutionary

pattern of 'brawn before bite' in endemic Asian mammals mirrors the endocranial and

jaw functional morphology patterns identified in their North American and European

counterparts [21,22], underscoring that an initial size-driven post-K-Pg recovery followed by

ecomorphological radiation was a global phenomenon, even as regional tectonic events

such as the initial collision of the Indian subcontinent with Asia and Deccan Traps

volcanism influenced local mammal evolution [40,41].

Topography-performance integration underlies mammal dental shifts

Within an overall pattern of increasing integration between dental topographic

traits and bite performance traits across the Paleocene time intervals (Fig. 3; Table S1),

topographic versus performance trait variability shifted markedly both from early to

middle and from middle to late Paleocene time bins (Figs S3-S4;). Early Paleocene to

middle Paleocene DTA-FEA intra- and inter-correlations remained stable (Fig. 3B), but

DTA-FEA inter-correlations strengthened while intra-DTA correlations weakened in the

Middle to Late Paleocene transition (Fig. 3C). This transition pattern is in part due to the

divergence of shape-performance linkages in CEP versus non-pantodont mammals.

Dental topography variability tracked bite performance variability in non-pantodonts

through time, peaking in the middle Paleocene (Fig S4D). By contrast, both intra- and

inter-partition comparisons of topography and performance trends in CEPs showed a

complete decoupling to shifts through time (Fig. S4G). These findings indicate that form-

function malleability, the coexistence of distinct topography-performance relationships in

each time and taxon partition while overall integration between the two trait groups

increases between time bins, was present throughout the Paleocene. Complex form-

function linkages generally promote evolutionary redundancy and can enhance

optimization of phenotypic traits when selective trade-offs are present [42–44]. The

presence of functional redundancies underlies the high levels of dental topographic

variability in Asian Paleocene mammals. Alternatively, varying degrees of independence

between the two performance traits and dental topographic traits analyzed could allow

the two aspects of the dentition to evolve in a decoupled manner (Fig. 3).

South Asia as a Paleogene 'Garden of Eden' for placental mammals?

Among the most consequential implications of accurately interpreting post-K-Pg
mammalian recovery dynamics in Asia is the ability to reconstruct the evolutionary
conditions during the biogeographic origination of modern placental orders. Current
knowledge and an overwhelming majority of data about the first post-dinosaur mammal
ecological communities center on North American localities [12,45], and secondarily from
European records [46]. Yet, the fossil records of those continents pinpoint a high turnover
of mammal faunas at the Paleocene-Eocene boundary, driven by the Paleocene
origination of major modern mammal clades elsewhere and subsequent dispersal of
those lineages to the European and North American continents, respectively. At least
five living orders (Primates, Rodentia, Lagomorpha, Perissodactyla, Eulipotyphla),
representing over half of mammalian species richness today, trace their evolutionary
origins to Asia [18,47–49]. There is also mounting evidence that other organisms, including
fish and plant lineages, followed a similar biogeographic pattern [50].

The abrupt appearance of early representatives of modern mammal lineages in
North America has been formally operationalized by the 'East of Eden' model, wherein
Asia is the 'Garden of Eden' and a biodiversity pump of living orders of placental
mammals [51] and other organisms. Biogeographic modeling analysis of modern
mammalian diversity also strongly identifies tropical south Asia, including the
geographic regions occupied today by southern China, India, and southeast Asian
countries, as the cradle of mammalian diversification since the Paleocene [52]. As
geographically proximate faunal assemblages to the epicenter of the 'Garden of Eden'
hypothesis, the endemic Paleocene mammals of the Asian paleotropics analyzed in this
study exhibited a high degree of dental topographic variability while strengthening
topography-performance integration during a period of biogeographic isolation and
regional and global climate changes. This finding favors the scenario whereby many
living orders of placental mammals were borne out of phenotypically and functionally
plastic ancestral assemblages, including those that lived in south China, during the
Paleocene [53].

We further infer from these findings that episodes of fluctuating global warming
during the first 10 Myr of the Cenozoic promoted the evolution of 'all-purpose'

mammalian dentitions rather than those with specialized functions (Fig. 3). This
prolonged dental topographic variability tracks extended post-K-Pg floral recovery times
[35,54], and suggests a 'brawn before brains' mode of placental mammal evolution may in
part represent a series of correlated evolutionary shifts in sync with the steadily
increasing complexity of Paleocene primary producers [20,36]. That a global signal would
be detected in the splendidly isolated south Asian Paleocene assemblage is
unexpected, and indicates that effects of climate forcing on post-K-Pg mammal recovery
may be ubiquitous, as has been predicted for Earth's ongoing rapid environmental shifts
[55]. In response, Paleocene mammal clades in south China reached maximum tooth size
disparity early and maximum integration between dental topography and bite
performance later, all the while maintaining high levels of variability in dental complexity
and convexity (Fig. 1). These preconditions set the stage for the subsequent taxonomic
turnover between archaic and crown lineages during the Paleogene to Neogene
modernization of mammal communities [19,24]. As a primary interface against shifting food
resources and the environment, the placental dentition was poised to play an outsized
role in the explosive diversification following the K-Pg extinctions, with the masticatory
complex having already completed the macroevolutionary transition to a mechanically
stiff but phenotypically flexible jaw during the Mesozoic Era [56,57]. The end-Cretaceous
extinctions and the climatic volatility that followed then set in motion the ecological
release that allowed mammals to explore dental form-function to a far fuller extent in 66
Myr of the Cenozoic Era than during the preceding 150 Myr under the reign of
dinosaurs. The transformational evolutionary sequence of 'brawn before bite' in tooth
size, topographic, and performance evolution was likely central to this placental success
story.

References

1.   O'Leary, M.A., Bloch, J.I., Flynn, J.J., Gaudin, T.J., Giallombardo, A., Giannini,
N.P., Goldberg, S.L., Kraatz, B.P., Luo, Z.-X., and Meng, J. (2013). The placental
mammal ancestor and the post–K-Pg radiation of placentals. Science (80-.). *339*,
662–667.

2.   Liu, L., Zhang, J., Rheindt, F.E., Lei, F., Qu, Y., Wang, Y., Zhang, Y., Sullivan, C.,
Nie, W., Wang, J., et al. (2017). Genomic evidence reveals a radiation of
placental mammals uninterrupted by the KPg boundary. Proc. Natl. Acad. Sci.
114, E7282–E7290. https://doi.org/10.1073/pnas.1616744114.

3.   Foley, N.M., Mason, V.C., Harris, A.J., Bredemeyer, K.R., Damas, J., Lewin, H.A.,
Eizirik, E., Gatesy, J., Karlsson, E.K., Lindblad-Toh, K., et al. (2025). A genomic
timescale for placental mammal evolution. Science (80-.). *380*, eabl8189.
https://doi.org/10.1126/science.abl8189.

4.   Quintero, I., Lartillot, N., and Morlon, H. (2024). Imbalanced speciation pulses
sustain the radiation of mammals. Science (80-.). *384*, 1007–1012.

5.   Upham, N.S., Esselstyn, J.A., and Jetz, W. (2021). Molecules and fossils tell
distinct yet complementary stories of mammal diversification. Curr. Biol. *31*,
4195–4206.

6.   Springer, M.S., Foley, N.M., Brady, P.L., Gatesy, J., and Murphy, W.J. (2019).
Evolutionary models for the diversification of placental mammals across the KPg
boundary. Front. Genet. *10*, 1241.

7.   Halliday, T.J.D., Upchurch, P., and Goswami, A. (2017). Resolving the
relationships of Paleocene placental mammals. Biol. Rev. *92*, 521–550.
https://doi.org/https://doi.org/10.1111/brv.12242.

8.   Vajda, V., and Bercovici, A. (2014). The global vegetation pattern across the
Cretaceous–Paleogene mass extinction interval: a template for other extinction
events. Glob. Planet. Change *122*, 29–49.

9.   Csiki-Sava, Z., Buffetaut, E., Ősi, A., Pereda-Suberbiola, X., and Brusatte, S.L.

(2015). Island life in the Cretaceous-faunal composition, biogeography, evolution,
and extinction of land-living vertebrates on the Late Cretaceous European
archipelago. Zookeys, 1–161.

10. De Bast, E., and Smith, T. (2017). The oldest Cenozoic mammal fauna of Europe:
implication of the Hainin reference fauna for mammalian evolution and dispersals
during the Paleocene. J. Syst. Palaeontol. *15*, 741–785.

11. Wang, Y., Meng, J., Ni, X., and Li, C. (2007). Major events of Paleogene mammal
radiation in China. Geol. J. *42*, 415–430.

12. Wing, S.L., Alroy, J., and Hickey, L.J. (1995). Plant and mammal diversity in the
Paleocene to early Eocene of the Bighorn Basin. Palaeogeogr. Palaeoclimatol.
Palaeoecol. *115*, 117–155.

13. Alroy, J. (1999). The fossil record of North American mammals: evidence for a
Paleocene evolutionary radiation. Syst. Biol. *48*, 107–118.

14. Longrich, N.R., Scriberas, J., and Wills, M.A. (2016). Severe extinction and rapid
recovery of mammals across the Cretaceous–Palaeogene boundary, and the
effects of rarity on patterns of extinction and recovery. J. Evol. Biol. *29*, 1495–
1512.

15. Lyson, T.R., Miller, I.M., Bercovici, A.D., Weissenburger, K., Fuentes, A.J., Clyde,
W.C., Hagadorn, J.W., Butrim, M.J., Johnson, K.R., Fleming, R.F., et al. (2019).
Exceptional continental record of biotic recovery after the Cretaceous–Paleogene
mass extinction. Science (80-.). *366*, 977–983.
https://doi.org/10.1126/science.aay2268.

16. Wilson, G.P. (2014). Mammalian extinction, survival, and recovery dynamics
across the Cretaceous-Paleogene boundary in northeastern Montana, USA. In
Through the End of the Cretaceous in the Type Locality of the Hell Creek
Formation in Montana and Adjacent Areas, G. P. Wilson, W. A. Clemens, J. R.
Horner, and J. H. Hartman, eds. (Geological Society of America).

17. Grossnickle, D.M., and Newham, E. (2016). Therian mammals experience an

ecomorphological radiation during the Late Cretaceous and selective extinction at
the K–Pg boundary. Proc. R. Soc. B Biol. Sci. *283*, 20160256.
https://doi.org/10.1098/rspb.2016.0256.

18. Gingerich, P.D. (2010). Mammalian faunal succession through the Paleocene-
Eocene Thermal MAximum (PETM) in western North America. Vertebr. Palasiat.
48, 308–327.

19. Bowen, G.J., Clyde, W.C., Koch, P.L., Ting, S., Alroy, J., Tsubamoto, T., Wang,
Y., and Wang, Y. (2002). Mammalian dispersal at the Paleocene/Eocene
boundary. Science (80-.). *295*, 2062–2065.

20. Wilson, G.P. (2013). Mammals across the K/Pg boundary in northeastern
Montana, U.S.A.: dental morphology and body-size patterns reveal extinction
selectivity and immigrant-fueled ecospace filling. Paleobiology *39*, 429–469.
https://doi.org/DOI: 10.1666/12041.

21. Bertrand, O., Shelley, S., Williamson, T., Wible, J., Chester, S., Flynn, J.,
Holbrook, L., Lyson, T., Meng, J., Miller, I., et al. (2022). Brawn before brains in
placental mammals after the end-Cretaceous extinction. Science (80-.). *376*, 80–
85. https://doi.org/10.1126/science.abl5584.

22. Benevento, G.L., Benson, R.B.J., and Friedman, M. (2019). Patterns of
mammalian jaw ecomorphological disparity during the Mesozoic/Cenozoic
transition. Proc. R. Soc. B Biol. Sci. *286*, 20190347.
https://doi.org/10.1098/rspb.2019.0347.

23. Clyde, W.C., Ting, S., Snell, K.E., Bowen, G.J., Tong, Y., Koch, P.L., Li, Q., and
Wang, Y. (2010). New paleomagnetic and stable-isotope results from the
Nanxiong Basin, China: implications for the K/T boundary and the timing of
Paleocene mammalian turnover. J. Geol. *118*, 131–143.

24. Ting, S.-Y., Tong, Y.-S., Clyde, W.C., and Koch, P.L. (2011). Asian early
Paleogene chronology and mammalian faunal turnover events. Vertebr. Palasiat.
49, 1–28.

25.   Clyde, W.C., Tong, Y., Snell, K.E., Bowen, G.J., Ting, S., Koch, P.L., Li, Q., Wang,
Y., and Meng, J. (2008). An integrated stratigraphic record from the Paleocene of
the Chijiang Basin, Jiangxi Province (China): Implications for mammalian turnover
and Asian block rotations. Earth Planet. Sci. Lett. *269*, 554–564.

26.   Chow, M., Zhang, Y., Wang, B., and Ding, S. (1977). Paleocene mammalian
fauna from the Nanxiong Basin, Guangdong Province. Paleontol. Sin. New Ser. C
20, 1–100.

27.   Wang, Y.-Q., Li, C.-K., Li, Q., and Li, D.-S. (2016). A synopsis of Paleocene
stratigraphy and vertebrate paleontology in the Qianshan Basin, Anhui, China.
Vertebr. Palasiat. *54*, 89–120.

28.   Lucas, P., Constantino, P., Wood, B., and Lawn, B. (2008). Dental enamel as a
dietary indicator in mammals. BioEssays *30*, 374–385.
https://doi.org/https://doi.org/10.1002/bies.20729.

29.   Yin, Y., Zhang, L., Gu, X., Yin, R., Wen, Y., Jin, T., and Wang, C. (2023). High
terrestrial temperature in the low-latitude Nanxiong Basin during the Cretaceous-
Paleogene boundary interval. Palaeogeogr. Palaeoclimatol. Palaeoecol. *617*,
111489. https://doi.org/https://doi.org/10.1016/j.palaeo.2023.111489.

30.   Huang, H., Ma, M., Wang, M., and Liu, X. (2024). The paleoclimatic significance
indicated by the Early Paleogene gray-black mudstone layers lies in Nanxiong
Basin. Quat. Sci. *44*, 1250–1261.

31.   Wang, Y., Li, X., Zhou, Y., and Liu, L. (2015). Paleoclimate indication of
terrigenous clastic rock's component during the Late Cretaceous–Early
Paleocene in the Nanxiong Basin. Acta Sedimentol. Sin. *33*, 116–123.

32.   Ma, M., Liu, X., and Wang, W. (2018). Palaeoclimate evolution across the
Cretaceous–Palaeogene boundary in the Nanxiong Basin (SE China) recorded by
red strata and its correlation with marine records. Clim. Past *14*, 287–302.
https://doi.org/10.5194/cp-14-287-2018.

33.   Hao, H., Ferguson, D.K., Feng, G.-P., Ablaev, A., Wang, Y.-F., and Li, C.-S.

(2010). Early Paleocene vegetation and climate in Jiayin, NE China. Clim.

Change *99*, 547–566.

34. Xie, Y., Wu, F., and Fang, X. (2020). A major environmental shift by the middle

Eocene in southern China: Evidence from palynological records. Rev. Palaeobot.

Palynol. *278*, 104226.

35. Ma, X., Jiang, H., Cheng, J., and Xu, H. (2012). Spatiotemporal evolution of

Paleogene palynoflora in China and its implication for development of the

extensional basins in East China. Rev. Palaeobot. Palynol. *184*, 24–35.

36. Su, T., Spicer, R.A., Li, S.-H., Xu, H., Huang, J., Sherlock, S., Huang, Y.-J., Li, S.-

F., Wang, L., and Jia, L.-B. (2019). Uplift, climate and biotic changes at the

Eocene–Oligocene transition in south-eastern Tibet. Natl. Sci. Rev. *6*, 495–504.

37. MacLeod, N., Ortiz, N., Fefferman, N., Clyde, W., Schulter, C., MacLean, J.,

Culver, S.J., and Rawson, P.F. (2000). Phenotypic response of foraminifera to

episodes of global environmental change. Biot. response to Glob. Environ. Chang.

last *145*, 51–78.

38. Keller, G., Mateo, P., Punekar, J., Khozyem, H., Gertsch, B., Spangenberg, J.,

Bitchong, A.M., and Adatte, T. (2018). Environmental changes during the

Cretaceous-Paleogene mass extinction and Paleocene-Eocene thermal maximum:

implications for the Anthropocene. Gondwana Res. *56*, 69–89.

39. Hollis, C.J., Naeher, S., Clowes, C.D., Naafs, B.D.A., Pancost, R.D., Taylor,

K.W.R., Dahl, J., Li, X., Ventura, G.T., and Sykes, R. (2022). Late Paleocene CO2

drawdown, climatic cooling and terrestrial denudation in the southwest Pacific.

Clim. Past *18*, 1295–1320. https://doi.org/10.5194/cp-18-1295-2022.

40. Hu, X., Garzanti, E., Moore, T., and Raffi, I. (2015). Direct stratigraphic dating of

India-Asia collision onset at the Selandian (middle Paleocene, 59±1 Ma). Geology

43, 859–862.

41. Ma, M., Wang, M., Huang, H., and Liu, X. (2024). Terrestrial records of two

hyperthermal events in the Cretaceous-Paleogene boundary suggest different

control mechanisms. Commun. Earth Environ. *5*, 1–7.

42.   Wainwright, P.C., Alfaro, M.E., Bolnick, D.I., and Hulsey, C.D. (2005). Many-to-
One Mapping of Form to Function: A General Principle in Organismal Design?
Integr. Comp. Biol. *45*, 256–262. https://doi.org/10.1093/icb/45.2.256.

43.   Polly, P.D., Stayton, C.T., Dumont, E.R., Pierce, S.E., Rayfield, E.J., and
Angielczyk, K.D. (2016). Combining geometric morphometrics and finite element
analysis with evolutionary modeling: towards a synthesis. J. Vertebr. Paleontol. *36*,
e1111225.

44.   Pollock, T.I., Deakin, W.J., Chatar, N., Milla Carmona, P.S., Rovinsky, D.S.,
Panagiotopoulou, O., Parker, W.M.G., Adams, J.W., Hocking, D.P., Donoghue,
P.C.J., et al. (2025). Functional optimality underpins the repeated evolution of the
extreme saber-tooth morphology. Curr. Biol. *35*, 455-467.e6.
https://doi.org/10.1016/j.cub.2024.11.059.

45.   Rose, K.D. (1981). Composition and species diversity in Paleocene and Eocene
mammal assemblages: an empirical study. J. Vertebr. Paleontol. *1*, 367–388.
https://doi.org/10.1080/02724634.1981.10011907.

46.   Hooker, J.J. (2000). Paleogene mammals: crises and ecological change. Biot.
response to Glob. Chang. last *145*, 333–349.

47.   Beard, K.C. (1998). East of Eden: Asia as an important center of taxonomic
origination in mammalian evolution. Bull. Carnegie Museum Nat. Hist. *34*, 5–39.

48.   Maas, M.C., Krause, D.W., and Strait, S.G. (1988). The decline and extinction of
Plesiadapiformes (Mammalia:? Primates) in North America: displacement or
replacement? Paleobiology *14*, 410–431.

49.   Lopatin, A. V (2006). Early Paleogene insectivore mammals of Asia and
establishment of the major groups of Insectivora. Paleontol. J. *40*, S205–S405.
https://doi.org/10.1134/S0031030106090012.

50.   Culver, S.J., and Rawson, P.F. (2006). Biotic response to global change: the last
145 million years (Cambridge University Press).

51.   Beard, C. (2002). East of Eden at the Paleocene/Eocene boundary. Science (80-
.). *295*, 2028–2029.

52.   Feijó, A., Ge, D., Wen, Z., Cheng, J., Xia, L., Patterson, B.D., and Yang, Q.
(2022). Mammalian diversification bursts and biotic turnovers are synchronous
with Cenozoic geoclimatic events in Asia. Proc. Natl. Acad. Sci. *119*,
e2207845119. https://doi.org/10.1073/pnas.2207845119.

53.   McKenna, M.C., Chow, M., Ting, S., and Luo, Z. (1989). Radinskya yupingae a
perissodactyl-like mammal from the Late Paleocene of China. In The evolution of
perissodactyls, pp. 24–36.

54.   Peppe, D.J. (2010). Megafloral change in the early and middle Paleocene in the
Williston Basin, North Dakota, USA. Palaeogeogr. Palaeoclimatol. Palaeoecol.
298, 224–234.

55.   Scheffers, B.R., De Meester, L., Bridge, T.C.L., Hoffmann, A.A., Pandolfi, J.M.,
Corlett, R.T., Butchart, S.H.M., Pearce-Kelly, P., Kovacs, K.M., Dudgeon, D., et al.
(2016). The broad footprint of climate change from genes to biomes to people.
Science (80-.). *354*, aaf7671. https://doi.org/10.1126/science.aaf7671.

56.   Tseng, Z.J., Garcia-Lara, S., Flynn, J.J., Holmes, E., Rowe, T.B., and Dickson, B.
V (2023). A switch in jaw form–function coupling during the evolution of mammals.
Philos. Trans. R. Soc. B Biol. Sci. *378*, 20220091.
https://doi.org/10.1098/rstb.2022.0091.

57.   Schultz, J.A. (2025). A perspective from the Mesozoic: Evolutionary changes of
the mammalian skull and their influence on feeding efficiency and high-frequency
hearing. Anat. Rec., 1–16. https://doi.org/https://doi.org/10.1002/ar.25652.

58.   Zachos, J., Pagani, M., Sloan, L., Thomas, E., and Billups, K. (2001). Trends,
rhythms, and aberrations in global climate 65 Ma to present. Science (80-.). *292*,
686–693.

59.   Speijer, R.P., Pälike, H., Hollis, C.J., Hooker, J.J., and Ogg, J.G. (2020). The
Paleogene Period (Chapter 28). In Geological Time Scale, F. M. Gradstein, J. G.

Ogg, M. D. Schmitz, and G. M. Ogg, eds. (Elsevier), pp. 1087–1140.

60.   Cande, S.C., and Kent, D. V (1995). Revised calibration of the geomagnetic
436   polarity timescale for the Late Cretaceous and Cenozoic. J. Geophys. Res. Solid
437   Earth *100*, 6093–6095.

438   61.   Pampush, J.D., Winchester, J.M., Morse, P.E., Vining, A.Q., Boyer, D.M., and Kay,
439   R.F. (2016). Introducing molaR: a New R Package for Quantitative Topographic
440   Analysis of Teeth (and Other Topographic Surfaces). J. Mamm. Evol. *23*, 397–
441   412. https://doi.org/10.1007/s10914-016-9326-0.

442   62.   Bunn, J.M., Boyer, D.M., Lipman, Y., St. Clair, E.M., Jernvall, J., and Daubechies,
443   I. (2011). Comparing Dirichlet normal surface energy of tooth crowns, a new
444   technique of molar shape quantification for dietary inference, with previous
445   methods in isolation and in combination. Am. J. Phys. Anthropol. *145*, 247–261.

446   63.   Waldman, E., Gonzalez, Y., Flynn, J.J., and Tseng, Z.J. (2023). Dental
447   topographic proxies for ecological characteristics in carnivoran mammals. J. Anat.
448   *242*, 627–641. https://doi.org/https://doi.org/10.1111/joa.13806.

449   64.   Berthaume, M.A., Lazzari, V., and Guy, F. (2020). The landscape of tooth shape:
450   Over 20 years of dental topography in primates. Evol. Anthropol. Issues, News,
451   Rev. *29*, 245–262.

452   65.   Evans, A.R., Wilson, G.P., Fortelius, M., and Jernvall, J. (2007). High-level
453   similarity of dentitions in carnivorans and rodents. Nature *445*, 78–81.

454   66.   DeMers, A.C., and Hunter, J.P. (2024). Dental complexity and diet in amniotes: A
455   meta-analysis. PLoS One *19*, e0292358.

456   67.   Crofts, S.B., Smith, S.M., and Anderson, P.S.L. (2020). Beyond Description: The
457   Many Facets of Dental Biomechanics. Integr. Comp. Biol. *60*, 594–607.
458   https://doi.org/10.1093/icb/icaa103.

459   68.   Crompton, A.W., and Hiiemae, K. (1969). How mammalian molar teeth work.
460   Discovery *5*, 23–34.

69. Bhullar, B.-A.S., Manafzadeh, A.R., Miyamae, J.A., Hoffman, E.A., Brainerd, E.L.,
Musinsky, C., and Crompton, A.W. (2019). Rolling of the jaw is essential for
mammalian chewing and tribosphenic molar function. Nature *566*, 528–532.
https://doi.org/10.1038/s41586-019-0940-x.

70. Simpson, J.J. (1936). Studies of the earliest mammalian dentitions. Dent. Cosm.
78, 791–800.

71. Yamanaka, A., Haider, Y., Morita, W., Corfe, I., Nakamura, N., and Goto, T.
(2024). Developmental process of the modern house shrew's molars: implications
for the evolution of the tribosphenic molar in Mesozoic mammals. Evolution (N. Y).
78, 463–479. https://doi.org/10.1093/evolut/qpad228.

72. Lucas, P.W., and van Casteren, A. (2014). The Wear and Tear of Teeth. Med.
Princ. Pract. *24*, 3–13. https://doi.org/10.1159/000367976.

73. van Casteren, A., and Crofts, S.B. (2019). The Materials of Mastication: Material
Science of the Humble Tooth. Integr. Comp. Biol. *59*, 1681–1689.
https://doi.org/10.1093/icb/icz129.

74. Crofts, S. (2015). Finite element modeling of occlusal variation in durophagous
tooth systems. J. Exp. Biol. *218*, 2705–2711. https://doi.org/10.1242/jeb.120097.

75. Pérez-Ramos, A., Romero, A., Rodriguez, E., and Figueirido, B. (2020). Three-
dimensional dental topography and feeding ecology in the extinct cave bear. Biol.
Lett. *16*, 20200792. https://doi.org/10.1098/rsbl.2020.0792.

76. Pollock, T.I., Panagiotopoulou, O., Hocking, D.P., and Evans, A.R. (2022). Taking
a stab at modelling canine tooth biomechanics in mammalian carnivores with
beam theory and finite-element analysis. R. Soc. Open Sci. *9*, 220701.
https://doi.org/10.1098/rsos.220701.

77. Villalobos-Chaves, D., and Santana, S.E. (2022). Craniodental traits predict
feeding performance and dietary hardness in a community of Neotropical free-
tailed bats (Chiroptera: Molossidae). Funct. Ecol. *36*, 1690–1699.
https://doi.org/https://doi.org/10.1111/1365-2435.14063.

78.   Sender, R.S., and Strait, D.S. (2023). The biomechanics of tooth strength: testing
490   the utility of simple models for predicting fracture in geometrically complex teeth. J.
491   R. Soc. Interface *20*, 20230195. https://doi.org/10.1098/rsif.2023.0195.

492   79.   Berthaume, M.A. (2016). On the Relationship Between Tooth Shape and
493   Masticatory Efficiency: A Finite Element Study. Anat. Rec. *299*, 679–687.
494   https://doi.org/https://doi.org/10.1002/ar.23328.

495   80.   Ungar, P.S. (2015). Mammalian dental function and wear: A review. Biosurface
496   and Biotribology *1*, 25–41.
497   https://doi.org/https://doi.org/10.1016/j.bsbt.2014.12.001.

498   81.   Dumont, E.R. (1995). Enamel Thickness and Dietary Adaptation among Extant
499   Primates and Chiropterans. J. Mammal. *76*, 1127–1136.
500   https://doi.org/10.2307/1382604.

501   82.   Shellis, R.P., Beynon, A.D., Reid, D.J., and Hiiemae, K.M. (1998). Variations in
502   molar enamel thickness among primates. J. Hum. Evol. *35*, 507–522.

503   83.   Dumont, E.R., Grosse, I.R., and Slater, G.J. (2009). Requirements for comparing
504   the performance of finite element models of biological structures. J. Theor. Biol.
505   *256*, 96–103. https://doi.org/10.1016/j.jtbi.2008.08.017.

506   84.   He, L.H., Fujisawa, N., and Swain, M. V (2006). Elastic modulus and stress–strain
507   response of human enamel by nano-indentation. Biomaterials *27*, 4388–4398.
508   https://doi.org/https://doi.org/10.1016/j.biomaterials.2006.03.045.

509   85.   Conith, A.J., Imburgia, M.J., Crosby, A.J., and Dumont, E.R. (2016). The
510   functional significance of morphological changes in the dentitions of early
511   mammals. J. R. Soc. Interface *13*, 20160713.
512   https://doi.org/10.1098/rsif.2016.0713.

513   86.   Lee, J.J.-W., Constantino, P.J., Lucas, P.W., and Lawn, B.R. (2011). Fracture in
514   teeth—a diagnostic for inferring bite force and tooth function. Biol. Rev. *86*, 959–
515   974. https://doi.org/https://doi.org/10.1111/j.1469-185X.2011.00181.x.

87.   Irschick, D.J., Meyers, J.J., Husak, J.F., and Le Galliard, J.-F. (2008). How does

selection operate on whole-organism functional performance capacities? A review

and synthesis. Evol. Ecol. Res. *10*, 177–196.

88. Team, R.C. (2025). R: A Language and Environment for Statistical Computing at

R Foundation for Statistical Computing, Vienna, Austria.

89. Adams, D.C., and Otárola-Castillo, E. (2013). geomorph: an r package for the

collection and analysis of geometric morphometric shape data. Methods Ecol.

Evol. *4*, 393–399. https://doi.org/https://doi.org/10.1111/2041-210X.12035.

90. Klingenberg, C.P. (2014). Studying morphological integration and modularity at

multiple levels: concepts and analysis. Philos. Trans. R. Soc. B Biol. Sci. *369*,

20130249. https://doi.org/10.1098/rstb.2013.0249.

91. Wei, T., and V, S. (2024). R package "corrplot": Visualization of a Correlation

Matrix.

Acknowledgments:

We thank I. Ruiz for assisting with generating 3D tooth models from CT data; X. Tseng

for assisting with data collection; P. Holroyd for assisting with locating specimen casts in

the collections of the University of California Museum of Paleontology; J. Liu for helpful

discussion and comments on earlier versions of the study. Z. Luo provided earnest and

constructive feedback on an earlier version of the study.

Funding: This study was funded in part by grant 213109 from the Nanjing

Institute of Geology and Paleontology, Chinese Academy of Sciences (to QL and

ZJT) and National Key Research and Development Project of China

(2024YFF0807603; to QL).

Author contributions:

Conceptualization: All authors.

Methodology: ZJT.

Formal analyses: ZJT.

Investigation: All authors.

Results acquisition: ZJT.

Raw data acquisition (surface and CT scan data): ZJT, QL.

Writing, original draft: ZJT.

Writing, review, and editing: All authors.

Visualization: ZJT, QL.

Competing interests: The authors declare that they have no competing interests.

Resource availability:

-    Requests for further information and resources should be directed to and will be
fulfilled by the lead contact, Jack Tseng (zjt@berkeley.edu).
-    Code associated with analyses is included with this article.
-    Raw data are available in the supplementary materials.
-    All original fossil specimens are accessioned in the Institute of Vertebrate
Paleontology and Paleoanthropology, Chinese Academy of Sciences.

Supplemental information

Document S1. Figures S1–S6, Tables S1 and S2, and supplemental references

Data S1

Data S2

Data S3

Data S4

Data S5

Data S6

Data S7

Data S8

Data S9

Data S10

Figure Legends

Fig. 1. Temperature and fossil mammal dental trait shifts during the first 10 Myr of the Cenozoic. (A) Dental topographic trait values (boxplots) and mean variance (red curves) during the first 10 Myr of the Cenozoic, signifying the time after the K-Pg mass extinctions and before the Paleocene-Eocene hyperthermal event. Global temperature curve based on Zachos et al. [58]. Dental traits measured include crown complexity (OPCR, orientation patch count rotated), curvature (DNE, Dirichlet normal energy), height (RFI, relief index), and slope. (B) Mammal tooth size distributions represented by log 10 square root tooth area, in units of log10 millimeters. (C) Variance of compressive bite performance based on tooth crown finite element simulations, in units of squared Joules. (D) Variance of shear bite performance based on tooth crown finite element simulations, in units of squared Joules. Examples of endemic Asian fossil specimens analyzed: (E) Lateral and ventral views of early Paleocene Chinese endemic pantodont (CEP) *Bemalambda nanhsiungensis* IVPP (Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences) V4116. (F) Lateral and ventral views of middle Paleocene CEP *Harpyodus decorus* IVPP 5035.1. (G) Lateral and occlusal views of late Paleocene CEP *Guichilambda zhaii* IVPP V12037.2 (dentary) and V12037.3 (maxillary fragment). Firey asteroid symbols indicate the end-Cretaceous asteroid impact in the Yucatán Peninsula; thermometer symbols indicate the Paleocene-Eocene hyperthermal event. Subdivisions of the Paleocene approximately correspond to the Shanghuan, Nongshanian, and Gashatan Asian Land Mammal Ages, respectively (see supplemental text for competing age boundary scenarios).

Fig. 2. Association of paleopalynological data from the Nanxiong Basin, south China, and late Paleocene niche expansion in endemic Asian fossil mammals. (A) Proportion of environmental humidity indicator taxa from early versus late Paleocene paleobotanical localities, respectively, in the Nanxiong Basin; data based on [34,35](Data S12). (B) Boxplots of dental complexity (OPCR, orientation patch count rotated) in the Chinese endemic pantodont (CEP) data partition across the three Paleocene time intervals examined. Note the concomitant increase in CEP tooth complexity (OPCR) and increased proportion of drought-tolerant plant species in the Nanxiong Basin during the late Paleocene. (C) Principal component morphospace of all tooth data analyzed;

convex hulls delineate overall morphospace occupation during each time interval.
Eigenvectors of the four dental topographic traits are indicated in blue. Late Paleocene
shift and expansion in dental topographic morphospace is statistically significant at the p
= 0.05 level (Table 1). Pantodont silhouette by S. Traver from phylopic.org.

Fig. 3. Correlation plots of dental topographic and bite performance traits in
endemic Asian Paleocene mammals. (A) Hypothetical correlation scenarios used to
interpret stasis, directional, verses decoupled change through time in specimen data.
(B) Pairwise ranked correlation coefficient estimated using Kendall's τ between early
and middle Paleocene dental topographic and performance traits in the main dataset.
(C) Correlation between middle and late Paleocene traits in the main dataset. (D)
Correlation between early and middle Paleocene traits in the Chines endemic pantodont
(CEP) data partition. (E) Correlation between middle and late Paleocene traits in the
CEP data partition. Topography-performance correlations are marked in red boxes.
Decoupled/reversed trait correlations are marked in gray boxes.

All data	Early-Middle Paleocene	Middle-Late Paleocene
DNE	0.7753	**0.003**
OPCR	0.0902	**0.0001**
RFI	0.605	**0.0128**
Slope	0.643	0.287
PCA Disparity	0.4562	**0.0001**
Tooth size	**<0.0001**	**<0.0001**
Chinese Endemic Pantodonts		
DNE	0.9264	**0.0002**
OPCR	0.17	**<0.0001**
RFI	1	1
Slope	1	1
Chinese Arctostylopids and Anagalids		
DNE	0.847	0.135
OPCR	0.47	0.18
RFI	0.2486	**0.0056**
Slope	0.642	0.269

Table 1. Pairwise t test of dental topographic trait and disparity differences across adjacent time bins. Dental topographic trait differences are assessed across time intervals in all-data, Chinese endemic pantodont, and non-pantodont partitions. Dental trait disparity was estimated based on all principal component axes using the outputs of PCA. Tooth size variance differences were calculated from tooth area or square root of tooth area in all-data and no-outlier partitions to assess effect of outliers on statistical significance (see Data S6 for details). Bolded font indicates p values < 0.05.

METHOD DETAILS

We focused sampling on three clades: Chinese endemic Pantodonta, Chinese

Arctostylopida, and Anagalida. Additional data were collected on other clades (e.g.,

Chinese Tillodontia) opportunistically when well-preserved specimens were available

(Data S1). These three main clades together represent >50% of the species found in

Paleocene faunas across China [11]. They also individually represent the most diverse

clades across the three Asian Land Mammal Ages of the Paleocene: Shanghuan,

Nongshanian, and Gashatan. By contrast, these clades are reduced to <25% of species

diversity in Eocene assemblages in China, finally disappearing altogether in the late

Oligocene. Therefore, we take the three clades to be representative of mammal

assemblage dynamics during the Paleocene time interval in China.

Historically, Paleocene fossil mammal faunas in China have been defined based

on biostratigraphic criteria and supplemented by magnetostratigraphic correlations

where available. Here we follow the assessment of [59] in generally correlating the

Shanghuan Asian Land Mammal Age (ALMA) with the Puercan and Torrejonian North

American Land Mammal Ages (NALMAs), the Nongshanian ALMA with early to middle

Tiffanian NALMAs, and the Gashantan ALMA with late Tiffanian and Clarkforkian

NALMAs, respectively (see [11] for an alternative interpretation). Additionally, the

Shanghuan and Nongshanian boundary was interpreted by [23] to be at the top of Chron

C27N, which is dated at 60.920 Myr ago in the Geomagnetic Polarity Time Scale [60] but

indicated as closer to ~62.3 Myr ago in [59]. Similarly, the Nongshanian-Gashatan

boundary has been variably defined at 59.24 Myr ago [59] or the base of Chron C26N,

which is 57.911 Myr ago [60]. In contrast, the end of the Gashatan coincides with the

Paleocene-Eocene boundary at 56 Myr ago and has been consistently defined as such

[59]. Given these existing uncertainties, we use the terms 'early Paleocene', 'middle

Paleocene', and 'late Paleocene' to refer generally to the Shanghuan, Nongshanian,

and Gashatan ALMAs, respectively, in this study.

We analyzed 200 individual teeth from 48 specimens, representing 37 species

(Data S1-S2). The specimens represent 2 upper first premolars, 4 upper second

premolars, 15 upper third premolars, 19 upper fourth premolars, 22 upper first molars,

23 upper second molars, 16 upper third molars, 3 lower first premolars, 5 lower second

premolars, 13 lower third premolars, 17 lower fourth premolars, 20 lower first molars, 22

lower second molars, and 19 lower third molars. The assigned geologic age of the

studied specimens spans the entire Paleocene, binned into early (n = 91 teeth), middle

(n = 46 teeth), and late Paleocene (n = 63 teeth) data partitions. To maximize sample

size while minimizing disturbance to more delicate specimens, a combination of original

specimens and previously produced high-fidelity specimen casts were used in our

sampling of dental crown morphology.

<u>Tooth shape estimates using Dental Topographic Analysis</u>

We digitized dental crown morphology via microCT either using a GE v|tome|x m

300/180 kV micro-computed-tomography system (GE Measurement & Control

Solutions, Wuntsdorf, Germany), housed at the Institute of Vertebrate Paleontology and

Paleoanthropology (IVPP), Chinese Academy of Sciences (CAS), or a GE Phoenix

Nanotom M in the Functional Anatomy and Vertebrate Evolution Laboratory, University

of California, Berkeley. Projection images (1,000 to 1,500 images depending on

specimen size) were acquired with an isotropic voxel size of 10 to 40 μm, at an energy

range of 120-150 kV, current of 100-150 μA. Additionally, surface 3D scans (generated

using IVPP's Artec Space Spider 3D scanner, with a precision of 0.05 mm and

resolution of 0.1 mm) were used for larger specimens to efficiently obtain surface

morphology data. The enamel caps (the crown portion of each tooth above the enamel-

dentine junction) were extracted from the specimen models using Geomagic Wrap

v2021; all crown surface holes (generated from scanning imperfections or digitally

removed sediment) were patched, the models were remeshed with triangles that have

aspect ratios (base:height or edge-edge) of <10 and decimated to ~10,000 triangles.

Finally, we standardized the spatial orientation of all tooth crowns before exporting them

as .ply files for dental topographic analysis (DTA). Individual tooth crown (enamel cap)

models have the Z axis normal to the occlusal plane, the X axis parallel to the mesial-

distal axis, and the Y axis parallel to the labial-lingual axis.

Exported meshes were further vetted and prepared for DTA within the R

programming environment. We used functions implemented in the *molaR* R package [61]

for all of the steps described next. First, all .ply files are further cleaned using

‘molarR_Clean’ to remove floating points/vertices and any triangular faces with zero

area. Then, the batch function ‘molarR_Batch’ was used to calculate four dental

topographic metrics: DNE, OPCR, Slope, and RFI.

Dirichlet Normal Energy (DNE) is a measure of occlusal sharpness, using a

quantification of surface energy in tooth crowns relative to a gently curving or flat mesh

surface. Steep, high, and/or shearing cusps tend to produce higher DNE values in DTA,

whereas bulbous cusps tend to produce lower DNE values [62,63]. DNE has been used to

successfully distinguish folivores, omnivores, and frugivores in euarchontan (primates,

colugos, tree shrews) mammals [62,64].

Orientation Patch Count Rotated (OPCR) is a measure of tooth crown

complexity, using quantification of distinct patches on the crown face that possess

unique orientations. Teeth with larger number of cusps and crenulations tend to produce

higher OPCR values, whereas teeth with fewer cusps and simpler ridges tend to

produce lower OPCR values. This metric has been used effectively to distinguish

convergently evolved carnivore versus herbivore species across multiple mammalian

orders [65]. DeMers and Hunter [66] suggest that OPCR does not currently provide a

reliable approximation for the degree of adaptation to herbivory, and that any

interpretation of ecomorphological differences should be made within a taxon-specific

context. Given the narrow set of clades targeted in our dataset, we interpret increasing

OPCR as an indication of increased functional capability to process vegetation. A

related measure, slope, quantifies the mean tilt of the tooth surface relative to the

occlusal plane. Teeth with lower and more gently curving cusps tend to have lower

slopes, whereas teeth with sharp and/or abruptly ridged crests tend to have higher

slopes.

Relief Index (RFI) is a measure of the relative height and complexity of the tooth

crown, using the values of the 3D surface and 2D ‘footprint’ areas of the tooth crown.

RFI has been shown to be effective in distinguishing frugivores from insect and leaf

specialists in euarchontan mammals [62]. Furthermore, RFI is also able to distinguish

carnivoran species at different trophic levels and dietary breadths [63].

All tooth crown models are provided as ply files and deposited in FigShare

(10.6084/m9.figshare.28611854).

<u>Tooth performance estimates using Finite Element Analysis</u>

The common inference that dental morphology can reflect dietary adaptations

makes an underlying connection between the two traits through biomechanical

performance; the mechanical performance imbued by a particular morphological

configuration accomplishes one or more food acquisition and/or processing tasks. Such

mechanical performance links can be tested using experimental and simulation

approaches on both theoretical grounds and actual tooth shapes [67]. Here we applied a

simulation approach to estimate two general performance traits commonly interpreted

for mammalian tribosphenic teeth: puncturing/compressing and shearing/grinding [68].

The ancestral therian tribosphenic stroke, or the chewing movement underlying

mammals with unfused mandibular symphysis and tribosphenic molars, has been

reconstructed to involve significant components of (1) long-axis rotation in each

hemimandible and (2) mortar-and-pestle grinding between upper and lower cusps [69].

These movements are associated with the 'lock-and-key' occlusion of a stereotypical

mammalian tribosphenic dentition, which exhibit tall crest-like cusps in the trigonid and

lower basin-like valleys in the talonid of lower molars that articulate with their

corresponding upper teeth (Fig. S3).

We used finite element analysis (FEA) to estimate the work efficiency of different

tooth crowns subjected to compressive/puncturing and shearing/grinding forces, the two

major masticatory actions of tribosphenic teeth [68,70,71]. Adequate compressive and

puncturing forces applied through individual or combined action of cusps are needed to

overcome the fracture strength of brittle and tough food such as seeds and invertebrate

exoskeletons in order for individuals to access soft and/or more nutritious tissues within.

Sufficient shearing and grinding forces applied through surfaces or cusp edges are

necessary to further break down food boli into smaller pieces to aid in the digestive

process. The combination of these two major functions of tribosphenic teeth is thought

to be a foundational adaptation that enabled the mammalian radiation [68,69,72]. We

focused on the dental enamel portion of the tooth crown in our biting simulations

because it is the most mineralized vertebrate tissue and typically the best-preserved

component of fossil teeth. The fracture mechanics of mammalian dental enamel is

thought to be intimately related to dietary function [73], including the evolution of different

enamel thickness in different feeding ecomorphologies [28]. It is important to note that

because the dental data available for this study included not only CT scans, but also

surface scans and specimen casts, enamel thickness was not quantifiable for most

specimens in our dataset (see below for a description of enamel thickness scaling

procedure applied in the simulations).

The two masticatory scenarios tested, compressing/puncturing versus

shearing/grinding, have previously been shown to be broadly connected to tooth

topology. Taller cusps (approximated in our study by Relief Index and partially by Slope)

and more convex crowns (approximated in our study by DNE) tend to exhibit higher

strain in finite element models of hypothetical tooth shapes [74]. However, there is by no

means a consensus on the presence of a tight form-function linkage across all tooth

types and/or clades. Whereas a dental topography to food mechanical property linkage

can be detected in some extant carnivores [75,76], bats [77], and primate-like tooth models

[78], a form-function relationship between dental topography and tooth/food mechanical

properties is absent in other model systems [64,79]. Therefore, we incorporated

biomechanical simulations as an explicit test of the mechanical performance

associations implied by the DTA-flora correlations established in the dental topography

and paleoenvironmental portion of our analyses. The main question addressed by these

simulations is not necessarily whether a form-function linkage exists in the Paleocene

mammal dataset, but whether there is a consistent relationship between DTA and FEA

traits across the three time bins of the Paleocene. If such consistent linkages are

observed, it would suggest a stasis in dental topography and compressive/shearing

performance through the first 10 million years of the Cenozoic in our dataset;

alternatively, any significant changes in the topography-performance relationship would

indicate substantial evolutionary change through the same time period.

We imported the same tooth crown models generated for DTA into the Strand7

finite element software (Strand7 Pty Ltd, Australia) to create finite element models of the

individual teeth. The tooth meshes are composed of two-dimensional, three-noded

triangular elements ranging between 9,000-10,000 elements, with assigned thickness

parameter (see below) so they behave similarly to single-layer 3-D elements. A

convergence test was done on two of the specimens (IVPP V5228, V5231) and

indicated that the <10,000 three-noded triangular plate elements (tri3) returned

simulation output values with less than 10% deviation from higher resolution meshes of

the same models (Fig. S3). Therefore, we deemed the default resolution of the tooth

meshes used in DTA to be sufficient for finite element simulations. Furthermore,

because the dataset contained digital models built from both CT and surface scan data,

not all specimens have enamel thickness information. Although increased enamel

thickness has been associated with durophagy in mammals, there appears to be taxon-

specific patterns of enamel thickness to tooth size and dietary ecology, in addition to

extensive variation in intra-tooth enamel thickness distributions [80–82]. In absence of

information on enamel thickness differences among the taxa studied, we standardized

the thickness of the tooth crown models isometrically to be 10% of the total surface area

of a given tooth model. As such, the simulation results should be treated as enamel cap

shape-derived performance traits, rather than those based on a fully parameterized

enamel model incorporating localized thickness and inter-specific allometric differences.

We scaled applied forces on the teeth, simulating compressive or shearing loads,

to be a value equivalent to total model surface area (but in units of Newtons instead of

mm^2). Compressive loads were represented on each tooth by a single nodal force

vector directed towards the base of the crown along the height axis of each tooth, on

the tallest cusp or cuspid (which is typically the protocone for upper molars and

protoconid for lower molars, or the main cusp in premolars). This configuration

simulated a compressive load directed into the tallest cusp (towards the root) on a given

tooth from a food item (Fig S3), as reconstructed for the mammalian tribisphenic bite

during a crushing or puncturing movement [68,69]. Shearing loads were represented on

each tooth by two nodal force vectors, one on each of the two tallest cusps or cuspids;

the two nodal forces are of equal magnitude but opposing directions, perpendicular to

the long axis of the tooth and in the occlusal plane. This configuration portrays a

shearing motion along the short axis of the tooth (Fig. S3), and simulates tooth-to-tooth

and tooth-to-food contact during the 'mortar and pestle' grinding rotation movement

reconstructed for the ancestral mammal tribosphenic bite [69]. This force magnitude

standardization procedure effectively generates identical force to area ratios across the

models, ensuring that the magnitudes of mechanical stress placed on the model are the

same across models of different sizes; furthermore, we adjusted the strain energy

values collected for each tooth model simulation using the volume and input force ratio-

based correction equation provided in [83]. Given our objective to assess the association

between relative performance traits and DTA metrics, with the latter being size-free

variables, only the relative magnitudes of bite performance, rather than absolute values,

were collected from the performance analyses.

After each tooth crown model was defined using the criteria outlined above,

homogenous material properties were assigned. All models were defined as plate

models (with thickness scaled to surface area) with a Young's (elastic) modulus of 80

GPa [84] and Poisson Ratio of 0.3. Next, each tooth was constrained from translation and

rotation using four nodal constraints distributed in the four corners of each tooth,

respectively (Fig. S3).

All bite scenarios were solved using the linear static solver implemented in

Strand7. A total number of 400 analyses were performed (one compressive bite and one

shearing bite simulation for each of the 200 tooth specimens in the dataset). We

extracted two traits from the models: compressive bite strain energy and shear bite

strain energy. Strain energy is defined as the area under a stress-strain curve of an

object under load; operationally it measures the amount of work done by the load to

deform the object (analogous to experimental work-to-fracture measurements)[85]. An

object that is more resistant to deformation would have lower strain energy than an

object that easily deforms under load. The use of work-to-fracture measures to assess

tooth performance is consistent with the understanding of mammalian tooth enamel as

a fracture-prone, yet fracture-resistant, biological tissue [86]. In such a framework, fracture

resistance is expected be directly related to the amount of force an individual is able to

exert through the tooth crown during mastication, and by extension the hardest or

toughest food item that can be processed without catastrophic damage to the tooth

crown itself. Fracture resistance (as approximated by strain energy in this study) is also

likely to be a stronger target of selection in mammals compared to those of other

toothed vertebrates, given the former's diphyodonty (having only two sets of tooth

instead of continuous dental replacement) and thus necessity to prolong usable tooth

lifespan[80]. Therefore, we used strain energy values under compressive and shear bite

simulation scenarios as a proxy for the effectiveness of each tooth crown model at

resisting deformation from the respective biting forces. A tooth that is more resistant

(i.e., have lower strain energy values) from compressive forces would be able to more

effectively crush/puncture harder food items, and a tooth more resistant from shear

forces would be able to cut/grind tougher food items. We modify Irschick et al.'s[87]

definition of whole-organism performance and define bite performance in the context of

this study as the capacity of individual tooth models to resist simulated compressive and

shearing forces, which represent ecologically relevant factors that influence masticatory

efficacy.

All finite element models are provided as NASTRAN/text files and deposited in

FigShare (10.6084/m9.figshare.28611854).

<u>Sensitivity validation of original versus cast specimen models</u>

The tooth dataset developed in this study contains a mixture of CT scans of

original and casts of fossil specimens. In order to assess the potential discrepancies in

DTA and FEA trait values collected from cast versus original specimen derived models,

either due to phenotypic details not captured in cast specimens or deterioration of epoxy

or resin-based casts over time, we randomly sampled two specimens for which both

original and cast based models were analyzed. Results from DTA and FEA of the CT-

derived models of lower first molars from IVPP V5228 (*Altilambda pactus*) and V5231

(*A. tenuis*) were compared to those conducted on digital models built from CT scans of

casts of those specimens.

We used the 'n-point' and 'global' alignment functions in Geomagic Wrap to first

align the models using arbitrary homologous landmarks visually identified on both tooth

models and then the automatic global alignment function to maximize overlap between

the two models in 3D coordinate space. The 'deviation' function in Geomagic Wrap was

then used to calculate summary statistics for linear deviations orthogonal to the surface

of the original model. The V5228 Lm1 model showed a maximum deviation of 0.58 mm

(5.8% of crown height) and average deviation of 0.04-0.09 mm (<1% of crown height),

with a standard deviation of 0.11 mm (1% of crown height). The V5231 Lm1 model

showed a maximum deviation of 0.38 mm (6% of crown height) and average of 0.03-

0.04 mm (<1% of crown height), with a standard deviation of 0.06 mm (1% of crown

height)(Fig. S3).

<u>Quantification of uncertainty ranges for bootstrap sampling</u>

We then subjected the cast-based models to DTA and FEA. In both cases, the

DTA values between cast and specimen models are the most different for DNE (30-40%

difference), followed by OPCR (18-32% difference), then RFI (2-13% difference), and

finally Slope (1-5% difference) (Data S4). Also in both cases, adjusted compressive SE

values differed by 14-41% and adjusted shear values differed by 20% (Data S3). Based

on these specimen-cast validation tests, we set a conservative +/- 40% uncertainty

range for all DTA and FEA values obtained from all cast-based tooth models. DTA and

FEA trait mean and variance estimates used for downstream statistical analyses were

then calculated using a bootstrap sampling scheme where 1,000 replicates of the DTA

and FEA datasets were passed through the statistical tests (see Quantification and

statistical analysis section, below). Overall, the early and middle Paleocene time bins

contain around 50% cast data, whereas the late Paleocene time bin contains 66% cast

data. If data uncertainty from original versus specimen models present a major signal,

we would expect the late Paleocene time interval to always show higher variance given

an abundance of cast data in that time bin. We did not observe any consistent trends of

high variance in the late Paleocene in the 1,000 bootstrap samples, suggesting that the

results are not significantly biased by data quality differences between cast and original

specimen derived models (Fig. S4).

<u>Time bin duration correction and tooth position ratios</u>

According to Ni et al. [6], the Shanghuan Asian Land Mammal Age (ALMA)

faunas in Guangdong and Anhui range from 66-61.6 Ma (or 4.4 Myr in duration), the

Nongshanian ALMA faunas in Guangdong, Jiangxi, and Anhui range from 61.6-59.2 Ma

(or 2.4 Myr in duration), and the Gashatan ALMA faunas in Anhui rangesfrom 59.2 to 56

Ma (or 3.2 Myr in duration). Given the different durations represented in each of the

three time bins used in our analyses, we corrected all variance estimates by dividing the

variance calculated in each of the bootstrapped samples by the time duration of the
respective time bin. In this manner, the variance values reported in the study represent
per-million-year values.

We used pie chart analysis to verify that the proportions of tooth positions in each
time bin are not substantially different from each other (Fig. S1). However, we caution
that time bin comparisons of individual tooth positions are unlikely to be statistically
robust because of small sample sizes. We only use aggregates of dental positions (all
teeth, molars, or premolars) in our data partition analyses and interpretation.

QUANTIFICATION AND STATISTICAL ANALYSIS

We used the following features of the dental topography and performance data
as the basis for assessing faunal assemblage trait shifts through the Paleocene: trait
mean, variance, trait-to-trait correlation, and partition-to-partition correlation. We used
ANOVA (Analysis of Variance) and pairwise t test to compare trait means, F test to
compare trait variance, linear regression analysis and Kendall's τ to compare trait-to-
trait correlation, and two-block partial least squares (2B-PLS) analysis to compare
partition-to-partition correlation. There are a total of six traits forming two main data
partitions: topographic partition (DNE, OPCR, Slope, RFI) and performance partition
(Compressive SE, and Shear SE). To assess the sensitivity of the results to subsets of
the data, all of the trait comparisons were done iteratively using the total dataset, by
time bin (early, middle, late Paleocene), taxon (Chinese endemic pantodonts versus
non-pantodonts), and/or by dental position (all teeth, molar teeth, premolar teeth). All
statistical tests were performed on 1,000 bootstrap resampled datasets that were
parameterized using results of sensitivity tests on original versus cast specimen models,
finite element mesh convergence tests, and time bin duration comparisons (see
previous section). In addition, we quantified tooth size using two measures: total 2D
surface area and square root of surface area. The R script for all statistical analyses
and plots are included in Data S9.

<u>Statistical resampling and tests</u>

We used bootstrap resampled variance of dental topographic metrics and dental

performance variables as a measure of tooth form-function disparity. 1,000 replicates of

the 200-specimen DTA-FEA trait datasets were generated by sampling each DTA and

FEA trait value from a uniform distribution defined using a +/-40% range of the original

model-derived values. The bootstrap resampling was done with replacement, so it was

possible for a given DTA/FEA trait value in a sampled tooth model to be repeated in a

different replicate, but very unlikely for entire datasets to share similar values with

another replicate. We used this resampling scheme to account for the uncertainty in trait

estimates introduced by the totality of modeling, specimen preservation, cast versus

original, and other potential sources of uncertainty in trait values. For each replicate,

variance was calculated for each metric, time interval, for Chinese endemic pantodont

(CEP) versus non-pantodont, and molar versus premolar data partitions. To assess

shifts through time, statistical differences between the variance of each dental

topographic metric sample in adjacent time intervals (A, early Paleocene; B, middle

Paleocene; C, late Paleocene) were evaluated using F tests. A null hypothesis of a

variance ratio of 1 between pairwise comparisons was tested using the var.test()

function in R [88]. The outputs of the variance tests on a give bootstrap sample were then

compiled for all 1,000 replicates, and the overall variance mean and variance test output

used as the statistical basis for detecting significant differences between time, taxon,

and tooth data partitions. Unless indicated otherwise, all tests described below were

also done using the 1,000 bootstrap samples.

We evaluated differences in central tendency (mean value) of dental topographic

metrics and dental performance variables through time using analysis of variance

(ANOVA) and pairwise t tests. Each dental topographic metric was tested against the

three time intervals defined above, using the aov() function in R. For statistically

significant (at the $p < 0.05$ level) results, we additionally evaluated the pairwise intervals

that contribute significant differences in mean dental topographic values. We assessed

pairwise differences using pairwise t tests implemented with a Holm correction for

multiple comparisons. The t test was conducted using pairwise.t.test() in R.

We constructed a tooth morphospace using all four dental topographic metrics

analyzed by principal components (PC) analysis. The first and second PC axes were

chosen to visualize the two-dimensional morphospace. Additionally, we quantified the

degree of morphological disparity and statistical differences in disparity between

adjacent time interval data partitions. All PC scores generated from the PCA were

included in the disparity analysis. The prcomp() function in R was used for PCA, and the

morphol.disparity() function implemented in the Geomorph R package [89] was used for

morphological disparity significance tests. Input data for the PCA were from the original

dataset values, not from the bootstrap samples.

Linear regression analysis between individual dental topographic metric (DNE,

OPCR, Slope, RFI) and dental performance metric (compressive bite strain energy,

shear bite strain energy) was performed to quantify the correlation between dental form

and function. Adjusted R^2 and p values generated from the lm() function in R were used

to evaluate the goodness of fit and statistical significance of the form-function

relationships. The distribution of R^2 values and their corresponding p values are

reported in Fig. S6.

In addition to pairwise form-function linear regression analyses, we also

evaluated the degree of correlation between the DTA and FEA data blocks as a means

to measure integration between dental topography and deformation resistance. We

used two-block partial least squares analysis[90] coupled with bootstrap resampling to

generate distributions of correlation coefficients (*r-pls*) for each of the three time bins,

and then tested for significant differences in the magnitude of DTA-FEA correlation

between adjacent time bins using Welch's two-sample t tests on the correlation

coefficient distributions. DTA-FEA correlation differences at $p < 0.05$ are interpreted as a

significant shift in the degree of integration of the two traits from one time bin to the next

time bin.

All R scripts used in the analyses are included as supplementary files (Data S8,

S9).

A tooth traits

70   60   50   Million years ago

Cret.   Paleocene   Eo.

Temperature (°C)
12°
8°
4°

OPCR
300
200
100
0

orientation

DNE
1500
1000
500
0

convex
concave

RFI
0.8
0.6
0.4

tall
flat

Slope

steep
gentle

B Log10 (sqrt area)
1.2
0.9
0.6
0.3
0

E.Paleo   M. Paleo   L.Paleo

C Compressive bite variance
1.5e-08
1.0e-08
0.5e-08
0

E.Paleo   M. Paleo   L.Paleo

D Shear bite variance
5e-07
4e-07
3e-07
2e-07
1e-07
0

E.Paleo   M. Paleo   L.Paleo

Upper molars   All premolars
Lower molars   All molars
All teeth

IVPP V4116
IVPP V12037.2

E

F

G