Exploring the genetic basis of cluster architecture-related traits in

grapevine through a Genome-wide Association Study

Geovani Luciano de Oliveira[1], Felipe Roberto Francisco[2], Yohans Alves de Moura[1], Guilherme

Francio Niederauer[1], Roberto Fritsche-Neto[3], Anete Pereira de Souza[1,4], Mara Fernandes Moura

Furlan[2*]

[1] Molecular Biology and Genetic Engineering Center (CBMEG), Universidade Estadual de Campinas

(UNICAMP), Campinas, Brazil

[2] Advanced Fruit Research Center, Instituto Agronômico (IAC), Jundiaí, SP, Brazil

[3] Horticulture Sciences, College of Agriculture and Life Sciences, North Carolina State University,

Raleigh, NC, 27695-7609, Campus Box 7609, USA

[4] Department of Plant Biology, Biology Institute, Universidade Estadual de Campinas (UNICAMP),

Campinas, Brazil

* Corresponding author

E-mail: mara.moura@sp.gov.br (MFMF)

Abstract

Berry and cluster size are pivotal determinants of grapevine productivity and consumer preferences and remain major targets in grapevine breeding. However, given their complexity as quantitative traits under polygenic control, a deeper understanding of their genetic determinants is needed. The gene pool of the Brazilian grapevine has made a significant contribution to enhancing grapevine performance in tropical and subtropical regions. In this study, we conducted a genome-wide association study (GWAS) using a diverse panel of 288 *Vitis* spp. accessions from the Instituto Agronômico Germplasm Bank, Brazil. This panel was phenotyped for six critical cluster architecture traits over 12 years and genotyped using the Vitis18k SNP array. Using two different algorithms, the GWAS identified 56 significant SNPs distributed across 17 chromosomes, validating previously identified QTLs and revealing novel associations. Four closely spaced markers on Chr1 suggest the presence of a QTL influencing five traits simultaneously. A strong association signal, explaining approximately 29–35% of the phenotypic variance (PVE), indicated a major QTL for berry length (BL) and width (BWi) on Chr14. Additionally, major-effect SNP loci were identified for cluster weight (CW) on Chr1, cluster length (CL) on Chr7 and 14, cluster width (CWi) on Chr6 and 18, and berry weight (BW) on Chr4, with PVE values ranging from 18–27%. Furthermore, 80 genes associated with berry traits and 52 genes associated with cluster traits were identified as putative candidate genes in the genomic regions associated with significant SNPs. These candidate genes are involved in the regulation of growth and development, hormone regulation, protein synthesis, stress response, and other physiological processes essential for cell health and functionality. Our results provide valuable insights into the genetic determinants of grape berry size and cluster architecture, offering critical data to support future functional studies and enhance the efficiency of related breeding programs.

Keywords: quantitative trait loci, subtropical viticulture, berry size, GWAS, genetic resources, grapevine breeding

Introduction

Grapevine (*Vitis* spp.) is a key fruit crop worldwide, with berries consumed fresh or used in the production of wine, juice, jellies, and other food products [1,2]. Each grape product demands specific berry traits, such as color, composition, taste, and size, to meet diverse market and industry requirements [3]. Over centuries of domestication and breeding, grapes have accumulated significant morphological and physiological diversity, resulting in thousands of varieties tailored for specific purposes. The variations in these traits hold considerable potential as valuable resources for grape breeding [4,5].

Grapevine, as a woody perennial fruit crop, presents challenges associated with its substantial size and prolonged juvenile phase, which make plant cultivation efforts and evaluation in breeding programs time-consuming and costly [6]. This issue is particularly pertinent because the assessment of fruit characteristics can only commence once the plant reaches maturity, typically in the fourth or fifth year of its life cycle [7]. To address this, early genotype selection technologies, including the use of genomic tools for marker-assisted selection (MAS), have been suggested to accelerate and improve the plant breeding process. These advances enable breeders to identify and select individuals with desired genotypes at an early stage, facilitating more efficient and targeted breeding programs. This promises a significant reduction in the breeding cycle by up to a decade, along with notable cost savings ranging from 16–34% [8,9]. Identifying genotype□phenotype associations is crucial for plant breeding. However, several fundamental issues, especially regarding complex traits, still need to be resolved before MAS can fully realize its potential in practical breeding programs [10]. Fruit yield and quality are the most commonly measured characteristics in grapevines but are poorly understood.

These traits are quantitatively extremely complex due to the number of segregating loci that control

all the components involved in productivity, as well as the influence of nongenetic factors [11–13].

Berry size is a crucial agronomic trait that strongly influences both quality and yield in

grapes. This feature is essential for table grapes, as it is critical for consumer preferences and

acceptance and directly influences market value [14,15]. Berry size also plays a key role in

determining juice composition and affects the overall quality of wine [16–19]. Cluster size is another

essential factor influencing fruit quality and production. Together with berry size, cluster size is vital

for defining cluster architecture and significantly affects disease pressure, sun exposure, and uniform

ripening [20,21]. Given the importance of these factors for breeding, numerous studies have utilized

quantitative trait locus (QTL) mapping in biparental progenies to elucidate the genetic architecture

underlying key berry size and cluster architecture traits in grapes and identify QTLs on various

chromosomes [15,22–30]. Notably, most major QTLs related to berry size have been found on

Chr18, closely linked to the well-characterized SDI (seed development inhibitor) locus [31–34].

Linkage mapping has significantly advanced our understanding of important berry and

cluster-related traits in grapes. However, the results have shown variability across different genotypes

and environments, underscoring the complexity of the genetic control of these traits [35,36].

Moreover, many of the biparental progenies analyzed in these studies were derived exclusively from

V. vinifera genotypes, which may have produced inconclusive results and an underestimation of the

genetic complexity of the crop. This approach also has inherent limitations, including restricted

allelic diversity in parental lines, low recombination frequency in offspring, underestimation of

polygenic contributions for predictive modeling, and reduced applicability across broader genetic

backgrounds [37].

To overcome these constraints, genome-wide association studies (GWASs) offer an

alternative approach in which genetic architecture and gene interactions influencing traits are

investigated without the need for a specifically designed mapping population [38–41]. The GWAS

methodology leverages the extensive history of recombination within diverse germplasms to identify

small haplotype blocks associated with phenotypes of interest across species-scale diversity [42].

These recombination events diminish the extent of linkage disequilibrium (LD) and ensure more

precise genome exploration, provided that marker density is sufficient [36,43]. In recent years,

several GWAS investigations have focused on traits related to berry and cluster traits in grapes

[37,39,40,44–46]. However, the genetic foundations of many important traits, such as berry width

and length, as well as cluster weight, width, and length, remain underexplored. The extensive

diversity within cultivated grapevines requires further investigation. Intra- and interspecific crosses

conducted throughout breeding cycles to introduce novelty, resistance, adaptability, and hybrid vigor

have resulted in hybrids with heterogeneous genetic backgrounds [38,47]. Analyzing a more diverse

germplasm in additional association studies may uncover new sources of variation for previously

characterized traits [43]. Moreover, integrating genomic technologies with traditional breeding

methods could significantly increase the ability to address current and future environmental

challenges. Under these circumstances, the conservation of germplasms across plant species and their

utilization, leveraging complementary approaches such as genetic and genomic resources, are vital

for achieving rapid advances in productivity [48].

One of the main objectives of grapevine breeding programs in Brazil is to develop cultivars

that combine adaptations to tropical and subtropical regions with fruit quality, productivity, and

resistance to biotic stresses [49]. To support this goal, the Instituto Agronômico (IAC) has, over

several years, established a diverse germplasm collection that includes key *V. vinifera* cultivars,

complex interspecific hybrids from both national and international sources, along with non-*Vinifera*

species [50], and the phenotypic performance of these accessions has been characterized under

Brazil's subtropical conditions. Leveraging the broad genetic base of this diverse panel of grapevine

resources, we carried out a GWAS analysis on a set of 288 preselected genotypes to investigate six

yield-associated traits to identify novel genomic regions (QTLs) associated with these traits and

confirm previously detected QTLs, as well as locate candidate genes within the identified genomic

regions. The results of this study provide valuable resources for grapevine breeding by offering

insights into the molecular mechanisms potentially involved in cluster architecture, establishing a

foundation for subsequent validation experiments, the functional characterization of candidate genes,

and the implementation of targeted gene editing strategies.

Materials and methods

Plant material and DNA extraction

The tested population consisted of 288 accessions of *Vitis* spp. from the IAC Grapevine

Germplasm Bank, located in Jundiaí, São Paulo (SP), Brazil (23°06'53"S, 46°57'38"W, elevation

745 m). In accordance with the Köppen classification [51], the climate is categorized as Cfa,

characterized by subtropical conditions featuring dry winters and hot summers, and the soil in the

area is classified as Cambisol Dystrophic Red [52]. The germplasm is established within the same

region, with the accessions distributed across three physically adjacent experimental fields (S1 Fig).

The accessions were previously characterized for their trueness-to-type and molecular

diversity [50,53], and only distinct genotypes were selected. The material consisted of *Vitis vinifera*

cultivars, interspecific hybrids, and non-*Vinifera Vitis* species (see S1 Table for detailed data on the

accessions). Each accession consisted of three or four clonally propagated plants grafted onto 'IAC

766 Campinas' rootstock and trained on a vertical shoot position (VSP), with spacings of 2.5 m

between rows and 1 m between vines. Production pruning was carried out annually in August,

leaving one or two buds per branch. After pruning, 2.5% hydrogen cyanamide was applied to induce

and stimulate a more uniform budburst. Additional culturing and phytosanitary management

practices were also performed according to the standard practices for local growers in São Paulo

(SP), Brazil. For sampling purposes, young leaves were collected from a single plant from each

accession.

Total genomic DNA was extracted from the leaf tissue with a DNeasy® Plant Mini Kit

(QIAGEN, Hilden, Germany) according to the manufacturer's instructions. DNA quality was

assessed via both agarose gel electrophoresis and a NanoDrop 8000 spectrophotometer (Thermo

Fisher Scientific, Wilmington, DE, United States). The DNA concentration was determined with a

Qubit 3.0 fluorometer (Thermo Fisher Scientific, Waltham, MA, United States).

Phenotypic data

Phenotypic characterization of clusters and berries was conducted over twelve production

cycles from 2010–2021. Harvests were carried out between December and February of each cycle,

and six traits related to the berries and clusters were evaluated. In general, three plants per genotype

were maintained. For each genotype, six clusters were collected, and cluster weight (CW, in grams),

cluster length (CL, in centimeters), and cluster width (CWi, in centimeters) were recorded. For berry-

related traits, measurements were taken from ten berries collected from the upper, middle, and lower

parts of the clusters in a 3:4:3 ratio, providing data on berry weight (BW, in grams), berry length

(BL, in centimeters), and berry width (BWi, in centimeters). Weight was measured using an

analytical balance, and length and width were determined using a metric scale.

Outliers from the phenotypic data were removed using the "outlierTest" function in the *car*

package in R [54], employing the Bonferroni outlier test with a statistical threshold of $p < 0.05$. The

best linear unbiased estimators (BLUEs) of grape genotype effects were estimated for each of the

phenotypic fruit yield traits. Genotypic BLUE values were obtained through spatial adjustment based

on rows and columns, using the 'mmec' function from the R package *sommer* [55]. The mixed model

employed was as follows:

$$Y_{ij} = X_g\beta_g + Z_{gxe}(G \ x \ E) + Z_r R + Z_c C + e$$

where Y_{ij} represents the phenotype of the *i*-th genotype in the *j*-th field environment (A, B or

C). The fixed effect of the genotypes is denoted by β_g, with the corresponding design matrix X_g. Z_{gxe}

corresponds to the random effect of the genotype × environment interaction, whereas Z_r and Z_c

represent the random row and column effects, respectively, within each environment. The residual

error term *e* is assumed to follow a normal distribution, $e \sim N (0, \sigma^2_e)$.

Variance components were derived from the output of the *sommer* package, and broad-sense

heritability (H²) was calculated as the ratio of total genetic variance to total phenotypic variance via a

modified version of the mixed model described earlier, with genotype effects treated as random. This

was accomplished using the formula H² = VG/(VG + VE), where VG represents the estimated

genetic variance and VE represents the estimated environmental variance. The estimated BLUE

values were used as phenotype values in the GWAS. Pearson correlations between traits were

estimated from adjusted BLUEs, and the correlation matrix results were plotted in R using *ggplot2*

and the *ggpairs* function in Ggally.

185   SNP genotyping

The 288 accessions were genotyped using the commercial Vitis18kSNP array (Illumina Inc.,

San Diego, CA, United States) at Neogen (Pindamonhangaba, Brazil), following the Infinium HD

Assay Ultra Protocol (Illumina Inc., San Diego, CA, USA). The raw SNP data were visualized and

clustered in GenomeStudio 2.0 software (Illumina Inc., San Diego, CA, USA) with a GenCall

threshold of 0.15. The SNP scoring tool ASSIsT 1.02 [56] was subsequently utilized, employing

default thresholds tailored for germplasm material, to filter the dataset. SNPs classified as

"Monomorphic", "Failed", or "NullAllele-Failed" were removed. SNPs that exceeded the previous

step with a minor allele frequency (MAF) higher than 0.05 were subsequently filtered out using

snpReady software [57], and the remaining SNPs were used as the high-quality subset for further

analysis. Imputation was performed via the k-nearest neighbor imputation (kNNi) algorithm [58].

Genome-wide association study

GWAS was performed for each trait independently via the R statistical genetics package

GAPIT version 3 [59]. The first three principal components from a principal component analysis

(PCA) were included as covariates in the mixed linear model to account for population structure

effects [60]. In contrast, the kinship matrix was used to reduce false positives in the GWAS results.

Individual relationships were estimated using the VanRaden method, as implemented in GAPIT3

software [61]. Two GWAS methods were employed, namely, fixed and random model circulating

probability unification (FarmCPU) [62] and Bayesian information and linkage disequilibrium

iteratively nested keyway (BLINK) [63]. Significant marker–trait associations were defined as those

with Benjamini–Hochberg (B&H) false discovery rate (FDR)-adjusted p values < 0.05, as calculated

by GAPIT3. The online tool MapGene2Chrom [64] was employed to generate a map of significant

SNPs.

Gene annotation

To identify candidate genes associated with significant SNPs, the grapevine reference genome

PN40024.v4 [65] was accessed using the JBrowse [66] feature of Grapedia

(https://grapedia.org/genomes). Considering the average LD window in *Vitis* [38,40,45,67,68],

candidate genes were identified within a 15 kb window upstream and downstream of each significant

SNP region in the genome browser. We utilized the grape gene reference catalog [69] and the

UniProt database [70] to obtain functional annotations for each gene identifier. Further functional

annotations for each gene identifier were obtained via Blast2GO (https://www.blast2go.com/) to

provide additional context.

Results

224
Phenotype analysis

Phenotypic data on the cluster and berry traits studied were collected over twelve consecutive

fruiting seasons. Substantial phenotypic variation was observed among the evaluated grapevine

genotypes (S2 Fig). The trait values were adjusted according to the mixed model from which the

BLUEs were extracted for further analysis (S2 Table). The variance components estimated for each

trait and broad-sense heritability (H^2) are presented in S3 Table. Broad-sense heritability (H^2) was

consistently high across all berry traits, with closely aligned values. The highest heritability was

observed for berry weight (BW, 0.95), followed by berry length (BL, 0.92) and berry width (BWi,

0.90). With respect to cluster traits, the H^2 values varied, with high heritability for cluster weight

(CW, 0.77), relatively high heritability for cluster length (CL, 0.66), and moderate heritability for

cluster width (CWi, 0.50).

Overall, all the correlation coefficients obtained were positive and significant (S3 Fig). BW

was strongly and equally correlated with BL and BWi (r > 0.95). The correlations between cluster

traits ranged from moderate to high, with the strongest correlation observed between CW and CWi (r

= 0.80) and the weakest correlation observed between CL and CWi (r = 0.61). Lower correlations

were detected between traits for the different organs (berry and cluster), with the highest correlation

occurring between CWi and BL (r = 0.56) and the lowest occurring between BW and CL (r = 0.17).

Genotyping

246   The grapevine diversity panel was genotyped using the Vitis18K SNP chip, and a total of

18,071 SNPs were scored in GenomeStudio. No poor-quality samples (call rate < 0.8) were detected,

and all the samples were retained. After quality filtering, ASSIsT classified 1,437 SNPs (8%) as

monomorphic, 1,520 SNPs (8.4%) as failed, 2,105 SNPs (11.6%) as shifted homozygotes, and 745

SNPs (4.1%) as failed null alleles. All these groups were subsequently excluded from the

downstream analysis. Additionally, the snpReady software removed 708 (4%) SNPs with a minor

allele frequency (MAF) of less than 0.05 and 441 (2.4%) SNPs with a call rate of less than 0.85.

Ultimately, a subset of 11,115 high-quality SNPs was obtained, with an imputation rate of 1.67% (S4

Table). This filtered SNP dataset was used for PCA to assess the clustering of genetic variation in the

grapevine accessions (Fig 1A). PC1 explained 12.38% of the variation in the genotypic data, whereas

PC2 and PC3 explained 4.86% and 4.47% of the variation, respectively (Fig 1B). Although PCA

revealed a low degree of population stratification, a subtle species distribution pattern among *V.*

vinifera genotypes could be deduced.

Fig 1. **Principal component analysis (PCA) of grapevine genotypes.** (A) Scatter plot with the

genomic data represented in two principal components (PC1 and PC2). Each point represents a

genotype from the diversity panel. (B) Genetic variation explained by the first ten PCs.

Genome-wide association study

GWAS was conducted using the BLINK and FarmCPU statistical models, which identified 41 and 32 SNP loci, respectively, with significant associations with the phenotypic traits of interest. Among the significantly associated SNP loci, 15 were commonly detected for the same trait by both models. Furthermore, six of the SNP loci were associated with two different traits and two with three different traits simultaneously, totaling 56 different SNP loci with associations in the study (S5 Table, Fig 2). The Manhattan plots summarize the association results obtained from the two statistical models for the studied traits (Fig 3). The QQ plots generated from the GWAS models with significant associations exhibited a pronounced deviation from the expected P value distribution in the tail area, indicating that the population structure and familial relatedness were effectively controlled by the selected covariates, leading to a reduction in false positive associations (Fig 4). The updated positions of the SNPs associated with the traits, mapped to the reference genome of the grapevine PN40024.v4, enabled the identification of the genes harboring these SNPs. Among the identified SNPs, 38 were located within a gene, and 16 were situated within 15 kb of a candidate gene. No gene was found within 90 kb of the SNP chr18_random_1146283_G_T (associated with BW), and the nearest gene to the SNP ae_C_T_chr14_16833185 (associated with BL and BWi) was 51.44 kb away. A total of 132 candidate genes were identified: 112 genes associated with a single trait, 14 genes associated with two traits, and six genes associated with three traits (S6 Table). When only markers with a phenotypic variance explained (PVE) greater than 5% were considered, 29 SNPs met this criterion in at least one of the tested models. Within the 15 kb flanking regions of these markers, 33 candidate genes were associated with berry-related traits, and 30 were associated with cluster-related traits (Table 1).

Fig 2. Distribution of SNPs associated with the traits according to the BLINK and FarmCPU models. The horizontal bars represent the total number of SNPs per trait, and the vertical bars

indicate SNPs that are either unique or shared among traits. Colors in the vertical bars distinguish

SNPs identified exclusively by one model (BLINK or FarmCPU) and those detected by both models

simultaneously. (A) SNPs associated with berry traits: BW, berry weight; BL, berry length; BWi,

berry width. (B) SNPs associated with cluster traits: CW, cluster weight; CL, cluster length; CWi,

cluster width.

Fig 3. Manhattan plots indicating single-nucleotide polymorphisms (SNPs) associated with the

traits targeted in this study, as predicted by the BLINK and FarmCPU models. The vertical

lines indicate SNPs commonly detected by both models. (A) BW, berry weight; (B) CW, cluster

weight; (C) BL, berry length; (D) CL, cluster length; (E) BWi, berry width; (F) CWi, cluster width.

Fig 4. Comparison of the QQ plots resulting from the different GWAS statistical models for

eight traits in the full grapevine diversity panel. The red diagonal lines indicate the null

hypothesis, where the observed and expected P values would be situated if there were no

associations. The shaded area indicates a 95% confidence interval; BW, berry weight; BL, berry

length; BWi, berry width; CW, cluster weight; CL, cluster length; CWi, cluster width.

Table 1. Summary of significant markers associated with berry and cluster traits identified through genome-wide association analysis (GWAS), including only those whose PVE was greater than 5% and the corresponding candidate genes located within a 15 kb flanking region of each significant marker.

Trait	Associated marker	Chr	Position	P .value	PVE (%)	Model	Gene	Gene Annotation
BL	chr1_5754629_C_T	1	5737661	1.19E-08	6.36	BLINK	Vitvi01g00511	Heat stress transcription factor (HSF) A-8
							Vitvi01g00512	Vacuolar iron transporter 1 (VIT1)
							Vitvi01g00513	Vacuolar iron transporter 1 (VIT1)
	lb_G_A_chr6_1714702[a]	6	1703187	2.44E-07–5.91E-10	0.58–11.02	FarmCPU, BLINK	Vitvi06g00141	Glycosyl hydrolase 1 family (GH1)
							Vitvi06g00143	Glycosyl hydrolase 1 family (GH1)
							Vitvi06g01615	Glycosyl hydrolase 1 family (GH1)
	chr8_21881697_G_T	8	21872602	5.02E-07	7.49	BLINK	Vitvi08g01906	Peptidase M16 family
							Vitvi08g01910	Peptidase M16 family
	chr11_1513738_G_T	11	1513466	1.48E-06	9.25	BLINK	Vitvi11g00146	NB-LRR family
							Vitvi11g00147	Acetyltransferase component of pyruvate dehydrogenase complex (DLAT/E2)
							Vitvi11g00148	Lycopene epsilon cyclase
	ae_C_T_chr14_16833185[a]	14	16527961	1.59E-09–4.33E-08	30.49–35.22	FarmCPU, BLINK	No hit	-
	chr19_23684078_C_T	19	24046863	1.94E-06–2.84E-06	0.19–10.54	FarmCPU, BLINK	Vitvi19g01761	General transcription factor 3C polypeptide
BWi	ae_A_G_chr1_12009877	1	12313223	1.29E-07	6.47	FarmCPU	Vitvi01g01010	TATA box-binding protein-associated factor RNA polymerase I subunit B
	chr2_3892599_A_G	2	3690239	4.09E-08	6.64	BLINK	Vitvi02g00402	Lysophospholipid acyltransferase 1
							Vitvi02g00403	VvMATE50 - MATE efflux family protein
							Vitvi02g00404	Uncharacterized protein
	be_T_G_chr3_12208763	3	11839072	3.23E-06	5.38	BLINK	Vitvi15g01767	Auxin response factor ARF6
							Vitvi15g04149	Protein kinase domain-containing protein
	chr5_15221260_C_T[c]	5	15456289	2.71E-09	5.42	BLINK	Vitvi05g01115	Myb-like protein X

	lb_G_A_chr6_1714702[a]	6	1703187	2.91E-08–1.12E-08	0.11–11.70	FarmCPU, BLINK	Described in BL	-
	chr6_14784507_A_G	6	14712992	3.59E-06–7.73E-11	1.24–12.66	FarmCPU, BLINK	Vitvi06g01132	Pseudouridine-5'-phosphate glycosidase
							Vitvi06g01133	GRAS/Scarecrow-like protein 3 (SCL3)
	chr11_6170462_C_T	11	6176021	1.11E-09	5.55	BLINK	Vitvi11g00570	Prolamin-like domain-containing protein
							Vitvi11g00571	Triphosphate tunnel metalloenzyme
	ae_C_T_chr14_16833185[b]	14	16527961	7.71E-09–2.27E-11	33.04–29.89	FarmCPU, BLINK	No hit	-
BW	chr4_2974988_A_C	4	2969932	2.78E-07	6.61	FarmCPU	Vitvi04g00314	Glycosyltransferases
							Vitvi04g00316	ER membrane protein complex subunit 1
							Vitvi04g00317	U-box domain-containing protein 26
	be_T_C_chr4_6544365	4	6502499	4.01E-11	19.23	BLINK	Vitvi04g00594	Chorismate mutase (CM2)
							Vitvi04g00595	Kinetochore protein NUF2
							Vitvi04g01927	DNA-directed RNA polymerase II subunit B
	chr5_15221260_C_T[c]	5	15456289	2.24E-12	7.50	BLINK	Described in BWi	-
	lb_G_A_chr6_1714702[a]	6	1703187	8.67E-10	14.23	BLINK	Described in BL	-
	chr11_5942540_A_G	11	5948097	1.70E-07	7.77	BLINK	Vitvi11g00552	Ubiquitin carboxyl-terminal hydrolase
							Vitvi11g00554	ARP2/3 complex 21 kDa subunit
	chr19_23949121_G_T	19	24311977	2.75E-10	6.26	BLINK	Vitvi19g01777	Uncharacterized protein
							Vitvi19g01778	NAC domain-containing protein
CL	chr3_16593567_A_G	3	18000659	4.42E-08–1.48E-07	1.58–7.55	FarmCPU, BLINK	Vitvi03g01151	Peptidyl-prolyl cis-trans isomerase (FKBP12)
	chrUn_7297439_A_G	7	1035840	5.30E-07	24.29	BLINK	Vitvi07g02106	Uncharacterized protein
							Vitvi07g02107	Ankyrin repeat-containing protein
							Vitvi07g02108	Uncharacterized protein
	chr8_18600161_C_T	8	18604552	2.75E-06–1.65E-07	0.61–5.86	FarmCPU, BLINK	Vitvi08g01585	Thaumatin-like protein 1
							Vitvi08g01586	Glutamine synthetase
							Vitvi08g02292	Uncharacterized protein
	lb_T_C_chr14_25461770	14	25894747	4.39E-07–1.65E-09	18.34–23.11	FarmCPU, BLINK	Vitvi14g01522	Reductase

Trait	SNP	Chr	Position	p-value	Effect	Method	Gene ID	Gene description
							Vitvi14g01523	VvWRKY43 - WRKY domain-containing protein
CWi	be_C_A_chr3_5463929	3	4044392	2.50E-06–3.78E-06	15.76–1.64	FarmCPU, BLINK	Vitvi03g00359	PMT5 - Sugar transporter family
							Vitvi03g00360	PLATZ transcription factor
							Vitvi03g00361	Pentatricopeptide (PPR) repeat-containing protein
	chr4_16706884_C_T	4	16951942	1.10E-07	11.43	BLINK	Vitvi04g01164	Seipin-1
	be_T_C_chr6_7091603	6	7035547	2.04E-08	21.90	BLINK	Vitvi06g00614	Cell division control protein (CDC6)
							Vitvi06g00615	Ribosomal protein P0 (RPP0A) acidic 60S
	chr7_19824302_C_T	7	28297095	1.9E-08	7.53	FarmCPU	Vitvi07g01988	Translation factor GUF1 homolog
							Vitvi07g01989	MACPF domain-containing protein
	chr15_18113172_A_G	15	19889585	3.62E-06–3.86E-08	12.26–8.43	FarmCPU, BLINK	Vitvi15g00994	Zinc finger CCCH domain-containing protein 1
							Vitvi15g00995	Lipase class 3
							Vitvi15g01618	Poly(A) RNA polymerase mitochondrial-like central palm domain-containing protein
	chr18_20491804_A_G	18	23547288	1.09E-06–2.98E-07	6.20–19.56	FarmCPU, BLINK	Vitvi18g01679	TOD1/MUCI70 glycosyltransferase-like domain-containing protein
							Vitvi18g01681	VvCPK16-Calcium-dependent protein kinase
CW	chr1_5933624_A_G	1	5916771	2.05E-06–3.11E-06	0.67–27.14	FarmCPU, BLINK	Vitvi01g00524	Endoplasmic reticulum transmembrane protein
							Vitvi01g00525	U-box domain-containing protein
							Vitvi01g00526	Protein kinase domain-containing protein
	chr9_290178_C_T	9	290018	1.04E-06–5.56E-07	0.09–10.56	FarmCPU, BLINK	Vitvi09g00022	Uncharacterized protein
							Vitvi09g00023	1-phosphatidylinositol-3-phosphate 5-kinase
							Vitvi09g00026	Translation initiation factor eIF2B subunit alpha
							Vitvi09g00027	Translation initiation factor eIF2B subunit alpha
	chr10_17990933_A_C	10	24330783	3.84E-10–1.98E-07	4.87–11.25	FarmCPU, BLINK	Vitvi10g01581	Katanin p80 (WD repeat containing) subunit B 1

309 BW, berry weight; (B) CW, cluster weight; (C) BL, berry length; (D) CL, cluster length; (E) BWi, berry width; (F) CWi, cluster width.

310 SNPs associated with more than one trait share the same superscript.

Berry traits

In the GWAS, a total of 10 SNPs were significantly associated with BL (Fig 2A). These markers are distributed across Chrs 1, 2, 6, 8, 11, 14, 17, 18, and 19 (Fig 5). Three of the markers, those on Chr 6 (lb_G_A_chr6_1714702), Chr 14 (ae_C_T_chr14_16833185), and Chr 19 (chr19_23684078_C_T), were identified by both models. The phenotypic variance of BL explained by the significant SNPs ranged from 0.19–35.2%. The marker ae_C_T_chr14_16833185 on Chr 14, identified by both models, accounted for the highest percentage of the phenotypic variance in BL (PVE = 35.2%). The GWAS identified thirteen SNPs with significant associations with BWi across ten different chromosomes (1, 2, 5, 6, 7, 10, 11, 14, 15, and 16). Three of these SNP markers were identified by both models and explained 11.7% (lb_G_A_chr6_1714702; on Chr 6), 12.6% (chr6_14784507_A_G; on Chr 6), and 33% (ae_C_T_chr14_16833185; on Chr 14) of the phenotypic variance in BWi. In this study, the greatest number of significant marker–trait associations was identified for BW. The models identified a total of 20 SNPs, but none were identified simultaneously by both (Fig 2A). These markers are distributed across fourteen different chromosomes, excluding only chromosomes 7, 9, 10, 12, and 13. The percentage of PVE by these markers ranged from 0.01% (chr18_7027167_A_G; on Chr 14) to 19.2% (be_T_C_chr4_6544365; on Chr 4). Considering that the three berry traits together reflect berry size, QTLs were found for all three on Chrs 1, 2, 6, 11, and 14. Notably, two significant SNPs (chr2_3892599_A_G and lb_G_A_chr6_1714702) were shared among BL, BW, and BWi. Additionally, two SNPs (chr1_5754629_C_T and chr18_7027167_A_G) were common to BL and BW, two SNPs (chr5_15221260_C_T and be_T_G_chr3_12208763) were common to BW and BWi, and one SNP (ae_C_T_chr14_16833185) was shared between BL and BWi (Fig 2A).

Fig 5. Chromosomal localization of significant SNPs associated with berry- and cluster-related

traits. The chromosome number is shown at the top of each chromosome, and chromosome sizes are

depicted on a vertical scale (Mb).

A set of 80 distinct genes was associated with at least one of the berry size traits (S6 Table).

Among these genes, 26 were associated with BL, 50 with BW, and 26 with BWi. Ten of these genes

were associated with two traits simultaneously, and interestingly, six genes (Vitvi02g00402,

Vitvi02g00403, Vitvi02g00404, Vitvi06g00141, Vitvi06g00143, and Vitvi06g01615) were shared

among the three evaluated berry traits. Among the 80 candidate genes, 72 (90%) had at least one

Gene Ontology (GO) annotation, resulting in a total of 273 GO terms (S6 Table). The GO terms were

classified into three main categories: cellular component (50 GO terms), molecular function (99 GO

terms), and biological process (124 GO terms). In general, the noted GOs associated with berry

characteristics are linked to the synthesis of organic monomers necessary for cellular structures

(carbohydrates, lipids, and pigments). Terms related to growth (cell division and distension), control

of oxidative stress, accumulation of substances characteristic of ripe fruit, such as small

carbohydrates and aromatic molecules, and pre- and posttranslational regulation are also present (S6

Table). BL specifically identifies significant terms associated with DNA replication and

transcription, phospholipid metabolism, carotenoids, protein recycling, and phosphorylation. BW is

associated with terms related to secondary wall synthesis, lipids, membrane transporters, aromatic

amino acids, transcriptional regulation, and proteolysis. BWi is associated with terms related to

responses to gibberellic acid (GA) and jasmonic acid (JA), lipid synthesis, nutrient and ATP

transport, fucosylation, and ribosomal synthesis. Annotated terms associated with more than one trait

simultaneously were also identified, including those related to transcriptional regulation, auxin

response, cellulose, and anthocyanin synthesis. Among the terms commonly associated with all three

berry traits, the annotated terms are related to the synthesis and transport of small carbohydrates,

phospholipids, and triglycerides (S6 Table).

Among the several candidate genes associated with berry traits, Vitvi06g00141,

Vitvi06g00143, and Vitvi06g01615, which are linked to all three traits, encode members of the

glycosyl hydrolase family 1 (GH1). Genes encoding proteins from the multidrug and toxic compound

extrusion (MATE) family were also associated with the three traits, with Vitvi02g00403

(VvMATE50) linked to BW, BL, and BWi. In contrast, Vitvi08g00085 (VvMATE22) and

Vitvi08g00086 (VvMATE23) were specifically associated with BW. Among the genes related to

both BW and BL, Vitvi01g00512 and Vitvi01g00513 encode vacuolar iron transporter 1 (VIT1),

whereas Vitvi01g00511 encodes heat stress transcription factor A-8 (HSFA8). With respect to genes

associated with both BW and BWi, Vitvi05g01115 encodes MYB-like protein X (MYB-X), and

Vitvi15g01767 encodes auxin response factor 6 (ARF6). With respect to genes exclusively

associated with BW, notable candidates include Vitvi19g01778, which encodes a NAC domain-

containing protein (NAC TF), and Vitvi11g00554, which encodes a subunit of the actin-related

protein 2/3 complex (ARP2/3 complex subunit). With respect to genes linked solely to BL, we

identified genes encoding enzymes from the peptidase M16 family (Vitvi08g01906 and

Vitvi08g01910); Vitvi11g00146, a member of the nucleotide-binding site leucine-rich repeat (NB-

LRR) family; and a gene encoding an acetyltransferase component of the pyruvate dehydrogenase

complex (DLAT/E2). Finally, among the genes associated with BWi, Vitvi06g01133 encodes a

gibberellin-mediated growth regulator from the GRAS/SCARECROW-LIKE 3 family

(GRAS/SCL3), and Vitvi01g00473 encodes jasmonate ZIM-domain protein 6B-like (JAZ/TIFY6B).

Cluster traits

Marker–trait association analyses revealed a total of eight SNP markers associated with CL

(Fig 2B), explaining 0.1–24.2% of the phenotypic variance (S5 Table). These SNPs are distributed

across Chrs 1, 3, 4, 7, 8, 14, 18, and 19 (Fig 5). Notably, four of these markers

(chr3_16593567_A_G, chrUn_28901725_A_G, chr8_18600161_C_T, and lb_T_C_chr14_25461770

on Chrs 3, 4, 8, and 14, respectively) were identified simultaneously by both models (Fig 2B). With

respect to CW, four SNPs were identified, three of which (chr1_5933624_A_G, chr9_290178_C_T,

and chr10_17990933_A_C) were identified simultaneously by both models. These SNPs are

distributed across Chrs 1, 5, 9, and 10, explaining 0.6–27.1% of the phenotypic variance. For CWi,

the GWAS revealed 11 significantly associated SNPs on Chrs 3, 4, 5, 6, 7, 9, 14, 15, and 18. Four

SNPs were consistently identified by both methods, with two of these located in different regions of

Chr 15. The PVE values of the significant SNPs ranged from 0.15–21.8% for CWi. Only one

significant SNP was associated with more than one cluster trait (Fig 2B). The SNP

chr9_290178_C_T on Chr 9 was common to both CW and CWi.

A group of 52 distinct genes was pinpointed as potentially associated with cluster size traits.

Among them, 20 are exclusively linked to CL, 6 to CW, and 22 to CWi. Notably, four genes

(Vitvi09g00022, Vitvi09g00023, Vitvi09g00026, and Vitvi09g00027) are implicated in both CW and

CWi. Furthermore, of these 52 genes, 42 (80.76%) have at least one Gene Ontology (GO) annotation,

collectively providing 173 distinct GO terms, with 74 classified as biological processes, 30 as cellular

components, and 69 as molecular functions (S6 Table). In general, the annotated terms associated

with cluster traits are related to ribosomal RNA processing, protein synthesis, transport, and

packaging. For CL, in addition to protein-related processes, terms related to glutamine biosynthesis

were identified. In the case of CW, the annotated terms were linked to JA biosynthesis, lipid

catabolism, and cytoskeleton depolymerization. CWi was enriched for terms related to DNA and

ribosome biosynthesis, lipid catabolism, and calcium ion transport. When CW and CWi are

considered together, the terms associated with both traits indicate a role in translation initiation and

calcium-mediated signaling (S6 Table).

Notable among the genes associated with CW and CWi are Vitvi09g00026 and

Vitvi09g00027, which encode translation initiation factor 2B subunit alpha (eIF2Bα). In addition,

Vitvi01g00572, encoding translation initiation factor 3 subunit 2 (eIF3-S2), was associated with CL.

Other interesting genes associated with CL include Vitvi01g00571, belonging to the nitrate

transporter 1/peptide transporter family (NRT1/PTR); Vitvi14g01523, encoding a WRKY domain-

containing protein 43 (VvWRKY43); Vitvi04g02280, encoding protein phosphatase 2C (PP2C); and

Vitvi03g01151, encoding peptidyl-prolyl cis-trans isomerase (FKBP12). With respect to the genes

linked to CWi, notable candidates include Vitvi03g00359, which encodes a member of the

polyol/monosaccharide transporter (PMT) family; Vitvi06g00614, encoding cell division control

protein 6 (CDC6); Vitvi07g01989, encoding membrane attack complex/perforin (MACPF) domain-

containing protein; Vitvi15g00994, encoding zinc finger CCCH domain-containing protein 1; and

Vitvi15g00915, encoding cytochrome P450 78A40 (CYP78A40) (Table 1 and S6 Table).

Discussion
423

Association panel design
425

427   Genome-wide association mapping is a robust technique that facilitates the identification of

428 genes that regulate traits of interest. The effectiveness of this method relies on the genetic diversity

429 present within the germplasm used as association panels [71]. To precisely capture the genome-wide

430 distribution of linkage disequilibrium and pinpoint markers associated with various traits, it is

431 essential to include a wide array of diverse accessions. In this study, we selected 288 grapevine

genotypes from different origins (S1 Table) to encompass a substantial portion of the genetic and phenotypic diversity found in Brazilian germplasms. The IAC grapevine germplasm encompasses approximately 420 accessions, 100 of which are exclusive to this germplasm and originate from the institution's breeding program [53]. In a previous study, this collection was examined with microsatellite (SSR) markers, revealing a broad spectrum based on the species and the intended use of the accessions [50]. To design our grapevine association panel, we considered the results of genetic structure analyses generated from SSR studies. We selected the key genotypes from each genetic cluster, avoiding closely related individuals (i.e., first-degree relatives or mutants).

The 120 accessions from the previously established core collection [50] were prioritized in the selection process to enhance the representation of different alleles of interest across the genetic pools. This selection strategy successfully limited the structuring of the germplasm. In this study, the first three principal components accounted for 21.71% of the genomic population structure, revealing only moderate clustering among the *V. vinifera* accessions. This low level of structure in our population reflects the diverse geographic origins of the grape cultivars and the presence of various intra- and interspecific hybrids, which strengthened the association analyses. In grapevines, the crossbreeding of cultivars or natural hybridization, including among different species, results in hybrids with heterogeneous genetic compositions, making it challenging to assign these genotypes to specific genetic groups [72–74]. The presence of these hybrids with complex pedigrees, derived from crosses among three or more species, certainly contributed to a weaker structure in the diversity panel. Conversely, *V. vinifera* genotypes, despite having different uses and origins, still belong to the same species, resulting in more evident clustering.

QTLs for berry size-related traits

Traits related to berry and cluster size are closely associated with grapevine productivity and

quality. Berry size stands out as one of the most economically significant agronomic traits for table

grapes, acting as a crucial quality factor that shapes consumer preferences and acceptance [15]. It is

also a key determinant of juice composition and final wine quality, particularly in red wines [17].

Furthermore, natural variation in berry size significantly influences the volatile profiles of grape

berries [18]. Fruit size is typically assessed by considering three dimensions: weight, length, and

width. In the case of grape berries, as observed in this study, these three characteristics are typically

reported to have a strong positive correlation [7,15,45,75]. This suggests that berries larger in one

dimension, such as length, also tend to be larger in other dimensions, including width and weight

[76]. The high broad-sense heritability (H^2) estimates for BW, BL, and BWi obtained in this study

indicate that these traits are under strong genetic control. This aligns with previously reported high

H^2 values (ranging from 50–97%) for berry size-related traits in other populations [15,24,37,75],

suggesting that phenotypic variation is predominantly derived from genetic variance. These findings

imply that association analyses to identify favorable alleles linked to molecular markers hold

promising potential for MAS.

Several QTLs for berry size-related traits have already been reported, and distinct QTLs have

been detected in different genetic backgrounds. An extensive literature review was conducted to

investigate potential overlaps between the QTLs identified in this study and previously reported

QTLs associated with any berry size-related traits [15,22,31,32,34,37,40,44–46,77,78,23–30]. A

major berry weight QTL on Chr18 colocalized with the seedless trait was previously found in several

studies near the SSR marker VMC7F2 [22,30–32]. The GWAS findings in this study validate the

presence of significantly associated SNPs for BW on Chr18. However, the locations of the two loci

identified here differ from those found through biparental QTL mapping. This discrepancy from

previous biparental mapping studies may be due to the use of segregating populations derived from

crosses involving seedless cultivars. In GWASs using diversity panels, Guo et al. (2019) and Flutre et

al. (2022) reported that approximately 5–6 Mbp of the SNPs identified in the present study were

QTLs on Chr18, which also diverged from the BW QTL colocalized with the seedless trait. Some

seedless cultivars exhibit reduced berry weight at harvest, necessitating exogenous applications of

gibberellic acid and cluster thinning to maximize berry growth potential. This reduction may be due

to the absence of growth regulators typically produced by seeds [22,79]. Consequently, the markers

colocalized with the QTLs for seed number and seed size indirectly affect berry size [80].

In this study, the SNP markers associated with BW overlapped with previously reported

QTLs on chromosomes 1, 2, 3, 4 [37], 5 [44], 6 [46], 8 [22,37,44], 11 [22,26], 14 [37], 17 [22,44],

and 19 [37,44] (S5 Table). On Chr1, we identified two potential colocalizations with the QTLs

reported by Flutre et al. (2022) for BW. The first is in the region containing the SNP shared by both

BW and BL traits (chr1_5754629_C_T) and the SNP identified for BWi (chr1_5295802_C_T),

approximately 0.8 Mbp away. The second colocalization is with the SNP for BW

(chr1_22609084_C_T), located approximately 2.8 Mbp from one of the QTLs reported by these

authors. Our results also confirmed a QTL previously identified on Chr17 by Doligez et al. (2013).

This QTL was reported as a stable and large-effect QTL for berry weight that did not colocalize with

QTLs for seed traits. However, this QTL was identified using biparental populations and low-

resolution linkage maps. Recently, Thorat et al. (2024) identified a marker associated with BW in the

same region. By integrating high-throughput genotyping tools and using a diversity panel, they likely

increased the precision of the confidence intervals of the QTLs, aligning more accurately with the

QTLs identified in this study.

The SNP located on Chr2 (chr2_3892599_A_G), associated simultaneously with the three

berry size traits in this study, was found to be less than 1 Mbp from the marker associated with BWi

in the study by García-Abadillo et al. (2024) and approximately 1.4 Mbp from the region related to

BW by Flutre et al. (2022). The simultaneous association of this region with all three berry size-

related traits observed in this study is supported by the strong phenotypic correlations between them

(r > 0.9). A similar situation was observed on Chr6. The SNP simultaneously associated with all

three traits (BW, BL, and BWi) was found to be approximately 3 Mbp from the QTL reported by

Marrano et al. (2018) for BW. Few studies have focused on identifying QTLs related to BL and BWi

compared with BW, likely because of the high correlation among these traits and the intensive labor

required for phenotyping BL and BWi. Additionally, SNPs exclusively associated with BWi had the

lowest overlap rate with previously reported QTLs. However, based on our findings, we can infer

that many QTLs reported solely for BW may also be associated with other berry size-related traits. In

other words, the QTLs identified for one of these traits are likely to influence all dimensions of berry

size, probably because the same genetic factors govern these characteristics.

Many regions identified in this study for the three berry traits colocalized with QTLs

previously reported for different berry size-related characteristics. On Chr8, three distinct regions

were   associated   with   BW.   Two   of   these   regions   (ae_A_G_chr8_11440804   and

chr8_17424579_A_G) had previously been reported for this trait [22,37,44] and were also associated

with berry shape and BWi by García-Abadillo et al. (2024), respectively. The third region

(chr8_1499673_G_T) exactly matched the QTLs identified by Wang et al. (2022a) for BL and BWi.

Additionally, on Chr8, the SNP associated with BL (chr8_21881697_G_T) overlapped with the

QTLs identified by Flutre et al. (2022) for BW and by García-Abadillo et al. (2024) for BWi. The

SNP on Chr16 (chr16_14671632_C_T) associated with BWi likely corresponds to the QTL identified

by García-Abadillo et al. (2024) for berry shape. Because this trait was derived by the authors from

the ratio of berry height (or BL) to BWi, it can be concluded that these traits are highly correlated and

might correspond to the same QTL.

Furthermore, new potential QTLs were identified on chromosomes that are already associated

with berry size but in regions that are not consistent with previous studies. Previously unreported

associations for berry size were detected in regions of chromosomes 1, 4, 6, 7, 10, 11, 14, 15, 16, 17,

18,   and   19.   For   example,   no   overlaps   were   identified   for   the   SNP   on   Chr14

(ae_C_T_chr14_16833185), which was simultaneously associated with both models for BL and BWi

and had a high PVE value (30–35%). The differences in the positions of some QTLs compared with

those in previous studies suggest that fruit size-related traits are polygenic and controlled by different

causal polymorphisms potentially segregating among various populations [22]. Additionally, factors

such as genotyping density, varying phenotyping methods, and the statistical models employed can

complicate the precise identification of overlaps with QTLs reported in other studies [37].

QTLs for cluster-related traits

In grape production, subtraits associated with cluster architecture, including cluster and berry

size, number of berries per cluster, and cluster compactness, are crucial. These traits not only impact

fruit quality and yield but also influence factors such as disease susceptibility, sunlight exposure, and

uniform ripening [27,44]. The weight, length, and width of clusters are among the subtraits

commonly used to objectively and quantitatively assess cluster compactness and size in research

studies. The genetic mechanisms underlying these traits in grapes are complex and influenced by

both viticultural practices and environmental factors, resulting in significant year-to-year variation

[11,44,81,82]. Consequently, the influence of numerous minor genetic contributions makes it

difficult for the literature to reach a consensus on the location of their QTL. Moreover, population or

environmental factors could cause these QTLs to be unstable [45]. Nonetheless, certain overlaps were

detected. The first concerns the SNP on Chr10 recognized by both models linked to CW, which

coincided with a QTL identified by Richter et al. (2019) connected with the same trait mapped in a

biparental population. The region identified on Chr15 associated with CWi corresponded to the QTL

found by Flutre et al. (2022) for the same trait. Finally, the SNP linked to CL on Chr14 overlapped

with the QTL documented by Thorat et al. (2024) for the same trait, as well as with the QTL

discovered by Marrano et al. (2018) related to CW.

In addition, several other instances of QTL colocalization were observed, where the QTLs

reported for one cluster differed from the trait associated in this study but were correlated (S5 Table).

We found moderate to high correlation levels between cluster length, width, and weight. However,

they have been variably reported in other studies, ranging from high to moderate and even low

[7,75,83,84]. In addition to being heavily influenced by nongenetic factors, these traits are also

affected by other characteristics, such as berry size, the number of berries per cluster, and rachis size,

which likely contribute to the variation in their correlation. Nevertheless, these QTL overlaps

associated with different cluster traits highlight the interconnected nature of these characteristics and

suggest that common genetic factors may influence multiple aspects of cluster architecture.

Additionally, as shown on Chr1, Chr7, Chr8, and Chr14, some regions in this study were

associated with both berry and cluster traits simultaneously (Fig 5). Despite being significant, we

identified a weak to moderate correlation between cluster and berry traits (S3 Fig). However, cluster

traits are indirectly linked to berry traits. For example, as cluster size increases, the likelihood of

larger berries may also increase, or larger and heavier berries will contribute more to the total cluster

weight [83]. Of course, numerous other factors influence the relationships between these traits, such

as the number of berries per cluster, berry density, environmental factors, and gene☐gene

interactions, which can also moderate the expression of these traits. These factors can result in lower

phenotypic correlations while still reflecting a shared genetic basis. It remains challenging to suggest

a pleiotropic effect of a locus on several phenotypic traits. However, the proximity of the QTLs for

berry- and cluster-related subtraits in these regions offers an opportunity for MAS. This situation

could be leveraged using a small range of molecular markers from these QTL regions to select for

larger or smaller berry volumes, accompanied by larger or medium-sized clusters, depending on the

intended use for table grapes or wine. By doing so, it may be possible to capture several coinherited

traits, potentially leading to a more efficient selection process that aligns with the desired

characteristics for each application [27].

Efforts have been made to study the genetic determinants of berry and cluster size through

QTL mapping using segregating populations [15,22,27,29–31,34], yielding valuable insights into the

genetic inheritance of these complex traits. However, these analyses used populations with relatively

limited genetic variability and often failed to capture all recombination events. Using the GWAS

approach, we identified many associated markers for each of the studied berry and cluster traits, with

some markers explaining substantial phenotypic variation. Despite new associations being identified

in this study, overlap with previously reported QTLs was observed (S5 Table). Most of these

overlaps were with recent studies that also conducted GWAS on diversity panels [37,44,45]. This

reproducibility stems from the technique's ability to leverage historical recombination events to

identify haplotype blocks associated with phenotypes of interest across the diversity of crops,

identifying stable QTLs that are not population specific.

Candidate genes for berry size traits

Fruit development is typically categorized into three distinct stages: fruit set, fruit growth, and

fruit maturation [85]. Fleshy fruiting species undergo phases of cell division and expansion following

fruit set, which allows the determination of fruit size before maturation and ripening. Plant hormones

regulate various aspects of growth and development, including fruit size, by responding to both

environmental and endogenous signals [86]. At the cellular and molecular levels, phytohormones can

influence gene expression and transcription, as well as cell division and growth [87]. In this study,

we identified several genes related to hormones that are associated with berry size. Most of these

genes were linked to the plant hormones auxin, gibberellic acid (GA), and JA. Studies have shown that auxin and GA are early indicators of fruit setting and growth [88–92]. Genes related to these hormones have been previously linked to berry size and shape traits [22,26,44,68,77]. Auxins play crucial roles in regulating nearly all aspects of plant growth and development by influencing cell division, cell growth, and cell differentiation [93,94]. Auxin response factors (ARFs) are key transcription factors that mediate auxin signaling in plants by binding to auxin-response elements in the promoter regions of early auxin-response genes, thereby regulating auxin-induced gene responses [95,96]. Class A ARFs can also respond to environmental signals through auxin-independent pathways, regulating plant growth and development [97]. QTLs associated with important traits have been increasingly linked to auxin, suggesting that auxin engineering holds tremendous potential for manipulating crop architecture in plant breeding [98].

GA is known to be essential for regulating berry expansion. Research has shown that exogenous GA can promote the division and expansion of grape pulp cells, increase sugar accumulation, and increase fruit size [86,99–101]. The gene Vitvi06g01133, associated with BWi, has been identified in grapevine as a GRAS gene belonging to the SCL3 subfamily and is predominantly expressed in the stem, seed, and berry flesh [102]. Proteins from the SCL3 subfamily are key components in the GA signaling pathway and cooperate to mediate cell elongation during root development in *Arabidopsis thaliana* [103]. The GRAS gene family is a plant-specific group of transcription factors that play crucial roles in various metabolic pathways, including plant growth, development, fruit ripening, and stress response [104,105]. The jasmonate–ZIM domain protein acts as an inhibitory factor in the JA pathway, which is involved in regulating plant development and the response of plants to biotic and abiotic stresses [106–109].

Additionally, several genes involved in responses to abiotic and biotic stresses were associated with berry size traits in this study. This associations likely occur because abiotic and biotic stresses can significantly affect plant growth, yield, and quality by disrupting physiological

processes, weakening defense mechanisms, and reducing overall productivity [110]. Heat stress is among the major threats in viticulture [111], significantly affecting berry ripening, quality, and yield when air temperatures exceed 35°C [112]. As a result, high temperatures directly impact berry size [113]. The gene Vitvi01g00511, associated with both BW and BL, encodes an HSF, the most important transcription factor family involved in the response of plants to heat [114]. To reduce the severity of destructive effects, plants use a range of defensive strategies. NB-LRR family genes encode intracellular resistance (R) proteins that directly or indirectly recognize and activate defense responses, including the hypersensitive response and systemic acquired resistance [115,116]. Several transcription factors play crucial roles in regulating the internal defense system of plants. The MYB and NAC families of transcriptional regulators play essential roles in regulating various vital biological processes during plant growth, development, and stress responses [117,118]. Specifically, genes encoding NAC domain-containing proteins have been associated with the early development of grape flowers and berries [119], with several studies suggesting their involvement in determining berry size variation [15,68,78,120,121].

In plants, members of the GH1 family have been identified as key players in various biological processes related to growth and development [122,123]. Notably, substantial evidence indicates that GH1 genes are also involved in the activation of defense compounds in response to stress and the release of active plant phytohormones [124–126]. Studies have demonstrated that in grapevines, β-glucosidase enzymes, a subclass of the GH1 family, participate in responses to waterlogging stress [127], low-temperature stress [128], and abscisic acid (ABA) biosynthesis [129].

Several transport-related genes were associated with the three berry traits across different chromosomes and play crucial roles in plant defense, growth, and development. The transport and conversion of photosynthetic products are closely tied to sugar accumulation in fruit, with total soluble sugars increasing during growth and peaking at maturity or ripening [130,131]. The genes Vitvi02g00403 (VvMATE50), Vitvi08g00085 (VvMATE22), and Vitvi08g00086 (VvMATE23)

belong to the MATE family, which is among the largest families of transporter genes in plants and is involved in various physiological processes throughout plant growth and development [132]. In grapevine, the gene VvMATE50 belongs to a group of genes involved in the transport or signaling of phytohormones such as ABA, auxin, and salicylic acid (SA) and plays a significant role in grape growth and defense responses. The genes VvMATE22 and VvMATE23 are part of a MATE group and function in the transport of flavonoids and alkaloids [133]. According to Watanabe et al. (2022), VvMATE22 is highly expressed in seed and berry tissues during early developmental stages, with expression levels decreasing as development progresses. These findings suggest that VvMATE22 plays a crucial role in regulating the transport of substances that impact berry growth and development. Among the transport-related genes identified in this study, two encode VIT1, which plays a crucial role in maintaining iron homeostasis and regulating various cellular functions essential for fruit development [134,135].

In plant cells, actin filaments indirectly influence cell shape by regulating the transport properties of organelles and cargo molecules, which in turn alter the mechanical properties of the cell wall [136]. The Arp2/3 complex serves as a critical regulator of actin filament dynamics across a wide range of eukaryotic cells [137]. This complex nucleates new actin filaments and stimulates the formation of branched actin networks, which are essential for various cellular processes, including cell shape modulation, cell wall assembly, intracellular trafficking, and cell expansion [138–142]. In the context of grape berry development, it is plausible that the Arp2/3 complex regulates the extent of cell expansion in the pericarp, directly controlling the volumetric increase that defines berry size. Moreover, its participation in vesicle trafficking may affect the deposition of cell wall components, influencing fruit hardness and elasticity, which are indirectly associated with berry development potential.

Additional genes identified through GWAS that may directly or indirectly affect the determination of final fruit size include members of the peptidase M16 family, which are involved in

protein processing and play critical roles in maintaining cellular viability by ensuring protein and

energy homeostasis [143–145]. In addition, a gene encoding DLAT/E2 was identified, which may

contribute to the regulation of cell expansion. An adequate supply of acetyl-CoA provided by this

complex is essential for the biosynthesis of membrane lipids, which are integral constituents of the

cell wall, and for energy production (ATP), processes that are critical during fruit growth and

development [146,147].

Candidate genes for cluster-related traits

Cluster compactness and size significantly influence the commercial value of both wine and

table grapes. These traits are highly complex, with various berry and cluster attributes contributing to

their variation across different varieties, each influenced by their genetic architecture [13,148].

Among these attributes, berry dimensions are key factors that directly affect cluster compactness and

size [149]. Notably, significant SNPs on Chr1 and 14 associated with CL are near genes encoding

proteins from the NRT1/PTR and WRKY domain-containing families. These families were identified

by García-Abadillo et al. (2024) as potential candidates for controlling both berry length and width.

NRT1/PTR genes, originally identified as nitrate or peptide transporters in plants, are involved in

transporting a wide range of substrates, such as amino acids, nitrate, auxin, JA, abscisic acid (ABA),

GA, and glucosinolates [150], which play essential roles in the molecular mechanisms influencing

fruit size determination [86]. Grape WRKY transcription factors play crucial regulatory roles in both

fruit development and overall grapevine growth. They are also vital for mediating plant responses to

various biotic and abiotic stresses [151]. Specifically, the gene encoding VvWRKY43 identified in

this study is potentially involved in regulating the stilbene biosynthetic pathway [152]. Stilbenes are

important compounds that protect plants from pests, pathogens, and abiotic stresses [153].

Plants have evolved a wide range of defense strategies to cope with biotic and abiotic

stresses; however, these responses are often energetically costly and divert resources away from

growth, development, and reproduction, including during fruit development [154]. In this study,

distinct genes associated with CL and CWi were identified, many of which are involved in stress-

related pathways. Members of the MACPF family play crucial roles in maintaining the balance

between plant growth and immunity [155], contributing to both biotic and abiotic stress responses,

likely through interactions with JA and SA signaling pathways [156–158]. Additionally, genes

encoding PP2C have been shown to participate in various stress responses by modulating growth

factors, regulating phytohormones, and controlling metabolic processes in several important plant

species [159]. In grapevine, PP2Cs are actively involved in the ABA signal transduction pathway

[160] and have been proposed as candidate genes affecting BL [44]. Zinc finger CCCH-type proteins

also play significant roles in plant adaptation to various abiotic stresses, including oxidative stress,

salinity, drought, flooding, and cold temperatures, across different species [161]. In grapevine, these

proteins are potentially involved in not only stress responses but also growth and developmental

processes [162]. Furthermore, they are involved in various developmental processes in plants,

including seed development [163], flowering and senescence [164], plant architecture [165], and

secondary wall synthesis [166].

Genes related to translation and protein synthesis were identified in genomic regions near

SNPs associated with all three cluster traits. In particular, genes encoding eIF3-S2 and eIF2Bα were

detected. These factors play essential roles in the initiation phase of protein synthesis, a critical step

for assembling the ribosomal machinery required to translate mRNA into functional proteins. Their

activity ensures the efficiency and fidelity of protein biosynthesis, which is fundamental for

supporting cellular processes such as proliferation, differentiation, and adaptation to environmental

cues [167,168]. The efficiency of translation, which is regulated by the eIF2Bα and eIF-3 complexes,

directly impacts the production of proteins necessary for cell division and expansion [169,170],

which are critical processes for fruit development. Other noteworthy genes involved in cell cycle

regulation and associated with cluster traits include FKBP12 and CDC6. Plant FKBPs perform

diverse functions in essential biological processes related to growth and development [171].

Specifically, FKBP12 has been reported to participate in cell cycle regulation and embryo

development, as well as in controlling the direction of pollen tube growth in *Arabidopsis thaliana*

and *Picea wilsonii* [172–174]. Moreover, CDC6 plays a central role in the initiation of DNA

replication, ensuring accurate and coordinated genome duplication during the cell cycle [175].

Modulation of the expression or activity of these genes may directly impact the rate of cell division

in developing tissues, thereby influencing fruit growth and ultimately impacting grape cluster size.

Cluster traits are strongly associated with flower development and fertilization, as cluster

growth depends on both cell proliferation and expansion during the development of floral and

peduncle structures [176,177]. Reproductive development is an energy-demanding process, from

floral initiation to fruit maturation [178]. The transport of soluble sugars across membranes is a

tightly regulated process mediated by sugar transporters belonging to multiple gene families

[130,179,180]. In grapevine, PMTs have been detected in different vegetative organs [181].

Specifically, the VvPMT5 gene is expressed at high levels in mature leaves, petioles, and tendrils,

with its expression peaking during the fruit set stage [181]. The observed expression pattern suggests

a potential role for the PMT5 gene in regulating the development and structural definition of the

cluster axis. Genes from the PLATZ transcription factor family have also been implicated in flower

development in grapevine [182]. They are also known to participate in various aspects of plant

growth and development across different species, including stress responses, cell proliferation, and

seed development [183–185]. Notably, in rice, the PLATZ gene *GL6* positively regulates cell

division by increasing the number of cells in the spikelet hull, ultimately leading to larger grains

[186].

Extensive research has demonstrated that several CYP78A genes are also involved in the

regulation of seed development and organ size [187]. Arabidopsis CYP78A9 plays a role in

controlling floral organ size and has redundant functions with those of CYP78A6 in integument

development [188,189]. Similarly, TaCYP78A3 influences wheat seed size by affecting the extent of

integument cell proliferation [190]. In maize, CYP78A1 stimulates leaf growth by extending the

duration of cell division [191]. In sweet cherry, PaCYP78A6 acts redundantly with PaCYP78A9 to

influence fruit size during growth and development [192]. These findings support the potential

involvement of the candidate gene CYP78A40 in determining the final size of the grape cluster.

Other candidate genes identified in this study may influence various grape cluster traits at

different levels. The regions identified in GWASs typically contain blocks of credible SNPs, and a

major challenge is to pinpoint the causal gene in that locus of interest [193]. Our association study

revealed numerous QTLs with multiple genes for various grape yield traits. Further investigation into

the expression patterns of these candidate genes at specific stages of fruit development, along with an

analysis of their regulatory networks, will provide deeper insights into how these genes contribute to

the observed variations in the studied berry and cluster traits.

Conclusion

This study demonstrated the feasibility of conducting a GWAS in a perennial fruit crop such

as grapevine for complex traits related to yield and quality, which are crucial for breeding. Using a

grapevine diversity panel with accessions phenotyped over 12 seasons under subtropical conditions,

we identified numerous SNP markers that are significantly associated with berry and cluster-related

traits. Notably, 67.8% of these markers were located within annotated genes, with some explaining a

substantial portion of the observed variation. Additionally, we confirmed several previously known

genomic regions and identified new regions linked to the traits studied. This study also connected

different traits to QTLs reported in earlier research, highlighting the pleiotropic nature of these regions. Several predicted genes have emerged as strong candidates for influencing berry size and cluster architecture, as they are involved in critical functions such as growth regulation, hormone control, protein synthesis, stress response, and other essential physiological processes. We anticipate that the identified marker–trait associations and candidate genes will significantly advance our understanding of the molecular basis underlying cluster architecture and berry size, paving the way for further validation, functional investigation, and gene editing approaches. Ultimately, these findings provide valuable tools for grapevine breeders, contributing to the development of elite cultivars and benefiting the grapevine research community.

Acknowledgments

The authors are grateful to the São Paulo Research Foundation (FAPESP), the Coordination for the Improvement of Higher Education Personnel (CAPES), and the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for their support of the project and the researchers.

References

1.   OIV. State of the World Vine and Wine Sector in 2022. 2023 [cited 22 May 2023]. Available: https://www.oiv.int/sites/default/files/documents/2023_SWVWS_report_EN.pdf

2.   Nanni A, Parisi M, Colonna M. Wine By-Products as Raw Materials for the Production of Biopolymers and of Natural Reinforcing Fillers: A Critical Review. Polymers (Basel). 2021;13: 381. doi:10.3390/polym13030381

3.   Alston JM, Sambucci O. Grapes in the World Economy. In: Cantu D, Walker MA, editors.

The Grape Genome. Cham: Springer International Publishing; 2019. pp. 1–24.

doi:10.1007/978-3-030-18601-2

4.   Terral JF, Tabard E, Bouby L, Ivorra S, Pastor T, Figueiral I, et al. Evolution and history of

grapevine (Vitis vinifera) under domestication: new morphometric perspectives to understand

seed domestication syndrome and reveal origins of ancient European cultivars. Ann Bot.

2010;105: 443–455. doi:10.1093/aob/mcp298

5.   Ferreira V, Pinto-Carnide O, Arroyo-García R, Castro I. Berry color variation in grapevine as

a source of diversity. Plant Physiol Biochem. 2018;132: 696–707.

doi:10.1016/j.plaphy.2018.08.021

6.   Gray DJ, Li ZT, Dhekney SA. Precision breeding of grapevine (Vitis vinifera L.) for improved

traits. Plant Sci. 2014;228: 3–10. doi:10.1016/j.plantsci.2014.03.023

7.   Migicovsky Z, Sawler J, Gardner KM, Aradhya MK, Prins BH, Schwaninger HR, et al.

Patterns of genomic and phenomic diversity in wine and table grapes. Hortic Res. 2017;4:

17035. doi:10.1038/hortres.2017.35

8.   Anand A, Subramanian M, Kar D. Breeding techniques to dispense higher genetic gains. Front

Plant Sci. 2023;13. doi:10.3389/fpls.2022.1076094

9.   Edge-Garza DA, Luby JJ, Peace C. Decision support for cost-efficient and logistically feasible

marker-assisted seedling selection in fruit breeding. Mol Breed. 2015;35: 223.

doi:10.1007/s11032-015-0409-z

10. Delrot S, Grimplet J, Carbonell-Bejerano P, Schwandner A, Bert P-F, Bavaresco L, et al.

Genetic and Genomic Approaches for Adaptation of Grapevine to Climate Change. Genomic

Designing of Climate-Smart Fruit Crops. Cham: Springer International Publishing; 2020. pp.

157–270. doi:10.1007/978-3-319-97946-5_7

11. Ohana-Levi N, Cohen Y, Munitz S, Michaelovsky R, Ferman Mintz D, Hagag N, et al. The

response of yield, number of clusters, and cluster weight to meteorological factors and

irrigation practices in grapevines: A multiexperiment study. Sci Hortic (Amsterdam).

2024;326: 112761. doi:10.1016/j.scienta.2023.112761

12. Wenter A, Zanotelli D, Montagnani L, Tagliavini M, Andreotti C. Effect of different timings

and intensities of water stress on yield and berry composition of grapevine (cv. Sauvignon

blanc) in a mountain environment. Sci Hortic (Amsterdam). 2018;236: 137–145.

doi:10.1016/j.scienta.2018.03.037

13. Tello J, Ibáñez J. Review: Status and prospects of association mapping in grapevine. Plant Sci.

2023;327: 0–2. doi:10.1016/j.plantsci.2022.111539

14. Wang Z, Zhou J, Xu X, Perl A, Chen S, Ma H. Adoption of table grape cultivars: An attribute

preference study on Chinese grape growers. Sci Hortic (Amsterdam). 2017;216: 66–75.

doi:10.1016/j.scienta.2017.01.001

15. Wang H, Yan A, Wang X, Zhang G, Liu Z, Xu H, et al. Identification of QTLs and candidate

genes controlling berry size in table grape by integrating QTL and transcriptomic analysis. Sci

Hortic (Amsterdam). 2022;305: 111403. doi:10.1016/j.scienta.2022.111403

16. Chen W-K, He F, Wang Y-X, Liu X, Duan C-Q, Wang J. Influences of Berry Size on Fruit

Composition and Wine Quality of Vitis vinifera L. cv. 'Cabernet Sauvignon' Grapes. South

African J Enol Vitic. 2018;39. doi:10.21548/39-1-2439

17. Martin D, Lindsay M, Kilmartin P, Dias Araujo L, Rutan T, Yvon M, et al. Grape berry size is

a key factor in determining New Zealand Pinot noir wine composition. OENO One. 2022;56:

389–411. doi:10.20870/oeno-one.2022.56.2.5436

18. Xie S, Tang Y, Wang P, Song C, Duan B, Zhang Z, et al. Influence of natural variation in

berry size on the volatile profiles of Vitis vinifera L. cv. Merlot and Cabernet Gernischt

grapes. Amancio S, editor. PLoS One. 2018;13: e0201374. doi:10.1371/journal.pone.0201374

19. Walker RR, Blackmore DH, Clingeleffer PR, Kerridge GH, Rühl EH, Nicholas PR. Shiraz

850  berry size in relation to seed number and implications for juice and wine composition. Aust J

851  Grape Wine Res. 2005;11: 2–8. doi:10.1111/j.1755-0238.2005.tb00273.x

852 20. Tello J, Ibáñez J. What do we know about grapevine bunch compactness? A state-of-the-art

853  review. Aust J Grape Wine Res. 2018;24: 6–23. doi:10.1111/ajgw.12310

854 21. Torres-Lomas E, Lado-Bega J, Garcia-Zamora G, Diaz-Garcia L. Segment Anything for

855  Comprehensive Analysis of Grapevine Cluster Architecture and Berry Properties. Plant

856  Phenomics. 2024;6. doi:10.34133/plantphenomics.0202

857 22. Doligez A, Bertrand Y, Farnos M, Grolier M, Romieu C, Esnault F, et al. New stable QTLs for

858  berry weight do not colocalize with QTLs for seed traits in cultivated grapevine (Vitis vinifera

859  L.). BMC Plant Biol. 2013;13: 217. doi:10.1186/1471-2229-13-217

860 23. Viana AP, Riaz S, Walker MA. Genetic dissection of agronomic traits within a segregating

861  population of breeding table grapes. Genet Mol Res. 2013;12: 951–964.

862  doi:10.4238/2013.April.2.11

863 24. Houel C, Chatbanyong R, Doligez A, Rienth M, Foria S, Luchaire N, et al. Identification of

864  stable QTLs for vegetative and reproductive traits in the microvine (Vitis vinifera L.) using the

865  18 K Infinium chip. BMC Plant Biol. 2015;15: 205. doi:10.1186/s12870-015-0588-0

866 25. Zhao YH, Guo YS, Lin H, Liu ZD, Ma HF, Guo XW, et al. Quantitative trait locus analysis of

867  grape weight and soluble solid content. Genet Mol Res. 2015;14: 9872–9881.

868  doi:10.4238/2015.August.19.21

869 26. Ban Y, Mitani N, Sato A, Kono A, Hayashi T. Genetic dissection of quantitative trait loci for

870  berry traits in interspecific hybrid grape (Vitis labruscana × Vitis vinifera). Euphytica.

871  2016;211: 295–310. doi:10.1007/s10681-016-1737-8

872 27. Richter R, Gabriel D, Rist F, Töpfer R, Zyprian E. Identification of colocated QTLs and

873  genomic regions affecting grapevine cluster architecture. Theor Appl Genet. 2019;132: 1159–

874  1177. doi:10.1007/s00122-018-3269-1

28. Underhill A, Hirsch C, Clark M. Image-based Phenotyping Identifies Quantitative Trait Loci

for Cluster Compactness in Grape. J Am Soc Hortic Sci. 2020;145: 363–373.

doi:10.21273/JASHS04932-20

29. Fanizza G, Lamaj F, Costantini L, Chaabane R, Grando MS. QTL analysis for fruit yield

components in table grapes (Vitis vinifera). Theor Appl Genet. 2005;111: 658–664.

doi:10.1007/s00122-005-2016-6

30. Correa J, Ravest G, Laborie D, Mamani M, Torres E, Muñoz C, et al. Quantitative trait loci for

the response to gibberellic acid of berry size and seed mass in tablegrape (V itis vinifera L.).

Aust J Grape Wine Res. 2015;21: 496–507. doi:10.1111/ajgw.12141

31. Cabezas JA, Cervera MT, Ruiz-García L, Carreño J, Martínez-Zapater JM. A genetic analysis

of seed and berry weight in grapevine. Gustafson JP, editor. Genome. 2006;49: 1572–1585.

doi:10.1139/g06-122

32. Mejía N, Gebauer M, Muñoz L, Hewstone N, Muñoz C, Hinrichsen P. Identification of QTLs

for Seedlessness, Berry Size, and Ripening Date in a Seedless x Seedless Table Grape

Progeny. Am J Enol Vitic. 2007;58: 499–507. doi:10.5344/ajev.2007.58.4.499

33. Costantini L, Battilana J, Lamaj F, Fanizza G, Grando M. Berry and phenology-related traits

in grapevine (Vitis vinifera L.): From Quantitative Trait Loci to underlying genes. BMC Plant

Biol. 2008;8: 38. doi:10.1186/1471-2229-8-38

34. Doligez A, Bouquet A, Danglot Y, Lahogue F, Riaz S, Meredith C, et al. Genetic mapping of

grapevine (Vitis vinifera L.) applied to the detection of QTLs for seedlessness and berry

weight. Theor Appl Genet. 2002;105: 780–795. doi:10.1007/s00122-002-0951-z

35. Vezzulli S, Doligez A, Bellin D. Molecular Mapping of Grapevine Genes. 1st ed. In: Cantu D,

Walker MA, editors. The Grape Genome. 1st ed. Cham: Springer Nature; 2019. pp. 103–136.

doi:10.1007/978-3-030-18601-2

36. Moura-furlan MF, de Oliveira GL, Rodrigues CS, Paioli-Pires EJ. The role of plant breeding

in grapevine production. Advances in Botanical Research. Elsevier; 2024. pp. 1–40.

doi:10.1016/bs.abr.2024.01.001

37.   Flutre T, Le Cunff L, Fodor A, Launay A, Romieu C, Berger G, et al. A genome-wide

association and prediction study in grapevine deciphers the genetic architecture of multiple

traits and identifies genes under many new QTLs. G3 Genes, Genomes, Genet. 2022;12.

doi:10.1093/g3journal/jkac103

38.   Myles S, Boyko AR, Owens CL, Brown PJ, Grassi F, Aradhya MK, et al. Genetic structure

and domestication history of the grape. Proc Natl Acad Sci. 2011;108: 3530–3535.

doi:10.1073/pnas.1009363108

39.   Laucou V, Launay A, Bacilieri R, Lacombe T, Andre MT De, Hausmann L, et al. Extended

diversity analysis of cultivated grapevine Vitis vinifera with 10K genome-wide SNPs. PLoS

One. 2018;13: 1–27. doi:https://doi.org/10.1371/journal.pone.0192540

40.   Guo DL, Zhao HL, Li Q, Zhang GH, Jiang JF, Liu CH, et al. Genome-wide association study

of berry-related traits in grape [Vitis vinifera L.] based on genotyping-by-sequencing markers.

Hortic Res. 2019;6. doi:10.1038/s41438-018-0089-z

41.   Zhang H, Fan X, Zhang Y, Jiang J, Liu C. Identification of favorable SNP alleles and

candidate genes for seedlessness in Vitis vinifera L. using genome-wide association mapping.

Euphytica. 2017;213: 1–13. doi:10.1007/s10681-017-1919-z

42.   Wang M, Qi Z, Thyssen GN, Naoumkina M, Jenkins JN, McCarty JC, et al. Genomic

interrogation of a MAGIC population highlights genetic factors controlling fiber quality traits

in cotton. Commun Biol. 2022;5: 1–12. doi:10.1038/s42003-022-03022-7

43.   Amyotte B, Bowen AJ, Banks T, Rajcan I, Somers DJ. Mapping the sensory perception of

apple using descriptive sensory evaluation in a genome wide association study. PLoS One.

2017;12. doi:10.1371/journal.pone.0171710

44.   Thorat KD, Upadhyay A, Samarth RR, Machchhindra SR, Jagtap MA, Kushwaha K, et al.

Genome-wide association analysis to identify genomic regions and predict candidate genes for

bunch traits in grapes (Vitis vinifera L.). Sci Hortic (Amsterdam). 2024;328.

doi:10.1016/j.scienta.2024.112882

45.   García-Abadillo J, Barba P, Carvalho T, Sosa-Zuniga V, Lozano R, Carvalho HF, et al.

Dissecting the Complex Genetic Basis of Pre and Postharvest Traits in Vitis vinifera L . using

Genome-Wide Association Studies . Hortic Res. 2024. doi:10.1093/hr/uhad283

46.   Marrano A, Micheletti D, Lorenzi S, Neale D, Grando MS. Genomic signatures of different

adaptations to environmental stimuli between wild and cultivated Vitis vinifera L. Hortic Res.

2018;5: 34. doi:10.1038/s41438-018-0041-2

47.   Foria S, Magris G, Jurman I, Schwope R, De Candido M, De Luca E, et al. Extent of wild-to-

crop interspecific introgression in grapevine (Vitis vinifera) as a consequence of resistance

breeding and implications for the crop species definition. Hortic Res. 2022;9.

doi:10.1093/hr/uhab010

48.   Morgil H, Can Gercek Y, Tulum I. Single Nucleotide Polymorphisms (SNPs) in Plant

Genetics and Breeding. The Recent Topics in Genetic Polymorphisms. IntechOpen; 2020.

doi:10.5772/intechopen.91886

49.   Maia JDG, Camargo UA, Tonietto J, Zanus MC, Quecini V, Ferreira ME, et al. Grapevine

breeding programs in Brazil. In: Reynolds A, editor. Grapevine Breeding Programs for the

Wine Industry. Elsevier; 2015. pp. 247–271. doi:10.1016/B978-1-78242-075-0.00011-9

50.   de Oliveira GL, de Souza AP, de Oliveira FA, Zucchi MI, de Souza LM, Moura MF. Genetic

structure and molecular diversity of Brazilian grapevine germplasm: Management and use in

breeding programs. Goel S, editor. PLoS One. 2020;15: e0240665.

doi:10.1371/journal.pone.0240665

51.   Alvares CA, Stape JL, Sentelhas PC, de Moraes Gonçalves JL, Sparovek G. Köppen's climate

classification map for Brazil. Meteorol Zeitschrift. 2013;22: 711–728. doi:10.1127/0941-

950    2948/2013/0507

951 52. Santos HG, Jacomine PKT, Anjos LHC, Oliveira VA, Lumbreras JF, Coelho MR, et al.

952    Sistema brasileiro de classificação de solos. 5th ed. Brasília, DF, Brazil: Embrapa; 2018.

953    Available: https://www.embrapa.br/busca-de-publicacoes/-/publicacao/1094003/sistema-

954    brasileiro-de-classificacao-de-solos

955 53. de Oliveira GL, Niederauer GF, de Oliveira FA, Rodrigues CS, Hernandes JL, de Souza AP, et

956    al. Genetic diversity, population structure and parentage analysis of Brazilian grapevine

957    hybrids after half a century of genetic breeding. Sci Hortic (Amsterdam). 2023;311.

958    doi:10.1016/j.scienta.2023.111825

959 54. Fox J, Weisberg S. An R Companion to Applied Regression. Third. Thousand Oaks, CA:

960    Sage; 2019.

961 55. Covarrubias-Pazaran G. Genome-Assisted Prediction of Quantitative Traits Using the R

962    Package sommer. Zhang A, editor. PLoS One. 2016;11: e0156744.

963    doi:10.1371/journal.pone.0156744

964 56. Di Guardo M, Micheletti D, Bianco L, Koehorst-van Putten HJJ, Longhi S, Costa F, et al.

965    ASSIsT: an automatic SNP scoring tool for in- and outbreeding species. Bioinformatics.

966    2015;31: 3873–3874. doi:10.1093/bioinformatics/btv446

967 57. Granato I, Fritsche-Neto R. snpReady: Preparing Genotypic Datasets in Order to Run

968    Genomic Analysis. 2018 [cited 15 May 2023]. Available: https://cran.r-

969    project.org/web/packages/snpReady/index.html

970 58. Hastie T, Tibshirani R, Narasimhan B. impute: Imputation for microarray data. 2020 [cited 10

971    Feb 2023] pp. 1–5. Available: https://git.bioconductor.org/packages/impute

972 59. Wang J, Zhang Z. GAPIT Version 3: Boosting Power and Accuracy for Genomic Association

973    and Prediction. Genomics, Proteomics Bioinforma. 2021;19: 629–640.

974    doi:10.1016/j.gpb.2021.08.005

60.　Challa S, Neelapu NRR. Genome-Wide Association Studies (GWAS) for Abiotic Stress

Tolerance in Plants. Biochemical, Physiological and Molecular Avenues for Combating

Abiotic Stress Tolerance in Plants. Elsevier; 2018. pp. 135–150. doi:10.1016/B978-0-12-

813066-7.00009-7

61.　VanRaden PM. Efficient Methods to Compute Genomic Predictions. J Dairy Sci. 2008;91:

4414–4423. doi:10.3168/jds.2007-0980

62.　Liu X, Huang M, Fan B, Buckler ES, Zhang Z. Iterative Usage of Fixed and Random Effect

Models for Powerful and Efficient Genome-Wide Association Studies. Listgarten J, editor.

PLOS Genet. 2016;12: e1005767. doi:10.1371/journal.pgen.1005767

63.　Huang M, Liu X, Zhou Y, Summers RM, Zhang Z. BLINK: a package for the next level of

genome-wide association studies with both individuals and markers in the millions.

Gigascience. 2019;8. doi:10.1093/gigascience/giy154

64.　Chao J, Li Z, Sun Y, Aluko OO, Wu X, Wang Q, et al. MG2C: a user-friendly online tool for

drawing genetic maps. Mol Hortic. 2021;1: 16. doi:10.1186/s43897-021-00020-x

65.　Velt A, Frommer B, Blanc S, Holtgräwe D, Duchêne É, Dumas V, et al. An improved

reference of the grapevine genome reasserts the origin of the PN40024 highly homozygous

genotype. Whiteman N, editor. G3 Genes|Genomes|Genetics. 2023.

doi:10.1093/g3journal/jkad067

66.　Skinner ME, Uzilov A V., Stein LD, Mungall CJ, Holmes IH. JBrowse: A next-generation

genome browser. Genome Res. 2009;19: 1630–1638. doi:10.1101/gr.094607.109

67.　Myles S, Chia J-M, Hurwitz B, Simon C, Zhong GY, Buckler E, et al. Rapid Genomic

Characterization of the Genus Vitis. Ingvarsson PK, editor. PLoS One. 2010;5: e8219.

doi:10.1371/journal.pone.0008219

68.　Zhang C, Cui L, Fang J. Genome-wide association study of the candidate genes for grape

berry shape-related traits. BMC Plant Biol. 2022;22: 42. doi:10.1186/s12870-022-03434-x

69.   Navarro-Payá D, Santiago A, Orduña L, Zhang C, Amato A, D'Inca E, et al. The Grape Gene

Reference Catalogue as a Standard Resource for Gene Selection and Genetic Improvement.

Front Plant Sci. 2022;12. doi:10.3389/fpls.2021.803977

70.   UniProtConsortium. UniProt: a worldwide hub of protein knowledge. Nucleic Acids Res.

2019;47: D506–D515. doi:10.1093/nar/gky1049

71.   Enyew M, Feyissa T, Carlsson AS, Tesfaye K, Hammenhag C, Seyoum A, et al. Genome-

wide analyses using multi-locus models revealed marker-trait associations for major

agronomic traits in Sorghum bicolor. Front Plant Sci. 2022;13: 1–19.

doi:10.3389/fpls.2022.999692

72.   Zecca G, Labra M, Grassi F. Untangling the Evolution of American Wild Grapes: Admixed

Species and How to Find Them. Front Plant Sci. 2020;10: 1–17. doi:10.3389/fpls.2019.01814

73.   Riaz S, Tenscher AC, Heinitz CC, Huerta-Acosta KG, Andrew Walker M. Genetic analysis

reveals an east-west divide within North American Vitis species that mirrors their resistance to

Pierce's disease. PLoS One. 2020;15: 7–10. doi:10.1371/journal.pone.0243445

74.   Cipriani G, Spadotto A, Jurman I, Gaspero G Di, Crespan M, Meneghetti S, et al. The SSR-

based molecular profile of 1005 grapevine (Vitis vinifera L.) accessions uncovers new

synonymy and parentages, and reveals a large admixture amongst varieties of different

geographic origin. Theor Appl Genet. 2010;121: 1569–1585. doi:10.1007/s00122-010-1411-9

75.   Rodrigues CS, Moura MF, de Oliveira GL, da Silva MJR, Tecchio MA. Best linear unbiased

prediction in combination with path analysis in processing grapes. Sci Agric. 2024;81.

doi:10.1590/1678-992X-2022-0218

76.   Abiri K, Rezaei M, Tahanian H, Heidari P, Khadivi A. Morphological and pomological

variability of a grape (Vitis vinifera L.) germplasm collection. Sci Hortic (Amsterdam).

2020;266: 109285. doi:10.1016/j.scienta.2020.109285

77.   Wu Y, Wang Y, Fan X, Zhang Y, Jiang J, Sun L, et al. QTL mapping for berry shape based on

a high-density genetic map constructed by whole-genome resequencing in grape. Hortic Plant

J. 2023;9: 729–742. doi:10.1016/j.hpj.2022.11.005

78.   Muñoz-Espinoza C, Di Genova A, Sánchez A, Correa J, Espinoza A, Meneses C, et al.

Identification of SNPs and InDels associated with berry size in table grapes integrating genetic

and transcriptomic approaches. BMC Plant Biol. 2020;20: 1–21. doi:10.1186/s12870-020-

02564-4

79.   Mejía N, Soto B, Guerrero M, Casanueva X, Houel C, de los Ángeles Miccono M, et al.

Molecular, genetic and transcriptional evidence for a role of VvAGL11 in stenospermocarpic

seedlessness in grapevine. BMC Plant Biol. 2011;11: 57. doi:10.1186/1471-2229-11-57

80.   Chialva C, Eichler E, Grissi C, Muñoz C, Gomez-Talquenca S, Martínez-Zapater JM, et al.

Expression of grapevine AINTEGUMENTA-like genes is associated with variation in ovary

and berry size. Plant Mol Biol. 2016;91: 67–80. doi:10.1007/s11103-016-0443-1

81.   Al-Saif AM, Fahmy MA, Baghdady GA, El-Razik AMA, Kabsha EA, Farouk MH, et al. The

Impact of Bud Load on Berry Quality, Yield, and Cluster Compactness in H4 Strain

Grapevines. Agronomy. 2023;13: 2431. doi:10.3390/agronomy13092431

82.   Pedro Júnior MJ, Hernandes JL, Bardin-Camparotto L, Blain GC. Plant parameters and must

composition of 'Syrah' grapevine cultivated under sequential summer and winter growing

seasons. Bragantia. 2017;76: 345–351. doi:10.1590/1678-4499.146

83.   Vafaee Y, Ghaderi N, Khadivi A. Morphological variation and marker-fruit trait associations

in a collection of grape (Vitis vinifera L.). Sci Hortic (Amsterdam). 2017;225: 771–782.

doi:10.1016/j.scienta.2017.08.007

84.   Khadivi-Khub A, Salimpour A, Rasouli M. Analysis of grape germplasm from Iran based on

fruit characteristics. Rev Bras Bot. 2014;37: 105–113. doi:10.1007/s40415-014-0054-5

85.   Azzi L, Deluche C, Gévaudant F, Frangne N, Delmas F, Hernould M, et al. Fruit growth-

related genes in tomato. J Exp Bot. 2015;66: 1075–1086. doi:10.1093/jxb/eru527

86.   Zhao X, Muhammad N, Zhao Z, Yin K, Liu Z, Wang L, et al. Molecular regulation of fruit

1051   size in horticultural plants: A review. Sci Hortic (Amsterdam). 2021;288.

1052   doi:10.1016/j.scienta.2021.110353

1053   87.   Hussain Q, Shi J, Scheben A, Zhan J, Wang X, Liu G, et al. Genetic and signalling pathways

1054   of dry fruit size: targets for genome editing☐based crop improvement. Plant Biotechnol J.

1055   2020;18: 1124–1140. doi:10.1111/pbi.13318

1056   88.   Zhao J, Li Y, Ding L, Yan S, Liu M, Jiang L, et al. Phloem transcriptome signatures underpin

1057   the physiological differentiation of the pedicel, stalk and fruit of cucumber (Cucumis sativus

1058   L.). Plant Cell Physiol. 2016;57: 19–34. doi:10.1093/pcp/pcv168

1059   89.   Pattison RJ, Csukasi F, Catalá C. Mechanisms regulating auxin action during fruit

1060   development. Physiol Plant. 2014;151: 62–72. doi:10.1111/ppl.12142

1061   90.   Liao X, Li M, Liu B, Yan M, Yu X, Zi H, et al. Interlinked regulatory loops of ABA

1062   catabolism and biosynthesis coordinate fruit growth and ripening in woodland strawberry.

1063   Proc Natl Acad Sci. 2018;115. doi:10.1073/pnas.1812575115

1064   91.   de Jong M, Mariani C, Vriezen WH. The role of auxin and gibberellin in tomato fruit set. J

1065   Exp Bot. 2009;60: 1523–1532. doi:10.1093/jxb/erp094

1066   92.   Godoy F, Kühn N, Muñoz M, Marchandon G, Gouthu S, Deluc L, et al. The role of auxin

1067   during early berry development in grapevine as revealed by transcript profiling from

1068   pollination to fruit set. Hortic Res. 2021;8: 1–15. doi:10.1038/s41438-021-00568-1

1069   93.   Benková E, Michniewicz M, Sauer M, Teichmann T, Seifertová D, Jürgens G, et al. Local,

1070   Efflux-Dependent Auxin Gradients as a Common Module for Plant Organ Formation. Cell.

1071   2003;115: 591–602. doi:10.1016/S0092-8674(03)00924-3

1072   94.   Scarpella E, Marcos D, Friml J, Berleth T. Control of leaf vascular patterning by polar auxin

1073   transport. Genes Dev. 2006;20: 1015–1027. doi:10.1101/gad.1402406

95.   Korasick DA, Westfall CS, Lee SG, Nanao MH, Dumas R, Hagen G, et al. Molecular basis for

AUXIN RESPONSE FACTOR protein interaction and the control of auxin response

repression. Proc Natl Acad Sci. 2014;111: 5427–5432. doi:10.1073/pnas.1400074111

96.   Li S-B, Xie Z-Z, Hu C-G, Zhang J-Z. A Review of Auxin Response Factors (ARFs) in Plants.

Front Plant Sci. 2016;7. doi:10.3389/fpls.2016.00047

97.   Yan M, Yan Y, Wang P, Wang Y, Piao X, Di P, et al. Genome-Wide Identification and

Expression Analysis of Auxin Response Factor (ARF) Gene Family in Panax ginseng

Indicates Its Possible Roles in Root Development. Plants. 2023;12: 3943.

doi:10.3390/plants12233943

98.   Li L, Chen X. Auxin regulation on crop: from mechanisms to opportunities in soybean

breeding. Mol Breed. 2023;43: 16. doi:10.1007/s11032-023-01361-9

99.   Ravest G, Mamani M, Giacomelli L, Moser C, Pastenes C, Hinrichsen P. Bioactive

Gibberellins Show Differential Abundance at Key Phenological Stages for Berry Growth in

Table Grapes. Am J Enol Vitic. 2017;68: 478–484. doi:10.5344/ajev.2017.16036

100.   Suehiro Y, Mochida K, Tsuma M, Yasuda Y, Itamura H, Esumi T. Effects of Gibberellic

Acid/Cytokinin Treatments on Berry Development and Maturation in the Yellow-green

Skinned 'Shine Muscat' Grape. Hortic J. 2019;88: 202–213. doi:10.2503/hortj.UTD-046

101.   Casanova L, Casanova R, Moret A, Agustí M. The application of gibberellic acid increases

berry size of "Emperatriz" seedless grape. Spanish J Agric Res. 2009;7: 919–927.

doi:10.5424/sjar/2009074-1105

102.   Grimplet J, Agudelo-Romero P, Teixeira RT, Martinez-Zapater JM, Fortes AM. Structural and

Functional Analysis of the GRAS Gene Family in Grapevine Indicates a Role of GRAS

Proteins in the Control of Development and Stress Responses. Front Plant Sci. 2016;7.

doi:10.3389/fpls.2016.00353

103.   Heo J-O, Chang KS, Kim IA, Lee M-H, Lee SA, Song S-K, et al. Funneling of gibberellin

signaling by the GRAS transcription regulator SCARECROW-LIKE 3 in the Arabidopsis root.

Proc Natl Acad Sci. 2011;108: 2166–2171. doi:10.1073/pnas.1012215108

104. Wang T-T, Yu T-F, Fu J-D, Su H-G, Chen J, Zhou Y-B, et al. Genome-Wide Analysis of the GRAS Gene Family and Functional Identification of GmGRAS37 in Drought and Salt Tolerance. Front Plant Sci. 2020;11. doi:10.3389/fpls.2020.604690

105. Neves C, Ribeiro B, Amaro R, Expósito J, Grimplet J, Fortes AM. Network of GRAS transcription factors in plant development, fruit ripening and stress responses. Hortic Res. 2023;10. doi:10.1093/hr/uhad220

106. Song H, Fu X, Li J, Niu T, Shen J, Wang X, et al. Phylogenetic analysis and expression profiles of jasmonate ZIM-domain gene family provide insight into abiotic stress resistance in sunflower. Front Plant Sci. 2022;13. doi:10.3389/fpls.2022.1010404

107. Taniguchi S, Hosokawa☐Shinonaga Y, Tamaoki D, Yamada S, Akimitsu K, Gomi K. Jasmonate induction of the monoterpene linalool confers resistance to rice bacterial blight and its biosynthesis is regulated by <scp>JAZ</scp> protein in rice. Plant Cell Environ. 2014;37: 451–461. doi:10.1111/pce.12169

108. Singh AP, Pandey BK, Mehra P, Heitz T, Giri J. OsJAZ9 overexpression modulates jasmonic acid biosynthesis and potassium deficiency responses in rice. Plant Mol Biol. 2020;104: 397–410. doi:10.1007/s11103-020-01047-2

109. Peethambaran PK, Glenz R, Höninger S, Shahinul Islam SM, Hummel S, Harter K, et al. Salt-inducible expression of OsJAZ8 improves resilience against salt-stress. BMC Plant Biol. 2018;18: 311. doi:10.1186/s12870-018-1521-0

110. Lal MK, Tiwari RK, Altaf MA, Kumar A, Kumar R. Editorial: Abiotic and biotic stress in horticultural crops: insight into recent advances in the underlying tolerance mechanism. Front Plant Sci. 2023;14. doi:10.3389/fpls.2023.1212982

111. Zha Q, Xi X, He Y, Jiang A. Comprehensive evaluation of heat resistance in 68 Vitis Germ Plasm resources. Vitis - J Grapevine Res. 2018;57: 75–81. doi:10.5073/vitis.2018.57.75-81

112. Teslić N, Zinzani G, Parpinello GP, Versari A. Climate change trends, grape production, and

potential alcohol concentration in wine from the "Romagna Sangiovese" appellation area

(Italy). Theor Appl Climatol. 2018;131: 793–803. doi:10.1007/s00704-016-2005-5

113. Greer DH, Weston C. Heat stress affects flowering, berry growth, sugar accumulation and

photosynthesis of Vitis vinifera cv. Semillon grapevines grown in a controlled environment.

Funct Plant Biol. 2010;37: 206. doi:10.1071/FP09209

114. Liu G, Chai F, Wang Y, Jiang J, Duan W, Wang Y, et al. Genome-wide Identification and

Classification of HSF Family in Grape, and Their Transcriptional Analysis under Heat

Acclimation and Heat Stress. Hortic Plant J. 2018;4: 133–143. doi:10.1016/j.hpj.2018.06.001

115. DeYoung BJ, Innes RW. Plant NBS-LRR proteins in pathogen sensing and host defense. Nat

Immunol. 2006;7: 1243–9. doi:10.1038/ni1410

116. van Wersch S, Tian L, Hoy R, Li X. Plant NLRs: The Whistleblowers of Plant Immunity.

Plant Commun. 2020;1: 100016. doi:10.1016/j.xplc.2019.100016

117. Biswas D, Gain H, Mandal A. MYB transcription factor: A new weapon for biotic stress

tolerance in plants. Plant Stress. 2023;10: 100252. doi:10.1016/j.stress.2023.100252

118. Nuruzzaman M, Sharoni AM, Kikuchi S. Roles of NAC transcription factors in the regulation

of biotic and abiotic stress responses in plants. Front Microbiol. 2013;4.

doi:10.3389/fmicb.2013.00248

119. Fernandez L, Ageorges A, Torregrosa L. A putative NAP homolog specifically expressed

during grapevine flower and berry development. Vitis. 2006;45: 51–52.

doi:10.5073/vitis.2006.45.51-52

120. Muñoz-Espinoza C, Di Genova A, Correa J, Silva R, Maass A, González-Agüero M, et al.

Transcriptome profiling of grapevine seedless segregants during berry development reveals

candidate genes associated with berry weight. BMC Plant Biol. 2016;16: 1–17.

doi:10.1186/s12870-016-0789-1

121. Tello J, Torres-Pérez R, Grimplet J, Carbonell-Bejerano P, Martínez-Zapater JM, Ibáñez J.

Polymorphisms and minihaplotypes in the VvNAC26 gene associate with berry size variation

in grapevine. BMC Plant Biol. 2015;15: 253. doi:10.1186/s12870-015-0622-2

122. Ketudat Cairns JR, Esen A. β-Glucosidases. Cell Mol Life Sci. 2010;67: 3389–3405.

doi:10.1007/s00018-010-0399-2

123. Ketudat Cairns JR, Mahong B, Baiya S, Jeon J-S. β-Glucosidases: Multitasking, moonlighting

or simply misunderstood? Plant Sci. 2015;241: 246–259. doi:10.1016/j.plantsci.2015.10.014

124. Kong H, Song J, Ma S, Yang J, Shao Z, Li Q, et al. Genome-wide identification and

expression analysis of the glycosyl hydrolase family 1 genes in Medicago sativa revealed their

potential roles in response to multiple abiotic stresses. BMC Genomics. 2024;25: 20.

doi:10.1186/s12864-023-09918-w

125. Morant AV, Bjarnholt N, Kragh ME, Kjærgaard CH, Jørgensen K, Paquette SM, et al. The β -

Glucosidases Responsible for Bioactivation of Hydroxynitrile Glucosides in Lotus japonicus.

Plant Physiol. 2008;147: 1072–1091. doi:10.1104/pp.107.109512

126. Baba SA, Vishwakarma RA, Ashraf N. Functional Characterization of CsBGlu12, a β-

Glucosidase from Crocus sativus, Provides Insights into Its Role in Abiotic Stress through

Accumulation of Antioxidant Flavonols. J Biol Chem. 2017;292: 4700–4713.

doi:10.1074/jbc.M116.762161

127. Wang X, Yan L, Wang B, Qian Y, Wang Z, Wu W. Comparative Proteomic Analysis of

Grapevine Rootstock in Response to Waterlogging Stress. Front Plant Sci. 2021;12.

doi:10.3389/fpls.2021.749184

128. Han X, Li Y-H, Yao M-H, Yao F, Wang Z-L, Wang H, et al. Transcriptomics Reveals the

Effect of Short-Term Freezing on the Signal Transduction and Metabolism of Grapevine. Int J

Mol Sci. 2023;24: 3884. doi:10.3390/ijms24043884

129. Griesser M, Savoi S, Supapvanich S, Dobrev P, Vankova R, Forneck A. Phytohormone

profiles are strongly altered during induction and symptom development of the physiological

ripening disorder berry shrivel in grapevine. Plant Mol Biol. 2020;103: 141–157.

doi:10.1007/s11103-020-00980-6

130. Li M, Feng F, Cheng L. Expression Patterns of Genes Involved in Sugar Metabolism and

Accumulation during Apple Fruit Development. Börnke F, editor. PLoS One. 2012;7: e33055.

doi:10.1371/journal.pone.0033055

131. Cao S, Yang Z, Zheng Y. Sugar metabolism in relation to chilling tolerance of loquat fruit.

Food Chem. 2013;136: 139–143. doi:10.1016/j.foodchem.2012.07.113

132. Du Z, Su Q, Wu Z, Huang Z, Bao J, Li J, et al. Genome-wide characterization of MATE gene

family and expression profiles in response to abiotic stresses in rice (Oryza sativa). BMC Ecol

Evol. 2021;21: 141. doi:10.1186/s12862-021-01873-y

133. Watanabe M, Otagaki S, Matsumoto S, Shiratake K. Genome-Wide Analysis of Multidrug and

Toxic Compound Extrustion Transporters in Grape. Front Plant Sci. 2022;13.

doi:10.3389/fpls.2022.892638

134. Gollhofer J, Timofeev R, Lan P, Schmidt W, Buckhout TJ. Vacuolar-Iron-Transporter1-Like

Proteins Mediate Iron Homeostasis in Arabidopsis. Witt SN, editor. PLoS One. 2014;9:

e110468. doi:10.1371/journal.pone.0110468

135. Kato T, Kumazaki K, Wada M, Taniguchi R, Nakane T, Yamashita K, et al. Crystal structure

of plant vacuolar iron transporter VIT1. Nat Plants. 2019;5: 308–315. doi:10.1038/s41477-

019-0367-2

136. Yanagisawa M, Zhang C, Szymanski DB. ARP2/3-dependent growth in the plant kingdom:

SCARs for life. Front Plant Sci. 2013;4. doi:10.3389/fpls.2013.00166

137. Harries PA, Pan A, Quatrano RS. Actin-Related Protein2/3 Complex Component ARPC1 Is

Required for Proper Cell Morphogenesis and Polarized Cell Growth in Physcomitrella patens.

Plant Cell. 2005;17: 2327–2339. doi:10.1105/tpc.105.033266

138.   Mathur J, Mathur N, Kernebeck B, Hülskamp M. Mutations in Actin-Related Proteins 2 and 3

1201   Affect Cell Shape Development in Arabidopsis. Plant Cell. 2003;15: 1632–1645.

1202   doi:10.1105/tpc.011676

1203   139.   Pratap Sahi V, Cifrová P, García-González J, Kotannal Baby I, Mouillé G, Gineau E, et al.

1204   Arabidopsis thaliana plants lacking the ARP2/3 complex show defects in cell wall assembly

1205   and auxin distribution. Ann Bot. 2017. doi:10.1093/aob/mcx178

1206   140.   Jiang K, Sorefan K, Deeks MJ, Bevan MW, Hussey PJ, Hetherington AM. The ARP2/3

1207   Complex Mediates Guard Cell Actin Reorganization and Stomatal Movement in Arabidopsis.

1208   Plant Cell. 2012;24: 2031–2040. doi:10.1105/tpc.112.096263

1209   141.   Yanagisawa M, Desyatova AS, Belteton SA, Mallery EL, Turner JA, Szymanski DB.

1210   Patterning mechanisms of cytoskeletal and cell wall systems during leaf trichome

1211   morphogenesis. Nat Plants. 2015;1: 15014. doi:10.1038/nplants.2015.14

1212   142.   García-González J, Kebrlová Š, Semerák M, Lacek J, Kotannal Baby I, Petrášek J, et al.

1213   Arp2/3 Complex Is Required for Auxin-Driven Cell Expansion Through Regulation of Auxin

1214   Transporter Homeostasis. Front Plant Sci. 2020;11. doi:10.3389/fpls.2020.00486

1215   143.   Garrido C, Wollman F-A, Lafontaine I. The Evolutionary History of Peptidases Involved in

1216   the Processing of Organelle-Targeting Peptides. Angert E, editor. Genome Biol Evol. 2022;14.

1217   doi:10.1093/gbe/evac101

1218   144.   Trösch R, Jarvis P. The Stromal Processing Peptidase of Chloroplasts is Essential in

1219   Arabidopsis, with Knockout Mutations Causing Embryo Arrest after the 16-Cell Stage.

1220   Tsiantis M, editor. PLoS One. 2011;6: e23039. doi:10.1371/journal.pone.0023039

1221   145.   Kmiec B, Teixeira PF, Glaser E. Shredding the signal: targeting peptide degradation in

1222   mitochondria and chloroplasts. Trends Plant Sci. 2014;19: 771–778.

1223   doi:10.1016/j.tplants.2014.09.004

146.   Tovar☐Méndez A, Miernyk JA, Randall DD. Regulation of pyruvate dehydrogenase complex

activity in plant cells. Eur J Biochem. 2003;270: 1043–1049. doi:10.1046/j.1432-

1033.2003.03469.x

147. Ke J, Behal RH, Back SL, Nikolau BJ, Wurtele ES, Oliver DJ. The Role of Pyruvate

Dehydrogenase and Acetyl-Coenzyme A Synthetase in Fatty Acid Synthesis in Developing

Arabidopsis Seeds. Plant Physiol. 2000;123: 497–508. doi:10.1104/pp.123.2.497

148. Tello J, Torres-Pérez R, Grimplet J, Ibáñez J. Association analysis of grapevine bunch traits

using a comprehensive approach. Theor Appl Genet. 2016;129: 227–242. doi:10.1007/s00122-

015-2623-9

149. Tello J, Aguirrezábal R, Hernáiz S, Larreina B, Montemayor MI, Vaquero E, et al.

Multicultivar and multivariate study of the natural variation for grapevine bunch compactness.

Aust J Grape Wine Res. 2015;21: 277–289. doi:10.1111/ajgw.12121

150. Chao H, He J, Cai Q, Zhao W, Fu H, Hua Y, et al. The Expression Characteristics of NPF

Genes and Their Response to Vernalization and Nitrogen Deficiency in Rapeseed. Int J Mol

Sci. 2021;22: 4944. doi:10.3390/ijms22094944

151. Wu W, Fu P, Lu J. Grapevine WRKY transcription factors. Fruit Res. 2022;2: 1–8.

doi:10.48130/FruRes-2022-0010

152. Vannozzi A, Wong DCJ, Hï¿½ll J, Hmmam I, Matus JT, Bogs J, et al. Combinatorial

Regulation of Stilbene Synthase Genes by WRKY and MYB Transcription Factors in

Grapevine (Vitis vinifera L.). Plant Cell Physiol. 2018;59: 1043–1059.

doi:10.1093/pcp/pcy045

153. Valletta A, Iozia LM, Leonelli F. Impact of Environmental Factors on Stilbene Biosynthesis.

Plants. 2021;10: 90. doi:10.3390/plants10010090

154. Karasov TL, Chae E, Herman JJ, Bergelson J. Mechanisms to Mitigate the Trade-Off between

Growth and Defense. Plant Cell. 2017;29: 666–680. doi:10.1105/tpc.16.00931

155. Wang H, Huang T, Ruan Z, Tu R, He Y, Liu Q, et al. A MACPF Protein OsCAD1 Balances

Plant Growth and Immunity Through Regulating Salicylic Acid Homeostasis in Rice. Plant Cell Environ. 2025. doi:10.1111/pce.15366

156. Zhang X, Dai Y-S, Wang Y-X, Su Z-Z, Yu L-J, Zhang Z-F, et al. Overexpression of the Arabidopsis MACPF Protein AtMACP2 Promotes Pathogen Resistance by Activating SA Signaling. Int J Mol Sci. 2022;23: 8784. doi:10.3390/ijms23158784

157. Yu L, Liu D, Chen S, Dai Y, Guo W, Zhang X, et al. Evolution and Expression of the Membrane Attack Complex and Perforin Gene Family in the Poaceae. Int J Mol Sci. 2020;21: 5736. doi:10.3390/ijms21165736

158. Chen P, Jian H, Wei F, Gu L, Hu T, Lv X, et al. Phylogenetic Analysis of the Membrane Attack Complex/Perforin Domain-Containing Proteins in Gossypium and the Role of GhMACPF26 in Cotton Under Cold Stress. Front Plant Sci. 2021;12. doi:10.3389/fpls.2021.684227

159. Akter N, Islam MSU, Rahman MS, Zohra FT, Rahman SM, Manirujjaman M, et al. Genome-wide identification and characterization of protein phosphatase 2C (PP2C) gene family in sunflower (Helianthus annuus L.) and their expression profiles in response to multiple abiotic stresses. Verma SK, editor. PLoS One. 2024;19: e0298543. doi:10.1371/journal.pone.0298543

160. Boneh U, Biton I, Schwartz A, Ben-Ari G. Characterization of the ABA signal transduction pathway in Vitis vinifera. Plant Sci. 2012;187: 89–96. doi:10.1016/j.plantsci.2012.01.015

161. Han G, Qiao Z, Li Y, Wang C, Wang B. The Roles of CCCH Zinc-Finger Proteins in Plant Abiotic Stress Tolerance. Int J Mol Sci. 2021;22: 8327. doi:10.3390/ijms22158327

162. Wang X, Zhong Y, Cheng Z (Max). Evolution and Expression Analysis of the CCCH Zinc Finger Gene Family in Vitis vinifera. Plant Genome. 2014;7. doi:10.3835/plantgenome2014.05.0019

163. Chen Y, Sun A, Wang M, Zhu Z, Ouwerkerk PBF. Functions of the CCCH type zinc finger protein OsGZF1 in regulation of the seed storage protein GluB-1 from rice. Plant Mol Biol.

1275   2014;84: 621–634. doi:10.1007/s11103-013-0158-5

1276   164.   Yan Z, Jia J, Yan X, Shi H, Han Y. Arabidopsis KHZ1 and KHZ2, two novel non-tandem

1277   CCCH zinc-finger and K-homolog domain proteins, have redundant roles in the regulation of

1278   flowering and senescence. Plant Mol Biol. 2017;95: 549–565. doi:10.1007/s11103-017-0667-8

1279   165.   Wang L, Xu Y, Zhang C, Ma Q, Joo S-H, Kim S-K, et al. OsLIC, a Novel CCCH-Type Zinc

1280   Finger Protein with Transcription Activation, Mediates Rice Architecture via Brassinosteroids

1281   Signaling. Ausubel FM, editor. PLoS One. 2008;3: e3521. doi:10.1371/journal.pone.0003521

1282   166.   Zhang D, Xu Z, Cao S, Chen K, Li S, Liu X, et al. An Uncanonical CCCH-Tandem Zinc-

1283   Finger Protein Represses Secondary Wall Synthesis and Controls Mechanical Strength in Rice.

1284   Mol Plant. 2018;11: 163–174. doi:10.1016/j.molp.2017.11.004

1285   167.   Raabe K, Honys D, Michailidis C. The role of eukaryotic initiation factor 3 in plant translation

1286   regulation. Plant Physiol Biochem. 2019;145: 75–83. doi:10.1016/j.plaphy.2019.10.015

1287   168.   Marintchev A, Ito T. eIF2B and the Integrated Stress Response: A Structural and Mechanistic

1288   View. Biochemistry. 2020;59: 1299–1308. doi:10.1021/acs.biochem.0c00132

1289   169.   Browning KS. Plant translation initiation factors: it is not easy to be green. Biochem Soc

1290   Trans. 2004;32: 589–591. doi:10.1042/BST0320589

1291   170.   Hinnebusch AG. Molecular Mechanism of Scanning and Start Codon Selection in Eukaryotes.

1292   Microbiol Mol Biol Rev. 2011;75: 434–467. doi:10.1128/MMBR.00008-11

1293   171.   Dong Q, Mao K, Duan D, Zhao S, Wang Y, Wang Q, et al. Genome-wide analyses of genes

1294   encoding FK506-binding proteins reveal their involvement in abiotic stress responses in apple.

1295   BMC Genomics. 2018;19: 707. doi:10.1186/s12864-018-5097-8

1296   172.   Aghdasi B, Ye K, Resnick A, Huang A, Ha HC, Guo X, et al. FKBP12, the 12-kDa FK506-

1297   binding protein, is a physiologic regulator of the cell cycle. Proc Natl Acad Sci. 2001;98:

1298   2425–2430. doi:10.1073/pnas.041614198

1299   173.   Vespa L, Vachon G, Berger F, Perazza D, Faure J-D, Herzog M. The Immunophilin-

Interacting Protein AtFIP37 from Arabidopsis Is Essential for Plant Development and Is

Involved in Trichome Endoreduplication. Plant Physiol. 2004;134: 1283–1292.

doi:10.1104/pp.103.028050

174. Yu Y, Li Y, Huang G, Meng Z, Zhang D, Wei J, et al. PwHAP5, a CCAAT-binding

transcription factor, interacts with PwFKBP12 and plays a role in pollen tube growth

orientation in Picea wilsonii. J Exp Bot. 2011;62: 4805–4817. doi:10.1093/jxb/err120

175. Castellano MM, del Pozo JC, Ramirez-Parra E, Brown S, Gutierrez C. Expression and

Stability of Arabidopsis CDC6 Are Associated with Endoreplication. Plant Cell. 2001;13:

2671–2686. doi:10.1105/tpc.010329

176. Shiri Y, Solouki M, Ebrahimie E, Emamjomeh A, Zahiri J. Unraveling the transcriptional

complexity of compactness in sistan grape cluster. Plant Sci. 2018;270: 198–208.

doi:10.1016/j.plantsci.2018.02.011

177. Díaz-Riquelme J, Martínez-Zapater JM, Carmona MJ. Transcriptional Analysis of Tendril and

Inflorescence Development in Grapevine (Vitis vinifera L.). Blazquez MA, editor. PLoS One.

2014;9: e92339. doi:10.1371/journal.pone.0092339

178. Lebon G, Wojnarowiez G, Holzapfel B, Fontaine F, Vaillant-Gaveau N, Clement C. Sugars

and flowering in the grapevine (Vitis vinifera L.). J Exp Bot. 2008;59: 2565–2578.

doi:10.1093/jxb/ern135

179. Reuscher S, Akiyama M, Yasuda T, Makino H, Aoki K, Shibata D, et al. The Sugar

Transporter Inventory of Tomato: Genome-Wide Identification and Expression Analysis. Plant

Cell Physiol. 2014;55: 1123–1141. doi:10.1093/pcp/pcu052

180. Akbar S, Hua X, Zhang Y, Liu G, Wang T, Shi H, et al. Genome-wide analysis of sugar

transporter gene family in Erianthus rufipilus and Saccharum officinarum, expression profiling

and identification of transcription factors. Front Plant Sci. 2025;15.

doi:10.3389/fpls.2024.1502649

181. Afoufa-Bastien D, Medici A, Jeauffre J, Coutos-Thévenot P, Lemoine R, Atanassova R, et al.

The Vitis vinifera sugar transporter gene family: phylogenetic overview and macroarray

expression profiling. BMC Plant Biol. 2010;10: 245. doi:10.1186/1471-2229-10-245

182. Iocco-Corena P, Chaïb J, Torregrosa L, Mackenzie D, Thomas MR, Smith HM. VviPLATZ1

is a major factor that controls female flower morphology determination in grapevine. Nat

Commun. 2021;12: 6995. doi:10.1038/s41467-021-27259-8

183. Zhang H, Liu L, Li Z, Wang S, Huang L, Lin S. PLATZ transcription factors and their

emerging roles in plant responses to environmental stresses. Plant Sci. 2025;352: 112400.

doi:10.1016/j.plantsci.2025.112400

184. Han X, Rong H, Tian Y, Qu Y, Xu M, Xu L. Genome-Wide Identification of PLATZ

Transcription Factors in *Ginkgo biloba* L. and Their Expression Characteristics During Seed

Development. Front Plant Sci. 2022;13. doi:10.3389/fpls.2022.946194

185. Baucher M, Guérin C, El Jaziri M, Behr M. Integration of the Plant-Specific PLATZ

Transcription Factors into Gene Regulatory Networks Controlling Developmental Processes.

CRC Crit Rev Plant Sci. 2024;43: 376–393. doi:10.1080/07352689.2024.2388013

186. Wang A, Hou Q, Si L, Huang X, Luo J, Lu D, et al. The PLATZ Transcription Factor GL6

Affects Grain Length and Number in Rice. Plant Physiol. 2019;180: 2077–2090.

doi:10.1104/pp.18.01574

187. Li Y, Wei K. Comparative functional genomics analysis of cytochrome P450 gene superfamily

in wheat and maize. BMC Plant Biol. 2020;20: 93. doi:10.1186/s12870-020-2288-7

188. Sotelo-Silveira M, Cucinotta M, Chauvin A-L, Chávez Montes RA, Colombo L, Marsch-

Martínez N, et al. Cytochrome P450 CYP78A9 Is Involved in Arabidopsis Reproductive

Development. Plant Physiol. 2013;162: 779–799. doi:10.1104/pp.113.218214

189. Fang W, Wang Z, Cui R, Li J, Li Y. Maternal control of seed size by EOD3/CYP78A6 in

Arabidopsis thaliana. Plant J. 2012;70: 929–939. doi:10.1111/j.1365-313X.2012.04907.x

190. Ma M, Wang Q, Li Z, Cheng H, Li Z, Liu X, et al. Expression of Ta CYP 78A3 , a gene

encoding cytochrome P450 CYP 78A3 protein in wheat (*Triticum aestivum* L.), affects seed

size. Plant J. 2015;83: 312–325. doi:10.1111/tpj.12896

191. Sun X, Cahill J, Van Hautegem T, Feys K, Whipple C, Novák O, et al. Altered expression of

maize PLASTOCHRON1 enhances biomass and seed yield by extending cell division

duration. Nat Commun. 2017;8: 14752. doi:10.1038/ncomms14752

192. Qi X, Liu C, Song L, Li M. Arabidopsis EOD3 homologue PaCYP78A6 affects fruit size and

is involved in sweet cherry (Prunus avium L.) fruit ripening. Sci Hortic (Amsterdam).

2019;246: 57–67. doi:10.1016/j.scienta.2018.10.041

193. Allayee H, Farber CR, Seldin MM, Williams EG, James DE, Lusis AJ. Systems genetics

approaches for understanding complex traits with relevance for human disease. Elife. 2023;12.

doi:10.7554/eLife.91004

1362

1363 Supporting information

1364 **S1 Fig. Geographical location of the IAC grapevine germplasm.** The three experimental fields

1365 that comprise the germplasm are highlighted in the figure with rectangles and labeled A, B, and C.

1366

1367 **S2 Fig. Box plots showing the distribution of raw phenotypic data for each trait and year**. BL,

1368 berry length; BWi, berry width; BW, berry weight; CL, cluster length; CWi, cluster width; CW,

1369 cluster weight.

1370

1371 **S3 Fig. Pearson correlation coefficients and frequency distribution histograms of BLUEs for**

1372 **different berry and cluster traits evaluated**. Significance: ** $p \leq 0.01$, *** $p \leq 0.001$.

1373   Abbreviations: BL, berry length; BWi, berry width; BW, berry weight; CL, cluster length; CWi,

1374   cluster width; CW, cluster weight.

1375

1376   **S1 Table. Genotype codes and passport data for 288 grapevine accessions used in the GWAS.**

1377

1378   **S2 Table. Best linear unbiased estimators (BLUEs) for six traits evaluated over 12 seasons.** The

1379   traits included berry length (BL), berry width (BWi), berry weight (BW), cluster length (CL), cluster

1380   width (CWi), and cluster weight (CW).

1381

S3 Table. Estimates of variance components and heritability (H²) for six evaluated traits,

obtained from mixed model analysis considering different model terms. ModelTerm indicates the

factor or interaction included in the model; VarianceComponent corresponds to the estimated

variance; SE is the standard error of the estimate; Zratio is the ratio between the estimate and its

standard error, used as an indicator of the relative magnitude of the component; and H² represents the

broad-sense heritability of the trait.

1388

1389   **S4 Table. SNP profiles of 288 grapevine accessions from the IAC grapevine germplasm,**

1390   **genotyped at 11,115 high-quality SNPs used for GWAS analysis**. "AA": homozygous for the

1391   dominant allele; "AB": heterozygous; "BB": homozygous for the recessive allele; "NA": missing

1392   data.

1393

1394   **S5 Table. List of significant markers associated with berry and cluster-related traits from a**

1395   **genome-wide association analysis of 288 grapevine accessions using 11,115 SNPs whose MAFs**

1396   **were greater than 0.05.**

1397

1398 **S6 Table. Details of the genes closest to significant marker–trait associations, including their**

1399 **physical positions, annotated functions, and GO terms.**