Discrete interneuron subsets participate in GluN1/GluN3A excitatory glycine receptor (eGlyR)-mediated regulation of hippocampal network activity throughout development and evolution.

June Hoan Kim[1], Anna Vlachos[1], Vivek Mahadevan[1], Adam P Caccavano[1], Tue Banke[4], Oliver C Crawley[2], Ana I Navarro[2], Xiaoqing Yuan[1], Daniel Abebe[1], Steven Hunt[1], Geoffrey A Vargish[1], Ramesh Chittajallu[1], Mark A G Eldridge[3,8], Reza Azadi[3], Alex C Cummins[3], Anne-Claire Tangen[3], Peyton Harmon[3], Anya Plotnikova[3], Arya Mohanty[3], Elisabetta Furlanis[6,7], Yating Wang[6,7], Min Dai[6,7], Brenda Leyva Garcia[6,7], Ding Liu[4], Zongjian Zhu[4], Hongjie Yuan[4], Samantha L Summer[5], Matthew P Epplin[5], Dennis C Liotta[5], James Pickel[3], Bruno B Averbeck[3], Isabel Perez-Otaño[2], Jordane Dimidschstein[7], Gord Fishell[6,7], Stephen F Traynelis[4], Chris J McBain[1], Kenneth A Pelkey[1]

[1] Section on Cellular and Synaptic Physiology, Eunice Kennedy Shriver National Institute of Child and Human Development, National Institutes of Health, Bethesda, MD 20892, USA
[2] Instituto de Neurociencias (CSIC-UMH), San Juan de Alicante 03550, Spain
[3] Laboratory of Neuropsychology, National Institute of Mental Health (NIMH), NIH, Bethesda, MD, 20892 USA
[4] Department of Pharmacology and Chemical Biology, Emory University School of Medicine, Atlanta, GA, 30322, USA
[5] Department of Chemistry, Emory University, Atlanta, GA, 30322, USA
[6] Department of Neurobiology, Harvard Medical School, Boston, MA, USA
[7] Stanley Center for Psychiatric Research, Broad Institute of Harvard and MIT, Cambridge, MA, USA
[8] Biosciences Institute, Newcastle University, Newcastle upon Tyne NE1 7RU, United Kingdom
Corresponding authors email: pelkeyk2@nih.gov (KAP), mcbainc@nih.gov (CJM)

Summary

Decades of studies implicating GluN3A *N*-methyl-D-aspartate receptor (NMDAR) subunits in physiological and pathological function have largely been interpreted through direct regulation of conventional glutamatergic NMDARs. However, emerging evidence indicates that GluN3A frequently assembles with GluN1 forming unconventional glutamate-insensitive NMDARs that operate as native excitatory glycine receptors (eGlyRs). Here we demonstrate that hippocampal somatostatin and neurogliaform interneurons (Sst-INs and NGFCs) express functional eGlyRs from early postnatal through adult ages. In the developing hippocampus eGlyR-mediated excitation of NGFCs with ambient glycine dramatically increases GABAergic tone, with consequences for the generation of giant depolarizing potentials (GDPs). In the mature hippocampus, eGlyR-mediated excitation of Sst-INs regulates sharp wave ripples (SWRs). Finally, we reveal evolutionary conservation of hippocampal Sst-IN eGlyRs and eGlyR-mediated SWR regulation in non-human primates confirming functional eGlyR availability for therapeutic potential in higher species. Our findings underscore that eGlyR mediated regulation of cell and circuit excitability through both cell autonomous and cell non-autonomous mechanisms must be considered to understand GluN3A roles in brain development, plasticity, and disease.

Introduction

For almost half a century NMDARs have been recognized as crucial mediators of excitatory glutamatergic synapse development, function, and plasticity in brain health and disease. The coincidence detecting features, kinetics, and calcium permeability of conventional GluN1/GluN2 subunit-containing NMDARs confer integrative properties uniquely suited to controlling synaptic maturation, integration, and activity-dependent plasticity [Hansen2021, Nicoll2017]. Interestingly, GluN3A NMDAR subunits are also strongly implicated in nascent synapse and circuit development/plasticity [Das1998, Gonzalez-Gonzalez2023, Hurley2024, Sasaki2002, Tong2008, Roberts2009, Henson2012, Mohamad2013, Kehoe2014, Perez-Otano2016, Crawley2022]. However, emerging evidence indicates that GluN3A subunits primarily co-assemble with GluN1 to yield unconventional glycine (Gly) gated "non-Hebbian NMDARs" that are glutamate insensitive and voltage-independent with low Ca^{2+} permeability [Chatterton2002, Pina-Crespo2010, Awobuluyi2007, Madry2007, Kvist2013a, Bossi2022]. As such GluN1/GluN3 NMDARs are not members of the glutamate receptor family and are referred to as eGlyRs to differentiate them from both conventional glutamate/glycine gated NMDARs and inhibitory chloride conducting pentameric glycine-gated receptors [Chatterton2002, Otsu2019, Crawley2022, Bossi2022, Hurley2024]. Curiously, eGlyRs are often expressed by neurons in forebrain regions essentially devoid of dedicated glycinergic afferents [Zeilhofer2005, Bossi2022, Bossi2023]. Thus, evidence for prototypical rapid, phasic communication through eGlyRs is lacking. Instead, evidence suggests that eGlyRs regulate neuronal excitability by sensing ambient levels of extracellular Gly consistent with observations that surface GluN3A subunits comprise a highly diffusive receptor pool that does not strongly localize to postsynaptic sites [Perez-Otano2006, Otsu2019, Bossi2022, Gonzalez-Gonzalez2023].

Gly gates eGlyRs by binding to GluN3 subunits, whereas Gly binding to GluN1 profoundly desensitizes eGlyRs [Awobuluyi2007, Madry2007, Kvist2013a, Cummings2017, Grand2018, Zhu2020, Rouzbeh2023]. While saturating levels of ambient Gly for GluN1 have historically confounded interrogation of eGlyRs, findings that the glycine site antagonist CGP-78608 (CGP) blocks desensitization and awakens eGlyRs has fostered a new era of cellular and network investigation into this notoriously enigmatic receptor subpopulation that has remained elusive since their discovery in recombinant systems over 20 years ago [Chatterton2002, Matsuda2002, Yao2006, Madry2007, Grand2018, Otsu2019, Zhu2020, Bossi2022, Rouzbeh2023, Hurley2024]. Genetic deletion of *Grin3a*, has revealed critical roles for eGlyRs in regulating diverse behaviors including sensorimotor gating, sleep, place aversion conditioning and fear memory stability [Otsu2019, Bossi2022, Brody2005, Sunagawa2016, Crawley2022]. Moreover, mutations or alterations in *GRIN3A/Grin3a* expression in humans and mice are linked to diverse neurological conditions such as schizophrenia, bipolar disorder, addiction, epilepsy, and Huntington's disease making GluN3A an attractive candidate for therapeutic intervention [Mueller2004, Gallinat2007, Marco2013, Yuan2013, Yang2015, Lee2016, Huang2017, Marco2018, Crawley2022]. The unique features of eGlyR ligand binding, channel activation, and gating can be successfully exploited to develop selective allosteric modulators [Kvist2013a, Kvist2013b, Zhu2020, Rouzbeh2023, Michalski2024, Jacobs2025]. However, determination of therapeutic potential requires thorough characterization of eGlyR function within discrete receptor expressing cell-types across divergent circuits throughout development and across species.

Grin3a expression is particularly high in the developing hippocampus but is not ubiquitous
exhibiting unique regional and cell type specific patterns with divergent temporal profiles
[Sucher1995, Wong2002, Roberts2009, Kehoe2014, Murillo2021, Hurley2024]. At early postnatal
stages *Grin3a* is highly expressed in CA1 pyramidal neurons (PNs) throughout the entire
rostrocaudal axis then dramatically declines in dorsal CA1 in adults but remains elevated in ventral
CA1 pyramidal neurons [Murillo2021, Hurley2024]. *Grin3a* is not expressed in developing or adult
PNs of CA2, CA3, or dentate gyrus indicating that any potential role for *Grin3a* in regulating circuit
maturation and function in these hippocampal substructures requires cell non-autonomous
mechanisms [Sucher1995, Murillo2021, Beesley2022, Pizzamiglio2025]. Hippocampal and
neocortical medial ganglionic eminence (MGE)-derived inhibitory GABAergic interneurons
express conspicuously high levels of *Grin3a* throughout development, and adult somatostatin-
expressing interneurons (Sst-INs) exhibit robust eGlyR-mediated currents [Pfeffer2013, Paul2017,
Murillo2021, Bossi2022, Pizzamiglio2025]. Individual local circuit interneurons widely innervate
large numbers of nearby PNs to critically instruct nascent circuit connectivity through early
postnatal depolarizing GABAergic drive [Ben-Ari2002, Ben-Ari2007, Cossart2022, Pelkey2017].
Moreover, in mature hippocampus GABAergic inhibition from divergent interneuron subtypes
imposes discrete spatiotemporal profiles of inhibition to entrain principal cell assemblies and
coordinate network oscillations critical for cognition [Klausberger2008, Stark2014, Pelkey2017,
Tzilivaki2023]. These central roles of GABAergic interneurons could be leveraged by ambient Gly
through eGlyRs to control network excitability for circuit maturation and rhythmogenesis. Here
using a multiparametric approach including transcriptional profiling, conditional loss of function
mice, immunohistochemistry, and *ex vivo* electrophysiology with optogenetics and pharmacology
we demonstrate that hippocampal Sst-INs and neurogliaform interneurons (NGFCs) express
functional eGlyRs from early postnatal through adult ages. Activation of eGlyRs on these
interneuron subsets has the capacity to regulate the dominant spontaneous network pacing
activities, GDPs and SWRs, in developing and mature hippocampus respectively. Finally, we
demonstrate evolutionary conservation of hippocampal Sst-IN eGlyR expression and eGlyR-
mediated SWR regulation in the nonhuman primate (NHP) hippocampus confirming functional
availability of this target for therapeutic potential in higher mammalian species.
119

Results

eGlyR modulation of inhibitory drive and network synchronization in the early postnatal hippocampus.

Detailed spatiotemporal profiling using multiplexed fluorescence in situ hybridization (mFISH) revealed remarkably high *Grin3a* expression within MGE-derived hippocampal interneurons, particularly CA1 Sst-INs, at early postnatal ages (~postnatal day (P) 6-9) [Murillo2021]. To initiate our study, we employed Nkx2.1[Cre/+]:Rpl22(RiboTag)[HA/HA] (MGE-Ribotag) mice to independently validate and extend these findings. This Ribotag-based sequencing strategy provides a robust quantitative platform to examine mRNAs associated with the translational machinery for protein production selectively within the population of MGE-derived interneurons [Mahadevan2020, Sanz2009]. We observed an enrichment of *Grin3a* within hippocampal MGE-derived interneurons relative to bulk hippocampal tissue from early postnatal ages through adulthood (Figure 1A). Notably, bulk hippocampal *Grin3a* levels dramatically declined by 2 months of age, as expected given the well-established reduction in dorsal CA1 PNs [Ciabarra1995, Sucher1995, Sun1998, Sasaki2002, Zhang2002, Perez-Otano2016, Wee2016, Murillo2021, Crawley2022, Bossi2023]. However, MGE-derived interneurons retained high expression of *Grin3a* at all time points examined consistent with findings in adult neocortex [Pfeffer2013, Paul2017, Yao2021, Bossi2022]. Examination of dorsal hippocampal sections from a newly developed mClover-GluN3A reporter mouse line confirmed persistent GluN3A expression in a sparse population of cells outside the PN layer, prominently Sst+ neurons of stratum oriens, at adult stages when CA1 PN labeling is dramatically attenuated (Figure 1B).

At early postnatal ages the hippocampal network is dominated by GABAergic tone that exerts an excitatory influence upon postsynaptic PNs prior to establishment of low intracellular chloride levels by upregulation of KCC2 [Ben-Ari2002, Ben-Ari2007]. Between the ages of P5-10 this depolarizing GABAergic input coordinates spontaneous, synapse-driven network synchronizations in the form of giant depolarizing potentials (GDPs) that propagate throughout the hippocampus to drive activity-dependent circuit maturation and MGE-derived interneurons, particularly Sst-INs, are strongly implicated in this pacing [Ben-Ari1989, Picardo2011, Wester2016, Flossmann2019, Cossart2022]. To probe for eGlyR-mediated regulation of GABAergic tone in the developing hippocampus we recorded spontaneous GABAergic postsynaptic currents (G-PSCs) and the GABAergic component of GDP-associated currents (GDP-Is) in acute hippocampal slices from P5.5–9.5 wild-type and *Grin3a* knockout (*Grin3a*-KO) mice while applying CGP-78608 to remove any eGlyR desensitization (Figure 1C-F). GABAergic synaptic events were electrically isolated by voltage-clamp at 0 mV where both G-PSCs and GDP-Is are detectable as outward GABA$_A$R-mediated currents without significant contamination from phasic glutamatergic transmission (alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor (AMPAR)-mediated) needed in concert with GABAergic activity to sustain GDPs. Recordings were performed in CA3 PNs which lack *Grin3a* (Figure 1B; Supplemental Figure 1) to limit confounds of pyramidal cell eGlyRs through network effects and conventional NMDARs were blocked to eliminate confounds of CGP-78608 effects at their glycine binding sites. Under these conditions, CGP-78608 robustly increased CA3 PN G-PSC frequencies in WT but not *Grin3a*-KO mice in a reversible fashion (Figure 1C-E). Concomitantly, we consistently observed that GDP-I frequencies

declined by the end of CGP-78608 application in WT mice without consistent effects in *Grin3a*-
KOs (Figure 1F). Our initial expression profiling and functional observations suggest that
immature hippocampal interneurons express eGlyRs that are liganded and desensitized by
ambient Gly levels. Moreover, engaging interneuron eGlyRs has the capacity to dramatically
increase their excitability to enhance GABAergic tone throughout the hippocampal circuit and
potentially disrupt spontaneous rhythmicity.
170
171 **Grin3a expression in hippocampal NGFCs and Sst-INs yields functional eGlyRs throughout**
172 **development that influence excitability.**
173
To obtain a more granular appreciation of *Grin3a* expression within specific interneuron subtypes
we analyzed single-cell RNA sequencing (scRNA-seq) data from adult mice when interneuron
transcriptional identities are well defined and segregated ([Harris2018, Hodge2019, Krienen2020,
Yao2021, Mahadevan2021]). Focusing on GABAergic interneurons from a multi-institute
integrated scRNA-seq dataset (see Methods) we confirmed that most Sst-INs express high levels
of *Grin3a* consistent with prior studies (Figure 2A;Supplemental Figure 1) [Pfeffer2013, Paul2017,
Murillo2021, Bossi2022]. Indeed, *Grin3a* serves as a secondary molecular marker for Sst-INs and
within these cells the *Grin3a* locus exhibits prominently open chromatin and low DNA
methylation states that are indicative of actively transcribed genes promoting cell identity
([Pfeffer2013, Yao2021]). We also discovered previously unappreciated *Grin3a* expression across
most NGFC subsets (Figure 2A; Supplemental Figure 1). Amongst NGFCs *Grin3a* is expressed in
both Lamp5$^+$/Lhx6$^+$ MGE-derived subsets, also known as Ivy cells (IvyCs), and Lamp5$^+$/Lhx6$^-$
caudal ganglionic eminence (CGE)-derived subsets that are both well represented in the rodent
hippocampus [Fuentealba2008, Tricoire2010, Tricoire2011]. Consistent with these scRNA-seq
profiles mClover-GluN3A labeled cells co-expressed Sst in stratum oriens and neuronal nitric oxide
synthase (nNOS) in putative NGFCs that comprise the dominant interneuron subtype in stratum
lacunosum moleculare (Figure 2B; Supplemental Figure 1).
To directly confirm functional eGlyR expression within hippocampal interneurons we
electrophysiologically measured their responses to brief, pressure applied Gly (10 mM) delivered
locally via a glass micropipette (Figure 2C-D). Sst-INs and NGFCs were targeted using Sst-Cre:Ai14
and NPY-Cre:Ai14 tdTomato reporter mice respectively. To pharmacologically isolate eGlyR-
mediated currents, we included a cocktail of receptor antagonists in the perfusing artificial
cerebrospinal fluid (ACSF) to block conventional ionotropic glycine (strychnine, 1 µM) glutamate
(DL-APV, 50 µM, DNQX, 10 µM), and GABA receptors (picrotoxin, 50 µM; bicuculline, 10 µM).
With this cocktail any initial current induced by Gly in regular ACSF was essentially abolished
indicating minimal eGlyR-mediated current availability (Figure 2C-D and see Supplemental Figure
2 for sequential antagonist example). However, application of CGP-78608 in the presence of the
antagonist cocktail unmasked robust Gly-evoked currents in both Sst-INs and NGFCs consistent
with mediation through eGlyRs that are desensitized under baseline conditions (Figure 2C-D).
Importantly, these Gly-evoked currents required *Grin3a*, as they were absent in either constitutive
or conditional *Grin3a* knockout mice (*Grin3a*-KO; Sst-*Grin3a*-cKO and NPY-*Grin3a*-cKO), though
ventral CA1 PNs retained CGP-78608 sensitive Gly-evoked currents in both interneuron
conditional KOs (Figure 2E). Similarly, recordings from stratum oriens Sst-INs in Nkx2.1-*Grin1*-
cKOs lacked CGP-78608 sensitive Gly-evoked currents consistent with the requirement for both

GluN1 and GluN3 to form functional eGlyRs (Figure 2E). Further pharmacological evidence that
the currents observed in hippocampal interneurons reflect bona fide eGlyR conductance includes
antagonism by the GluN1/3A Gly binding site antagonist 5,7-dichlorokynurenic acid (DCKA) and
suppression by the negative allosteric modulator EU1180-438 (Figure 2C-D, [Chatterton2002,
Awobuluyi2007, Zhu2020, Bossi2022]). In addition, we found that CGP-78608 unmasked Gly-
evoked currents in both Sst-INs and NGFCs displayed significant potentiation in response to a
newly developed GluN3-selective positive allosteric modulator of eGlyRs, EU1180-490, identified
via focused library screening of human GluN1/GluN3A (Figure 2C-D, [Nguyen2025]). In sum,
these findings reveal robust eGlyR expression within hippocampal Sst-INs and NGFCs providing
potential cellular mediators for the regulation of GABAergic tone onto CA3 PNs by CGP-78608 in
the developing hippocampus. Indeed, a more detailed temporal developmental profiling of
pressure applied Gly-evoked responses confirmed functional eGlyR expression within Sst-INs and
NGFCs at early postnatal ages with Sst-INs exhibiting a modest reduction in current amplitudes
with age and NGFCs displaying increased current amplitudes (Supplemental Figure 3).

Building on these findings, we next investigated whether ambient Gly working through
eGlyRs has the capacity to directly modulate excitability of *Grin3a*-expressing hippocampal
interneurons as suggested by the ability of CGP-78608 to drive GABAergic tone. First, we assessed
whether eGlyRs provide a tonic current in hippocampal interneurons by examining the effects of
DCKA on membrane holding/leak currents in the presence of the conventional ionotropic
glycinergic, glutamatergic, and GABAergic receptor antagonist cocktail. Application of DCKA
modestly reduced inward holding current at -70 mV in NGFCs but not Sst-INs (Figure 3A-B). This
differential influence of tonic eGlyR currents in the two hippocampal interneuron subtypes is
reminiscent of differences reported between amygdala principal cells and neocortical Sst-INs and
may reflect regional and/or cell-type specific differences in ambient *ex vivo* Gly levels, as well as
eGlyR redox and desensitization states [Grand2018, Bossi2022]. Although recorded in the same
gross *ex vivo* structure under the same recording conditions NGFCs and Sst-INs were targeted in
very deep or superficial CA1 respectively potentially with unique microenvironmental influences.
Conversely, both NGFCs and Sst-INs consistently displayed an increase in holding current in
response to application of CGP-78608 indicating sufficient endogenous ambient Gly levels to
ligand and largely desensitize the eGlyRs on both interneuron populations (Figure 3C-D). The
effects of CGP-78608 on holding current were reversed by DCKA and enhanced by EU1180-490
consistent with mediation through eGlyRs (Figure 3D and Supplemental Figure 4). Moreover, in
the current-clamp recording configuration CGP-78608 depolarized both NGFCs and Sst-INs
sufficiently to induce or increase action potential (AP) firing (Figure 3E-F). Interestingly, EU1180-
490 alone could also drive increased firing of Sst-INs, suggesting that this positive allosteric
modulator may relieve eGlyR desensitization directly (Figure 3F). Together our findings
conclusively demonstrate that ambient Gly levels have the capacity to strongly influence NGFC
and Sst-IN excitability, and hence circuit GABAergic tone, throughout development via eGlyRs
within these interneuron subpopulations.

An unexpectedly outsized contribution from NGFCs/IvyCs in early postnatal GABAergic tone
and rhythmicity under control and CGP-78608 conditions.

A significant body work has implicated Sst-INs as major contributors to early postnatal
hippocampal and neocortical network GABAergic tone, coordination of population synchrony,
and circuit connectivity/maturation [Picardo2011, Anastasiades2016, Marques-Smith2016,
Oh2016, Tuncdemir2016, Modol2017, Flossmann2019, Wang2019, Su2020, Cossart2022,
Dard2022]. However, similar studies regarding the role of NGFCs/IvyCs in nascent network
function are lacking despite these cells representing perhaps the most abundant dendrite
targeting interneurons of the hippocampus [Bezaire2013, Overstreet-Wadiche2015]. To dissect
the contributions of Sst-INs and NGFCs to GABAergic tone and pacing within the developing
hippocampus under control and CGP-78608 conditions we employed an optogenetic strategy
using conditional expression of the light-activated proton pump Achaerhodopsin-3 (Arch) to
silence distinct interneuron populations (Figure 4). First, to determine the maximal effectiveness
of this approach we used pan-GABAergic neuron Arch mice generated by driving conditional Arch
expression with Cre under control of the vesicular inhibitory amino acid transporter gene (VGAT-
Cre:ArCh). In control conditions (normal ACSF) light driven suppression of all GABAergic
interneurons immediately and persistently blocked the majority of G-PSCs and GDP-Is recorded
in early postnatal CA3 PNs of VGAT-Cre:ArCh mice (Figure 4A, C-F). Any residual G-PSCs and GDP-
Is in light could reflect AP independent spontaneous GABA release and/or stochastic escapes from
Arch-mediated suppression by interneurons with high input resistance during prolonged light
epochs (for example, by a large spontaneous AMPAR-mediated glutamatergic input). The effects
of Arch activation on G-PSC and GDP-I frequencies reversed upon light termination which typically
yielded a bout of "rebound light-off GDPs" as expected for the synchronous release of
interneurons from hyperpolarization that can drive them to AP threshold triggering a window of
reverberant network activity due to the excitatory nature of GABA at this stage of development
(Figure 4A, G and see [Wester2016, Flossmann2019]). Critically, light driven suppression of
interneuron activity in VGAT-Cre:ArCh mice significantly inhibited G-PSCs in the presence of CGP-
78608 confirming that our optogenetic approach continues to restrain interneurons under
conditions of eGlyR-mediated enhanced excitability (Figure 4B-D).

In control conditions, selective silencing of Sst-INs with light while recording from early
postnatal CA3 PNs yielded highly inconsistent effects that on average modestly depressed G-PSC
frequencies but did not significantly alter GDP-Is (Figure 4A, C-F). Successful light-mediated
control of the Sst-IN population was evidenced by the high probability of light-off rebound GDPs
observed in Sst-Cre:ArCh mice (Figure 4A, G). Surprisingly, Sst-IN silencing failed to significantly
depress G-PSCs in the presence of CGP-78608, suggesting that the massive recruitment of
GABAergic tone by eGlyRs occurs largely independent of Sst-IN output (Figure 4B-D). In contrast,
optogenetic silencing in NPY-Cre:ArCh mice more closely matched our observations from pan-
GABAergic neuron silencing with robust inhibition of G-PSC and GDP-I frequencies in control
conditions, generation of rebound light-off GDP-Is, and continued inhibition of G-PSCs in the
presence of CGP-78608 (Figure 4). Though NPY is strongly expressed in both NGFCs and Sst-INs,
potentially confounding interpretation of our optogenetic results, subtractive reasoning points to
an outsized role for NGFCs/IvyCs in providing GABAergic tone and rhythmicity within the early
postnatal CA3 hippocampus. Indeed, the inability of selective Sst-IN silencing to match the impact
of NPY-IN silencing on control G-PSCs and GDP-Is requires consideration of an NPY$^+$/Sst$^-$ IN cohort,
with NGFCs/IvyCs being the most logical candidates. This interpretation is further bolstered by
considering the limited possible sources of GABAergic input recruited by eGlyRs. As NPY-IN, but

not Sst-IN, silencing significantly blocks G-PSCs recruited by eGlyR activation and *Grin3a*
expression is largely limited to NGFCs/IvyCs and Sst-INs, our findings implicate NGFCs/IvyCs as
the dominant source of CGP-78608 recruited GABAergic tone in the developing CA3 hippocampal
circuit.
To independently test whether NGFCs/IvyCs are largely responsible for the increased
GABAergic tone recruited by eGlyR activation in early postnatal tissue, we evaluated CGP-78608
effects on G-PSCs and GDP-Is in CA3 PNs of conditional *Grin3a* knockout mice (Figure 5). For these
experiments we separated the CGP-78608 effects on G-PSCs and GDP-Is into early and late phases
since closer temporal inspection of our recordings in wild type mice revealed that while G-PSC
frequencies continually increase and plateau during drug application, GDP-Is undergo biphasic
effects first increasing in frequency then decreasing (Figure 5). The initial increase in GDP-I
frequency is accompanied by decreasing amplitudes until ultimately events cannot be reliably
resolved from the increased G-PSCs (eg. Figure 5A-B). This profile for GDP-Is is consistent with
the ability of CGP-78608 to increase IN excitability and ultimately drive them beyond threshold
(Figure 3). A transient increase in GDP frequency could reflect enhanced excitability of INs
participating in GDP generation and propagation; however, as greater numbers of INs are driven
to threshold out of phase with the rhythmic network, and perhaps even into depolarization block,
the ability to sustain synchronization may be compromised prompting degradation of GDP-Is into
uncoordinated unitary events. Importantly, in constitutive *Grin3a*-KO mice all early and late phase
CGP-78608 effects on G-PSCs and GDP-Is are eliminated confirming the pharmacological
specificity for *Grin3a*-containing receptors in mediating transient and sustained changes in
GABAergic tone and circuit pacing (Figure 5Ca,b,f-Da,b,f). However, selective removal of *Grin3a*
across the entire IN population (VGAT-*Grin3a*-cKO mice) eliminated only the CGP-78608 induced
increase in G-PSCs and the transient increase in GDP-Is (Figure 5Cc,f-Dc,f). Thus, late-stage GDP
inhibition by CGP-78608 could reflect network changes triggered by eGlyRs in PNs, for example
through polysynaptic interactions between eGlyR expressing CA1 PNs to the CA3 network that
enhance network excitability to disrupt rhythmicity [Sik1994, Kim2015, Bocchio2020].
Surprisingly, but consistent with our optogenetic findings, selective removal of *Grin3a* from Sst-
INs had no effect on early or late phase effects of GCP-78608 on CA3 PN G-PSCs or GDP-Is (Figure
5Ce,f-De,f). In contrast, NPY-*Grin3a*-cKO mice failed to exhibit persistent or transient increases in
CA3 PN G-PSCs or GDP-Is respectively in response to CGP-78608, essentially phenocopying the
VGAT-*Grin3a*-cKO mice (Figure 5Cd,f-Dd,f). These findings are entirely consistent with our
optogenetic results further indicating that the majority of GABAergic recruitment onto CA3 PNs
by removal of eGlyR desensitization arises from NGFCs/IvyCs highlighting a previously
unappreciated critical role for this class of interneuron in nascent hippocampal circuit activity.
330
331 **eGlyR-mediated excitation of Sst-INs regulates sharp wave ripples in adult hippocampal circuits.**
332
In the mature hippocampus the dominant spontaneous synchronous network signature is that of
sharp wave ripples (SWRs) that are typically observed *in vivo* during "off-line" brain states such as
quiet wakefulness, consummatory behaviors, and non-REM sleep [Buzsáki2015]. These high-
frequency synchronized oscillatory events play a critical role in memory consolidation and
cognition by allowing rapid network replay of exploratory driven activity patterns over
compressed time periods and are characterized by large-amplitude sharp waves with

superimposed high-frequency ripple oscillations [Girardeau2009, Jadhav2012, Buzsáki2015,
Joo2018]. SWRs are temporally and spatially coordinated by stereotyped activation and silencing
of discrete interneuron subtypes, including Sst-INs with oriens-lacunosum moleculare (OLM) Sst-
INs exhibiting suppression or excitation depending on experimental preparation and long-range
projecting Sst-INs undergoing strong recruitment during SWRs [Royer2012, Varga2012,
Somogyi2014, Buzsáki2015, Katona2017, Szabo2022]. Interestingly, interference with *Grin3a*
expression can disrupt cognition and consolidation of episodic memories; however, the influence
of eGlyRs on mature hippocampal network oscillations has not been previously investigated
[Gallinat2007, Roberts2009, Papenberg2014]. As our findings confirm continued expression of
functional eGlyRs in adult hippocampal Sst-INs we directly evaluated whether eGlyRs have the
capacity to influence SWR rhythmicity (Figure 6).
Spontaneous SWRs were readily detected with local field potential (LFP) recordings in the
CA3 PN layer of acute hippocampal slices from adult mice maintained at interface (Figure 6A,
[Hajos2009, Pelkey2015]). Importantly, spontaneous SWRs persist, and may even be enhanced,
in the absence of conventional NMDAR-mediated transmission, and thus, our recordings were
performed with conventional NMDARs blocked to eliminate pharmacological confounds to data
interpretation of findings with eGlyR-related compounds [Behrens2005, Colgin2005]. In slices
from wild type mice application of CGP-78608 or EU1180-490 significantly depressed the
frequency of SWR occurrence in a reversible fashion (Figure 1A-B,E). Inhibition of SWRs by CGP-
78608 or EU1180-490 was absent in *Grin3a*-KO mice confirming the pharmacological specificity
of these reagents for eGlyRs (Figure 6C,E). Most interestingly, selective removal of *Grin3a* from
Sst-INs in Sst-*Grin3*-cKO mice greatly attenuated SWR sensitivity to CGP-78608 compared to
control animals leaving modest inhibition that was not significantly different from data obtained
in *Grin3a*-KOs (Figure 6D-E). Collectively, these findings illustrate that eGlyRs, particularly those
expressed by Sst-INs, have the capacity to influence coordinated network rhythmicity in the adult
hippocampus.
365
366 **Evolutionary conservation of eGlyR function in hippocampal interneurons.**
367
Genetic mutations or altered expression levels of *GRIN3A* in humans are implicated in cognitive
deficits and diverse neurological/neuropsychiatric conditions such as schizophrenia, bipolar
disorder, addiction, epilepsy, and Huntington's Disease [Mueller2004, Gallinat2007, Marco2013,
Yuan2013, Yang2015, Lee2016, Huang2017, Marco2018, Crawley2022]. Historically,
interpretation and posited roles of dysregulated *GRIN3A* in disease progression have focused on
potential influences over conventional NMDARs (eg. triheteromeric GluN1/2/3 contributions) and
phasic synaptic transmission [Kehoe2013, Perez-Otano2016, Crawley2022]. The emerging
picture in mice that GluN3A readily assembles as eGlyRs in diverse neurons expressing *Grin3a* to
tonically regulate excitability through ambient Gly provides a new lens through which to consider
GluN3A-associated disease etiology [Bossi2023]. However, this demands functional confirmation
that eGlyRs are evolutionarily conserved regulators of neuronal excitability. Thus, in our final
series of experiments we aimed to validate our rodent findings in higher species by functionally
interrogating hippocampal interneurons and network rhythmogenesis in adult non-human
primates (NHPs).

Analysis of publicly available scRNA-seq data sets online indicate conserved *GRIN3A*
expression within neocortical/hippocampal Sst-INs and NGFCs in both NHPs and humans
(Supplemental Figure 5). By mFISH we confirmed that *GRIN3A* exhibits a high degree of
colocalization with *SST* and *LAMP5* throughout the adult NHP (rhesus macaque) hippocampus
(Figure 7A-B). To facilitate Sst-IN identification for functional evaluation of eGlyRs we
intracranially injected adult macaques with novel enhancer based viral tools designed to limit
transgene expression to Sst-INs, pAAV-PHP.eB-BiSSTe10-dTomato and pAAV-PHP.eB-BiSSTe4
[Furlanis2025]. Then 5-6 weeks after hippocampal AAV introduction we made acute hippocampal
slices and targeted fluorescently labeled cells for electrophysiological interrogation (Figure 7C).
Consistent with our mouse findings NHP Sst-INs exhibited CGP-78608 sensitive Gly-evoked
excitatory currents in the cocktail of conventional glutamate, GABA, and glycine receptor
antagonists (Figure 7D). Importantly, these Gly-evoked currents were facilitated by EU1180-490
consistent with mediation by eGlyRs (Figure 7D). Parallel recordings in NHP anterior hippocampal
CA1 PNs similarly yielded CGP-78608 and EU1180-490 sensitive Gly-evoked currents (Figure 7D).
Finally, as SWRs are evolutionarily conserved network oscillations observed across multiple
species, including NHPs and humans [Buzsáki2015] we assessed whether engaging eGlyRs by
removal of desensitization/allosteric enhancement could regulate rhythmicity in NHPs.
Congruent with our rodent findings, we found that either CGP-78608 or EU1180-490 significantly
depressed the incidence of SWRs recorded in the CA3 PN layer of adult NHP slices in a reversible
fashion (Figure 7E). In total our expression and functional profiling confirm strong conservation
of eGlyR function over approximately 70-100 million years of evolution validating its relevance as
a potential novel therapeutic target to regulate neuronal excitability across a constellation of
human neurological and neuropsychiatric disorders.

405
406
407

Discussion
Emerging evidence indicates that GluN3A subunits frequently assemble with GluN1 to
form unconventional NMDARs that serve as eGlyRs in diverse neurons throughout the brain,
consistent with initial cloning studies [Perez-Otano2001, Chatterton2002, Grand2018, Otsu2019,
Zhu2020, Bossi2022, Rouzbeh2023, Pizzamiglio2025]. Our present findings unequivocally
demonstrate that *Grin3a*[+] hippocampal Sst-INs and NGFCs/IvyCs express functional eGlyRs from
early postnatal to adult ages and across evolution from mouse to macaque. Excitation of
NGFCs/IvyCs and Sst-INs with endogenous ambient glycine through eGlyRs has the capacity to
regulate spontaneous synchronized hippocampal network activities throughout development.
Putative roles for GluN3A in synapse development and circuit function inferred from preclinical
knockout/overexpression studies and genetic mutation linkage clinical studies are often modeled
on cell autonomous influences of GluN3A over conventional NMDARs through formation of
triheteromeric receptors or by serving as dominant negative subunits. Our findings, combined
with others, highlight that eGlyR mediated regulation of cell and circuit excitability through both
cell autonomous and cell non-autonomous mechanisms must also be considered to understand
the role of *Grin3a*/GluN3A in brain development, plasticity, and disease [Otsu2019, Bossi2022,
Hurley2024, Pizzamiglio2025].

The gating of eGlyRs is unusual with glycine serving as both agonist through high affinity GluN3A binding and functional antagonist by driving profound desensitization upon binding the lower affinity GluN1 subunit [Chatterton2002, Yao2006, Awobuluyi2007, Madry2007, Bossi2022]. Thus, eGlyRs are well suited to provide tonic excitation at low levels of ambient glycine with an auto-inactivation mechanism at higher glycine levels to limit receptor mediated hyperexcitability/excitotoxicity. Interestingly, higher ambient glycine levels leading to GluN1 binding will promote a switch to increased phasic glutamatergic excitation by saturating co-agonist GluN1 binding sites of conventional NMDARs concomitant with the suppression of tonic eGlyR mediated excitation. Though we observed a modest influence of tonic eGlyR activation on NGFC resting membrane potential, the robust depolarizing influence of CGP-78608 on both Sst-INs and NGFCs indicates that ambient glycine levels throughout our acute hippocampal slices are by and large sufficient to bind GluN1 subunits and drive eGlyR desensitization. This is consistent with decades of electrophysiological interrogation of conventional NMDARs in acute hippocampal slices across many labs without the need for extracellular glycine supplementation. Our findings mirror observations in amygdala and neocortical Sst-INs that exhibit eGlyR desensitization under basal slice conditions [Bossi2022]. In contrast, under the same conditions eGlyRs in amygdala PNs exhibit minimal desensitization and provide significant tonic depolarizing influence that is titrated through neuromodulation of glycine transporter activity [Bossi2022]. Thus, even within the same gross brain structure under the same conditions eGlyR glycine-binding site occupancies may differ across neuronal subtypes according to differential extracellular ambient glycine levels. A very recent report provided evidence for tonic Gly liganding and depolarizing influence of eGlyRs on interneurons with somatic localization to stratum oriens but not PNs and interneurons residing in CA1 stratum pyramidale [Pizzamiglio2025]. While these findings do not entirely align with our observations, perhaps due to recording conditions and animal ages, they further highlight potential for local Gly microenvironments in modulating cell excitability through eGlyRs. It will be important in future investigations to probe both sources and spatiotemporal plasticity of ambient hippocampal glycine *in vivo* across development. In addition, the physiological conditions and extent to which Sst-IN and NGFC/IvyC eGlyR functions are subject to activity dependent modulation through redox regulation, acidification, and zinc should also be evaluated [Madry2007, Cummings2017]. Moreover, it remains possible that there exists an as yet undiscovered endogenous neuroactive molecule that promotes eGlyR function through removal of desensitization or allosteric potentiation mechanistically analogous to CGP-78608 and 1180-490.

GABAergic INs throughout the neocortex, amygdala, and hippocampus have common developmental origins in the ganglionic eminences and discrete IN subtypes in each of these structures display considerable genetic, physiological, and synaptic connectivity/function homologies [Zeisel2015, Tremblay2016, Pelkey2017, Gouwens2020, Hajos2021, Yao2021]. However, granular investigation can reveal regional differences in receptor expression profiles for homologous IN subpopulations across brain regions [Caccavano2025]. Expression of eGlyRs within Sst-INs is universal across the cortical structures thus far examined ([Bossi2022, Pizzamiglio2025], and present findings). In contrast, parvalbumin (PV) IN expression of eGlyRs appears inconsistent with functional expression demonstrated in amygdala and hippocampal PV INs but not their neocortical counterparts [Bossi2022, Pizzamiglio2025]. Curiously, *Grin3a* is minimally expressed within transcriptionally defined rodent hippocampal and neocortical PV INs,

though increases through evolution (Supplemental figure 5).   These findings highlight the
importance of careful cell-type specific evaluation across brain regions and species in mapping
eGlyR expressing neurons to understand their circuit influences and potential for therapeutic
leverage. In addition to the utility of emerging nervous system wide transcriptional data sets
across species, identification of rodent GluN3A/eGlyR expressing neurons throughout the brain
will be greatly facilitated by the newly developed mClover-GluN3A reporter mouse line.

Here we further identify hippocampal MGE- and CGE-derived NGFC/IvyC cohorts as neuronal
elements through which ambient Gly can regulate circuit function by eGlyR signaling.
Transcriptional profiling indicates conserved strong *Grin3a* expression levels in neocortical NGFCs
defined by *Id2* and *Lamp5* expression (Supplemental Figure 5A). In evaluating NGFCs/IvyCs
through the lens of eGlyRs we discovered a previously unappreciated capacity for these
interneurons to dramatically shape GABAergic tone and network pacing in the developing
hippocampus. Past studies have largely implicated Sst-INs as major contributors to early
postnatal hippocampal and neocortical network GABAergic tone, coordination of population
synchrony, and circuit connectivity/maturation [Picardo2011, Anastasiades2016, Marques-
Smith2016, Oh2016, Tuncdemir2016, Modol2017, Flossmann2019, Wang2019, Su2020,
Cossart2022, Dard2022]. However, while synchronization of Sst-IN activation has the capacity to
generate GDP-like events (eg. light off events in Sst-Cre: ArCh) minimizing Sst-IN activity fails to
consistently depress spontaneous GABAergic tone or intrinsic GDP pacing in the developing
hippocampus (see also [Flossmann2019]). In contrast, the same optogenetic strategy to silence
NGFC/IvyC activity consistently suppressed spontaneous GABAergic tone and GDPs in CA3 PNs.
Moreover, NGFC/IvyC inputs were found to dominate the excess inhibitory drive recruited by
eGlyR activation with CGP-78608 in the immature hippocampus. These findings, together with
recent evidence implicating immature parvalbumin interneurons in early network GABAergic
pacing [Caccavano2025], highlight the need for further investigation of multiple cardinal IN
classes in coordinating developing hippocampal circuit dynamics.

The increasing availability of publicly accessible single cell transcriptomic data sets has greatly
facilitated genetic profiling of discrete neuronal populations across species as a starting point for
validating translational relevance of rodent centric studies to higher species.   However,
opportunities for functional validation in neuronal cells and circuits of higher species remains
limited by tissue availability and genetic identification of discrete cell types. Here we employed
recently developed enhancer based AAV tools to facilitate selective targeting of Sst-INs in live NHP
tissue for direct functional interrogation of the evolutionary conservation of eGlyR expression in
this neuronal subpopulation [Furlanis2025]. Further development of such tools for cell-type
specific targeting will greatly expand the discrete neuronal subtypes throughout the brain for
targeting across species in preclinical studies requiring selective cell labeling/actuation and for
future therapeutic intervention or gene therapy [Tasic2025]. Our findings confirmed that
expression of *GRIN3A* is indeed predictive of operational eGlyRs in NHPs as in rodents. The
emerging picture from our current findings combined with others [Grand2018, Zhu2020,
Bossi2022], necessitate consideration of eGlyR dysregulation in addition to conventional NMDARs
as an underlying feature of neurological disorders associated with genetic mutations or
alterations in *GRIN3A* expression such as schizophrenia, bipolar disorder, addiction, epilepsy, and
Huntington's Disease [Mueller2004, Gallinat2007, Marco2013, Yuan2013, Yang2015, Lee2016,
Huang2017, Marco2018, Crawley2022]. Importantly, NHP hippocampal Sst-IN and CA1 PN eGlyRs

displayed similar pharmacological profiles to those in rodents with CGP-78608 "awakening" the
receptors from desensitization and the newly described allosteric modulator EU1180-490 able to
potentiate their responses. Moreover, as in rodents, both compounds worked independent of
conventional NMDARs to depress *in vitro* SWR activity, illustrating the ability to leverage eGlyRs
expressed in circuit minority cell populations for regulation of network population activities
across species. These findings provide a foundation for further native eGlyR evaluation across
species in health and disease and confirm their potential as promising therapeutic targets using
small molecule selective allosteric modulators.
521
522

Acknowledgements

This research was supported [in part] by the Intramural Research Program of the National Institutes of Health (NIH). The contributions of the NIH author(s) are considered Works of the United States Government. The findings and conclusions presented in this paper are those of the author(s) and do not necessarily reflect the views of the NIH or the U.S. Department of Health and Human Services. We thank Apratim Mitra, Mira Sohn and Ryan Dale (NICHD Bioinformatics and Scientific Programming Core) for assistance with downloading and analyzing scRNAseq datasets. These analyses utilized the computational resources of the NIH Biowulf cluster (http://hpc.nih.gov).

Author contributions

JHK was responsible for project conceptualization, experimental design, electrophysiology experiments, and imaging, analyses. KAP and CJM conceptualized, designed, and supervised the project. JHK, AV GAV, TB, RC, and KAP contributed to electrophysiology experiments. AV and GAV conducted AAV injections and surgery for mice. XY and SH conducted IHC, ISH, morphological reconstructions, and imaging. VM and APV conducted RNAseq analyses, RiboTag experiments and analysis. OCC, AIN and IPO provided Sst-Cre::$Grin3a^{fl/fl}$ and mClover-GluN3A reporter IHC analysis. JHK, DA, KP generated Npy-/VGat-Cre:: $Grin3a^{fl/fl}$ mice and maintained mouse colonies. DL, ZZ, HY, SFT, SLS, MPE, and DCL provided GluN3a specific allosteric modulators (EU1180-490 and EU1180-438). MAGE, ACC, ACT, PH, AP, AM, JP, and BBA conducted NHP handling, injections, and surgeries. EF, YW, MD, BLG, JD, and GF provided support and design of enhancer AAV constructs. All authors participated in drafting the manuscript with JHK and KAP serving as principal writers. MAGE, BBA, GF, IPO, SFT and CJM provided critical funding support for the study.

Declaration of interests

SFT is a member of the SAB for Eumentis Therapeutics and Neurocrine Biosciences, and a member of the Medical Advisory Board for the GRIN2B Foundation, CureGRIN, and CombinedBrain, co-founder of NeurOp Inc and Agrithera Inc, and a consultant for GRIN Therapeutics and Seyltx. SFT is a PI on a research grant from GRIN Therapeutics to Emory University School of Medicine. SLS, MPE, DCL, and SFT are co-inventors on Emory-owned Intellectual Property that includes allosteric modulators of NMDA receptor function.
The authors declare no further competing interests.

 STAR★METHODS

 KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit anti-Sst	Thermo Fisher Scientific	Cat# PA5-85759, RRID:AB_ 2792896
Guinea Pig anti-RFP	Synaptic Systems	Cat# 390005, RRID:AB_2737051
Rabbit anti-GFP	Abcam	Cat# ab290, RRID:AB_303395
Rabbit anti-RFP	Antibodies-Online	Cat# ABIN129578, RRID:AB_10781500
AF488 Goat anti-Rabbit IgG (H+L)	Thermo Fisher Scientific	Cat# A-11008, RRID:AB_143165
AF555 Goat anti-Guinea Pig IgG (H+L)	Thermo Fisher Scientific	Cat# A-21435, RRID:AB_2535856
AF488 Goat anti-Rabbit IgG (H+L)	Thermo Fisher Scientific	Cat# A-11034, RRID:AB_2576217
AF555 Goat anti-Rabbit IgG (H+L)	Thermo Fisher Scientific	Cat# A-21429, RRID:AB_2535850
Bacterial and virus strains		
pAAV(AAV5)-pCAG-FLEX-EGFP-WPRE	Addgene[Chen2014]	RRID:Addgene_51502-AAV5
pAAV-PHP.eB-BiSSTe4-dTomato	Addgene [Furlanis2025]	RRID:Addgene_213944
pAAV-PHP.eB-BiSSTe10-dTomato	Addgene [Furlanis2025]	RRID:Addgene_213814
Chemicals, peptides, and recombinant proteins		
DL-APV sodium salt	Abcam	Cat# ab120271
DNQX disodium salt	Abcam	Cat# ab120169
(-)-Bicuculline methobromide	Abcam	Cat# ab120109
Picrotoxin	Abcam	Cat# ab120315
Strychnine	Millipore Sigma	Cat# S0532
CGP-78608 hydrochloride	Tocris	Cat# 1493
EU1180-438	Stephen F Traynelis [Zhu2020]	
EU1180-490	Stephen F Traynelis [Nguyen2025]	
5,7-Dichlorokynurenic Acid, Sodium Salt (DCKA)	Millipore Sigma	Cat# 5.04841
Biocytin	Millipore Sigma	Cat# B4261
AF488 streptavidin	Thermo Fisher Scientific	Cat# S11223
AF555 streptavidin	Thermo Fisher Scientific	Cat# S21381
AF633 streptavidin	Thermo Fisher Scientific	Cat# S21375
DAPI	Millipore Sigma	Cat# D9542
RNAscope Multiplex Fluorescent v2 Assay	Advanced Cell Diagnostic	Cat# 323110
RNAscope mouse probe mmu-Sst	Advanced Cell Diagnostic	Cat# 461681
RNAscope mouse probe mmu-*Grin3a*-C2	Advanced Cell Diagnostic	Cat# 1225361-C2
Opal 520	AKOVA	Cat# OP-001001
Opal 650	AKOVA	Cat# OP-001006
ProLong Gold Antifade Mountant	Thermo Fisher Scientific	Cat# P10144
RNeasy Plus Micro Kit	Qiagen	Cat# 74034
RNA 6000 Pico kit	Agilent	Cat# 5067-1513
SMARTer Stranded Total RNA-Seq. Kit v2	Takara Bio	Cat# 634411

Experimental models: Organisms/strains		
Mouse: B6;129X1-*Grin3a*^{tm1Nnk}/J NR3a KO	The Jackson Laboratory [Das1998]	RRID:IMSR_JAX:029974
Mouse: *Grin3a*$^{fl/fl}$	Isabel Perez-Otano	
Mouse: *Grin1*$^{fl/fl}$	The Jackson Laboratory [Tsien1996]	RRID:IMSR_JAX:005246
Mouse: Sst-Cre::*Grin3a*$^{fl/fl}$	Isabel Perez-Otano Lab	
Mouse: Sst$^{tm2.1(cre)Zjh}$/J Sst-IRES-Cre	The Jackson Laboratory [Taniguchi2011]	RRID:IMSR_JAX:013044
Mouse: B6.Cg-NPY$^{tm1(cre)Zman}$/J NPY-IRES-Cre	The Jackson Laboratory [Milstein2015]	RRID:IMSR_JAX:027851
Mouse: C57BL/6J-Tg(Nkx2-1-cre)2Sand/J Nkx2.1-Cre	The Jackson Laboratory [Xu2008]	RRID:IMSR_JAX:008661
Mouse: Slc32a1$^{tm2(cre)Lowl}$/J VGAT-IRES-Cre	The Jackson Laboratory [Vong2011]	RRID:IMSR_JAX:016962
Mouse: B6.Cg-Gt(ROSA)26Sor$^{tm9(CAG-tdTomato)Hze}$/J Ai9(RCL-tdT)	The Jackson Laboratory [Madisen2010]	RRID:IMSR_JAX:007909
Mouse: B6.Cg-Gt(ROSA)26Sor$^{tm14(CAG-tdTomato)Hze}$/J Ai14(RCL-tdT)-D	The Jackson Laboratory [Madisen2010]	RRID:IMSR_JAX:007914
Mouse: B6.Cg-Gt(ROSA)26Sor$^{tm40.1(CAG-aop3/EGFP)Hze}$/J Ai40(RCL-ArchT/EGFP)-D	The Jackson Laboratory	RRID:IMSR_JAX:021188
Publicly available transcriptomic data, integrated and reanalysed		
scRNAseq data – processed count file and sample metadata (Allen 2019)	[Hodge2019]	https://transcriptomic-viewer-downloads.s3-us-west-2.amazonaws.com/mouse/transcriptome.zip
scRNAseq data – processed count file and sample metadata (Allen 2021)	[Yao2021]	https://assets.nemoarchive.org/dat-jb2f34y
scRNAseq data – processed count file and sample metadata (Harvard 2020)	[Krienen2020]	GSE151761
scRNAseq data – processed count file and sample metadata (Karolinska 2018)	[Harris2018]	GSE99888
scRNAseq data – processed count file and sample metadata (NIH 2021)	[Mahadevan2021]	GSE156201
snRNAseq human data	[Siletti2023]	https://alleninstitute.github.io/abc_atlas_access/descriptions/WHB_dataset.html
snRNAseq and MERFISH mouse data	[Yao2023]	https://alleninstitute.github.io/abc_atlas_access/descriptions/WMB_dataset.html
Software and algorithms		
pClamp 10.6	Molecular Devices	RRID:SCR_011323
Multiclamp 700B	Molecular Devices	RRID:SCR_018455
FIJI (ImageJ)	NIH	RRID:SCR_002285
Excel	Microsoft	RRID:SCR_016137
ZEN Microscopy Software	Zeiss	RRID:SCR_013672
SPOT 5.2	SPOT Imaging	RRID:SCR_014313

MATLAB 2023b	MathWorks	RRID:SCR_001622
R v3.5.0 and greater	R Foundation	https://www.R-project.org
Seurat v3.4	[Stuart2019]	https://github.com/satijalab/seurat
RStudio IDE	RStudio	https://www.rstudio.com
DESeq2	Bioconductor [Love2014]	RRID:SCR_015687

561
562 ### RESOURCE AVAILABILITY

563 #### Lead Contact

564 Further information and requests for resources and reagents should be directed to and will be
565 fulfilled by the lead contact, Kenneth Pelkey (pelkeyk2@nih.gov).

566
567 #### Materials Availability

568 This study did not generate novel, unique reagents.

569
570 #### Data and Code Availability

571 Data generated during this study are available upon request.

572
573 ### EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

574 #### Mice

575 All mice experiments were conducted in accordance with animal protocols approved by animal
576 care and use committee (ACUC) at the National Institute of Child Health and Human
577 Development (ASP# 20-045) and at Emory University.

578
Targeted neuronal subpopulations were achieved by fluorescently labeling, using Cre-
recombinase driven expression of floxed reporters. Experimental offspring were maintained as
heterozygous crosses from homozygous Cre-recombinase driver Sst-IRES-Cre
(RRID:IMSR_JAX:013044)[Taniguchi2011], NPY-IRES-Cre (RRID:IMSR_JAX:027851)[Milstein2015],
Nkx2.1-Cre (RRID:IMSR_JAX:008661)[Xu2008], *Grin1* (RRID:IMSR_JAX:005246) [Tsien1996]
mouse lines and homozygous floxed reporters Ai9(RCL-tdT)
(RRID:IMSR_JAX:007909)[Madisen2010], Ai14(RCL-tdT)-D (RRID:IMSR_JAX:007914)
[Madisen2010], Ai40(RCL-ArchT/EGFP)-D (RRID:IMSR_JAX:021188) lines.

587
588
Non-Human Primates (NHPs)

Adult rhesus macaque and marmoset tissue was obtained as described in [Caccavano2025]. All
experiments were performed in accordance with the ILAR Guide for the Care and Use of
Laboratory Animals and were conducted under Animal Study Protocols approved by the ACUC
at the National Institute of Mental Health. All procedures adhered to the applicable Federal and

local laws, regulations, and standards, including the Animal Welfare Act and Regulations and
Public Health Service policy (PHS2002).
596
597

METHOD DETAILS

Viral Injections (mouse)

For perinatal injections as described in [Caccavano2025], P0 mice were anesthetized by
hypothermia. Mice were bilaterally injected free-hand with 1 µL of solution containing
pAAV(AAV5)-pCAG-FLEX-EGFP-WPRE (Addgene 51502-AAV5, $\geq 7 \times 10^{12}$ vg/mL) in the lateral
ventricles: 0.8 mm anterior to lambda and 0.8-1 mm lateral to the sagittal suture, using a 32G
microliter Neuros Syringe (Hamilton).
605
606

Viral Injections and brain harvests in NHPs

As described in [Caccavano2025], briefly intracranial macaque injections were targeted using
stereotaxic coordinates derived from MRI and delivered using a needle guide for enhanced
accuracy [Fredericks2020]. Surgeries were performed under aseptic conditions in a fully
equipped operating suite. pAAV-PHP.eB-BiSSTe4-dTomato (Addgene 213944) or pAAV-PHP.eB-
BiSSTe10-dTomato (Addgene 213814) ($\geq 1\times10^{13}$ vg/mL) was injected into HPC and M1 of 6
rhesus macaques. Within HPC, 30-50 µL of total virus was injected, with 15-25 µL of virus
injected at each of 2 locations spaced approximately 2 mm apart in the antero-posterior plane,
caudal to the level of the uncus. Within M1, 30-40 µL of total virus was injected, with 10 µL of
virus injected at each of 3-4 locations spaced approximately 2 mm apart, targeted via direct
visualization.
For NHP brain extraction, (6-8 weeks after virus injection for macaque), animals were sedated
with ketamine/midazolam (ketamine 5-15 mg/kg, midazolam 0.05-0.3 mg/kg) and maintained
on isoflurane. A deep level of anesthesia was verified by an absence of toe-pinch and corneal
reflex response. Prior to brain removal and blocking, NHPs were transcardially perfused with
ice-cold sucrose-substituted artificial cerebrospinal fluid (SSaCSF) containing in mM: 90 Sucrose,
80 NaCl, 3.5 KCl, 1.25 NaH_2PO_4, 24 $NaHCO_3$, 10 Glucose, 0.5 CaCl, 4.5 $MgCl_2$ saturated with
carbogen (95% O_2, 5% CO_2), with osmolarity 310-320 Osm. Skulls were opened and brain was
removed and blocked in cold oxygenated SSACSF then sectioned as described below for mouse
tissue.
627

Acute brain slice preparation

Postnatal (p5–p10) or juvenile (p11–p21) mice were anesthetized with isoflurane and then
decapitated. The brain was dissected in ice-cold high-Mg^{2+} low-Na^+ ACSF for postnatal mice,
sucrose-substituted ACSF for juvenile or adults. The high-Mg^{2+} low-Na^+ ACSF contains the
following (in mM): 130 NaCl, 3.5 KCl, 24 $NaHCO_3$, 1.25 NaH_2PO_4, 4.5 MgCl, 0.5 $CaCl_2$, and 10
glucose with osmolarity 300-310 mOsm. The sucrose-substituted ACSF contains in mM 90

sucrose, 80 NaCl, 3.5 KCl, 24 NaHCO$_3$, 1.25 NaH$_2$PO$_4$, 4.5 MgCl, 0.5 CaCl$_2$, and 10 glucose.
Horizontal hippocampal slices (300 μm) were sectioned using a VT-1200S vibratome (Leica
Microsystems), then transferred and incubated in high-Mg^{2+} low-Na$^+$ ACSF at 32°C at least 1
hour before recording. Both ACSFs were saturated with 95% O$_2$ and 5% CO$_2$.

Ex vivo Electrophysiology recording

The brain slices were transferred to an upright microscope (Zeiss Axioskop 2), perfused at 2-3
ml/min with oxygenated ACSF containing in mM 130 NaCl, 3.5 KCl, 24 NaHCO$_3$, 1.25 NaH$_2$PO$_4$,
1.5 MgCl, 2.5 CaCl$_2$, and 10 glucose, and maintained at a temperature of 32°C-34°C. Electrodes
were pulled from borosilicate glass (World Precision Instruments) to a resistance of 2.5–4
Mohm using a vertical pipette puller (PP-830, Narishige). Whole-cell patch-clamp recordings
were made using a Multiclamp 700B amplifier (Molecular Devices), filtered at 3 kHz (Bessel
filter), and digitized at 10-25 kHz (Digidata 1550, Molecular Devices) and pClamp 10.6 software;
Molecular Devices). Uncompensated series resistance during recordings ranged from 7 to 30
Mohm and was monitored continuously throughout with 5mV voltage steps. In current-clamp
mode, cells were hold to -70 mV; in voltage-clamp mode, holding potentials were set to -70 mV
for measuring holding current and 0-10 mV for G-PSC and GDP-I. For whole-cell recordings,
internal solutions used were as follows. For recording glycine puff response of NGF-INs in
[Figure 2C] and action potentials and holding currents in [Figure 3], the internal solution
contains the following (in mM): 130 K-gluconate, 5 KCl, 2 MgCl$_2$, 2 MgATP, 0.3 NaGTP, 10 HEPES,
0.6 EGTA, and 0.2% biocytin. In other experiments, the internal solution contains in mM 123
CsCH$_3$SO$_4$, 4.5 NaCl, 4 MgATP, 0.4 NaGTP, 10 HEPES, 1 QX-314, 10 Bapta, and 0.2 % biocytin. The
pH of internal solutions was adjusted to 7.4, and osmolarity was adjusted to 285-290 mOsm.

For synaptically isolated conditions, NMDA, AMPA and GABA$_A$ receptors blocked by APV 50 μM,
DNQX 10 μM, BIC 10 μM and PTX 50 μM respectively. Binding CGP-78608 to the GluN1 subunit
does not only inhibit desensitization of eGlyRs, but also inhibits conventional GluN2-containing
NMDAR, influencing (excitatory) post-synaptic current. Thus, for recording network activity
changes by CGP-78608 application, the brain slices were incubated in 100 μM APV containing
ACSF for at least 1 hour prior to and during recording, to exclude influence of conventional
NMDAR-mediated current change by CGP on the GDP generation.

NGFCs express Neuro peptide Y (NPY) [Price2005, Karagiannis2009, Tricoire2010], and the
combination of the NPY-Cre mouse line and the cell location in the stratum lacunosum moleculare
(SLM) enables targeted recording of NGFCs. The labeled NPY-NGF-INs showed late spiking, which
is the electrophysiological signature of NGF-INs. For targeting Sst-INs, Sst-IRES-Cre::Ai9-tdTomato
or Sst-IRES-Cre::Ai14-tdTomato mice were adopted. In some experiments, putative OLM neurons
expressing Sst were targeted based on their morphology such as a horizontally oriented cell body
shape, and a deep layer location in the SO near the alveus. Their identification was confirmed by
extending the axons to reach the SLM through post hoc cell morphology reconstruction using
biocytin staining.

Local field potential recording

Local field potential (LFP) recordings were obtained on an interface recording set-up (Scientific
System Design Inc.). Brains of adult (>P60) mice were dissected as mentioned above in ice-cold
sucrose-substituted ACSF. Horizontal hippocampal slices were sectioned at 350 µm, then
transferred to an interface holding chamber with the slices suspended on the surface level of
oxygenated 2 mM $MgCl/CaCl_2$ ACSF containing the following (in mM): 130 NaCl, 3.5 KCl, 24
$NaHCO_3$, 1.25 NaH_2PO_4, 2 MgCl, 2 $CaCl_2$, and 10 glucose with an osmolarity of 300-310 mOsm.
Slices were incubated at 32°C in the holding chamber for 30 minutes, then held at room
temperature until recording. Slices were transferred to an upright microscope (Zeiss Stemi 2000-
C) and interface chamber, perfused at 2 ml/min with 32°C-34°C oxygenated 2 mM $MgCl/CaCl_2$
ACSF containing 50 µM of APV. 3-5 MOhm tip electrodes were filled with 2 mM $MgCl/CaCl_2$
ACSF. A Multiclamp 700B amplifier (Molecular Devices) was used to make LFP recordings,
filtered at 3 kHz (Bessel filter), and digitized at 10-25 kHz (Axon Digidata 1550, Molecular
Devices) to pClamp 10.6 software; Molecular Devices). LFP recordings were conducted in
hippocampal CA3 pyramidal cell layer. Spontaneous sharp wave-ripples (SWRs) were recorded in
10-minute epochs for baseline, bath-applied drug (1 µM CGP-78608 or 40 µM EU1180-490), and
washout conditions. Recording traces were filtered at 1 Hz (high pass). The frequency of SWRs
was determined by the number of SWRs detected during a 2-minute period using Clampfit 10
software (Molecular Devices). Traces were filtered 150-250 Hz to confirm ripples.
695
Post hoc morphological Reconstruction

Slices containing biocytin filled cells were fixed in 4 % paraformaldehyde overnight at 4 °C. Then
fixed slices were washed in PBS, permeabilized and incubated with Alexa-488, Alexa-555, or
Alexa-633 conjugated streptavidin overnight (Molecular Probes). After incubation, slices were
washed multiple times and then were resected to 70µm using a freezing microtome (Microm,
Whaltham, MA). Slices were mounted on glass slides (Fisherbrand Superfrost Plus, Fisher
Scientific) using Mowiol as a mounting medium. Cells were visualized using epifluorescence
microscopy (AX70, Olympus). Confocal images of labelled cells were attained on a Zeiss LSM 780
microscope (NICHD Imaging and Microscopy Core).
705
Immunohistochemistry (IHC)

Mice of different ages (P6, 13, and 6 weeks) were deeply anesthetized with isoflurane and
transcardially perfused with 4% paraformaldehyde (PFA) in phosphate buffered saline (PBS) pH
7.4. Brains were removed, post-fixed 1 hour in the same fixative at 4°C and transferred to 30%
sucrose in PBS at 4°C for at least 24 hours before sectioning. 30 µm coronal slices were cut and
stored at -20°C in cryoprotectant of PBS, 30% ethylene glycol and 30% glycerol in PBS.
Free-floating brain sections were briefly washed in PBS to remove cryoprotectant and then
incubated with blocking solution containing 4% normal goat serum (NGS), 1% BSA, 0.1-0.3%
Triton X-100 in PBS for 2 hours. Then sections were incubated with primary antibody in 1% NGS,
1% BSA, 0.1-0.3% Triton X-100 in PBS overnight at 4ºC. The following primary antibodies were
used: rabbit anti-GFP (Synaptic Systems 132002, 1:1000), chicken anti-GFP (Aves labs GFP-1020,
1:1000), rabbit anti-SST (Invitrogen PA585759, 1:500), rabbit anti-nNOS (Abcam ab5586, 1:200).

Sections were then washed extensively with PBS and incubated with fluorophore conjugated
secondary antibody in 1% BSA, 0.1% Triton X-100 in PBS for 2-3 hours at room temperature
(overnight at 4°C for dendritic analysis). Secondary antibodies used were: anti-rabbit Alexa 488
(Invitrogen A11008 1:750), anti-chicken Alexa 488 (Invitrogen A11039, 1:750) and anti-rabbit
Alexa 555 (Invitrogen A21428, 1:500). After PBS washes, sections were counterstained with
DAPI, mounted onto Superfrost Plus slides, air-dried, and coverslipped with fluorescence
mounting medium.
725
In situ Hybridization (ISH) RNAscope

To prepare snap frozen tissue for ISH, freshly dissected macaque hippocampal tissues were
submerged into 2-methylbutane that was pre-chilled on a dry ice/ethanol bath for 5 minutes.
Tissue was removed, wrapped in foil, and stored at -80 °C. 10 µm frozen sections were made
using a Leica Cryostat, mounted on Superfrost Plus slides (Fisher Scientific Cat#12-550-15) and
stored at -80 °C.
Target probes were designed and manufactured by Advanced Cell Diagnostic (ACD): mmu-
SST(Cat#461681), mmu-G*rin3a*-C2(Cat#1225361-C2). ISH was performed following the
RNAscope Multiplex Fluorescent v2 Assay instructions provided by ACD. Briefly, frozen thin
sections were post-fixed in 4% PFA and dehydrated sequentially in 50%, 70% and 100% ethanol.
After H_2O_2 and Protease IV treatment, sections were incubated with probes at 40 °C for 2
hours. Probed signals were detected using RNAscope Multiplex Detection v2 kit (ACD, Cat#
323110). Opal 520 (AKOYA, Cat# OP-001001) and Opal 650 (AKOYA, Cat# OP-001006) were used
to detect C1 (sst) and C2 (*Grin3a*) probes, respectively. After DAPI staining, sections were
covered using ProLong Gold Antifade Mountant (ThermoFisher Scientific, Cat# P10144) and
cured in darkness before imaging.
RNAscope images were acquired on a SLIDEVIEW VS200 slide scanner (Olympus) mounted with
an ORCA-Fusion Digital Camera (HAMAMATSU). Hippocampal regions were scanned at 20x and
40x for counting and signal colocalization analysis.
745
RiboTag

This assay was performed as previously described[Sanz2009, Mahadevan2020,
Mahadevan2021]. RNA bound with anti-HA immunoprecipitates and RNA from bulk tissue were
purified using RNeasy Plus Micro Kit (Qiagen, Cat# 74034) and the quality of RNA was measured
using RNA 6000 Pico kit (Agilent, Cat# 5067-1513) and 2100 Bioanalyzer system (Agilent, Cat#
G2939BA). cDNA libraries were constructed from 250 pg RNA using the SMARTer Stranded Total
RNA-Seq. Kit v2 (Takara Bio, Cat# 634411) from samples with RNA Integrity Numbers > 6.
Sequencing of the libraries were performed on the Illumina HiSeq 2500, at 50 million 2 × 100 bp
paired-end reads per sample. ~75% of reads were uniquely mapped to genomic features in the
reference genome. Bioconductor package DESeq2[Love2014] was used to identify differentially
expressed genes (DEG). This package allows for statistical determination of DEGs using a
negative binomial distribution model. The resulting values were then adjusted using the
Benjamini and Hochberg's method for controlling the false discovery rate.
759

Quantification and statistical analysis

Data analysis and visualization was conducted by homemade MATLAB codes. GDP-I was
detected using simple threshold method after gaussian filtering, and G-PSC detection was
achieved by template event detection in Clampfit 10.6. Paired t-test was selected for change of
activity by chemical or light stimuli, and Wilcoxon rank sum test was used for comparison
between different conditions (*: 0.01 to 0.05, **: 0.001 to 0.01, ***: < 0.001). Summary data
are presented as mean ± SEM.
766

Multi-institute single-cell transcriptome analyses of GABAergic interneurons

Single-nucleus (sn) / single-cell (sc) transcriptomes from mouse cortical and hippocampal
regions were downloaded from studies spanning research institutes (Allen Institute:
[Hodge2019, Yao2021], Harvard Medical School: [Krienen2020], Karolinska Institute:
[Harris2018], NIH: [Mahadevan2021]). HDF5/raw files from these sources were downloaded
and converted into Seurat v3-compatible format using custom scripts in R package as described
earlier [Chittajallu2020, Ekins2020]. Basic integration, processing and visualization of the scRNA-
seq data were performed with the Seurat v3 [Butler2018, Stuart2019] in R environment using R
Studio. Briefly, data was preprocessed by removing cells with less than 200 detectable genes
and with reads mapping to mitochondrial genes comprising more than 0.05% of total reads.
Gene expression counts were then log normalized and the resulting expression matrix was
scaled to 10,000 transcripts per cell. Next, the expression values of each gene across all the cells
(z-score transformation) were utilized for dimensionality reduction. These were performed
using the NormalizeData() and ScaleData() functions in Seurat with default parameters. Next,
we identified the top 2000 genes that exhibited high variability across cells and therefore
represent unique features of the cell clusters, for downstream analysis using the
FindVariableGenes() function, by applying selection.method = "vst" feature. Next, principal
component analysis (PCA) was performed, and the top 20 principal components were stored.
We clustered the cells using the FindClusters() with a clustering resolution set to 2.0. Using the
FindAllMarkers() function at cutoff of genes expressed in 25% of cells, at a logfc.threshold =
0.25, we identified the genes that are expressed in each cluster. For each cell type we used the
well-established marker genes that uniquely established the cell identity, as previously
described in the respective studies. Subsetting only the Gad1 clusters expressing a combination
of Lhx6, Pvalb, Sst, Id2 were annotated as MGE-derived GABAergic interneurons, and a
combination of Lhx6-negative, Sncg, Vip, Calb2, Meis2, Cnr1, Ntng1, Ndnf were annotated as
CGE-derived GABAergic interneurons. The Neurogliaform interneurons that are
Lhx6$^+$/Lamp5$^+$/Id2$^+$ are annotated as MGE-derived (NGFC.M); and those that are Lhx6$^-$
/Lamp5$^+$/Id2$^+$/Ndnf$^+$ are annotated as CGE-derived (NGFC.C). In total we used 86,852 GABA-
subset sc-transcriptomes in this study. Expression levels of various GABAergic inteneuron
positive control genes and expression of *Grin3a* across the GABAergic interneuron clusters are
plotted using DotPlot function in Seurat [Figure 2A].
799

Analyses of publicly available transcriptomic data

For cross-species comparison of *GRIN1* and *GRIN3A* expression, we utilized a publicly available single-nucleus RNA sequencing (snRNA-seq) dataset comparing across neocortical INs in mouse, marmoset, macaque, and human [Krienen2020]. We probed the mean expression across four neocortical IN subpopulations (PV, SST, ID2, VIP) with a gene panel including *GRIN1*, *GRIN3A*, and distinguishing markers *PVALB*, *SST*, *LAMP5*, and *VIP*.

For further analysis into human single cell expression, we utilized the Allen Institute whole human brain dataset of the single-cell transcriptomes of ~3.4 million cells across the entire human brain [Siletti2023]. To probe our genes and regions of interest, we adapted Python notebooks provided for accessing the Allen Brain Cell (ABC) Atlas (https://alleninstitute.github.io/abc_atlas_access/intro.html). The log2-normalized expression matrices were downloaded and filtered through several criteria to reach the cell types of interest. First, only neurons were considered (~2.5m cells), which were limited to the superclusters "MGE interneuron", "CGE interneuron", "LAMP5-LHX6 and Chandelier", "Hippocampal CA1-3", "Hippocampal CA4", and "Hippocampal dentate gyrus" (~648k cells), and finally to the hippocampal anatomical division (~204k cells). Analysis of PNs was limited to the supercluster level, while INs were further analyzed down to the cluster level to meaningfully group clusters into identifiable interneuron subpopulations (e.g. PV, SST, LAMP5). We calculated the mean log2-normalized expression values across clusters within this subset of cells for a gene panel including: *GAD1*, *GAD2*, *SLC17A7* (VGluT1), *SLC17A6* (VGluT2), *LHX6*, *PROX1*, *PVALB*, *SST*, *VIP*, *LAMP5*, *CCK*, *SNCG*. Using a mean threshold = 2, MGE-PV clusters were defined as those expressing *LHX6+*, *PVALB+* (MGE_261/263, LLC_264/266, 1480 cells), MGE-SST clusters as *LHX6+*, *SST+* (MGE_263/239-254/258, LLC_271, 12935 cells), MGE-LAMP5 as *LHX6+*, *LAMP5+* (LLC_266-275, 7663 cells), CGE-LAMP as CGE *LAMP5+* (CGE_286-288, 6497 cells), CGE-SNCG as CGE *SNCG+* (CGE_276/277/279/281/285/291, 6604 cells), and CGE-VIP as CGE *VIP+* (CGE_276/280/281/284/285/289-296, 16007 cells). We next downloaded the single-cell gene expression across these defined groups for a gene panel including *GRIN1* and *GRIN3A* to create violin and dot plots via Graphpad Prism. All adapted Python notebooks are available on Github (https://github.com/acaccavano/ABC_Atlas_Kim2025).

References

Anastasiades, P. G., Marques-Smith, A., Lyngholm, D., Lickiss, T., Raffiq, S., Katzel, D., Miesenbock, G. and Butt, S. J. (2016) GABAergic interneurons form transient layer-specific circuits in early postnatal neocortex. Nat Commun **7**: 10584.
Awobuluyi, M., Yang, J., Ye, Y., Chatterton, J. E., Godzik, A., Lipton, S. A. and Zhang, D. (2007) Subunit-specific roles of glycine-binding domains in activation of NR1/NR3 N-methyl-D-aspartate receptors. Mol Pharmacol **71**(1): 112-122.
Beesley, S., Sullenberger, T., Lee, C. and Kumar, S. S. (2022) GluN3 subunit expression correlates with increased vulnerability of hippocampus and entorhinal cortex to neurodegeneration in a model of temporal lobe epilepsy. J Neurophysiol **127**(6): 1496-1510.
Behrens, C. J., van den Boom, L. P., de Hoz, L., Friedman, A. and Heinemann, U. (2005) Induction of sharp wave-ripple complexes in vitro and reorganization of hippocampal networks. Nat Neurosci **8**(11): 1560-1567.

Ben-Ari, Y., Cherubini, E., Corradetti, R. and Gaiarsa, J. L. (1989) Giant synaptic potentials in immature rat CA3 hippocampal neurones. J Physiol **416**: 303-325.

Ben-Ari, Y. (2002) Excitatory actions of gaba during development: the nature of the nurture. Nat Rev Neurosci **3**(9): 728-739.

Ben-Ari, Y., Gaiarsa, J. L., Tyzio, R. and Khazipov, R. (2007) GABA: a pioneer transmitter that excites immature neurons and generates primitive oscillations. Physiol Rev **87**(4): 1215-1284.

Bezaire, M. J. and Soltesz, I. (2013) Quantitative assessment of CA1 local circuits: knowledge base for interneuron-pyramidal cell connectivity. Hippocampus **23**(9): 751-785.

Bocchio, M., Gouny, C., Angulo-Garcia, D., Toulat, T., Tressard, T., Quiroli, E., Baude, A. and Cossart, R. (2020) Hippocampal hub neurons maintain distinct connectivity throughout their lifetime. Nat Commun **11**(1): 4559.

Bossi, S., Dhanasobhon, D., Ellis-Davies, G. C. R., Frontera, J., de Brito Van Velze, M., Lourenco, J., Murillo, A., Lujan, R., Casado, M., Perez-Otano, I., Bacci, A., Popa, D., Paoletti, P. and Rebola, N. (2022) GluN3A excitatory glycine receptors control adult cortical and amygdalar circuits. Neuron **110**(15): 2438-2454 e2438.

Bossi, S., Pizzamiglio, L. and Paoletti, P. (2023) Excitatory GluN1/GluN3A glycine receptors (eGlyRs) in brain signaling. Trends Neurosci **46**(8): 667-681.

Brody, S. A., Nakanishi, N., Tu, S., Lipton, S. A. and Geyer, M. A. (2005) A developmental influence of the N-methyl-D-aspartate receptor NR3A subunit on prepulse inhibition of startle. Biol Psychiatry **57**(10): 1147-1152.

Butler, A., Hoffman, P., Smibert, P., Papalexi, E. and Satija, R. (2018) Integrating single-cell transcriptomic data across different conditions, technologies, and species. Nat Biotechnol **36**(5): 411-420.

Buzsáki, G. (2015) Hippocampal sharp wave-ripple: A cognitive biomarker for episodic memory and planning. Hippocampus **25**(10): 1073-1188.

Caccavano, A. P., Vlachos, A., McLean, N., Kimmel, S., Kim, J. H., Vargish, G., Mahadevan, V., Hewitt, L., Rossi, A. M., Spineux, I., Wu, S. J., Furlanis, E., Dai, M., Leyva Garcia, B., Wang, Y., Chittajallu, R., London, E., Yuan, X., Hunt, S., Abebe, D., Eldridge, M. A. G., Cummins, A. C., Hines, B. E., Plotnikova, A., Mohanty, A., Averbeck, B. B., Zaghloul, K. A., Dimidschstein, J., Fishell, G., Pelkey, K. A. and McBain, C. J. (2025) Divergent opioid-mediated suppression of inhibition between hippocampus and neocortex across species and development. Neuron **113**(11): 1805-1822 e1807.

Chatterton, J. E., Awobuluyi, M., Premkumar, L. S., Takahashi, H., Talantova, M., Shin, Y., Cui, J., Tu, S., Sevarino, K. A., Nakanishi, N., Tong, G., Lipton, S. A. and Zhang, D. (2002) Excitatory glycine receptors containing the NR3 family of NMDA receptor subunits. Nature **415**(6873): 793-798.

Chen, B. C., Legant, W. R., Wang, K., Shao, L., Milkie, D. E., Davidson, M. W., Janetopoulos, C., Wu, X. F. S., Hammer, J. A., Liu, Z., English, B. P., Mimori-Kiyosue, Y., Romero, D. P., Ritter, A. T., Lippincott-Schwartz, J., Fritz-Laylin, L., Mullins, R. D., Mitchell, D. M., Bembenek, J. N., Reymann, A. C., Bohme, R., Grill, S. W., Wang, J. T., Seydoux, G., Tulu, U. S., Kiehart, D. P. and Betzig, E. (2014) Lattice light-sheet microscopy: Imaging molecules to embryos at high spatiotemporal resolution. Science **346**(6208): 1257998.

Chittajallu, R., Auville, K., Mahadevan, V., Lai, M., Hunt, S., Calvigioni, D., Pelkey, K. A., Zaghloul,
K. A. and McBain, C. J. (2020) Activity-dependent tuning of intrinsic excitability in mouse and
human neurogliaform cells. Elife **9**.

Ciabarra, A. M., Sullivan, J. M., Gahn, L. G., Pecht, G., Heinemann, S. and Sevarino, K. A. (1995)
Cloning and characterization of chi-1: a developmentally regulated member of a novel class of
the ionotropic glutamate receptor family. J Neurosci **15**(10): 6498-6508.

Colgin, L. L., Jia, Y., Sabatier, J. M. and Lynch, G. (2005) Blockade of NMDA receptors enhances
spontaneous sharp waves in rat hippocampal slices. Neurosci Lett **385**(1): 46-51.

Cossart, R. and Khazipov, R. (2022) How development sculpts hippocampal circuits and function.
Physiol Rev **102**(1): 343-378.

Cossart, R. and Garel, S. (2022) Step by step: cells with multiple functions in cortical circuit
assembly. Nat Rev Neurosci **23**(7): 395-410.

Crawley, O., Conde-Dusman, M. J. and Perez-Otano, I. (2022) GluN3A NMDA receptor subunits:
more enigmatic than ever? J Physiol **600**(2): 261-276.

Cummings, K. A., Belin, S. and Popescu, G. K. (2017) Residues in the GluN1 C-terminal domain
control kinetics and pharmacology of GluN1/GluN3A N-methyl-d-aspartate receptors.
Neuropharmacology **119**: 40-47.

Dard, R. F., Leprince, E., Denis, J., Rao Balappa, S., Suchkov, D., Boyce, R., Lopez, C., Giorgi-Kurz,
M., Szwagier, T., Dumont, T., Rouault, H., Minlebaev, M., Baude, A., Cossart, R. and Picardo, M.
A. (2022) The rapid developmental rise of somatic inhibition disengages hippocampal dynamics
from self-motion. Elife **11**.

Das, S., Sasaki, Y. F., Rothe, T., Premkumar, L. S., Takasu, M., Crandall, J. E., Dikkes, P., Conner, D.
A., Rayudu, P. V., Cheung, W., Chen, H. S., Lipton, S. A. and Nakanishi, N. (1998) Increased NMDA
current and spine density in mice lacking the NMDA receptor subunit NR3A. Nature **393**(6683):
377-381.

Ekins, T. G., Mahadevan, V., Zhang, Y., D'Amour, J. A., Akgul, G., Petros, T. J. and McBain, C. J.
(2020) Emergence of non-canonical parvalbumin-containing interneurons in hippocampus of a
murine model of type I lissencephaly. Elife **9**.

Flossmann, T., Kaas, T., Rahmati, V., Kiebel, S. J., Witte, O. W., Holthoff, K. and Kirmse, K. (2019)
Somatostatin Interneurons Promote Neuronal Synchrony in the Neonatal Hippocampus. Cell
Rep **26**(12): 3173-3182 e3175.

Fredericks, J. M., Dash, K. E., Jaskot, E. M., Bennett, T. W., Lerchner, W., Dold, G., Ide, D.,
Cummins, A. C., Der Minassian, V. H., Turchi, J. N., Richmond, B. J. and Eldridge, M. A. G. (2020)
Methods for mechanical delivery of viral vectors into rhesus monkey brain. J Neurosci Methods
339: 108730.

Fuentealba, P., Begum, R., Capogna, M., Jinno, S., Marton, L. F., Csicsvari, J., Thomson, A.,
Somogyi, P. and Klausberger, T. (2008) Ivy cells: a population of nitric-oxide-producing, slow-
spiking GABAergic neurons and their involvement in hippocampal network activity. Neuron
57(6): 917-929.

Furlanis, E., Dai, M., Leyva Garcia, B., Tran, T., Vergara, J., Pereira, A., Gorissen, B. L., Wills, S.,
Vlachos, A., Hairston, A., Dwivedi, D., Du, S., McMahon, J., Huang, S., Morabito, A., Vazquez, A.,
Kim, S., Lee, A. T., Chang, E. F., Razzaq, T., Qazi, A., Vargish, G., Yuan, X., Caccavano, A., Hunt, S.,
Chittajallu, R., McLean, N., Hewitt, L., Paranzino, E., Rice, H., Cummins, A. C., Plotnikova, A.,
Mohanty, A., Tangen, A. C., Shin, J. H., Azadi, R., Eldridge, M. A. G., Alvarez, V. A., Averbeck, B. B.,

Alyahyay, M., Vallejo, T. R., Soheib, M., Vattino, L. G., MacGregor, C. P., Chatain, C. P., Banks, E.,
Olah, V. J., Naskar, S., Hill, S., Liebergall, S., Badiani, R., Hyde, L., Hanley, E., Xu, Q., Allaway, K. C.,
Goldberg, E. M., Rowan, M. J. M., Nowakowski, T. J., Lee, S., Favuzzi, E., Kaeser, P. S., Sjulson, L.,
Batista-Brito, R., Takesian, A. E., Ibrahim, L. A., Iqbal, A., Pelkey, K. A., McBain, C. J.,
Dimidschstein, J., Fishell, G. and Wang, Y. (2025) An enhancer-AAV toolbox to target and
manipulate distinct interneuron subtypes. Neuron **113**(10): 1525-1547 e1515.
Gallinat, J., Gotz, T., Kalus, P., Bajbouj, M., Sander, T. and Winterer, G. (2007) Genetic variations
of the NR3A subunit of the NMDA receptor modulate prefrontal cerebral activity in humans. J
Cogn Neurosci **19**(1): 59-68.
Girardeau, G., Benchenane, K., Wiener, S. I., Buzsaki, G. and Zugaro, M. B. (2009) Selective
suppression of hippocampal ripples impairs spatial memory. Nat Neurosci **12**(10): 1222-1223.
Gonzalez-Gonzalez, I. M., Gray, J. A., Ferreira, J., Conde-Dusman, M. J., Bouchet, D., Perez-Otano,
I. and Groc, L. (2023) GluN3A subunit tunes NMDA receptor synaptic trafficking and content
during postnatal brain development. Cell Rep **42**(5): 112477.
Gouwens, N. W., Sorensen, S. A., Baftizadeh, F., Budzillo, A., Lee, B. R., Jarsky, T., Alfiler, L., Baker,
K., Barkan, E., Berry, K., Bertagnolli, D., Bickley, K., Bomben, J., Braun, T., Brouner, K., Casper, T.,
Crichton, K., Daigle, T. L., Dalley, R., de Frates, R. A., Dee, N., Desta, T., Lee, S. D., Dotson, N.,
Egdorf, T., Ellingwood, L., Enstrom, R., Esposito, L., Farrell, C., Feng, D., Fong, O., Gala, R., Gamlin,
C., Gary, A., Glandon, A., Goldy, J., Gorham, M., Graybuck, L., Gu, H., Hadley, K., Hawrylycz, M. J.,
Henry, A. M., Hill, D., Hupp, M., Kebede, S., Kim, T. K., Kim, L., Kroll, M., Lee, C., Link, K. E.,
Mallory, M., Mann, R., Maxwell, M., McGraw, M., McMillen, D., Mukora, A., Ng, L., Ng, L., Ngo,
K., Nicovich, P. R., Oldre, A., Park, D., Peng, H., Penn, O., Pham, T., Pom, A., Popovic, Z.,
Potekhina, L., Rajanbabu, R., Ransford, S., Reid, D., Rimorin, C., Robertson, M., Ronellenfitch, K.,
Ruiz, A., Sandman, D., Smith, K., Sulc, J., Sunkin, S. M., Szafer, A., Tieu, M., Torkelson, A., Trinh, J.,
Tung, H., Wakeman, W., Ward, K., Williams, G., Zhou, Z., Ting, J. T., Arkhipov, A., Sumbul, U.,
Lein, E. S., Koch, C., Yao, Z., Tasic, B., Berg, J., Murphy, G. J. and Zeng, H. (2020) Integrated
Morphoelectric and Transcriptomic Classification of Cortical GABAergic Cells. Cell **183**(4): 935-
953 e919.
Grand, T., Abi Gerges, S., David, M., Diana, M. A. and Paoletti, P. (2018) Unmasking
GluN1/GluN3A excitatory glycine NMDA receptors. Nat Commun **9**(1): 4769.
Hajos, N., Ellender, T. J., Zemankovics, R., Mann, E. O., Exley, R., Cragg, S. J., Freund, T. F. and
Paulsen, O. (2009) Maintaining network activity in submerged hippocampal slices: importance
of oxygen supply. Eur J Neurosci **29**(2): 319-327.
Hajos, N. (2021) Interneuron Types and Their Circuits in the Basolateral Amygdala. Front Neural
Circuits **15**: 687257.
Hansen, K. B., Wollmuth, L. P., Bowie, D., Furukawa, H., Menniti, F. S., Sobolevsky, A. I., Swanson,
G. T., Swanger, S. A., Greger, I. H., Nakagawa, T., McBain, C. J., Jayaraman, V., Low, C. M.,
Dell'Acqua, M. L., Diamond, J. S., Camp, C. R., Perszyk, R. E., Yuan, H. and Traynelis, S. F. (2021)
Structure, Function, and Pharmacology of Glutamate Receptor Ion Channels. Pharmacol Rev
73(4): 298-487.
Harris, K. D., Hochgerner, H., Skene, N. G., Magno, L., Katona, L., Bengtsson Gonzales, C.,
Somogyi, P., Kessaris, N., Linnarsson, S. and Hjerling-Leffler, J. (2018) Classes and continua of
hippocampal CA1 inhibitory neurons revealed by single-cell transcriptomics. PLoS Biol **16**(6):
e2006387.

Henson, M. A., Larsen, R. S., Lawson, S. N., Perez-Otano, I., Nakanishi, N., Lipton, S. A. and
Philpot, B. D. (2012) Genetic deletion of NR3A accelerates glutamatergic synapse maturation.
PLoS One **7**(8): e42327.
Hodge, R. D., Bakken, T. E., Miller, J. A., Smith, K. A., Barkan, E. R., Graybuck, L. T., Close, J. L.,
Long, B., Johansen, N., Penn, O., Yao, Z., Eggermont, J., Hollt, T., Levi, B. P., Shehata, S. I.,
Aevermann, B., Beller, A., Bertagnolli, D., Brouner, K., Casper, T., Cobbs, C., Dalley, R., Dee, N.,
Ding, S. L., Ellenbogen, R. G., Fong, O., Garren, E., Goldy, J., Gwinn, R. P., Hirschstein, D., Keene,
C. D., Keshk, M., Ko, A. L., Lathia, K., Mahfouz, A., Maltzer, Z., McGraw, M., Nguyen, T. N., Nyhus,
J., Ojemann, J. G., Oldre, A., Parry, S., Reynolds, S., Rimorin, C., Shapovalova, N. V.,
Somasundaram, S., Szafer, A., Thomsen, E. R., Tieu, M., Quon, G., Scheuermann, R. H., Yuste, R.,
Sunkin, S. M., Lelieveldt, B., Feng, D., Ng, L., Bernard, A., Hawrylycz, M., Phillips, J. W., Tasic, B.,
Zeng, H., Jones, A. R., Koch, C. and Lein, E. S. (2019) Conserved cell types with divergent features
in human versus mouse cortex. Nature **573**(7772): 61-68.
Huang, X., Chen, Y. Y., Shen, Y., Cao, X., Li, A., Liu, Q., Li, Z., Zhang, L. B., Dai, W., Tan, T., Arias-
Carrion, O., Xue, Y. X., Su, H. and Yuan, T. F. (2017) Methamphetamine abuse impairs motor
cortical plasticity and function. Mol Psychiatry **22**(9): 1274-1281.
Hurley, E. P., Mukherjee, B., Fang, L. Z., Barnes, J. R., Barron, J. C., Nafar, F., Hirasawa, M. and
Parsons, M. P. (2024) GluN3A and Excitatory Glycine Receptors in the Adult Hippocampus. J
Neurosci **44**(42).
Jacobs, L., Nguyen, T. T., McCallum, A., Epplin, M. P., Perszyk, R. E., Myers, S. J., Liu, K., Yang, Y.,
Westhuyzen, A. E. V. d., Nocilla, K. A., Allen, J., Ullman, E. Z., Liu, D., Du, Y., Yuan, H., Liotta, D. C.
and Traynelis, S. F. (2025) A Subunit-Selective Negative Allosteric Modulator of GluN1/GluN3A
Glycine-Activated Receptor. Mol Pharmacol(manuscript submitted).
Jadhav, S. P., Kemere, C., German, P. W. and Frank, L. M. (2012) Awake hippocampal sharp-wave
ripples support spatial memory. Science **336**(6087): 1454-1458.
Joo, H. R. and Frank, L. M. (2018) The hippocampal sharp wave-ripple in memory retrieval for
immediate use and consolidation. Nat Rev Neurosci **19**(12): 744-757.
Karagiannis, A., Gallopin, T., David, C., Battaglia, D., Geoffroy, H., Rossier, J., Hillman, E. M.,
Staiger, J. F. and Cauli, B. (2009) Classification of NPY-expressing neocortical interneurons. J
Neurosci **29**(11): 3642-3659.
Katona, L., Micklem, B., Borhegyi, Z., Swiejkowski, D. A., Valenti, O., Viney, T. J., Kotzadimitriou,
D., Klausberger, T. and Somogyi, P. (2017) Behavior-dependent activity patterns of GABAergic
long-range projecting neurons in the rat hippocampus. Hippocampus **27**(4): 359-377.
Kehoe, L. A., Bernardinelli, Y. and Muller, D. (2013) GluN3A: an NMDA receptor subunit with
exquisite properties and functions. Neural Plast **2013**: 145387.
Kehoe, L. A., Bellone, C., De Roo, M., Zandueta, A., Dey, P. N., Perez-Otano, I. and Muller, D.
(2014) GluN3A promotes dendritic spine pruning and destabilization during postnatal
development. J Neurosci **34**(28): 9213-9221.
Kim, J. H., Lee, H. J., Min, C. H. and Lee, K. J. (2015) Coherence resonance in bursting neural
networks. Phys. Rev. E **92**(4): 042701.
Klausberger, T. and Somogyi, P. (2008) Neuronal diversity and temporal dynamics: the unity of
hippocampal circuit operations. Science **321**(5885): 53-57.
Krienen, F. M., Goldman, M., Zhang, Q., R, C. H. D. R., Florio, M., Machold, R., Saunders, A.,
Levandowski, K., Zaniewski, H., Schuman, B., Wu, C., Lutservitz, A., Mullally, C. D., Reed, N., Bien,

E., Bortolin, L., Fernandez-Otero, M., Lin, J. D., Wysoker, A., Nemesh, J., Kulp, D., Burns, M.,
Tkachev, V., Smith, R., Walsh, C. A., Dimidschstein, J., Rudy, B., L, S. K., Berretta, S., Fishell, G.,
Feng, G. and McCarroll, S. A. (2020) Innovations present in the primate interneuron repertoire.
Nature **586**(7828): 262-269.
Kvist, T., Greenwood, J. R., Hansen, K. B., Traynelis, S. F. and Brauner-Osborne, H. (2013a)
Structure-based discovery of antagonists for GluN3-containing N-methyl-D-aspartate receptors.
Neuropharmacology **75**: 324-336.
Kvist, T., Steffensen, T. B., Greenwood, J. R., Mehrzad Tabrizi, F., Hansen, K. B., Gajhede, M.,
Pickering, D. S., Traynelis, S. F., Kastrup, J. S. and Brauner-Osborne, H. (2013b) Crystal structure
and pharmacological characterization of a novel N-methyl-D-aspartate (NMDA) receptor
antagonist at the GluN1 glycine binding site. J Biol Chem **288**(46): 33124-33135.
Lee, J. H., Wei, L., Deveau, T. C., Gu, X. and Yu, S. P. (2016) Expression of the NMDA receptor
subunit GluN3A (NR3A) in the olfactory system and its regulatory role on olfaction in the adult
mouse. Brain Struct Funct **221**(6): 3259-3273.
Love, M. I., Huber, W. and Anders, S. (2014) Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol **15**(12): 550.
Madisen, L., Zwingman, T. A., Sunkin, S. M., Oh, S. W., Zariwala, H. A., Gu, H., Ng, L. L., Palmiter,
R. D., Hawrylycz, M. J., Jones, A. R., Lein, E. S. and Zeng, H. (2010) A robust and high-throughput
Cre reporting and characterization system for the whole mouse brain. Nat Neurosci **13**(1): 133-
140.
Madry, C., Mesic, I., Bartholomaus, I., Nicke, A., Betz, H. and Laube, B. (2007) Principal role of
NR3 subunits in NR1/NR3 excitatory glycine receptor function. Biochem Biophys Res Commun
354(1): 102-108.
Mahadevan, V., Peltekian, A. and McBain, C. J. (2020) Translatome Analyses Using Conditional
Ribosomal Tagging in GABAergic Interneurons and Other Sparse Cell Types. Curr Protoc Neurosci
92(1): e93.
Mahadevan, V., Mitra, A., Zhang, Y., Yuan, X., Peltekian, A., Chittajallu, R., Esnault, C., Maric, D.,
Rhodes, C., Pelkey, K. A., Dale, R., Petros, T. J. and McBain, C. J. (2021) NMDARs Drive the
Expression of Neuropsychiatric Disorder Risk Genes Within GABAergic Interneuron Subtypes in
the Juvenile Brain. Front Mol Neurosci **14**: 712609.
Marco, S., Giralt, A., Petrovic, M. M., Pouladi, M. A., Martinez-Turrillas, R., Martinez-Hernandez,
J., Kaltenbach, L. S., Torres-Peraza, J., Graham, R. K., Watanabe, M., Lujan, R., Nakanishi, N.,
Lipton, S. A., Lo, D. C., Hayden, M. R., Alberch, J., Wesseling, J. F. and Perez-Otano, I. (2013)
Suppressing aberrant GluN3A expression rescues synaptic and behavioral impairments in
Huntington's disease models. Nat Med **19**(8): 1030-1038.
Marco, S., Murillo, A. and Perez-Otano, I. (2018) RNAi-Based GluN3A Silencing Prevents and
Reverses Disease Phenotypes Induced by Mutant huntingtin. Mol Ther **26**(8): 1965-1972.
Marques-Smith, A., Lyngholm, D., Kaufmann, A. K., Stacey, J. A., Hoerder-Suabedissen, A.,
Becker, E. B., Wilson, M. C., Molnar, Z. and Butt, S. J. (2016) A Transient Translaminar GABAergic
Interneuron Circuit Connects Thalamocortical Recipient Layers in Neonatal Somatosensory
Cortex. Neuron **89**(3): 536-549.
Matsuda, K., Kamiya, Y., Matsuda, S. and Yuzaki, M. (2002) Cloning and characterization of a
novel NMDA receptor subunit NR3B: a dominant subunit that reduces calcium permeability.
Brain Res Mol Brain Res **100**(1-2): 43-52.

Michalski, K. and Furukawa, H. (2024) Structure and function of GluN1-3A NMDA receptor excitatory glycine receptor channel. Sci Adv **10**(15): eadl5952.

Milstein, A. D., Bloss, E. B., Apostolides, P. F., Vaidya, S. P., Dilly, G. A., Zemelman, B. V. and Magee, J. C. (2015) Inhibitory Gating of Input Comparison in the CA1 Microcircuit. Neuron **87**(6): 1274-1289.

Modol, L., Sousa, V. H., Malvache, A., Tressard, T., Baude, A. and Cossart, R. (2017) Spatial Embryonic Origin Delineates GABAergic Hub Neurons Driving Network Dynamics in the Developing Entorhinal Cortex. Cereb Cortex **27**(9): 4649-4661.

Mohamad, O., Song, M., Wei, L. and Yu, S. P. (2013) Regulatory roles of the NMDA receptor GluN3A subunit in locomotion, pain perception and cognitive functions in adult mice. J Physiol **591**(1): 149-168.

Mueller, H. T. and Meador-Woodruff, J. H. (2004) NR3A NMDA receptor subunit mRNA expression in schizophrenia, depression and bipolar disorder. Schizophr Res **71**(2-3): 361-370.

Murillo, A., Navarro, A. I., Puelles, E., Zhang, Y., Petros, T. J. and Perez-Otano, I. (2021) Temporal Dynamics and Neuronal Specificity of Grin3a Expression in the Mouse Forebrain. Cereb Cortex **31**(4): 1914-1926.

Nguyen, T. T., Zhu, Z., Epplin, M. P., Summer, S. L., Kelsey A, N., Mohan, A., Yuan, H., Hansen, K. B., Liotta, D. C. and Traynelis, S. F. (2025) Selective Positive Allosteric Modulators of GluN1/GluN3A Receptors. (Manuscript in preparation).

Nicoll, R. A. (2017) A Brief History of Long-Term Potentiation. Neuron **93**(2): 281-290.

Oh, W. C., Lutzu, S., Castillo, P. E. and Kwon, H. B. (2016) De novo synaptogenesis induced by GABA in the developing mouse cortex. Science **353**(6303): 1037-1040.

Otsu, Y., Darcq, E., Pietrajtis, K., Matyas, F., Schwartz, E., Bessaih, T., Abi Gerges, S., Rousseau, C. V., Grand, T., Dieudonne, S., Paoletti, P., Acsady, L., Agulhon, C., Kieffer, B. L. and Diana, M. A. (2019) Control of aversion by glycine-gated GluN1/GluN3A NMDA receptors in the adult medial habenula. Science **366**(6462): 250-254.

Overstreet-Wadiche, L. and McBain, C. J. (2015) Neurogliaform cells in cortical circuits. Nat Rev Neurosci **16**(8): 458-468.

Papenberg, G., Li, S. C., Nagel, I. E., Nietfeld, W., Schjeide, B. M., Schroder, J., Bertram, L., Heekeren, H. R., Lindenberger, U. and Backman, L. (2014) Dopamine and glutamate receptor genes interactively influence episodic memory in old age. Neurobiol Aging **35**(5): 1213 e1213-1218.

Paul, A., Crow, M., Raudales, R., He, M., Gillis, J. and Huang, Z. J. (2017) Transcriptional Architecture of Synaptic Communication Delineates GABAergic Neuron Identity. Cell **171**(3): 522-539 e520.

Pelkey, K. A., Barksdale, E., Craig, M. T., Yuan, X., Sukumaran, M., Vargish, G. A., Mitchell, R. M., Wyeth, M. S., Petralia, R. S., Chittajallu, R., Karlsson, R. M., Cameron, H. A., Murata, Y., Colonnese, M. T., Worley, P. F. and McBain, C. J. (2015) Pentraxins coordinate excitatory synapse maturation and circuit integration of parvalbumin interneurons. Neuron **85**(6): 1257-1272.

Pelkey, K. A., Chittajallu, R., Craig, M. T., Tricoire, L., Wester, J. C. and McBain, C. J. (2017) Hippocampal GABAergic Inhibitory Interneurons. Physiol Rev **97**(4): 1619-1747.

Perez-Otano, I., Schulteis, C. T., Contractor, A., Lipton, S. A., Trimmer, J. S., Sucher, N. J. and Heinemann, S. F. (2001) Assembly with the NR1 subunit is required for surface expression of NR3A-containing NMDA receptors. J Neurosci **21**(4): 1228-1237.

Perez-Otano, I., Lujan, R., Tavalin, S. J., Plomann, M., Modregger, J., Liu, X. B., Jones, E. G., Heinemann, S. F., Lo, D. C. and Ehlers, M. D. (2006) Endocytosis and synaptic removal of NR3A-containing NMDA receptors by PACSIN1/syndapin1. Nat Neurosci **9**(5): 611-621.

Perez-Otano, I., Larsen, R. S. and Wesseling, J. F. (2016) Emerging roles of GluN3-containing NMDA receptors in the CNS. Nat Rev Neurosci **17**(10): 623-635.

Pfeffer, C. K., Xue, M., He, M., Huang, Z. J. and Scanziani, M. (2013) Inhibition of inhibition in visual cortex: the logic of connections between molecularly distinct interneurons. Nat Neurosci **16**(8): 1068-1076.

Picardo, M. A., Guigue, P., Bonifazi, P., Batista-Brito, R., Allene, C., Ribas, A., Fishell, G., Baude, A. and Cossart, R. (2011) Pioneer GABA cells comprise a subpopulation of hub neurons in the developing hippocampus. Neuron **71**(4): 695-709.

Pina-Crespo, J. C., Talantova, M., Micu, I., States, B., Chen, H. S., Tu, S., Nakanishi, N., Tong, G., Zhang, D., Heinemann, S. F., Zamponi, G. W., Stys, P. K. and Lipton, S. A. (2010) Excitatory glycine responses of CNS myelin mediated by NR1/NR3 "NMDA" receptor subunits. J Neurosci **30**(34): 11501-11505.

Pizzamiglio, L., Morice, E., Cardoso, C., Bossi, S., Mailhes-Hamon, C., Heimendahl, M. V., Girardeau, G. and Paoletti, P. (2025) Excitatory glycine receptors control ventral hippocampus synaptic plasticity and anxiety-related behaviors. Proc Natl Acad Sci U S A **122**(37): e2501118122.

Price, C. J., Cauli, B., Kovacs, E. R., Kulik, A., Lambolez, B., Shigemoto, R. and Capogna, M. (2005) Neurogliaform neurons form a novel inhibitory network in the hippocampal CA1 area. J Neurosci **25**(29): 6775-6786.

Roberts, A. C., Diez-Garcia, J., Rodriguiz, R. M., Lopez, I. P., Lujan, R., Martinez-Turrillas, R., Pico, E., Henson, M. A., Bernardo, D. R., Jarrett, T. M., Clendeninn, D. J., Lopez-Mascaraque, L., Feng, G., Lo, D. C., Wesseling, J. F., Wetsel, W. C., Philpot, B. D. and Perez-Otano, I. (2009) Downregulation of NR3A-containing NMDARs is required for synapse maturation and memory consolidation. Neuron **63**(3): 342-356.

Rouzbeh, N., Rau, A. R., Benton, A. J., Yi, F., Anderson, C. M., Johns, M. R., Jensen, L., Lotti, J. S., Holley, D. C. and Hansen, K. B. (2023) Allosteric modulation of GluN1/GluN3 NMDA receptors by GluN1-selective competitive antagonists. J Gen Physiol **155**(6).

Royer, S., Zemelman, B. V., Losonczy, A., Kim, J., Chance, F., Magee, J. C. and Buzsaki, G. (2012) Control of timing, rate and bursts of hippocampal place cells by dendritic and somatic inhibition. Nat Neurosci **15**(5): 769-775.

Sanz, E., Yang, L., Su, T., Morris, D. R., McKnight, G. S. and Amieux, P. S. (2009) Cell-type-specific isolation of ribosome-associated mRNA from complex tissues. Proc Natl Acad Sci U S A **106**(33): 13939-13944.

Sasaki, Y. F., Rothe, T., Premkumar, L. S., Das, S., Cui, J., Talantova, M. V., Wong, H. K., Gong, X., Chan, S. F., Zhang, D., Nakanishi, N., Sucher, N. J. and Lipton, S. A. (2002) Characterization and comparison of the NR3A subunit of the NMDA receptor in recombinant systems and primary cortical neurons. J Neurophysiol **87**(4): 2052-2063.

Sik, A., Ylinen, A., Penttonen, M. and Buzsaki, G. (1994) Inhibitory CA1-CA3-hilar region feedback in the hippocampus. Science **265**(5179): 1722-1724.

Siletti, K., Hodge, R., Mossi Albiach, A., Lee, K. W., Ding, S. L., Hu, L., Lonnerberg, P., Bakken, T., Casper, T., Clark, M., Dee, N., Gloe, J., Hirschstein, D., Shapovalova, N. V., Keene, C. D., Nyhus, J.,

Tung, H., Yanny, A. M., Arenas, E., Lein, E. S. and Linnarsson, S. (2023) Transcriptomic diversity of
cell types across the adult human brain. Science **382**(6667): eadd7046.
Somogyi, P., Katona, L., Klausberger, T., Lasztoczi, B. and Viney, T. J. (2014) Temporal
redistribution of inhibition over neuronal subcellular domains underlies state-dependent
rhythmic change of excitability in the hippocampus. Philos Trans R Soc Lond B Biol Sci
369(1635): 20120518.
Stark, E., Roux, L., Eichler, R., Senzai, Y., Royer, S. and Buzsaki, G. (2014) Pyramidal cell-
interneuron interactions underlie hippocampal ripple oscillations. Neuron **83**(2): 467-480.
Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W. M., 3rd, Hao, Y.,
Stoeckius, M., Smibert, P. and Satija, R. (2019) Comprehensive Integration of Single-Cell Data.
Cell **177**(7): 1888-1902 e1821.
Su, J., Basso, D., Iyer, S., Su, K., Wei, J. and Fox, M. A. (2020) Paracrine Role for Somatostatin
Interneurons in the Assembly of Perisomatic Inhibitory Synapses. J Neurosci **40**(39): 7421-7435.
Sucher, N. J., Akbarian, S., Chi, C. L., Leclerc, C. L., Awobuluyi, M., Deitcher, D. L., Wu, M. K.,
Yuan, J. P., Jones, E. G. and Lipton, S. A. (1995) Developmental and regional expression pattern
of a novel NMDA receptor-like subunit (NMDAR-L) in the rodent brain. J Neurosci **15**(10): 6509-
6520.
Sun, L., Margolis, F. L., Shipley, M. T. and Lidow, M. S. (1998) Identification of a long variant of
mRNA encoding the NR3 subunit of the NMDA receptor: its regional distribution and
developmental expression in the rat brain. FEBS Lett **441**(3): 392-396.
Sunagawa, G. A., Sumiyama, K., Ukai-Tadenuma, M., Perrin, D., Fujishima, H., Ukai, H.,
Nishimura, O., Shi, S., Ohno, R. I., Narumi, R., Shimizu, Y., Tone, D., Ode, K. L., Kuraku, S. and
Ueda, H. R. (2016) Mammalian Reverse Genetics without Crossing Reveals Nr3a as a Short-
Sleeper Gene. Cell Rep **14**(3): 662-677.
Szabo, G. G., Farrell, J. S., Dudok, B., Hou, W. H., Ortiz, A. L., Varga, C., Moolchand, P., Gulsever,
C. I., Gschwind, T., Dimidschstein, J., Capogna, M. and Soltesz, I. (2022) Ripple-selective
GABAergic projection cells in the hippocampus. Neuron **110**(12): 1959-1977 e1959.
Taniguchi, H., He, M., Wu, P., Kim, S., Paik, R., Sugino, K., Kvitsiani, D., Fu, Y., Lu, J., Lin, Y.,
Miyoshi, G., Shima, Y., Fishell, G., Nelson, S. B. and Huang, Z. J. (2011) A resource of Cre driver
lines for genetic targeting of GABAergic neurons in cerebral cortex. Neuron **71**(6): 995-1013.
Tasic, B. and Fishell, G. (2025) Exploring brain circuits, one cell type-or more- at a time. Neuron
113(10): 1469-1473.
Tong, G., Takahashi, H., Tu, S., Shin, Y., Talantova, M., Zago, W., Xia, P., Nie, Z., Goetz, T., Zhang,
D., Lipton, S. A. and Nakanishi, N. (2008) Modulation of NMDA receptor properties and synaptic
transmission by the NR3A subunit in mouse hippocampal and cerebrocortical neurons. J
Neurophysiol **99**(1): 122-132.
Tremblay, R., Lee, S. and Rudy, B. (2016) GABAergic Interneurons in the Neocortex: From Cellular
Properties to Circuits. Neuron **91**(2): 260-292.
Tricoire, L., Pelkey, K. A., Daw, M. I., Sousa, V. H., Miyoshi, G., Jeffries, B., Cauli, B., Fishell, G. and
McBain, C. J. (2010) Common origins of hippocampal Ivy and nitric oxide synthase expressing
neurogliaform cells. J Neurosci **30**(6): 2165-2176.
Tricoire, L., Pelkey, K. A., Erkkila, B. E., Jeffries, B. W., Yuan, X. and McBain, C. J. (2011) A
blueprint for the spatiotemporal origins of mouse hippocampal interneuron diversity. J Neurosci
31(30): 10948-10970.

Tsien, J. Z., Chen, D. F., Gerber, D., Tom, C., Mercer, E. H., Anderson, D. J., Mayford, M., Kandel, E.
R. and Tonegawa, S. (1996) Subregion- and cell type-restricted gene knockout in mouse brain.
Cell **87**(7): 1317-1326.
Tuncdemir, S. N., Wamsley, B., Stam, F. J., Osakada, F., Goulding, M., Callaway, E. M., Rudy, B.
and Fishell, G. (2016) Early Somatostatin Interneuron Connectivity Mediates the Maturation of
Deep Layer Cortical Circuits. Neuron **89**(3): 521-535.
Tzilivaki, A., Tukker, J. J., Maier, N., Poirazi, P., Sammons, R. P. and Schmitz, D. (2023)
Hippocampal GABAergic interneurons and memory. Neuron **111**(20): 3154-3175.
Varga, C., Golshani, P. and Soltesz, I. (2012) Frequency-invariant temporal ordering of
interneuronal discharges during hippocampal oscillations in awake mice. Proc Natl Acad Sci U S
A **109**(40): E2726-2734.
Vong, L., Ye, C., Yang, Z., Choi, B., Chua, S., Jr. and Lowell, B. B. (2011) Leptin action on
GABAergic neurons prevents obesity and reduces inhibitory tone to POMC neurons. Neuron
71(1): 142-154.
Wang, C. Z., Ma, J., Xu, Y. Q., Jiang, S. N., Chen, T. Q., Yuan, Z. L., Mao, X. Y., Zhang, S. Q., Liu, L. Y.,
Fu, Y. and Yu, Y. C. (2019) Early-generated interneurons regulate neuronal circuit formation
during early postnatal development. Elife **8**.
Wee, K. S., Tan, F. C., Cheong, Y. P., Khanna, S. and Low, C. M. (2016) Ontogenic Profile and
Synaptic Distribution of GluN3 Proteins in the Rat Brain and Hippocampal Neurons. Neurochem
Res **41**(1-2): 290-297.
Wester, J. C. and McBain, C. J. (2016) Interneurons Differentially Contribute to Spontaneous
Network Activity in the Developing Hippocampus Dependent on Their Embryonic Lineage. J
Neurosci **36**(9): 2646-2662.
Wong, H. K., Liu, X. B., Matos, M. F., Chan, S. F., Perez-Otano, I., Boysen, M., Cui, J., Nakanishi, N.,
Trimmer, J. S., Jones, E. G., Lipton, S. A. and Sucher, N. J. (2002) Temporal and regional
expression of NMDA receptor subunit NR3A in the mammalian brain. J Comp Neurol **450**(4):
303-317.
Xu, Q., Tam, M. and Anderson, S. A. (2008) Fate mapping Nkx2.1-lineage cells in the mouse
telencephalon. J Comp Neurol **506**(1): 16-29.
Yang, J., Wang, S., Yang, Z., Hodgkinson, C. A., Iarikova, P., Ma, J. Z., Payne, T. J., Goldman, D. and
Li, M. D. (2015) The contribution of rare and common variants in 30 genes to risk nicotine
dependence. Mol Psychiatry **20**(11): 1467-1478.
Yao, Y. and Mayer, M. L. (2006) Characterization of a soluble ligand binding domain of the NMDA
receptor regulatory subunit NR3A. J Neurosci **26**(17): 4559-4566.
Yao, Z., van Velthoven, C. T. J., Nguyen, T. N., Goldy, J., Sedeno-Cortes, A. E., Baftizadeh, F.,
Bertagnolli, D., Casper, T., Chiang, M., Crichton, K., Ding, S. L., Fong, O., Garren, E., Glandon, A.,
Gouwens, N. W., Gray, J., Graybuck, L. T., Hawrylycz, M. J., Hirschstein, D., Kroll, M., Lathia, K.,
Lee, C., Levi, B., McMillen, D., Mok, S., Pham, T., Ren, Q., Rimorin, C., Shapovalova, N., Sulc, J.,
Sunkin, S. M., Tieu, M., Torkelson, A., Tung, H., Ward, K., Dee, N., Smith, K. A., Tasic, B. and Zeng,
H. (2021) A taxonomy of transcriptomic cell types across the isocortex and hippocampal
formation. Cell **184**(12): 3222-3241 e3226.
Yao, Z., van Velthoven, C. T. J., Kunst, M., Zhang, M., McMillen, D., Lee, C., Jung, W., Goldy, J.,
Abdelhak, A., Aitken, M., Baker, K., Baker, P., Barkan, E., Bertagnolli, D., Bhandiwad, A., Bielstein,
C., Bishwakarma, P., Campos, J., Carey, D., Casper, T., Chakka, A. B., Chakrabarty, R., Chavan, S.,

Chen, M., Clark, M., Close, J., Crichton, K., Daniel, S., DiValentin, P., Dolbeare, T., Ellingwood, L.,
Fiabane, E., Fliss, T., Gee, J., Gerstenberger, J., Glandon, A., Gloe, J., Gould, J., Gray, J., Guilford,
N., Guzman, J., Hirschstein, D., Ho, W., Hooper, M., Huang, M., Hupp, M., Jin, K., Kroll, M.,
Lathia, K., Leon, A., Li, S., Long, B., Madigan, Z., Malloy, J., Malone, J., Maltzer, Z., Martin, N.,
McCue, R., McGinty, R., Mei, N., Melchor, J., Meyerdierks, E., Mollenkopf, T., Moonsman, S.,
Nguyen, T. N., Otto, S., Pham, T., Rimorin, C., Ruiz, A., Sanchez, R., Sawyer, L., Shapovalova, N.,
Shepard, N., Slaughterbeck, C., Sulc, J., Tieu, M., Torkelson, A., Tung, H., Valera Cuevas, N.,
Vance, S., Wadhwani, K., Ward, K., Levi, B., Farrell, C., Young, R., Staats, B., Wang, M. M.,
Thompson, C. L., Mufti, S., Pagan, C. M., Kruse, L., Dee, N., Sunkin, S. M., Esposito, L., Hawrylycz,
M. J., Waters, J., Ng, L., Smith, K., Tasic, B., Zhuang, X. and Zeng, H. (2023) A high-resolution
transcriptomic and spatial atlas of cell types in the whole mouse brain. Nature **624**(7991): 317-
332.
Yuan, T., Mameli, M., O'Connor, E. C., Dey, P. N., Verpelli, C., Sala, C., Perez-Otano, I., Luscher, C.
and Bellone, C. (2013) Expression of cocaine-evoked synaptic plasticity by GluN3A-containing
NMDA receptors. Neuron **80**(4): 1025-1038.
Zeilhofer, H. U., Studler, B., Arabadzisz, D., Schweizer, C., Ahmadi, S., Layh, B., Bosl, M. R. and
Fritschy, J. M. (2005) Glycinergic neurons expressing enhanced green fluorescent protein in
bacterial artificial chromosome transgenic mice. J Comp Neurol **482**(2): 123-141.
Zeisel, A., Munoz-Manchado, A. B., Codeluppi, S., Lonnerberg, P., La Manno, G., Jureus, A.,
Marques, S., Munguba, H., He, L., Betsholtz, C., Rolny, C., Castelo-Branco, G., Hjerling-Leffler, J.
and Linnarsson, S. (2015) Brain structure. Cell types in the mouse cortex and hippocampus
revealed by single-cell RNA-seq. Science **347**(6226): 1138-1142.
Zhang, X. Y., Liu, A. P., Ruan, D. Y. and Liu, J. (2002) Effect of developmental lead exposure on the
expression of specific NMDA receptor subunit mRNAs in the hippocampus of neonatal rats by
digoxigenin-labeled in situ hybridization histochemistry. Neurotoxicol Teratol **24**(2): 149-160.
Zhu, Z., Yi, F., Epplin, M. P., Liu, D., Summer, S. L., Mizu, R., Shaulsky, G., XiangWei, W., Tang, W.,
Burger, P. B., Menaldino, D. S., Myers, S. J., Liotta, D. C., Hansen, K. B., Yuan, H. and Traynelis, S.
F. (2020) Negative allosteric modulation of GluN1/GluN3 NMDA receptors. Neuropharmacology
176: 108117.
1268
1269

1270
Figure 1. Developmental expression of *Grin3a* transcripts and G-PSC/GDP modulation by
functional eGlyRs in early postnatal hippocampus. A. Developmental profile of *Grin3a* single
RNA-seq data hippocampus using Ribo-tag. **B.** Developmental change of GluN3a expression
images from mClover3-GluN1a mice demonstrating significant co-expression with somatostatin
(Sst). **C-D.** Single trace example of spontaneous GABAergic postsynaptic currents (G-PSCs) and
GDPs in CA3 pyramidal neurons (PNs) from WT (C), *Grin3a*-KO mouse brain slices (D). **E.** G-PSC
frequency change in (a) WT, (b) *Grin3a*-KO, and (c) comparison of G-PSC frequency change ratio
by CGP. **F.** Giant depolarizing potential (GDP) frequency change in WT (a), *Grin3a*-KO (b), and
normalized comparison of GDP frequency change by CGP in WT and *Grin3a*-KO mice (c). (WT: 62
cells/slices from 26 mice, P4.5-8.5), *Grin3a*-KO: 32 cells/slices from 13 mice, P5.5-12.5 *pre-
incubated in APV 100 μM.

1282
Figure 2. *Grin3a* expression and functional eGlyR-mediated currents in adult hippocampal INs.
A. Survey of *Grin3a* expression across major GABAergic interneuron classes (Sst, somatostatin-
expressing cells; PV, Parvalbumin-expressing cells; NGFC, neurogliaform cells; VIP, vaso-intestinal
peptide-expressing cells) from mice cortex and hippocampus. **B.** Co-expression of *Grin3a* with
Sst (left), and nNOS (right) respectively. **C.**(a) Whole cell patch-clamping an NPY-IN with glycine
puff. Glycine puff response of NPY-IN in (b) desensitization reducer (CGP), (c) allosteric
modulators positive (490), (d) negative (438), and (e) normalized glycine puff responses. **D.**(a) A
sample image of Sst-IN (OLM-IN) (b) Glycine puff response of Sst-IN in CGP, (c-d) allosteric
positive/negative modulators, respectively, (e) competitive eGlyR blocker (DCKA), and (f)
normalized glycine puff responses. **E.** Glycine puff responses of (a) Sst-IN in *Grin3a*-KO mice, (b)
Sst-IN and PN in Sst-*Grin3a*-cKO, (c) NPY-IN and PN in NPY-*Grin3a*-cKO, and (d) Sst-IN in Nkx2.1-
Grin1-cKO, and (e) average glycine puff responses.
1295

1296
1297 **Figure 3. eGlyRs are tonically liganded by endogenous glycine in hippocampal SST and NPY-**
1298 **NGF-INs. A-B.** eGlyR tonic current blocked by DCKA in NPY-INs, but no tonic current in Sst-INs. **C-**
1299 **D.** Holding current increase by desensitization reduction by CGP and/or EU1180-490 in NPY-INs
1300 and Sst-IN respectively. **E-F.** Firing activity increase by CGP or EU1180-490 in NPY-INs and Sst-INs
1301 respectively.

1302

Figure 4. Relative contributions of NPY-NGF and SST INs in eliciting early hippocampal inhibition and network synchrony. A. Whole-cell recording samples of CA3-PN in brain slices from VGAT-ArCh (top), NPY-ArCh (middle), and Sst-ArCh (bottom) mice with light-stimulation. **B.** Examples of G-PSC change by light-on in presence of CGP-78608 in CA3-PN in hippocampal slices from VGAT-ArCh (top), NPY-ArCh (middle), and Sst-ArCh (bottom) mice **C.** G-PSC frequency change ratio by light-on in VGAT-Arch, NPY-Arch, and Sst-Arch mouse hippocampus. **D.** G-PSC frequency change proportion; decrease (red), change less than +- 10 % (green), and increase (yellow). **E.** GDP frequency change ratio by light-on. **F.** Light-on evoked GDP frequency change proportion; decrease (red), no change (green), increase (yellow). **G.** Probability of light-off-evoked rebound GDP (rGDP). (VGAT-Arch: 20 cells/slices from 5 mice, P6.5-8.5, NPY-Arch: 39 cells/slices from 10 mice, P5.5-8.5, Sst-Arch: 30 cells/slices from 10 mice, P5.5-8.5).

1314
Figure 5. Recruitment of GABAergic tone and GDP generation is predominantly mediated by
activation of eGlyRs expressed on NPY-NGF-INs. A. An example recording of GDP with CGP-
78608 (upper), G-PSC frequency (middle), and GDP frequency (bottom). **B.** Enlarged traces in
baseline (upper)/CGP-Early(bottom) phases from A. **C.** G-PSC frequency change before,
immediate (early) and stable (late) phase of CGP application in (a) WT, (b) *Grin3a*-KO, (c) VGAT-
Grin3a-cKO, (d) NPY-*Grin3a*-cKO, (e) Sst-*Grin3a*-cKO mouse hippocampal slices, and (f)
normalized immediate G-PSC frequency change by CGP. **D.** GDP frequency change before, early
and late CGP application in (a) WT, (b) *Grin3a*-KO, (c) VGAT-*Grin3a*-cKO, (d) NPY-*Grin3a*-cKO, (e)
Sst-*Grin3a*-cKO mouse hippocampal slices, and (f) normalized immediate GDP frequency change
by CGP.

1325
1326 **Figure 6. Activation of eGlyRs shape adult hippocampal sharp wave ripple dynamics. A.** An
1327 example of sharp wave ripples. **B-D.** Sharp wave frequency change by CGP-78608/EU1180-490
1328 in WT, *Grin3a*-KO, and Sst-*Grin3a*-cKO, respectively. **E.** SPW change rate in drug normalized by
1329 frequency in ACSF. All recordings were performed with APV (50 µM) in bath solution.

1330

Figure 7. Evolutionary conservation of hippocampal interneuron eGlyR functional expression and role in network rhythmogenesis. A. Co-expression of Sst and *Grin3a* in a Macaque CA3 (stratum oriens) slice. **B.** Co-expression of Lamp5 and *Grin3a*. **C.** An example image of Sst-IN reported by pAAV-PHP.eB-BiSSTe4-dTomato (red) and post-hoc cell morphology recovery (green) in Macaque CA1 hippocampus. **D.** Glycine puff response of Sst-INs and PNs with CGP-78608 (middle) and CGP-78608 + EU1180-490 (right) in macaque hippocampus. **E.** SWRs frequency change by CGP-78608 / EU1180-490 in macaque (n=3, N=2) and marmoset (n=5, N=2). (n: number of slices, N: number of animals)

1339 **S1**

1340

***Grin3a* is absent in CA3 PNs and enriched in *Sst* and *Lamp5* expressing cells.** Publicly available
spatial transcriptomics MERFISH data from the Allen Brain Cell Atlas with imputed gene
expression [Yao2023]. Visualized gene expression restricted to the hippocampal formation,
showing (top) *Grin3a*, (middle) *Sst*, and (bottom) *Lamp5*. Insets highlight expression of putative
Sst-INs in stratum oriens (or) and Lamp5-INs in stratum lacunosum-moleculare (lm). Arrows
denote significant expression of Grin3A transcript in SST or Lamp5 expressing INs.

S2

1348
Experimental procedure employed to isolate strychnine-insensitive glycine evoked eGlyR
mediated depolarizing currents. (a) Example current traces from a CA1 Sst-IN with glycine puff
responses in presence of blockers and CGP-78608, filled circles in (b). Strychnine sensitive
eGlyR-mediated current prominently contributed the glycine response current in ACSF. (b)
Glycine puff currents in antagonists and CGP-78608. * STR 1 μM, APV 50 μM, DNQX 10 μM, BIC
10 uM and PTX 50 μM.

S3

1356

eGlyRs mediate functional responses in SST and NPY-NGFC INs throughout development. A.
Sample images of postnatal (top)/adulthood (bottom), NPY-IRES-Cre (left)/Sst-IRES-Cre (right)
mouse hippocampal slices. **B.** (left) Glycine puff response currents in NPY-INs, Sst-INs, and PNs
from postnatal and adulthood mice. (right) Glycine puff response charge transfer in NPY-INs, Sst-
INs, and PNs from postnatal and adulthood mice. **C.** Glycine puff responses in NPY -INs, Sst-INs
and PNs in different ages. Thick lines are average puff responses of cells and gray lines are
average puff responses of trials in each cell.

1364

1365 **S4**

1366
DCKA sensitivity of CGP78608-induced change holding current in hippocampal interneurons.
Shown are the time course (left) and traces (middle) from a representative sample NGFC
recording illustrating increased holding current (V_h=-70mV) in response to CGP78608 and
subsequent reduction of holding current upon DCKA application. At right, is the group data
summarizing the effects of CGP and subsequent DCKA application upon holding currents for Sst-
INs and NGFCs combined.

S5

1374

GRIN1 and GRIN3A expression cross species. A. Publicly-available snRNA-seq mean expression

in neocortical IN subpopulations PVALB (orange), SST (cyan), ID2/LAMP5 (red) and VIP (purple)

across mouse, marmoset, macaque, and human [Krienen2020], for (left) *GRIN1* and *GRIN3A* and

(right) distinguishing markers *PVALB*, *SST*, *LAMP5*, and *VIP*. **B.** Publicly-available snRNA-seq of

human hippocampal neurons [Siletti2023], with (left) *GRIN1* and *GRIN3A* expression in

glutamatergic (Glu) superclusters and grouped MGE and CGE clusters (see Methods), and (right)

dot plot of distinguishing marker genes and *GRIN1*, *GRIN3A*.

1382