Nanopore genome skimming with Illumina polishing yields highly accurate mitogenome sequences: a case study of *Niphargus* amphipods

Alice Salussolia[1], Ninon Lecoquierre[1], Fabio Stoch[1], and Jean-François Flot[1,2,3]

Abstract

With over 430 species currently described, the amphipod genus *Niphargus* Schiödte, 1849 is the most species-rich crustacean genus in subterranean waters. Previous phylogenetic studies of this genus have relied mainly on mitochondrial COI and nuclear 28S sequences, which do not resolve all the nodes in its phylogeny. As a first step towards a mitogenome-based phylogeny of niphargids, we present here the first complete mitogenome sequence of *Niphargus*. To obtain high-accuracy mitogenome sequences and annotations, genome skims of three individuals of the *Niphargus dolenianensis* Lorenzi, 1898 species complex were generated using both short, accurate reads (Illumina) and long, noisier reads (Nanopore). Whereas direct assembly of Illumina sequences yielded structurally incorrect mitogenome sequences, assembly of Nanopore reads produced highly accurate sequences that were corroborated by mapping of Illumina reads. Polishing of the Nanopore consensus using Illumina reads corrected a handful of errors at the homopolymer level. The resulting mitogenome sequences ranged from 14,956 to 15,199 bp and shared the same arrangement of 13 protein-coding genes, two ribosomal RNA genes, 22 transfer RNA genes, and a putative control region. Phylogenetic analyses based on protein-coding genes confirmed that the Niphargidae family is sister to Pseudoniphargidae, resolving their relationships with other amphipod families. This highlights the utility of mtDNA genome sequences for studying the evolution, diversification, and taxonomy of this groundwater genus.

Keywords: niphargids, Niphargidae, Amphipoda, Groundwater, mtDNA, Phylogeny

[1]Evolutionary, Biology & Ecology, Université libre de Bruxelles (ULB), Brussels, Belgium, [2]Interuniversity Institute of Bioinformatics in Brussels – (IB)², Brussels, Belgium, [3]Brussels Laboratory of the Universe – BLU, Brussels, Belgium

Correspondence
alice.salussolia@ulb.be, jean-francois.flot@ulb.be

Introduction

Groundwater constitutes an extensive and globally prevalent ecosystem that contains the majority of accessible freshwater resources (Saccò et al., 2024). Animals inhabiting groundwater play a crucial role in enhancing the biodiversity of freshwater ecosystems and are essential for processes such as nutrient cycling and bioturbation (Bardgett and Van Der Putten, 2014). Groundwater offers numerous ecosystem services, ranging from supporting terrestrial and surface freshwater ecosystems to supplying drinking water (Griebler and Avramov, 2015). Therefore, investigating groundwater animals is vital for advancing our understanding of groundwater community structure and processes, ecosystem services, and their reactions to environmental changes (Maurice and Bloomfield, 2012).

With a remarkable diversity of over 430 identified species (Horton et al., 2023), the amphipod genus *Niphargus* Schiödte, 1849 is the most species-rich crustacean genus in subterranean waters. The distribution of this genus spans the Iberian Peninsula to the west and Iran to the east, covering European regions south of the limits of Quaternary glaciations (Zagmajster et al., 2014). Studies of *Niphargus* are hampered by the lack of phylogenetic information on most species (except in a few well-studied areas) (Zagmajster et al., 2014) and by the frequent occurrence of complexes of cryptic and pseudocryptic species *Niphargus* (Stoch et al., 2022). To date, studies on the phylogenetics of the genus *Niphargus* have relied mainly on mitochondrial COI and nuclear 28S rRNA marker genes, and the resulting phylogenies remain poorly resolved, particularly for deeper nodes (Stoch et al., 2024b). Complete mitochondrial genome sequences of animals have been reported to yield well-resolved phylogenies (Cameron et al., 2007), but so far, not a single complete mitogenome sequence of *Niphargus* has been published.

To fill this gap and start investigating the potential of niphargid complete mitochondrial genome sequences to shed new light on amphipod phylogenetic relationships, we focused on the Alpine species *Niphargus dolenianensis* Lorenzi, 1898 and used genome skimming (Dodsworth, 2015) to assemble its complete mitogenome. We aimed to obtain a highly reliable *Niphargus dolenianensis* mitogenome for future comparative studies. For this purpose, we sequenced three different individuals using both Nanopore and Illumina platforms, and used Illumina reads to polish the assembly of Nanopore reads. Finally, to propose guidelines for future studies targeting niphargid mitogenomes, we tested whether Illumina-only and Nanopore-only genome skimming would have been sufficient to obtain high-quality mitogenome sequences required for downstream analyses.

Material and methods

DNA extraction and sequencing

Specimens of *Niphargus dolenianensis* Lorenzi, 1898 (Fig.1) were collected in 2024 from two springs and a brook using a hand-net. Information on the specimens used in the analyses, collection sites, and DNA vouchers is provided in Table 2. Fresh samples were immediately preserved in 96% EtOH and then stored at -20°C at the Evolutionary Biology & Ecology unit of the Université libre de Bruxelles (ULB), Belgium. DNA extraction was performed from one or two pereopods (depending on the size of the specimen) using Macherey-Nage's NucleoSpin Tissue kit, following the manufacturer's protocol. The eluted DNAs was stored at-20°C. Illumina sequencing (2 x 151 bp paired-end reads) was performed at BRIGHTcore (Brussels, Belgium) on an Illumina NovaSeq 6000 machine. The same three DNA extracts were also used for long-read Nanopore sequencing using the Rapid PCR Barcoding kit SQK-RPB114.24 on a PromethION R10.4.1 flow cell.

Mitochondrial genome annotation and analysis

Illumina data were trimmed using Skewer (Jiang et al., 2014) in paired-end mode. Nanopore data were basecalled using Guppy v3.8 (in super-high-quality mode) to generate fastq reads from the raw fast5. The Nanopore reads were filtered to keep only reads with an average quality

score of at least 14 using the script split_qscore.py available as part of the Buttery-eel package (Samarakoon et al., 2023) and assembled using Flye v2.9.6 (Kolmogorov et al., 2019), as well as with hifiasm v0.25.0-r726 (Cheng et al., 2021) for comparison. The resulting graphical fragment assembly (GFA) files were examined using Bandage (Wick et al., 2015) to identify and extract the circular contig of the mitochondrial genome. The first automatic Illumina polishing of the nanopore-assembled mitochondrial contig was performed using Polypolish (Wick and Holt, 2022). The resulting assemblies were checked closely for structural and base-level errors by mapping the Nanopore and Illumina reads on them using minimap2 (Li, 2018), converting the resulting SAM file into BAM and sorting it using SAMtools (Li et al., 2009), and then visualizing the resulting read pileups using Tablet (Milne et al., 2010). The few remaining errors detected in Tablet were corrected manually in the contig FASTA files using AliView (Larsson, 2014). For comparison, Illumina-only assemblies of the three mitochondrial genomes were attempted using the NOVOplasty assembler (Dierckxsens et al., 2017), using as seed the previously published COI sequence (Genbank accession KY706720) of a *Niphargus dolenianensis* individual (Eme et al., 2018).

A quick annotation of each Nanopore-assembled, Illumina-polished mitogenome sequence was performed using GeSeq (Tillich et al., 2017), with tRNAscan-SE v2.0.7 (Chan et al., 2021) to locate tRNAs; we used the available mitogenomes of *Pseudoniphargus stocki* and *Pseudoniphargus carpalis* (Stokkan et al., 2018) as "3rd Party References". A second annotation was conducted using MITOS2 (Bernt et al., 2013), specifying the invertebrate mitochondrial genetic code 5 and the reference dataset RefSeq63 Metazoa.

Considering that no *Niphargus* genome was available in GenBank, we manually checked and refined all annotations, including the stop codons of the protein-coding genes and the predicted secondary structure of tRNA genes (using the RNAfold tool available on the ViennaRNA web service; Gruber et al., 2015). The resulting annotated sequences are available in GenBank (accession numbers PV534570, PV534571 and PV534572).

The program mtSVG (available at https://github.com/odethier-ulb/mtSVG) was used to create a visual summary of the final annotated mitogenome of *N. dolenianensis* based on the three mitogenome sequences obtained in the present study. Read coverage depth graphs were created from BAM files using Grace-5.1.17 (Vaught, 1996). Nucleotide diversity analyses of the 13 protein-coding genes and two ribosomal RNA genes were conducted using the packages pegas (Paradis, 2010) and seqinR (Charif and Lobry, 2007) in RStudio 4.4.1.

Phylogenetic reconstruction

Together with our three Niphargidea mitogenome sequences, we included a selection of representative complete mitogenomes from the amphipod families Pseudoniphargidae (Stokkan et al., 2018), Gammaridae (Cormier et al., 2018; Macher et al., 2017; Mamos et al., 2021), Metacrangonyctidae (Bauzà-Ribot et al., 2009, 2012), Crangonyctidae (Benito et al., 2021), Talitridae (Kumar Patra et al., 2019), and Hyalellidae, plus two isopod species (Kilpert et al., 2012; Kilpert and Podsiadlowski, 2006) as outgroups (all GenBank accession numbers available in Table3. The translated protein sequences of all protein-coding genes were concatenated and used to build a maximum-likelihood (ML) phylogenetic tree using the program IQtree2 (Nguyen et al., 2015) with the invertebrate mitochondrial amino acid substitution model (MtInv) (Le et al., 2017), and node support was assessed using 50,000 ultrafast bootstrap replicates (Hoang et al., 2018). A linear representation of the order of protein-coding genes in each mitogenome was generated using the program mtSVG and these images were added next to the leaves of the ML phylogenetic tree.

Results

The final mitochondrial genome sequences of the three individuals of textitNiphargus dolenianensis ranged from 14,964 to 15,097 bp in length (Fig.4). Details of their annotation are reported in Table 5. Their global GC content was 24.6% for FS_24.009, 23.5% for FS_24.010 and 23.0% for FS_24.011.

The coverage depth obtained for the assembled mitogenomes generated in NovaSeq 6000 Illumina short reads ranged from approximately 40 to 250X; similarly, the coverage depth for Nanopore long reads ranged from 40 to 300X (Fig.6). Nanopore coverage depth profiles were less even than for Illumina, possibly because of biases caused by the PCR amplification step in the Rapid PCR Barcoding Kit protocol.

The predicted secondary structures of the tRNA genes are shown in Fig.7. Only for tRNA-Gly and tRNA-Leu2 were the sequences identical across all three individuals, whereas all other tRNA sequences had few base differences, which are highlighted in pink in Fig.7. These differences with mostly substitution (most often located outside of stem structures, except for a few mutations in stems that either did not disturb base pairing or, in the case of tRNA-Met, that was compensated by a mutation in the facing position). Of the 22 tRNAs annotated in the mitochondrial genome of *Niphargus dolenianensis*, 19 had the typical 3-arm structure, whereas tRNA-Ser1 and tRNA-Val lacked the dihydrouridine (DHU) arm and tRNA-Phe lacked the T-arm. The structure of tRNA-Val of FS_24.011 could not be predicted *de novo* from its sequence using the Vienna server, because of the extra pairing G-A pairing of this sequence compared with the other two. Therefore, we resorted to constraning its folding into the structure represented in Fig.7 using the results obtained from the two other individuals.

Nucleotide diversity (π) values for each protein-coding and ribosomal RNA gene are reported in Fig.8. The values range from 0.06 to 0.09 for the protein-coding genes, whereas it is only 0.03-0.04 for the two ribosomal RNA genes.

There were no significant differences between Flye and Hifiasm mitogenome assemblies obtained from the Nanopore long reads: for FS_24.009 the two were perfectly identical, whereas in the case of FS_24.010 and FS_24.011 there were only two differences between the Flye and Hifiasm mitogenome assemblies, in both short indels at the level of long homopolymers. These indels and several others (always located in log homopolymers) in the Nanopore-assembled sequences were corrected through polishing using Illumina reads. In the mitogenome of FS_24.009, the differences between the Flye Nanopore assembly before and after Illumina polishing were: a one-base indel in *nad4L* and a three-base indel in the *rrnS*; in the mitogenome of FS_24.010, a one-base indel in *atp6*, another one in *nad4L* and a two-base indel in *rrnS*; in the mitogenome of FS_24.011, a one-base indel in *nad4L* and a two-base indel in *rrnS*.

By contrast, the Illumina-only assemblies obtained using NOVOplasty had many discrepancies with the Nanopore-assembled, Illumina-polished highly accurate sequences: the majority of the differences were detected in the control regions with various insertions, and few indels and base substitutions were also found in tRNAs as well as in some protein-coding genes; for the individual FS_24.009 we detect four substitutions in the trnY and one substitutions in *nad3* and *nad4L*, for the individual FS_24.010 we detect four substitutions in the trnY, a 43 bases deletion between the trnQ and trnM, one base substitution in *nad3*, for the individual FS_24.011 we detect four substitutions in the trnY, a 55 bases deletion between the trnI and trnY, a 5 bases deletion between in the trnG and one base substitution in *nad3*

The ML tree obtained from this analysis is shown in Fig.9. The newly annotated mitogenome of *Niphargus* forms a clade with *Pseudoniphargus* supported with a 99% ultrafast bootstrap value. No difference was found in the protein-coding gene order between Niphargidae and Pseudoniphargidae, but some differences were detected between more distant amphipod families. Most Gammaridae and all Crangonyctidae had the same gene order as Niphargidae and Pseudoniphargidae, with the exception of *Marinogammarus marinus* for which *nad5*, *rrnS* and *rrnL* were translocated. In the case of Metacrangonyctidae, *cytb* changed position and orientation; whereas in Hyalellidae, *nad1* changed orientation but remained in the same location. The talitrid *Trinorchestia longiramus* had *nad3* located before *cox1* and its *cytb* and *nad6* were swapped.

Discussion

Like other metazoans, the mitochondrial genome of crustaceans typically consists of a circu-
lar double-stranded DNA molecule ranging from 12 to 20 kb, with highly conserved gene con-
tent. It contains 13 protein-coding genes (PCGs), two ribosomal RNA genes (rRNAs), 22 trans-
fer RNA genes (tRNAs), and a large non-coding region where replication of the mitochondrial
genome is initiated, called the D-loop or control region (CR) (Boore, 1999). Nonetheless, the ar-
rangement of genes varies significantly among higher taxonomic levels in amphipods, such as
families and above, owing to gene rearrangements such as transpositions and reversals (Macher
et al., 2023). In this study, we present the results obtained from the first three mitogenome
annotations of the genus *Niphargus*.
Nanopore assembly followed by careful Illumina polishing (first automatically, then manually)
yielded genome annotations that were highly consistent with published amphipod mitogenomes,
particularly with those of Pseudoniphargidae, which is the sister family to Niphargidae (Weber
et al., 2021). Most mt-tRNAs folded into the same cloverleaf secondary structure as nuclear-
encoded tRNA sequences, made of four stems and three loops (Jühling et al., 2012). However,
as reported in the genus *Pseudoniphargus* Stokkan et al., 2016, the tRNA-Ser1 and tRNA-Val of
Niphargus dolenianensis lacked the dihydrouridine (DHU) arm, also common in nearly all meta-
zoans (Ki et al., 2010), while its tRNA-Phe lacked the T-arm.
The tRNA secondary structures generally contained wobble base pairs (G–U, I–U, I–A, and
I–C); however, we also noticed some non-canonical paring (e.g. U–U, G–G, U–C, A–A) as fre-
quently observed in other species (Ki et al., 2010).
Nucleotide diversity values were higher for protein-coding genes than for ribosomal RNA
genes. Among the protein-coding genes, a higher diversity value was observed for *nad6*, whereas
cox1 had a comparatively lower value, even though the latter gene is commonly used to identify
arthropod species.
By contrast, both Nanopore-only and Illumina assemblies were imperfect, with many struc-
tural errors for Illumina-only assemblies (mainly in the control region and in ribosomal RNA genes,
although adjacent tRNA regions were also affected), whereas the Nanopore-only assemblies
were plagued by one-based artifactual indels in coding regions (caused by long homopolymers),
resulting in reading-frame shifts and erroneous protein predictions.
The fact that the Nanopore-only assembly contained artifactual indels that, for some of them,
disrupted protein sequence prediction was unexpected, given that Nanopore R10.4.1 data have
been reported to yield error-free bacterial genome sequences that do not require Illumina pol-
ishing (Sereika et al., 2022). As mitochondria are alpha-proteobacteria (Fan et al., 2020), the
same approach would have been expected to work on mitogenomes as well. However, the mi-
togenome of *Niphargus dolenianensis* contained many long stretches of polyA and polyT (up to a
length of 25 identical nucleotides in a row) that are not typically observed in bacterial genomes
and were responsible for the artifactual indels observed. This is most probably caused by the ex-
tremely low GC content of the mitochondrial genome of *Niphargus dolenianensis*, which, being
lower than 25% GC, falls outside the range of GC content of bacteria and archaea, except for a
very few cases such as *Carsonella* (Mann and Chen, 2010).
In such case, both Nanopore and Illumina data are required to obtain a highly accurate, error-
free mitogenome sequence. This is particularly important when producing the first reference
mitogenome sequence for a previously unexplored genus or family, as was the case here. Subse-
quent resequencing of other species closely related to the one for which a reference sequence
was generated could, however, probably rely on Nanopore only, since all the artifactual errors
in the resulting genome assembly will be located at the level of long homopolymers and will be
easily detected and corrected by hand.
Sequencing the mitogenomes of three closely related specimens, as we did, allowed us to
cross-compare and refine our gene annotations and was also instrumental in obtaining properly
folded sequences for all tRNAs, as the tRNA-Val of FS_24.011 could not be folded properly with-
out relying on information from the other two. Therefore, if we had only sequenced FS_24.011,
we could have incorrectly concluded that tRNA-Val was absent in *Niphargus dolenianensis*.

The phylogenetic tree of amphipods using the complete set of protein-coding genes in the mitochondrial genomes yielded a well-resolved phylogeny where all nodes had very strong bootstrap support and confirmed the close evolutionary relationship between Niphargidae and Pseudoniphargidae. This highlights the usefulness of complete mitochondrial genome sequences to conduct studies at a deeper phylogenetic level (Bauzà-Ribot et al., 2009; Sun et al., 2018). It is well known that the saturation problem of the *cox1* gene in amphipods (Stoch et al., 2024a, 2022) leads to poor resolution of ancient splits. By contrast, the strongly supported, highly resolved phylogenetic tree yielded by complete mitogenome sequences of amphipods will make it possible to resolve these basal nodes and to obtain a complete picture of the evolution and diversification of amphipods.

Acknowledgements

The authors wish to thank Laurent Grumiau and Florence Rodriguez Gaudray for their assistance with lab work.

AS's PhD was supported by a Université libre de Bruxelles (ULB) seed grant and by DarCo (The vertical dimension of conservation: A cost-effective plan to incorporate subterranean ecosystems in post-2020 biodiversity and climate change agendas, BIODIV21_0006).

NL's PhD was supported by a seed grant for ULB-VUB collaborative research.

FS and JFF were supported by the Belgian Fond de la Recherche Scientifique (F.R.S.-FNRS) via 'Chargé de recherches' fellowship no. FC43267 to FS and 'Projet de Recherches' grant no. T.0078.23 to JFF.

This study is part of the DarCo project funded by Biodiversa+, the European Biodiversity Partnership under the 2021–2022 BiodivProtect joint call for research proposals, co-funded by the European Commission (GA no. 101052342) and the Ministry of Universities and Research (Italy), Agencia Estatal de Investigación—Fundación Biodiversidad (Spain), Fundo Regional para a Ciência e Tecnologia (Portugal), Suomen Akatemia—Ministry of the Environment (Finland), Belgian Science Policy Office (Belgium), Agence Nationale de la Recherche (France), Deutsche Forschungsgemeinschaft e.V. (Germany), Schweizerischer Nationalfonds (grant no. 31BD30_209583, Switzerland), Fonds zur Förderung der Wissenschaftlichen Forschung (Austria), Ministry of Higher Education, Science and Innovation (Slovenia), and the Executive Agency for Higher Education, Research, Development and Innovation Funding (Romania).

Conflict of interest disclosure

The authors declare no financial or personal interest that could appear to have influenced the work reported in this study.

Author contributions

AS conducted most of the laboratory work, contributed to data analysis and interpretation, and drafted the manuscript with input from all authors. NL contributed to the laboratory work. FS collected all samples and contributed to data analysis, mitogenome annotation, and interpretation. JFF conducted most of the data analyses and contributed to their interpretation and validation. All authors reviewed and approved the final manuscript.

Figure 1 – *Niphargus dolenianensis*, habitus (a): male; (b): female

Voucher	Accession number	Locality	Latitude (WGS84)	Longitude (WGS 84)	Altitude (m a.s.l.)
FS_24.009	PV534572	San Giovanni al Natisone, spring Villa Trento, Dolegnano, Italy	13.420386	45.986732	80
FS_24.010	PV534571	Gorizia, spring along Vallone dell'Acqua road, Italy	13.597421	45.954605	163
FS_24.011	PV534570	Gorizia, Cormons, Forest of Plessiva, springbrook, Italy	13.488482	45.977869	97

Figure 2 – List of specimens used in the study with the GenBank accession numbers of the deposited mitogenome sequences. All specimens were collected on 08/04/2024 by F. Stoch F. and G. Tomasin then identified morphologically.

Species:	GenBank code:
Asellus aquaticus	GU130252
Echinogammarus berilloni	BK059223
Gammarus fossarum	NC_034937
Gammarus lacustris	NC_044469
Gammarus roeselii	NC_037481
Hyalella azteca	MH542433
Ligia oceanica	DQ442914
Marinogammarus marinus	BK059224
Metacrangonyx dominicanus	HE860499
Metacrangonyx ilvanus	NC_019656
Metacrangonyx longipes	NC_013032
Metacrangonyx repens	HE860495
Pectenogammarus veneris	BK059233
Pseudoniphargus grandis	MH592128
Pseudoniphargus morenoi	MH592132
Pseudoniphargus sp. 2-Canaries	MH592142
Pseudoniphargus stocki	NC_039354
Stygobromus allegheniensis	NC_046511
Stygobromus indentatus	NC_030261
Stygobromus pizzinii	NC_046510
Stygobromus tenuis potomacus	KU869712
Trinorchestia longiramus	MH542431

Figure 3 – List of all the complete mitochondrial sequences downloaded from GenBank used for mitochondrial genome analysis and integrated with new ones used to build the ML tree.

Figure 4 – Organization of the mitochondrial genome of *Niphargus dolenianensis* showing the arrangement of the 13 protein-coding genes, the twelve tRNAs, the two rRNA genes and the putative control region (in black).

Region	FS_24.009 Position	AA	Start	Stop	FS_24.010 Position	AA	Start	Stop	FS_24.011 Position	AA	Start	Stop
trnI	1-61				1-61				1-60			
trnY	64-122				64-122				113-171			
trnQ	119-176				119-176				168-225			
trnM	227-287				274-334				277-337			
trnC	288-346				335-393				338-396			
nad2	**348-1338**	**330**	**ATT**	**T--**	**395-1385**	**330**	**ATA**	**T--**	**398-1388**	**330**	**ATT**	**T--**
trnW	1339-1401				1386-1448				1389-1451			
trnG	1406-1469				1452-1515				1455-1518			
cox1	**1470-3006**	**512**	**ATA**	**T--**	**1516-3052**	**512**	**ATA**	**T--**	**1519-3055**	**511**	**ATT**	**T--**
trnL2	3007-3068				3053-3114				3056-3117			
cox2	**3069-3741**	**224**	**ATA**	**T--**	**3115-3787**	**224**	**ATA**	**T--**	**3118-3790**	**224**	**ATA**	**T--**
trnK	3742-3802				3788-3848				3791-3851			
trnD	3803-3862				3849-3908				3852-3911			
atp8	**3863-4024**	**53**	**ATT**	**TAA**	**3909-4070**	**53**	**ATT**	**TAA**	**3912-4073**	**53**	**ATC**	**TAA**
atp6	**4018-4686**	**222**	**ATG**	**TAA**	**4064-4732**	**222**	**ATG**	**TAA**	**4067-4735**	**222**	**ATG**	**TAA**
cox3	**4687-5475**	**262**	**ATG**	**TAA**	**4733-5524**	**263**	**ATG**	**TAA**	**4736-5524**	**262**	**ATG**	**TAA**
nad3	**5484-5837**	**117**	**ATG**	**TAG**	**5530-5883**	**117**	**ATG**	**TAG**	**5533-5886**	**117**	**ATG**	**TAG**
trnA	5836-5896				5882-5942				5885-5945			
trnS1	5896-5947				5942-5993				5945-5996			
trnN	5947-6008				5993-6054				5996-6057			
trnE	6006-6069				6052-6115				6055-6118			
trnR	6064-6122				6110-6168				6113-6171			
trnF	6121-6180				6167-6225				6170-6229			
nad5	**6181-7888**	**569**	**ATA**	**T--**	**6226-7933**	**569**	**ATA**	**T--**	**6230-7937**	**569**	**ATA**	**T--**
trnH	7889-7948				7934-7993				7938-7997			
nad4	**7949-9263**	**438**	**ATG**	**T--**	**7994-9308**	**439**	**ATG**	**T--**	**7998-9312**	**438**	**ATG**	**T--**
nad4L	**9257-9547**	**96**	**ATG**	**TAA**	**9302-9592**	**96**	**ATG**	**TAA**	**9306-9596**	**96**	**ATG**	**TAA**
trnT	9551-9610				9595-9654				9599-9658			
trnP	9609-9670				9653-9714				9657-9718			
nad6	**9673-10170**	**165**	**ATG**	**TAA**	**9729-10214**	**161**	**ATT**	**TAA**	**9733-10218**	**161**	**ATT**	**TAA**
cob	**10170-11309**	**379**	**ATG**	**TAA**	**10214-11353**	**379**	**ATG**	**TAA**	**10218-11357**	**379**	**ATG**	**TAA**
trnS2	11308-11367				11352-11411				11356-11415			
nad1	**11365-12276**	**303**	**ATT**	**TAA**	**11409-12320**	**303**	**ATT**	**TAA**	**11413-12324**	**303**	**ATT**	**TAA**
trnL1	12286-12347				12330-12391				12334-12395			
rrnL	12325-13359				12369-13404				12373-13408			
trnV	13384-13429				13428-13473				13433-13478			
rrnS	13430-14487				13474-14528				13479-14531			
CR	14488-15097				14529-14964				14532-15069			

Figure 5 – Position of all coding and non-coding genes, number of aminoacids in PCGs and their start and stop codons. The underlined numbers represent the position where genes overlap.

Figure 6 – Read coverage graph comparing Illumina and Nanopore data for the three *Niphargus* individual. Graphs with logarithmic y-axis.

Figure 7 – Predicted secondary structure of the 22 tRNA genes.

Figure 8 – Nucleotide diversity of the three *Niphargus* mitogenomes for protein-coding genes and rRNA genes.

Figure 9 – Maximum likelihood phylogenetic tree obtained from translated protein sequences of all mitochondrial protein-coding genes. Different families are indicated by black lines. Labels of nodes with a probability below 95 % were removed. 16S, 12S and CR represent the position of the *rrnL* and *rrnS* genes as well as the control region, when available from the annotation on Genbank.

Command lines

- Assembly of Nanopore reads using Flye:

```
mkdir Q14min
gzip -c -d *.fastq.gz|split_qscore.py -q 14 -f fastq - Q14min
cd Q14min/
rm reads.fail.fastq
flye --nano-hq reads.pass.fastq --threads 10 -o flye
```

- Assembly of Nanopore reads using hifiasm:

```
hifiasm --ont reads.pass.fastq -t10 -o hifiasm
```

- Checking the Nanopore coverage:

```
minimap2 -a -t 10 -x map-ont mtgenome.fasta reads.fastq > checkNanopore.sam
samtools view -b checkNanopore.sam -o checkNanopore.bam
samtools sort checkNanopore.bam -o checkNanopore_sorted.bam
samtools index checkNanopore_sorted.bam
```

- Comparison with Illumina assembly and polishing:

```
skewer -m pe R1_001.fastq.gz "R2_001.fastq.gz
mv R1_001.fastq-trimmed-pair1.fastq R1trimmed.fastq
mv "R1_001.fastq-trimmed-pair2.fastq R2trimmed.fastq
micromamba activate polypolish
module load BWA
bwa index mtDNA_circular.fasta
bwa mem -t 16 mtDNA_circular.fasta R1trimmed.fastq > R1trimmed.sam
bwa mem -t 16 _mtDNA_circular.fasta R2trimmed.fastq > R2trimmed.sam
polypolish filter --in1 R1trimmed.sam --in2 R2trimmed.sam \
        --out1 R1trimmed_filtered.sam --out2 R2trimmed_filtered.sam
polypolish polish mtDNA_circular.fasta R1trimmed_filtered.sam \
        R2trimmed_filtered.sam > mtDNA_circular_polished.fasta
```

- Plotting a circular map of the genome:

```
mtSVG.py --gff FS\_24.010.gff3 --size 14964 --species "Niphargus dolenianensis" --circular
```

- Plotting coverage graphs:

```
genomeCoverageBed -ibam checkNanopore_sorted.bam -d > checkNanopore.cov
genomeCoverageBed -ibam checkIllumina_sorted.bam -d > checkIllumina.cov
cut -f 2,3  checkIllumina.cov|xmgrace -
cut -f 2,3  checkNanopore.cov|xmgrace -
cut -f 2,3 checkNanopore.cov > Nanopore.cov
cut -f 2,3 checkIllumina.cov > Illumina.cov
xmgrace Illumina.cov Nanopore.cov
```

- Nucleotide diversity using Rstudio 4.4.1:

```
library(pegas)
library(seqinr)
file_paths <- c("atp6.fasta",
"atp8.fasta",
"cob.fasta",
"cox1.fasta",
"cox2.fasta",
"cox3.fasta",
"nad1.fasta",
"nad2.fasta",
"nad3.fasta",
"nad4L.fasta",
"nad4.fasta",
```

```
294  "nad5.fasta",
295  "nad6.fasta",
296  "rrnL.fasta",
297  "rrnS.fasta")
298
299  nuc_div_results <- list()
300  for (file in file_paths) {
301    # Read the gene sequence
302    gene_data <- read.dna(file, format = "fasta")
303
304    # Calculate nucleotide diversity
305    nuc_div_result <- nuc.div(gene_data)
306
307    # Store the result with the file name (gene name)
308    gene_name <- basename(file)  # Extract the file name to use as the gene name
309    nuc_div_results[[gene_name]] <- nuc_div_result
310  }
311  print(nuc_div_results)
312
313  write.csv(nuc_div_df, "C:/Users/Alice/Desktop/nucleotide_diversity_results.csv", row.names =
314  nuc_div_df <- data.frame(Gene = character(), Nucleotide_Diversity = numeric(), stringsAsFac
315  for (file in file_paths) {
316    gene_data <- read.dna(file, format = "fasta")
317    nuc_div_result <- nuc.div(gene_data)
318    gene_name <- basename(file)
```

319 • ML phylogenetic tree using IQtree2:

```
320  iqtree2 -s Mitogenomes_PCGs.phy -m mtInv+I+G4 -nt auto -B 50000
```

References

Bardgett RD, Van Der Putten WH (2014). Belowground biodiversity and ecosystem functioning. *Nature* **515**, 505–511. https://doi.org/10.1038/nature13855.

Bauzà-Ribot MM, Jaume D, Juan C, Pons J (2009). The complete mitochondrial genome of the subterranean crustacean *Metacrangonyx longipes* (Amphipoda): A unique gene order and extremely short control region. *Mitochondrial DNA* **20**, 88–99. https://doi.org/10.1080/19401730902964417.

Bauzà-Ribot MM, Juan C, Nardi F, Oromí P, Pons J, Jaume D (2012). Mitogenomic phylogenetic analysis supports continental-scale vicariance in subterranean thalassoid crustaceans. *Current Biology* **22**, 2069–2074. https://doi.org/10.1016/j.cub.2012.09.012.

Benito JB, Porter ML, Niemiller ML (2021). The mitochondrial genomes of five spring and groundwater amphipods of the family Crangonyctidae (Crustacea: Amphipoda) from eastern North America. *Mitochondrial DNA Part B* **6**, 1662–1667. https://doi.org/10.1080/23802359.2021.1926350.

Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritzsch G, Pütz J, Middendorf M, Stadler PF (2013). MITOS: Improved de novo metazoan mitochondrial genome annotation. *Molecular Phylogenetics and Evolution* **69**, 313–319. https://doi.org/10.1016/j.ympev.2012.08.023.

Boore JL (1999). Animal mitochondrial genomes. *Nucleic Acids Research* **27**, 1767–1780. https://doi.org/10.1093/nar/27.8.1767.

Cameron SL, Lambkin CL, Barker SC, Whiting MF (2007). A mitochondrial genome phylogeny of Diptera: whole genome sequence data accurately resolve relationships over broad timescales with high precision. *Systematic Entomology* **32**, 40–59. https://doi.org/10.1111/j.1365-3113.2006.00355.x.

Chan PP, Lin BY, Mak AJ, Lowe TM (2021). tRNAscan-SE 2.0: improved detection and functional classification of transfer RNA genes. *Nucleic Acids Research* **49**, 9077–9096. https://doi.org/10.1093/nar/gkab688.

Charif D, Lobry JR (2007). SeqinR 1.0-2: A contributed package to the R project for statistical computing devoted to biological sequences retrieval and analysis. In: *Structural Approaches to Sequence Evolution*. Springer, Berlin, Heidelberg, pp. 207–232. https://doi.org/10.1007/978-3-540-35306-5_10.

Cheng H, Concepcion GT, Feng X, Zhang H, Li H (2021). Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm. *Nature Methods* **18**, 170–175. https://doi.org/10.1038/s41592-020-01056-5.

Cormier A, Wattier R, Teixeira M, Rigaud T, Cordaux R (2018). The complete mitochondrial genome of *Gammarus roeselii* (Crustacea, Amphipoda): insights into mitogenome plasticity and evolution. *Hydrobiologia* **825**, 197–210. https://doi.org/10.1007/s10750-018-3578-z.

Dierckxsens N, Mardulyn P, Smits G (2017). NOVOPlasty: *de novo* assembly of organelle genomes from whole genome data. *Nucleic Acids Research* **45**, e18. https://doi.org/10/gfws9t.

Dodsworth S (2015). Genome skimming for next-generation biodiversity analysis. *Trends in Plant Science* **20**, 525–527. https://doi.org/10.1016/j.tplants.2015.06.012.

Eme D, Zagmajster M, Delić T, Fišer C, Flot JF, Konecny-Dupré L, Pálsson S, Stoch F, Zakšek V, Douady CJ, Malard F (2018). Do cryptic species matter in macroecology? Sequencing European groundwater crustaceans yields smaller ranges but does not challenge biodiversity determinants. *Ecography* **41**, 424–436. https://doi.org/10.1111/ecog.02683.

Fan L, Wu D, Goremykin V, Xiao J, Xu Y, Garg S, Zhang C, Martin WF, Zhu R (2020). Phylogenetic analyses with systematic taxon sampling show that mitochondria branch within Alphaproteobacteria. *Nature Ecology & Evolution* **4**, 1213–1219. https://doi.org/10.1038/s41559-020-1239-x.

Griebler C, Avramov M (2015). Groundwater ecosystem services: a review. *Freshwater Science* **34**, 355–367. https://doi.org/10.1086/679903.

Gruber AR, Bernhart SH, Lorenz R (2015). The ViennaRNA web services. In: *RNA Bioinformatics*. Humana Press, New York, NY, pp. 307–326. https://doi.org/10.1007/978-1-4939-2291-8_19.

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS (2018). UFBoot2: improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution* **35**, 518–522. https://doi.org/10.1093/molbev/msx281.

Horton T, De Broyer C, Bellan-Santini D, Coleman CO, Copilaș-Ciocianu D, Corbari L, Daneliya ME, Dauvin JC, Decock W, Fanini L, Fišer C, Gasca R, Grabowski M, Guerra-García JM, Hendrycks EA, Hughes LE, Jaume D, Kim YH, King RA, Lo Brutto S, et al. (2023). The World Amphipoda Database: history and progress. *Records of the Australian Museum* **75**, 329–342. https://doi.org/10.3853/j.2201-4349.75.2023.1875.

Jiang H, Lei R, Ding SW, Zhu S (2014). Skewer: a fast and accurate adapter trimmer for next-generation sequencing paired-end reads. *BMC Bioinformatics* **15**, 182. https://doi.org/10.1186/1471-2105-15-182.

Jühling F, Pütz J, Bernt M, Donath A, Middendorf M, Florentz C, Stadler PF (2012). Improved systematic tRNA gene annotation allows new insights into the evolution of mitochondrial tRNA structures and into the mechanisms of mitochondrial genome rearrangements. *Nucleic Acids Research* **40**, 2833–2845. https://doi.org/10.1093/nar/gkr1131.

Ki JS, Hop H, Kim SJ, Kim IC, Park HG, Lee JS (2010). Complete mitochondrial genome sequence of the Arctic gammarid, *Onisimus nanseni* (Crustacea; Amphipoda): Novel gene structures and unusual control region features. *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics* **5**, 105–115. https://doi.org/10.1016/j.cbd.2010.02.002.

Kilpert F, Held C, Podsiadlowski L (2012). Multiple rearrangements in mitochondrial genomes of Isopoda and phylogenetic implications. *Molecular Phylogenetics and Evolution* **64**, 106–117. https://doi.org/10.1016/j.ympev.2012.03.013.

Kilpert F, Podsiadlowski L (2006). The complete mitochondrial genome of the common sea slater, *Ligia oceanica* (Crustacea, Isopoda) bears a novel gene order and unusual control region features. *BMC Genomics* **7**, 241. https://doi.org/10.1186/1471-2164-7-241.

Kolmogorov M, Yuan J, Lin Y, Pevzner PA (2019). Assembly of long, error-prone reads using repeat graphs. *Nature Biotechnology* **37**, 540–546. https://doi.org/10/gfzbrd.

Kumar Patra A, Kim MS, Jung TW, Cho IY, Yoon M, Choi JH, Yang Y (2019). The complete mitochondrial genome of the sand-hopper *Trinorchestia longiramus* (Amphipoda: Talitridae). *Mitochondrial DNA Part B* **4**, 2104–2105. https://doi.org/10.1080/23802359.2019.1623100.

Larsson A (2014). AliView: a fast and lightweight alignment viewer and editor for large datasets. *Bioinformatics* **30**, 3276–3278. https://doi.org/10.1093/bioinformatics/btu531.

Le VS, Dang CC, Le QS (2017). Improved mitochondrial amino acid substitution models for metazoan evolutionary studies. *BMC Evolutionary Biology* **17**, 136. https://doi.org/10.1186/s12862-017-0987-y.

Li H (2018). Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**. Ed. by Inanc Birol, 3094–3100. https://doi.org/10/gdhbqt.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, Genome Project Data Processing S (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079. https://doi.org/10.1093/bioinformatics/btp352.

Lorenzi A (1898). Prime osservazioni zoologiche sulle acque freatiche del Friuli. *In Alto - Cronaca della Societa Alpina Friulana* **9**, 35–37.

Macher JN, Leese F, Weigand AM, Rozenberg A (2017). The complete mitochondrial genome of a cryptic amphipod species from the *Gammarus fossarum* complex. *Mitochondrial DNA Part B* **2**, 17–18. https://doi.org/10.1080/23802359.2016.1275844.

Macher JN, Šidagytė-Copilas E, Copilaș-Ciocianu D (2023). Comparative mitogenomics of native European and alien Ponto-Caspian amphipods. *NeoBiota* **87**, 27–44. https://doi.org/10.3897/neobiota.87.105941.

Mamos T, Grabowski M, Rewicz T, Bojko J, Strapagiel D, Burzyński A (2021). Mitochondrial genomes, phylogenetic associations, and SNP recovery for the key invasive Ponto-Caspian amphipods in Europe. *International Journal of Molecular Sciences* **22**, 10300. https://doi.org/10.3390/ijms221910300.

Mann S, Chen YPP (2010). Bacterial genomic G+C composition-eliciting environmental adaptation. *Genomics* **95**, 7–15. https://doi.org/10.1016/j.ygeno.2009.09.002. (Visited on 12/30/2021).

Maurice L, Bloomfield J (2012). Stygobitic invertebrates in groundwater — a review from a hydrogeological perspective. *Freshwater Reviews* **5**, 51–71. https://doi.org/10/gfkjr5.

Milne I, Bayer M, Cardle L, Shaw P, Stephen G, Wright F, Marshall D (2010). Tablet—next generation sequence assembly visualization. *Bioinformatics* **26**, 401–402. https://doi.org/10/cp76wb.

Nguyen LT, Schmidt HA, Von Haeseler A, Minh BQ (2015). Iq-tree: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution* **32**, 268–274. https://doi.org/10.1093/molbev/msu300.

Paradis E (2010). pegas: an R package for population genetics with an integrated-modular approach. *Bioinformatics* **26**, 419–420. https://doi.org/10/dx52rx.

Saccò M, Mammola S, Altermatt F, Alther R, Bolpagni R, Brancelj A, Brankovits D, Fišer C, Gerovasileiou V, Griebler C, Guareschi S, Hose GC, Korbel K, Lictevout E, Malard F, Martínez A, Niemiller ML, Robertson A, Tanalgo KC, Bichuette ME, et al. (2024). Groundwater is a hidden global keystone ecosystem. *Global Change Biology* **30**, e17066. https://doi.org/10.1111/gcb.17066.

Samarakoon H, Ferguson JM, Gamaarachchi H, Deveson IW (2023). Accelerated nanopore basecalling with SLOW5 data format. *Bioinformatics* **39**, btad352. https://doi.org/10.1093/bioinformatics/btad352.

Sereika M, Kirkegaard RH, Karst SM, Michaelsen TY, Sørensen EA, Wollenberg RD, Albertsen M (2022). Oxford Nanopore R10.4 long-read sequencing enables the generation of near-finished bacterial genomes from pure cultures and metagenomes without short-read or reference polishing. *Nature Methods* **19**, 823–826. https://doi.org/10.1038/s41592-022-01539-7.

Stoch F, Citoleux J, Weber D, Salussolia A, Flot JF (2024a). New insights into the origin and phylogeny of Niphargidae (Crustacea: Amphipoda), with description of a new species and synonymization of the genus *Niphargellus* with *Niphargus*. *Zoological Journal of the Linnean Society* **202**, zlae154. https://doi.org/10.1093/zoolinnean/zlae154.

Stoch F, Knüsel M, Zakšek V, Alther R, Salussolia A, Altermatt F, Fišer C, Flot JF (2024b). Integrative taxonomy of the groundwater amphipod *Niphargus bihorensis* Schellenberg, 1940 reveals a species-rich clade. *Contributions to Zoology* **93**, 371–395. https://doi.org/10.1163/18759866-bja10064.

Stoch F, Salussolia A, Flot JF (2022). Polyphyly of the *Niphargus stygius* species group (Crustacea, Amphipoda, Niphargidae) in the Southern Limestone Alps. https://doi.org/10.1101/2022.04.28.489871.

Stokkan M, Jurado-Rivera JA, Juan C, Jaume D, Pons J (2016). Mitochondrial genome rearrangements at low taxonomic levels: three distinct mitogenome gene orders in the genus *Pseudoniphargus* (Crustacea: Amphipoda). *Mitochondrial DNA Part A* **27**, 3579–3589. https://doi.org/10.3109/19401736.2015.1079821.

Stokkan M, Jurado-Rivera JA, Oromí P, Juan C, Jaume D, Pons J (2018). Species delimitation and mitogenome phylogenetics in the subterranean genus *Pseudoniphargus* (Crustacea: Amphipoda). *Molecular Phylogenetics and Evolution* **127**, 988–999. https://doi.org/10.1016/j.ympev.2018.07.002.

Sun S, Hui M, Wang M, Sha Z (2018). The complete mitochondrial genome of the alvinocaridid shrimp *Shinkaicaris leurokolos* (Decapoda, Caridea): insight into the mitochondrial genetic basis of deep-sea hydrothermal vent adaptation in the shrimp. *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics* **25**, 42–52. https://doi.org/10.1016/j.cbd.2017.11.002.

Tillich M, Lehwark P, Pellizzer T, Ulbricht-Jones ES, Fischer A, Bock R, Greiner S (2017). GeSeq – versatile and accurate annotation of organelle genomes. *Nucleic Acids Research* **45**, W6–W11. https://doi.org/10.1093/nar/gkx391.

Vaught A (1996). Graphing with Gnuplot and Xmgr: Two graphing packages available under Linux. *Linux Journal* **1996**, 7.

Weber D, Stoch F, Knight LR, Chauveau C, Flot JF (2021). The genus *Microniphargus* (Crustacea, Amphipoda): evidence for three lineages distributed across northwestern Europe and transfer from Niphargidae to Pseudoniphargidae. *Belgian Journal of Zoology* **151**, 169–191. https://doi.org/10.26496/bjz.2021.92.

Wick RR, Holt KE (2022). Polypolish: Short-read polishing of long-read bacterial genome assemblies. *PLOS Computational Biology* **18**, e1009802. https://doi.org/10.1371/journal.pcbi.1009802.

Wick RR, Schultz MB, Zobel J, Holt KE (2015). Bandage: interactive visualization of *de novo* genome assemblies. *Bioinformatics* **31**, 3350–3352. https://doi.org/10/gb5g64.

Zagmajster M, Eme D, Fišer C, Galassi D, Marmonier P, Stoch F, Cornu JF, Malard F (2014). Geographic variation in range size and beta diversity of groundwater crustaceans: insights from habitats with low thermal seasonality. *Global Ecology and Biogeography* **23**, 1135–1145. https://doi.org/10.1111/geb.12200.