Dimeric Cin8 motors have an inherent plus-end bias and weak inter-head coordination

Running title: Cin8 dimer motility

Himanshu Pandey[1], Tzu-Chen Ma[1], Eric Bonventre[2], Luke M. Rice[2], Larisa Gheber[3], and William O. Hancock[1,4]*

[1] Department of Biomedical Engineering, Pennsylvania State University, University Park, Pennsylvania, USA
[2] Departments of Biophysics and Biochemistry, University of Texas Southwestern Medical Center, Dallas, Texas, USA
[3] Department of Chemistry, Ben-Gurion University of the Negev, Israel
[4] Department of Chemistry, Pennsylvania State University, University Park, Pennsylvania, USA

* Corresponding author:
William O. Hancock
Email: woh1@psu.edu

Abstract:
Kinesin-5 motors are bipolar tetramers that crosslink and slide antiparallel microtubules during mitotic spindle assembly. Fungal kinesin-5 motors, such as Cin8, exhibit bidirectional motility, switching between minus- and plus-end-directed stepping in response to environmental conditions; however, the molecular basis of this directional switching remains unclear. To better understand the origin of this bidirectional behavior, we investigated the motility and ATPase kinetics of two Cin8 dimers, created by fusing the motor domains to a stable coiled-coil domain from kinesin-1. To investigate the role of the proximal neck coiled-coil region in coordinating motor activity, we compared Cin8 dimers that included or lacked the first four heptads of the Cin8 neck-coil domain. By analyzing the stepping kinetics, microtubule residence times, and directional switching dynamics, we found that these Cin8 dimers move processively with a net plus-end directionality along

with undirected movements, behaviors that mimic the plus-ended motility state of wild-
type Cin8. However, fast minus-ended motility seen in wild-type Cin8 tetramers was not
observed in the dimers. The instantaneous velocity distributions and ATPase rates were
inconsistent with the undirected movement being solely due to passive diffusion,
suggesting that they reflect random bidirectional stepping. Fewer undirected movements
were seen on yeast microtubules, their native physiological substrate, compared to on
bovine microtubules. Replacing the Cin8 neck-coil domain with a stable coiled-coil led to
faster plus-end stepping, fewer undirected movements, a reduction in the microtubule
binding duration, and enhanced coupling between ATP hydrolysis and plus-end stepping.
Our results suggest that the native Cin8 neck coil confers flexibility between the two motor
domains that contributes to bidirectional stepping, and that sustained minus-end
movement requires regions outside the motor domain.
48

49 Statement of Significance

The kinesin-5 family of motors, which contain two pairs of heads located on either end of a
long stalk domain, power mitotic spindle formation by sliding antiparallel microtubules.
The yeast kinesin-5, Cin8 moves to microtubule minus-ends under specific conditions,
breaking the dogma that N-terminal kinesins move to microtubule plus-ends. To gain
insight into how Cin8 changes its walking direction, we analyzed engineered Cin8 dimers,
models of each half of full-length Cin8. The dimers step erratically, consistent with
stepping in both directions, but have a plus-end bias. We find evidence that the two heads
poorly coordinate, which contrasts with other kinesins, and suggest that the stepping
direction may be regulated by altering the degree of inter-head coordination.
59

60 Introduction

Kinesins are a diverse family of microtubule-based motor proteins that play
essential roles in intracellular transport and mitotic spindle dynamics. Canonically, kinesin
directionality depends on the position of the motor domain: kinesins with N-terminal motor
domains move toward the microtubule plus end, whereas C-terminal kinesins move
toward the minus end (1,2). However, this paradigm is challenged by members of the
kinesin-5 and kinesin-14 families of mitotic motors that exhibit bidirectional motility (3-7).
During elongation of the mitotic spindle, kinesin-5 motors use their two pairs of motor
domains to slide apart antiparallel microtubules, with each pair of heads moving toward
microtubule plus-ends (8,9). However, fungal kinesin-5s, such as Cin8 and Kip1 from
Saccharomyces cerevisiae and Cut7 from *Schizosaccharomyces pombe*, have also been

shown to switch to minus-end directionality under certain conditions (3-5,10). Consistent
with this minus-end directionality, Cin8 localizes near the spindle pole bodies at
microtubule minus ends in early mitotic cells with monopolar spindles, and this minus-end
localization is proposed to enhance microtubule capture from the neighboring spindle pole
bodies to set up proper spindle organization for metaphase to proceed (4,10,11).
Understanding how bidirectional motility is controlled, particularly in the context of spindle
assembly where precise regulation of motor protein activity is critical, remains a major
challenge. Furthermore, the lack of a clear mechanism by which kinesins change their
stepping direction represents a significant hole in our understanding of kinesin
mechanochemistry.

In N-terminal kinesins such as kinesin-1, stepping is driven by ATP-dependent
docking of the neck linker (NL) domain, a ~14-18-residue segment at the C-terminal of the
motor domain (12). Upon ATP binding and hydrolysis, the neck linker transitions from an
undocked, unstructured conformation to a docked beta-sheet conformation that is
stabilized along the catalytic core by the N-terminal 'cover strand' (13-15). This
conformational change positions the partner (tethered) head toward the microtubule plus
end where it then binds to complete a plus-end directed (forward) step (13,16). In the
resulting two-head bound state, the now trailing head has its neck linker docked forward,
while the leading head has its neck linker stretched backward, and this inter-head tension
transmitted through the neck domains is thought to coordinate the ATPase cycles of the
two heads to ensure processive, hand-over-hand stepping (16-19). Consistent with this,
increasing the length and flexibility of the neck linker domain has been shown to disrupt
head-head coordination and reduce processivity (18,20-22).

A dominant theme that emerges from Cin8 work to date is that perturbations that
enhance microtubule affinity bias the motor toward slow plus-end motility, whereas
perturbations that reduce the microtubule affinity tend to bias the motor toward fast
minus-end directionality. For instance, motor clustering and crosslinking of antiparallel
microtubules by Cin8 bias it toward plus-end-directed motility (3,23). In contrast,
increasing the ionic strength or deleting the N-terminal extension, loop-8, or the C-terminal
tail, all of which can interact with microtubules, bias the motor toward minus-end
directionality (3,4,24-26). The underlying mechanism, however, remains unexplained. A
recent model proposes that Cin8 inherently steps in both directions, with minus-end
stepping dominating at zero load but having a stronger dependence on load such that plus-
end stepping dominates under load (23). An alternative model for the related fungal
kinesin-5, Cut7, posits that stepping is inherently minus-end directed and crowding on the
microtubule surface promotes plus-end stepping by a steric blocking mechanism (11).

There is also evidence that directionality of Cin8 is regulated allosterically by regions
outside the motor domain such as the tail or the N-terminal non-motor region (24-26).

In the present study, we investigated the role of inter-head coordination in the
bidirectional stepping of dimeric Cin8. We hypothesized that directional switching is
enabled by loose mechanical coupling between the motor domains. To test this, we used
the stable coiled-coil of kinesin-1 to dimerize Cin8 motor domains with and without their
native proximal neck-coil dimerization domain. We compared the stepping speed and
directionality on both bovine and yeast microtubules and analyzed the undirected
movements to determine the relative contributions of Brownian diffusion and ATP-driven
stepping. Our results demonstrate that Cin8 dimers exhibit slow net plus-end motility
along with undirected movements, and that flexibility in the proximal neck-coil reduces
inter-head coordination and leads to partial uncoupling between ATP hydrolysis and plus-
end directed stepping. These findings support a model in which loose mechanical
coordination between motor heads, modulated by structural elements of the neck region,
plays a critical role in enabling bidirectional motility in fungal kinesin-5.

Methods

Protein Constructs

Plasmids to express Cin8 variants were generated using standard PCR and Gibson
assembly and verified by DNA sequencing. All constructs are dimers comprising the *S.*
cerevisiae Cin8 motor and neck domains (residues 39–530 for Cin8Kin and 39–558 for
Cin8$_{+28NC}$Kin), fused to the Drosophila KHC neck-coil and dimerization coiled-coil domain
(residues 345–560; Fig. 1A). Constructs were made by inserting the Cin8 motor domains
into the K560-AviN plasmid described previously (27). All constructs begin at Cin8 residue
39 and include a C-terminal AviTag™ and His$_6$ tag for purification.

Purification and Quantification

Bacterial expression procedures were adapted from previously described protocol (25,27).
Briefly, constructs were expressed in E. coli BL21-CodonPlus (DE3) grown in 1 L of terrific
broth, in-house. 24 mg/l of biotin were added along with the isopropyl 1-thio-β-d-
galactopyranoside (IPTG) at the time of induction, followed by nickel gravity column
chromatography purification with an 50 mM HEPES based elution buffer containing 2 μM
mant-ADP along with 0.25 mM TCEP, 5% glycerol, 500 mM KCl, 10 mM MgCl$_2$, 0.01% Triton
X-100, pH 7.2. The elution fraction containing motors were supplemented with 10% (w/v)
sucrose and 2 mM DTT, flash-frozen and stored at -80 °C. The active motor concentration

was determined by measuring the total mant-ADP concentration (356 nm excitation/450

nm emission) in a Shimadzu 3000 spectrofluorometer and correcting for the free mant-

ADP, and using a fluorescence enhancement factor of 1.8 for motor-bound mant-ADP (28).

Stock active dimeric motor concentrations were in the range of 1–10 µM dimer.

Gliding assay

Flow chambers were assembled and motors adhered to the surface by flowing in 20 µg/ml

NeutrAvidin followed by C-terminal biotinylated motors. Taxol-stabilized microtubules were

then added to the flow cell in motility buffer (BRB80 adjusted to pH 7.2, 1 mM ATP, 10 µM

Taxol, 2 mM $MgCl_2$, 5 mM DTT, and 2 mg/ml casein, 0.5 mg/ml BSA). Microtubule

movements were monitored with the microscopy setup described for the single-molecule

motility assay and recorded at 5 fps. Microtubule velocities were measured by creating

kymographs along the MT path using ImageJ-Fiji software and measuring the slope. Periods

when microtubules paused or stopped moving were not included in the velocity analysis.

Single-molecule motility assay

Motility assays were performed on a Nikon TE2000 TIRF microscope equipped with

interference reflection microscopy, at room temperature, as described previously (29,30).

Dimeric Cin8 motors were labeled with streptavidin-labeled Quantum dots (Qdot 525;

Thermo Fisher) through the C-terminal AviTagTM bound biotin. For labelling, 100 nM

dimeric motors were mixed with Qdots at 1:5 ratio of motor:Qdots and incubated in ice for

20 min, such that each dimeric motor molecule is labeled with one Qdot. Flow cells were

functionalized by flowing in 0.5 mg/ml casein, followed by full-length rigor kinesin (27) in

the motility buffer (BRB80 adjusted to pH 7.2, 1 mM ATP/ADP, 10 µM Taxol, 2 mM $MgCl_2$, 5

mM DTT, and 2 mg/ml casein, 1 mg/ml BSA, 10 mM glucose, 100 µg/ml glucose oxidase, 80

µg/ml catalase 40, 10 mM phosphocreatine, and 50 µg/ml creatine phosphokinase). Taxol-

stabilized microtubules, polymerized from bovine or yeast tubulin, were then introduced

into the flow cell, where they attached to surface-immobilized rigor kinesin. Subsequently,

Qdot labeled motors were introduced and imaged. To accurately measure run times,

motors were introduced into the flow cell immediately upon initiating image acquisition,

ensuring that no motors were bound prior to the start of imaging. Cin8 motility on surface-

attached microtubules was recorded at either 20 fps or 5 fps and analyzed using FIESTA

software (31) to determine velocity and run durations. When measuring run durations,

motors that landed during the final 20% of the total acquisition time were excluded from

the analysis to prevent censoring by termination of the video. Microtubule polarities were

determined by flushing in fluorescent KIF1A at the end of the experiment (28).

Data analysis

Single-molecule tracking movies of Cin8 dimers were analyzed with FIESTA (31) to generate

displacement along microtubule versus time trajectories. From the tracked traces two

types of motility behaviors were evident: directed motility traces where motors move

processively towards plus ends of microtubules, and undirected traces, which we defined

as those with average velocities < 20 nm/s and net displacements < 1 µm (the majority of

which were < 0.5 µm). For mean-squared displacement (MSD) analysis, we calculated the

squared displacement during each time lag (τ) for each tracked particle and averaged the

squared displacements over all particles. To estimate the diffusion coefficients (D) we fit

MSD data with $v^2\tau^2 + 2D\tau + s^2$, where v is mean velocity obtained from the 5 s

instantaneous velocity distribution and s set to 10 nm based on the point spread function

measurement error in FIESTA. For Gaussian Mixture Model fitting, we used Matlab built-in

function "fitgmdist" to fit the PDF distribution of 5-s instantaneous velocities with 3

Gaussian functions. The maximum iteration was set to 1000 to ensure convergence of the

fit. All the fitting and plotting were performed using Matlab. Kymographs were generated

using ImageJ-FIJI (32).

Steady state ATPase assay

Cin8 ATPase activity at varying [Mt] was quantified by an enzyme-coupled assay following a

protocol adapted from Huang and Hackney (28,33). BRB80 buffer supplemented with 1 mM

Mg-ATP, 2 mm phosphoenolpyruvate, 1 mm $MgCl_2$, 0.2 mg/ml casein, 10 µM Taxol, 0.25 mM

NADH, and 1.5% volume % of Pyruvate Kinase/Lactate dehydrogenase (Sigma P-0294) was

used for ATPase assays at 21 °C. Dimeric motor concentrations between 50 and 100 nM

were used. NADH absorbance at 340 nm was measured over time on a Molecular Devices

FlexStation3 microplate reader, converted to an ATPase rate by linear fit to regions of

steady-state absorbance decrease, and divided by the active dimer-motor concentration to

give the total hydrolysis cycle rate. k_{cat} and K_M (MT) were determined from the fit to a

Michaelis-Menten equation of ATP turnover rates at various microtubule concentrations.

Mean ± SD values were obtained from five independent assays for each Cin8 variant with

motors from at least two different purification preparations.

Simulations

Cin8 stepping simulations used a Gillespie stochastic stepping model based on an

algorithm developed in previous work (34,35). Plus-end directed stepping was the only

reaction considered, with the unloaded stepping rate, $k_{forward}^0$ taken from the mean velocity

on yeast microtubules, assuming an 8 nm step size. To model diffusion, a diffusive term

was added to each step based on sampling from a normal distribution with mean of zero

and standard deviation $\sigma = \sqrt{2D\tau}$, where D is the chosen diffusion coefficient and τ is the

transition time for that step. All simulations were run 1000 times, and each run was
recorded for 50 seconds.

Results

Fig. 1: Design and microtubule gliding velocities of dimeric Cin8 constructs. (A) Domain
organization of Cin8 constructs used in this study. The Cin8 catalytic motor domain (residues 76–
512, blue) is flanked by the N-terminal non-motor region (residues 39–75, yellow) and the C-
terminal neck-linker, NL_{18} (residues 513–530, light blue). In Cin8Kin, the Kinesin-1 neck-coil and
dimerization domain (residues 345–560, gray) are fused downstream of the Cin8 head, followed by
a C-terminal Avi Tag (white) and His_6 tag. The $Cin8_{+28NC}$Kin construct includes additional 28 residues
from the Cin8 neck coil (residues 531–558, cyan), inserted between the Cin8 head and the Kinesin-
1 dimerization domain. Diagrams are not to scale **(B)** Distribution of microtubule gliding velocities
for both motors with 100 nM motors introduced into flow cell. **(C)** Microtubule gliding velocities at
varying motor concentrations in the flow cell.

Cin8 dimers are plus-end steppers in microtubule gliding assays

228 To investigate the role of inter-head tension in the directional switching and motility of
229 Cin8, we used a simplified dimeric construct lacking the second pair of motor domains, the
230 tetrameric stalk, and the native tail. To improve solubility, the first 38 amino acids of the
231 unstructured N-terminal domain were deleted. Stable dimers were created by fusing the
232 Cin8 motor domain (amino acids 39–530) to the neck coil and coiled-coil-1 of *Drosophila*

kinesin-1 (amino acids 345-560), followed by a biotinylation tag and His$_6$tag (Fig. 1A). This dimerization strategy has been used successfully for motors from the kinesin-1, 2, 3, 5, and 7 families and thus allows for direct comparison to previous work (18,28,36-38). The first dimeric construct, Cin8Kin, consists of the Cin8 catalytic core and 18-residue neck linker domain (amino acids 39-530) fused to the kinesin-1 dimerization domain. This construct is similar to one used previously in multimotor gliding assays (25), but lacks N-terminal residues 1-38 and four C-term residues compared to a Cin8Kin construct (residues 1-534) used previously in single-molecule assays (24).

Processive kinesin stepping requires a mechanical connection between the two motor domains, which is achieved through dimerization of the neck-coil domain. In contrast to kinesin-1 (39), the stability of the proximal neck-coil domain of Cin8 is not known. Coiled-coil prediction analyses indicate that the proximal neck coil of Cin8 is less stable than kinesin-1, as evidenced by lower coiled-coil stabilization scores and a reduced number of stabilizing interactions (Fig. S1). Because the inter-head tension that coordinates the hand-over-hand stepping in processive kinesins relies on the proximal neck-coil remaining tightly dimerized, we designed a second construct, Cin8$_{+28NC}$Kin, that included the first four heptads (28-residues) of the native Cin8 neck coil. These constructs were bacterially expressed and purified following standard procedures.

To confirm that the Cin8 dimers are functional, we tested their activity in multi-motor microtubule gliding assays. When biotinylated motors (100 nM solution concentration) were adsorbed to a neutravidin-coated coverslip, Cin8$_{+28NC}$Kin and Cin8Kin steadily drove bovine microtubules in a unidirectional plus-ended manner with no evidence of directional switching. Notably, the Cin8$_{+28NC}$Kin construct exhibited a microtubule gliding velocity approximately half that of Cin8Kin, providing evidence that the native Cin8 neck-coil lacks the stability of kinesin-1 (Fig. 1B). Previous work using full-length tetrameric Cin8 showed that microtubule gliding switched to minus-end directionality as the surface motor density was decreased (4). To test this, we decreased the motor concentration in the gliding assay and found no change in directionality. Instead, there was a slight increase in plus-end velocity for both motors at reduced motor densities (Fig. 1C). Thus, in multi-motor assays, Cin8 dimers displayed plus-end directionality, and dimerizing the heads just distal to their neck linker doubled the gliding speed compared to dimers that retained the first four heptads of the native Cin8 neck-coil.

Fig. 2: Single-molecule Cin8 dimer motility on bovine microtubules. Representative **(A)** kymographs and **(B)** displacement versus time traces for Cin8$_{+28NC}$Kin (orange) and Cin8Kin (blue) in the presence of 1 mM ATP on surface-immobilized bovine microtubules. Motors were labeled with quantum dots and movies acquired at 5 frames/s. Microtubule polarities, determined by introducing fluorescent KIF1A at the end of the experiment (Fig. S2), are marked at the bottom of kymograph. **(C)** Distribution of whole trace velocities for Cin8$_{+28NC}$Kin and Cin8Kin. Mean velocities were 43.8 ± 7.3 nm/s for Cin8$_{+28NC}$Kin and 62.1 ± 6.4 nm/s for Cin8Kin (mean ± SEM for n= 75 and 63, respectively). **(D)** 1 s and **(E)** 5 s instantaneous velocity distributions for Cin8$_{+28NC}$Kin and Cin8Kin. Average velocities from the 5 s velocity distributions were 32.9 ± 6.2 nm/s for Cin8$_{+28NC}$Kin and 58.6 ± 6.3 nm/s for Cin8Kin (mean ± SEM for N = 516 and 148 independent 5 s windows, respectively). Gray (Cin8$_{+28NC}$Kin) and green (Cin8Kin) shaded areas represent minus-end directed velocity segments. **(F)** Mean squared displacement (MSD) analysis in ATP. Data at varying lag time (τ) were fit to MSD = $v^2\tau^2$+2Dτ (black lines), with velocities (v) fixed at 32.9 nm/s (Cin8$_{+28NC}$Kin) and 58.6 nm/s (Cin8Kin). Apparent diffusion coefficients (D) were 14338 ± 172 nm^2/s and 10921 ± 40 nm^2/s, respectively (fit ± 95% CI). Fit to MSD = $a\tau^{\alpha}$ + b gave α of 1.29 for Cin8$_{+28NC}$Kin and 1.38 for Cin8Kin (Table S1).

Single Cin8 dimers move processively with net plus-end directionality

To investigate the speed and directionality of individual Cin8 dimers, we labeled the motors with streptavidin-coated quantum dots and observed their movements along immobilized microtubules polymerized from bovine brain tubulin (Fig. 2A & B). Consistent with the microtubule gliding assays, both Cin8 constructs displayed net plus-end directed movement (Fig. 2A-C & S2), with average trace velocities of 43.8 ± 7.3 nm/s for Cin8$_{+28NC}$Kin

and 62.1 ± 6.4 nm/s for Cin8Kin (mean ± SEM; Fig. 2C). However, in contrast to the smooth

movements in the multi-motor gliding assay, there were durations of undirected

movement, fluctuations in velocity during plus-end directed segments, and frequent short

segments with minus-end directionality (Fig. 2A & B). To more comprehensively compare

the single-molecule motility, we calculated instantaneous velocities for every trace using a

1 s or 5 s window for both motors and plotted the resulting instantaneous velocity

distributions (Fig. 2D & E). The 1 s and 5 s results were similar, but due to the long tails in

the 1 s distributions we focused our analysis on the 5 s distributions. Both variants had a

clear plus-end bias, with Cin8Kin having a distinct peak around ~100 nm/s and Cin8$_{+28NC}$Kin

having a broader distribution that was centered closer to zero. Notably, even with this

relatively long 5 s window, 33% of Cin8$_{+28NC}$Kin and 18% of Cin8Kin segments were minus-

end directed (Fig. 2E, shaded region), consistent with the considerable undirected

movements seen in the raw traces (Fig. 2A&B).

As an alternative way to quantify the degree of undirected motility, we carried out a mean

squared displacement (MSD) analysis using the mean velocity from the instantaneous

velocity distributions along with a diffusive term (Fig. 2F). Consistent with the wider

instantaneous velocity distribution, the apparent 1D diffusion coefficient was 14338 ± 172

nm^2/s for Cin8$_{+28NC}$Kin and 10921 ± 40 nm^2/s for Cin8Kin (fit ± 95% confidence interval, CI).

Fitting to a model MSD = aτ^{α} + b, where α = 1 denotes pure diffusion and α = 2 denotes

processive directional movement gave α of 1.29 for Cin8$_{+28NC}$Kin and 1.38 for Cin8Kin (Table

S1). Thus, consistent with its broader velocity distribution, Cin8$_{+28NC}$Kin displayed more

diffusion-like undirected movement than Cin8Kin.

Previous work on tetrameric Cin8 found that weakening motor-microtubule interactions by

increasing the buffer ionic strength resulted in a shift to minus-end directionality (3,4,23).

To test whether increasing ionic strength caused a similar directionality shift for Cin8

dimers, we repeated the experiments in BRB80 buffer supplemented with 110 mM KCl. In

contrast to tetramers, clear net plus-end motility of Cin8$_{+28NC}$Kin and Cin8Kin dimers was

retained in the presence of added KCl, with slightly higher whole trace velocities of 47.3 ±

7.0 and 84.1 ± 7.6 nm/s, respectively (mean ± SEM) (Fig. S3). Thus, engineered Cin8 dimers

display more inherent plus-end bias than native tetramers.

Fig. 3: Cin8 dimer motility on yeast microtubules. (A) Representative kymographs and (**B**) displacement versus time traces for Cin8$_{+28NC}$Kin (orange) and Cin8Kin (blue) motility in the presence of 1 mM ATP on yeast microtubules. Microtubule polarities are marked at the bottom of kymographs. **(C)** Distribution of whole trace velocities, with mean velocities 46.7 ± 4.4 for Cin8$_{+28NC}$Kin and 76.6 ± 4.2 nm/s for Cin8Kin (mean ± SEM for 95 and 106 traces). **(D)** 1 s and **(E)** 5 s instantaneous velocity distributions. Gray (Cin8$_{+28NC}$Kin) and green (Cin8Kin) shaded areas represent minus-end directed velocity segments. Average velocities from the 5 s velocity distributions were 29.4 ± 2.0 nm/s for Cin8$_{+28NC}$Kin and 59.8 ± 2.5 nm/s for Cin8Kin (mean ± SEM for N = 726 and 179 independent 5 s windows, respectively). **(F)** Mean squared displacement (MSD) analysis in ATP. Data were fit to, MSD = $v^2\tau^2+2D\tau$ (black lines), with velocities taken from Fig. 3E, yielding diffusion coefficients of 7194 ± 94 nm^2/s and 6915 ± 220 nm^2/s, respectively (fit ± 95% CI). Fit to MSD = $a\tau^\alpha + b$ yielded α of 1.29 for Cin8$_{+28NC}$Kin and 1.64 for Cin8Kin (Table S1). **(G)** Comparison of velocities (summarized from Figs. 2E & 3E). **(H)** Comparison of diffusion coefficients and **(I)** motor landing rates for Cin8$_{+28NC}$Kin (0.06 ± 0.007 and 0.036 ± 0.006 μm^{-1} s^{-1} nM^{-1} for bovine vs yeast, respectively; mean ± SEM) and Cin8Kin (0.068 ± 0.02 and 0.027 ± 0.002 μm^{-1} s^{-1} nM^{-1} for

bovine vs yeast, respectively; mean ± SEM) on bovine and yeast microtubules (2 and 3 fields,
respectively).

Cin8 dimers move more directionally on yeast microtubules

To test the role of the microtubule track on Cin8 motility, we carried out single-molecule
assays using recombinant *S. cerevisiae* tubulin, matching the species of the *S. cerevisiae*
Cin8. To the best of our knowledge, only one previous study investigated the motility of
Cin8 on yeast microtubule in vitro, and that study used full-length Cin8 tetramers from
yeast lysates (40). On yeast microtubules, both Cin8 dimer constructs exhibited similar net
plus-end directed motility as on bovine microtubules, though by visual inspection there
were fewer fluctuations in directionality on the yeast microtubules (Fig. 3A&B; see also Fig.
S4A). To quantify this undirected movement, we compared the fraction of traces that
showed clear processive movements, defined as traces with average velocities >20 nm/s
and net displacements >1 µm. Compared to bovine microtubules, the overall fraction of
directed traces on yeast microtubules increased from 28% to 41% for Cin8$_{+28NC}$Kin and
from 26% to 68% for Cin8Kin (Fig. S4B). Similar to the bovine microtubules, replacing the
proximal neck-coil of Cin8 with the stable kinesin-1 dimerization domain increased the
mean trace velocity from 46.7 ± 4.4 nm/s for Cin8$_{+28NC}$Kin to 76.6 ± 4.2 nm/s for Cin8Kin
(mean ± SEM for 95 and 106 traces, respectively; Fig. 3C).

To quantify the relative proportions of directional and nondirectional movements on yeast
microtubules, we carried out instantaneous velocity and mean squared displacement
analyses. The instantaneous velocity analyses (Fig. 3D&E) revealed clear minus-end
movements, particularly for Cin8$_{+28NC}$Kin – from the 5 second instantaneous velocity
distributions, 26.8% of segments were minus-end directed for Cin8$_{+28NC}$Kin versus 13.7%
for Cin8Kin (shaded regions). Cin8$_{+28NC}$Kin also had a broader instantaneous velocity
distribution than Cin8Kin with a standard deviation of 53.9 nm/s versus 33.4 nm/s,
respectively (Fig. 3E). From the MSD analysis (Fig. 3F&H), the apparent diffusion constants
on yeast microtubules were also similar to one another (7194 ± 94 nm^2/s for Cin8$_{+28NC}$Kin
and 6915 ± 220 nm^2/s for Cin8Kin); however fitting to MSD = aτ^α + b gave α of 1.29 for
Cin8$_{+28NC}$Kin and 1.64 for Cin8Kin, consistent with a greater degree of processive
movement for Cin8Kin (Table S1). Interestingly, despite the shift toward more consistent
plus-ended motility on yeast microtubules, motor landing rates on yeast microtubules
were lower than on bovine microtubules (Fig. 3I). Overall, on yeast microtubules the mean
velocities for each motor matched those on bovine microtubules (Fig. 3G); however, there

was less undirected motility for both motors on yeast microtubules, which revealed more

clearly the greater degree of undirected motility of Cin8$_{+28NC}$Kin relative to Cin8Kin.

Instantaneous velocity fit parameters									
	Zero centered peak			Broad peak			Dominant peak		
All nm/s	μ	σ	%	μ	σ	%	μ	σ	%
Cin8$_{+28NC}$Kin	0.6	5.0	21.8	42.5	80.2	34.3	33.5	35.6	43.8
Cin8Kin	9.7	14.3	24.7	40.5	69.5	24.3	93.2	38.9	51.0

Figure 4. Separating diffusion from bidirectional stepping. (A and B) Gaussian Mixture Model

analysis of 5 s instantaneous velocity distributions. Velocity distributions on yeast microtubules

(from Fig. 3E) were fit by three Gaussians (gray lines), with values given in table. (**C and D**)

Comparison of experimental results with simulations of a motor with a unidirectional velocity plus

random diffusion. Motor velocities were taken from mean of instantaneous velocity distributions in

Fig. 3E. Initial diffusion constants (purple line) were taken from MSD analysis in Fig. 3F and varied

up and down by two-fold. The experimental velocity distribution from Fig. 3D&E are shown in gray,

whereas the distributions shown by cyan, purple and magenta represent simulated data with

diffusion coefficients values given in figure inset. In the table, μ, σ and % represent mean, standard

deviation, and weight of each Gaussian distribution, respectively.

385
386

Instantaneous velocity distributions are not consistent with simple diffusion

To gain further insight into the degree to which the undirected movements reflected bidirectional stepping versus thermally driven diffusion, we analyzed the instantaneous velocity distributions using a Gaussian Mixture Model. We first asked whether the velocity distributions on yeast microtubules could be decomposed into sum of two normal distributions representing processive stepping and undirected diffusion or bidirectional stepping. A sum of two Gaussian did not fit the data well (Fig. S5). Instead, the velocity distributions for both Cin8Kin and Cin8$_{+28NC}$Kin were well fit by the sum of three Gaussians (Fig. 4A&B). For each variant there was a principal mode (dominant peak) corresponding to plus-ended stepping; there was a narrow mode centered around zero (zero-centered peak), and there was a third mode with a very wide distribution (broad peak). The first difference between the two dimers was that the dominant velocity peak, which accounted for approximately 50% of the total distribution in both cases, exhibited markedly different mean velocities: 33.5 nm/s for Cin8$_{+28NC}$Kin compared to 93.2 nm/s for Cin8Kin. The second difference was the width of the broad velocity peak, with Cin8$_{+28NC}$Kin having a larger standard deviation (80.2 nm/s) than Cin8Kin (69.5 nm/s) and accounting for a greater proportion of the distribution (34% versus 24%). Together, these fits reiterate the observation that Cin8$_{+28NC}$Kin exhibited greater stepping variability and reduced net plus-end velocity than Cin8Kin. In principle, the narrow modes centered around zero could reflect unbiased thermal diffusion while the broad peaks could represent bidirectional stepping with a plus-end bias. However, these data alone are not sufficient to support that interpretation. In sum, the presence of three modes in the velocity distribution argues against model consisting only of plus-end stepping and simple diffusion.

As a second test of whether simple diffusion can account for the observed undirected movements, we carried out stochastic simulations of Cin8 motility that included motor stepping and diffusion. We took the stepping rate from the mean velocity of the 5 s velocity distribution for each motor and added a diffusive term ($\langle x(t)\rangle^2 = 2Dt$), where D is the apparent diffusion constant. We generated simulated motility traces and analyzed them in the same manner as the experimental data to generate a 5 second instantaneous velocity distribution (Fig. 4 C&D). When we chose a value of D from the MSD analysis in ATP (7194 nm^2/s and 6915 nm^2/s), the resulting distributions were wider than the experimental and they also failed to capture the height of the peaks (cyan curves in Fig. 4 C&D). For both variants, a smaller D was able to reproduce the tails of the distribution, but the peaks were not aligned with the experimental peaks (purple curves in Fig. 4 C&D)). Finally, a larger D failed to capture either the peak or the tails of the distribution (magenta curves in Fig. 4 C&D). The failure of a simple plus-end directed stepping plus diffusion model to recapitulate the experimental data, together with the finding that the distributions were fit

424   by three Gaussians (Fig. 4 A&B) strongly suggest that the undirected movements result

425   from bidirectional stepping and not solely due to thermally driven diffusion of the motors.

Fig. 5. ATP turnover kinetics and corresponding stepping velocities. (A) Microtubule-stimulated
ATP hydrolysis of Cin8 dimers in BRB80 buffer at various bovine microtubule concentrations (n = 5
trials per point). Data were fit to the Michaelis-Menten equation, using 1/SEM weighing for each
point, to obtain the k_{cat} and a K_m (± 95% CI) presented in the table. Shaded area represents fit ± 95%
CI. **(B)** Comparison of k_{cat} (fit ± 95% CI; left axis, green) to the 5 s mean instantaneous velocity
(mean ± SEM; right axis, blue) for Cin8$_{+28NC}$Kin and Cin8Kin (from Fig. 2D).

**ATPase kinetics indicate that flexibility of the Cin8 neck coil diminishes
mechanochemical coupling**

Understanding how ATP hydrolysis is coupled to mechanical stepping is central to
understanding the kinesin stepping mechanism. For kinesin-1, the stepping rate (k_{step})
matches the ATP hydrolysis rate (k_{cat}), consistent with the motor hydrolyzing one ATP per
step, defined as tight mechanochemical coupling (41,42). In our single-molecule assays,
both Cin8 dimers showed net plus-end directed processive motility along with undirected
fluctuations, which in principle could result from either pure diffusion with no ATP
hydrolysis (the consensus model for processive monomeric KIF1A motors (43)) or from
bidirectional stepping (with each step consuming one ATP). Cin8$_{+28NC}$Kin, which contains its
native proximal neck-coil domain had a slower net plus-end velocity and more undirected
movement. If this undirected movement was simply diffusion, the Cin8$_{+28NC}$Kin ATPase rate
would be expected to be lower than that of Cin8Kin, reflecting the time the motor spends in

a diffusive state. In contrast, if the greater undirected movement of Cin8$_{+28NC}$Kin was due to
bidirectional stepping, the ATPase rate would be expected to be higher than Cin8Kin. To
examine the degree of mechanochemical coupling for the two Cin8 dimers, we measured
the microtubule-stimulated ATPase rates at varying concentrations of bovine microtubules
using an NADH-coupled assay (Fig. 5A). Cin8Kin had a mean plus-end velocity of 58.6 nm/s
on bovine microtubules (Fig. 2E), corresponding to 7 steps/s, and the measured k_{cat} was 5.1
± 1.5 s^{-1} (± 95% CI; Fig. 5B). This agreement is consistent with tight coupling between ATP
hydrolysis and stepping. However, the data are also consistent with the motor sharing time
between a diffusive state with no ATPase and bidirectional stepping state with an elevated
ATPase rate. In contrast, Cin8$_{+28NC}$Kin had a mean plus-end velocity of 23.9 ± 6.9 nm/s on
bovine microtubules (Fig. 2E), corresponding to 3 steps/s, and a k_{cat} of 6.9 ± 2.3 s^{-1} (± 95%
CI). The ~two-fold faster ATPase rate for Cin8$_{+28NC}$Kin compared to its stepping rate suggests
that instability of the native Cin8 neck coil leads to uncoupling between stepping and
hydrolysis. This uncoupling also suggests that the greater proportion of undirected
movement for Cin8$_{+28NC}$Kin (Fig. 2 & 3) results from bidirectional stepping and not from
enhanced diffusion.

Fig. 6. Cin8 remains bound to microtubules in the presence of ADP. (A) Representative
kymographs in the presence of 1 mM ADP on yeast microtubules. **(B)** 1 s and **(C)** 5 s instantaneous
velocity distributions for Cin8$_{+28NC}$Kin (orange) and Cin8Kin (blue). **(D)** Mean squared displacement
(MSD) comparison in ADP. Data were fit to MSD = 2Dτ, yielding D = 6314 ± 17 nm^2/s for
Cin8$_{+28NC}$Kin and 8565 ± 58 nm^2/s for Cin8Kin (fit ± 95% CI). **(E)** Comparison of apparent diffusion
coefficients for Cin8$_{+28NC}$Kin and Cin8Kin in ATP and ADP on yeast microtubules.

Cin8 dimers diffuse on MTs for long durations in the ADP-bound state

In a further effort to separate out thermally-driven diffusion from bidirectional
stepping, we examined the behavior of the Cin8 dimers on yeast microtubules in the
presence of 1 mM ADP, which generates a diffusive weak-binding state. Unlike most
kinesins, which bind weakly to microtubules in ADP (28,44,45), Cin8 tetramers have been
shown to bind for long durations in ADP (3,23,25,26). We found that both Cin8 dimers
remained bound to microtubules for long durations in ADP (Fig. 6A). Thus, consistent with
previous observations (24), Cin8 motor domains have inherently high microtubule affinity
in ADP independent of their tetrameric structure or their tail domains.

To quantify motor diffusion in the ADP-bound state, we plotted the instantaneous
velocity distributions for both motors and carried out a mean squared displacement (MSD)
analysis. Both 1 s and 5 s instantaneous velocity distributions are centered around zero, as
expected, indicating that directional motility is ATP hydrolysis-dependent (Fig. 6B&C). The
apparent diffusion constants in ADP were 6314 ± 17 nm^2/s for Cin8$_{+28NC}$Kin and 8565 nm^2/s
± 58 nm^2/s Cin8Kin (fit $\pm$ 95% CI; Fig. 6D&E), which are in the same range of the
corresponding diffusion constants in ATP on yeast microtubules (Fig. 6E). Thus, the data
don't exclude the possibility that the nondirected movements in ATP are 1D diffusion of the
motors during phases when the motors are not stepping.

Fig. 5. Microtubule binding durations of Cin8 dimers on yeast microtubules. Distribution of
bound durations (τ) from single-molecule motility traces of Cin8$_{+28NC}$Kin and Cin8Kin in **(A)** ATP and
(B) ADP. Data are plotted as 1 - Cumulative distribution function (1-CDF) and fit with an exponential
(black lines). **(C)** Comparison of microtubule detachment rates (k_{off}^{Mt}), calculated as $k_{off}^{Mt} = 1/\tau$.
Sample sizes (n) for Cin8$_{+28NC}$Kin were 246 (ATP) and 190 (ADP); for Cin8Kin, 172 (ATP) and 59 (ADP).
Error bars represent 95% CI. Processive and diffusive traces in ATP were analyzed separately and
found to have similar binding durations (Fig. S6).

500 **Cin8 neck-coil domain enhances microtubule binding duration in ATP but not ADP**

501   To investigate the influence of inter-head tension on the overall microtubule affinity of
502   Cin8$_{+28NC}$Kin and Cin8Kin, we quantified the residence times of Cin8 dimers in ATP and ADP
503   on yeast microtubules. If flexibility of the Cin8 neck-coil reduces inter-head coordination
504   needed for processive stepping, then Cin8$_{+28NC}$Kin should have shorter binding durations
505   than Cin8Kin. In contrast, if flexibility of the Cin8 neck-coil reduces steric constraints and
506   allows both heads to bind to the microtubule more readily, then Cin8$_{+28NC}$Kin should have
507   longer binding durations than Cin8Kin. Mean binding durations in ATP were calculated by
508   fitting an exponential function to the distribution of run durations. We found that
509   Cin8$_{+28NC}$Kin remained bound to microtubules ~3-fold longer than Cin8Kin (26.3 ± 0.32 s
510   and 8.9 ± 0.27 s, respectively (fit ± 95% CI); Fig. 7A). Thus, the data are consistent with
511   flexibility of the Cin8 neck coil reducing steric constraints between the two heads and
512   promoting a two-head binding state in ATP. To test whether similar behavior occurs in ADP,
513   we repeated the analysis in 1 mM ADP and found the opposite result, with the Cin8$_{+28NC}$Kin
514   binding duration (10.6 ± 0.61 s) being shorter than that for Cin8Kin (15.8 ± 0.67 s; Fig. 7B &
515   C). Thus, in ATP the proximal Cin8 neck domain prolongs the microtubule binding duration
516   in ATP, whereas in ADP it shortens the binding duration.

517

Discussion

518

519   For transport kinesins, processivity relies on mechanochemical coordination
520   between the two motor domains; however, the degree to which the two heads of Cin8 are
521   coordinated and whether this coordination plays a role in the bidirectional motility of Cin8
522   are not clear. We find that engineered Cin8 dimers move with a plus-end bias and exhibit
523   undirected movements that result from a combination of ATP-driven bidirectional stepping
524   and thermally-driven diffusion. Including the native neck-coil of Cin8 leads to more
525   undirected movements and a slower net plus-end speed, no change in the ATP turnover
526   rate, and a longer residence time on the microtubule. On yeast microtubules, Cin8 moved
527   at a similar speeds as on bovine microtubules but with fewer undirected movements.
528   These results suggest that plus-ended movement of Cin8 requires little coordination
529   between the activities of its motor domains and instead relies on the high affinity of Cin8 in
530   the ADP state.

531 **Cin8 dimer motility is consistent with previous work**

532   Motility of the two Cin8 dimer variants studied here is broadly consistent with
533   previous work. Duselder et al constructed a Cin8Kin containing the full N-terminal, 4
534   residues more in the neck domain, and a shorter kinesin-1 dimerization domain (24). On

porcine microtubules in BRB80 buffer without added salt, the dimer showed bidirectional movements with a small plus-ended mean velocity (18 nm/s) and an MSD exponent of 1.24 consistent with directional movements. Our Cin8Kin dimer had a faster net velocity (62 nm/s on bovine microtubules from Fig. 2E), a slightly higher MSD exponent of 1.38, and clear plus-end directed segments not observed in that work. Although there are differences in the motility analysis between the two studies, differences in the truncation points of the constructs are the most likely explanation for the differences in motility. Consistent with this, previous work using dimers having the same motor domain length as our Cin8Kin (38-530) with a different dimerization strategy found similar plus-end gliding velocities in multi-motor gliding assays (25). One consistent result from these dimer studies was that the fast minus-end motility seen in Cin8 tetramers was absent, consistent with regions outside the motor domain being required for fast minus-end velocities seen in tetramers.

Diffusion and bidirectional stepping are not easily separated

Because bidirectional stepping is a form of a random walk and random walks inherently generate diffusion-like behavior (46), it is challenging to separate out contributions of ATP-driven bidirectional stepping from simple thermally-driven diffusion. The first argument in favor of bidirectional stepping is the substantial fraction of minus-end velocity segments in the instantaneous velocity distributions (Fig. 2E and 3E). When we analyzed whether a model consisting of plus-ended stepping together with thermal diffusion could recapitulate the instantaneous velocity histograms, we failed to find a diffusion constant that matched both the height of the velocity peak and the width of the overall distribution. Although we can't rule out the possibility that more complex stepping and diffusion models could better match the data, our analysis argues against diffusion alone accounting for the undirected motility.

The second argument supporting bidirectional stepping is the fit of a three-component Gaussian Mixture Model to the instantaneous velocity histograms (Fig. 4A). In all cases the fits included a dominant peak (roughly half of the total) centered around the mean plus-end velocity, consistent with active stepping. There was also a peak near zero (~20-25% of total), consistent with thermally-driven diffusion. The third peak was broad and centered at a positive velocity, properties consistent with it representing bidirectional stepping with a plus-end bias. The fact that this broad peak represented 34% of the data for Cin8$_{+28NC}$Kin but only 24% for Cin8Kin is consistent with the native Cin8 neck coil leading to more bidirectional stepping.

The third argument supporting bidirectional stepping as the dominant source of undirected movement is the relationship of the motor velocities to their ATPase rates. For Cin8Kin, the ATPase rate (k_{cat} of 5 s^{-1}) was similar to the net stepping rate calculated from

the mean velocity (7 s^{-1}). This match could be interpreted as ATP hydrolysis being tightly
coupled to forward stepping; however, the substantial undirected movements suggest a
more complicated picture involving some combination of bidirectional stepping and
diffusion. The more important result was that the k$_{cat}$ of Cin8$_{+28NC}$Kin (7 s^{-1}) was more than
twice the net stepping rate (3 s^{-1}). This discrepancy suggests that the observed net velocity
results from a combination ATP-driven plus- and minus-end directed steps. For instance, if
one ATP was associated with every step, the results could be explained by a model with a 5
s^{-1} plus-end stepping rate and a 2 s^{-1} minus-end stepping rate. However, the undirected
movements we observed, which lasted for multiple seconds for some motors and the
entire duration for a subset of motors, suggest that the Cin8 dimers alternate between
phases of ATP-driven stepping and thermally-driven diffusion where little or no stepping
occurs. In this case, diffusive periods with zero ATPase could be balanced by periods of
bidirectional stepping at rates substantially above the average.

Two pieces of evidence argue that at least some portion of the undirected
movements are due to thermally-driven diffusion. The first is that comparing the behavior
in ATP versus ADP, both the apparent diffusion constants and the microtubule off-rates
differ very little (Fig. 6E and 7C). Because the ADP state is generally the lowest affinity state
in the kinesin ATP hydrolysis cycle (44,47), both the diffusion constant and the dissociation
rate would be expected to be highest in ADP. One interpretation is that Cin8 spends most of
its ATP hydrolysis cycle in the ADP state, similar to what has been suggested for the
superprocessive kinesin-3 motor, KIF1A (28,48). Unfortunately, the high affinity of Cin8 for
mant-ADP (Fig. S7) prevented us from measuring the ADP off-rate directly. The second
argument that diffusion makes up part of the undirected movement is the different
behaviors on yeast versus bovine microtubules. Consistent with the general conservation
of tubulin from fungi to mammals, the mean velocities observed on the two substrates
were similar for both motors. However, there were more undirected movements on bovine
microtubules than on yeast microtubules, particularly for Cin8Kin. Motor landing rates
were also higher on bovine microtubules. We tentatively interpret the enhanced undirected
movements on bovine microtubules as resulting from more electrostatic interactions
between Cin8 and the bovine C-terminal tails, which shift the motor from a stepping state
to a diffusive state. The importance of the C-terminal tail to Cin8 motility was
demonstrated by a recent study that found that deleting the yeast β-tubulin C-terminal tail
reduces the on-rate, run length and velocity of full-length Cin8 (40).

Cin8 lacks tight inter-head coordination

In contrast to kinesin-1 where processivity is achieved by tight coordination between
the mechanochemical cycles of the two heads, we favor a model for Cin8 in which there is

little inter-head coordination and processivity is instead achieved by the inherent high
affinity of the motor domains for the microtubule. We found that in the presence of ADP,
Cin8 dimers bound to microtubules for long durations, consistent with previous
observations for tetrameric, dimeric, and monomeric Cin8 (3-5,24,25). This property differs
from the relatively fast microtubule off-rate of kinesin-1 in ADP state (44), and it relieves the
constraint inherent in kinesin-1 that the mechanochemical cycles of the heads remain out
of phase to minimize time in the ADP state (16). Notably, we found that Cin8$_{+28NC}$Kin moved
at a slower mean velocity than Cin8Kin and had more undirected motility, but it remained
bound to the microtubule nearly three-fold longer than Cin8Kin (Fig. 7A). We interpret this
to mean that the relative instability of the native Cin8 neck-coil domain diminishes
mechanical coupling between the two motor domains, resulting in Cin8$_{+28NC}$Kin spending a
greater fraction of time in the two-heads-bound state than Cin8Kin. It was shown
previously that disrupting the mechanical connection between the two heads of kinesin-1
by inserting six or more residues into the neck linker domain resulted in significant
backstepping under hindering loads, longer residence times under no load, and the motor
losing the requirement of ATP binding to occupy the two-heads-bound state (21,22). Thus,
there is precedence for a loosening of the mechanical connection between the two heads
of kinesin enabling the two-head-bound state.

Further evidence supporting a lack of interhead coupling in Cin8 come from
comparisons of our dimers to previous work on Cin8 monomers. It was found previously
that in multimotor gliding assays, monomeric Cin8 motor domains (residues 1-528 linked
to gelsolin for surface immobilization) moved microtubules at 20-40 nm/s, similar to our
gliding speeds for dimeric Cin8$_{+28NC}$Kin (Fig. 1) (49). Furthermore, in ATPase assays Cin8
monomers (residues 38-530) were found to have a k_{cat} of 4.1 s^{-1} (25), which is roughly half of
our dimer ATPase rates (meaning the rates are the same on a per-head basis). Thus,
multimotor gliding speeds and ATPase rates appear to be equivalent whether the heads are
connected through the Cin8 neck-coil or not. Notably, in Cin8Kin where we connected two
Cin8 heads directly through the stable kinesin-1 neck-coil domain, the motor moved faster
and with more plus-end directionality than when the Cin8 neck-coil was present. These
results suggest that features that regulate Cin8 directionality, such as the tail domain (24),
N-terminal extension (26), loop-8 (25) and phosphorylation (50,51) could exert their effects
by altering inter-head coordination. Similarly, geometries that promote plus-end
directionality such as motor clustering (4,23) and crosslinking of antiparallel microtubules
(3,10) could be thought of as mechanically coordinating the motor domains.

Author contributions

H.P., W.O.H. and L.G. developed original ideas for the project. H.P. and W.O.H. designed research. H.P. and T-C.M. carried out experiments and analyzed data. T-C.M. carried out simulations. H.P., T-C.M., L.G. and W.O.H. interpreted data. E.B. and L.M.R. generated yeast tubulin and helped design experiments. H.P. and W.O.H. wrote the manuscript and all authors edited the manuscript.

Declaration of interests

The authors declare no competing interests.

Acknowledgments

This work was funded by NIH R35 GM139568 awarded to W.O.H., NIH R35 GM156385 awarded to L.M.R., and ISF grant 629/23 awarded to L.G. The authors thank members of the Hancock and Gheber labs for helpful suggestions.

References

1. Henningsen, U., and Schliwa, M. (1997) Reversal in the direction of movement of a molecular motor. *Nature* **389**, 93-96
2. Endow, S. A., and Waligora, K. W. (1998) Determinants of kinesin motor polarity. *Science* **281**, 1200-1202
3. Gerson-Gurwitz, A., Thiede, C., Movshovich, N., Fridman, V., Podolskaya, M., Danieli, T., Lakamper, S., Klopfenstein, D. R., Schmidt, C. F., and Gheber, L. (2011) Directionality of individual kinesin-5 Cin8 motors is modulated by loop 8, ionic strength and microtubule geometry. *EMBO J* **30**, 4942-4954
4. Roostalu, J., Hentrich, C., Bieling, P., Telley, I. A., Schiebel, E., and Surrey, T. (2011) Directional switching of the kinesin Cin8 through motor coupling. *Science* **332**, 94-99
5. Fridman, V., Gerson-Gurwitz, A., Shapira, O., Movshovich, N., Lakamper, S., Schmidt, C. F., and Gheber, L. (2013) Kinesin-5 Kip1 is a bi-directional motor that stabilizes microtubules and tracks their plus-ends in vivo. *J Cell Sci* **126**, 4147-4159
6. Endow, S. A., and Higuchi, H. (2000) A mutant of the motor protein kinesin that moves in both directions on microtubules. *Nature* **406**, 913-916
7. Popchock, A. R., Tseng, K. F., Wang, P., Karplus, P. A., Xiang, X., and Qiu, W. (2017) The mitotic kinesin-14 KlpA contains a context-dependent directionality switch. *Nature communications* **8**, 13999

8.   Mann, B. J., and Wadsworth, P. (2019) Kinesin-5 Regulation and Function in Mitosis.
Trends Cell Biol **29**, 66-79
9.   Pandey, H., Popov, M., Goldstein-Levitin, A., and Gheber, L. (2021) Mechanisms by
Which Kinesin-5 Motors Perform Their Multiple Intracellular Functions. *Int J Mol Sci*
22
10.   Shapira, O., Goldstein, A., Al-Bassam, J., and Gheber, L. (2017) A potential
physiological role for bi-directional motility and motor clustering of mitotic kinesin-5
Cin8 in yeast mitosis. *J Cell Sci* **130**, 725-734
11.   Britto, M., Goulet, A., Rizvi, S., von Loeffelholz, O., Moores, C. A., and Cross, R. A.
(2016) Schizosaccharomyces pombe kinesin-5 switches direction using a steric
blocking mechanism. *Proc Natl Acad Sci U S A* **113**, E7483-E7489
12.   Rice, S., Lin, A. W., Safer, D., Hart, C. L., Naber, N., Carragher, B. O., Cain, S. M.,
Pechatnikova, E., Wilson-Kubalek, E. M., Whittaker, M., Pate, E., Cooke, R., Taylor, E.
W., Milligan, R. A., and Vale, R. D. (1999) A structural change in the kinesin motor
protein that drives motility. *Nature* **402**, 778-784
13.   Vale, R. D., and Milligan, R. A. (2000) The way things move: looking under the hood of
molecular motor proteins. *Science* **288**, 88-95
14.   Hwang, W., Lang, M. J., and Karplus, M. (2008) Force generation in kinesin hinges on
cover-neck bundle formation. *Structure* **16**, 62-71
15.   Khalil, A. S., Appleyard, D. C., Labno, A. K., Georges, A., Karplus, M., Belcher, A. M.,
Hwang, W., and Lang, M. J. (2008) Kinesin's cover-neck bundle folds forward to
generate force. *Proc Natl Acad Sci U S A* **105**, 19247-19252
16.   Hancock, W. O. (2016) The Kinesin-1 Chemomechanical Cycle: Stepping Toward a
Consensus. *Biophys J* **110**, 1216-1225
17.   Gennerich, A., and Vale, R. D. (2009) Walking the walk: how kinesin and dynein
coordinate their steps. *Curr Opin Cell Biol* **21**, 59-67
18.   Shastry, S., and Hancock, W. O. (2011) Interhead tension determines processivity
across diverse N-terminal kinesins. *Proc Natl Acad Sci U S A* **108**, 16253-16258
19.   Niitani, Y., Matsuzaki, K., Jonsson, E., Vale, R. D., and Tomishige, M. (2025) Kinetic
regulation of kinesin's two motor domains coordinates its stepping along
microtubules. *eLife* **14**
20.   Shastry, S., and Hancock, W. O. (2010) Neck linker length determines the degree of
processivity in Kinesin-1 and Kinesin-2 motors. *Current Biology* **20**, 939-943
21.   Yildiz, A., Tomishige, M., Gennerich, A., and Vale, R. D. (2008) Intramolecular strain
coordinates kinesin stepping behavior along microtubules. *Cell* **134**, 1030-1041
22.   Clancy, B. E., Behnke-Parks, W. M., Andreasson, J. O., Rosenfeld, S. S., and Block, S.
M. (2011) A universal pathway for kinesin stepping. *Nat Struct Mol Biol* **18**, 1020-
1027
23.   Pandey, H., Reithmann, E., Goldstein-Levitin, A., Al-Bassam, J., Frey, E., and Gheber,
L. (2021) Drag-induced directionality switching of kinesin-5 Cin8 revealed by cluster-
motility analysis. *Sci Adv* **7**
24.   Duselder, A., Fridman, V., Thiede, C., Wiesbaum, A., Goldstein, A., Klopfenstein, D.
R., Zaitseva, O., Janson, M. E., Gheber, L., and Schmidt, C. F. (2015) Deletion of the

Tail Domain of the Kinesin-5 Cin8 Affects Its Directionality. *J Biol Chem* **290**, 16841-
16850

25.   Pandey, H., Singh, S. K., Sadan, M., Popov, M., Singh, M., Davidov, G., Inagaki, S., Al-
Bassam, J., Zarivach, R., Rosenfeld, S. S., and Gheber, L. (2021) Flexible microtubule
anchoring modulates the bi-directional motility of the kinesin-5 Cin8. *Cell Mol Life*
Sci **78**, 6051-6068

26.   Singh, S. K., Siegler, N., Pandey, H., Yanir, N., Popov, M., Goldstein-Levitin, A.,
Sadan, M., Debs, G., Zarivach, R., Frank, G. A., Kass, I., Sindelar, C. V., Zalk, R., and
Gheber, L. (2024) Noncanonical interaction with microtubules via the N-terminal
nonmotor domain is critical for the functions of a bidirectional kinesin. *Sci Adv* **10**,
eadi1367

27.   Mickolajczyk, K. J., Deffenbaugh, N. C., Ortega Arroyo, J., Andrecka, J., Kukura, P.,
and Hancock, W. O. (2015) Kinetics of nucleotide-dependent structural transitions
in the kinesin-1 hydrolysis cycle. *Proc Natl Acad Sci U S A* **112**, E7186-7193

28.   Zaniewski, T. M., Gicking, A. M., Fricks, J., and Hancock, W. O. (2020) A kinetic
dissection of the fast and superprocessive kinesin-3 KIF1A reveals a predominant
one-head-bound state during its chemomechanical cycle. *J Biol Chem* **295**, 17889-
17903

29.   Cleary, J. M., Kim, T., Cook, A. S. I., McCormick, L. A., Hancock, W. O., and Rice, L.
M. (2022) Measurements and simulations of microtubule growth imply strong
longitudinal interactions and reveal a role for GDP on the elongating end. *eLife* **11**,
e75931

30.   Mahamdeh, M., Simmert, S., Luchniak, A., Schaffer, E., and Howard, J. (2018) Label-
free high-speed wide-field imaging of single microtubules using interference
reflection microscopy. *J Microsc* **272**, 60-66

31.   Ruhnow, F., Zwicker, D., and Diez, S. (2011) Tracking single particles and elongated
filaments with nanometer precision. *Biophys J* **100**, 2820-2828

32.   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J.,
Hartenstein, V., Eliceiri, K., Tomancak, P., and Cardona, A. (2012) Fiji: an open-
source platform for biological-image analysis. *Nat Methods* **9**, 676-682

33.   Huang, T. G., and Hackney, D. D. (1994) Drosophila kinesin minimal motor domain
expressed in Escherichia coli. Purification and kinetic characterization. *J Biol Chem*
269, 16493-16501

34.   Ohashi, K. G., Han, L., Mentley, B., Wang, J., Fricks, J., and Hancock, W. O. (2019)
Load-dependent detachment kinetics plays a key role in bidirectional cargo
transport by kinesin and dynein. *Traffic* **20**, 284-294

35.   Ma, T. C., Gicking, A. M., Feng, Q., and Hancock, W. O. (2023) Simulations suggest
robust microtubule attachment of kinesin and dynein in antagonistic pairs. *Biophys*
J **122**, 3299-3313

36.   Chen, Y., and Hancock, W. O. (2015) Kinesin-5 is a microtubule polymerase. *Nature*
communications **6**, 8160

37. Chen, G. Y., Mickolajczyk, K. J., and Hancock, W. O. (2016) The Kinesin-5
Chemomechanical Cycle Is Dominated by a Two-heads-bound State. *J Biol Chem*
291, 20283-20294
38. Chen, G. Y., Cleary, J. M., Asenjo, A. B., Chen, Y., Mascaro, J. A., Arginteanu, D. F. J.,
Sosa, H., and Hancock, W. O. (2019) Kinesin-5 Promotes Microtubule Nucleation
and Assembly by Stabilizing a Lattice-Competent Conformation of Tubulin. *Curr Biol*
29, 2259-2269 e2254
39. Tripet, B., Vale, R. D., and Hodges, R. S. (1997) Demonstration of coiled-coil
interactions within the kinesin neck region using synthetic peptides. Implications for
motor activity. *J Biol Chem* **272**, 8946-8956
40. Thomas, E. C., and Moore, J. K. (2025) Selective regulation of kinesin-5 function by
beta-tubulin carboxy-terminal tails. *J Cell Biol* **224**
41. Coy, D. L., Wagenbach, M., and Howard, J. (1999) Kinesin takes one 8-nm step for
each ATP that it hydrolyzes. *J Biol Chem* **274**, 3667-3671
42. Schnitzer, M. J., and Block, S. M. (1997) Kinesin hydrolyses one ATP per 8-nm step.
Nature **388**, 386-390
43. Okada, Y., and Hirokawa, N. (2000) Mechanism of the single-headed processivity:
diffusional anchoring between the K-loop of kinesin and the C terminus of tubulin.
Proc Natl Acad Sci U S A **97**, 640-645
44. Hancock, W. O., and Howard, J. (1999) Kinesin's processivity results from
mechanical and chemical coordination between the ATP hydrolysis cycles of the
two motor domains. *Proc Natl Acad Sci U S A* **96**, 13147-13152
45. Chen, G. Y., Arginteanu, D. F., and Hancock, W. O. (2015) Processivity of the kinesin-
2 KIF3A results from rear head gating and not front head gating. *J Biol Chem* **290**,
10274-10294
46. Berg, H. C. (1993) *Random Walks in Biology*, 2nd ed., Princeton University Press,
Princeton, NJ
47. Romberg, L., and Vale, R. D. (1993) Chemomechanical cycle of kinesin differs from
that of myosin. *Nature* **361**, 168-170
48. Zaniewski, T. M., and Hancock, W. O. (2023) Positive charge in the K-loop of the
kinesin-3 motor KIF1A regulates superprocessivity by enhancing microtubule affinity
in the one-head-bound state. *J Biol Chem* **299**, 102818
49. Yamagishi, M., Maruyama, Y., Sugawa, M., and Yajima, J. (2021) Characterization of
the motility of monomeric kinesin-5/Cin8. *Biochem Biophys Res Commun* **555**, 115-
120
50. Avunie-Masala, R., Movshovich, N., Nissenkorn, Y., Gerson-Gurwitz, A., Fridman, V.,
Koivomagi, M., Loog, M., Hoyt, M. A., Zaritsky, A., and Gheber, L. (2011) Phospho-
regulation of kinesin-5 during anaphase spindle elongation. *J Cell Sci* **124**, 873-878
51. Shapira, O., and Gheber, L. (2016) Motile properties of the bi-directional kinesin-5
Cin8 are affected by phosphorylation in its motor domain. *Sci Rep* **6**, 25597

Supporting information

Dimeric Cin8 motors have an inherent plus-end bias and weak inter-head coordination

Running title: Cin8 dimer motility

Himanshu Pandey[1], Tzu-Chen Ma[1], Eric Bonventre[2], Luke M. Rice[2], Larisa Gheber[3], and William O. Hancock[1,4] [*]

[1] Department of Biomedical Engineering, Pennsylvania State University, University Park, Pennsylvania, USA
[2] Department of Biophysics, University of Texas Southwestern Medical Center, Dallas, Texas, USA
[3] Department of Chemistry, Ben-Gurion University of the Negev, Israel
[4] Department of Chemistry, Pennsylvania State University, University Park, Pennsylvania, USA

[*] Corresponding author:
William O. Hancock
Email: woh1@psu.edu

Fig. S1. Cin8 is predicted to have a less stable neck coiled-coil than kinesin-1. (A, B) Coiled-coil prediction models for the proximal neck coil domains of kinesin-1 (A) and Cin8 (B) generated using CCbuilder 2.0 (1). The predicted coiled-coil stability scores were −81.8 for kinesin-1 and −51.1 for Cin8, with more negative values indicating greater stability. The backbones of the interacting helices are shown in blue and red, with side chains depicted in grey. **(C, D)** Predicted stabilizing interactions within the coiled-coil domains of kinesin-1 (C) and Cin8 (D). For kinesin-1, three strong (solid lines) and three intermediate (dashed lines) stabilizing interactions were identified, whereas only three intermediate interactions were predicted for Cin8. Coiled-coil predictions and alignment were performed using CCbuilder 2.0, COcoNat (2), T-Coffee (3), and Waggawagga (4).

Fig. S2: Microtubule polarity dermination. To define microtubule polarity, fluorescent KIF1A (5) was introduced at the end of the experiment (top). Kymographs at bottom are same microtubule showing Qdot-labeled Cin8$_{+28NC}$Kin or Cin8Kin motility, with microtubule polarity marked at the bottom of kymograph.

Fig. S3: Single-molecule motility of Cin8 dimers on bovine microtubules in BRB80 buffer plus 110 mM KCl. Net plus-end motility was retained even at elevated ionic strength with whole trace velocities of 47.3 ± 7.0 and 84.1 ± 7.6 nm/s for Cin8$_{+28NC}$Kin (n = 27) and Cin8Kin (n = 30), respectively (mean ± SEM).

Fig. S4: Comparison of Cin8 dimer motility on bovine and yeast microtubules. (A) Representative kymographs for Cin8$_{+28NC}$Kin and Cin8Kin motility on bovine or yeast microtubules as indicated at the top. On yeast microtubules Cin8 dimers move with more consistent plus-end directionality. **(B)** Fraction of processive traces for each condition, defined as traces with overall velocity >20 nm/s and overall displacement >1 µm. The grey area at the top of each bar represents fraction of motors with undirected motility.

Fig. S5. Velocity distributions for Cin8$_{+28NC}$Kin (A) and Cin8Kin (B) could not be fit well by the sum of two Gaussians. The velocity distributions are depicted in orange (Cin8$_{+28NC}$Kin) or blue (Cin8Kin) and the Gaussian Mixture Model (GMM) fit in black. The Bayesian Information Criterion (BIC) of 3-Gaussian and 2-Gaussian models for Cin8$_{+28NC}$Kin are $1.0132*10^5$ and $1.0180*10^5$, respectively. For Cin8Kin, the BIC of 3-Gaussian and 2-Gaussian models are $9.2865*10^4$ and $9.3623*10^4$, respectively. These results suggest that the distributions are better explained by three Gaussians.

Fig. S6. Microtubule binding durations of processive and undirected Cin8 dimers on yeast microtubules in ATP. Processive molecules were defined as those with average velocities >20 nm/s and net displacements >1 μm. Cumulative distribution functions (1- CDF) of run durations for **(A)** processive and **(B)** undirected motility traces for Cin8$_{+28NC}$Kin and Cin8Kin were fit by single exponential functions to obtain the mean run duration, τ, for each.

Fig. S7. Cin8 has a high affinity for mant-ADP. (A) To measure microtubule-stimulated mant-ADP release, mant-ADP labeled Cin8Kin dimers were flushed against microtubules in the presence of varying concentrations of free mant-ADP. Experiments were carried out at 21 °C in BRB80 buffer using an Applied Photophysics SX20 spectrofluorometer at 356-nm excitation with a 395 long-pass emission filter. The conditions in the chamber are 5 μM microtubules, 0.2 μM Cin8, and mADP or

ATP as noted. Note that the fall in fluorescence, corresponding to mantADP release is small and concentration dependent, denoting incomplete mantADP release. Full mantADP release was achieved by flushing mADP labelled motors against 5 µM microtubules + 1 mM ATP (chamber concentrations). To determine the mantADP affinity of microtubule-bound Cin8, the fall in the fluorescence amplitude was plotted against the free mant-ADP concentration and fit using $f(\text{mADP}) = A * \left(1 - \frac{[\text{mADP}]}{[\text{mADP}] + K_D}\right) + c$, where A is the maximum amplitude, c is an offset, and K_D is the dissociation constant. The fit estimated K_D^{mADP} = 1 µM for microtubule-bound Cin8. **(B)** A competition assay between mant-ADP and ATP in the absence of microtubules was performed by flushing microtubule-bound mant-ADP labeled Cin8 monomer against varying concentrations of ATP. The amplitude of the fall in fluorescence is plotted against [ATP] and fit using $f(\text{ATP}) = A *$

$$\left(1 - \frac{[\text{mADP}]}{[\text{mADP}] + \frac{K_D^{mADP}}{K_D^{ATP}} * [ATP]}\right) + c,$$ where A is the maximum amplitude and c is the offset. The resulting

fit gave $K_D^{ATP} = 122 * K_D^{mant-ADP}$. Thus, the affinity for mantADP is 122 times tighter than the affinity for ATP. This complicates mantADP stopped flow experiments with Cin8.

Fit: MSD = aτ^α + b	Bovine MTs			Yeast MTs		
	a	α	b	a	α	b
Cin8Kin	$2.03*10^4$ $\pm0.04*10^4$	**1.38** **±0.01**	5034 ±550	$1.16*10^4$ $\pm0.03*10^4$	**1.64** **±0.01**	1727 ±435
Cin8$_{+28NC}$Kin	$2.15*10^4$ $\pm0.04*10^4$	**1.29** **±0.01**	2278 ±550	$1.17*10^4$ $\pm0.03*10^4$	**1.29** **±0.01**	1844 ±323

Table S1. Fits to MSD results in Fig. 2F and 3F where α=1 represents diffusion and α=2 represents processive movement.

Supplementary references:

1. Wood, C. W., and D. N. Woolfson. 2018. CCBuilder 2.0: Powerful and accessible coiled-coil modeling. Protein Sci 27:103-111.
2. Madeo, G., C. Savojardo, M. Manfredi, P. L. Martelli, and R. Casadio. 2023. CoCoNat: a novel method based on deep learning for coiled-coil prediction. Bioinformatics 39.
3. Notredame, C., D. G. Higgins, and J. Heringa. 2000. T-Coffee: A novel method for fast and accurate multiple sequence alignment. J Mol Biol 302:205-217.
4. Simm, D., K. Hatje, and M. Kollmar. 2015. Waggawagga: comparative visualization of coiled-coil predictions and detection of stable single alpha-helices (SAH domains). Bioinformatics 31:767-769.
5. Zaniewski, T. M., A. M. Gicking, J. Fricks, and W. O. Hancock. 2020. A kinetic dissection of the fast and superprocessive kinesin-3 KIF1A reveals a predominant one-head-bound state during its chemomechanical cycle. J Biol Chem 295:17889-17903.