Dehydrating microhabitats increase mite activity and intensify ectoparasitism of

Drosophila

Joshua B. Benoit[1], Gabrielle LeFevre, Joy Bose, Ann Miller, David Lewis, Hailie Talbott,

Chandrima Das, Emily Susanto, Lyn Wang, Oluwaseun M. Ajayi, Shyh-Chi Chen, and

Michal Polak

Department of Biological Sciences, University of Cincinnati, Cincinnati, OH 45211

[1]Author for correspondence: Joshua B. Benoit, Department of Biological Sciences,

University of Cincinnati, OH 45221, USA.

E-mail address: joshua.benoit@uc.edu (J.B. Benoit).

Abstract – Parasites interact with their host in variable environments that are often

subject to water scarcity and dehydration. *Drosophila* fruit flies and associated

ectoparasitic mites interact across a range of microhabitats, typically in decaying organic

matter, such as fallen fruit and cactus tissue, that dries out and deteriorates over periods

ranging from days to months. Here, we report that mite parasitism of *Drosophila* increases

with exposure to increasingly dry conditions for two fly-mite systems (*D. nigrospiracula-*

Macrocheles and *D. melanogaster-Gamasodes*). In *D. melanogaster*, artificial selection

for behavioral resistance did not eliminate this effect, as previously selected lines

remained relatively more resistant than non-selected controls even under dry conditions.

Water balance assays confirmed that mites became dehydrated when held under dry

conditions, which was also associated with increased mite activity. Exposure of *D.*

melanogaster to mites dehydrated by exposure to low relative humidity increased

parasitism, further supporting that mite infestation intensifies under dry conditions. The

results indicate that ectoparasitism in this system is affected by the water content of the

mites. The increased motivation of mites to parasitise flies under dry conditions may serve

to replenish mite water stores and facilitate dispersal to more favorable microhabitats.

Keywords: fly, mite, dehydration, parasitism, activity, resistance

Introduction

A diverse array of parasites exploit *Drosophila* fruit flies as hosts, and many of these associations are well-suited for experimental studies of host-parasite interactions. In particular, naturally occurring associations between *Drosophila* and ectoparasitic mites (Polak and Markow 1995; Halliday 2000; Halliday et al. 2005; Perez-Leanos et al. 2017; Yao et al. 2020) serve as tractable models for investigating the behavioral mechanisms insects use to resist infestation–mechanisms that are well characterized in vertebrate animals (Hart and Hart 2018). In general, mites are known to benefit from attaching to insect hosts through acquiring nutrients, increasing reproduction, and facilitating dispersal to new habitats (Walter and Proctor 2013). In *Drosophila*, mite attachment harms the host through direct damage to the cuticle and resource extraction, triggering wound repair, melanotic responses, and other immune responses (Benoit et al. 2020; Polak et al. 2025). The damage caused by parasitism can be severe with negative consequences for host reproduction and lifespan (Polak and Markow 1995; Polak 1996). Parasitism by mites is also associated with a suite of behavioral and physiological changes in the host, including upregulation of male reproductive effort (Polak and Starmer 1998), diminished copulation success owing to direct mechanical interference by attached mites (Polak 2003), and increased metabolic rate (Brophy and Luong 2021).

Drosophila melanogaster* and other fruit fly species are known to be parasitised by *Gamasodes* mites, with field records from Asia (Taiwan and Thailand) and Australia (Halliday et al. 2005; Yao et al. 2020; Polak 2025; Polak et al. 2025). In North America, the mite *Macrocheles subbadius* parasitizes *Drosophila nigrospiracula* in the Sonoran Desert (Polak and Markow 1995). This broad geographic distribution of fly–mite

associations, together with their occurrence across diverse ecological settings, suggests

that ectoparasitic mites exert strong selective pressures on multiple aspects of fly biology.

Flies avoid infestation by ectoparasitic mites by deploying rapid bursts of flight from the

substrate, evasive locomotor maneuvers, and vigorous grooming once mite contact is

made (Polak 2003; Polak et al. 2023). Behavioral resistance to mites is significantly

heritable (Polak 2003; Luong and Polak 2007) and can be subject to strong selection

(Polak and Markow 1995; Polak 1996; Benoit et al. 2025). Little is known about how

environmental conditions experienced by mites and flies influence the outcome of these

interactions.

Exposure to dry conditions and the prevention of dehydration are significant factors

influencing the survival and proliferation of terrestrial arthropods (Benoit et al. 2023).

When flies and mites interact, the water content of the local substrate, such as decaying

fruits, flowers, cacti, or mushrooms, often declines, eventually reaching levels that cannot

sustain many small arthropods, such as *Gamasodes* or *Macrocheles,* either due to this

declining water content of the substrate or the availability of prey. These observations

suggest that mites may increase their rates of parasitism on fruit flies when chronically

exposed to dry off-host conditions, both to obtain water and nutrients from the fly and to

disperse to more favorable microhabitats. For example, one previous study of the

cactophilic fruit fly species *D. nigrospiracula*, found that both the intensity and prevalence

of *M. subbadius* mite parasitism increased with the age and deterioration of necrotic

saguaro cactus (*Carnegiea gigantea*) tissue, likely due to a combination of desiccation of

the necrosis and declining habitat suitability for mite foraging and reproduction (Polak and

Markow 1995). Several other studies suggest that ectoparasites and parasitoids increase

their feeding and parasitism rates when exposed to dry conditions (Hagan et al. 2018;

Bezerra Da Silva et al. 2019; Abu et al. 2024; Holmes et al. 2025). However, such

interactions have yet to be directly explored in *Drosophila*–mite systems.

The present study investigated how dry conditions influence interactions between

Drosophila fruit flies and their ectoparasitic mites. Our experiments, which examined two

distinct fly–mite systems, *D. nigrospiracula-Macrocheles* and *D. melanogaster-*

Gamasodes, demonstrated that dehydration increases mite parasitism in both systems.

In *D. melanogaster*, flies that were previously artificially selected for parasite resistance

(Polak et al. 2023; Benoit et al. 2025) exhibited greater resistance to mite attachment than

non-selected flies under both dry and moist conditions, and both groups became more

heavily parasitized when mites had been exposed to dry conditions. Thus, dry conditions

overall significantly increased parasitism rate. Water balance assays revealed that mites

held under dry conditions exhibited reduced water content, which correlated with

heightened mite activity levels. Importantly, exposure of *D. melanogaster* to dehydrated

mites resulted in increased parasitism of flies, supporting a link between substrate quality,

mite physiological state, and parasitism. Our findings suggest that dry conditions directly

increase mite motivation to parasitize fruit flies, which are likely to obtain water and

nutrients from their hosts and be carried to more favorable microhabitats.

Materials and methods

<u>Flies, mites and selection for mite resistance</u>

A population of *Drosophila melanogaster* Meigen (Canton-S, number 64349) was

obtained from the Bloomington *Drosophila* Stock Center and used in all experiments with

this species described here, except those involving lines artificially selected for mite

resistance. The selected lines were derived from a single outbred base population

established with field-collected flies from Cape Tribulation, Australia (Polak et al. 2023).

Artificial selection for increasing resistance, described in detail elsewhere (Polak 2003;

Polak et al. 2023), was conducted using mites from a laboratory culture established with

individuals harvested from flied-caught flies at Cape Tribulation (Polak et al. 2023). The

selection protocol generated three replicate lines of *D. melanogaster*, each showing

significantly greater resistance than its paired unselected control line (Polak et al. 2023).

For the present study, flies were used after 22–24 generations of selection. Cultures of

Drosophila nigrospiracula Patterson and Wheeler and their ectoparasitic mites

Macrocheles subbadius Berlese were established from collections on *Carnegiea gigantea*

cacti near Phoenix, Arizona, USA.

Drosophila melanogaster was cultured in 4 half-pint glass bottles per generation

containing a cornmeal-agar food substrate (Webster and Polak 2025), while *D.*

nigrospiracula was cultured in bottles with a food substrate composed of powdered potato

flakes, instant *Drosophila* medium, and necrotic cactus (Polak 1998). All fly cultures were

maintained in an environmental chamber under standard 12h light (24°C):12h dark (22°C)

conditions. Both mite species were cultured in 5 L plastic jugs with a specialized medium

(Polak 2003), and maintained separately in environmental chambers at 12h light

(25°C):12h dark (24°C) conditions. The culture medium for both mite species consisted

of a rich organic substrate of autoclaved wheat bran and wood shavings, supplemented

with autolyzed yeast and bacteriophagic nematodes as a food source for the mites (Polak

2003; Benoit et al. 2020; Polak et al. 2023). The culture medium was moistened with

sufficient deionized water to keep it damp without puddling. Dry stock medium used in the

experiments described below consisted of autoclaved wheat bran, wood shavings, and

autolyzed yeast. Male flies were used at 7-10 days of age.

Dehydration treatment

To generate "dry" and "wet" culture conditions, a portion of the mite-containing culture

medium was replaced with either dry or hydrated stock medium. This was accomplished

by first removing ~30-40% of the culture medium and replacing it with stock medium that

was either dry or wet. The media were mixed by gently shaking the container for one

minute until the mixture was homogeneous. After 24 hours, the medium was observed

for the presence of mites. If mites were present, medium and mites were collected to

determine their water contents. Rehydrated medium was generated for a subset of the

dry treatments by adding water to the container, allowing the mites to rehydrate for 24

hours before conducting mite infestation assays.

Media water content determination

A standard soil moisture content assay (Basimike and Mutinga, 1990; Su and Puche,

2003) was used to determine the proportional change in water content of the mite media

as a result of making them either drier or wetter. One day after the addition of dry or wet

media, the top 2 cm was removed and immediately weighed on aluminum pans. Following

the determination of the initial mass (wet mass), the media were dried at 0% RH and 50°C

within a drying oven for two weeks. Following drying, they were weighed on three

consecutive days to determine that the water lost from the soil had reached the dry mass

(no additional decline in mass). Water content was determined as a percentage of the

initial mass.

<u>Mite infestation experiments</u>

Mite infestation assays followed established methods to test for differences in

susceptibility between experimental groups of flies (Polak 1996, 2003; Luong and Polak

2007; Benoit et al. 2020). Briefly, flies were placed in infestation chambers (300 mL glass

jars) containing culture medium with mites (Polak 1996, 2003; Benoit et al. 2020), similar

to the chambers used during artificial selection for increased resistance in *D.*

melanogaster (Polak et al. 2023). For both *D. melanogaster* and *D. nigrospiracula*, groups

of 40-50 flies were added to each infestation chamber; *D. melanogaster* was exposed to

G. queenslandicus mites, whereas *D. nigrospiracula* was exposed to *M. subbadius* mites.

Only male flies were used in the study.

To assess the effect of the moisture content of the substrate on infestation, assays

compared ectoparasitism rates under dry versus wet conditions across replicate

infestation chambers for each host species. Infestation chambers containing dry or wet

media were run in parallel to ensure exposure times were similar between treatments.

Approximately the same volume of media was added to the chambers to help maintain

similar numbers of mites. After exposure, flies were extracted from chambers with an

aspirator, and under CO_2, the presence of mites and scars was assessed for each fly.

Prevalence of parasitism ((scarred plus parasitized flies)/total flies exposed) was

calculated for each group.

For the *D. melanogaster* selection lines, two groups of flies were exposed to mites

in a common infestation chamber, one group from a given selected line and other from

its paired non-selected line. Group sizes were approximately equal, with a total of 40–50

flies per chamber. To distinguish the two groups, a minute wing clip was administered to

the tip of either the right or left wing (≤ 3% of wing area) while flies were under CO_2.

Prior to aspirating them into chambers, flies were allowed to recover from the CO_2 for 24

hours. Wing clips have been previously shown to have no significant effect on fly

susceptibility to mites (Polak 2003; Benoit et al. 2020). The wing that received a clip

was alternated between groups across chambers. Flies were aspirated into infestation

chambers and allowed to interact with mites for 6–8 hours in darkness, depending on

mite density in the medium. After exposure, flies were extracted from chambers with an

aspirator, and under CO_2 anesthesia, their identities ascertained by wing clips, and the

prevalence of parasitism determined as described above.

In separate assays, mites were exposed to dehydrating conditions in the absence

of culture medium. Briefly, mites were removed from the media and individually placed in

mesh-covered 0.5 ml microcentrifuge tubes. The tubes were placed at either 33% relative

humidity (RH, maintained with saturated $MgCl_2$ solution in water) or 93% RH (maintained

with saturated KNO_3 solution in water) in glass desiccators for 6-8 hours before exposing

the flies to the mites. One fly and one mite were put together in a standard *Drosophila*

vial (25 x 95 mm), which was then kept in a desiccator at 93% RH. Incidence of parasitism

was evaluated after 30 minutes.

<u>Mite water content determination</u>

Mite water content was determined using protocols previously applied to other mite

species (Benoit et al. 2008; Yoder et al. 2012). Mites (deutonymphs) were collected

directly from the media, and their initial mass was determined within five minutes after

freezing at -20°C. A Cahn microbalance was used to measure the mass of individual

mites to the nearest 0.1 μg. Dry mass was determined by drying mites at 0% relative

humidity (using $CaSO_4$, Drierite) at 50°C until stable weights were obtained over three

consecutive days; the final value was taken as the dry mass. Water content was defined

as the difference between wet and dry mass.

<u>Activity assays</u>

Mite activity was assessed following methods developed for small arthropods (Fieler et

al. 2021; Bailey et al. 2024; Benoit et al. 2025). As before, mites were individually placed

in mesh-covered 0.5 ml microcentrifuge tubes and held either at 75% RH or 93% RH for

6 hours. Groups of three mites were collected and transferred to 10-cm glass tubes, which

were sealed at both ends with foam stoppers. Tubes were placed horizontally into a

locomotor activity monitor (LAM, Trikinetics). Groups of three mites, rather than

individuals, were used to assess activity to ensure that enough mites passed through the

infrared beams that serve to detect movement (Fieler et al. 2021; Benoit et al. 2025). The

activity monitor was placed in darkness at 95-98% RH within a Percival incubator. After

a two-hour acclimation period, mite behavior was recorded for six hours. Activity was

quantified as the number of beam crossings per hour. To minimize circadian effects,

measurements were taken nightly from 8 PM to 8 AM, as mites show marked differences

in activity between day and night (Benoit et al. 2025).

Statistical analyses

R software (Version 4.3.1) was used for statistical analyses and figure development.

General linear models (binomial for proportion and gamma for non-proportion data) were

used to examine statistical differences between treatments (dry vs. wet conditions) for

each species. A Generalized Linear Mixed Model (GLMM - binomial) was used to

compare the results of selection lines in relation to hydration status. In the models, the

infestation chamber was added as a random effect to account for differences among

chambers, due, for example, to mite density variation. In all cases, chamber effects were

not significant (P > 0.2 for all models). Statistical outputs and sample sizes are provided

within the figure legends.

Results

<u>Dry conditions increase mite parasitism in two fly-mite systems</u>

When we examined the water content of wet vs. dry media, there was a significant 30-

40% decline in the water content of the media (Fig. 1). When *D. melanogaster* was

exposed to *G. queenslandicus*, a little over half of the flies became parasitized with mites

(Fig. 2A). Dehydrating conditions resulted in a significant increase in parasitism, with 25-

30% more flies harboring mites (Fig. 2A, P < 0.0001). A similar effect of medium

dehydration was noted for *D. nigrospiracula* interacting with *M. subbadius* mites (Fig. 2B;

P = 0.0002), but the only difference was that overall parasite levels were lower compared

to *D. melanogaster–G. queenslandicus* interactions. The addition of water to the dry

media resulted in a reduction in mite parasitism for *D. melanogaster* (Fig. 3, P < 0.0005),

confirming that dry conditions contributed to increased mite infestation of the flies.

Selection for behavioral resistance functions under dry conditions

A previous study performed artificial selection for increased behavioral resistance of *D.*

melanogaster against *G. queenslandicus* mites (Polak et al. 2023; Benoit et al. 2025),

generating the selected and non-selected lines used in the present study. When selected

and non-selected flies were exposed to mites in the same chamber, dry conditions

increased parasitism in both groups compared to wet conditions (Fig. 4, P < 0.005).

Selected flies in both environments maintained their resistance superiority, with non-

selected flies exhibiting 12.8% and 11.8% higher parasitism under wet and dry conditions,

respectively (Fig. 4).

Dry conditions lead to a decrease in water content in mites

We evaluated whether the water content of *Gamasodes* mites differed between wet and

dry conditions (Fig. 5). Mites exposed to dry conditions within infestation chambers had

significantly lower initial (wet) mass compared to those sampled from wet conditions (Fig.

5A; P = 0.002), but dry mass did not differ between the two groups (Fig. 5B; P = 0.374).

Mites from the dry medium had a 20–25% reduction in water content relative to those

from the wet medium (Fig. 5C; P = 0.0003). This water balance assessment confirms that

G. queenslandicus mites become dehydrated when exposed to dry conditions.

<u>Dehydrated mites exhibit increased activity and are more likely to parasitize flies.</u>

Dehydrated mites exhibited significantly higher activity levels than those from moist

media, showing a 20–30% increase in activity (Fig. 6A, P = 0.006). When mites were

exposed to the flies in a media-free chamber, the dehydrated mites were nearly 2-fold

more likely to parasitize the flies compared to hydrated mites (Fig. 6B, P = 0.012).

Discussion

This study demonstrated that dry substrate conditions are a significant factor

driving parasitism by mites. In two independent fly-mite systems, dehydration

significantly increased rates of parasitism. Interestingly, even after artificial selection for

resistance in *D. melanogaster* (Polak et al., 2023a), desiccation still enhanced

parasitism, although selected flies retained elevated resistance when compared to their

non-selected counterparts. Water balance measurements in *G. queenslandicus* mites

confirmed reduced body water content under dry conditions. Dehydrated mites also

exhibited increased locomotor activity, which likely contributed to heightened host

encounter and attachment rates. However, this increased activity is perhaps best

interpreted as part of a broader suite of mite behaviors. These include heightened host-

seeking, which directly contributes to higher parasitism rates, as well as dispersal efforts

to escape desiccating conditions through independent movement to wetter

microhabitats. Together, these findings indicate that desiccating conditions can intensify

host–parasite interactions by affecting off-host parasite behavior.

The impact of dry conditions on arthropod pests, especially ectoparasites has been

a recent focus of several studies. In mosquitoes and ticks, dry conditions increase activity,

host seeking, and feeding (Rosendale et al. 2016, 2019; Hagan et al. 2018; Manzano-

Alvarez et al. 2023; Abu et al. 2024), as well as even drive male mosquitoes to ingest

blood (Bozic et al. 2024). In addition to increased feeding, ectoparasites often exhibit

behavioral shifts in response to dehydration, including elevated general activity and

heightened movement after a bloodmeal (Rosendale et al., 2016; Holmes et al., 2025).

Similar patterns have been observed beyond vertebrate–arthropod systems; for instance,

water deprivation in insects has been linked to increased parasitism by parasitoid wasps

(Bezerra Da Silva et al., 2019). Our results align with these previous studies, showing

that two species of ectoparasitic mites that naturally parasitize fruit flies exhibited

increased rates of parasitism under dry conditions that demonstrably promote mite

dehydration. These results support the hypothesis that dehydration is a general

environmental factor capable of intensifying ectoparasitism. Importantly, the observed

increase in ectoparasitism is unlikely to be the result of mite starvation. The exposure

period was relatively short, and prior work indicates that more extended deprivation is

required to induce increased feeding due to starvation (Luong et al. 2017). Additionally,

mites likely retained access to nematode prey within the dry and wet medium, as fresh

media was added to both for only a single day. Thus, the increase in parasitism may more

plausibly be attributed to dehydration-induced increases in mite activity, which likely

enhanced host encounter rates. Further work will be required to distinguish the relative

importance of dehydration versus starvation in driving elevated parasitism.

Resistance to mites confers fitness benefits to flies by reducing the probability of

infestation (Luong and Polak 2007; Polak et al. 2023; Benoit et al. 2025). However,

resistance is not cost-free: in the absence of parasites, elevated resistance may incur

fitness costs, such as reduced stress tolerance and diminished fecundity, as documented

in this and other host–parasite systems (Ewald 1995; Windsor 1998; Fitze et al. 2004;

Luong and Polak 2007). These costs are usually more muted than the direct fitness

consequences of parasitism, which can lead to strong physiological declines and

increased probability of death (Polak and Markow 1995; Benoit et al. 2020; Polak et al.

2023). However, little is known about how the outcomes of host-ectoparasite interactions

shift under suboptimal conditions. In the present study, we demonstrated that dry

conditions elevated infestation rates in both selected and non-selected flies, indicating an

effect of substrate humidity on the prevalence of parasitism. Interestingly, selected lines

retained an advantage over their non-selected counterparts under both wet and dry

conditions, indicating stability of the genetic effect on behavioral immunity across these

environmental conditions.

The increased parasitism by mites when held under dry conditions we documented

could represent a critical mechanism for mite dispersal to more optimal habitats. Flies

utilize a wide range of substrates, ranging from fermenting fruit, cactus, and fungi, among

other organic materials (Markow and O'Grady 2006). As these organic substrates are

consumed, decompose, and become dry, particularly in areas of seasonal periods of

limited rainfall, they not only become increasingly suboptimal for fly survival and

reproduction, but these environmental effects are also felt by the mites themselves.

Depletion of available nematodes and other small invertebrate prey for the mites may

lead to mite starvation, which has previously been shown to increase mite motivation to

parasitise flies (Luong et al., 2017). Furthermore, egg viability is expected to decline under

desiccating conditions, a pattern commonly observed in mite eggs exposed to dry

environments (Benoit et al. 2007; Yoder et al. 2012), thus reducing mite survival in

microhabitats that dry as they deteriorate. Given that mites commonly parasitize flies for

many hours (Webster and Polak 2025), and that parasitism by mites improves their

fecundity (Luong and Subasinghe 2017), movement with the flies likely enhances mite

fecundity and facilitates locating new, more optimal habitats (Polak and Markow 1995).

Acknowledgments

The research was supported by National Science Foundation (NSF) grant 1654417 (to

M.P. and J.B.B.). Partial funding for reusable equipment was provided indirectly by the

National Institute of Allergy and Infectious Diseases, award numbers R01AI148551 and

R21AI166633 to J.B.B.

Figure legends

Figure 1 - Water content of mite media under wet and dry conditions. Water content of mite media in dry or wet conditions, along with the water content of the dry stock medium (N = 3 samples for each media type). $F_{2,6}$ = 648.5, P < 0.0001, all media types are significantly different (P < 0.005).

Figure 2 - Dehydration increases mite parasitism on fruit flies. A. *Drosophila melanogaster* (Canton S) exposed to mites (*Gamasodes queenslandicus*) under wet and dry conditions. N = 8 chambers per condition, P < 0.0001. B. *Drosophila nigrospiracula* exposed to mites (*Macrocheles subbadius*) under wet and dry conditions. N = 8 chambers per condition, P = 0.0002. There was a general effect where both species showed increased parasitism under dry conditions, $F_{1,30}$ = 43.64, P < 0.0001.

Figure 3 - Rehydration of media reduces parasitism to levels in wet media. *Drosophila melanogaster* (Canton S) exposed to mites (*Gamasodes queenslandicus*) under wet and dry conditions, along with those exposed to dry medium that was rehydrated for 24 hours. N = 8 chambers per treatment. An increase in parasitism was noted only under dry conditions compared to both wet and rehydrated conditions, $F_{2,21}$ = 10.81, P < 0.0005.

Figure 4 - Dehydration increases mite parasitism for mite-resistant and non-resistant lines. Both non-selected (A) and selected (B) lines were parasitized at higher rates under dry conditions compared to wet conditions (P < 0.002 for both cases).

Selected lines were more resistant than non-selected lines under both wet (P=0.001)

and dry (P=0.003) conditions. N = 12 chambers per treatment.

Figure 5 - Water content mites from wet and dry media conditions. A. Initial mass.

$F_{1,28}$ = 23.26, p = P < 0.001. B. Dry mass. $F_{1,28}$ = 0.8178, P = 0.0374. C. Water content

of mites. $F_{1,28}$ = 17.19, P = 0.001. N = 15 for mites held in wet and dry media.

Figure 6 - Activity levels and parasitism are increased in dehydrated mites. A.

Activity levels were higher in dehydrated mites compared to hydrated mites, N = 30 for

hydrated and N =28 for dehydrated mites. $F_{1,56}$ = 8.01, p = 0.006. B. Direct assessment

of mite-fly interactions confirms that dehydration triggers increased parasitism. N = 6

exposures of flies to the mites for hydrated and dehydration treatments. $F_{1,10}$ = 9.31, p =

0.012.

References

Abu AEI, Becker M, Accoti A, et al (2024) Low humidity enhances Zika virus infection and dissemination in *Aedes aegypti* mosquitoes. mSphere 9:e0040124

Bailey ST, Kondragunta A, Choi HA, et al (2024) Dehydration and tomato spotted wilt virus infection combine to alter feeding and survival parameters for the western flower thrips, *Frankliniella occidentalis*. Curr Res Insect Sci 6:100086

Benoit JB, Bose J, Ajayi OM, et al (2025) Shifted levels of sleep and activity during the night as mechanisms underlying ectoparasite resistance. NPJ Biol Timing Sleep 2:15

Benoit JB, Bose J, Bailey ST, Polak M (2020) Interactions with ectoparasitic mites induce host metabolic and immune responses in flies at the expense of reproduction-associated factors. Parasitology 147:1196–1205

Benoit JB, McCluney KE, DeGennaro MJ, Dow JAT (2023) Dehydration dynamics in terrestrial arthropods: from water sensing to trophic interactions. Annu Rev Entomol 68:129–149

Benoit JB, Yoder JA, Lopez-Martinez G, et al (2008) Adaptations for the maintenance of water balance by three species of Antarctic mites. Polar Biol 31:539–547

Benoit JB, Yoder JA, Lopez-Martinez G, et al (2007) Habitat requirements of the seabird tick, *Ixodes uriae* (Acari: Ixodidae), from the Antarctic Peninsula in relation to water balance characteristics of eggs, nonfed and engorged stages. J Comp Physiol B 177:205–215

Bezerra Da Silva CS, Price BE, Walton VM (2019) Water-deprived parasitic wasps (*Pachycrepoideus vindemmiae*) kill more pupae of a pest (*Drosophila suzukii*) as a water-intake strategy. Sci Rep 9:3592

Bozic J, Joseph RE, Krizek RS, et al (2024) Revisiting the paradigm of anhematophagy in male mosquitoes. bioRxiv. https://doi.org/10.1101/2024.10.08.617226

Brophy T, Luong LT (2021) Ectoparasite-induced increase in *Drosophila* host metabolic rate. Physiological Entomology 46:1–7

Ewald PW (1995) The evolution of virulence: a unifying link between parasitology and ecology. J Parasitol 81:659–669

Fieler AM, Rosendale AJ, Farrow DW, et al (2021) Larval thermal characteristics of multiple ixodid ticks. Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology 257:110939

Fitze PS, Tschirren B, Richner H (2004) Life history and fitness consequences of

ectoparasites. J Anim Ecol 73:216–226

Hagan RW, Didion EM, Rosselot AE, et al (2018) Dehydration prompts increased
activity and blood feeding by mosquitoes. Sci Rep 8:6804

Halliday RB (2000) The Australian species of *Macrocheles* (Acarina : Macrochelidae).
Invertebr Syst 14:273–326

Halliday RB, Walter DE, Polak M (2005) A new species of *Gamasodes* Oudemans from
Australia (Acari: Parasitidae). Zootaxa 1001:17

Hart BL, Hart LA (2018) How mammals stay healthy in nature: the evolution of
behaviours to avoid parasites and pathogens. Philos Trans R Soc Lond B Biol Sci
373:20170205

Holmes CJ, Chakraborty S, Ajayi OM, et al (2025) Multiple blood feeding bouts in
mosquitoes allow for prolonged survival and are predicted to increase viral
transmission during dry periods. iScience 28:111760

Luong LT, Brophy T, Stolz E, Chan SJ (2017) State-dependent parasitism by a
facultative parasite of fruit flies. Parasitology 144:1468–1475

Luong LT, Polak M (2007) Environment-dependent trade-offs between ectoparasite
resistance and larval competitive ability in the *Drosophila–Macrocheles* system.
Heredity 99:632–640

Luong LT, Subasinghe D (2017) A facultative ectoparasite attains higher reproductive
success as a parasite than its free-living conspecifics. Exp Appl Acarol 71:63–70

Manzano-Alvarez J, Terradas G, Holmes CJ, et al (2023) Dehydration stress and
Mayaro virus vector competence in *Aedes aegypti*. J Virol 97:e0069523

Markow TA, O'Grady P (2006) Drosophila. Academic Press, San Diego, CA

Perez-Leanos A, Loustalot-Laclette MR, Nazario-Yepiz N, Markow TA (2017)
Ectoparasitic mites and their *Drosophila* hosts. Fly 11:10–18

Polak M (2003) Heritability of resistance against ectoparasitism in the *Drosophila-*
Macrocheles system. J Evol Biol 16:74–82

Polak M (1996) Ectoparasitic effects on host survival and reproduction: the *Drosophila-*
Macrocheles association. Ecology 77:1379–1389

Polak M (2025) Personal observations

Polak M (1998) Effects of ectoparasitism on host condition in the *Drosophila-*
Macrocheles system. Ecology 79:1807

Polak M, Bose J, Benoit JB, Singh H (2023) Heritability and preadult survivorship costs

of ectoparasite resistance in the naturally occurring *Drosophila-Gamasodes* mite
system. Evolution 77:2068–2080

Polak M, Das C, Schaefer P (2025) *Drosophila* melanotic lesions caused by
ectoparasitic mites *Gamasodes pachysetis* from southern Thailand. figshare.
https://doi.org/10.6084/M9.FIGSHARE.28216868.V1

Polak M, Markow TA (1995) Effect of ectoparasitic mites on sexual selection in a
Sonoran desert fruit fly. Evolution 49:660–669

Polak M, Starmer WT (1998) Parasite–induced risk of mortality elevates reproductive
effort in male *Drosophila*. Proceedings of the Royal Society of London Series B:
Biological Sciences 265:2197–2201

Rosendale AJ, Dunlevy ME, McCue MD, Benoit JB (2019) Progressive behavioural,
physiological and transcriptomic shifts over the course of prolonged starvation in
ticks. Mol Ecol 28:49–65

Rosendale AJ, Romick-Rosendale LE, Watanabe M, et al (2016) Mechanistic
underpinnings of dehydration stress in the American dog tick revealed through
RNA-Seq and metabolomics. J Exp Biol 219:1808–1819

Walter DE, Proctor HC (2013) Mites and Biological Diversity. In: Walter DE, Proctor HC
(eds) Mites: Ecology, Evolution & Behaviour: Life at a Microscale. Springer
Netherlands, Dordrecht, pp 447–459

Webster AL, Polak M (2025) Priming for protection: Inducible attachment-resistance to
ectoparasitic mites in *Drosophila*. Parasitology 1–41

Windsor DA (1998) Controversies in parasitology, most of the species on Earth are
parasites. Controversies in parasitology, Most of the species on Earth are parasites
International Journal for Parasitology 12:1939–1941

Yao M-Y, Guo J-J, Michal P, et al (2020) A new species and new record of *Gamasodes*
(Mesostigmata: Parasitidae) from China. Systematic and Applied Acarology
25:1299–1318

Yoder JA, Hedges BZ, Benoit JB (2012) Water balance of the American dog tick,
Dermacentor variabilis, throughout its development with comparative observations
between field-collected and laboratory-reared ticks. Int J Acarology 38:334–343

Figure 1

Figure 2

A. *Drosophila melanogaster*

B. *Drosophila nigrospiracula*

Figure 3

Figure 4

A. Non-selected

B. Selected

Figure 5

Figure 6

A. Activity

B. Infection