Neurotype matching in monogamous rodents is modulated by early-life sleep experience

Lezio S. Bueno-Junior[1]*, Noah E. P. Milman[2,3], Carolyn E. Jones-Tinsley[2,3], Anjesh Ghimire[1], Peyton T. Wickham[2,3], Yujia Hu[4,5], Bing Ye[4], Miranda M. Lim[2,3], Brendon O. Watson[1]*

[1]Department of Psychiatry, University of Michigan Medical School, Ann Arbor, MI, USA

[2]Portland VA Research Foundation, Portland, OR, USA

[3]Department of Neurology, Oregon Health and Science University, Portland, OR, USA

[4]Life Sciences Institute and Department of Cell and Developmental Biology, University of Michigan, Ann Arbor, MI, USA

[5]Lerner Research Institute, Cleveland Clinic, Cleveland, OH, USA

*Correspondence: Lezio S. Bueno-Junior (soaresbl@umich.edu), Brendon O. Watson (brendonw@umich.edu)

Abstract

Studies of human sociability indicate stronger social affinity in matched-neurotype dyads (e.g., two individuals with autism or two without) compared to mixed-neurotype dyads (e.g., one individual with autism paired with one without). Is this neurotype matching phenomenon also quantifiable in non-human animals? Using deep learning tools, we analyzed dyadic male-female interactions in prairie voles, a highly social rodent species. To simulate "neurotypes", voles were exposed to either control conditions or early-life sleep disruption (ELSD) during a critical neurodevelopmental period (postnatal days 14-21), recapitulating two features of human autism: developmental sleep disruption and later-life atypical sociability. Analogous to human studies, voles showed signs of reduced social affinity in mixed dyads compared to matched dyads, including sex-specific changes in aggression and body orientation toward the conspecific. These findings advance our understanding of social affinity, providing a framework for new studies in both animal models and humans.

Introduction

During social interactions, we may experience feelings of closeness and affinity or, conversely, awkwardness and lack of affinity. Researchers have sought to quantify these feelings of "social chemistry" in humans, often through questionnaires to assess subjective perceptions about the social peers [1–6]. There have also been attempts to track these social properties more objectively in humans by measuring story recall, speech fluency, motion energy, facial expressions, mutual gaze, task cooperation and inter-brain synchrony [7–20].

Most of these human studies have focused on the smallest social unit – the dyad – to assess how the similarity between individuals affects social affinity, or rapport. A common strategy in these studies is to mix and match individuals with different "neurotypes", especially those with autism spectrum disorders (ASD) or neurotypical controls, forming dyads with matched neurotypes (neurotypical-neurotypical or ASD-ASD) or mixed neurotypes (neurotypical-ASD or vice-versa). Interestingly, mixed dyads have been associated with lower affinity than matched dyads, suggesting that it is not simply the number of neurodivergent individuals in the dyad that modulates sociability [2–6,13,15,17–19]. This aligns with the growing awareness that social difficulties in neuropsychiatric spectra may partly result from interpersonal mismatches, rather than solely from individual traits [21–23].

To explore neurotype matching biologically, we can use rodent models. However, neurotype matching research has limited correspondence in the rodent model literature, despite the existence of established rodent models in social neuroscience, as well as computational ethology tools for tracking groups of animals. Rodent models include ASD-like phenotypes or genotypes (e.g., prenatal valproic acid rat model and Shank3 knockout mouse model) [24,25], which capture some components of sociability but are often limited by the focus on just one animal per assay. In turn, computational ethology tools include whole-body or body-part keypoint trackers (e.g., JAABA, idtracker.ai, DeepLabCut, SLEAP) [26–29], which capture multi-

animal dynamics but fail to identify nuances, like the social role of each individual. Our group

has contributed innovations to both areas. For rodent modeling, we have described that early-

life sleep disruption (ELSD) impairs adult pair bonding in a monogamous rodent species: the

prairie vole (*Microtus ochrogaster*), mimicking the link between developmental sleep issues and

altered sociability in human ASD [30]. For computational ethology, we initially published a

DeepLabCut-based system to measure body orientation and distance between two spatially-

restricted prairie voles [31]. Then, more recently, we co-developed LabGym2: a postural motif

tracker that identifies social roles (e.g., "chasing" versus "being chased") in groups of freely-

moving animals, including prairie vole dyads [32].

In the present study, we integrated these innovations to quantify social roles in matched

versus mixed prairie vole dyads, drawing a parallel to neurotype matching studies in humans.

Specifically, male and female prairie vole pups were exposed to control conditions or ELSD

from postnatal days 14-21, a critical neurodevelopmental window. Upon reaching adulthood,

these voles were paired into matched or mixed neurotype combinations. Behaviors were then

analyzed using long-duration recordings, lasting several hours to days. We found that mixed

dyads exhibit altered body orientation behaviors and higher incidence of aggressive bouts,

comparable to the lower affinity found previously in mixed human dyads. Furthermore, each

behavior in prairie voles varied by sex and followed a different timeline, adding both sex

specificity and temporal resolution to the available knowledge from human dyad studies. These

findings have implications for both the basic and clinical understanding of social affinity, in

addition to offering an analytical framework for future studies in different species.

Results

Overview of animal pairing and behavioral analysis

We utilized two behavioral phenotypes: early-life sleep disrupted (ELSD) prairie voles and age-matched controls (Ctrl), including both sexes. This resulted in four dyad combinations: Ctrl-Ctrl, ELSD-ELSD, Ctrl-ELSD, ELSD-Ctrl (first and second acronyms represent male and female, respectively). The ELSD method was initially validated by our group [30] and is illustrated in **Fig 1A**. Briefly, home cages containing both parents and their pups – postnatal days (P) 14-21 – were placed on orbital shakers for gentle, intermittent agitation (see Methods). The pups were then weaned and housed with same-sex siblings under standard conditions until adulthood, generating cohorts of male and female ELSD voles. Ctrl voles underwent the same conditions, except for sleep disruption with the orbital shaker (**Fig 1A**). Our previous studies [30,33] have shown that ELSD reduces REM sleep time and fragments NREM sleep in prairie vole pups while preserving their body weight, corticosterone levels, and parental care. Subsequent to these acute changes, we have shown that ELSD triggers long-term changes in social affiliation (e.g., partner-directed huddling during tests for pair bond formation), cognitive flexibility, object interactions and neocortical cytoarchitecture, forming a phenotype relevant to human ASD [30,34].

Here, ESLD effects on dyadic behavior are explored more deeply thanks to computational ethology. We describe two strategies. Experiment 1: body-part keypoint tracking (DeepLabCut) [28] of mutually naïve male-female pairs separated by a wire mesh divider (**Fig 1B**). Experiment 2: whole-body behavioral classification (LabGym2) [32] of the same male-female pairs, but without the divider (**Fig 1C**).

Experiment 1: initial divided cohabitation (**Fig 1B**; **S1 Movie**). This experiment consisted of 72-h overhead video recordings where voles interacted through a wire-mesh divider at the

center of the cage. This was the first encounter these animals had with each other. While this

setup imposed a physical limitation on social interactions, it yielded uniform postural and spatial

measures throughout recordings. This allowed us to decompose these measures into three

DeepLabCut-derived variables: body direction to divider, distance to divider and locomotion

speed (**Fig 1B**). In a previous study [31], we characterized the sex-specific impacts of ELSD to

these variables, including alterations in male body orientation and female circadian activity.

However, we did not examine neurotype matching effects in that study; such effects are

reported here.

Experiment 2: subsequent undivided cohabitation (**Fig 1C**; **S2 Movie**). This experiment

consisted of 4-h overhead video recordings using the same voles, three to four days after

Experiment 1, meaning that the opposite-sex mates had already been exposed to each other at

the time of Experiment 2, i.e., dyad assignments were preserved between experiments (see

Methods). In this setup, the animals were allowed to engage in freely moving partner-referenced

behaviors without a divider. This resulted in an enriched behavioral repertoire, which is more

difficult to analyze with DeepLabCut but amenable to newer LabGym2-based behavioral

classification (**Fig 1C**). One novel aspect of LabGym2 – a software that we co-developed [32] –

is the ability to categorize the social role of each individual (e.g., "chasing" versus "being

chased"). We utilized this capability in combination with manual validation to ensure reliable

identity tracking and sex specification per behavior. This resulted in the labeling of five behavior

categories: *chasing*, *attacking*, *pointing*, *behaving alone*, and *huddling* (**Fig 1C**). Here, "*pointing*"

combines sniffing behaviors with a freezing-like behavior that we often observed after each

aggressive bout (see **S2 Movie**). See Methods for details.

Therefore, our two experiments yielded very different analyses. Nevertheless, neurotype

matching effects emerged in both experiments, as reported below.

Experiment 1: Mixed dyads showed exacerbated sex differences in body orientation

In Experiment 1, male and female interacted through a wire-mesh divider. **Fig 2**

describes these interactions over 72 h (1-h bins, see x-axes), with graph columns displaying

DeepLabCut-derived behavioral decomposition variables. **Fig 2A** starts by comparing sexes

irrespective of dyad grouping. When analyzing direction to divider, we found that male voles

spent significantly more time facing toward the divider than females throughout the recording.

For distance to divider, both sexes preferred areas near the divider (4-6 cm) during the initial 12

h of recording; this variable later plateaued at around 6 cm (halfway between divider and

opposite cage wall), suggesting habituation. For locomotion speed, both sexes exhibited higher

locomotion in the initial 12 h of recording; this variable later stabilized to lower basal levels, also

suggesting habituation (**Fig 2A**). These results corroborate our previous findings [31] that sex-

specific body orientation patterns persist even after voles habituate to the environment, as

indicated by the stabilization of locomotion and area preference after the initial 12 h. See **Table**

1 for statistics and **S1 Movie** for representative behaviors.

By parsing these sex differences into dyad types, we begin to reveal the novel findings

of this study. **Fig 2B** shows that while sex differences in direction to divider were significant in

both matched and mixed dyads, these differences were more pronounced in mixed dyads and

persisted for 72 h (**Table 1**). **Fig 2C** further divides male and female voles into the four juvenile

sleep status combinations. In this case, sex differences in direction to divider were significant

only in mixed dyads (Ctrl-ELSD, ELSD-Ctrl). Matched dyads, including Ctrl-Ctrl and ELSD-

ELSD, showed no significant sex differences (**Table 1**). These results indicate that body

orientation behaviors did not change linearly with the number of ELSD individuals in the dyad;

rather, neurotype matching was the main factor modulating body orientation.

Additional sex differences were specific to mixed dyads and confined to more localized

time periods. We observed that ELSD females in Ctrl-ELSD dyads maintained a greater

distance from the divider more than 36 hours into the session, while Ctrl females in ELSD-Ctrl

dyads exhibited higher locomotion speed during the initial 12 hours compared to males (**Fig**

2C). However, these sex differences were modest compared to the prominent results in

direction to divider (**Table 1**). **Fig 2D** further highlights direction to divider as a sensitive

variable, this time comparing matched and mixed dyads within each sex. Males spent

significantly more time toward the divider in mixed compared to matched dyads, indicating

heightened female-directed orientation in mixed-dyad males (**Table 1**).

Overall, **Fig 2** shows that a supra-individual factor (neurotype matching) preferentially

modulated one behavioral component (body direction) in one sex (males). Next, we describe

that neurotype matching remained an important factor in a different environment.

Experiment 2: Mixed dyads showed higher female-to-male aggression

In Experiment 2, the wire mesh divider was not present, and animals were able to

interact in an unrestricted fashion. **Fig 3** is structured similarly to the results above, with two

differences: the recordings lasted 4 h (3-min bins, see x-axes) and the graph columns show

LabGym2-based behavior categories. We again start by comparing sexes irrespective of dyad

grouping in **Fig 3A**. The social roles of *chasing* and *attacking* were significantly more prevalent

in female voles, especially in the first hour of recording (insets), after which these variables

decayed to lower basal levels. The social role of *pointing*, which included sniffing and other

behaviors (see Methods), was evenly distributed across sexes and showed a more gradual

decline over the 4-h period (see lower F values in effects of time in **Table 2**). The role-unspecific

categories of *behaving alone* and *huddling* (i.e., behaviors that considered the full dyad, see

Methods) also showed gradual time courses, but in opposite directions: *behaving alone*

decreased, while *huddling* proportionally increased over time (**Fig 3A**). In addition, *behaving*

alone and *huddling* were the most frequent behaviors across **Fig 3**, as indicated by the

probability axes. See **Table 2** for statistics.

Despite their lower occurrence rate, *chasing* and *attacking* are the behaviors with the

greatest differential trajectories across types of dyads. As shown in **Fig 3B**, female voles were

more likely to chase and attack the males in mixed than matched dyads (**Table 2**). Consistently,

in dyad subgroups (**Fig 3C**), female-to-male *chasing* and *attacking* were more frequent in both

Ctrl-ELSD and ELSD-Ctrl dyads (**Table 2**). Matched ELSD-ELSD dyads showed no sex

differences in *chasing* and a comparatively small difference in *attacking* (**Table 2**), reinforcing

that these behaviors did not change linearly with the number of ELSD individuals in the dyad,

similar to the findings in dyads behaving across a divider. Additional sex differences were

observed in *attacking* (Ctrl-Ctrl) and *pointing* (ELSD-ELSD) (**Fig 3C**), but these were also

modest compared to the sex differences in *chasing* and *attacking* behaviors found in mixed

dyads (**Table 2**). Furthermore, **Fig 3D** shows within-sex comparisons between dyad types,

confirming that mixed-dyad females were more likely to chase and attack than matched-dyad

females. No dyad type differences were observed in the other behaviors, i.e., *pointing*, *behaving*

alone and *huddling* (**Fig 3D-E**; **Table 2**), again highlighting female-to-male *chasing* and

attacking as the most sensitive behaviors in this analysis.

Therefore, neurotype matching preferentially modulated female-to-male aggression in

the undivided cage. Below we link Experiments 1 and 2 via correlations within dyads.

Linking divided cohabitation to undivided cohabitation: Neurotype matching affected

correlations among behaviors

The same dyad assignments were used in Experiment 1 (first male-female encounter

and 72-h divided cohabitation) and Experiment 2 (second encounter and 4-h undivided

cohabitation), with an interval of three to four days between experiments. This gave us the

opportunity to quantitatively link these experiments. Thus, we averaged all behavioral variables

per animal (three variables from Experiment 1, five from Experiment 2) and explored linear

correlations among such variables. **Fig 4A-B** illustrates correlations between direction to divider

in Experiment 1 and all behavioral categories of Experiment 2. Each data point in **Fig 4A-B** is

an individual animal. Separate analyses were performed per sex and major dyad type (see

Methods).

Significant correlations were restricted to female voles and differed between matched

versus mixed dyads (**Fig 4B**; see plots with gray background). *Chasing*: positively correlated

with direction to divider in mixed. *Attacking*: negatively correlated with direction to divider in

matched. *Behaving alone*: negatively correlated with direction to divider in matched. *Huddling*:

positively correlated with direction to divider in matched (**Fig 4B**). Thus, a female that oriented

more toward the divider in Experiment 1 was more likely to engage in different behaviors in

Experiment 2 depending on the dyad type: affiliative behaviors in matched, chasing behaviors in

mixed. This suggests that, depending on the dyad type, the same behavior in Experiment 1

(body orientation) may have been driven by a different underlying state. **Fig 4C** presents

correlation matrices for all variable pairs, providing a comprehensive view of both sex and dyad

type differences. The leftmost columns in each matrix of **Fig 4C** (direction to divider) contain the

correlations emphasized above. Other correlations were found within each experiment,

including negative correlations between alone and huddling, consistent with their opposing

trends in Experiment 2 (**Fig 3**).

Female-to-male behavior transitions differed between matched and mixed dyads

The analyses above ignore within-dyad actions and reactions at finer temporal scales.

How likely is behavior B in vole 2 after behavior A in vole 1? Do these likelihoods change with

dyad type and/or sex directionality (male-to-female, female-to-male)? In **Fig 5** we explore these

questions using behavior transition probabilities, thus capturing temporal sequence patterns

from each dyad type and sex directionality. Data from the video frames were divided into 1-s

time bins; this temporal resolution was deemed optimal to capture all transitions between

behaviors, including to and from aggression bouts – the quickest behaviors in our dataset (**S2**

Movie). We then conducted this analysis separately per major dyad type (matched, mixed) and

sex directionality (male-to-female, female-to-male).

Fig 5A-B shows data from Experiment 1 (divided cohabitation). The most informative

variables in that dataset – direction and distance to divider – were continuous and therefore

required discretization prior to transition analysis. Thus, we discretized direction to divider into

"toward" or "opposite", and distance to divider into "near" or "away", resulting in four categories:

toward-near, *opposite-near*, *toward-away*, *opposite-away* (see Methods). The directed graphs in

Fig 5A illustrate transitions among these categories, averaged across dyads. The edges

represent probabilities, coded by line width and grayscale (**Fig 5A**). For this analysis, we

focused on the initial 12 h of the 72-h recording, when the animals were more active (**Fig 2**).

The main visual pattern in all graphs is the convergence to, and recurrence of, *toward-*

near (black edges): these represent the highest transition probabilities in both sex

directionalities (**Fig 5A**). Particularly in the male-to-female graphs, the convergence to, and

recurrence of, *opposite-near* (dark gray edges) also showed relatively high probabilities. All

other transitions were less likely (light gray edges) (**Fig 5A**). The more distributed patterns in the

male-to-female graphs indicate higher variation: female voles were more active and rotated

themselves more (**Fig 2**, **S1 Movie**), thus increasing the chances of male behaviors being

followed by both body orientation categories in the females. In contrast, the female-to-male

graphs indicate lower variation: all female behaviors were more likely followed by *toward-near* in

the males, simply because this behavior was predominant in males (**S1 Movie**).

Of interest to this study, we found differences between matched and mixed dyads,

specifically in female-to-male transitions. For quantification, transition probability matrices were

calculated per dyad, reshaped into dyad x edge matrices (edges were used as repeated

measures), and compared between matched and mixed dyads, resulting in **Fig 5B** (self-loops

were excluded; see Methods). In mixed dyads, two female-to-male transitions were more likely:

opposite-near to *toward-near* and *toward-away* to *toward-near* (**Fig 5B**). We interpret that this

was driven by the tendency of mixed-dyad males to spend more time toward the divider, as

reported earlier (**Fig 2**). In matched dyads, three female-to-male transitions were more likely:

toward-near to *opposite-away*, *opposite-away* to *opposite-near*, and vice-versa for the latter (**Fig**

5B). These re-orientation patterns between male and female were infrequent (see light gray

edges in **Fig 5A**) but could be speculated as body posture signals. **Fig 5B** additionally contains

examples of the distributions behind each behavior pair, shown here as mean +/- standard

deviations (each data point represents a dyad). See **Table 3** for statistics.

Fig 5C-D shows the same analysis, but from the full 4 h of Experiment 2 (undivided

cohabitation), with five labeled behaviors during free interaction. As illustrated by the average

directed graphs in **Fig 5C**, the highest transition probabilities were found in the recurrence of

behaving alone and *huddling* (black self-loops). This was expected, as *behaving alone* and

huddling were both prevalent and self-sustaining, as shown previously (**Fig 3**). For

quantification, we excluded such loops to focus on the behavior pairs, as described above,

resulting in **Fig 5D**. In mixed dyads, two female-to-male transitions were more likely: *huddling-*

to-chasing to *alone-to-pointing* (**Fig 5D**). This could reflect higher activity overall in mixed dyads,

including shorter and more intermittent periods of huddling and more frequent alone-to-pointing

re-encounters. However, these transitions showed very low probabilities (hardly visible in **Fig**

5C; see also y-axes in **Fig 5D**) and thus we interpret them cautiously. A more robust

observation was that female-to-male *chasing-to-huddling* was more likely in matched dyads (**Fig**

5D). Thus, female chasing in matched dyads was possibly more affiliative, or huddling-seeking

(**S2 Movie**). This suggests the existence of different states underlying the same behaviors

depending on dyad type, echoing an interpretation we proposed in the previous subsection.

Therefore, aside from the specificities of each behavior pair, **Fig 5** shows that neurotype

matching influences actions and reactions on the scale of seconds, reinforcing the time series

and correlation analyses reported earlier.

Discussion

We utilized algorithms for annotation and decomposition of social behaviors to examine

neurotype matching in rodents. Opposite-sex prairie voles were assigned to either matched or

mixed-neurotype dyads, resembling human studies involving ASD. The voles were then filmed

in two experiments: initial divided and subsequent undivided cohabitation recordings. Dyad type

effects emerged from both experiments in a correlated manner, suggesting robust underlying

mechanisms. Our observations varied by sex and spanned multiple temporal scales, including

body orientation patterns over days, incidence of aggression over hours, and behavioral

transitions over seconds. These findings might influence further development of behavioral

paradigms and studies integrating social behavior and physiology.

Juvenile sleep and atypical sociability

Neurotype matching studies in humans often involve mixing and matching neurotypical

and neurodivergent individuals, especially those with ASD [2–6,13,15,17–19]. Here,

"neurodivergent" corresponds to adult prairie voles exposed to ELSD during post-natal days 14-

21, a method described previously by our group [30,33]. ELSD causes fragmentation of non-

rapid eye movement sleep (NREM) and reduces both duration and gamma power of rapid eye

movement sleep (REM), as recorded electrographically from prairie vole pups. Body weight,

corticosterone levels, and parental care are all unaffected by ELSD, indicating that this method

selectively targets juvenile sleep physiology [30,33]. Despite this selectivity, ELSD provokes

long-lasting behavioral and neurobiological alterations into adulthood. Behavioral alterations

include impaired cued fear extinction and reduced male-to-female partner preference

expression. Neurobiological alterations include cytoarchitectural markers of excitatory-inhibitory

imbalance in brain regions involved in sensory and cognitive processing, such as increased

expression of inhibitory neurons in somatosensory cortex, as well as increased dendritic spine

density and decreased presynaptic glutamate labeling in prefrontal cortex [30,34]. Similar

neurobiological changes are present in human ASD, making prairie vole ELSD relevant for

investigating ASD and other disorders linked to poor juvenile sleep [35].

As reviewed previously [35–37], sleep disruption during infancy has been associated

with ASD in humans. Brain maturation processes in humans, such as axonal myelination and

synaptic pruning, depend on healthy sleep during development. Disrupted sleep may interfere

with these processes and increase vulnerability to ASD-like symptoms, including social

difficulties [35–37]. Since neurotype matching modulates behavior in both prairie vole ELSD

(present study) and human ASD [2–6,13,15,17–19], we can speculate that some of these social

properties may trace back to sleep quality during development. This is, of course, a more

complex question. First, the relationship between sleep disruption and ASD is recognized as a

bidirectional and multifaceted comorbidity, with sleep disturbances both amplifying and being

amplified by ASD [36,37]. Second, human ASD is linked with other factors not directly related to

sleep, such as heritability and perinatal exposure to toxicants [38]. And third, juvenile sleep is

sensitive to circumstances not directly related to ASD, such as respiratory problems and

household conditions [39,40]. All these factors can lead to (and interact with) ASD [41], making

it difficult to isolate the role of juvenile sleep in ASD, let alone dyad type properties. Still, our

results suggest that juvenile sleep does play a role in these social properties – a role that we

found to be discernible in non-human animals, despite the elusive mechanisms. Future studies

could disentangle poor juvenile sleep from other environmental and genetic factors using

different rodent models, furthering the study of non-human neurotype matching and

complementing human ASD research.

Markers of social affinity across neuropsychiatric spectra

Human ASD encompasses a myriad of features, including subtle presentations of ASD,

which can be challenging to detect and often go undiagnosed [42,43]. Likewise, ELSD prairie

voles do not show overt behavioral changes. In our experience, these changes can only be

detected with quantification, whether using manual [30] or computerized [31] tracking. Thus,

ELSD prairie voles might represent a nuanced ASD-like spectrum, analogous to nuanced

neuropsychiatric spectra in humans. Our experiments add complexity to this question. Despite

the elusive nature of these spectra, their phenotypes can still supply us with controllable

"reagents" to manipulate "social chemistry". The "purity" of these reagents (i.e., phenotypes)

certainly vary among species or between laboratory and naturalistic settings, yet they still seem

to bring about some form of chemistry, as suggested by the literature on human dyads and our

study in prairie voles. Furthermore, dyadic behavior in humans has been investigated beyond

ASD, including studies on dysphoria, attention deficit disorder, and decision making [1,7–9], all

of which reinforcing that mixing or matching different phenotypes (or social styles) can modulate

behavioral performance and sociability. This indicates that social matching properties are

measurable across species and phenotypes, with implications for basic science, mental health,

and a better understanding of reciprocity and empathy [21–23].

These questions warrant further investigation using data-intensive metrics of social

affinity. Recent human ASD studies involving matched and mixed dyads used objective metrics

from video, audio and electroencephalography, including motion energy, mutual gaze,

backchanneling, verbosity, smile synchrony, and inter-brain synchrony [12,13,15,17,18].

Generally, these studies also support neurotype matching properties, though some metrics are

inconsistent with this trend. Such inconsistency might be attributed to low temporal resolution,

as these studies used aggregated metrics across sessions when comparing dyad types. Time

series analyses, like those here, could be employed in future studies to extract detailed

information from each metric, potentially identifying optimal metrics for real-time sociability

tracking and translatability across species. Other human dyad studies, unrelated to ASD or

neurotype matching, used additional metrics, including task cooperation, motor coordination,

smell-induced arousal, and functional brain imaging or spectroscopy [14,20,44–49]. This variety

of methods suggests promising avenues for studying dyadic behavior across species and

neuropsychiatric disorders.

Sex differences and ethological nuance

In our male-female grouping design (Ctrl-Ctrl, ELSD-ELSD, Ctrl-ELSD, ELSD-Ctrl) we

considered both sex and dyad type. This revealed dyad type effects both between and within

sexes in prairie voles. Between-sex and within-sex interactions have also been recognized in

human ASD. Core ASD traits seem to vary equivalently between sexes (e.g., communication

impairments, repetitive behaviors), whereas associated ASD traits seem to vary independently

within each sex (e.g., hyperactivity in males, depression in females) [50–52]. To our knowledge,

these sex-related variables have not been experimentally manipulated in studies on neurotype

matching and ASD. As shown here with prairie voles, sex differences in body direction (males

spent more time toward the divider) and aggression (females chased and attacked more) were

exacerbated in mixed dyads. We can now speculate that neurotype matching could also amplify

sex differences in humans and other species. Investigating these questions could provide

deeper insight into sex differences in human ASD or ASD-like models.

Moreover, we found that female body orientation in divided cohabitation (Experiment 1)

led to different outcomes in undivided cohabitation (Experiment 2) depending on the dyad type:

increased affiliation in matched dyads (less attacking, more huddling) and decreased affiliation

in mixed dyads (more chasing, less chasing-to-huddling transitions). Therefore, a female

behavior that looked the same – body direction – may have been driven by different underlying

states depending on the dyad type, e.g., female body direction was possibly more huddling-

oriented in matched dyads. Interestingly, these states in females may have influenced (and

been influenced by) the males. In mixed dyads, males spent more time toward the divider in

Experiment 1, and once the divider was removed in Experiment 2 the females attacked more

frequently. Thus, females may have reacted proportionally to the males' approaches upon

removal of the divider, which could be another explanation why aggression was more frequent

in mixed dyads. Aggressivity, affiliation and other behavior types (e.g., courting, foraging) are

indeed recognized to vary with sex, social hierarchy and environmental context [53–55],

suggesting an interplay between internal and expressed states. We propose that matching and

mixing rodent "neurotypes" like we did here is a promising approach to investigate these states.

Concluding remarks

This study reveals that neurotype matching is a quantifiable phenomenon in rodents. As

computational ethology continues to optimize social behavior tracking, the emergent properties

of social groups may become more tangible. This progress may reshape behavioral paradigms,

from dyad-based studies to freely moving partner preference studies, strengthening the link

between animal models and human sociability.

Materials and methods

Subjects

Prairie voles (n = 28 each sex) were sourced from a genetically diverse pool of 13

breeders in an accredited, fully-contained laboratory facility (VA Portland Health Care System).

Litters containing both male and female sibling pups (2-9 pups each breeder pair) were raised

with their parents (no cross fostering) until post-natal day 21 (P21). During the P14-21 period,

specifically, the litters were subjected to either control (Ctrl) or early-life sleep disruption (ELSD)

conditions (details below). At P21, pups were weaned into same-sex sibling groups and

remained at the same vivarium until approximately P90. The sibling groups were then shipped

to the University of Michigan Medical School for the primary recordings of this study.

Consistent housing conditions were maintained across both institutions, including 14:10

h light/dark cycle (lights on at 5:00 am), temperature (20-23 °C), ventilation (double-filtered

outside air, negative pressure), humidity (30-70%), bedding (rolled paper pellets), environmental

enrichment (cotton nestlets and wooden blocks/sticks) and *ad libitum* water and food (rabbit

chow, corn, cracked oats). Each week, the animals were transferred to a new cage with fresh

bedding and enrichment. All procedures in this study were approved by the Institutional Animal

Care and Use Committees (IACUC) of VA Portland Health Care System (protocols 6069 and

3652) and University of Michigan Medical School (protocol 00011423).

Our animals originated from a colony at Emory University, itself derived from wild-

captured animals in the State of Illinois, USA. To preserve genetic diversity, bi-annual

exchanges of animals occur among several institutions, including North Carolina State

University, University of California Davis, University of Colorado Boulder, and Florida State

University.

Early-life sleep disruption

Early-life sleep disruption

Prairie vole ELSD phenotypes of both sexes were generated by placing home cages

with P14-21 pups and their parents on an orbital shaker for intermittent gentle agitation (10

seconds on, 100 seconds off, 110 rotations per minute). ELSD animals were provided hydrogel

instead of water bottles to prevent water spillage during agitation. Ctrl animals experienced the

same conditions except for agitation (**Fig 1A**). As previously described by our group [30,33,34],

ELSD disturbs sleep in prairie vole pups without affecting their hormonal markers of stress, or

the parental care they receive. The acute effects of ELSD are, therefore, confined to sleep

electroencephalography (EEG) parameters, including fragmentation of NREM sleep and

reduction of REM sleep time by more than 20% over a 24-hour period [30]. Given this support

from our previous studies, here we used identical cage agitation parameters without EEG

validation, resulting in a non-invasive (purely behavioral) experimentation timeline.

Dyad assignment and age at recording

Following the ELSD or Ctrl procedures at P21, the pups were weaned into same-sex

groups of 2-4 siblings per cage until adulthood, resulting in four types of sibling groups (Male-

Ctrl, Male-ELSD, Female-Ctrl, Female-ELSD). Sexually naïve opposite-sex adults were then

randomly assigned into matched dyads [Male-Ctrl/Female-Ctrl (n=7), Male-ELSD/Female-ELSD

(n=8)] or mixed dyads [Male-Ctrl/Female-ELSD (n=7), Male-ELSD/Female-Ctrl (n=6)] prior to

recordings, for a total of 28 dyads (**Fig 1A**).

Our recording setup could accommodate up to four dyads at a time (details below).

Since there were 28 dyads in total, the recordings were divided into seven cohorts of four

dyads. Furthermore, an interval of 2-3 weeks was necessary between cohorts due to the

duration of the experiments combined with preparatory activities, like cage cleaning and file

management. This resulted in a four-month period to record all 28 dyads. Thus, the ages of the

animals at the time of recording varied with a mean +/- standard error of 185.8 +/- 6.5 days of

age. However, the individuals of each dyad were still age-matched, with an age difference of

11.1 +/- 2.2 days, randomly distributed across dyad types.

Recording setup

For recording, we developed a vibration-free optomechanical assembly using metal

parts from a vendor (ThorLabs), as described previously [31]. This setup holds two infrared-

sensitive grayscale cameras (Basler, acA1300-60gm) positioned at 90° overhead angle. Each

camera was mounted to a fixed focal length lens (Edmund Optics, 6 mm UC Series) and

surrounded by four infrared LED illuminators. Each camera/lens was able to frame two home

cages, hence the ability to record four dyads at a time. This system was installed in a room with

light/dark phases, which however did not affect imaging brightness, as it relied on infrared

reflectance. The cameras were connected to a computer via a network adapter (Intel Pro

1000/PT), and data were stored on a redundant array of independent disks (RAID). Recording

settings were configured via Pylon software (Basler) with the following parameters: 8-bit

grayscale depth, frame width x height of 800 x 896 pixels, no binning, 1328 kbps data rate, 20

Hz frame rate. Exposure and brightness were adjusted prior to each recording by manipulating

the lenses and infrared illuminators, without digital adjustments. Data were acquired into mp4

files (H.264 coded) using StreamPix software (NorPix). For the 72-h recordings of Experiment 1,

specifically, recordings were acquired into 6-h mp4 files to ensure file readability, including 2-s

margins at the beginning and end of each 6-h file.

Experiments 1 and 2

Experiment 1 consisted of divided cohabitation for 72 h starting at 8 am, i.e., 3 h after the

onset of the light phase. Each male-female pair was placed in a bedded home cage (48.3 cm

length, 25.4 cm width, 20.3 cm height) with a lab-made divider at the center of the cage. Thus,

each animal was spatially restricted to a quadrant, producing uniform behavioral variables

across recordings (**Fig 1B**). The dividers were made of metal wire mesh (square mesh, 6.5 mm

aperture, 22 cm wide, 28 cm high) covered on both sides with clear polycarbonate sheet (0.5

mm thick) to prevent animals from climbing. The bottom section of the metal mesh, measuring

6.5 cm in height, was left exposed without the polycarbonate sheet, allowing animals to

exchange bedding, smell each other and engage in face-to-face and body-to-body interactions.

Chow and gel were placed at consistent locations away from the divider, on the left and right

sides of the animal, respectively. No environmental enrichment objects were provided,

encouraging the animals to interact. A handwritten label was placed outside each quadrant

displaying animal metadata (sex, ELSD vs. Ctrl, quadrant number), clearly visible in each

recording for extra documentation. Cardboard barriers were placed between adjacent cages,

preventing different dyads from seeing each other during the multi-dyad recordings.

Experiment 2 consisted of undivided cohabitation for 4 h, also beginning at 8 am, 3-4

days after the end of Experiment 1; animals were single-housed at the vivarium between

experiments. Before each recording of Experiment 2, the male and female were placed in the

same home-cage quadrants, with the same handwritten label, same bedding and without chow

or gel. We then removed the divider *while recording*, ensuring animal traceability throughout the

recording, as male and female lacked physical differentiators (**Fig 1C**). See the next section for

further identity verification methods. The male and female were then allowed to interact freely.

At the end of the recording, each animal was visually inspected to confirm sex, resulting in a

100% match with the original metadata.

Data pre-processing and animal tracking

For Experiment 1, each camera recorded two divided cages and, therefore, four

quadrants, one animal per quadrant. Thus, we decided to crop each video into four smaller

videos (384 x 416 pixels), one per quadrant, using Adobe Premiere and Media Encoder. By

doing this we eliminated the need for identity verification in Experiment 1 (unlike Experiment 2,

as described below). The cropped videos were then processed through DeepLabCut, a

supervised keypoint tracker [28]. A resnet-50 network was used to label seven keypoints per

animal: nose, left/right ears, three locations along the "spine", and tail base (**Fig 1B**; **S1 Movie**).

The network was trained with video frames from three males and three females, each frame

representing varying levels of difficulty: from clear imaging of the animal (without obstruction of

body parts) to challenging situations (e.g., curled posture when sleeping or eating, tail base

hidden under bedding, nose hidden by the mesh divider). The network was trained using a

laboratory server (CPU: Intel Xeon E5-2640 v3 @ 2.60 GHz. RAM memory: 512 GB. GPU:

NVIDIA GeForce GTX 1080 Ti) and deployed to unseen (held-out) data. Once keypoint

coordinates were generated for each recording, we proceeded to behavioral decomposition

analysis using MATLAB (MathWorks), as detailed in the next section.

For Experiment 2, all video processing and behavior annotation steps were performed

using LabGym2, an open-source machine vison tool that we co-developed (including authors:

Y.H., B.Y., and L.S.B-J.) [32]. First, each raw video was cropped into two rectangular videos

(800 x 450 pixels), one per cage, so that each video contained one male-female dyad. The

cropped videos were then contrast-enhanced to highlight the animals against the background

(bedded cage). These videos were then analyzed using two neural networks that we trained

during our previous study: *Detector* and *Categorizer* [32]. Step 1: Segmentation and masking of

each animal using the *Detector*. Step 2: Mask-based foreground extraction to obtain the raw

pixels of each animal; background pixels were set to zero (black). Step 3: Unsupervised tracking

538 of masks over consecutive frames based on foreground appearance and distance. Step 4:

539 Generation of spatiotemporal patterns as inputs for the *Categorizer*, including body contour

540 outlines around the animals. Specifically for the focal animal, these outlines were color-coded

541 from blue to red to indicate the temporal progression of movements within groups of frames, i.e.,

542 time bins totaling 0.67 s per bin. The non-focal (or reference) animal was color coded in a fixed

543 tone of gray (**Fig 1C**). The focal and reference roles were alternated per animal by processing

544 each recording twice, enabling social role identification [32]. Time bin size (0.67 s) was

545 determined to match the timescale of prairie vole behavior, including quick bouts of chasing and

546 aggression. Step 5: Categorization of spatiotemporal patterns into behaviors (**Fig 1C**). This step

547 involved manual sorting of behavioral examples – the primary supervised aspect of LabGym2 –

548 followed by deep learning of the spatiotemporal features of these behaviors using a custom

549 neural network structure, as detailed in our software publication [32]. LabGym2 was run

550 remotely on the University of Michigan Great Lakes cluster. The sessions on this cluster were

551 configured with the following resources: 2.9 GHz Intel Xeon Gold 6226R with 372 GB RAM,

552 NVIDIA A40 with 48 GB VRAM.

553  LabGym2's publication included a proof-of-concept study for prairie vole dyads,

554 demonstrating 100% accurate individual animal tracking throughout each recording [32].

555 However, the recordings in that study were relatively short (10 min), whereas the recordings in

556 the present study were longer (4 h), which we found to increase susceptibility to identity

557 switching. Tracking identities over extended periods (e.g., several hours) indeed remains a

558 challenging task in the field of computational ethology, despite the development of various

559 approaches to multi-animal tracking [26–29]. In our experience, species-typical huddling

560 behaviors exacerbated the risk of identity switching, especially 2-3 hours into the recording,

561 when the voles tended to huddle more. Thus, we chose to identify male and female *manually*,

562 independently on LabGym2's behavioral categorization. Additionally, we leveraged this manual

563 inspection effort to annotate "attacking" and "being attacked" events (**S2 Movie**), which we could

not reliably automate in the current dataset. The annotation of "chasing" and "being chased", on

the other hand, was fully automated, serving as a control for our findings on sex-specific

aggression roles. For the manual annotation of animal identities and aggression roles we used

the graphical user interface (GUI) of another software tool: DeepEthogram [56].

In summary, we employed two distinct machine vision pipelines in this study, each

tailored to a recording environment (divided and undivided cohabitation). However, our study

also involved significant human supervision, including manual validation of sex identities and

aggression roles, mitigating the risk of over-automation [57]. In fact, our results on "chasing" and

"attacking" supported each other (**Fig 3**), which is an important control, as "chasing" and

"attacking" were annotated by computer and human, respectively. We believe that human-in-

the-loop strategies like these will continue to offer a balance between effort cost and ethological

validity, revealing meaningful patterns like the neurotype matching effects reported here.

Behavioral decomposition and categorization

<u>Experiment 1</u>: Keypoints from DeepLabCut represented each animal as an arrow-

shaped "skeleton", one per video frame (**Fig 1B**). The coordinates for these keypoints were

imported into MATLAB, converted from pixels to centimeters, and processed into three

behavioral variables.

1) Direction to divider: a line was fitted between the extreme keypoints (nose and tail

584   base), and the angle of this line relative to the horizontal plane of the frame was

585   calculated. This angle was then rescaled from -1 (opposite from the divider) to +1

586   (toward the divider), with clockwise and counterclockwise directions treated equally.

587   2) Distance to divider: horizontal distance between the center of the animal and the

588   divider, regardless of its position along the vertical axis.

3) Locomotion speed: difference (hypotenuse) between the center of the animal in

frame n and frame n+1.

These measures were timestamped according to clock time, in number of seconds x

number of frames. The clock was not restarted at midnight, assigning each frame to a unique

temporal identifier across the 72 hours. This allowed us to trim and align the recordings to a

common 72 h axis, as safety margins of a few minutes were recorded before and after each

session. Clock time information was obtained from file naming metadata through the acquisition

software (StreamPix, NorPix). These procedures replicate those used in a previous publication

[31]. See **S1 Movie**.

Experiment 2: LabGym2's outputs include the likelihood of each behavioral category per

video frame on a 0-1 scale. The categories were initially the same as our proof-of-concept study

in prairie voles [32]: aggression, chasing, being chased, approaching, sniffing, being sniffed,

moving alone, idling alone, huddling. For the present study, aggression was ignored (replaced

by manual annotations, as described below) and adaptations to other categories were made

using MATLAB.

1) Sniffing into Pointing: This category merged three types of sniffing (face-to-face,

face-to-body, anogenital) and approaching behaviors. This was renamed to a generic

term, "pointing", as freezing-like behaviors that typically occurred after aggressive

bouts (**S2 Movie**) were often misclassified as "sniffing" under LabGym2's

spatiotemporal patterning. Neural network refinement and deeper ethological

exploration were outside our scope (e.g., sub-classification of sniffing, sequencing of

peri-aggression postures).

2) Behaving alone: This category merged "moving alone" and "idling alone" (this latter

included self-grooming). In-depth exploration of these solitary behaviors was outside

our scope, hence the creation of this generic category. Frames were set to zero

likelihood of "behaving alone" if only one animal was classified under this category,

making it a full-dyad category (see purple curves in **Fig 3**).

3)   Huddling: Same as the previous proof-of-concept study in 10-min recordings [32], but

with one correction that we found necessary in the present 4-h recordings. During

intense body contact, the identities of both animals sometimes merged into one,

causing "huddling" to be misclassified as "idling alone". Frames with this issue we re-

classified as huddling. Frames were set to zero likelihood of "huddling" if only one

animal was classified under this category, making it a full-dyad category (see purple

curves in **Fig 3**).

After these adaptations, likelihood values were converted into a vector of integer labels

using a winner-takes-all approach. Finally, frames manually classified as "attacking" and "being

attacked" were superimposed on that vector of labels. For final quantification (explained next),

the focal roles of chasing, attacking, and pointing were analyzed without their mirroring

counterparts (i.e., being chased, being attacked, being pointed at), thus minimizing redundancy

in the figures.

Final data processing and statistical analysis per figure

Figs 2 and 3: Time series data obtained from the behavioral decomposition (Experiment

1) and behavioral annotation (Experiment 2) methods above were analyzed, with each data

point corresponding to one video frame of a single animal. These data points were binned and

smoothed on a per-animal basis: the 72-h recordings of Experiment 1 were averaged into 1-h

bins (total of 72 bins) and the 4-h recordings of Experiment 2 were averaged into 3-min bins

(total of 80 bins), both followed by a 6-bin moving mean window. The binned data were then

plotted as +/- standard error curves comparing males vs. females overall, males vs. females

within major dyad groups (matched or mixed), males vs. females within dyad sub-groups (Ctrl-

Ctrl, ELSD-ELSD, Ctrl-ELSD, ELSD-Ctrl), and matched vs. mixed dyads within each sex. These comparisons were performed comprehensively across behavioral variables, allowing to identify sex or dyad type differences with both: (1) behavioral variable specificity (in which behavioral variables the dyad type precipitated or amplified differences?); and (2) temporal specificity (at which time bins did these differences emerge?). Statistical comparisons were made using two-way ANOVA with time bins as repeated measures, followed by Tukey's post-hoc tests per time bin. The main statistical effects (sex, time, interaction) are reported in **Tables 1 and 2** and significant post-hoc differences are depicted as bold lines along the x-axes in **Figs 2 and 3**. Distributions per repeated measure passed the Kolmogorov-Smirnov normality test.

Fig 4: Experiments 1 and 2 were performed in different environments (divided and undivided cohabitation), 3-4 days apart. The animals and dyad assignments were the same between experiments, allowing for correlation analyses among all eight variables from the two experiments. To investigate this, behavioral variables were averaged across the entirety of each experiment, resulting in eight averages per animal. These averages were then grouped by sex and major dyad type (males or females from matched or mixed), forming four animal x variable arrays. Linear correlations within each sex and dyad type group were calculated, producing matrices with correlation coefficients (R) and corresponding P values. Key correlations were illustrated as scatter plots with linear fits in **Fig 4A-B**, while the R and P value matrices themselves were illustrated with a color code in **Fig 4C**. Distributions per variable passed the Kolmogorov-Smirnov normality test.

Fig 5A-B: Behavior transition probabilities were computed using 1-s time bins, matching the approximate timescale of prairie vole behaviors, including chasing and aggression bouts. For Experiment 1, continuous DeepLabCut-based variables of direction and distance to divider were averaged per time bin and discretized into categorical vectors (number of elements = number of time bins), enabling transition probability analysis. For discretization, direction to divider was converted from a -1 to 1 scale into "toward" vs. "opposite", whereas distance to

divider was converted from a centimeter scale into "near" vs. "away". These categories were

combined into a four-category vector (toward-near, opposite-near, toward-away, opposite-

away), one per animal. Then each dyad had a "time t" animal and a "time t+1" animal

assignment, one iteration per sex directionality (male-then-female, female-then-male). Each of

these iterations resulted in a 4 x 4 adjacency matrix (4 = number of behavioral categories), with

rows representing "time t" behaviors, columns representing "time t+1" behaviors, and values

indicating probabilities. Multiple such adjacency matrices were created, grouped by both sex

directionality and dyad type. Specifically for the illustrations in **Fig 5A**, adjacency matrices were

averaged across the dyads of each group and plotted as directed graphs. The edges of these

graphs were coded by both weight and grayscale: the thicker and darker the edge, the higher

the transition probability. For quantification (**Fig 5B**) no averaging was applied. Instead, each

adjacency matrix (one per dyad) was reshaped into a row vector containing all behavior pairs

("behavior in time t" and "behavior in time t+1"). These rows were then concatenated across

dyads, creating dyad x behavior pair matrices. Finally, dyad x behavior pair matrices from

matched and mixed dyads were statistically compared per sex directionality using two-way

ANOVA with behavior pairs as repeated measures, followed by Tukey's post-hoc tests per

behavior pair. The main statistical effects (dyad type, behavior pair, interaction) are reported in

Table 3 and significant post-hoc differences are depicted as bold edges in **Fig 5B**. Distributions

per repeated measure passed the Kolmogorov-Smirnov normality test.

Fig 5C-D: Similar methods were used for Experiment 2 with three differences: (1)

LabGym2-based data were already in categorical form prior to analysis, eliminating the need for

categorization; (2) each time bin (1 s) was assigned its most frequent category (winner-takes-

all); and (3) this analysis included five behavioral categories (chasing, attacking, pointing,

behaving alone, huddling), leading to 5 x 5 adjacency matrices.

Data and code availability

The behavior tracking software (LabGym) and the training dataset for prairie voles are both available on GitHub (https://github.com/umyelab/LabGym). Processed post-tracking data, as well as code for figure plotting and statistical analysis are available to peer reviewers via private link on figshare.

Supporting information

S1 Movie. Decomposing behavioral variables in the divided cage

S2 Movie. Examples of behaviors in the undivided cage

Acknowledgements

This research was supported by the National Institute of Mental Health to B.O.W., M.M.L., L.S.B-J., C.E.J-T. (R01MH131592) and N.E.P.M. (F31MH136684), and the National Institute of Neurological Disorders and Stroke to B.Y. (R01NS137222). We thank all lab colleagues for discussions.

Author contributions

Conceptualization: Lezio S. Bueno-Junior, Noah E. P. Milman, Carolyn E. Jones-Tinsley, Miranda M. Lim, Brendon O. Watson.

Data curation: Lezio S. Bueno-Junior, Brendon O. Watson.

Formal analysis: Lezio S. Bueno-Junior.

Funding acquisition: Brendon O. Watson, Miranda M. Lim, Lezio S. Bueno-Junior, Carolyn E.

Jones-Tinsley.

Investigation: Lezio S. Bueno-Junior.

Methodology: Lezio S. Bueno-Junior, Carolyn E. Jones-Tinsley, Noah E. P. Milman.

Project administration: Brendon O. Watson, Miranda M. Lim.

Software: Lezio S. Bueno-Junior, Yujia Hu, Bing Ye, Anjesh Ghimire.

Resources: Carolyn E. Jones-Tinsley, Peyton T. Wickham, Brendon O. Watson, Miranda M.

Lim.

Supervision: Brendon O. Watson, Miranda M. Lim.

Visualization: Lezio S. Bueno-Junior.

Writing – original draft: Lezio S. Bueno-Junior.

Writing – review & editing: Noah E. P. Milman, Brendon O. Watson, Miranda M. Lim, Carolyn

E. Jones-Tinsley, Anjesh Ghimire, Peyton T. Wickham, Yujia Hu, Bing Ye.

Declaration of interests

The authors declare no competing interests.

References

1.   Locke KD, Horowitz LM. Satisfaction in interpersonal interactions as a function of similarity

in level of dysphoria. J Pers Soc Psychol. 1990;58: 823–831. doi:10.1037/0022-

3514.58.5.823

2.   Bauminger N, Solomon M, Aviezer A, Heung K, Brown J, Rogers SJ. Friendship in high-

functioning children with Autism Spectrum Disorder: Mixed and non-mixed dyads. J Autism

Dev Disord. 2008;38: 1211–1229. doi:10.1007/s10803-007-0501-2

3.   Petrina N, Carter M, Stephenson J, Sweller N. Perceived friendship quality of children with

Autism Spectrum Disorder as compared to their peers in mixed and non-mixed dyads. J

Autism Dev Disord. 2016;46: 1334–1343. doi:10.1007/s10803-015-2673-5

4.   Crompton CJ, Sharp M, Axbey H, Fletcher-Watson S, Flynn EG, Ropar D. Neurotype-

matching, but not being autistic, influences self and observer ratings of interpersonal

rapport. Front Psychol. 2020;11: 586171. doi:10.3389/fpsyg.2020.586171

5.   Morrison KE, DeBrabander KM, Jones DR, Faso DJ, Ackerman RA, Sasson NJ. Outcomes

of real-world social interaction for autistic adults paired with autistic compared to typically

developing partners. Autism. 2020;24: 1067–1080. doi:10.1177/1362361319892701

6.   Bolis D, Lahnakoski JM, Seidel D, Tamm J, Schilbach L. Interpersonal similarity of autistic

traits predicts friendship quality. Soc Cogn Affect Neurosci. 2021;16: 222–231.

doi:10.1093/scan/nsaa147

7.   Cunningham CE, Siegel LS. Peer interactions of normal and attention-deficit-disordered

boys during free-play, cooperative task, and simulated classroom situations. J Abnorm

Child Psychol. 1987;15: 247–268. doi:10.1007/BF00916353

8.   Clark ML, Cheyne JA, Cunningham CE, Siegel LS. Dyadic peer interaction and task

orientation in attention-deficit-disordered children. J Abnorm Child Psychol. 1988;16: 1–15.

doi:10.1007/BF00910496

9.   Schuldt JP, Chabris CF, Woolley AW, Hackman JR. Confidence in dyadic decision making:

The role of individual differences. J Behav Decis Mak. 2017;30: 168–180.

doi:10.1002/bdm.1927

10. Crompton CJ, Ropar D, Evans-Williams CV, Flynn EG, Fletcher-Watson S. Autistic peer-

to-peer information transfer is highly effective. Autism. 2020;24: 1704–1712.

doi:10.1177/1362361320919286

11. Crompton CJ, Foster SJ, Wilks CEH, Dodd M, Efthimiou TN, Ropar D, et al. Information

transfer within and between autistic and non-autistic people. Nat Hum Behav. 2025 [cited

22 May 2025]. doi:10.1038/s41562-025-02163-z

12. Georgescu AL, Koeroglu S, Hamilton AFDC, Vogeley K, Falter-Wagner CM, Tschacher W.

Reduced nonverbal interpersonal synchrony in autism spectrum disorder independent of

partner diagnosis: a motion energy study. Mol Autism. 2020;11: 11. doi:10.1186/s13229-

019-0305-1

13. Rifai OM, Fletcher-Watson S, Jiménez-Sánchez L, Crompton CJ. Investigating markers of

rapport in autistic and nonautistic interactions. Autism Adulthood. 2022;4: 3–11.

doi:10.1089/aut.2021.0017

14. Estrugo Y, Bar Yehuda S, Bauminger-Zviely N. Pathways to peer interaction in ASD and

TD through individual and dyadic joint-action motor abilities. Front Psychol. 2023;14:

1234376. doi:10.3389/fpsyg.2023.1234376

15. Geelhand P, Papastamou F, Jaspard S, Kissine M. Autistic adults display different verbal

behavior only in mixed-neurotype interactions: Evidence from a referential communication

task. Autism. 2024; 13623613241298376. doi:10.1177/13623613241298376

16.   Koehler JC, Dong MS, Bierlich AM, Fischer S, Späth J, Plank IS, et al. Machine learning

classification of autism spectrum disorder based on reciprocity in naturalistic social

interactions. Transl Psychiatry. 2024;14: 76. doi:10.1038/s41398-024-02802-5

17.   McNaughton KA, Moss A, Yarger HA, Redcay E. Smiling synchronization predicts

interaction enjoyment in peer dyads of autistic and neurotypical youth. Autism. 2024;28:

2754–2767. doi:10.1177/13623613241238269

18.   Moreau Q, Brun F, Ayrolles A, Nadel J, Dumas G. Distinct social behavior and inter-brain

connectivity in Dyads with autistic individuals. Soc Neurosci. 2024;19: 124–136.

doi:10.1080/17470919.2024.2379917

19.   Oates M, Bean A, Kickbusch R, Sauer S. Extending double empathy: Effects of neurotype-

matching on communication success in an expository context. Am J Speech Lang Pathol.

2024;33: 2364–2377. doi:10.1044/2024_AJSLP-23-00393

20.   Zhao H, Zhang C, Tao R, Wang M, Yin Y, Xu S. Dyadic similarity in social value orientation

modulates hyper-brain network dynamics during interpersonal coordination: An fNIRS-

based hyperscanning study. Brain Topogr. 2025;38: 15. doi:10.1007/s10548-024-01092-w

21.   Milton DEM. On the ontological status of autism: the 'double empathy problem.' Disabil

Soc. 2012;27: 883–887. doi:10.1080/09687599.2012.710008

22.   Anderson C, Wood CI, Franklin L, Iampieri A, Sarsony C. "Getting Autism": Educators,

parents, and autistic adults and teens reflect on the importance of teachers who

understand. J Autism Dev Disord. 2024 [cited 2 Apr 2025]. doi:10.1007/s10803-024-

06353-y

23.   Rachanska PE. The Double Empathy Problem: An exploration of trans-neurotype attitudes

on autistic–non-autistic social interactions and communication. Adv Neurodev Disord. 2025

[cited 2 Apr 2025]. doi:10.1007/s41252-024-00435-9

24.   Nicolini C, Fahnestock M. The valproic acid-induced rodent model of autism. Exp Neurol.

2018;299: 217–227. doi:10.1016/j.expneurol.2017.04.017

25.   Delling JP, Boeckers TM. Comparison of SHANK3 deficiency in animal models:

phenotypes, treatment strategies, and translational implications. J Neurodevelop Disord.

2021;13: 55. doi:10.1186/s11689-021-09397-8

26.   Kabra M, Robie AA, Rivera-Alba M, Branson S, Branson K. JAABA: interactive machine

learning for automatic annotation of animal behavior. Nat Methods. 2013;10: 64–67.

doi:10.1038/nmeth.2281

27.   Romero-Ferrero F, Bergomi MG, Hinz RC, Heras FJH, De Polavieja GG. idtracker.ai:

tracking all individuals in small or large collectives of unmarked animals. Nat Methods.

2019;16: 179–182. doi:10.1038/s41592-018-0295-5

28.   Lauer J, Zhou M, Ye S, Menegas W, Schneider S, Nath T, et al. Multi-animal pose

estimation, identification and tracking with DeepLabCut. Nat Methods. 2022;19: 496–504.

doi:10.1038/s41592-022-01443-0

29.   Pereira TD, Tabris N, Matsliah A, Turner DM, Li J, Ravindranath S, et al. SLEAP: A deep

learning system for multi-animal pose tracking. Nat Methods. 2022;19: 486–495.

doi:10.1038/s41592-022-01426-1

30. Jones CE, Opel RA, Kaiser ME, Chau AQ, Quintana JR, Nipper MA, et al. Early-life sleep

disruption increases parvalbumin in primary somatosensory cortex and impairs social

bonding in prairie voles. Sci Adv. 2019;5: eaav5188. doi:10.1126/sciadv.aav5188

31. Bueno-Junior LS, Jones-Tinsley CE, Milman NEP, Wickham PT, Watson BO, Lim MM.

Early-life sleep disruption impairs subtle social behaviours in prairie voles: a pose-

estimation study. R Soc Open Sci. 2023;10: 230700. doi:10.1098/rsos.230700

32. Goss K, Bueno-Junior LS, Stangis K, Ardoin T, Carmon H, Zhou J, et al. Quantifying social

roles in multi-animal videos using subject-aware deep-learning. 2024.

doi:10.1101/2024.07.07.602350

33. Jones-Tinsley CE, Olson RJ, Mader M, Wickham PT, Gutowsky K, Wong C, et al. Early life

sleep disruption has long lasting, sex specific effects on later development of sleep in

prairie voles. Neurobiology of Sleep and Circadian Rhythms. 2023;14: 100087.

doi:10.1016/j.nbscr.2022.100087

34. Jones CE, Chau AQ, Olson RJ, Moore C, Wickham PT, Puranik N, et al. Early life sleep

disruption alters glutamate and dendritic spines in prefrontal cortex and impairs cognitive

flexibility in prairie voles. Curr Res Neurobiol. 2021;2: 100020.

doi:10.1016/j.crneur.2021.100020

35. Milman NEP, Tinsley CE, Raju RM, Lim MM. Loss of sleep when it is needed most –

Consequences of persistent developmental sleep disruption: A scoping review of rodent

models. Neurobiol Sleep Circadian Rhythms. 2023;14: 100085.

doi:10.1016/j.nbscr.2022.100085

36.   Bruni O, Breda M, Mammarella V, Mogavero MP, Ferri R. Sleep and circadian

846   disturbances in children with neurodevelopmental disorders. Nat Rev Neurol. 2025;21:

847   103–120. doi:10.1038/s41582-024-01052-9

848   37.   Vargas C, Paoletti D, De Stasio S, Berenguer C. Sleep disturbances in autistic children

849   and adolescents: A systematic review and meta-analysis of randomized controlled trials.

850   Autism. 2025;29: 1661–1673. doi:10.1177/13623613251319391

851   38.   Pugsley K, Scherer SW, Bellgrove MA, Hawi Z. Environmental exposures associated with

852   elevated risk for autism spectrum disorder may augment the burden of deleterious de novo

853   mutations among probands. Mol Psychiatry. 2022;27: 710–730. doi:10.1038/s41380-021-

854   01142-w

855   39.   Covington LB, Patterson F, Hale LE, Teti DM, Cordova A, Mayberry S, et al. The

856   contributory role of the family context in early childhood sleep health: A systematic review.

857   Sleep Health. 2021;7: 254–265. doi:10.1016/j.sleh.2020.11.010

858   40.   Reiter J, Ramagopal M, Gileles-Hillel A, Forno E. Sleep disorders in children with asthma.

859   Pediatr Pulmonol. 2022;57: 1851–1859. doi:10.1002/ppul.25264

860   41.   Miot S, Chancel R, Peries M, Crepiat S, Couderc S, Pernon E, et al. Multimorbidity

861   patterns and subgroups among autistic adults with intellectual disability: Results from the

862   EFAAR study. Autism. 2023;27: 762–777. doi:10.1177/13623613221121623

863   42.   Sharma SR, Gonda X, Tarazi FI. Autism Spectrum Disorder: Classification, diagnosis and

864   therapy. Pharmacol Ther. 2018;190: 91–104. doi:10.1016/j.pharmthera.2018.05.007

865   43.   Hirota T, King BH. Autism Spectrum Disorder: A Review. JAMA. 2023;329: 157.

866   doi:10.1001/jama.2022.23661

867   44.   Poncela-Casasnovas J, Gutiérrez-Roig M, Gracia-Lázaro C, Vicens J, Gómez-Gardeñes J,

868   Perelló J, et al. Humans display a reduced set of consistent behavioral phenotypes in

869   dyadic games. Sci Adv. 2016;2: e1600451. doi:10.1126/sciadv.1600451

870   45.   Zapata-Fonseca L, Dotov D, Fossion R, Froese T. Time-series analysis of embodied

871   interaction: Movement variability and complexity matching as dyadic properties. Front

872   Psychol. 2016;7. doi:10.3389/fpsyg.2016.01940

873   46.   Endevelt-Shapira Y, Perl O, Ravia A, Amir D, Eisen A, Bezalel V, et al. Altered responses

874   to social chemosignals in autism spectrum disorder. Nat Neurosci. 2018;21: 111–119.

875   doi:10.1038/s41593-017-0024-x

876   47.   Parkinson C, Kleinbaum AM, Wheatley T. Similar neural responses predict friendship. Nat

877   Commun. 2018;9: 332. doi:10.1038/s41467-017-02722-7

878   48.   Hyon R, Youm Y, Kim J, Chey J, Kwak S, Parkinson C. Similarity in functional brain

879   connectivity at rest predicts interpersonal closeness in the social network of an entire

880   village. Proc Natl Acad Sci U S A. 2020;117: 33149–33160. doi:10.1073/pnas.2013606117

881   49.   Ravreby I, Snitz K, Sobel N. There is chemistry in social chemistry. Sci Adv. 2022;8:

882   eabn0154. doi:10.1126/sciadv.abn0154

883   50.   Robinson EB, Lichtenstein P, Anckarsäter H, Happé F, Ronald A. Examining and

884   interpreting the female protective effect against autistic behavior. Proc Natl Acad Sci U S

885   A. 2013;110: 5258–5262. doi:10.1073/pnas.1211070110

886   51.   Werling DM, Geschwind DH. Sex differences in autism spectrum disorders. Curr Opin

887   Neurol. 2013;26: 146–153. doi:10.1097/WCO.0b013e32835ee548

52.   Hull L, Mandy W, Petrides K. Behavioural and cognitive sex/gender differences in autism

spectrum condition and typically developing males and females. Autism. 2017;21: 706–

727. doi:10.1177/1362361316669087

53.   Brusman LE, Protter DSW, Fultz AC, Paulson MU, Chapel GD, Elges IO, et al. Emergent

intra-pair sex differences and organized behavior in pair bonded prairie voles (Microtus

ochrogaster). Genes Brain Behav. 2022;21: e12786. doi:10.1111/gbb.12786

54.   Flavell SW, Gogolla N, Lovett-Barron M, Zelikowsky M. The emergence and influence of

internal states. Neuron. 2022;110: 2545–2570. doi:10.1016/j.neuron.2022.04.030

55.   Vahaba DM, Halstead ER, Donaldson ZR, Ahern TH, Beery AK. Sex differences in the

reward value of familiar mates in prairie voles. Genes Brain Behav. 2022;21: e12790.

doi:10.1111/gbb.12790

56.   Bohnslav JP, Wimalasena NK, Clausing KJ, Dai YY, Yarmolinsky DA, Cruz T, et al.

DeepEthogram, a machine learning pipeline for supervised behavior classification from raw

pixels. eLife. 2021;10: e63377. doi:10.7554/eLife.63377

57.   Kenkel W. Automated behavioral scoring: Do we even need humans? Ann N Y Acad Sci.

2023;1527: 25–29. doi:10.1111/nyas.15041

 Figure legends

Fig 1. Overview of animal pairing and behavioral analysis. A. Early-life sleep disruption

(ELSD) in prairie vole pups (post-natal days 14-21) generates phenotypes relevant to autism, as

indicated by behavioral and cytoarchitectural evidence [30,34]. ELSD subjects and their controls

were raised into adulthood in same-sex sibling groups under standard conditions. The adults

were then assigned into matched and mixed opposite-sex dyads. **B.** Experiment 1: Male and

female were filmed for 72 h while they interacted through a wire-mesh divider at the center of

the cage. Cameras were positioned at an overhead angle and the animals were tracked using

DeepLabCut [28], yielding cage-referenced behavioral variables. A similar method was utilized

in a previous study [31] but without directly investigating dyad type. **C.** Experiment 2: After 3-4

days of single housing in the vivarium, the same dyads were placed back together into the

recording setup, this time for 4 h and without the divider. This resulted in a different kind of

dataset comprising partner-referenced social role categories, annotated using LabGym2 [32]

and manual validation (see Methods).

Fig 2. Experiment 1: Mixed dyads showed exacerbated sex differences in body

orientation. A. Time series data (72 h, 1-h bins) showing sex differences in direction to divider

across all dyads. Males spent more time toward the divider than females throughout the

recordings. The curves represent mean +/- standard error aggregated across animals,

regardless of individual dyad assignments. The black bars on the x-axes represent posthoc

differences, after two-way ANOVA with time bins as repeated measures. **B.** Sex differences in

direction to divider were more pronounced in mixed dyads (highlighted with gray background).

Distance to divider and locomotion speed were less sensitive to dyad type, illustrating that

behavioral decomposition was important to capture the effects shown here. **C.** Further

specification into the four dyad subtypes. Sex differences in direction to divider were confined to

the mixed-dyad subtypes (Ctrl-ELSD, ELSD-Ctrl). Additional smaller effects were observed in

distance to divider and locomotion speed, specific to each mixed-dyad subtype. **D.** Same

groupings as panel B but comparing matched and mixed dyads within each sex. The tendency

to behave toward the divider was higher in males and further increased in mixed-dyad males.

See **Table 1** for statistics and **S1 Movie**.

Fig 3. Experiment 2: Mixed dyads showed higher female-to-male aggression. A. Time

series data (4 h, 3-min bins) showing sex differences in the likelihood of chasing and attacking

across all dyads. Females were more likely to chase and attack than males, especially in the

initial hour of recording. The curves represent mean +/- standard error aggregated across

animals, regardless of individual dyad assignments. The black bars on the x-axes represent

posthoc differences, after two-way ANOVA with time bins as repeated measures. Inset plots

show probability within the first hour. **B.** Sex differences in the likelihood of chasing and

attacking were more pronounced in mixed dyads (highlighted by gray background). Pointing

behaviors, including face-to-face, face-to-body and anogenital sniffing, showed no sex or dyad

type specificity, making them useful comparators (see Methods for details). The role-unspecific

categories of alone and huddling were more dominant and mutually exclusive, illustrating the

transition from high activity (periods of moving and idling alone punctuated by chasing,

aggression and sniffing bouts) to low activity (periods of continuous huddling). **C.** Further

specification into the four dyad subtypes. Sex differences in the likelihood of chasing and

attacking were stronger in the mixed-dyad subtypes (Ctrl-ELSD, ELSD-Ctrl). **D.** Same groupings

as panel B but comparing matched and mixed dyads within each sex. Mixed-dyad females were

more likely to chase and attack than matched-dyad females. See **Table 2** for statistics and **S2**

Movie.

Fig 4. Linking divided cohabitation to undivided cohabitation: Neurotype matching

affected correlations among behaviors. A. Correlations between the most sensitive variable

from Experiment 1 (72-h divided cohabitation), direction to divider (x-axes), and behavior

probabilities from Experiment 2 (4-h undivided cohabitation; y-axes), averaged per male

individual (data points). **B.** Same but for females. Significant correlations revealed further

distinctions between dyad types, highlighted by gray backgrounds. Matched-dyad females

formed an affiliation-driven spectrum: a higher tendency to behave toward the divider in

Experiment 1 correlated with higher sociability 3-4 days later, in Experiment 2 (lower likelihood

to attack and behave alone, higher likelihood to huddle). This pattern was absent in mixed-dyad

females; instead, they exhibited a positive correlation between direction to divider and chasing

behavior. **C.** Comprehensive view of all correlation pairs. R and P values are coded by color and

square size, respectively. Note that the scatter plots in panels A-B correspond to the leftmost

column of each correlation matrix.

Fig 5. Female-to-male behavior transitions differed between matched and mixed dyads.

A. Data from Experiment 1 (divided cohabitation). Direction and distance to divider were time-

binned (1-s), categorized (toward-near, opposite-near, toward-away, opposite-away), and

analyzed as male-to-female and female-to-male transition probability matrices (see Methods).

Directed graphs (averaged across dyads) illustrate patterns that varied with both dyad type and

sex directionality. **B.** Quantitative comparisons between dyad types within each sex

directionality, using one transition matrix per dyad, rather than the averages in panel A.

Behavior pairs were used as repeated measures in this analysis (see Methods). Self-loops were

excluded to focus on inter-behavior relationships. The directed graphs highlight which

transitions differed statistically between dyad types. The scatter plots exemplify the distributions

underlying each transition edge (each data point is a dyad). Mean +/- standard deviation bars

accompany the scatter plots. **C-D.** Same as panels A-B but from Experiment 2 (undivided

cohabitation). Overall, these patterns suggest that matched dyads showed more variation in

body re-orientation behaviors between the females and males, as well as a higher likelihood of

female chasing leading to huddling, among other interpretations. See Results and Discussion.

See **Table 3** for statistics.

Fig 1

a early-life sleep disruption (ELSD)

P14-P21 · prairie voles · undisturbed juvenile sleep

P14-P21 · disturbed juvenile sleep · **shaker**

raised into adulthood under standard conditions
(same-sex sibling cages)

♂ Ctrl ♀ Ctrl   ♂ ELSD ♀ ELSD

sexually naïve → **matched dyads** Ctrl-Ctrl, ELSD-ELSD   **mixed dyads** Ctrl-ELSD, ELSD-Ctrl

b Experiment 1: **DeepLabCut**-based behavioral decomposition

10 cm

← wire-mesh divider

toward, near | toward, near

toward, near | opposite, away

variables: direction to divider, distance to divider, locomotion speed

c Experiment 2 *(same dyads, 3-4 days later)*: **LabGym2**-based social role annotation

male-female identified | chasing | attacking | pointing | behaving alone | huddling

divider removed

identities manually verified throughout dataset

0   0.33   0.67   time from behavior onset (s)   ▪ reference animal

Fig 2

Fig 3

Fig 4

a

x-axes: <u>Experiment 1</u>, y-axes: <u>Experiment 2</u>

males *from* <u>*matched*</u> *dyads (n = 15)*

males *from* <u>*mixed*</u> *dyads (n = 13)*

b

females *from* <u>*matched*</u> *dyads (n = 15)*

females *from* <u>*mixed*</u> *dyads (n = 13)*

c

Fig 5