Title

Central carbon metabolism switching in lytic versus temperate coral reef viral communities

Authors

Jacob Kelman[1,2*], Meena Khan[1,2*], Chibundu Umunna[1,2*], Russell Brainard[3], Grace Donohue[1,2], Rob Edwards[4], Natalie A Falta[1,2], Emma George[5,6], Eleanor Gorham[1,2], Juris Grasis[7], Kevin Green[5], Andreas Haas[5,8], Kimberly Halsey[9], Eric Hester[5,10], Summer Jacob[1], Aydin Loid Karatas[1,2,11], Yan Wei Lim[5], Mark Little[5,12], Stuart Sandin[13], Jessie Segnitz[1], Maya Serota[1], Natalia Shahwan[1], Giselle Simmons[1], Jennifer E Smith[13], Isha Tripathi[1,2], Linda Wegley Kelly[13], Lauren Woodward[1], Nickie Yang[1,2], Charles Young[14], Brian Zgliczynski[13], Forest Rohwer[5], Ben Knowles[1,2,15,16†]

* These authors contributed equally to this work.

Affiliations

[1] Department of Ecology and Evolutionary Biology, University of California, Los Angeles, California 90095, USA

[2] Institute for Quantitative and Computational Biosciences, University of California, Los Angeles, California 90095, USA

[3] R Ocean LLC, Kāne'ohe, Hawaii 96744, USA

[4] Flinders Accelerator for Microbiome Exploration, Flinders University, Bedford Park, South Australia 5042, Australia

[5] Biology Department, San Diego State University, San Diego, California 92182, USA

[6] Integrative Oceanography Division, Scripps Institution of Oceanography, University of California, San Diego, La Jolla, California 92093, USA

[7] School of Natural Sciences, University of California, Merced, Merced, California 95343, USA

[8] Department of Marine Microbiology and Biogeochemistry, NIOZ Royal Netherlands Institute for Sea Research, 1797 SZ 't Horntje (Texel), Netherlands

[9] Department of Microbiology, Oregon State University, Corvallis, Oregon 97331, USA

[10] NIZO Food Research B.V., Kernhemseweg 2, 6718 ZB Ede, Netherlands

[11] Department of Quantitative and Computational Biology, University of Southern California, Los Angeles, California, USA

[12] Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, Massachusetts 02138, USA

[13] Marine Biology Research Division, Scripps Institution of Oceanography,
University of California, San Diego, La Jolla, California 92093, USA
[14] Pacific Islands Ocean Observing System, University of Hawaii at Manoa,
Honolulu, Hawaii 96822, USA
[15] California NanoSystems Institute, University of California, Los Angeles, California
90095, USA
[16] Institute of the Environment and Sustainability, University of California, Los
Angeles, California 90095, USA

Correspondence
[†] Ben Knowles: bwknowles@ucla.edu

Main text length (Introduction, Methods, Results, and Discussion): 4743 words

List of Tables and Figures

Table 1: Central carbon metabolism genes shown in Figures 2, 3, and 4.

Figure 1: Correlations between metabolic function and community temperateness across the breadth of cellular metabolism (SEED Level 1).

Figure 2: Correlation between central carbon metabolism genes and viral community temperateness.

Figure 3: Pathway map of changes in the frequency of central carbon metabolism genes across the lytic to temperate spectrum.

Figure 4: Correlations between viral community temperateness and central carbon gene frequency across pathways and gene function.

Figure 5: Schematic model incorporating viral metabolic effects on resilience of healthy coral reefs and exacerbation of reef degradation.

List of Supplementary Tables and Figures

Supplementary Table 1: Summary of samples analyzed, including sample accession numbers, the total number of base pairs, reads, and locations of viral metagenomes.

Supplementary Table 2: Parameters and values used in the qualitative ecosystem model.

Supplementary Table 3: The frequency of photosynthesis genes across the lytic-temperate spectrum.

Supplementary Figure 1: Temperate viral communities are more cell-like, and have higher gene diversity and evenness, but not richness.

Supplementary Figure 2: Bootstrapped distribution of ρ_s from 1,000 iterations correlating the frequency of central carbon metabolism genes with viral community temperateness.

Significance (114 words)

Coral reefs are collapsing worldwide due to "microbialization," where algae-fueled bacteria overgrow corals. Viruses that infect these bacteria can suppress this process through lysis, but on degraded reefs they often switch to nonlethal temperate lifestyles, accelerating decline. Here we show that similar shifts occur in virus-encoded metabolism. On healthy reefs, lytic viruses carry genes that drain host metabolites to fuel virus production, which likely enhances infection and lysis rates and limits bacterial overgrowth. On degraded reefs, temperate viruses encode reactions that expand host metabolite pools, supporting bacterial proliferation. Thus, viral metabolism can either reinforce reef resilience or exacerbate collapse, making it a hidden driver of ecosystem fate and a potential target for conservation strategies.

Abstract (235 words)

Coral reefs are declining globally due in part to bacterial overgrowth, a process known as microbialization. However, the role of bacteriophages that may inhibit microbialization by infecting and killing these bacteria remains poorly understood, especially their metabolic impacts on bacterial proliferation. To address this, we analyzed central carbon metabolism gene frequencies in viral communities from healthy (lytic-dominated) and degraded (temperate-dominated) Central Pacific coral reefs. We found that viral metabolism shifted broadly from being dominated by metabolism that builds up pools of central intermediates on degraded reefs dominated by temperate viral infection ('anaplerotic' reactions) to metabolism that consumes these pools to prioritize production of metabolic precursors for virion construction on healthy reefs dominated by lytic infection ('cataplerotic' reactions). This switch was shown by the over-representation of Entner-Doudoroff (ED) glycolysis genes on degraded, temperate-dominated reefs and of pentose phosphate pathway (PPP) and reductive tricarboxylic acid cycle (TCA) genes on healthy, lytic-dominated reefs. As a result of this metabolic dichotomy, our qualitative compartment modeling revealed two distinct ecosystem states: (i) healthy reefs, where lytic viral metabolism enhances viral production and suppresses bacterial overgrowth, and (ii) degraded reefs, where temperate viral metabolism accelerates bacterial proliferation. Because viral switching between lytic and temperate lifestyles is a known function of host physiological state, these findings position viral metabolism as both a driver of reef decline and a conservation lever, with metabolically mediated 're-viralization' offering a novel strategy to restore reef resilience.

Introduction (704 words)

Ecosystem overgrowth by microbes, a process known as microbialization, is a major threat to coral reefs worldwide[1–3]. Microbialized reefs are subject to a positive feedback loop in which proliferating algae generate excess dissolved organic carbon (i.e., sugar) that fuels the growth of increasingly pathogenic and copiotrophic microbes that thrive in nutrient-rich environments[1,4,5]. This process exacerbates coral diseases and clears space for further algal growth[2,6–8].

Bacterial metabolism on degraded coral reefs is directed towards high-throughput rather than efficient carbohydrate utilization[1,9]. For instance, while bacterial metabolism on healthy reefs is dominated by genes for the efficient but multi-step Embden-Meyerhof-Parnas (EMP) pathway, degraded reefs exhibit greater representation of genes for the inefficient but short Entner-Doudoroff (ED) glycolysis pathway [1,10–12]. This shift on degraded reefs enable bacteria to readily exploit ecosystem organic carbon enrichment to build up cellular stockpiles of central metabolites like glyceraldehyde-3-phopshate (G3P) and pyruvate ('anaplerotic' reactions) as well as use those central metabolite pools to produce cellular energy such as ATP, reductants like NADH, and amino and nucleic acid biosynthetic precursors required for bacterial over-proliferation ('cataplerotic' reactions)[1,7,11].

Viral predation could, in principle, preclude microbialization by killing proliferating microbes, as lytic infection and lysis rates are predicted by established predator-prey models to rise with host densities and growth rates[13–16]. However, viral communities switch from lytic to temperate infection during eutrophication and microbialization[14]. This transition towards non-lethal, temperate infection can facilitate further increases in microbial biomass due to suppressed lysis[14,17]. Further, temperate viral communities encode higher frequencies of pathogenicity genes that could protect bacteria from consumption by eukaryotic grazers[14,18–20]. These genes could facilitate

microbialization by limiting protist predation, the major non-viral predation guild

in the ocean[21,22]. Altogether, rather than controlling bacterial populations,

viruses may form a second 'Temperate Ratchet' feedback loop that traps

ecosystems in persistent microbialization through suppressed lysis rates and

protist grazing on pathogenic copiotrophs[14,23].

In addition to these population-level effects, viruses may also exert gene-level

effects on microbialization by reshaping bacterial metabolism. Viruses shuttle a

broad array of metabolic genes between hosts via horizontal gene transfer,

thereby altering bacterial metabolism and ecology[24–27]. These include central

carbon metabolism genes implicated in the metabolic switching of bacterial

communities on microbialized reefs, such as the pentose phosphate pathway

(PPP) and the ED and EMP glycolysis pathways[1,28]. In lytic viruses, these pathways

support virion synthesis by converting general cellular metabolic pools like G3P

and pyruvate to specialized metabolic precursors (i.e., they are cataplerotic

reactions). For example, the PPP provides nucleic acid and amino acid

precursors, whereas EMP generates cellular energy like ATP and NADH[27,29,30].

However, these genes can also be co-opted by bacteria during infection. As a

result, while viral metabolic genes can combat microbialization by promoting

the production of lytic viruses that would suppress bacterial abundance, they

are also capable of exacerbating reef decline by enhancing bacterial

proliferation. Together, these processes introduce a novel viral-mediated

metabolic mechanism of reef decline or resilience beyond lytic-to-lysogenic

switching.

However, the role of viral-encoded metabolic genes in coral reef

microbialization remains unexplored. To address this gap, we analyzed the

frequencies of metabolic genes in 19 metagenomes of virus-like particles (VLPs),

concentrates containing primarily viruses but also including virus-like gene

transfer agents[31,32], from Pacific coral reefs spanning a gradient from healthy to

degraded states and from lytic to temperate infection dominance. We found

that temperate viral communities were enriched in anaplerotic central carbon

metabolism genes, particularly those associated with copiotrophic growth like

Entner-Doudoroff glycolysis and the generation of central metabolite pools that

can be used for diverse cellular processes[33]. Lytic communities had higher levels

of cataplerotic non-oxidative pentose phosphate pathway genes and

oxidoreductases that generate specialized metabolic precursors, cellular energy

and redox potential. Incorporating our findings into a conceptual model

revealed that lytic viral metabolism can enhance viral production and microbial

surveillance on healthy reefs ("viralization")[34], while temperate viral metabolism

may accelerate microbialization in degraded systems. These findings implicate

viral metabolism in feedback loops that either enhance reef resilience or

exacerbate decline. Because viral lifestyle switching is a known function of

cellular physiological state, with elevated nutrient availability driving non-lethal

temperate infection, while nutrient limitation promotes lytic infection, our work

suggests that metabolic management, such as limiting nutrient inputs, could

promote rapid recovery of degraded reefs.

Methods (1764 words)

Metagenome sampling and processing: Viral metagenomes were sampled from coral reefs across the central Pacific Ocean (n = 19; see **Supplementary Table 1** for sample details) and samples processed as described previously[14,35]. Note that viruses and gene transfer agents (GTAs) likely co-purify using this technique, making our analysis primarily focused on viruses but also encompassing GTAs, that together comprise virus-like particles (VLPs)[31]. For each viral metagenome, approximately 60-100 L of seawater were pumped from the reef surface into acid-washed, triple-rinsed low-density polyethylene bags and concentrated to < 500 mL using a 100 kDa tangential flow filter. Concentrates were then 0.45 μm-filtered to remove bacteria and 0.5 % final concentration chloroform was added to destroy any residual cells and extracellular vesicles[36]. Samples were stored at 4 °C until subsequent processing. The presence of concentrated VLPs was confirmed by epifluorescence microscopy of Sybr Gold stained viruses on anodisc filters[35].

VLPs were purified from concentrates using a cesium chloride step gradient approach[14,35,37]. The absence of cells in step gradient outputs was confirmed by microscopy. Purified samples were then DNAse I-treated at 37 °C for 2 hours and denatured at 65 °C for 15 minutes to remove extra-virion free DNA (peDNA; protected environmental DNA)[32]. Viral DNA was extracted using the formamide/chloroform isoamyl alcohol technique[14,35,37]. The absence of contaminating host DNA was confirmed by the lack of PCR amplification of the universal 16S rDNA gene using primers 27F (AGA GTT TGA TCC TGG CTC AG) and 1492R (TAC GGY TAC CTT GTT ACG ACT T) on all samples[14,35,37]. Note that our analysis is based on correlations across many samples, minimizing the impact of any contamination on our findings. DNA was amplified using the Linker Amplified Sequencing Library approach[19] prior to sequencing on an Illumina MiSeq platform.

Bioinformatic processing: Low-quality reads < 100 bp in length and with mean

PHRED quality scores of < 25 were excluded using Prinseq[38]. Samples were then

dereplicated with TagCleaner[39], human contaminants were removed using

DeconSeq[40], and sequences were posted to MG-RAST (www.mg-rast.org; see

Data Availability section). Accession numbers and summary statistics for the viral

metagenomes used in this paper are shown in **Supplementary Table 1**. All the

viromes were published as part of the Piggyback-the-Winner project (n = 17)[14]

except Pearl and Hermes Reef and Caroline Atoll Wreck samples (n = 2; PHR

Wreck and CWS), which were unpublished prior to this work.

Annotating metagenomes: Viral metagenomes were annotated by BLAST

against the SEED database on MG-RAST using the default MG-RAST parameters

as a very commonly used balance of accuracy and specificity[41]: e-value = -5

(annotated in MG-RAST as e-value = 5), 60 % identity, > 15 base length, a

minimum abundance of 1, and all analyses conducted using representative

hits[14,42]. Level 1 Gene categories with average frequencies below 0.5 % were

omitted. Temperateness of each viral metagenome was assessed as the

frequency of genes annotated in the Level 3 SEED category of

"Phage_integration_and_excision" (i.e., how common genes involved in

integration and excision were in each metagenome)[14]. The frequency of genes

in high-order classification (e.g., SEED level 1 with categories shown in **Figure 1**)

was calculated by summing the frequency of the genes making up each

category. All analyses were normalized to account for different sequencing

depth between samples. This was done by calculating gene frequency as the

number of hits to a given gene divided by all known genes in that sample

("percent known")[15], making all analyses comparable and unaffected by the

effects of varied sequencing depth between samples.

Pathway annotation: Pathway visualizations were guided by KEGG maps

(www.genome.jp/), pathway-specific Wikipedia pages (www.wikipedia.org/),

and enzyme Wikigenes entries (www.wikigenes.org/). Pathway maps were

drawn using EC Number names of each enzyme (**Table 1**).

Anaplerotic and cataplerotic gene categorization: Genes were categorized

based on whether they (i) generated central metabolite pools like

glyeraldehyde-3-phopshate (G3P) and pyruvate ('anaplerotic' or 'filling up'

reactions) that are flexibly used in multiple pathways by cells, or (ii) used these

central metabolite pools to generate cellular energy or metabolic precursor

products that are used for specific purposes (i.e., the products are 'used'

metabolically; known together as 'cataplerotic' or 'drawing down' reactions),

or (iii) were neither anaplerotic nor cataplerotic ('neither'). Note that although

anaplerotic and cataplerotic reactions often refer to steps in the TCA, we are

using the term more generally here[33].

Statistical analyses: The correlation between gene frequency, either at a Level 1

SEED category or individual gene level, and viral community temperateness

were assessed using Spearman's Rho (ρ_s), a non-parametric rank-based

measure of how monotonic the relationship between x- and y-values is (i.e., it is

a non-parametric correlation coefficient). For supplementary diversity and

photosynthesis gene analyses we estimated ρ_s using the R cor.test() function

with exact = FALSE to avoid ties. For our main analyses shown in **Table 1** and in

Figures 1, **2**, **3**, and **4**, we estimated ρ_s values using the stat.spearmanr() function

in the Python scipy.stats package with bootstrapping to ensure robust median ρ_s

assessment over 1000 iterations of bootstrapping with replacement (see

Supplementary Figure 2 for resulting distributions for central carbon metabolism

genes). Confidence intervals of the median were assessed as the values

corresponding to the 2.5th and 97.5th percentiles of the median values

obtained from bootstrapping (95 % CI). For clarity, 95 % confidence intervals are

listed in square brackets, as ρ_s [95 % CI minimum, 95 % CI maximum].

Bootstrapping code can be found at https://tinyurl.com/viral-metabolic-ratchet-

code. Diversity metrics were assessed using the R vegan package, with the

diversity() command used for Shannon diversity (natural log-transformed; H').

Median ρ_s at the pathway level were calculated as the median value of all of

the ρ_s values for each enzyme in the pathway (i.e., they are a simple overall

median).

Qualitative model: We created a conceptual model to demonstrate qualitative

trends in bacterial, viral, coral, and algae abundance stemming from the

changes in viral metabolism found here (**Figure 5**). Note that this model is not

intended to be quantitative and was designed aid our understanding of trends

in organismal abundance in response to the incorporation of viral metabolic

capacity. Parameter values were implemented as high or low relative values to

reflect known processes on healthy and degraded reefs (**Supplementary Table 2**

for parameter values). For example, high rates of disease[8,43] and bacterial

metabolism[1] were used in degraded coral reef simulations, while high lysis rates

were used in healthy systems[14,17].

Bacterial, viral, coral, and algae populations were initialized at 0.5 arbitrary units.

The model was implemented using an iterative loop over six iterations (i.e., time

in arbitrary units). This number of iterations was chosen as a balance of having

enough iterations to see qualitative population trends, but without the need to

make a more complicated model to generate long-term dynamics. At the

beginning of each time step, new abundance values for each species (B_{new},

C_{new}, V_{new}) except algae were calculated based on the previous values (A_{new}).

These updated values were applied at the end of each time step. Algae

abundance was calculated last, occupying any remaining share of the benthos

not occupied by coral. Once a population reached or fell below zero, it was

excluded from further dynamics and not allowed to re-enter the ecosystem.

Code can be found at https://tinyurl.com/viral-metabolic-ratchet-code

For the 'no viruses' model shown in **Figures 5a** and **5d**, we used the following

equations, where terms with changed parameter values between **Figures 5a**

and **5d** are in bold (see **Supplementary Table 2**):

$B_{new} = B + a * A + \mathbf{m * B}$ Equation 1

$C_{new} = C + c * C - i * A - \mathbf{d * B}$ Equation 2

$A_{new} = 1 - C_{new}$ Equation 3

In **Equation 1**, bacteria (B) proliferate due to carbon provided by algae, with

carbon supply itself a product of algal population (A) as well as the rate of

delivery (a). To make the model as conservative as possible, we did not vary the

rate of carbon delivered to bacteria by the algae, although it likely is higher on

degraded reefs and enhances bacterial grow more than shown by the model[10].

Bacterial proliferation is also enhanced by metabolic changes (m) reinforced by

bacterial population size on degraded reefs[1]. Note that, in the absence of

viruses the bacteria have no loss processes, including natural mortality, even

though their proliferation may be constrained by a lack of algae to support

growth.

In **Equation 2**, coral populations (C) proliferate as a product of coral population

size and growth rate (c). Coral growth is limited by algal population size and

allelochemical inhibition (i)[44] as well as by bacterial pathogenicity causing coral

disease (d)[8,45]. To make the model as conservative as possible, we did not alter

the direct inhibitory effect of algae on coral across healthy and degraded reefs,

although it is enhanced on unhealthy reefs[44,45].

In **Equation 3**, algal population size (A) is solved in the last step of each iteration

as the remainder of the benthos not occupied by coral (1 – C) to account for

the fact that the benthos is a finite space.

Note that **Equations 2** and **3** are used for all iterations of the model with or

without viral lysis and metabolism.

For the 'with lytic viruses' model shown in **Figures 5b** and **5e** we used the

following equations, with bold terms reflecting varied parameter values and with

new terms compared to **Equations 1** and **Equation 2** underlined:

B_{new} = B + a * A + **m * B** $\underline{- \textbf{k} * \textbf{V}}$ Equation 4

$\underline{V_{new}}$ = V + **k * B**   Equation 5

Building on **Equation 1**, lytic viruses are introduced in **Equation 4** as a mortality

term for bacteria as an outcome of viral population size (V) and lysis rate (k).

A viral population (V) is added to the model in **Equation 5**, where viruses

proliferate at burst size (k) from lysis if bacteria. Note that k changes between

healthy and degraded reef scenarios, consistent with Piggyback-the-Winner

dynamics suppressing lysis on degraded reefs[14].

For the 'with viral lysis and metabolism' model shown in **Figures 5c** and **5f** we

used the following equations, with bold terms reflecting varied parameter values

and with new terms compared to **Equations 4** and **5** underlined:

B_{new} = B + a * A + **m * B** – k * V $\underline{+ \textbf{t} * \textbf{V}}$ Equation 6

V_{new} = V + **k * B** $\underline{+ \textbf{l} * \textbf{V}}$ Equation 7

This model builds on **Equation 4** by adding viral metabolic function (t) that

enhances bacterial proliferation in degraded reef scenarios associated with

temperate Piggyback-the-Winner dynamics in **Equation 6**[14]. **Equation 7** builds on

Equation 5 by adding lytic functions that enhance virion production in healthy

reefs (l).

Results (1724 words)

Hallmarks of viral temperateness: To assess how viral metabolism across the lytic-temperate spectrum could affect coral reefs, we examined how the fraction of sequenced reads that mapped to known genes in broad metabolic categories (SEED Database Level 1 annotation) changed with community temperateness. Building on prior work[14], we used the frequency of integrase and excisionase genes as a proxy for community temperateness (i.e., hallmarks of temperateness) in viral communities across the Central Pacific Ocean (**Supplementary Table 1**). As a measure of how consistently or monotonously gene frequency and sample temperateness covary, Spearman Rho (ρ_s) values reflect the strength and sign of their correlation. In this analysis, temperate-associated genes have positive correlations ($\rho_s > 0$) while lytic-associated genes have negative correlations ($\rho_s < 0$. For clarity, 95 % confidence intervals are listed in square brackets as ρ_s [95 % CI minimum, 95 % CI maximum].

Gene names and corresponding pathways, EC numbers, median ρ_s and 95 % CIs, gene category (anaplerotic vs cataplerotic), and reports of the gene in viruses are shown in **Table 1**. **Figure 1** and **Figure 2** display the sample-level patterns of temperateness vs gene frequency in all Level 1 SEED categories and in central carbon metabolism functions, respectively. The correlations shown in **Figure 2** are represented in a metabolic pathway map in **Figure 3** and analyzed according to **Table 1** categories in **Figure 4**. Finally, the implications of our central findings are incorporated into a schematic ecosystem model shown in **Figure 5**.

Temperate viral communities are enriched in metabolic genes: The frequency of genes in the Phages, Prophages, Transposable Elements, and Plasmids category declined with increasing temperateness of viral communities (**Figure 1**; $\rho_s = -0.81$ [-0.95, -0.58]). In contrast, the frequency and diversity of 'bacterial' metabolic genes that encode known cellular functions increased in temperate viral

communities (**Supplementary Figure 1**; ρ_s = 0.63 and 0.81, respectively; c.f.,

genetic diversity declines with temperateness when unknown genes are

included[14]). Altogether, this analysis indicates that temperate viral communities

are more metabolically similar to bacteria than are lytic communities.

Functional homologues as viral lifestyle markers: Analogous to taxonomic

markers in cells, where the prevalence of class I and class II homologues of

fructose bisphosphate aldolase EC4.1.2.13 genes relates to bacterial

taxonomy[46], the divergent correlations in these genes across a viral

temperateness spectrum may reflect viral life strategies (EC 4.1.2.13 I ρ_s = -0.65 [-

0.86, 0.33] while EC4.1.2.13 II ρ_s = 0.56 [-0.18, 0.83]; see **Table 1** for summary

statistics).

Photosynthesis genes in lytic communities: The decline in 'Phages, Prophages,

Transposable Elements, and Plasmid' genes from ~ 50 % of the known genes in

lytic viromes to ~ 5 % in temperate viromes caused the gene frequency in almost

all other categories to increase (as total gene frequencies sum to 100 %; 17 of 22

categories had ρ_s > 0.5; **Figure 1**). A notable exception is Photosynthesis genes:

these had higher frequencies in lytic communities vs. temperate communities in

the Level 1 analysis (ρ_s = -0.78 [-0.92, -0.53]; **Figure 1**). Photosynthesis genes,

including rhodopsins, were also lytic-associated at the individual-gene level,

where 16 out of 20 photosynthesis genes had negative ρ_s values (**Supplementary**

Table 3). Finally, photosynthesis-related electron transport (Pet) genes that

couple Photosystems II and I are absent from the viromes (**Supplementary Table**

3).

Carbohydrate metabolism genes: Given the high prevalence of carbohydrate

genes in the viromes, their elevated frequency in temperate communities

(**Figure 1;** ρ_s > 0), and their ecological importance, we next focused on genes

comprising central carbon metabolism[26,27]: Embden-Meyerhof-Parnas (EMP) and

Entner-Doudoroff (ED) glycolysis pathways, the pentose phosphate pathway

(PPP) and the Tricarboxylic Acid Cycle (TCA; **Figure 2**). Note that for this and

subsequent analyses we retained only sequences with hits to SEED database

entries with high-confidence Enzyme Commission (EC) annotations[47].

Entner-Doudoroff genes are enriched in temperate viromes: Few patterns were

readily apparent from the ρ_s values for genes in each pathway (**Figure 2;**

Supplementary Figure 2). When viewed as a pathway map, though, the ED

pathway was more clearly enriched in temperate viral communities (i.e., the red

arrows running through the center of **Figure 3**). This map identified the ED

pathway as the only pathway that changed consistently across the lytic-

temperate spectrum (**Figure 4a**), and was exclusively over-represented in

temperate viral communities (i.e., $\rho_s > 0$; EC 1.1.1.49 $\rho_s = 0.26$ [-0.24, 0.68],

EC3.1.1.31 $\rho_s = 0.43$ [0.03, 0.77], EC4.1.2.14 $\rho_s = 0.67$ [0.36, 0.90], and EC4.2.1.12 ρ_s

$= 0.71$ [0.47, 0.89]). Altogether, this gave the ED pathway an overall median $\rho_s =$

0.55 in contrast to overall median $\rho_s = 0.03$ for EMP glycolysis, $\rho_s = 0.175$ for

gluconeogenesis (GNG), $\rho_s = 0.03$ for the PPP, and $\rho_s = -0.10$ for the TCA (see

Figure 4a).

Anaplerotic temperate metabolism: Given the role of ED glycolysis in generating

central metabolite pools for use in diverse pathways (i.e., ED is an anaplerotic

pathway), we investigated whether temperate viral communities were enriched

in anaplerotic genes more generally (see **Table 1** for a list of anaplerotic and

cataplerotic genes)[33]. Temperate viral communities were, indeed, enriched in

anaplerotic genes, with an overall median $\rho_s = 0.44$ for anaplerotic reactions

and only one gene even weakly lytic associated (6-phosphogluconate

dehydrogenase EC 1.1.1.44 from the PPP; $\rho_s = -0.06$ [-0.54, 0.45]; **Figure 4b**).

Amino acid precursor genes are enriched in lytic viromes: In contrast to the ED-

enrichment of temperate communities, the pathway map does not show many

areas of contiguous lytic-associated genes (**Figure 3**). However, the enrichment

of non-oxidative PPP genes in lytic communities, especially transaldolase EC

2.2.1.2 (ρ_s = -0.49 [-0.87, 0.02]). This enzyme consumes the central metabolite pool of glutaraldehyde-3-phosphate (G3P) to produce fructose 6-phosphate and erythrose 4-phosphate. Given that other fructose-6-phosphate producing genes are not similarly enriched in lytic communities (e.g., EC 2.2.1.1 in the PPP ρ_s = -0.08 [-0.52, 0.35], EC5.3.1.9 in EMP ρ_s = 0.45 [0.12, 0.73], and EC3.1.3.11 in gluconeogenesis ρ_s = 0.24 [-0.20, 0.66]), this suggests that the primary outcome of the non-oxidative PPP in lytic viral communities is the cataplerotic consumption of G3P for the generation of erythrose-4-phosphate, a precursor for amino acid biosynthesis[48].

Redox genes are over-represented in lytic communities: Given this cataplerotic focus in lytic communities, we investigated whether lytic communities were more strongly associated with individual cataplerotic reactions regardless of pathway rather than with whole pathways. Focusing on the distribution of anabolic redox steps across all the central carbon metabolism pathways, we found that NADPH-generating reactions were not consistently associated with either lytic or temperate communities (EC1.1.1.44 ρ_s = -0.06, EC1.1.1.42 ρ_s = -0.19, and EC1.1.1.49 ρ_s = 0.26, giving a median ρ_s = -0.06). In contrast, NADH-producing oxidoreductase genes were universally over-represented in lytic communities (i.e., all ρ_s < 0; median ρ_s = -0.13 across all NADH-producing reactions from EC1.1.1.37 ρ_s = -0.19 [-0.61, 0.29], EC1.1.1.41 ρ_s = -0.13 [-0.54, 0.34], EC1.2.1.12 ρ_s = -0.20 [-0.60, 0.26], EC1.2.4.1 ρ_s = -0.12 [-0.60, 0.36], EC1.2.4.2 ρ_s = -0.10 [-0.59, 0.42]; **Table 1**; **Figures 2**, **3,** and **4**). This included all redox genes in the TCA being lytic associated (i.e., ρ_s < 0, with an overall median ρ_s = -0.13) in contrast to the other, non-oxidoreductase genes which had a median ρ_s = 0.15, highlighting the specific overrepresentation of redox genes in lytic communities (**Figure 4a**).

Cataplerotic lytic metabolism: Given the association between lytic communities and cataplerotic reactions observed in the PPP and the TCA, we then

investigated whether lytic communities are broadly enriched in cataplerotic

genes. We observed that the overall median ρ_s of cataplerotic reactions was -

0.125 (i.e., these genes were enriched in lytic communities) compared to ρ_s =

0.44 and ρ_s = 0.11 for anaplerotic reactions and reactions that are neither

cataplerotic nor anaplerotic, respectively. This result highlights the association

between lytic viral communities and cataplerotic metabolism.

Implications for Healthy Coral Reefs: To qualitatively describe the ecosystem

effects of viral metabolism, we constructed a schematic model of algal-

bacterial-viral-coral interactions, with all population sizes measured in illustrative

arbitrary units (**Figure 5**). Parameterizing the model to healthy reef values (**Figure**

5a; note that parameterization of the healthy reef scenario largely insulates

coral from the effects of bacterial overgrowth due to the small d value used;

see **Supplementary Table 2** for parameters and values)[1,8,9,14,43], the system

became coral-dominated by t = 5, with coral-bacterial coexistence (coral and

bacteria had increased from 0.5 a.u. initial values to 1.0 a.u. and 0.89 a.u.,

respectively, by t = 6). While the addition of viral lysis (**Figure 5b**) did not

markedly change coral or algal dynamics, it led to the extinction of bacteria by

t = 6 and to an ecosystem dominated by coral and viruses (coral and viruses

had increased from 0.5 a.u. to 1.0 a.u. and 0.97 a.u., respectively, by the end of

the simulation). Finally, adding in viral metabolism led to an earlier extinction of

bacteria (by t = 5) and enhanced viral population size (viruses were 2.29 a.u. by

t = 6), although coral and algal population dynamics were unaffected (**Figure**

5c). Altogether, the model recaptured the coral-dominated status of healthy

coral reefs and showed that viral metabolism likely exerts a profound negative

effect on bacterial populations and a strongly positive effect on viral

populations on healthy reefs.

Implications for Degraded Coral Reefs: In contrast to healthy systems, our model

suggests that degraded coral reefs would be overgrown with algae rather than

coral (**Figure 5**). Note that the parameterization of the degraded reef scenarios

leads to enhanced deleterious bacterial effects on coral health. Coral went

extinct by $t = 5$ in the degraded scenarios without viruses and with viral lysis only

(**Figure 5d and e**), as algal overgrew the benthos and bacterial proliferated to

levels much higher than in any of the healthy scenarios ($B = 2.75$ a.u. and 2.64

a.u. in the no virus and lysis-only models, respectively, by $t = 6$). In contrast to

bacteria, viral populations did not proliferate markedly in the viral-lysis model,

increasing from 0.5 a.u. to only 0.64 a.u. by $t = 6$ (**Figure 5e**). Adding in viral

metabolism that enhanced bacterial proliferation (t) allowed bacteria to

proliferate even further (reaching 3.80 a.u.), supporting a modest rise in viruses

(to 0.75 a.u.), and driving the coral to extinction sooner (by $t = 4$; **Figure 5f**).

Altogether, our qualitative model suggests that viral metabolism supporting

bacterial proliferation can enhance microbial overgrowth of reefs and promote

shifts towards algal dominance of the benthos.

Discussion (551 words)

The role of microbial overgrowth in coral reef decline is well established[1–3,8]. It is further intensified by shifts in bacterial metabolism and viral lifestyles that promote bacterial proliferation[14,17]. Here, we show that this process is further exacerbated by viral-encoded metabolic genes, whereas healthy ecosystems may be protected by viral lysis and metabolic genes that enhance virion production. Because temperate viral infection is tightly linked to host physiology and environmental conditions such as nutrient availability[14,49,50], targeting bacterial metabolic states through manipulation of nutrients could enable a shift in viral effects from degradative to protective, presenting a promising new conservation strategy.

Viruses become increasingly temperate on degraded reefs, following Piggyback-the-Winner (PtW) dynamics[14,23], in which abundant, rapidly proliferating bacterial hosts select for non-lethal, lysogenic infection. This switch in viral lifestyle suspends the top-down control exerted by lytic viruses, allowing bacterial populations to expand unchecked by lethal infection[14,50]. Temperate infection not reduces lysis but also promotes horizontal gene transfer of pathogenicity genes[51], which can help bacteria evade predation by eukaryotic grazers and potentially compromise coral health[14,23]. This forms a positive feedback loop wherein bacterial growth triggers viral lifestyle switching, which in turn facilitates further bacterial growth, establishing a 'Temperate Ratchet'[23] that locks ecosystems into ever-exacerbating bacterial overgrowth and coral disease[14]. Worse, here we show that the temperate viruses on degraded reefs also harbor metabolic genes, such as those for Entner-Doudoroff glycolysis, that can directly support bacterial proliferation and overgrowth of reefs[1], thereby enhancing the Temperate Ratchet[23].

Healthy reefs, in contrast, appear to be guarded by lytic viruses. Lytic infection rates are driven by virus-host predator-prey encounter frequencies that are the outcome of host and viral abundances[13–15]. As a result, as bacterial populations grow, so should lytic infections and lysis, precluding bacterial overgrowth[14,17]. Further, the viral communities on healthy reefs do not harbor the high frequency of pathogenicity genes seen on degraded reefs, meaning that grazing and coral health are not being hindered by viral-encoded genes, which also helps to keep bacterial densities low[8]. Here we show that lytic viral communities on healthy reefs are enriched in metabolic genes associated with efficient energy production and biosynthesis, such as those for the Embden-Meyerhof-Parnas glycolysis pathway[1,10] and oxidoreductases involved in ATP generation and NADH reduction[52]. These metabolic features support robust virion production that can support reef 'viralization' through high viral abundances and high rates of virus-host encounters, infection, and lysis. In this way, lytic viral metabolism and reef viralization acts as a counterbalance to the Temperate Ratchet, helping to stabilize healthy reef ecosystems[34].

Viral metabolism associated with the lytic-to-temperate shift likely exacerbates coral reef decline, as shown here. Building on ecosystem-scale processes like viral lifestyle switching[14,17], the metabolic capacity of temperate and lytic viruses can further drive reefs into two stable states: healthy reefs protected by metabolically enhanced lytic control of bacteria, and degraded reefs ratcheted into persistent microbialization under temperate infection and metabolism. Ultimately, degraded coral reefs experience a metabolic disorder, where excess nutrients trigger ecosystem-scale dysregulation through microbial and viral lifestyle and metabolic switching[1–3,7]. However, the physiological basis of this switch also suggests that conservation measures targeting nutrient supply could undercut copiotrophic bacterial and viral metabolism[4,10,11]. Our findings suggest that doing so could lead to a 're-viralization' of the reef system,

555 harnessing viral metabolic and lifestyle switching to enhance reef restoration

556 and resilience.

Data Availability

Viral metagenomes can be found on MG-RAST under the accession numbers 4683670.3, 4683674.3, 4683677.3, 4683680.3, 4683683.3, 4683684.3, 4683686.3, 4683703.3, 4683704.3, 4683712.3, 4683718.3, 4683720.3, 4683731.3, 4683733.3, 4683745.3, 4683746.3, and 4683747.3 (see **Supplementary Table 1** for library data summaries; www.mg-rast.org). Fasta files for each sample can also be found at https://tinyurl.com/viral-metabolic-ratchet. Statistical and modeling code can be found at https://tinyurl.com/viral-metabolic-ratchet-code.

Acknowledgments

This work was supported by a BIG Summer stipend from Simons Foundation (awarded to Alexander Hoffmann for CU and NY) as well as by NSF Award OCE-2201645 to BK that provided Hopeful Monsters Summer stipends to JK, MK, GD, NF, EG, AK, and IT. LWK was supported by NSF award OCE-2118617. RAE was supported by an award from the NIH NIDDK RC2DK116713, an award from the Australian Research Council DP220102915, and an award from the Gordon and Betty Moore Foundation number 9871 (Perpetual Viral Origins). JG was supported by NSF Biology Integration Institute award 2119968. ML was supported by an NSF Postdoctoral Research Fellowship in Biology award 2209377. The authors would like to thank the captain, crew, and staff of the *NOAAS Hi'ialakai* (R 334) and National Oceanic and Atmospheric Administration (NOAA) shore support, as well as the members of the Pacific Reef Assessment and Monitoring Program of NOAA's National Coral Reef Monitoring Program.

References

1. Haas, A. F. *et al.* Global microbialization of coral reefs. *Nat. Microbiol.* **1**,
16042 (2016).

2. Kelly, L. W., Haas, A. F., Nelson, C. E. & Rohwer, F. Microbialization in coral
reefs: insights into the mechanisms and consequences of a changing
benthos. *mBio* **5**, e00080-14 (2014).

3. McDole, T. *et al.* Assessing Coral Reefs on a Pacific-Wide Scale Using the
Microbialization Score. *PLoS ONE* **7**, e43233 (2012).

4. Nelson, C. E., Wegley Kelly, L. & Haas, A. F. Microbial interactions with
dissolved organic matter are central to coral reef ecosystem function and
resilience. *Annu. Rev. Mar. Sci.* **15**, 431–460 (2023).

5. Haas, A. F. *et al.* Influence of coral and algal exudates on microbially
mediated reef metabolism. *PeerJ* **1**, e108 (2013).

6. McDole Somera, T. *et al.* Energetic differences between bacterioplankton
trophic groups and coral reef resistance. *Proc. R. Soc. B Biol. Sci.* **283**,
20160467 (2016).

7. Smith, J. E. *et al.* Indirect effects of algae on coral: algae-mediated,
microbe-induced coral mortality. *Ecol. Lett.* **9**, 835–845 (2006).

8. Barott, K. L. & Rohwer, F. L. Unseen players shape benthic competition on
coral reefs. *Trends Microbiol.* **20**, 621–628 (2012).

9. Roach, T. N. *et al.* Microbial bioenergetics of coral-algal interactions. *PeerJ* **5**,
e3423 (2017).

10. Nelson, C. E. *et al.* Coral and macroalgal exudates vary in neutral sugar
composition and differentially enrich reef bacterioplankton lineages. *ISME J.*
7, 962–979 (2013).

11. Silveira, C. B. *et al.* Microbial processes driving coral reef organic carbon
flow. *FEMS Microbiol. Rev.* **41**, 575–595 (2017).

12. Flamholz, A., Noor, E., Bar-Even, A., Liebermeister, W. & Milo, R. Glycolytic
strategy as a tradeoff between energy yield and protein cost. *Proc. Natl.*
Acad. Sci. **110**, 10039–10044 (2013).

13. Thingstad, T. F. Elements of a theory for the mechanisms controlling
abundance, diversity, and biogeochemical role of lytic bacterial viruses in
aquatic systems. *Limnol. Oceanogr.* **45**, 1320–1328 (2000).

14. Knowles, B. *et al.* Lytic to temperate switching of viral communities. *Nature*
531, 466–470 (2016).

15. Suttle, C. A. Marine viruses--major players in the global ecosystem. *Nat. Rev.*
Microbiol. **5**, 801–12 (2007).

16. Fuhrman, J. A. Marine viruses and their biogeochemical and ecological
effects. *Nature* **399**, 541–8 (1999).

17. Silveira, C. B. *et al.* Viral predation pressure on coral reefs. *BMC Biol.* **21**, 77
(2023).

18. Friman, V.-P. & Buckling, A. Phages can constrain protist predation-driven
attenuation of Pseudomonas aeruginosa virulence in multienemy
communities. *ISME J.* **8**, 1820–1830 (2014).

19. Aijaz, I. & Koudelka, G. B. Cheating, facilitation and cooperation regulate the
effectiveness of phage-encoded exotoxins as antipredator molecules.
Microbiologyopen **8**, e00636 (2019).

20. Brüssow, H. Bacteria between protists and phages: from antipredation
strategies to the evolution of pathogenicity. *Mol. Microbiol.* **65**, 583–589
(2007).

21. Cram, J., Parada, A. & Fuhrman, J. Dilution reveals how viral lysis and grazing
shape microbial communities. *Limnol. Oceanogr.* **61**, 889–905 (2016).

22. Fuhrman, J. A. & Noble, R. T. Viruses and protists cause similar bacterial
mortality in coastal seawater. *Limnol. Oceanogr.* **40**, 1236–1242 (1995).

23. Knowles, B. & Rohwer, F. Knowles & Rohwer Reply. *Nature* **549**, E3–E4 (2017).

24. Breitbart, M., Bonnain, C., Malki, K. & Sawaya, N. A. Phage puppet masters of
the marine microbial realm. *Nat. Microbiol.* **3**, 754–766 (2018).

25. Brum, J. R. *et al.* Patterns and ecological drivers of ocean viral communities.
Science **348**, 1261498 (2015).

26. Hurwitz, B. L. & U'Ren, J. M. Viral metabolic reprogramming in marine
ecosystems. *Curr. Opin. Microbiol.* **31**, 161–168 (2016).

27. Howard-Varona, C. *et al.* Phage-specific metabolic reprogramming of
virocells. *ISME J.* **14**, 881–895 (2020).

28. Roach, T. N. *et al.* A multiomic analysis of in situ coral–turf algal interactions.
Proc. Natl. Acad. Sci. **117**, 13588–13595 (2020).

29. Thompson, L. R. *et al.* Phage auxiliary metabolic genes and the redirection of
cyanobacterial host carbon metabolism. *Proc. Natl. Acad. Sci. U. S. A.* **108**,
E757-64 (2011).

30. Thompson, L. R., Zeng, Q. & Chisholm, S. W. Gene expression patterns during
light and dark infection of Prochlorococcus by cyanophage. *PloS One* **11**,
e0165375 (2016).

31. Fogg, P. C. M. Gene transfer agents: The ambiguous role of selfless viruses in
genetic exchange and bacterial evolution. *Mol. Microbiol.* **123**, 124–131
(2025).

32. Lücking, D., Mercier, C., Alarcón-Schumacher, T. & Erdmann, S. Extracellular
vesicles are the main contributor to the non-viral protected extracellular
sequence space. *ISME Commun.* **3**, 112 (2023).

33. Lehninger, A. L. Bioenergetics. The molecular basis of biological energy
transformations. New York: W.A. Benjamin (1965).

34. Baer, J. *et al.* Viralization as a microbial approach for enhancing coral reef
restoration. *ISME J.* **19**, wraf110 (2025).

35. Haas, A. F. *et al.* Unraveling the unseen players in the ocean-a field guide to
water chemistry and marine microbiology. *JoVE J. Vis. Exp.* e52131 (2014).

36. Sambrook, J., Fritsch, E. F. & Maniatis, T. *Molecular Cloning: A Laboratory*
Manual. 2nd ed. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory
Press (1989).

37. Thurber, R. V., Haynes, M., Breitbart, M., Wegley, L. & Rohwer, F. Laboratory
procedures to generate viral metagenomes. *Nat. Protoc.* **4**, 470–483 (2009).

38. Schmieder, R. & Edwards, R. Quality control and preprocessing of
metagenomic datasets. *Bioinformatics* **27**, 863–864 (2011).

39. Schmieder, R., Lim, Y. W., Rohwer, F. & Edwards, R. TagCleaner: Identification
and removal of tag sequences from genomic and metagenomic datasets.
BMC Bioinformatics **11**, 1–14 (2010).

40. Schmieder, R. & Edwards, R. Fast identification and removal of sequence
contamination from genomic and metagenomic datasets. *PloS One* **6**,
e17288 (2011).

41. Meyer, F. *et al.* The metagenomics RAST server – a public resource for the
automatic phylogenetic and functional analysis of metagenomes. *BMC*
Bioinformatics **9**, 386 (2008).

42. Kelly, L. W. *et al.* Black reefs: iron-induced phase shifts on coral reefs. *ISME J.*
6, 1–12 (2011).

43. Barott, K. L. *et al.* Microbial to reef scale interactions between the reef-
building coral Montastraea annularis and benthic algae. *Proc. R. Soc. B Biol.*
Sci. **279**, 1655–1664 (2012).

44. George, E. E. *et al.* Space-filling and benthic competition on coral reefs.
PeerJ **9**, e11213 (2021).

45. Barott, K. *et al.* Hyperspectral and physiological analyses of coral-algal
interactions. *PloS One* **4**, e8043 (2009).

46. Pirovich, D. B., Da'dara, A. A. & Skelly, P. J. Multifunctional fructose 1, 6-
bisphosphate aldolase as a therapeutic target. *Front. Mol. Biosci.* **8**, 719678
(2021).

47. Webb, E. C. *Enzyme Nomenclature 1992. Recommendations of the*
Nomenclature Committee of the International Union of Biochemistry and
Molecular Biology on the Nomenclature and Classification of Enzymes.
Academic Press, San Diego (1992).
48. Srinivasan, P., Katagiri, M. & Sprinson, D. B. The conversion of
phosphoenolpyruvic acid and d-erythrose 4-phosphate to 5-dehydroquinic
acid. *J. Biol. Chem.* **234**, 713–715 (1959).
49. Knowles, B. *et al.* Variability and host density independence in inductions-
based estimates of environmental lysogeny. *Nat. Microbiol.* **2**, 17064 (2017).
50. Knowles, B. *et al.* Temperate infection in a virus–host system previously known
for virulent dynamics. *Nat. Commun.* **11**, 1710 (2020).
51. Howard-Varona, C., Hargreaves, K. R., Abedon, S. T. & Sullivan, M. B.
Lysogeny in nature: mechanisms, impact and ecology of temperate phages.
ISME J. **11**, 1511 (2017).
52. Hurwitz, B. L., Hallam, S. J. & Sullivan, M. B. Metabolic reprogramming by
viruses in the sunlit and dark ocean. *Genome Biol.* **14**, 1–14 (2013).
53. Devoto, A. E. *et al.* A new family of "megaphages" abundant in the marine
environment. *Nat. Microbiol.* **6**, 123–131 (2021).
54. Anantharaman, K. *et al.* Sulfur oxidation genes in diverse deep-sea viruses.
Science **344**, 757–760 (2014).
55. Moniruzzaman, M., Martinez-Gutierrez, C. A., Weinheimer, A. R. & Aylward, F.
O. Dynamic genome evolution and complex virocell metabolism of globally-
distributed giant viruses. *Nat. Commun.* **11**, 1710 (2020).
56. Ha, A. D., Moniruzzaman, M. & Aylward, F. O. High transcriptional activity and
diverse functional repertoires of hundreds of giant viruses in a coastal marine
system. *mSystems* **6**, e0074321 (2021).
57. Rodrigues, R. A. L., Arantes, T. S., Oliveira, G. P., Silva, L. K. & Abrahão, J. S. The
complex nature of Tupanviruses. *Adv. Virus Res.* **103**, 135–162 (2019).
58. Schroeder, D. C. et al. Viruses infecting a warm water picoeukaryote shed
light on spatial and temporal patterns in virus community ecology. *ISME J.* **15**,
3621–3634 (2021).
722

Tables and Figures.

Table 1: Central Carbon Metabolism genes shown in Figures 2 and 3. Genes are categorized by pathway and gene function, identified by EC number and gene name, and annotated with prior references if they have been observed in viruses before. The Spearman's correlation between the frequency of each gene and viromes temperateness is indicated by bootstrapped mean ρ_s values and 95 % confidence intervals from 1,000 iterations are shown. Please refer to **Supplementary Figure 2** for bootstrap ρ_s distributions. Genes shared between pathways are shown with a slash. Note, although anaplerotic and cataplerotic reactions often refer to steps in the TCA, we are using the term more generally here to refer respectively to the generation of central intermediate metabolites ('filling up'; anaplerotic) and the consumption of central metabolites to generate specialized pre-cursor or cellular energy pools ('drawing down'; cataplerotic) more generally[33]. Bootstrapping code can be found at https://tinyurl.com/viral-metabolic-ratchet-code.

EC number	SEED database gene name	Pathway	Gene activity	ρ_s	95 % CI	Observed in viruses
1.1.1.37	Malate dehydrogenase	Tricarboxylic Acid	Cataplerotic	-0.19	-0.61, 0.29	
1.1.1.41	Isocitrate dehydrogenase [NAD]	Tricarboxylic Acid	Cataplerotic	-0.13	-0.54, 0.34	
1.1.1.42	Isocitrate dehydrogenase [NADP]	Tricarboxylic Acid	Cataplerotic	-0.19	-0.63, 0.29	52,53
1.1.1.44	6-phosphogluconate dehydrogenase	Pentose Phosphate	Anaplerotic	-0.06	-0.54, 0.45	29,52,54
1.1.1.49	Glucose-6-phosphate 1-dehydrogenase	ED glycolysis/ Pentose Phosphate	Anaplerotic	0.26	-0.24, 0.68	
1.2.1.12	NAD-dependent glyceraldehyde-3-phosphate dehydrogenase	EMP glycolysis/ Gluconeogenesis	Cataplerotic	-0.20	-0.60, 0.26	
1.2.4.1	Pyruvate dehydrogenase	Tricarboxylic Acid	Cataplerotic	-0.12	-0.60, 0.36	52
1.2.4.2	2-oxoglutarate dehydrogenase	Tricarboxylic Acid	Cataplerotic	-0.10	-0.59, 0.42	
2.2.1.1	Transketolase	Pentose Phosphate		-0.08	-0.52, 0.35	29,52,54
2.2.1.2	Transaldolase	Pentose Phosphate	Cataplerotic	-0.49	-0.87, 0.02	7,45,47
2.3.3.1	Citrate synthase	Tricarboxylic Acid		0.15	-0.32, 0.56	
2.3.3.9	Malate synthase	Tricarboxylic Acid	Anaplerotic	0.61	0.20, 0.90*	55
2.7.1.11	6-phosphofructokinase	EMP glycolysis	Cataplerotic	-0.03	-0.48, 0.45	
2.7.1.40	Pyruvate kinase	EMP glycolysis	Cataplerotic	0.42	0.04, 0.74*	
2.7.2.3	Phosphoglycerate kinase	EMP glycolysis/ Gluconeogenesis	Cataplerotic	0.03	-0.41, 0.47	
3.1.1.31	6-phosphogluconolactonase	ED glycolysis/ Pentose Phosphate	Anaplerotic	0.43	0.03, 0.77*	
3.1.3.11	Fructose-1,6-bisphosphatase	Gluconeogenesis	Anaplerotic	0.24	-0.20, 0.66	55
4.1.1.31	Phosphoenolpyruvate carboxylase	Gluconeogenesis	Anaplerotic	0.45	0.00, 0.77*	
4.1.1.32	Phosphoenolpyruvate carboxykinase [GTP]	Gluconeogenesis		0.11	-0.39, 0.57	55,56
4.1.2.13 I	Fructose-bisphosphate aldolase class I	EMP glycolysis/ Gluconeogenesis	Cataplerotic	-0.65	-0.86, 0.33	
4.1.2.13 II	Fructose-bisphosphate aldolase class II	EMP glycolysis/ Gluconeogenesis	Cataplerotic	0.56	-0.18, 0.83	
4.1.2.14	2-dehydro-3-deoxyphosphogluconate aldolase	ED glycolysis	Anaplerotic	0.67	0.36, 0.90*	

4.1.3.1	Isocitrate lyase	Tricarboxylic Acid		-0.10	-0.54, 0.37	
4.2.1.11	Enolase	EMP glycolysis/ Gluconeogenesis		0.43	0.02, 0.76*	[55]
4.2.1.12	Phosphogluconate dehydratase	ED glycolysis	Anaplerotic	0.71	0.47, 0.89*	
4.2.1.2	Fumarate hydratase class II	Tricarboxylic Acid		0.29	-0.15, 0.68	
4.2.1.3	Aconitate hydratase	Tricarboxylic Acid		-0.35	-0.71, 0.13	[55,57]
5.1.3.1	Ribulose-phosphate 3-epimerase	Pentose Phosphate		0.03	-0.45, 0.49	[58]
5.3.1.1	Triosephosphate isomerase	EMP glycolysis/ Gluconeogenesis		-0.02	-0.44, 0.41	[55]
5.3.1.6	Ribose 5-phosphate isomerase	Pentose Phosphate		0.32	-0.14, 0.68	[54]
5.3.1.9	Glucose-6-phosphate isomerase	EMP glycolysis/ Gluconeogenesis		0.45	0.12, 0.73*	
5.4.2.11	Phosphoglycerate mutase	EMP glycolysis/ Gluconeogenesis		-0.46	-0.80, 0.01	[55,56]
5.4.2.12	2,3-bisphosphoglycerate-independent phosphoglycerate mutase	EMP glycolysis/ Gluconeogenesis		0.41	-0.01, 0.73	
6.2.1.5a	Succinyl-CoA ligase [ADP-forming] alpha chain	Tricarboxylic Acid		0.17	-0.33, 0.61	
6.2.1.5b	Succinyl-CoA ligase [ADP-forming] beta chain	Tricarboxylic Acid		-0.04	-0.52, 0.41	

Figure 1: Correlations between metabolic function and community temperateness across the breadth of cellular metabolism. Spearman's rho estimates (ρ_s) are shown with 95 % confidence intervals (CIs) in parentheses from 1,000 bootstrapped simulations. Panels are ordered by mean gene frequency. The y-axis shows the gene frequency as a percentage of known genes, an internally normalized and relative metric where all of the frequencies for a given sample across all panels sums to 100 %. Gene categories with average frequencies below 0.5 % were omitted. Viral community temperateness was calculated as the percentage of integrase and excision genes in the known genes in each virome. Bootstrapping code can be found at https://tinyurl.com/viral-metabolic-ratchet-code.

748
Figure 2: Correlation between central carbon metabolism genes and viral community temperateness. Spearman's rho correlation coefficients (ρ_s) are shown from 1,000 bootstrapped simulations correlating central carbon metabolism gene frequency as a percentage of known genes vs viral community temperateness. Panels are ordered by pathway, including Embden-Meyerhof-Parnas glycolysis (EMP), Entner-Doudoroff glycolysis (ED), pentose phosphate (PP), and tricarboxylic acid cycle pathways. Viral community temperateness was calculated as the percentage of integrase and excision genes in the known genes in each virome. Enzymes with rho values significantly different to zero are indicated by asterisks. Genes are labeled by EC numbers; see **Table 1** for gene names and summary statistics.

758
Figure 3: Pathway map of changes in the frequency of central carbon metabolism genes across
the lytic to temperate spectrum. Map of central carbon metabolism with each enzymatic step
colored by the change in enzymatic metagenomic gene frequency across lytic to temperate
viral communities. Red arrows show higher frequency in temperate communities while blue
arrows are lytic associated, reflecting ρ_s values for each enzyme as seen in **Table 1** and **Figure 2**.
Enzymes are labeled by EC numbers (see **Table 1** for gene names). Single-headed and double-
headed arrows represent irreversible and reversible reactions, respectively. Black arrows show
genes that were absent from all viromes. Central intermediate metabolites glyceraldehyde-3-
phosphate (G3P) and pyryvate are shown in bold. *Note that fructose-6-phosphate is found in
the EMP/gluconeogenesis pathways as well as the PPP, ** that oxaloacetate is produced in both
the TCA and gluconeogenesis, and that EC1.1.1.42 in the TCA has an oxalosuccinate
intermediate. Code can be found at https://tinyurl.com/viral-metabolic-ratchet-code.

771
Figure 4: Correlations between viral community temperateness and central carbon gene
frequency across pathways and gene function. (a) Spearman's rho correlation coefficients (ρ_s)
for genes in the Embden- Meyerhof-Parnas glycolysis (EMP), gluconeogenesis (GNG), pentose
phosphate (PP), Entner-Doudoroff glycolysis (ED, and tricarboxylic acid cycle (TCA) pathways
are shown with median values marked by horizontal bars in each pathway. (b) Coefficients are
also shown for genes involved in building up central intermediate metabolite pools like pyruvate
and glyceraldehyde-3-phosphate (anaplerotic reactions) or consuming general metabolites to
make specialized pools of precursors and cellular energy (cataplerotic reactions) in central
carbon metabolism, compared with genes involved in neither anaplerosis nor cataplerosis, also
with median values shown. Oxidoreductase genes are colored by their electron acceptor: blue
for NADP[+] electron acceptors involved in biomass generation, and red for NAD[+] electron
acceptors involved in cellular energy generation. Dashed horizontal lines indicate ρ_s = 0. Note,
although anaplerotic and cataplerotic reactions often refer to steps in the TCA, we are using the
term more generally here to refer respectively to the generation ('filling up'; anaplerotic) of
flexible general intermediate metabolites and the consumption of general metabolites towards
precursor and cellular energy production ('drawing down'; cataplerotic) more generally[33].
788

789
**Figure 5: Schematic model incorporating viral metabolic effects on resilience of healthy coral
reefs and exacerbation of reef degradation.** The predicted effects of viral lysis and metabolic
capability on algal (A; green), coral (C; coral), bacteria (B; blue) and viral (V; violet) population
sizes in healthy and degraded reef scenarios with no viruses (a and d, respectively), with viral lysis
(b and e, respectively), and with viral lysis and metabolism (c and f, respectively). Compartment
models showing how the model was implemented for each panel are shown (left) as well as
resulting dynamics (right). Dynamics show changes in populations over time, in arbitrary units
(a.u.), reflecting the qualitative nature of the model. In each iteration, algal population size A
was solved as the 1 – C to reflect the finite space available on the benthos. Parameter values,
shown as line weights, qualitatively reflect known processes on healthy vs degraded reefs, and
can be found in **Supplementary Table 2**. Rates include: (i) coral-algal inhibition, (c) coral
proliferation, (a) algal-mediated DOC supply to bacteria, (m) bacterial anabolic metabolism,
(d) bacterial-mediate coral disease, (k) viral lysis of bacteria, (l) lytic viral metabolism, and (t)
temperate viral metabolism. Note that panels **(b)** and **(e)** depict the Kill-the-Winner and
Piggyback-the-Winner models, respectively. Algal, coral, bacteria, and viral pools were initialized
at 0.5 a.u. for all simulations where present. To make the model as conservative as possible,
rates of algal carbon delivery (a), and direct coral-algal interference (i) were not changed
across scenarios; doing so would lead to enhanced loss of coral. Note that flat lines at minimal
values reflect functional population extinction.

Legends only

Table 1: Central Carbon Metabolism genes shown in Figures 2 and 3. Genes are categorized by pathway and gene function, identified by EC number and gene name, and annotated with prior references if they have been observed in viruses before. The Spearman's correlation between the frequency of each gene and viromes temperateness is indicated by bootstrapped mean ρ_s values and 95 % confidence intervals from 1,000 iterations are shown. Please refer to **Supplementary Figure 2** for bootstrap ρ_s distributions. Genes shared between pathways are shown with a slash. Note, although anaplerotic and cataplerotic reactions often refer to steps in the TCA, we are using the term more generally here to refer respectively to the generation of central intermediate metabolites ('filling up'; anaplerotic) and the consumption of central metabolites to generate specialized pre-cursor or cellular energy pools ('drawing down'; cataplerotic) more generally[33]. Bootstrapping code can be found at https://tinyurl.com/viral-metabolic-ratchet-code.

Figure 1: Correlations between metabolic function and community temperateness across the breadth of cellular metabolism. Spearman's rho estimates (ρ_s) are shown with 95 % confidence intervals (CIs) in parentheses from 1,000 bootstrapped simulations. Panels are ordered by mean gene frequency. The y-axis shows the gene frequency as a percentage of known genes, an internally normalized and relative metric where all of the frequencies for a given sample across all panels sums to 100 %. Gene categories with average frequencies below 0.5 % were omitted. Viral community temperateness was calculated as the percentage of integrase and excision genes in the known genes in each virome. Bootstrapping code can be found at https://tinyurl.com/viral-metabolic-ratchet-code.

Figure 2: Correlation between central carbon metabolism genes and viral community temperateness. Spearman's rho estimates (ρ_s) are shown from 1,000

bootstrapped simulations correlating central carbon metabolism gene

frequency as a percentage of known genes vs viral community temperateness.

Panels are ordered by pathway, including Embden- Meyerhof-Parnas glycolysis

(EMP), Entner-Doudoroff glycolysis (ED), pentose phosphate (PP), and

tricarboxylic acid cycle pathways. Viral community temperateness was

calculated as the percentage of integrase and excision genes in the known

genes in each virome. Enzymes with rho values significantly different to zero are

indicated by asterisks. Genes are labeled by EC numbers; see **Table 1** for gene

names and summary statistics.

Figure 3: Figure 3: Pathway map of changes in the frequency of central carbon

metabolism genes across the lytic to temperate spectrum. Map of central

carbon metabolism with each enzymatic step colored by the change in

enzymatic metagenomic gene frequency across lytic to temperate viral

communities. Enzymes are labeled by EC numbers (see **Table 1** for gene names).

Red arrows show higher frequency in temperate communities while blue arrows

are lytic associated, reflecting ρ_s values for each enzyme as seen in **Table 1** and

Figure 2. Single-headed and double-headed arrows represent irreversible and

reversible reactions, respectively. Black arrows show genes that were absent

from all viromes. Central intermediate metabolites glyceraldehyde-3-phosphate

(G3P) and pyryvate are shown in bold. *Note that fructose-6-phosphate is found

in the EMP/gluconeogenesis pathways as well as the PPP, ** that oxaloacetate is

produced in both the TCA and gluconeogenesis, and that EC1.1.1.42 in the TCA

has an intermediate oxalosuccinate product/substrate. Code can be found at

https://tinyurl.com/viral-metabolic-ratchet-code.

Figure 4: Correlations between viral community temperateness and central

carbon gene frequency across pathways and gene function. (a) Spearman's

rho correlation coefficients (ρ_s) for genes in the Embden- Meyerhof-Parnas

glycolysis (EMP), gluconeogenesis (GNG), pentose phosphate (PP), Entner-

Doudoroff glycolysis (ED, and tricarboxylic acid cycle (TCA) pathways are shown with median values marked by horizontal bars in each pathway. (b) Coefficients are also shown for genes involved in building up central intermediate metabolite pools like pyruvate and glyceraldehyde-3-phosphate (anaplerotic reactions) or consuming general metabolites to make specialized pools of precursors and cellular energy (cataplerotic reactions) in central carbon metabolism, compared with genes involved in neither anaplerosis nor cataplerosis, also with median values shown. Oxidoreductase genes are colored by their electron acceptor: blue for $NADP^+$ electron acceptors involved in biomass generation, and red for NAD^+ electron acceptors involved in cellular energy generation. Dashed horizontal lines indicate $\rho_s = 0$. Note, although anaplerotic and cataplerotic reactions often refer to steps in the TCA, we are using the term more generally here to refer respectively to the generation ('filling up'; anaplerotic) of flexible general intermediate metabolites and the consumption of general metabolites towards precursor and cellular energy production ('drawing down'; cataplerotic) more generally.

Figure 5: Schematic model incorporating viral metabolic effects on resilience of healthy coral reefs and exacerbation of reef degradation. The predicted effects of viral lysis and metabolic capability on algal (A; green), coral (C; coral), bacteria (B; blue) and viral (V; violet) population sizes in healthy and degraded reef scenarios with no viruses (a and d, respectively), with viral lysis (b and e, respectively), and with viral lysis and metabolism (c and f, respectively). Compartment models showing how the model was implemented for each panel are shown (left) as well as resulting dynamics (right). Dynamics show changes in populations over time, in arbitrary units (a.u.), reflecting the qualitative nature of the model. In each iteration, algal population size A was solved as the 1 – C to reflect the finite space available on the benthos. Parameter values, shown as line weights, qualitatively reflect known processes on healthy vs degraded reefs, and can be found in **Supplementary Table 2.**

Rates include: (i) coral-algal inhibition, (c) coral proliferation, (a) algal-mediated DOC supply to bacteria, (m) bacterial anabolic metabolism, (d) bacterial-mediate coral disease, (k) viral lysis of bacteria, (l) lytic viral metabolism, and (t) temperate viral metabolism. Note that panels **(b)** and **(e)** depict the Kill-the-Winner and Piggyback-the-Winner models, respectively. Algal, coral, bacteria, and viral pools were initialized at 0.5 for all simulations where present. To make the model as conservative as possible, rates of algal carbon delivery (a), and direct coral-algal interference (i) were not changed across scenarios; doing so would lead to enhanced loss of coral. Note that flat lines at minimal values reflect functional population extinction. Code can be found at https://tinyurl.com/viral-metabolic-ratchet-code.

Figures only

Figure 1

 Figure 2

Figure 3

914 **Figure 4**

Figure 5

Supplementary Tables and Figures

Supplementary Table 1: Summary of the 19 samples analyzed, including sample accession numbers, the total number of base pairs, reads sequenced, and locations of viral metagenomes. Note that all metagenomes have been previously published except Pearl and Hermes Reef Wreck and Caroline Wreck Site samples indicated by an asterisk. Caroline Island is now named Millenium Atoll. Fasta files for all metagenomes can be found at https://tinyurl.com/viral-metabolic-ratchet.

Sample Location	MG-RAST Accession Number	bp count	Sequence count	Latitude	Longitude
Millennium Atoll	4683670.3	47624434	173813	-9.9138	-150.2038
Flint Atoll	4683674.3	7294624	26673	-11.43	-151.8192
Starbuck	4683677.3	28130343	102666	-5.6384	-155.8811
Maui	4683680.3	48918703	178536	20.86447	-156.13911
Oahu	4683683.3	28708747	104972	21.4389	-158
Pearl & Hermes Reef	4683684.3	211441921	771690	27.8557	-175.8318
Kaui	4683686.3	14722075	53831	22.0964	-159.5261
Starbuck	4683703.3	51673568	188592	-5.6384	-155.8811
Niihau	4683704.3	17098607	62505	21.85115	-160.18327
Molokai	4683712.3	31626104	115424	21.17481	-156.8214
Lisianski	4683718.3	51289619	187189	25.98702	-173.99438
Malden Atoll	4683720.3	68368950	249522	-4.0123	-154.9178
French Frigate Shoals	4683731.3	12353528	45131	23.87748	-166.29115
Hawaii	4683733.3	30728163	112147	20.00324	-155.83318
French Frigate Shoals	4683745.3	64763334	236363	23.6417	-155.8778
Lanaii	4683746.3	38160746	139273	20.76167	-156.8334
Millenium Atoll	4683747.3	71166319	259736	-9.9138	-150.2038
Pearl & Hermes Reef Wreck*	4574481.3	24339533	88980	27.8666	-175.7468
Caroline Wreck Site*	4574475.3	11555298	42262	-9.9957	-150.2161

Supplementary Table 2: Parameters and values used in the qualitative
ecosystem model. To isolate the effect of bacterial and viral proliferative genes
on algal, coral, bacteria, and viral populations, as many parameters as possible
were held constant, and only those relating directly to bacterial and viral
proliferation and pathogenicity changed (listed in **bold**). All populations, shown
with arbitrary units to show relative change in Figure 5, were initialized at 0.5.
Note that, to account for finite space on the benthos, algal population size A is
solved from coral population size C as A = 1 – C each iteration. Code can be
found at https://tinyurl.com/viral-metabolic-ratchet-code.

Parameter	Symbol	Healthy reef without viruses	Degraded reef without viruses	Healthy reef with viruses	Degraded reef with viruses	Healthy reef with viruses and viral metabolism	Degraded reef with viruses and viral metabolism
Bacterial replication	m	0.02	0.2	0.02	0.2	0.02	0.2
Bacterial inhibition of coral	d	0.02	0.2	0.02	0.2	0.02	0.2
Coral replication	c	0.2	0.2	0.2	0.2	0.2	0.2
Algal inhibition of coral	i	0.02	0.02	0.02	0.02	0.02	0.02
Algal support of bacteria	a	0.2	0.2	0.2	0.2	0.2	0.2
Viral replication	l	-	-	-	-	0.2	0.02
Viral lysis of bacteria	k	-	-	0.2	0.02	0.2	0.02
Bacterial support of viral replication	k	-	-	0.2	0.02	0.2	0.02
Viral support of bacterial replication	t	-	-	-	-	0.02	0.2
Bacterial initial population	B	0.5	0.5	0.5	0.5	0.5	0.5
Coral initial population	C	0.5	0.5	0.5	0.5	0.5	0.5
Algal initial population	A	0.5	0.5	0.5	0.5	0.5	0.5
Viral initial population	V	-	-	0.5	0.5	0.5	0.5

940 **Supplementary Table 3: The frequency of photosynthesis genes across the lytic-**
941 **temperate spectrum.** Genes were identified using SEED Level 2 categorization
942 and are organized according to the KEGG photosynthesis map. Spearman's
943 Rho values indicate whether genes are over-represented (value < 0) or under-
944 represented (value > 0) in lytic versus temperate viral communities.

Level 2 Category	Photosystem	Function	Gene	Rho
Electron transport and photophosphorylation	Photosystem_II	photosystem II protein D1	PsbA	-0.845344
Electron transport and photophosphorylation	Photosystem_II	photosystem II protein D2	PsbD	-0.7327776
Electron transport and photophosphorylation	Photosystem_II	Cytochrome b559 beta chain	PsbF	-0.3442652
Electron transport and photophosphorylation	Photosystem_II	Photosystem II 10 kDa phosphoprotein	PsbH	0.04303315
Electron transport and photophosphorylation	Photosystem_II	Photosystem II protein	PsbN	-0.400914
Electron transport and photophosphorylation	Photosystem_II	Photosynthetic reaction center H subunit		-0.1290994
Electron transport and photophosphorylation	Photosystem_II	Putative photosynthetic complex assembly protein		0.2581989
Electron transport and photophosphorylation	Photosystem_I	photosystem I P700 chlorophyll a apoprotein subunit Ia	PsaA	-0.5610839
Electron transport and photophosphorylation	Photosystem_I	photosystem I P700 chlorophyll a apoprotein subunit Ib	PsaB	-0.6837071
Electron transport and photophosphorylation	Photosystem_I	photosystem I iron-sulfur center subunit VII	PsaC	-0.4461846
Electron transport and photophosphorylation	Photosystem_I	photosystem I subunit II	PsaD	-0.5711162
Electron transport and photophosphorylation	Photosystem_I	photosystem I subunit IV	PsaE	-0.502189
Electron transport and photophosphorylation	Photosystem_I	photosystem I subunit III plastocyanin docking protein	PsaF	-0.04303315
Electron transport and photophosphorylation	Photosystem_I	photosystem I subunit IX	PsaJ	-0.3442652
Electron transport and photophosphorylation	Photosystem_I	photosystem I subunit X	PsaK	-0.2484191
Light-harvesting complexes	Bacterial_light-harvesting_proteins	Light-harvesting LHI, beta subunit		0.1290994
Light-harvesting complexes	Chlorosome	Chlorosome protein I, 2Fe-2S ferredoxin		-0.2151657
Rhodopsins	Bacteriorhodopsin	Brp-like protein		-0.1844696
Rhodopsins	Proteorhodopsin	Octaprenyl-diphosphate synthase		0.1290994
Rhodopsins	Proteorhodopsin	Proteorhodopsin		-0.4655338

Supplementary Figure 1: Temperate viral communities are more cell-like and are more genetically diverse in known genes. (a) The percent of genes with hits against the SEED database (i.e., percent known genes), and (b) the diversity of those known genes in the viral communities across the lytic to temperate spectrum. Note that the percent known genes is indicative of how metabolically similar the viruses are to cells given that known genes are commonly of identified in the study of cells. Viral community temperateness was calculated as the percentage of integrase and excision genes in the known genes in each virome.

956
Supplementary Figure 2: Bootstrapped distribution of ρ_s from 1,000 iterations
correlating the frequency of central carbon metabolism genes with viral
community temperateness (number of observations vs ρ_s value). Genes are
grouped by pathway. Histogram shows the distribution of bootstrapped ρ_s
values, with pink lines showing the median bootstrapped value and shaded
grey areas showing the 95 % confidence interval of the median ρ_s. Dashed
vertical lines show $\rho_s = 0$. Bootstrapping code can be found at
https://tinyurl.com/viral-metabolic-ratchet-code.