**Effective sequence-to-expression prediction for membrane proteins using machine learning and computational protein design**

Yuxin Shen[1], Juno Underhill[2], Adrian J Mulholland[2], Diego A Oyarzún[1,3]* and Paul Curnow[4]*

[1]School of Biological Sciences, University of Edinburgh, UK. [2]School of Chemistry, University of Bristol, UK. [3]School of Informatics, University of Edinburgh, UK. [4]School of Biochemistry, University of Bristol, UK.

*To whom correspondence should be addressed. Diego Oyarzún: d.oyarzun@ed.ac.uk; Paul Curnow: p.curnow@bristol.ac.uk

The recombinant expression of integral membrane proteins is notoriously challenging. One way to address this challenge is via computational genotype-to-phenotype models that determine how particular sequence features correlate with protein expression levels. The potential of such approaches is yet to be fully realised, at least partly because so few expression datasets are available. Here, we study the sequence-to-expression relationships of a library of 12,248 membrane proteins derived from combinatorial computational design. The expression phenotype of the entire library is assessed in the widely-used recombinant host *Escherichia coli*. We employed selected phenotypic data to train a sequence-to-expression predictor using supervised machine learning, which achieved high classification accuracy on held-out test sequences. This model was then used to infer the expression of >10,000 unmeasured sequences, and validation of the top predictions of both high and low expressers achieved 100% success rate. Using tools from explainable AI, we identified specific sequence positions and substitutions that are most important in dictating cellular expression levels. This analysis was validated by model-guided protein engineering that achieved an 8-fold increase in the purification yield of a poorly-expressing variant. We find that cells accumulate elevated levels of transcript mRNA for high-expressing proteins, and speculate that this arises from efficient translation-coupled membrane insertion which minimises transcript degradation. Our results show that computational protein design in tandem with supervised learning leads to effective models for the discovery of protein variants with improved

34    expression phenotypes, and can decode the molecular basis of membrane protein

35    expression.

## Introduction

Integral membrane proteins constitute ~20% of all proteins and are critically important for cellular life. However, the study of membrane proteins remains plagued by a well-attested and long-standing problem: Why are so few of these proteins tolerated by recombinant hosts? The inadequate (or entirely failed) recombinant production of many membrane proteins remains a major bottleneck for both fundamental research and industrial applications. The interlinked stages of recombinant production - gene expression, membrane localization, bilayer insertion and protein folding - are all potential points of failure, and structural genomics programmes have reported expression failure rates of >90% [1]. Yet the host organism must routinely biosynthesise its own membrane proteins, including some at high abundance, and high-level recombinant expression can certainly be achieved in some cases [1, 2]. This implies that there is specific information encoded within a gene or protein sequence that somehow underpins successful membrane protein production. Structural biology, enzymology, synthetic biology, biomanufacturing, and bioprocessing would all benefit from an improved understanding of the sequence factors that underpin the consistent, reproducible and high-level recombinant production of integral membrane proteins.

Membrane protein expression is a complex and coordinated process [3, 4]. In bacteria such as *Escherichia* (*E.*) *coli* – a common workhorse for recombinant expression [5], and the system used in this study - proteins of the inner (cytoplasmic) membrane are comprised of membrane-spanning $\alpha$-helices of about 21 amino acids in length. These transmembrane helices predominantly feature hydrophobic amino acids so that the protein surface matches the hydrophobic interior of the lipid bilayer. Some proteins feature multiple $\alpha$-helical segments within their sequence, and these transmembrane domains are linked by water-soluble loops to give a multipass topology. Briefly, the main pathway of inner membrane protein biogenesis involves the cotranslational insertion of transmembrane segments into the inner membrane as they emerge from the ribosome exit tunnel. Hydrophobic regions within the nascent polypeptide chain are recognised by the signal recognition particle (SRP) which directs the ribosome-nascent chain complex to the membrane-integral Sec translocon. The growing nascent chain is transferred to the translocon as it exits the

docked ribosome and transmembrane sections then move from the translocon into
the membrane bilayer, where the final stages of folding and assembly can occur.

In principle, the intrinsic sequence code controlling recombinant membrane protein
production could be deciphered by machine learning [6]. This would ultimately
produce a predictive model that could, for example, pre-select expression-
compatible homologues from genomic databases and evaluate the likely impact of
specific mutations on protein abundance. Although a general solution is not yet
available [7], sequence-to-expression models for some integral membrane proteins
have been developed based on multiple sequence features [8], biophysical-structural
and evolutionary properties [9, 10] and language models for codon usage [11]. One
substantial issue is that many of the available training sets specific to membrane
proteins are relatively small and do not necessarily have data characteristics that are
compatible with machine learning techniques [6].

Here, we introduce an alternative strategy that exploits large variant libraries
generated by computational sequence design. This approach draws upon our recent
description of a *de novo* integral membrane protein [12]. The goal of this prior work
was to design a small, heme-binding membrane protein inspired by the heme
centres that facilitate electron transport in respiration and photosynthesis. To this end
we used computational tools to transform a soluble di-heme four-helix bundle [13]
into a membrane protein by 'surface-swapping'; that is, the computational
substitution of solvent-facing residues to render the protein exterior hydrophobic.
During this sequence design the residues involved in critical protein-protein and
protein-heme interactions were strictly maintained, while 54 exterior residues were
allowed to sample a restricted hydrophobic amino acid alphabet within an implicit
membrane energy function. This surface-swapping approach generated thousands
of individual sequence designs (decoys) and a few of these protein sequences were
selected for genetic encoding and further characterisation. We found that one of
these designer sequences, which we named CytbX, was successfully expressed in
recombinant *E. coli* and could be isolated from the cell inner membrane in the
hemoprotein form [12]. Other variants from the same design run were similarly well-
folded but were expressed at lower levels.

We reasoned that this novel design library could be used to build genotype-
phenotype models of membrane protein expression, because the controlled
combinatorial diversity within these 'designologues' is balanced by tight constraints
over variables such as sequence length, codon usage, number of transmembrane
(TM) helices and loop composition. To provide an expression readout these *de novo*
proteins can be genetically fused to green fluorescent protein (GFP), an approach
that has been deployed at scale to rapidly phenotype and sort very large protein
libraries [14] and is widely used to measure the recombinant expression of diverse
sets of integral membrane proteins [2, 15-22]. While *functional* expression can be
different from *total* expression [10, 23], high-expressing GFP fusions are generally a
good starting point for molecular studies [24].

Here we show that for this novel dataset, assumption-free and mechanism-agnostic
machine learning can discern the sequence basis of membrane protein expression
and accurately infer the expression levels of related unseen proteins. Experimental
validation reveals that the expression code learned by the model is context-
dependent and, because of the nature of the training data, is sensitive to the
presence of the GFP fusion. Explainability analysis and rational mutagenesis
disclose that expression can be determined by just a few key residues and appears
to be closely linked to co-translational membrane insertion. Overall, we show that the
integration of computational protein design and accessible machine learning tools
can provide highly accurate sequence-to-expression models of recombinant
membrane proteins.

**Methods**

*Library design*

The 18,339 original design decoys produced in Hardy et al [12] were unduplicated using the seqkit function rmdup. The resulting 12,248 unique protein sequences were backtranslated to DNA sequences with the EMBOSS tool backtranseq using the *E. coli* codon usage table for high-expressing proteins (*Eecoli_high.cut*). Universal adapters were added to the upstream end (5'-ACTTTAAGAAGGAGATATACCATG) and downstream end (GCGGCCGCACTCGAGCTGGTGCCGCGCGGCAGCA-3') of each sequence to introduce a transcriptional start codon and to facilitate cloning and amplification. All 12,248 gene sequences were ordered as a combined oligo pool from Twist Bioscience.

For diversification of the supplied oligo pool, a mixed-template PCR reaction was performed with the degenerate primers 5'-ACTTTAAGAAGGAGATATACCATG-3' (Forward) and 5'-GGCACCAGCTCGAGTG-3' (Reverse) targeting the universal adaptor sequences at either end of all pool genes. Phusion polymerase (New England Biolabs), which has a processivity of ~100nt, was used in the supplied high-fidelity buffer with 0.2 mM dNTPs, 0.5μM each primer, 3% DMSO and approximately 1ng of the oligo pool. The reaction mixture was transferred from ice to a pre-heated block at 98°C for 30s prior to 10 cycles of 98°C for 10 s, 59°C for 30s, 72°C for 15s, with a final extension at 72°C for 5 min. The 383 bp amplicon was excised from a 2% Tris-acetate-EDTA (TAE)-buffered agarose gel and recovered using the QIAgen gel extraction kit.

*Library construction*

Library cloning used restriction/ligation. The diversified oligo pool (12 ng) was digested with 10 units each of NcoI-HF and XhoI (New England Biolabs) in the manufacturer's SureCut buffer at 37°C for 1h. This digested fragment was recovered using the NEB Monarch PCR purification kit using an elution volume of 10μl. An in-house pET28-GFP plasmid [12, 25] was prepared with NcoI-HF/XhoI using 2μg plasmid and 20U of each enzyme. After 30 min at 37°C, the digested plasmid was dephosphorylated with 5U NEB Antarctic Phosphatase for 30 min at 37°C. The

digested plasmid was recovered from a 1.8% agarose TAE gel using a QIAgen gel
extraction kit, eluting in 30μl volume. DNA concentrations were determined with a
Qubit 4 instrument (ThermoFisher). 40 ng of the digested dephosphorylated vector
and 8 ng of the digested oligo pool, constituting 20:60 fmol ends or 1:3 plasmid:insert
mole ratio, were ligated overnight on ice with T4 ligase (NEB M0202). The entire
ligation mix was dialysed on the shiny side of a 25 mm MCE 0.025 μm filter paper
(Whatman VSWP02500) floating in a petri dish of deionised water for 90 mins. The
dialysed sample was recovered and 10 μl was transformed into 80 μl 10Beta
ultracompetent *Escherichia coli* cells (New England Biolabs) via electroporation
using a BioRad MicroPulser instrument with a 0.1cm cuvette on the instrument
setting 'Ec1' (1.8 kV).  After 1h outgrowth in 975 μl of the recovery broth supplied
with the competent cells, the library transformation was transferred to 100 ml LB
broth plus 50 μg/ml kanamycin (LB/Kan) for selection of transformed cells. A sample
of the transformed cells was plated onto LB/Kan agar and colony counting
determined ~80,000 CFUs from the transformation. Subsequent DNA sequencing of
selected colonies gave an approximate library size of ~27,000 unique sequences.
After overnight growth of the 100 ml broth culture the library was recovered by
plasmid miniprep and transformed into the *E. coli* expression strain BL21(DE3) C43
[26] with at least 500,000 transformants at this stage. The library was stored in 1 ml
aliquots in 15% glycerol at -80°C.
178
*Sanger DNA sequencing*
Sanger sequencing of random colonies from the cloned library was performed in a
single direction only with the commercial standard sequencing primer *petup* (5'-
ATGCGTCCGGCGTAGA-3'; EurofinsGenomics). Reads were of sufficient length to
cover the entire library insert and confirm reading frame congruence with the GFP
fusion.
185
*Cytometry analysis and cell sorting.*
For flow cytometry, 1 ml of frozen library stock was used to inoculate 100 ml of
LB/Kan in 250 ml baffled glass flask with a foam bung. Cultures were grown to A600
of 0.8 at 37°C with rotary shaking at 220 rpm before induction for 2 h with 25 mg/l
isopropyl-β-D-thiogalactopyranoside (IPTG). The media was also supplemented with

25 mg/l of the heme precursor δ-aminolevulinic acid (ALA) upon induction. A 100 μl

sample was removed from the culture and diluted into 10 ml filter-sterile PBS to give

~$10^6$ cells/ml. Cytometry was performed at the Flow Cytometry facility of the

University of Bristol using a FACSAria instrument with a 70 μm nozzle. DRAQ7 was

used to determine cell viability and side-scatter was recorded with a violet laser

(405/20). A negative control (CytbX without GFP) and a positive control (CytbX-GFP

as published [12]) were cultured alongside the library and used for comparison.

Cytometry analysis was performed with 500,000 DRAQ7-negative GFP-positive

singlet cells (~6x library coverage). Two gates were defined for the unsorted library

to incorporate approximately 8% of the total population at either tail of the

distribution. This deliberately excluded a small proportion of very bright clones that

were GFP-only cloning artefacts. 10,000 individual events were collected from each

gate. The selected cell populations were re-grown and sorting was repeated twice.

The size and diversity of the sorted libraries was determined by colony counting and

sequencing. The presumed low-expressing 'dim' and higher-expressing 'bright'

libraries constituted ~3000 and ~8000 viable colony-forming units (CFUs)

respectively after sorting.   For calculating next-generation sequencing (NGS)

coverage this was conservatively assumed to be the size of the library *i.e.* 3000 or

8000 individual sequences, although it was expected that most library members

would be represented more than once among the CFUs. The sorted libraries were

independently sequenced with an Oxford Nanopore MinION device using a rapid

sequencing kit (SQK-RAD114) and R10.4.1 flow cell (FLO-MIN114.001). Sequencing

over 21h was sufficient for >2 Gbp, giving at last 30x library coverage. Postrun

basecalling was performed with Dorado 0.7.1 in duplex mode on the University of

Bristol high-performance computing cluster BluePebble1, using the super-accurate

model dna_r10.4.1_e8.2_400bps_sup@v5.0.0 with default trimming of sequencing

adapters. Sequencing calls for the 'dim' and 'bright' library featured 11% and 16%

duplex sequences, respectively, with N50 at the full plasmid length of 6.4k. As

expected, a proportion of plasmid dimers were evident to the same extent in both

sorted libraries. After removing the first 150nt of every read, which were found to be

of lower average quality, the sequencing data were filtered to select full-length

plasmid reads with read quality ≥Q30 using seqkit [27] and fastq-filter
(https://github.com/LUMC/fastq-filter). Open reading frames for the library-GFP
genes were extracted by EMBOSS getorf [28] and truncated to remove the GFP
fusion. This resulted in 9407 library gene sequences for the 'dim' library and 7529
sequences for the 'bright' library. As expected, at least 70% of these individual gene
sequences were duplicated at least once within each library. Deduplication and
manual removal of a small number of cloning artefacts provided a final set of 1054
unique low-expressing sequences and 1122 unique high-expressing sequences. For
data analysis, FastQC and Unique.seq were accessed via the GalaxyEU webserver
(https://usegalaxy.eu) [29]. Only 4 sequences were found to be shared between the
two libraries. About one-third of the sequences in each library population came from
the original designed oligo pool, meaning that about two-thirds were diversified
sequences produced via mixed-template PCR.
237

*Bulk fluorescence measurements*
For bulk expression measurements, individual library members in strain C43 were
grown overnight at 37°C/220 rpm in LB/Kan in 10 ml glass flat-bottomed universal
vials. 1 ml of each overnight culture was inoculated into 100 ml LB/Kan in a 250 ml
baffled glass flask with a foam bung and grown on as above. After reaching an A600
of 0.7 cultures were induced with IPTG with ALA supplementation as above. After 3h
post-induction growth, cultures were diluted 17x into PBS and the GFP fluorescence
at 512 nm was determined immediately in a Cary Eclipse fluorimeter with excitation
at 490 nm.
247

*Protein purification*
In all cases expression was in strain C43. A single colony was picked from a
transformant plate into 100 ml LB/Kan in a baffled 250 ml glass flask with a foam
bung. This primary culture was grown overnight at 37°C in a shaking incubator at
220 rpm. For secondary expression cultures, 10 ml of the primary culture was used
to inoculate each of three volumes of 1 l LB/Kan in 2.5 l baffled glass flasks. This
was grown on at 37°C/220 rpm to an A600 ~0.7, at which point protein expression
was induced with 25 mg/l IPTG/ALA as above. After 3h post-induction growth the
cells were harvested by centrifugation and purification of individual proteins was

performed as described [12]. Briefly, harvested cell pellets were resuspended in total
volume 100ml PBS and lysed in a continuous flow cell disruptor at 25 KPSI.
Membrane fragments were isolated from the cell lysate at 170,000 x g and
solubilised with 1% 5-Cyclohexyl-1-Pentyl-β-D-maltopyranoside (Cymal-5) in 50 mM
sodium phosphate buffer pH 7.4, 150 mM NaCl and 5% glycerol. After removing
insoluble material via centrifugation at 170,000 x g the detergent-solubilised protein
was purified in the same buffer with 0.24% Cymal-5 on a 5 ml Ni-NTA column at a
flow rate of 3 ml/min. Sample loading was at 20 mM imidazole, column washing used
4 column volumes with 75 mM imidazole, and elution was performed with 2 column
volumes of 0.5 M imidazole at a flow rate of 1 ml/min. The sample was adjusted to
2.5 ml and immediately desalted using a CentriPure Zetadex-25 column under
gravity flow. The concentration of the purified sample was determined by absorbance
spectroscopy, using an extinction coefficient for the oxidised heme peak at 418 nm of
155,200 $M^{-1}cm^{-1}$ [12]. Cellular protein abundance was estimated assuming ~$3x10^6$
proteins per cell [30] and ~$5x10^5$ inner membrane proteins per cell [31].
272
*In-gel fluorescence*
For in-gel fluorescence, induced cultures were harvested and resuspended at 20x in
1:2 PBS:BugBuster reagent plus DNAse. After 45 min at room temperature, SDS-
PAGE gel loading buffer (ThermoFisher NP0007) was added to 1x and samples
were either used immediately or boiled at 95°C for 5 mins before loading. Sample
loading was corrected for the total lysate protein concentration determined using a
BCA assay in 2% SDS. Gel imaging used 488 nm excitation on a Typhoon
instrument (GE Healthcare).
281
*RT-qPCR*
Methods for RT-qPCR are reported according to the MIQE guidelines [32]. A total
RNA cell extract from ~$1x10^9$ *E. coli* cells was isolated from induced and uninduced
cultures with the Monarch Total RNA Miniprep kit (New England Biolabs, T2110),
incorporating lysozyme for cell lysis and DNase treatment according to the
manufacturer's instructions. The extract was immediately frozen at -80°C and stored
for <1 month, only being thawed on ice immediately prior to use. Total RNA was
determined with a Qubit 4 instrument (ThermoFisher Scientific; HS RNA, Q32852),

also used to confirm <5% DNA in each extract, and 50ng total RNA was used in each

reaction. Biological replicates of two duplicates each were performed using the same

primer pair (Forward: 5'-ATGGGTTCTCCGTGGCTG-3', Reverse: 5'-

AGACCGGTCAGGAAAACCAG-3'; EurofinsGenomics) generating an amplicon of

239 bp that differs by only two nucleotides between the two variants studied. Data

were normalized against *cysG*, *idnT* and *hcaT* as suggested by Zhou et al [33], using

the primers from their study, with all reference reactions performed in parallel to the

test genes. The reaction was performed with SYBR fluorescent probe detection

using a commercial one-step RT-qPCR kit (New England Biolabs Luna Universal kit,

E3005) according to the default cycling parameters and aliquoting scheme

recommended by the manufacturer. Reactions were 20 µl volume in 96-well plates

sealed with adhesive film (MicroAmp N8010560 and 4360954, Applied Biosystems).

Thermal cycling was 10 min at 55°C, 1 min at 95°C, 40 cycles of 10 s at 95°C and

30-60 s at 60°C, immediately followed by a melt of 60°C-95°C at 0.15 °C/s. All

pipetting was performed manually by Paul Curnow. Data were collected on a

QuantStudio3 instrument in fast mode with provided analysis software v. 1.5.3

(Applied Biosystems). Amplification specificity was determined by post-reaction

agarose gel, which included no-mRNA and no-transcriptase controls. An uninduced

culture of the low-expressing variant was used as the reference for calculating ΔΔCt,

with Ct <30 in all samples, a linear dynamic range between 3-1500 ng total RNA,

and efficiency ≈1. The baseline and threshold values were calculated by the

instrument and used without manual adjustment.

*Machine learning*

A random forest model was implemented using scikit-learn (https://scikit-learn.org)

for expression classification. The amino acid sequences were assigned binary

categorical labels according to their expression level (1=high-expressing, 0=low-

expressing), and randomly shuffled prior to training. The amino acid sequences went

through one-hot encoding and were split into training (80%) and testing (20%) sets.

The random forest model was trained with number of estimator = 100, minimal

sample split = 2 and mean sample leaf = 1. The trained model was used to infer the

expression level of 11,503 unseen protein sequences encoded by the original oligo

library. For each sequence, the model outputs the predicted probability for the

sequence to be in the bright (high-expressing) class, which was used as an indicator of predicted sequence performance.

Feature importance values were obtained directly from the random forest model based on the mean decrease in impurity during training. For each sequence position, importance scores of different amino acids were aggregated to yield a single positional importance value across the 113 positions. Additionally, Shapley additive explanations (SHAP) analysis was performed on the trained model using 1,000 of the unseen protein sequences to assess the contribution of each amino acid at each position.

*Synthesis and purification of selected library sequences*

Sequences with predicted high or low expression characteristics were obtained as individual synthetic genes from Twist Bioscience, retaining the codon usage of the original library and incorporating a His10 tag at the C-terminus. For subcloning to the GFP vector, these genes were amplified by PCR using the universal reverse primer 5'-ATGATGCTCGAGTGCGGCCG-3' and one of three forward primers as appropriate: 5'-CTTCGACCATGGGTTCTCCGATCCTGCGT-3', 5'-CTTCGACCATGGGTTCTCCGTGGCTGCGT-3' or 5'-CTTCGACCATGGGTTCTCCGATCATCCG-3'. The individual amplicons and the GFP destination vector were digested with NcoI-HF/NotI and the vector was dephosphorylated prior to ligation with a rapid ligase (New England Biolabs M2200) for 10 min at room temperature before transforming into chemically-competent TOP10 *E. coli* cells (ThermoFisher). All constructs were confirmed by sequencing.

**Results**

An overview of the experimental approach used here is shown in Figure 1. In

summary, a library of synthetic oligonucleotides was obtained for 12,248 *de novo*

sequences derived from computational protein design, with all of the designed

sequences corresponding to a diheme membrane cytochrome comprising four

transmembrane $\alpha$-helices [12]. Codon usage across the oligonucleotide pool was

standardised to the usage pattern of high-expressing *E. coli* proteins. These

sequences were further diversified by mixed-template PCR and cloned upstream of

*gfp* to allow phenotypic selection by fluorescence cytometric cell sorting. Selected

clones were sequenced, labelled according to their expression phenotype, and used

to train a predictive classifier.

*Characteristics of the design library*

The design process allowed for multiple simultaneous amino acid substitutions at 54

sites distributed over the protein surface. Residues at 13 of these sequence

positions were allowed to sample a small set of alternative residues while the

remaining 41 sites sampled from a minimal amino acid alphabet of FAILVWGST; the

Rosetta resfile from ref. [12] specifying the positional alphabet is included here as

supplementary data. Evaluation of the resulting 12,248 designed protein sequences

revealed that amino acid substitutions occurred throughout the protein (Fig. 2), with

some positions showing greater relative diversity (although with low absolute

diversity). Analysing a subset of the designed sequences showed a typical between-

sequence distance of about 12 residues (Supplementary Figure S1).

**Figure 1: Overview of the experimental approach used in this study**. Each of 12,248 computationally-designed membrane proteins were genetically encoded and synthesised commercially as an oligo pool. The combined oligo pool was controllably diversified by PCR and cloned upstream of the open reading frame for GFP. The resulting library was expressed in recombinant bacteria and after flow cytometry analysis, fluorescence-activated cell sorting (FACS) was used to isolate 'Dim' and 'Bright' cell populations. These two populations were sequenced and the corresponding amino acid sequences used to train a predictive classifier.

**Figure 2: General characteristics of the designed protein library.** (**a**) Computationally-designed amino acid substitutions are distributed throughout the diheme four-helix bundle. Some positions showing higher relative diversity, calculated as Shannon entropy. *N*, amino terminus; *TM*, transmembrane; *L*, loop. (**b**) Protein model showing the surface location of the designable positions. (**c**) Illustrative multiple sequence alignment of five sequences highlighting the controlled diversity of the library. Boxed regions correspond to TM helices 1-4 as shown.

The obtained oligo pool was then controllably diversified (shuffled) via mixed-
template PCR with a low-processivity polymerase [34] before cloning into a bacterial
expression plasmid upstream of the open reading frame for superfolder GFP. This
resulted in a fusion protein library with ~27,000 individual variants. Sequencing a
random sample of the library suggested that about 1/3 of the library sequences were
from the original oligo pool and the remainder were shuffled versions of the original
sequences.

*Phenotypic analysis and sorting*
The entire library was transformed into *E. coli* strain BL21(DE3) C43 and cultured for
membrane protein expression. Induced cultures were analysed by flow cytometry
and sorted using GFP florescence as a selectable phenotype. After three successive
rounds of selection, two populations of dim (low expression) and bright (high
expression) phenotypes were isolated from the original library with limited
fluorescence overlap between these two library subsets (Fig. 3a). Nanopore
sequencing recovered 1122 unique gene sequences (encoding 1066 unique protein
sequences) from the sorted bright library and 1054 unique gene sequences (989
unique proteins) from the sorted dim library. Only four gene sequences found in
common between the bright and dim sequence libraries. The sequence diversity
within the selected library sets was comparable to that of the unselected original
oligo pool (Supplementary Figure S1).

**Figure 3. Sorting library-GFP fusion proteins by GFP phenotype and experimental validation of selected proteins**. (**a**) Two libraries enriched in low-GFP (dim) or high-GFP (bright) expression phenotypes are selected by three rounds of fluorescence-activated cell sorting. (**b**) Bulk expression measurements of individual selected clones confirm the phenotype associated with each library population. (‡) and (*) denote the same protein across panels b-d. (**c**) Confocal fluorescent microscopy of *E. coli* cells expressing two representative sequences from the dim and bright libraries show that these proteins are localised to the cell membrane. Image contrast is optimised for display and is not representative of the true fluorescence intensity *i.e.* cells expressing the bright protein showed higher GFP under imaging. (**d**) Selected dim and bright individuals are purified by affinity chromatography with different protein yields. (**e**) Purification yields of individual library-GFP proteins correlate directly with their bulk GFP signal. (**f**) Absorbance spectroscopy of purified individual fusion proteins confirms the expected heme cofactor loading and GFP chromophore absorption. Inset photo shows the intense red colouration of the purified protein. (**g**) Size-exclusion chromatography shows that purified dim and bright fusion proteins are monodisperse and have the same apparent size. (**h**) SDS-PAGE analysis confirms the expression of full-length fusion protein. The purified fusion protein remains folded under ambient conditions and a visible coloured band is observed without staining (*Visible*). Sample boiling unfolds GFP, evidenced by a band shift and loss of in-gel fluorescence (*Fluor.*). The heme signal is retained in the visible gel since the designer proteins are thermostable. Coomassie staining (*Coomassie*) shows the high purity of the sample. Theoretical molecular weight of the fusion protein is 42 kDa.

To validate the sorted libraries, a random subset of 8 individual clones from each
library population were tested in bulk fluorescence assays, where the GFP signal
from induced cultures was measured in a fluorimeter. Single clones from the bright
library did generally show higher GFP signal than those from the dim library. The
dynamic range of these measurements was less pronounced than in cytometry,
presumably because of instrument differences and inner filter effects (Fig. 3b). Cell
imaging confirmed that the fusion proteins were localized to the cell membrane, and
that this membrane-localized protein was the phenotypic readout (Fig. 3c). Individual
clones were also purified from cell membrane fractions to confirm that the
fluorescence signal correlated directly to the yield of purifiable protein (Fig. 3d and
e). This purification used the nonionic surfactant Cymal-5, a mild detergent that does
not disaggregate misfolded protein. The yield of the selected library-GFP fusion
protein purified from the bright library was an exceptional 16 mg protein per litre of
bacterial culture – equivalent to ~70,000 copies per cell, ~2% of all cell protein or
~10% of all inner membrane protein – confirming that the library harbours ultra-high-
fitness sequences. Absorbance spectroscopy of purified samples determined that the
library proteins were fully loaded with heme (Fig. 3f) and size-exclusion
chromatography showed that proteins were not disposed to aggregation (Fig. 3g),
indicating that aggregation propensity *per se* could not explain the difference in
expression levels. Finally, gel electrophoresis confirmed the purity and integrity of the
isolated protein (Fig. 3h and Supplementary Figure S2).
454

*Predictive sequence-to-expression model*
To train predictive models of expression, amino acid sequences from the selected
dim and bright variant libraries were featurized by one-hot encoding, assigned
categorical labels of either 1 (bright) or 0 (dim) according to expression, and
employed to train a Random Forest classifier with a stratified 80:20 data split of
training and test set respectively. Model evaluation on the held-out test set showed a
high classification accuracy >0.9 and the area under the Receiver Operating
Characteristic (ROC) curve was 0.96 (Fig 4a). The model was then queried with
11,503 unlabelled sequences encoded by the original oligo pool to predict the
probability of each variant having the bright phenotype (Fig. 4b). A UMAP
representation of the prediction scores for both the training set and the unlabelled
sequences (Fig. 4c) suggests the existence of two major sequence clusters, but with

bright and dim variants distributed across the sequence space. The model
predictions did not correlate with simple biophysical metrics such as the overall
Rosetta score, sequence hydrophobicity, or the insertion $\Delta G$ of TM1 (Supplementary
Fig S3).

For experimental validation of the model predictions we selected a set of 12
unstudied variants with predicted expression scores of <0.02 (high probability of dim)
or >0.9 (high probability of bright) and obtained the corresponding genes as
individual clones. The observed expression level of these proteins was consistent
with the classifier prediction in all cases (Fig. 4d).

To determine the sensitivity of these variants to the phenotypic marker, we replaced
GFP with a short His10 affinity tag and purified each of these His-tagged proteins by
affinity chromatography (Fig 4e). The His-tagged proteins were expressed at
significantly lower levels overall and without the expression differences of the
equivalent GFP fusions. This result reflects the nature of the training set, which is
phenotyped on the full-length fusion protein, and highlights the influence that soluble
domains can exert on membrane protein biosynthesis. The difference between
GFP/His constructs probably indicates the loss of co-translational insertion
mechanisms when the C-terminus is truncated by removal of the GFP [35, 36], and
correlates with our own observation that the expression of these *de novo* membrane
proteins is enhanced by an extended soluble domain at the end of the last
transmembrane helix [37].

**Figure 4: A machine learning model for membrane protein expression**. (**a**) Sequence libraries corresponding to either the bright or dim phenotype are labelled discretely, one-hot encoded and used to train a random forest classifier. Testing with 20% of the dataset excluded from training reveals excellent classifier performance. (**b**) The trained model was used to predict the expression phenotype of 11,503 unseen and unlabelled protein sequences encoded by the original oligo pool. (**c**) Dimensionality reduction by UMAP shows two major clusters in the data but no distinct clustering of high- and low-expressing sequences. (**d**) Bulk fluorescence measurements provide experimental confirmation of the phenotypic predictions made by the model. Control samples are from Fig. 1b-d. (**e**) The expression yield and expression profile are sensitive to the C-terminal domain.

*Expression engineering via explainability analysis*

We next used feature importance analysis to identify residues that strongly
contribute to expression. Sequence positions 12 and 113 were found to be
prominently affecting the model predictions (Fig 5a). Analysis of these positions with
the Shapley additive explanations (SHAP) algorithm for model explainability revealed
that position 12 was either F/I/L with I12 being strongly associated with high-
expression prediction by our model (Fig 5b). At position 113, library sequences were
almost exclusively K or Q and almost all high-expressing sequences featured K113
(Fig. 5b). Much weaker preferences were observed in high-expressing sequences for
I/L6, I/F11, I/L19, W/F76, Y91, and I98; low-expressing sequences were marginally
enriched in F92, L95 and N/R112.

We reasoned that the predominance of K113 in high-expressing sequences might
reflect a specific interaction of the Lys sidechain with the GFP fusion, perhaps
allowing the GFP to act as a folding chaperone. To probe this further we purified one
of the high-expressing variants in the presence and absence of GFP and directly
compared the size-exclusion profile of these two constructs (Supplementary Fig. S4).
While the expression yields were dramatically different - being about 30x higher in
the presence of GFP – there was no general increase in the aggregation propensity
or heme binding of the His-tagged protein. Thus while the presence of the GFP
fusion might support folding in the cell, it is not required to maintain the fold of
purified library proteins. This conclusion was supported by molecular dynamics
simulations, which found that GFP was highly mobile relative to CytbX due to the
flexibility of the connecting linker. These simulations could not identify substantive
interactions between the two fused proteins but did find salt bridges between Lys113
and lipid headgroups (Supplementary Figure S5).

We directly tested the feature importance scores from Fig. 5a-b by investigating two
variants with opposite expression outcomes but with very similar sequences,
differing by only 7/113 amino acids and 13/339 nucleotides (Fig.5c; Supplementary
Fig S6). Both sequences featured Ile at position 12 but were different for K/Q at
position 113. Culture growth curves showed that recombinant strains carrying the
low-expressing variant (termed PredLow) exhibited a more rapid and pronounced
growth arrest after induction versus the high-expressing variant PredHigh (Fig. 5d).

We hypothesised that if position 113 was a critical determinant of expression, then

we should be able to substantially improve the expression of PredLow by introducing

mutation Q113K (CAG→AAA). This approach was successful, with the rationally

engineered variant PredLow$^{Q113K}$ showing minimal growth arrest (Fig. 5d), high bulk

fluorescence (Fig. 5e) and an 8-fold increase in purification yield (Fig. 5e).

Finally, to dissect the possible mechanistic reasons for the observed expression

disparity we used RT-qPCR to assess the abundance of PredLow and PredHigh

transcript mRNA in recombinant cells. PredHigh mRNA was found to be present at

levels ~10x that of PredLow (Fig. 5f and Supplementary Figure S7). Transcript

abundance within the cell is dictated by a balance of transcription and mRNA decay.

We assume that transcription rates are generally similar between these constructs

given the close similarity of the two gene sequences. This suggests that PredLow

mRNA is more efficiently degraded in the cell. Notably, high transcript abundance is

recovered in PredLow$^{Q113K}$ (Fig. 5f) and thus appears to be a hallmark of highly-

expressing membrane proteins.

552
**Figure 5: Variant optimization with explainability analysis.** (**a**) The classifier
identifies two sequence positions, 12 and 113, as being the most important features
for determining expression. (**b**) Residue-level SHAP analysis shows that Ile at
position 12 and Lys at position 113 are associated with high-expressing sequences;
in contrast, Gln 113 is strongly associated with sequences that have low predicted
expression. (**c**) Overlay of the AlphaFold structures of two variants that differ by only
the 7 amino acids shown but have strongly High or Low predicted expression
outcomes, termed *PredHigh* and *PredLow* respectively. (**d**) Growth curves of *E. coli*
strains producing PredHigh or PredLow. Cells carrying PredLow show a more
pronounced growth arrest after induction. This metabolic burden is completely
relieved by a single mutation Q113K. (**e**) Bulk fluorescence measurements confirm
expression recovery in the engineered variant PredLow[Q113K]. Inset photograph of
purified protein confirms the associated difference in purification yields as shown. (**f**)
RT-qPCR shows an association between protein expression and transcript
abundance. For additional controls and efficiency curve see Fig. S7.

**Discussion**

The results presented here show that the fitness landscape of an integral membrane protein can be successfully described by tractable and explainable machine learning tools. We exploit a novel protein library where multiple simultaneous mutations are introduced computationally at designated surface sites throughout a small *de novo* protein, using a restricted amino acid alphabet and biased by a membrane-specific energy function (Fig. 2). The resulting membrane cytochromes can be fused to a phenotypic marker, are localized to the *E. coli* inner membrane, are straightforward to purify and characterise, and retain a strong heme absorbance signal that reports directly on successful protein folding in the membrane (Fig. 3). The computational search of a structured sequence space provides an alternative to library generation by systematic mutational scanning and random mutagenesis and to the greater diversity of natural protein collections.

Our results confirm that such computationally-generated protein libraries are ideal for machine learning and offer a suitable trade-off between feature diversity and library size (Fig 4). The sequence data are directly compatible with standard off-the-shelf bioinformatics packages and the analysis is of low computational expense, with all classifier training, prediction and feature analysis running in less than 3 minutes on a standard laptop computer. Collections of natural membrane proteins show no obvious link between specific sequence features and cellular protein expression [8, 20]. We show that, for our computational library, assumption-free and mechanism-agnostic sequence representations can reveal non-obvious features that exert strong control over membrane protein expression but are not necessarily captured by biophysical encoding schemes (Fig. 5). While encodings such as the one-hot representation used here can provide excellent model performance on the specific dataset being studied – often surpassing more intuitive biophysical representations of protein sequence [38, 39] – they do not generalize well [40]. Further work will explore whether hybrid models incorporating mechanistic features will allow extrapolation from our design library to other datasets.

We show that membrane protein expression can be strongly influenced by fusion tags (Fig. 4d-e) and can vary at least 8-fold based on just one or two key residues (Fig 5). In particular, the ultra-high expression of library variants depends upon the

presence of Lys at position 113, sited at the end of the last transmembrane helix.
There do not appear to be persistent contacts between K113 and the soluble GFP
domain. Instead, we propose that K113 is important for specifying protein folding in
the cell, and perhaps dictates co-translational membrane insertion and topology
according to the 'positive-inside rule' [4, 41-43]. This aligns with previous work that
has demonstrated the importance of insertion efficiency on protein expression yield,
with the length and composition of the protein C-terminus found to be particularly
important [44, 45].

How could these differences in insertion efficiency be linked to mRNA abundance
(Fig. 5f)? We assume that, like most integral membrane proteins, the library variants
are synthesised cotranslationally via a ribosome-translocon complex [3, 4]. This
means that membrane insertion is directly coupled to transcript utilization by the
ribosome. The cellular abundance of a particular transcript depends upon the rate at
which it is transcribed and the rate at which it is degraded. Since transcription ought
to be at a relatively constant and high level for these recombinant genes, we suggest
that mRNA decay is responsible for the differences in transcript abundance. The
cellular turnover of mRNA is not completely understood, but it is clear that several
pathways exist in *E. coli* to degrade poorly-translated mRNA [46-48]. We speculate
that slow membrane insertion of Q113 variants, perhaps caused by weak topological
definition and inefficient helix packing within the membrane, could stall translation
and lead to ribosome pausing and collisions. This would activate rescue pathways
leading to mRNA decay [47-52] and lead to the premature termination of translation;
the associated cellular stress may contribute to the observed growth arrest (Fig. 5d).
In contrast, more efficient or 'low-friction' membrane insertion and folding of K113
variants would allow unhindered translational elongation and avoid quality-related
transcript decay. Under such a model, rapidly-translated transcripts would be
completely translated with greater frequency and would be recycled to the cellular
mRNA pool, potentially available for multiple rounds of translation. These ideas are
summarised in Fig. 6. The rate of cotranslational membrane helix insertion is
probably slightly faster than the speed of translational elongation [3, 53, 54]; this
implies that even a subtle slowdown to the insertion kinetics may be enough to
cause a translational logjam. We note that multiple other factors might explain the
differential decay of transcript mRNA including secondary structure and localization

[46, 47], but it is challenging to apply these ideas to the transcripts studied here
which are very close in sequence but have radically different cellular fates.

**638**

**639**    **Figure 6. Speculative expression model integrating membrane protein folding**

**640**    **with mRNA decay.** (**a**) A library variant carrying a positive charge at the end of TM4

**641**    has strong topological definition according to the positive-inside rule. Once

**642**    translation is complete the mRNA is released back to the transcript pool. (**b**) A library

**643**    variant without a charged residue in TM4 is slower to adopt the topology required for

**644**    folding of the 4-helix bundle. This 'logjam' causes ribosome stalling or collision and

**645**    leads to mRNA decay. This means that fewer of these transcripts are completely

**646**    translated and the mRNA is less frequently recycled back to the cellular pool. Other

**647**    factors associated with low expression yields e.g. inclusion body formation are not

**648**    considered                                                                                              here.

We also find that an earlier codon corresponding to Ile at position 12 in TM helix 1 is
favoured in, but not exclusively limited to, high-expressing sequences (Fig. 5a).
Codon usage, nucleotide composition, amino acid identity and mRNA structure
within the early part of the transcript can all be important for prokaryotic gene
expression [14, 55, 56], including for membrane proteins [21, 57]. However,
whatever positive impact I12 may have on protein expression is evidently secondary
to the effect of K113 (Fig. 5). The bias towards I12 in high-expressing sequences
remains to be fully understood.

Overall, we have established a new membrane protein expression dataset that will
serve as a useful testbed for the development of specific and general sequence-to-
expression algorithms. The expression phenotype of these proteins can be
interpreted using classical pathways of membrane protein biogenesis, and ultra-high-
expression phenotypes may result from successfully matching the kinetics of
membrane protein insertion and folding to the rate of translational elongation. This
concept implies that tuning the integration kinetics might be a fruitful route to
engineering membrane proteins for desirable expression characteristics. Extending
the computational mutagenesis method developed here to a range of other
membrane proteins, and to residues within the protein core, should enable the
mapping of both local and global sequence parameters that control recombinant
expression. Our results also raise the prospect of applying computational resurfacing
to natural proteins to achieve enhanced recombinant expression while retaining
native structure and function.

**Data availability**
The sequence data and code used in this study is available online at
https://github.com/Curnow-Lab-University-of-Bristol/Membrane-protein-library-ML.

**Conflict-of-interest statement**
The oligo pool used here was obtained from Twist Bioscience at a heavily reduced
price as part of an academic pre-market trial for an enhanced synthesis method. The
authors have no affiliation with, or financial interest in, Twist Bioscience. Twist
Bioscience was not involved in any other aspect of the research described here or in
the preparation of this manuscript.

**Acknowledgements**

PC & AJM were supported by the UK Biotechnology and Biological Sciences Research Council (BBSRC) grant number BB/W003449/1. YS was supported by BBSRC grant number BB/T00875X/1. JU was supported by Engineering and Physical Sciences Research Council Doctoral Training Partnership EP/W524414/1. This research used the Flow Cytometry Facility and the computational facilities of the Advanced Computing Research Centre of the University of Bristol (http://www.bristol.ac.uk/acrc/). Thanks to Ben Hardy and Ross Anderson for helpful discussions.

**Author contributions**

Conceptualization: PC, DAO. Funding acquisition: PC, DAO, AJM. Investigation: YS, JU, PC. Methodology: YS, JU. Resources: PC, DAO, AJM. Supervision: PC, DAO, AJM. Visualization: YS, JU, PC. Writing – original draft: PC. Writing – review and editing: YS, AJM, DAO, PC. All authors have read and approved the final manuscript.

**References**

1. Love, J., F. Mancia, L. Shapiro, M. Punta, B. Rost, M. Girvin, et al., *The New York Consortium on Membrane Protein Structure (NYCOMPS): a high-throughput platform for structural genomics of integral membrane proteins.* J Struct Funct Genomics, 2010. **11**(3): p. 191-9.
2. Hammon, J., D.V. Palanivelu, J. Chen, C. Patel, and D.L. Minor, Jr., *A green fluorescent protein screen for identification of well-expressed membrane proteins from a cohort of extremophilic organisms.* Protein Sci, 2009. **18**(1): p. 121-33.
3. Mercier, E., X. Wang, L.A.K. Bogeholz, W. Wintermeyer, and M.V. Rodnina, *Cotranslational Biogenesis of Membrane Proteins in Bacteria.* Front Mol Biosci, 2022. **9**: p. 871121.
4. Luirink, J., G. von Heijne, E. Houben, and J.W. de Gier, *Biogenesis of inner membrane proteins in Escherichia coli.* Annu Rev Microbiol, 2005. **59**: p. 329-55.
5. Hattab, G., D.E. Warschawski, K. Moncoq, and B. Miroux, *Escherichia coli as host for membrane protein structure determination: a global analysis.* Sci Rep, 2015. **5**: p. 12097.
6. Nikolados, E.M. and D.A. Oyarzun, *Deep learning for optimization of protein expression.* Curr Opin Biotechnol, 2023. **81**: p. 102941.
7. Baranowski, C., M. Hector Garcia, D.A. Oyarzun, E.M. Nikolados, S. Jaaks-Kraatz, A. Gaber, et al., *Can protein expression be 'solved'?* 2024.
8. Saladi, S.M., N. Javed, A. Müller, and W.M. Clemons Jr, *A statistical model for improved membrane protein expression using sequence-derived features.* J Biol Chem, 2018. **293**(13): p. 4913-4927.

9.    Kuntz, C.P., H. Woods, A.G. McKee, N.B. Zelt, J.L. Mendenhall, J. Meiler, et al.,
*Towards generalizable predictions for G protein-coupled receptor variant expression.*
Biophys J, 2022. **121**(14): p. 2712-2720.

10.   Bedbrook, C.N., K.K. Yang, A.J. Rice, V. Gradinaru, and F.H. Arnold, *Machine learning*
*to design integral membrane channelrhodopsins for efficient eukaryotic expression*
*and plasma membrane localization.* PLoS Comput Biol, 2017. **13**(10): p. e1005786.

11.   Naghipourfar, M., S. Chen, M.K. Howard, C.B. Macdonald, A. Saberi, T. Hagen, et al., *A*
*Suite of Foundation Models Captures the Contextual Interplay Between Codons.*
bioRxiv, 2024.

12.   Hardy, B.J., A. Martin Hermosilla, D.K. Chinthapalli, C.V. Robinson, J.L.R. Anderson,
and P. Curnow, *Cellular production of a de novo membrane cytochrome.* Proc Natl
Acad Sci U S A, 2023. **120**(16): p. e2300137120.

13.   Hutchins, G.H., C.E.M. Noble, H.A. Bunzel, C. Williams, P. Dubiel, S.K.N. Yadav, et al.,
*An expandable, modular de novo protein platform for precision redox engineering.*
Proc Natl Acad Sci U S A, 2023. **120**(31): p. e2306046120.

14.   Cambray, G., J.C. Guimaraes, and A.P. Arkin, *Evaluation of 244,000 synthetic*
*sequences reveals design principles to optimize translation in Escherichia coli.* Nat
Biotechnol, 2018. **36**(10): p. 1005-1015.

15.   Drew, D.E., G. von Heijne, P. Nordlund, and J.W. de Gier, *Green fluorescent protein as*
*an indicator to monitor membrane protein overexpression in Escherichia coli.* FEBS
Lett, 2001. **507**(2): p. 220-4.

16.   Geertsma, E.R., M. Groeneveld, D.J. Slotboom, and B. Poolman, *Quality control of*
*overexpressed membrane proteins.* Proc Natl Acad Sci U S A, 2008. **105**(15): p. 5722-
7.

17.   Drew, D., S. Newstead, Y. Sonoda, H. Kim, G. von Heijne, and S. Iwata, *GFP-based*
*optimization scheme for the overexpression and purification of eukaryotic membrane*
*proteins in Saccharomyces cerevisiae.* Nat Protoc, 2008. **3**(5): p. 784-98.

18.   Goehring, A., C.H. Lee, K.H. Wang, J.C. Michel, D.P. Claxton, I. Baconguis, et al.,
*Screening and large-scale expression of membrane proteins in mammalian cells for*
*structural studies.* Nat Protoc, 2014. **9**(11): p. 2574-85.

19.   Kermani, A.A., *Applications of fluorescent protein tagging in structural studies of*
*membrane proteins.* FEBS J, 2024. **291**(13): p. 2719-2732.

20.   Daley, D.O., M. Rapp, E. Granseth, K. Melén, D. Drew, and G. von Heijne, *Global*
*topology analysis of the Escherichia coli inner membrane proteome.* Science, 2005.
**308**: p. 1321-1323.

21.   Fluman, N., S. Navon, E. Bibi, and Y. Pilpel, *mRNA-programmed translation pauses in*
*the targeting of E. coli membrane proteins.* Elife, 2014. **3**.

22.   Hsieh, J.M., G.M. Besserer, M.G. Madej, H.Q. Bui, S. Kwon, and J. Abramson, *Bridging*
*the gap: a GFP-based strategy for overexpression and purification of membrane*
*proteins with intra and extracellular C-termini.* Protein Sci, 2010. **19**(4): p. 868-80.

23.   Mathieu, K., W. Javed, S. Vallet, C. Lesterlin, M.P. Candusso, F. Ding, et al.,
*Functionality of membrane proteins overexpressed and purified from E. coli is highly*
*dependent upon the strain.* Sci Rep, 2019. **9**(1): p. 2654.

24.   Bill, R.M., P.J. Henderson, S. Iwata, E.R. Kunji, H. Michel, R. Neutze, et al., *Overcoming*
*barriers to membrane protein structure determination.* Nat Biotechnol, 2011. **29**(4):
p. 335-40.

25.    Lalaurie, C.J., V. Dufour, A. Meletiou, S. Ratcliffe, A. Harland, O. Wilson, et al., *The de*
*novo design of a biocompatible and functional integral membrane protein using*
*minimal sequence complexity.* Sci Rep, 2018. **8**: p. 14564.

26.    Miroux, B. and J.E. Walker, *Over-production of proteins in Escherichia coli: mutant*
*hosts that allow synthesis of some membrane proteins and globular proteins at high*
*levels.* J Mol Biol, 1996. **260**(3): p. 289-98.

27.    Shen, W., S. Le, Y. Li, and F. Hu, *SeqKit: A Cross-Platform and Ultrafast Toolkit for*
*FASTA/Q File Manipulation.* PLoS One, 2016. **11**(10): p. e0163962.

28.    Rice, P., I. Longden, and A. Bleasby, *EMBOSS: the European Molecular Biology Open*
*Software Suite.* Trends Genet, 2000. **16**(6): p. 276-7.

29.    The_Galaxy_Community_2022, *The galaxy platform for accessible, reproducible and*
*collaborative biomedical analyses: 2022 update.* Nucleic Acids Research, 2022.
**50**(15): p. 8999.

30.    Milo, R., *What is the total number of protein molecules per cell volume? A call to*
*rethink some published values.* Bioessays, 2013. **35**(12): p. 1050-5.

31.    Facey, S.J. and A. Kuhn, *Membrane integration of E. coli model membrane proteins.*
Biochim Biophys Acta, 2004. **1694**(1-3): p. 55-66.

32.    Bustin, S.A., V. Benes, J.A. Garson, J. Hellemans, J. Huggett, M. Kubista, et al., *The*
*MIQE guidelines: minimum information for publication of quantitative real-time PCR*
*experiments.* Clin Chem, 2009. **55**(4): p. 611-22.

33.    Zhou, K., L. Zhou, Q. Lim, R. Zou, G. Stephanopoulos, and H.P. Too, *Novel reference*
*genes for quantifying transcriptional responses of Escherichia coli to protein*
*overexpression by quantitative PCR.* BMC Mol Biol, 2011. **12**: p. 18.

34.    Kalle, E., M. Kubista, and C. Rensing, *Multi-template polymerase chain reaction.*
Biomol Detect Quantif, 2014. **2**: p. 11-29.

35.    Kalinin, I.A., H. Peled-Zehavi, A.B.D. Barshap, S.A. Tamari, Y. Weiss, R. Nevo, et al.,
*Features of membrane protein sequence direct post-translational insertion.* Nat
Commun, 2024. **15**(1): p. 10198.

36.    Sun, S. and M. Mariappan, *C-terminal tail length guides insertion and assembly of*
*membrane proteins.* J Biol Chem, 2020. **295**(46): p. 15498-15510.

37.    Hardy, B.J., H.C. Ford, M. Rudin, J.L.R. Anderson, and P. Curnow, *Polymer nanodiscs*
*support the functional extraction of an artificial transmembrane cytochrome.* BBA:
Biomembranes, 2025. **1867**(1): p. 184392.

38.    Raimondi, D., G. Orlando, W.F. Vranken, and Y. Moreau, *Exploring the limitations of*
*biophysical propensity scales coupled with machine learning for protein sequence*
*analysis.* Sci Rep, 2019. **9**(1): p. 16932.

39.    Wittmann, B.J., Y. Yue, and F.H. Arnold, *Informed training set design enables efficient*
*machine learning-assisted directed protein evolution.* Cell Syst, 2021. **12**(11): p. 1026-
1045 e7.

40.    Shen, Y., G. Kudla, and D.A. Oyarzun, *Improving the generalization of protein*
*expression models with mechanistic sequence information.* Nucleic Acids Res, 2025.
**53**(3).

41.    von Heijne, G., *The distribution of positively charged residues in bacterial inner*
*membrane proteins correlates with the trans-membrane topology.* EMBO J, 1986.
**5**(11): p. 3021-7.

42.   Seppala, S., J.S. Slusky, P. Lloris-Garcera, M. Rapp, and G. von Heijne, *Control of*
*membrane protein topology by a single C-terminal residue.* Science, 2010. **328**(5986):
p. 1698-700.
43.   Lerch-Bader, M., C. Lundin, H. Kim, I. Nilsson, and G. von Heijne, *Contribution of*
*positively charged flanking residues to the insertion of transmembrane helices into*
*the endoplasmic reticulum.* PNAS, 2008. **105**(11): p. 4127-32.
44.   Marshall, S.S., M.J.M. Niesen, A. Muller, K. Tiemann, S.M. Saladi, R.P. Galimidi, et al.,
*A Link between Integral Membrane Protein Expression and Simulated Integration*
*Efficiency.* Cell Rep, 2016. **16**(8): p. 2169-2177.
45.   Niesen, M.J.M., S.S. Marshall, T.F. Miller, 3rd, and W.M. Clemons, Jr., *Improving*
*membrane protein expression by optimizing integration efficiency.* J Biol Chem, 2017.
**292**(47): p. 19537-19545.
46.   Laalami, S., L. Zig, and H. Putzer, *Initiation of mRNA decay in bacteria.* Cell Mol Life
Sci, 2014. **71**(10): p. 1799-828.
47.   Mohanty, B.K. and S.R. Kushner, *Regulation of mRNA Decay in Bacteria.* Annu Rev
Microbiol, 2016. **70**: p. 25-44.
48.   Roux, C., T.A. Etienne, E. Hajnsdorf, D. Ropers, A.J. Carpousis, M. Cocaign-Bousquet,
et al., *The essential role of mRNA degradation in understanding and engineering E.*
*coli metabolism.* Biotechnol Adv, 2022. **54**: p. 107805.
49.   Dreyfus, M., *Killer and protective ribosomes.* Prog Mol Biol Transl Sci, 2009. **85**: p.
423-66.
50.   Hayes, C.S. and R.T. Sauer, *Cleavage of the A site mRNA codon during ribosome*
*pausing provides a mechanism for translational quality control.* Mol Cell, 2003. **12**(4):
p. 903-11.
51.   Sunohara, T., K. Jojima, H. Tagami, T. Inada, and H. Aiba, *Ribosome stalling during*
*translation elongation induces cleavage of mRNA being translated in Escherichia coli.*
J Biol Chem, 2004. **279**(15): p. 15368-75.
52.   Saito, K., H. Kratzat, A. Campbell, R. Buschauer, A.M. Burroughs, O. Berninghausen, et
al., *Ribosome collisions induce mRNA cleavage and ribosome rescue in bacteria.*
Nature, 2022. **603**(7901): p. 503-508.
53.   Min, D., *Folding speeds of helical membrane proteins.* Biochem Soc Trans, 2024.
**52**(1): p. 491-501.
54.   Mercier, E., W. Wintermeyer, and M.V. Rodnina, *Co-translational insertion and*
*topogenesis of bacterial membrane proteins monitored in real time.* EMBO J, 2020.
**39**(15): p. e104054.
55.   Tuller, T. and H. Zur, *Multiple roles of the coding sequence 5' end in gene expression*
*regulation.* Nucleic Acids Res, 2015. **43**(1): p. 13-28.
56.   Allert, M., J.C. Cox, and H.W. Hellinga, *Multifactorial determinants of protein*
*expression in prokaryotic open reading frames.* J Mol Biol, 2010. **402**(5): p. 905-18.
57.   Norholm, M.H., S. Toddo, M.T. Virkki, S. Light, G. von Heijne, and D.O. Daley,
*Improved production of membrane proteins in Escherichia coli by selective codon*
*substitutions.* FEBS Lett, 2013. **587**(15): p. 2352-8.
856