Title

Rats spontaneously show categorical responses toward familiar or unfamiliar conspecifics in a habituation-dishabituation task using multiple habituation stimuli

Authors

Shiomi Hakataya[1,2,3,4], Makiko Kamijo[1,5], Genta Toya[6,2,7], Hiroki Koda[1,8] and Kazuo Okanoya[2,1]*

Affiliations

[1] Graduate School of Arts and Sciences, The University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo, 153-8902, Japan

[2] Advanced Comprehensive Research Organization, Teikyo University, 2-21-1 Kaga, Itabashi-ku, Tokyo, 173-0003, Japan

[3] Japan Society for the Promotion of Science, Kojimachi Business Center, 5-3-1 Kojimachi, Chiyoda-ku, Tokyo, 102-0083, Japan

[4] Faculty of Science, University of the Ryukyus, 1 Sembaru, Nishihara-cho, Nakagami-gun, Okinawa, 903-0213, Japan

[5] Faculty of Liberal Arts, Teikyo University, 359 Otsuka, Hachioji-shi, Tokyo, 192-0395, Japan

[6] Research Center for Advanced Science and Technology, The University of Tokyo, 4-6-1 Komaba, Meguro-ku, Tokyo, Japan

[7] School of Environment and Society, Institute of Science Tokyo, 2-12-1 Ookayama, Meguro-ku, Tokyo, 152-8552, Japan

[8] Center for Evolutionary Cognitive Science, The University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo, 153-8902, Japan

* Corresponding author

Kazuo Okanoya

Email: kazuookanoya@gmail.com

Address: Advanced Comprehensive Research Organization, Teikyo University, 2-21-1 Kaga, Itabashi-ku, Tokyo, 173-0003, Japan

Phone number: +81-3-3964-1211

Highlights

- We tested familiarity discrimination in rats by habituation-dishabituation task
- Multiple cage-mate or non-cage-mate stimuli were presented during habituation
- Exploratory responses decreased as habituation sessions progressed
- Dishabituation occurred in individuals who exhibited sufficient habituation
- Rats showed categorical responses toward familiarity or unfamiliarity

Abstract

Discriminating between familiar and unfamiliar conspecifics is a fundamental aspect of social recognition. Previous studies have shown that familiarity strongly influences rat social behavior, and that rats can be trained to discriminate between olfactory cues from cage-mates and non-cage-mates in a digging task. However, such discrimination did not occur in a lever-pressing task using live conspecifics as stimuli. In the present study, we investigated whether rats could discriminate social familiarity in a simple, less-demanding and naturalistic context using whole-animal stimuli. We employed a habituation-dishabituation paradigm in which multiple cage-mate or non-cage-mate individuals were presented during the habituation phase. We examined whether rats would habituate to the familiarity category among these multiple stimuli and subsequently show dishabituation when the stimulus class switched (cage-mate to non-cage-mate or vice versa). During the habituation phase, exploratory responses declined over successive sessions, suggesting habituation to the common element among multiple cage-mates or non-cage-mates, i.e., familiarity/unfamiliarity. When all subjects were considered together, clear dishabituation was not observed, indicating individual variation in habituation. However, a post-hoc analysis restricted to subjects showing sufficient habituation revealed significant dishabituation to the stimulus class switch. These findings suggest that rats can spontaneously classify multiple conspecifics according to their familiarity without training. Rats may flexibly employ different forms of social recognition and information such as individual identity or familiarity-based heuristics to guide adaptive social behavior in varying contexts.

Keywords
Social familiarity, Rat, Habituation-dishabituation paradigm, Social recognition, Cage-mate

1. Introduction

In social recognition, animals recognize sets of individuals (class-level cognition) or each individual separately (individually recognition) (Wiley et al., 2013; Yorzinski, 2017). The ability to discriminate between familiar and unfamiliar conspecifics known as social familiarity discrimination is an essential form of class-level recognition. Because it is cognitively less demanding than remembering and recognizing each conspecific individually, social familiarity discrimination can be viewed as an adaptive heuristic for managing the complexity of the social environment (Coulon et al., 2011; Jones et al., 2014; Suwandschieff et al., 2023; Wilkinson et al., 2010). Experimental evidence shows some animals can be trained to classify images of conspecifics according to their familiarity (cattle: Coulon et al., 2011; pigeon: Wilkinson et al., 2010).

Laboratory rats can be a good model for studying social familiarity discrimination. Familiarity with a partner can modulate various social behaviors in rats (e.g., Alberts & Galef, 1973; Hackenberg et al., 2021; Kiyokawa et al., 2014; Rogers-Carter et al., 2018; Smith et al., 2015). Jones et al. (2014) demonstrated that rats can be trained to discriminate between olfactory cues from familiar cage-mates and unfamiliar non-cage-mates in a digging task, which uses natural digging behavior into media containing odors as response. However, rats failed to reach the learning criterion in a lever-pressing task using live conspecific stimuli. While learning tasks such as digging can clearly demonstrate stimulus discrimination, they are often time-consuming. Also, the collection and preparation of suitable odor samples require careful handling. Development of a simpler task could broaden opportunities for future research (e.g., Wada et al., 2025), including investigations into mechanisms underlying familiarity discrimination. We therefore aimed to test whether social familiarity discrimination could be achieved in a simpler and semi-natural task without specific training using whole-animal stimuli.

The habituation-dishabituation paradigm is widely used to assess olfactory discrimination and social recognition in rodents (Slotnick et al., 2004). In this task, repeated exposure to stimuli from the same class typically induces a gradual decline in behavioral responses during habituation. A subsequent recovery of responses to stimuli from a different class during dishabituation is generally interpreted as evidence of discrimination. However, if only a single stimulus is repeatedly presented during habituation, it remains possible that the recovered responses during dishabituation reflect discrimination between two individual stimuli rather than between two stimulus classes. Presenting multiple stimuli from the same class during habituation allows assessment of discrimination based on shared features across stimuli (Kondo et al., 2005). Accordingly, in our paradigm, we presented multiple cage-mates or non-cage-mates during the habituation phase to examine whether rats would show habituation and dishabituation to the common factor of familiarity versus unfamiliarity.

In this study, we presented cage-mate and non-cage-mate stimuli and measured the
duration of subjects' social exploration. In the cage-mate habituation condition, animals were
presented with multiple cage-mate stimuli during habituation and non-cage-mates during
dishabituation; vice versa for the non-cage-mate habituation condition. We predicted that rats
would habituate to the common factor among cage-mates or non-cage-mates (i.e., familiarity or
unfamiliarity), and would show dishabituation when the stimulus class switched to the other one.
107

2. Materials and Methods
More details are provided in Supplementary Methods.
2.1. Ethical statements
All procedures were conducted in accordance with the experimental implementation
regulations of the University of Tokyo and were approved by the Animal Experimental Committee
at the University of Tokyo, Graduate School of Arts and Sciences (Permission Number: 2022-5).
2.2. Animals
Subjects were 32 male Sprague-Dawley rats (SLC Japan Inc., Hamamatsu, Japan).
Individuals housed in the same cage (four per cage) were designated as cage-mates. Experiments
were conducted 5 weeks after the start of co-housing, when the rats were 17 weeks old. Prior to
this study, all rats experienced handling, a series of behavioral tests and group tracking in a large
cage. Because none of these prior procedures resembled the experimental settings or protocols
used here, they were unlikely to have affected the results.
Subjects were kept in a room under a 12:12 h light/dark cycle (lights on at 07:00). All
experiments were conducted during the light period. Each group received 125 g of standard
laboratory chew daily (Lab Diet, PMI Nutrition International, St. Louis, USA). Water was freely
available.
2.3. Apparatus
Experiments were conducted in a custom-made open field (80 × 80 × 50 cm). Stimulus
animals were presented in black acrylic cages (18 cm length × 25.5 cm width × 52 cm height).
The lower half of the front side of each cage was covered with 5 × 5 mm acrylic rods, allowing
interactions between the subject and the stimulus animal. The stimulus cage was placed
centrally along the back wall of the arena (**Figure 1a**). A camera (HERO9 BLACK, GoPro) was
mounted above the arena for video recording.
2.4. Experimental procedures
The habituation-dishabituation task included the two conditions: the cage-mate
habituation (CMH) and the non-cage-mate habituation (NCMH) condition. Each subject was
randomly assigned to one of these conditions. In the CMH condition, animals were presented
with cage-mate stimuli during habituation and non-cage-mates during dishabituation; vice versa

in the NCMH condition. In both conditions, the cage-mate stimuli were three individuals from
the subject's own group, and the non-cage-mate stimuli were three individuals from a different
group. All individuals served as both subjects and stimulus animals.

The behavioral task consisted of an arena acclimation phase followed by stimulus
presentation (6 habituation sessions and 3 dishabituation sessions) (**Figure 1b**). During arena
acclimation, subjects were introduced to the arena and allowed to explore freely for 10 minutes.
In each stimulus presentation session, a cage-mate or non-cage-mate stimulus was presented
for 3 minutes. The inter-session interval was 5 minutes, during which the apparatus was cleaned
using 75% ethanol.

In habituation sessions, the three cage-mate or non-cage-mate stimuli were each
presented twice in a pseudo-random order, ensuring that the same stimulus was not presented
in two consecutive sessions. In dishabituation sessions, the three cage-mate or non-cage-mate
stimuli were each presented once in a random order. The testing order of subjects was
randomized.

2.5. Analysis

Exploratory responses to stimuli were quantified as the total time the subject's nose
remained on the front of the stimulus cage or within 1.5 cm of it (nose length: approximately 1.5
cm). This was measured using DeepLabCut 2.3 (Mathis et al., 2018; Nath et al., 2019). Due to
mechanical troubles, all sessions for two individuals and four habituation sessions from four
different individuals were excluded from analysis.

We analyzed the data by fitting generalized linear mixed models (GLMMs) in R 4.5.0 (R
Core Team, 2025; glmmTMB: Brooks et al., 2017). All models included the duration of responses
as the response variable, with session, condition and their interaction as fixed effects, and
subject ID as a random effect.

3. Results

We analyzed whether exploratory responses changed during habituation sessions
using GLMM (**Figure 2a**). The parameter coefficient of session was estimated as a negative value
($\beta \pm SE$ = -0.161 ± 0.0308, z = -5.22, p < 0.001, 95% CI [-0.221, -0.0996]). The fixed effect of
condition ($\beta \pm SE$ = -0.0845 ± 0.196, z = -0.430, p = 0.667) and the session × condition interaction
($\beta \pm SE$ = 0.0190 ± 0.0426, z = 0.445, p = 0.656) were not significant. These results indicate that
exploratory responses decreased as habituation sessions progressed and that subjects
habituated to stimuli within the same class (cage-mate or non-cage-mate).

We next compared exploratory responses between the last habituation session and
first dishabituation session to examine whether dishabituation occurred (**Figure 2a**). Neither the
fixed effect of session ($\beta \pm SE$ = 0.182 ± 0.215, z = 0.847, p =0.397) nor condition ($\beta \pm SE$ = 0.106

± 0.219, z = 0.484, p = 0.629) nor the session × condition interaction ($β ± SE$ = 0.0678 ± 0.299, z

= 0.227, p = 0.821) was significant. These results suggest subjects did not show dishabituation in

response to the stimulus class switch.

Further inspection of the data revealed substantial individual variability in responses

during habituation sessions, suggesting that habituation had not been fully induced in some

individuals. Therefore, as an additional post-hoc analysis, we examined whether dishabituation

occurred in those that had habituated (N =21/30 in total; N = 11/15 from the CM condition, N =

10/15 from the NCM condition) (**Figure 2b**). We operationally defined individuals as sufficiently

habituated if they met both of the following criteria: (1) the response in the last habituation

session was less than two-thirds of that in the first habituation session, and (2) the average

response during the fourth to sixth habituation sessions was lower than that during the first to

third habituation sessions.

When responses were compared between the last habituation session and the first

dishabituation session in this subset, the fixed effect of session was significant ($β ± SE$ = 0.554 ±

0.226, z = 2.45, p = 0.0145, 95% CI [0.0967, 1.01]), whereas the fixed effect of condition ($β ± SE$

= 0.0618 ± 0.232, z = 0.267, p = 0.790) and the session × condition interaction ($β ± SE$ = 0.0649 ±

0.324, z = 0.200, p = 0.841) were not significant. These results suggest that exploratory responses

increased from the last habituation session to the first dishabituation session, indicating that

dishabituation occurred in individuals that had been adequately habituated during the

habituation phase.

4. Discussion

In this study, we examined social familiarity discrimination in rats using a habituation-

dishabituation paradigm. We presented multiple cage-mate or non-cage-mate stimuli during the

habituation phase and then tested whether rats would show dishabituation when the stimulus

class switched to the other one. During the habituation phase, exploratory responses decreased

as sessions progressed. Clear dishabituation to the stimulus class switch was not observed when

all individuals were pooled. However, the post-hoc analysis restricted to subjects that habituated

sufficiently revealed significant dishabituation.

Our results suggest that rats spontaneously exhibited a categorical response toward

familiarity or unfamiliarity in the current experimental setting and that rats can classify

conspecifics according to their familiarity. Insufficient habituation in some subjects may have

resulted from using live conspecific stimuli, whose activity or social relationships with the

subjects could have obscured responses. To control for these factors, it may be useful to restrain

stimulus animals' movement and increase the variety of stimulus animals. Also, in the

habituation-dishabituation paradigm, individual differences in the degree of habituation can be

large (Baciadonna et al., 2019; Charlton et al., 2007). In this study, we kept the number of
habituation sessions consistent across subjects to control the amount of stimulus exposure.
Future studies could increase exposure to habituation stimuli to ensure sufficient habituation in
all individuals.

The habituation-dishabituation paradigm with multiple conspecific stimuli is easy to
implement and may be a promising task for investigating social familiarity recognition and its
underlying mechanisms (c.f. mouse: Wolf et al., 2024; sheep: Kendrick et al., 2001). The specific
cues by which rats differentiate cage-mates from non-cage-mates remain unknown. Jones et al.
(2014) proposed that there may be group-specific odors (c.f. chemical cues of familiarity; ground
squirrel: Hare, 1994; bat: Rizzi et al., 2025), or that rats may form categories of familiarity and
unfamiliarity. Regarding odor cues, extended cohabitation may lead to convergence of odors
among group members through social interactions and the sharing of resources such as food and
a home base. One possible mechanism includes the transmission of gut microbiota among group
members through interactions (Raulo et al., 2021), as bacteria are known to influence the
discriminability of urine odors in rats (Schellinck et al., 1991).

It remains to be investigated whether rats utilize different types of social recognition,
such as individual recognition (Gheusi et al., 1997; Hopp et al., 1985) and the use of familiarity
heuristics, both of which may guide adaptive social behavior depending on the situation. For
instance, rats may rely on individual recognition when directly interacting with a small number
of individuals (Hakataya et al., 2023; Proops et al., 2021). However, in the wild, where there may
be many neighboring individuals (Davis, 1953) and odor cues may be degraded, familiarity
discrimination could be a more efficient way for managing frequent encounters.

Data availability Statement

Data and analysis codes are available from the following OSF project: https://osf.io/apngb/?view_only=5dd27745b1f54c92bc75f42f4e8b0360.

CRediT authorship contribution statement

- Shiomi Hakataya: Conceptualization, Methodology, Formal analysis, Investigation, Resources, Writing - Original Draft, Visualization, Funding acquisition
- Makiko Kamijo: Conceptualization, Methodology, Writing - Review & Editing, Funding acquisition
- Genta Toya: Methodology, Software, Writing - Review & Editing, Funding acquisition
- Hiroki Koda: Resources, Writing - Review & Editing, Funding acquisition
- Kazuo Okanoya: Writing - Review & Editing, Supervision, Funding acquisition

Acknowledgements

We thank Reo Wada and Kosuke Yoshida for their assistance in animal care. We are also grateful to the members of Okanoya laboratory at Teikyo University for stimulating discussion.

Funding

This work was supported by the Japan Society for the Promotion of Science Grant-in-Aid for JSPS Fellows (#22KJ0702, #24KJ0179) to SH, Early-Career Scientists (#21K13746) to MK, Transformative Research Areas (A, #24H01428; A, #21H05175) to GT, Scientific Research (S, #23H05428) to KO and (A, #21H04421; B, #22H03914) to HK. The funder had no role in study design, collection, analysis and interpretation of data, writing of the report or decision to submit the article for publication.

Declaration of Generative AI and AI-assisted Technologies in the Writing Process

During the preparation of this work, the first author used ChatGPT in order to improve the readability and language of the manuscript. After using this tool, the authors reviewed and edited the content as needed and take full responsibility for the content of the published article.

References

Alberts, J. R., & Galef, B. G., Jr. (1973). Olfactory cues and movement: stimuli mediating intraspecific aggression in the wild Norway rat. *Journal of Comparative and Physiological Psychology*, *85*(2), 233-242. https://doi.org/10.1037/h0035050

Baciadonna, L., Briefer, E. F., Favaro, L., & McElligott, A. G. (2019). Goats distinguish between positive and negative emotion-linked vocalisations. *Frontiers in Zoology*, *16*(1), 25. https://doi.org/10.1186/s12983-019-0323-z

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J., Maechler, M., & Bolker, B. M. (2017). glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R Journal*, *9*(2), 378–400. https://doi.org/doi:10.32614/RJ-2017-066

Charlton, B. D., Reby, D., & McComb, K. (2007). Female perception of size-related formant shifts in red deer, Cervus elaphus. *Animal Behaviour*, *74*(4), 707-714. https://doi.org//10.1016/j.anbehav.2006.09.021

Coulon, M., Baudoin, C., Heyman, Y., & Deputte, B. L. (2011). Cattle discriminate between familiar and unfamiliar conspecifics by using only head visual cues. *Animal Cognition*, *14*(2), 279-290. https://doi.org/10.1007/s10071-010-0361-6

Davis, D. E. (1953). The characteristics of rat populations. *The Quarterly Review of Biology*, *28*(4), 373-401. https://doi.org/10.1086/399860

Gheusi, G., Goodall, G., & Dantzer, R. (1997). Individually distinctive odours represent individual conspecifics in rats. *Animal Behaviour*, *53*(5), 935-944. https://doi.org//10.1006/anbe.1996.0314

Hackenberg, T. D., Vanderhooft, L., Huang, J., Wagar, M., Alexander, J., & Tan, L. (2021). Social preference in rats. *Journal of the Experimental Analysis of Behavior*, *115*(3), 634-649. https://doi.org /10.1002/jeab.686

Hakataya, S., Katsu, N., Okanoya, K., & Toya, G. (2023). An exploratory study of behavioral traits and the establishment of social relationships in female laboratory rats. *PLoS ONE*, *18*(12), e0295280. https://doi.org/10.1371/journal.pone.0295280

Hare, J. F. (1994). Group member discrimination by Columbian ground squirrels via familiarity with substrate-borne chemical cues. *Animal Behaviour*, *47*(4), 803-813. https://doi.org/10.1006/anbe.1994.1112

Hopp, S., Owren, M., & Marion, J. (1985). Olfactory discrimination of individual littermates in rats (*Rattus norvegicus*). *Journal of Comparative Psychology*, *99*(2), 248-251. https://doi.org/10.1037//0735-7036.99.2.248

Jones, S., Burman, O., & Mendl, M. (2014). Social discrimination of cage-mates and non-cage-mates by rats. *Behav Processes*, *106*, 130-140.

https://doi.org/10.1016/j.beproc.2014.05.007

Kendrick, K. M., da Costa, A. P., Leigh, A. E., Hinton, M. R., & Peirce, J. W. (2001). Sheep don't forget a face. *Nature*, *414*(6860), 165-166. https://doi.org/10.1038/35102669

Kiyokawa, Y., Honda, A., Takeuchi, Y., & Mori, Y. (2014). A familiar conspecific is more effective than an unfamiliar conspecific for social buffering of conditioned fear responses in male rats. *Behavioural Brain Research*, *267*, 189-193. https://doi.org/10.1016/j.bbr.2014.03.043

Kondo, S., Taniuchi, T., & Kondo, Y. (2005). Olfactory discrimination of male sexual odor in female rats using habituation-dishabituation paradigm. *The Japanese Journal of Animal Psychology*, *55*(1), 11-16.

Mathis, A., Mamidanna, P., Cury, K. M., Abe, T., Murthy, V. N., Mathis, M. W., & Bethge, M. (2018). DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nature Neuroscience*, *21*(9), 1281-1289. https://doi.org/10.1038/s41593-018-0209-y

Nath, T., Mathis, A., Chen, A. C., Patel, A., Bethge, M., & Mathis, M. W. (2019). Using DeepLabCut for 3D markerless pose estimation across species and behaviors. *Nature Protocols*, *14*(7), 2152-2176. https://doi.org/10.1038/s41596-019-0176-0

Proops, L., Troisi, C. A., Kleinhappel, T. K., & Romero, T. (2021). Non-random associations in group housed rats (*Rattus norvegicus*). *Scientific Reports*, *11*(1), 15349. https://doi.org/10.1038/s41598-021-94608-4

R Core Team (2025). *R: A language and environment for statistical computing*. In R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Raulo, A., Allen, B. E., Troitsky, T., Husby, A., Firth, J. A., Coulson, T., & Knowles, S. C. L. (2021). Social networks strongly predict the gut microbiota of wild mice. *The ISME Journal*, *15*(9), 2601-2613. https://doi.org/10.1038/s41396-021-00949-3

Rizzi, S., Schäfer, F., Schirmer, S., & Kerth, G. (2025). The smell of familiarity: conspecific faeces influence roost exploration and occupation in wild Bechstein's bats. *Animal Behaviour*, *224*, 123182. https://doi.org/10.1016/j.anbehav.2025.123182

Rogers-Carter, M. M., Djerdjaj, A., Culp, A. R., Elbaz, J. A., & Christianson, J. P. (2018). Familiarity modulates social approach toward stressed conspecifics in female rats. *PLoS ONE*, *13*(10), e0200971. https://doi.org/10.1371/journal.pone.0200971

Schellinck, H. M., Brown, R. E., & Slotnick, B. M. (1991). Training rats to discriminate between the odors of individual conspecifics. *Animal Learning & Behavior*, *19*(3), 223-233. https://doi.org/10.3758/BF03197880

Slotnick, B., Schellinck, H., & Brown, R. (2004). Olfaction. In I. Q. Whishaw & B. Kolb (Eds.), *The Behavior of the Laboratory Rat: A Handbook with Tests* (pp. 90-104). Oxford University

Press. https://doi.org/10.1093/acprof:oso/9780195162851.003.0009

Smith, C. J., Wilkins, K. B., Mogavero, J. N., & Veenema, A. H. (2015). Social Novelty Investigation

in the Juvenile Rat: Modulation by the μ-Opioid System. *Journal of Neuroendocrinology*,

27(10), 752-764. https://doi.org/10.1111/jne.12301

Suwandschieff, E., Mundry, R., Kull, K., Kreuzer, L., & Schwing, R. (2023). 'Do I know you?'

Categorizing individuals on the basis of familiarity in kea (*Nestor notabilis*). *Royal*

Society Open Science, *10*(6), 230228. https://doi.org/doi:10.1098/rsos.230228

Wada, R., Tomatsuri, T., Yoshida, K., Hakataya, S., Toya, G., Okanoya, K., & Koda, H. (2025). Rats

can spontaneously discriminate different rhythms: An investigation using an

automated habituation-dishabituation paradigm. *Behavioural Processes*, *231*, 105257.

https://doi.org/10.1016/j.beproc.2025.105257

Wiley, R. H. (2013). Specificity and multiplicity in the recognition of individuals: implications for

the evolution of social behaviour. *Biological Reviews*, *88*(1), 179-195. https://doi.org/

10.1111/j.1469-185X.2012.00246.x

Wilkinson, A., Specht, H. L., & Huber, L. (2010). Pigeons can discriminate group mates from

strangers using the concept of familiarity. *Animal Behaviour*, *80*(1), 109-115.

https://doi.org/ 10.1016/j.anbehav.2010.04.006

Wolf, D., Hartig, R., Zhuo, Y., Scheller, M. F., Articus, M., Moor, M., Grinevich, V., Linster, C., Russo,

E., Weber-Fahr, W., Reinwald, J. R., & Kelsch, W. (2024). Oxytocin induces the formation

of distinctive cortical representations and cognitions biased toward familiar mice.

Nature Communications, *15*(1), 6274. https://doi.org/10.1038/s41467-024-50113-6

Yorzinski, J. L. (2017). The cognitive basis of individual recognition. *Current Opinion in Behavioral*

Sciences, *16*, 53-57. https://doi.org/ 10.1016/j.cobeha.2017.03.009

Figure 1. Experimental procedures

(a) The experimental apparatus. The apparatus was a black acrylic open field. Stimulus

individuals were presented in black acrylic cages placed centrally along the back wall of the arena.

The lower half of the front side of each cage was covered with acrylic rods and allowed social

interactions between the subject and the stimulus animal. (b) Habituation-dishabituation

paradigm. During the arena acclimation phase, subjects were introduced to the arena and

allowed to explore freely for 10 minutes. Each stimulus presentation session lasted for 3 minutes.

The inter-session interval was 5 min. During habituation sessions (H1-H6), three cage-mate or

non-cage-mate stimuli were each presented twice in a pseudo-random order. In dishabituation

sessions (D1-D3), three different cage-mate or non-cage-mate stimuli were each presented once

in a random order.

Figure 2. Exploratory responses toward cage-mates and non-cage-mates

Box plots represent exploratory responses toward cage-mate and non-cage-mate stimuli. Blue indicates the cage-mate habituation condition (CMH) and red, the non-cage-mate habituation condition (NCMH), respectively. Black circles represent mean values. (a) whole data (N = 30; N = 15 from the CMH condition, N = 15 from the NCMH condition). (b) only individuals who showed sufficient habituation (N =21/30; N = 11/15 from the CM condition, N = 10/15 from the NCM condition). * indicates $p < 0.05$.

380