1

Pseudomonas aeruginosa lasR Mutants Resist Phagocytosis and Alter Inflammatory

Cytokine Production by Cystic Fibrosis Macrophages

Daniel S. Aridgides[1], Diane L. Mellinger[1], Lorraine L. Gwilt[1], Thomas H. Hampton[2], Dallas L.

Mould[3], Deborah A. Hogan[2], and Alix Ashare[1,2*]

[1]Section of Pulmonary and Critical Care Medicine, Dartmouth-Hitchcock Medical Center,

Lebanon, NH

[2]Department of Microbiology and Immunology, Geisel School of Medicine, Hanover, NH

[3]Department of Ecology and Evolutionary Biology, Yale University, New Haven, CT

* Correspondence: daniel.s.aridgides@hitchcock.org

Abstract

Cystic Fibrosis is characterized by chronic muco-obstructive lung disease and infection. People

with CF (pwCF) are often colonized with *Pseudomonas aeruginosa* for years to decades,

allowing for evolutionary adaptation. In chronic *P. aeruginosa* lung isolates from pwCF, the

quorum sensing regulator LasR frequently is nonfunctional, however the factors enabling *lasR*

loss-of-function (LOF) mutant selection are incompletely understood. We hypothesized that LOF

mutations in *lasR* could allow *P. aeruginosa* to resist the selective pressure of phagocytosis. We

found that in multiple strain backgrounds, LasR LOF decreased phagocytosis by both model

THP-1 and primary monocyte-derived macrophages. While exogenous administration of the

quorum-sensing autoinducer 3-oxo-C12-homoserine-lactone (3OC12HSL) that is made by an

enzyme regulated by LasR activity inhibited phagocytosis and mitochondrial respiration, the

phagocytosis resistance seen with *lasR* mutants appears to be bacterial cell intrinsic rather than

due to secreted factors. Finally, we found that *lasR* LOF mutations altered the inflammatory

profile upon infection of CF macrophages, with a shift from IL-1 family cytokine expression

towards canonical inflammatory markers including IL-6 and TNFα. Collectively these data

provide a potential explanation for both the prevalence of *lasR* mutants in the CF lung as well as

their association with worse outcomes.

Importance

Cystic Fibrosis (CF) is a genetically inherited disease that leads to chronic lung infections.

Pseudomonas aeruginosa is often implicated with worsening of lung disease, and it evolves in

the lung over time to resist eradication. One of the most commonly disrupted gene targets

identified in *Pseudomonas aeruginosa* isolates from chronically infected CF lungs is *lasR*, which

encodes a transcription factor which regulates multiple virulence factors. What contributes to the

apparent fitness of *lasR* mutants in the CF lung is not well known. Our study shows that *lasR*

loss-of-function (LOF) mutants resist phagocytosis by macrophages, one of the fundamental

mechanisms of clearance by the immune system. We identify mechanisms promoting

resistance to phagocytosis, and explore the downstream consequences on inflammatory

responses. Understanding why *lasR* mutations arise could inform strategies to eradicate them

from the CF lung and improve outcomes.

Introduction

Cystic Fibrosis (CF) is a multisystem disorder caused by mutations in the Cystic Fibrosis Transmembrane Conductance Regulator (*CFTR*) gene, encoding a chloride and bicarbonate channel (1). In the lungs of people with CF (pwCF), CFTR mutations lead to dehydrated, sticky mucus and subsequent colonization with bacteria often including *Pseudomonas aeruginosa* (2). Novel highly effective modulator compounds which improve trafficking and function of the CFTR protein have markedly improved outcomes for the subset of pwCF harboring CFTR mutations that are responsive to them (3–12), however CFTR modulators cannot restore sterilizing immunity to the lungs, particularly when there is pre-existing structural damage (13, 14). Given their relatively recent availability, the long-term trajectory of lung function in pwCF remains unknown, even for those who are clinically improved on modulators.

Lung function over time is impacted by bacterial colonization, with cycles of infection and dysregulated inflammation leading to lung damage (1). The specific composition of the lung microbiota as well as the host response to them play a role in determining whether lung function worsens or remains stable. Low bacterial diversity, *P. aeruginosa* colonization, and detection of *P. aeruginosa* with disruptions to the LasR/I quorum sensing system have all been shown to correlate with accelerated lung function decline in pwCF (15–17).

LasR loss-of-function (LOF) mutants are one of the most commonly isolated morphotypes found in CF lungs (18). LasR LOF mutants have also been isolated from environments such as pond water, and will arise de novo when *P. aeruginosa* are cultured in vitro (18–21). Their common emergence across various settings argues for a broad (or common) competitive advantage over LasR intact (LasR+) comparators, however the reasons for selection *in vivo* are not entirely clear. Social cheating whereby *lasR* LOF mutants can utilize secreted products from other *P. aeruginosa* isolates without incurring the metabolic cost of producing them has been proposed as a mechanism for *lasR* mutant selection. *lasR* mutation has also been shown increase

microoxic fitness and phosphate scavenging (22–24), providing additional possibilities for its

evolutionary advantage. Indeed *lasR* LOF mutants are selected for in vitro cultures due to a

growth advantage in post exponential phase (25–27) however these experimental systems lack

the host component. Interestingly, the phenotype of lack of protease activity has been shown to

increase in *P. aeruginosa* isolates that persist in pwCF on highly effective modulators,

suggesting *lasR* mutation may also be increased post modulator therapy (28).

In the CF lung *P. aeruginosa* must coexist with host immune cells which are attempting to

eradicate it, yet the effects of *lasR* LOF mutations on *P. aeruginosa* interactions with host

immune cells remain comparatively unstudied. Studies on interaction with airway epithelial cells

(29, 30), neutrophils (31, 32), and macrophages (33) have revealed variable effects on

inflammatory responses. Macrophages are responsible for orchestrating the host immune

response in the lung, phagocytosis of pathogens and debris, and have been shown to have

altered inflammatory and effector activity in CF (34). Although CFTR modulators have been

shown to improve CF macrophage function (35–37), an improved mechanistic understanding of

how modulators affect the host immune response could benefit the majority of pwCF who are

now eligible for highly effective modulator treatment and and lead to novel interventions that

might reduce bacterial persistence.

We hypothesized that loss of LasR function would alter interactions with host macrophages as a

mechanism of *lasR* mutant persistence within the CF lung, and that CFTR modulators would

further impact host response. We found that *lasR* LOF mutants indeed resist phagocytosis in

multiple *P. aeruginosa* backgrounds including laboratory and clinical isolates. Phagocytosis

resistance was maintained in mixed cultures with LasR+ relatives. *lasR* mutant resistance to

phagocytosis by primary CF monocyte derived macrophages (MDMs) was maintained despite

the presence of highly effective CFTR modulators. Finally, *lasR* mutation shifted the

inflammatory response of MDMs to *P. aeruginosa* infection away from IL-1 family cytokines

towards canonical proinflammatory cytokines IL-6 and TNFα. Collectively these results highlight

the importance of understanding *P. aeruginosa lasR* LOF in the context of immune cells and

provide potential explanations for its increased prevalence and virulence in CF. Some of these

results have previously been presented in the form of abstracts (38–42).

Results

***lasR* loss-of-function mutations confer phagocytosis resistance independent of**

background *P. aeruginosa* strain

To test the hypothesis that loss-of-function mutations in *lasR* confer phagocytosis resistance we

utilized model THP-1 monocytes which can be differentiated into macrophage-like cells by the

addition of the protein kinase C activator phorbol 12-myristate 13-acetate (43). We performed

gentamicin-protection assays with three pairs of LasR+ and LasR- *P. aeruginosa* to compare

rates of bacterial uptake. We chose a 20 minute timepoint to study phagocytosis as this

provides an investigation of uptake alone, whereas longer timepoints may see a mixture of

phagocytosis and killing (36). In strain PA14Δ*lasR*-THP-1 co-cultures, fewer cells were

phagocytosed then the in the otherwise isogenic wild-type (WT) strain (**Figure 1A**). We then

utilized two closely related pairs of LasR+/LasR- clinical isolates for which *lasR* LOF evolved

over the course of infection to confirm this effect with *P. aeruginosa* derived from the CF lung.

LasR LOF continued to confer phagocytosis resistance in both clinical isolate backgrounds

(**Figure 1B,C**) demonstrating that this was not a strain-specific effect.

LasR but not RhlR autoinducer homoserine lactone inhibits phagocytosis

One of the primary signaling molecules regulated by LasR is the autoinducer N-(3-

oxododecanoyl)-L-homoserine lactone (3OC12HSL) that binds to the LasR regulator to induce a

positive feedback cycle. 3OC12HSL has been shown to have direct effects on eukaryotic

metabolism and signaling (44–55), as well as enhancing phagocytosis of yeast (56). We

therefore tested whether 3OC12HSL would increase phagocytosis of *P. aeruginosa* as an

explanation for the decreased *lasR* mutant phagocytosis. Contrary to this hypothesis, we found

that 3OC12HSL inhibited phagocytosis of both LasR+ and LasR- PA14 at concentrations of 50

μM or higher in a dose-dependent manner (**Figure 2A**). By comparison, the RhlR autoinducer

C4-homoserine lactone (C4HSL) had no appreciable effect (**Figure 2B**). Under all

concentrations of HSLs tested there continued to be a reproducible difference between LasR+

and LasR- phagocytosis. These results suggest that 3OC12HSL is not responsible for the

effects seen in **Figure 1**.

Phagocytosis resistance of LasR- strains is maintained in mixed cultures

We therefore sought to ascertain whether *lasR* mutation effects were due to secreted or cell

intrinsic factors. We performed phagocytosis assays with a mixture of LasR+/LasR- PA14

derivatives at varying percentages (0/100, 10/90, 30/70, 50/50, 70/30, 90/10 and 100/0). The

LasR- competitor in the PA14 background bore a *lacZ* reporter gene and plating onto X-gal

plates allowed for post-assay determination of strain identities when mixed with an untagged

wild type. While we found bacterial uptake was generally proportional to the percent input for

each strain (**Figure 3A**), there was a persistently lower amount of LasR- *P. aeruginosa*

phagocytosed. **Figure 3B** displays the same data as in **3A** graphed relative to percent input for

each strain such that the decreased phagocytosis of LasR- strains is evident. These data

suggest that the effects of *lasR* mutation on phagocytosis is due to cell intrinsic factors.

Primary human macrophages augment phagocytosis with CFTR modulators, but LasR

phagocytosis resistance is maintained

To determine if the differences between phagocytosis of LasR+ and LasR- cells seen in THP-1

cells were also observed when using primary human cells, we isolated peripheral blood derived

mononuclear cells (PBMCs) from volunteers and differentiated them into monocyte-derived

macrophages (MDMs) in vitro for seven days. We first tested cells from CF volunteers. We and

others have demonstrated that CFTR modulators have a positive effect on macrophage and

monocyte phagocytosis (36, 37, 57–63), and as a majority of pwCF are now eligible for CFTR

modulators it is important to understand whether they may play an additional role in modifying

the response to *lasR* mutants. We tested one pair of laboratory isolates and one pair of clinical

isolates (DH1137/6) in MDMs that were treated for 48 hours with CFTR modulators or DMSO as

a control. The PA14Δ*lasR* (LasR-) strain was less efficiently phagocytosed by primary CF MDMs

relative to its LasR+ progenitor (**Figure 4A**). Tezacaftor/ivacaftor (TI) and

elexacaftor/tezacaftor/ivacaftor (ETI) improved phagocytosis of both strains in the PA14

background (**Figure 4A**) however there was no statistically significant interaction with *lasR*

status. We found similar results using paired clinical *P. aeruginosa* isolates, where *lasR* mutants

were less efficiently phagocytosed and CFTR modulators improved phagocytosis of both strains

with no difference in response to modulators between isolates (**Figure 4B**). We have previously

demonstrated that nonCF MDMs respond similarly to CFTR modulators when compared with

CF MDMs (37), therefore we tested whether any of the effects on *lasR* mutant phagocytosis

might be modified by *CFTR* genotype. NonCF MDMs displayed a similar pattern, with

decreased phagocytosis of *lasR* mutants, improved phagocytosis in the presence of CFTR

modulators, and no statistically significant interaction between modulators and *lasR* mutation

(**Figure 4C**). With the LasR+/LasR- clinical isolates, there continued to be an effect of *lasR*

mutation (**Figure 4D**) however double CFTR modulator treatment only displayed a non-

significant trend towards enhanced phagocytosis, with triple modulators still proving effective

(**Figure 4D**). Collectively these results confirm that *lasR* mutants exhibit reduced phagocytosis

by primary human CF and nonCF macrophages, and that while CFTR modulator treatment is

still beneficial, it fails to restore phagocytosis to the level of LasR+ strains.

***lasR* mutation abrogates the inhibitory effect of *P. aeruginosa* conditioned medium on**

macrophage mitochondrial respiration

Macrophage metabolism impacts their inflammatory predilection, with activated macrophages

relying on glycolysis for ATP generation whereas pro-resolution utilized aerobic respiration (64,

65). We have previously demonstrated that secreted products from *P. aeruginosa* inhibit aerobic

respiration in MDMs (37), and we hypothesized that *lasR* mutation may impact this effect. In

addition, we saw that ETI inhibits aerobic respiration without significantly affecting glycolysis.

We therefore performed Seahorse metabolic flux analysis on CF MDMs pre-treated with CFTR

modulators and then then conditioned media from either LasR+ or LasR- PA14 strains was

added acutely. Oxygen consumption rate (OCR) was measured as a proxy for mitochondrial

respiration and extracellular acidification rate (ECAR) as a proxy for lactate

generation/glycolysis. **Figure 5A-C** represent example readouts of OCR from a single CF

subject pre-treated with DMSO, TI or ETI, respectively and **Figure 5D-F** are the corresponding

readouts for ECAR. Addition of oligomycin, FCCP and rotenone/antimycin A allow for

measurement of additional parameters related to mitochondrial respiration. In CF MDMs, ETI

decreased basal respiration similarly to prior experiments whereas TI had no effect (**Figure 5G**).

Upon acute injection of LasR+ conditioned medium, there was a decrease in mitochondrial

respiration (**Figure 5H**), however LasR- conditioned medium completely lost this effect. In the

setting of low basal mitochondrial respiration with ETI, the effect of *P. aeruginosa* conditioned

medium (either LasR+ or LasR-) was lost (**Figure 5H**). Corresponding decreases in ATP-linked

respiration were seen with LasR+ conditioned medium and ETI pre-treatment (**Figure 5G**).

Basal glycolysis was not affected by CFTR modulators or different prior to injection (**Figure 5J**)

however LasR+ but not LasR- conditioned medium caused an acute rise in glycolysis (**Figure**

5K). This effect was lost upon ETI pre-treatment (**Figure 5K**). OCR/ECAR ratio is a way of

gauging relative contributions of each pathway to ATP-generation, and in line with the effects

seen on ATP-linked respiration and glycolysis, OCR/ECAR was decreased with LasR+ but not

LasR- conditioned media, as well as by ETI (**Figure 5L**). Measurements of maximal respiration,

spare respiratory capacity, non-mitochondrial respiration and proton leak are shown in

Supplemental Figure 1. These results collectively demonstrate that *lasR* mutation eliminates

the mitochondrial effects of *P. aeruginosa* exoproducts. Additionally, CFTR modulators inhibit

basal mitochondrial activity and there is no additive effect of *P. aeruginosa* exoproducts. Both of

these observations have potential downstream implications for induction of inflammation.

We next examined nonCF MDMs to determine whether there may be any differences in how

they respond to *lasR* mutants. Similarly to CF MDMs, we found the ETI inhibits basal respiration

(**Figure 5M**) with no effect of TI, in agreement with our prior published work (37). LasR+ but not

LasR- conditioned medium inhibited mitochondrial respiration (**Figure 5N**) and decreased ATP-

linked respiration (**Figure 5O**). These effects were decreased or eliminated by ETI-pretreatment.

Modulators had no effect on basal glycolysis (**Figure 5P**), with increased glycolysis secondary

to LasR+ but not LasR- conditioned medium (**Figure 5Q**). In contrast to CF MDMs, differences

in OCR/ECAR were not statistically significant (**Figure 5R**). Measurements of maximal

respiration, spare respiratory capacity, non-mitochondrial respiration and proton leak are shown

in **Supplemental Figure 2**.

LasR autoinducer homoserine lactone inhibits macrophage mitochondrial respiration

and induces glycolysis

The divergence between LasR+ and LasR- conditioned medium effects on MDM metabolism led

us to hypothesize that 3OC12HSL might underlie these effects. To test this, we returned to THP-

1 derived model macrophages and performed Seahorse metabolic analysis with acute injection

of 3OC12HSL, with C4HSL as a control. After first confirming no differences in basal respiration

prior to injection (**Figure 6A**), we found a dose responsive effect of 3OC12HSL injection (**Figure**

6B) causing inhibition of mitochondrial respiration whereas C4-HSL had no effect. This was

partially reflected in differences in ATP-linked respiration (**Figure 6C**) although the effects were

not statistically significant. Simultaneous measurement of glycolytic rates confirmed no basal

differences in glycolysis (**Figure 6D**), with a dose-dependent increase in glycolysis upon

3OC12HSL injection (**Figure 6E**) and no effect of C4HSL. The shift in metabolic profile from

mitochondrial respiration to glycolysis by 3OC12HSL was confirmed with decreased OCR/ECAR

(**Figure 6F**). Measurements of maximal respiration, spare respiratory capacity, non-

mitochondrial respiration and proton leak are shown in **Supplemental Figure 3**. These results

suggest that 3OC12HSL may underlie the effects of LasR+ *P. aeruginosa* exoproducts on

mitochondrial respiration.

***lasR* mutation shifts inflammatory cytokine profiles of primary human macrophages**

away from IL-1 family cytokines towards IL-6 and TNFα irrespective of CFTR modulators

and *CFTR* mutation status

Finally, we investigated whether *lasR* mutation impacted inflammatory profiles of MDMs, and the

interplay with CFTR modulators. We pretreated MDMs with CFTR modulators for 48 hours, then

infected MDMs for two hours with LasR+ or LasR- clinical isolates of *P. aeruginosa* and

performed cytokine multiplex analysis on the supernatants. Unsurprisingly we found broad

induction of multiple inflammatory cytokines upon *P. aeruginosa* infection of CF MDMs (**Figure**

7A). There was little appreciable effect of CFTR modulators, consistent with our prior work and

others' (37, 66). NonCF MDMs had a similar inflammatory response to CF MDMs, with no

statistically significant differences between the groups (**Figure 7B**). We then combined results

and tested specifically for differences between LasR+ and LasR- infection. IL-1 family cytokines

IL-1α, IL-1β and IL-18 were higher with LasR+ infection, whereas IL-6, IL-10, MCP-1, MIP-1α,

and MIP-1β were higher with LasR- infection (**Figure 7C**), demonstrating a specific alteration in

the inflammatory profile after macrophage response to *lasR* mutants.

Discussion

Loss-of-function mutations in *lasR* have been demonstrated to be important for *P. aeruginosa*

persistence and virulence within the CF lung (15, 18). How this impacts interactions with host

immune cells and specifically in the presence of highly effective CFTR modulators is

incompletely understood (29–33, 67). We have identified for the first time a robust and

reproducible resistance to phagocytosis by macrophages that is hypothesized to contribute to

persistence in the CF lung. While the difference was modest (0.3 to 0.5 log depending on the

assay), this represents a relative difference of 2-3 fold uptake in 20 minutes, therefore over

years this could represent a significant evolutionary pressure. The phagocytosis resistance was

present in multiple background strains including a common laboratory strain (PA14) and two

pairs of clinical isolates from the lungs of pwCF (**Figure 1**), increasing the generalizability of our

findings.

We also identified potential factors by which LasR mutants resist phagocytosis. The LasR

autoinducer 3OC12HSL inhibited *P. aeruginosa* phagocytosis (**Figure 2**), in contrast to prior

studies of yeast phagocytosis (56). If 3OC12HSL were responsible for differences in

phagocytosis however, it would be expected that LasR mutants would exhibit *higher*

phagocytosis rates, which was the opposite of what was seen. Differences in phagocytosis rates

were also maintained between LasR+ and LasR- strains when they were mixed in different

ratios (**Figure 3**), arguing that the responsible factors are cell intrinsic rather than secreted.

We also explored the interplay between CFTR modulators and *lasR* mutation with phagocytosis

and inflammation in primary MDMs. While CFTR modulators were able to enhance

phagocytosis of all strains tested consistent with our prior work (37), the decrement with *lasR*

mutation was maintained (**Figure 4**). Given the known persistence of *P. aeruginosa* in the lungs

of pwCF despite highly effective modulator therapy (14), this is relevant to understanding *P.*

aeruginosa colonization. We previously saw that *P. aeruginosa* secreted products inhibited

mitochondrial respiration in MDMs (37), as did the triple modulator combination ETI. Intriguingly

we found that *lasR* mutation essentially eliminated this effect (**Figure 5**). In combination with the

effects of exogenous 3OC12HSL (**Figure 6**), this suggests that the acyl-homoserine lactone is

the responsible factor, in line with previous studies of epithelial cells and fibroblasts (45, 46, 48–

51, 68).

Unsurprisingly given the variable rates of phagocytosis and differences in effects on

mitochondrial function in MDMs, we found that *lasR* mutation impacted inflammatory cytokine

profiles upon infection of CF and nonCF MDMs (**Figure 7**). While infection generally induced a

broad inflammatory cytokine profile, there was a shift from IL-1 family cytokines IL-1α, IL-1β, IL-

1RA and IL-18 towards more classical inflammatory markers IL-6 and TNFα, as well as

macrophage inflammatory protein 1α, 1β and macrophage chemoattractant protein 1. There

was a trend towards increased IL-8 for *lasR* mutant infection that was not statistically significant

after adjustment for multiple comparisons (**Figure 7C**). A shift towards IL-1 family cytokines may

reflect increased inflammasome activation versus direct proteolytic activation of IL-1 family

precursor proteins. The LasR-regulated elastase LasB has previously been demonstrated to

directly activate pro-IL-1β in mice and human macrophages (69, 70), consistent with our data

showing less mature IL-1β for infections with *lasR* loss-of-function mutants (**Figure 7C**). Early

CF lung *P. aeruginosa* isolates have been shown to be more potent inflammasome activators

when compared to later isolates (71), however LasR functional status was not tested in the

study by Phuong et al. Decreased protease expression has been seen in persistent *P.*

aeruginosa isolates in those on ETI, also suggesting *lasR* mutation may be increased post

modulators (28). It is important to note that in our system significant LasR quorum sensing

activation would not necessarily be expected, as bacteria are sub-cultured at low density prior to

addition to cells, and cells were infected for 2 hours. In contrast to its effects increasing

processed IL-1β, LasB can decrease IL-8 levels upon infection of airway epithelial cells via

direct digestion of IL-8 (72). Future experiments will help to elucidate the exact mechanism

whereby *lasR* mutation alters the inflammatory cytokine milieu in the shorter term. We are also

currently conducting transcriptomic studies on macrophages infected in parallel with LasR+ or

LasR- isolates which should shed light on whether the effects are transcriptional versus post-

transcriptional.

Strengths of our work include analysis of *lasR* mutations in multiple backgrounds including a

common laboratory strain as well as naturally occurring clinical isolates to improve the

generalizability. We began to investigate the mechanisms by which *lasR* loss-of-function

mutants may confer phagocytosis resistance although work remains to further elucidate the full

mechanism. We confirmed the results found in a human cell line (THP-1 cells) with primary

human macrophages from multiple subjects and investigated the interplay with CFTR

modulators and genotype (mutant vs wild-type CFTR). Studies performed in parallel on cells

from the same subject allowed for a close interrogation of the effects of CFTR modulators and

LasR mutation on phagocytosis, metabolism and inflammation.

Our work also carries several limitations. MDMs, while they are a commonly used surrogate for

primary lung macrophages due to their ease of procurement, do not behave identically to lung

macrophages (73). We plan to expand our work with primary lung macrophages isolated via

bronchoscopy (74), and compare them with classically generated MDMs as well as the novel

differentiation protocol proposed by Pahari et al. Additionally, while our ex vivo studies provided

the opportunity to directly compare effects of LasR status and modulators in paired samples

from the same subject, they are necessarily reductive and cannot replicate in vivo conditions.

Mouse models differ from human CF in chronicity and acuity of lung infection, however there are

data suggesting *lasR* mutants are more virulent in mice as well (31, 72). Finally, we found no

direct effect of CFTR modulators including ETI on inflammatory responses by MDMs, in contrast

to human studies pre- and post-modulator which demonstrate decreased inflammation (14, 60,

60, 75–78). In vivo studies however cannot separate direct effects on immune cells from

decreased mucus and pathogen burden leading to a systemic decrease in inflammation.

In summary, *lasR* mutation confers reproducible resistance to phagocytosis, altered metabolic

effects on macrophages, and a change in inflammatory cytokine repertoire upon infection.

These effects were independent of background strain of *P. aeruginosa*, as well as CFTR

genotype and CFTR modulator therapy. Our work adds to the body of literature on how and why

lasR mutants are problematic in the lungs of pwCF and will likely continue to be so in the highly

effective modulator era.

Materials and methods

Source of key chemicals and bacterial strains

	Source of materials	**Catalog number**
M-CSF	Miltenyi	130-096-491
Pan monocyte isolation kit	Miltenyi	130-096-537
3-oxo-C12-homoserinelactone (3OC12HSL)	Sigma	O9139
C4-homoserinelactone (C4HSL)	Chemodex	B0267
Ivacaftor	Selleckchem	S1144
Tezacaftor	Selleckchem	S7059
Elexacaftor	Selleckchem	S8851
Seahorse mitostress kit	Agilent	
THP-1 cells	ATCC	TIB-202
PMA	Sigma	P1585

PA14 WT	(79)	Hogan Lab (DH122)
PA14 ΔlasR	(80)	Hogan Lab (DH164)
PA14 ΔlasR att::PrhlI-lacZ	(81)	Hogan Lab
DH1137 (LasR+)	(82)	Hogan Lab (DH1137)
DH1136 (LasR-)	(82)	Hogan Lab (DH1136)
DH2417 (LasR+), NC-AMT0101-1-2 (parent of NC-AMT0101-1)	(82)	Hogan Lab (DH2417)
DH2415 (LasR-), NC-AMT0101-1-1, related to DH2417 with LasR LOF (frameshift) allele	(82)	Hogan Lab (DH2415)

THP-1 cell culture

THP-1 cells were cultured in RPMI with 10% FBS, 50 µM β-mercaptoethanol, and 50 µg/mL

gentamicin in a 37°C humidity-controlled incubator with 5% CO_2. The initial vial was expanded

and frozen down in aliquots which were stored in liquid nitrogen with DMSO. Cultures from

individual aliquots were maintained in vitro for ~2 months prior to being discarded. THP-1

monocytes were differentiated into macrophages by addition of 50 nM Phorbol 12-myristate 13-

acetate (PMA) for 48 hours prior to experiments.

Gentamicin protection phagocytosis assays with THP-1 derived macrophages

THP-1 cells were differentiated at 5 x 10^5/mL in 24-well plates. They were then washed with

PBS and medium was replaced with identical medium without antibiotics. Overnight LB cultures

of bacteria of indicated strains were sub-cultured 1:10 into LB for 60 minutes, centrifuged at

8000 x g for 2 minutes, washed twice with antibiotic-free THP-1 medium, then resuspended and

bacterial density was determined by spectrophotometry at 600 nm. OD600 of 1.0 was

empirically determined to represent 10^9 bacteria/mL media. After dilution, bacteria were added to

THP-1 cells at a multiplicity of infection (MOI) of 10 bacteria per macrophage. Plates were

incubated at 37°C for 20 minutes, then cells were washed once with PBS, then THP-1 medium

with 10x gentamicin (500 µg/mL) was added for 15 minutes. Cells were then washed twice with

PBS at 4°C followed by lysis with 250 µL 0.1% Triton X-100. Lysates were diluted 1:3 in PBS

and then 10 µL was then plated overnight onto LB agar plates for colony forming unit (CFU)

determination. Colonies were counted the next day and bacteria per 10^6 macrophages was

calculated. MOI was also determined empirically from the stock bacterial solutions for each

assay as an internal control. Average of at least three technical replicates per assay was then

log-transformed and results were combined from at least 5 assays performed on separate days

(n for each experiment listed in figure legends).

Phagocytosis competition assay

PA14 Δ*lasR* with a *lacZ* reporter gene integrated at a neutral site on the chromosome was used

along with PA14 wild type (WT). Assay was identical to above except fixed percentages of PA14

WT and the Δ*lasR* strain were used at 100/0, 90/10, 70/30, 50/50, 30/70, 10/90, and 0/100 such

that the total MOI of both strains together was constant at 10. Lysates were plated onto LB agar

plates with 5-bromo-4-chloro-3-indolyl-β-D-galactoside (X-gal) for blue-white colony

discrimination. 100/0 samples were entirely white and 0/100 samples were entirely blue,

confirming the fidelity of the reporter. Total PA14 WT and Δ*lasR* colonies were counted for each

well. Each experiment was performed in triplicate and the means of technical replicates for each

assay were log-transformed and combined.

Human subjects

Our blood draw protocol has been approved by the Dartmouth Health Institutional Review

Board. Subjects were recruited from our CF clinic population if they were 18-65 years old,

F508del homozygous or heterozygous, at clinical baseline without symptoms of exacerbation,

and had normal hemoglobin levels. NonCF control subjects were recruited by IRB-approved

flyers; smokers or those with respiratory disease, immunologic disorders, or pulmonary

medication use were excluded. Phlebotomy was performed (100 mL whole blood isolated) for

monocyte isolation by our research nurse and taken directly to the laboratory for downstream

processing.

Monocyte derived macrophage (MDM) generation and phagocytosis

MDMs were generated according to previously published protocols (37). Briefly, monocytes

were isolated from peripheral blood by Ficoll gradient separation. They were quantified and

plated directly onto plates as per intended assay to avoid the need for detachment and re-

plating. MDMs were generated by treatment with 100 ng/mL M-CSF for seven days in

RPMI/10% FCS with 50 µg/mL gentamicin. Medium was exchanged every 2-3 days. After seven

days of differentiation, MDMs were cultured in RPMI/10% FCS + gentamicin, and CFTR

modulators were added for 48 hours prior to the experiment. CFTR modulators were replaced

with all media exchanges. Gentamicin protection assay was performed as above. Three to four

technical replicates per subject were averaged and log-transformed. Data on paired *lasR*

mutants were combined with previously published data on *lasR*-WT strains (37).

Seahorse extracellular flux assay

Monocytes were plated initially onto a Seahorse 96-well plate at 50,000 per well and

differentiated into MDM directly in the wells for seven days as above. At that point, differentiation

medium was removed and CFTR modulators were added for 48 hours. Pa-conditioned medium

was prepared in advance by culturing bacteria overnight in Seahorse mitostress assay medium,

normalizing cultures to an OD600 of 0.500, centrifuging to pellet bacteria then passing

supernatant through a 0.22 µm filter. This was defined as 100% conditioned medium. Aliquots

were then frozen at -20° C until use.

On the day of assay, cell culture medium was exchanged for mitostress assay medium as per

manufacturer's instructions. A mitostress assay with acute injection was run on an Agilent

Seahorse XFe96 platform, with 20 µL conditioned medium in injection port A (final concentration

10%), plain mitostress medium was included for control wells. Oligomycin, FCCP, and

Rotenone/Antimycin A were placed into ports B-D respectively. Four to eight technical replicates

per condition were run. Parameters including response to injection, maximal respiration, and

others were calculated as per manufacturer's instructions. Data were combined with previously

published results with a *lasR*-WT strain only (37).

THP-1 Seahorse extracellular flux assay

THP-1 macrophages were differentiated directly on a Seahorse 96-well plate with 50 nM PMA at

50,000 cells/well for 48 h. Seahorse extracellular flux assay was then run with varying

concentrations of 3OC12HSL or C4HSL in injection port A. Five to six technical replicates were

run per condition for each assay and averaged. Five independent experiments were run.

Cytokine analysis

MDMs were differentiated for seven days as above then treated for 48 hours with CFTR

modulators or DMSO. Bacteria were prepared in an identical manner to the phagocytosis

assays. After exchange for antibiotic-free media, MDMs were infected with indicated strains at

an MOI of 10 for 2 hours, then washed and placed in media with gentamicin for an additional 2

hours. Supernatants were then collected, centrifuged at 14,000 x g at 4°C for 10 minutes,

transferred to a new tube, and frozen at -20°C until needed. Samples were batched and

cytokine multiplex analysis was run by the Dartmouth Immunology and Flow Cytometry Core.

One replicate per condition was run. Samples with levels below the limit of the detection of the

assay were set to a concentration of 0, and data were log-transformed prior to analysis.

Figure generation and statistical analysis

Figures were generated with GraphPad Prism 10. Statistical analyses were performed

separately in R, with mixed model linear analysis using nlme package version 3.1. Models

include all variables of interest as fixed variables and assay date (for THP-1 experiments) or

subject (for MDM experiments) as random variables.

Acknowledgments

Funding for DSA provided by the CF Foundation (ARIDGI19B0, ARIDGI20L0, STANTO19R0).

THH is supported by DartCF (NIH grant P30-DK117469) and Cystic Fibrosis Foundation

Research Development Program (BOMBER24G0). AA is supported by the NIH (R56HL158923,

R01HL174700) and CFF (ASHARE21G0). Dartmouth Immunology and Flow Cytometry Core is

supported by NIH 5P30CA023108. The authors declare no competing interests and the funders

had no direct role in the design or conduct of these studies.

1.   Elborn JS. 2016. Cystic fibrosis. Lancet 388:2519–2531.

2.   Patient Registry | Cystic Fibrosis Foundation. https://www.cff.org/medical-

professionals/patient-registry. Retrieved 1 June 2022.

3.   Barry PJ, Mall MA, Álvarez A, Colombo C, de Winter-de Groot KM, Fajac I, McBennett

KA, McKone EF, Ramsey BW, Sutharsan S, Taylor-Cousar JL, Tullis E, Ahluwalia N, Jun

LS, Moskowitz SM, Prieto-Centurion V, Tian S, Waltz D, Xuan F, Zhang Y, Rowe SM,

Polineni D, VX18-445-104 Study Group. 2021. Triple Therapy for Cystic Fibrosis

Phe508del-Gating and -Residual Function Genotypes. N Engl J Med 385:815–825.

4.   Bower JK, Volkova N, Ahluwalia N, Sahota G, Xuan F, Chin A, Weinstock TG, Ostrenga

J, Elbert A. 2023. Real-world safety and effectiveness of elexacaftor/tezacaftor/ivacaftor in

people with cystic fibrosis: Interim results of a long-term registry-based study. J Cyst

Fibros S1569-1993(23)00066–8.

5.   Bass R, Stalvey M, Solomon G, Rowe S, Nichols D, Schwarzenberg SJ, Freedman S,

Walega R, Kelly A. 2025. Cholesterol and triglyceride concentrations following 12-

18 months of clinically prescribed elexacaftor-tezacaftor-ivacaftor-PROMISE sub-study. J

Clin Transl Endocrinol 40:100391.

6.   Burgel P-R, Durieu I, Chiron R, Ramel S, Danner-Boucher I, Prevotat A, Grenet D,

Marguet C, Reynaud-Gaubert M, Macey J, Mely L, Fanton A, Quetant S, Lemonnier L,

Paillasseur J-L, Da Silva J, Martin C, Andrejak C, Becourt A, Mounard J, Poulet C, Rames

C, Talleux M, Chevalier M-C, Darviot E, Jouvenot M, Marien C, Paris A, Pelatan C,

Person C, Priou P, Troussier F, Urban T, Dalphin M-L, Dalphin J-C, Ladaurade A, Pernet

D, Richaud-Thiriez B, Roux-Claude P, Blanc N, Boisserie-Lacroix V, Bui S, Collet C,

Debelleix S, Macey J, Bergot E, Brouard J, Campbell K, Laurans M, Ribault V, Borderon

C, Heraud M-C, Labbe G, Montcouquiol S, Petit I, Ruivard M, Delestrain C, Douvry B,

Epaud R, Maitre B, Remus N, Beltramo G, Fanton A, Houzel A, Huet F, Perez S, Boldron-

Ghaddar A, Scalbert M, Bouzioukh R, Mely L, Simon C, Camara B, Hamidfar R, Llerena

C, Pin I, Quetant S, Deschildre A, Gicquello A, Le Rouzic O, Leroy C, Paris N, Perez T,

Prevotat A, Thumerelle C, Turck D, Wizla N, Dupuy-Grasset M, Languepin J, Masson-

Rouchaud A, Menetrey C, Durieu I, Durupt S, L'Excellent S, Nove-Josserand R, Ohlmann

C, Reix P, Reynaud Q, Werck-Gallois M-C, Baravalle M, Coltey B, Desmazes-Dufeu N,

Dubus J-C, Gautier C, Rey J-B, Reynaud-Gaubert M, Stremler N, Caimmi D, Chiron R,

Devrait M, Moreau J, Billon Y, Blondé A, Guillaumot A, Kiefer S, Peretti L, Tatopoulos A,

Tiotiu A, Benhamida M, Bihouee T, Danner-Boucher I, Eschapasse E, Tissot A,

Giannantonio M, Leroy S, Piccini-Bailly C, Pradelli J, Messika J, Boussaud V, Burgel P-R,

Carlier N, Honoré I, Hubert D, Kanaan R, Martin C, Bailly-Botuha C, Chedevergne F, De

Blic J, Delacourt C, Drummond D, Fauroux B, Karila C, Le Bourgeois M, Sermet I, Delaisi

B, Gerardin M, Houdouin V, Leclainche L, Aubertin G, Berdah L, Clement A, Corvol H,

Nathan N, Prevost B, Richard N, Tamalet A, Taytard J, Thouvenin G, Tourniaire B, Abely

M, Bessaci-Kabouya K, Dury S, Ravoninjatovo B, Dabadie A, Dagorne M, Deneuville E,

Jamin M, Ribault M, Vigier C, Belleguic C, Brinchault G, Desrues B, Barzic A, Dirou-

Prigent A, Le Bihan J, Ramel S, Revert K, Ropars T, Couderc L, Dominique S, Marguet C,

Morisse-Pradier H, Pramil S, Thiberville L, Allou N, Enaud L, Gachelin E, Huchot E, Payet

A, Perisson C, Piyaraly S, Valois S, Herzog A, Kessler R, Porzio M, Weiss L, Beaumont L,

Brugiere O, Colin S, Verdiere de, Cuquemelle E, De Miranda S, Grenet D, Monem Hamid

A, Parquin F, Picard C, Roux A, Roy C, Bremont F, Didier A, Dupuis M, Faviez G,

Labouret G, Mittaine M, Murris-Espin M, Roditis L, Cosson L, Diot P, Flament T, Giraut

C, Mankikian J, Arnouat B, Mousset G, Storni V, Vigneron P, Coirier-Duet E, Gabsi A.

2021. Rapid Improvement after Starting Elexacaftor–Tezacaftor–Ivacaftor in Patients with

Cystic Fibrosis and Advanced Pulmonary Disease. Am J Respir Crit Care Med 204:64–73.

7. Daines CL, Polineni D, Tullis E, Costa S, Linnemann RW, Mall MA, McKone EF, Quon

BS, Ringshausen FC, Selvadurai H, Taylor-Cousar JL, Withers NJ, Sawicki GS, Lee T,

Ahluwalia N, Morlando Geiger J, Jennings M, Tan YV, Waltz D, Ramsey B, Griese M,

VX17-445-105 Study Group. 2025. Long-Term Safety and Efficacy of

Elexacaftor/Tezacaftor/Ivacaftor in Adults and Adolescents with Cystic Fibrosis and at

Least One F508del Allele: A Phase 3, Open-Label Extension Study. Am J Respir Crit Care

Med https://doi.org/10.1164/rccm.202411-2231OC.

8. Daines CL, Tullis E, Costa S, Linnemann RW, Mall MA, McKone EF, Polineni D, Quon

BS, Ringshausen FC, Rowe SM, Selvadurai H, Taylor-Cousar JL, Withers NJ, Ahluwalia

N, Moskowitz SM, Prieto-Centurion V, Tan YV, Tian S, Weinstock T, Xuan F, Zhang Y,

Ramsey B, Griese M, VX17-445-105 Study Group. 2023. Long-term safety and efficacy of

elexacaftor/tezacaftor/ivacaftor in people with cystic fibrosis and at least one F508del

allele: 144-week interim results from a 192-week open-label extension study. Eur Respir J

62:2202029.

9. Heijerman HGM, McKone EF, Downey DG, Van Braeckel E, Rowe SM, Tullis E, Mall

MA, Welter JJ, Ramsey BW, McKee CM, Marigowda G, Moskowitz SM, Waltz D, Sosnay

PR, Simard C, Ahluwalia N, Xuan F, Zhang Y, Taylor-Cousar JL, McCoy KS, VX17-445-

103 Trial Group. 2019. Efficacy and safety of the elexacaftor plus tezacaftor plus ivacaftor

combination regimen in people with cystic fibrosis homozygous for the F508del mutation: a

double-blind, randomised, phase 3 trial. Lancet 394:1940–1948.

10. McNally P, Lester K, Stone G, Elnazir B, Williamson M, Cox D, Linnane B, Kirwan L,

Rea D, O'Regan P, Semple T, Saunders C, Tiddens HAWM, McKone E, Davies JC,

RECOVER Study Group. 2023. Improvement in Lung Clearance Index and Chest

Computed Tomography Scores with Elexacaftor/Tezacaftor/Ivacaftor Treatment in People

with Cystic Fibrosis Aged 12 Years and Older - The RECOVER Trial. Am J Respir Crit

Care Med 208:917–929.

11. Middleton PG, Mall MA, Dřevínek P, Lands LC, McKone EF, Polineni D, Ramsey BW,

Taylor-Cousar JL, Tullis E, Vermeulen F, Marigowda G, McKee CM, Moskowitz SM, Nair

N, Savage J, Simard C, Tian S, Waltz D, Xuan F, Rowe SM, Jain R, VX17-445-102 Study

Group. 2019. Elexacaftor-Tezacaftor-Ivacaftor for Cystic Fibrosis with a Single Phe508del

Allele. N Engl J Med 381:1809–1819.

12. Nichols DP, Paynter AC, Heltshe SL, Donaldson SH, Frederick CA, Freedman SD,

Gelfond D, Hoffman LR, Kelly A, Narkewicz MR, Pittman JE, Ratjen F, Rosenfeld M,

Sagel SD, Schwarzenberg SJ, Singh PK, Solomon GM, Stalvey MS, Clancy JP, Kirby S,

Van Dalfsen JM, Kloster MH, Rowe SM, PROMISE Study Group. 2021. Clinical

Effectiveness of Elexacaftor/Tezacftor/Ivacaftor in People with Cystic Fibrosis. Am J

Respir Crit Care Med https://doi.org/10.1164/rccm.202108-1986OC.

13. Heltshe SL, Mayer-Hamblett N, Burns JL, Khan U, Baines A, Ramsey BW, Rowe SM,

GOAL (the G551D Observation-AL) Investigators of the Cystic Fibrosis Foundation

Therapeutics Development Network. 2015. Pseudomonas aeruginosa in cystic fibrosis patients with G551D-CFTR treated with ivacaftor. Clin Infect Dis 60:703–712.

14. Nichols DP, Morgan SJ, Skalland M, Vo AT, Van Dalfsen JM, Singh SB, Ni W, Hoffman LR, McGeer K, Heltshe SL, Clancy JP, Rowe SM, Jorth P, Singh PK, PROMISE-Micro Study Group. 2023. Pharmacologic improvement of CFTR function rapidly decreases sputum pathogen density, but lung infections generally persist. J Clin Invest 133:e167957.

15. Hoffman LR, Kulasekara HD, Emerson J, Houston LS, Burns JL, Ramsey BW, Miller SI. 2009. PSEUDOMONAS AERUGINOSA lasR MUTANTS ARE ASSOCIATED WITH CYSTIC FIBROSIS LUNG DISEASE PROGRESSION. J Cyst Fibros 8:66–70.

16. Delhaes L, Monchy S, Fréalle E, Hubans C, Salleron J, Leroy S, Prevotat A, Wallet F, Wallaert B, Dei-Cas E, Sime-Ngando T, Chabé M, Viscogliosi E. 2012. The Airway Microbiota in Cystic Fibrosis: A Complex Fungal and Bacterial Community—Implications for Therapeutic Management. PLOS ONE 7:e36313.

17. Zhao K, Yang X, Zeng Q, Zhang Y, Li H, Yan C, Li JS, Liu H, Du L, Wu Y, Huang G, Huang T, Zhang Y, Zhou H, Wang X, Chu Y, Zhou X. 2023. Evolution of lasR mutants in polymorphic Pseudomonas aeruginosa populations facilitates chronic infection of the lung. Nat Commun 14:5976.

18. Feltner JB, Wolter DJ, Pope CE, Groleau M-C, Smalley NE, Greenberg EP, Mayer-Hamblett N, Burns J, Déziel E, Hoffman LR, Dandekar AA. 2016. LasR Variant Cystic Fibrosis Isolates Reveal an Adaptable Quorum-Sensing Hierarchy in Pseudomonas aeruginosa. MBio 7.

19. Hammond JH, Hebert WP, Naimie A, Ray K, Van Gelder RD, DiGiandomenico A, Lalitha

P, Srinivasan M, Acharya NR, Lietman T, Hogan DA, Zegans ME. 2016. Environmentally

Endemic Pseudomonas aeruginosa Strains with Mutations in lasR Are Associated with

Increased Disease Severity in Corneal Ulcers. mSphere 1.

20. Scribner MR, Stephens AC, Huong JL, Richardson AR, Cooper VS. 2022. The nutritional

environment is sufficient to select coexisting biofilm and quorum-sensing mutants of

Pseudomonas aeruginosa. J Bacteriol JB0044421.

21. Groleau M-C, Taillefer H, Vincent AT, Constant P, Déziel E. 2021. Pseudomonas

aeruginosa isolates defective in function of the LasR quorum sensing regulator are frequent

in diverse environmental niches. Environ Microbiol https://doi.org/10.1111/1462-

2920.15745.

22. Mould DL, Stevanovic M, Ashare A, Schultz D, Hogan DA. 2022. Metabolic basis for the

evolution of a common pathogenic Pseudomonas aeruginosa variant. Elife 11:e76555.

23. Mould DL, Finger CE, Conaway A, Botelho N, Stuut SE, Hogan DA. 2024. Citrate cross-

feeding by Pseudomonas aeruginosa supports lasR mutant fitness. mBio 15:e0127823.

24. Clay ME, Hammond JH, Zhong F, Chen X, Kowalski CH, Lee AJ, Porter MS, Hampton

TH, Greene CS, Pletneva EV, Hogan DA. 2020. Pseudomonas aeruginosa lasR mutant

fitness in microoxia is supported by an Anr-regulated oxygen-binding hemerythrin. PNAS

https://doi.org/10.1073/pnas.1917576117.

25. Feng X, Kostylev M, Dandekar AA, Greenberg EP. 2019. Dynamics of cheater invasion in

a cooperating population of Pseudomonas aeruginosa. Sci Rep 9:10190.

26.   Chen R, Déziel E, Groleau M-C, Schaefer AL, Greenberg EP. 2019. Social cheating in a

Pseudomonas aeruginosa quorum-sensing variant. PNAS 201819801.

27.   D'Argenio DA, Wu M, Hoffman LR, Kulasekara HD, Déziel E, Smith EE, Nguyen H,

Ernst RK, Larson Freeman TJ, Spencer DH, Brittnacher M, Hayden HS, Selgrade S,

Klausen M, Goodlett DR, Burns JL, Ramsey BW, Miller SI. 2007. Growth phenotypes of

Pseudomonas aeruginosa lasR mutants adapted to the airways of cystic fibrosis patients.

Mol Microbiol 64:512–533.

28.   Durfey SL, Kapnadak SG, Pena T, Willmering MM, Godwin JD, Teresi ME, Heltshe SL,

Vo AT, Villacreses RA, Aliukonyte I, Boyken L, Stroik MR, Morgan SJ, Wang GM, Betts

HL, Zhang S, Goss CH, Clancy JP, Aitken ML, Steele C, Feder AF, Esther CR, Tiddens

HAWM, Woods JC, Stoltz DA, Singh PK. 2025. Pseudomonas infections persisting after

CFTR modulators are widespread throughout the lungs and drive lung inflammation. Cell

Host Microbe 33:1428-1445.e4.

29.   Pseudomonas aeruginosa Modulates the Antiviral Response of Bronchial Epithelial Cells -

PubMed. https://pubmed.ncbi.nlm.nih.gov/32117250/. Retrieved 15 June 2022.

30.   Dulon S, Leduc D, Cottrell GS, D'Alayer J, Hansen KK, Bunnett NW, Hollenberg MD,

Pidard D, Chignard M. 2005. Pseudomonas aeruginosa elastase disables proteinase-

activated receptor 2 in respiratory epithelial cells. Am J Respir Cell Mol Biol 32:411–419.

31.   Hennemann LC, LaFayette SL, Malet JK, Bortolotti P, Yang T, McKay GA, Houle D,

Radzioch D, Rousseau S, Nguyen D. 2021. LasR-deficient Pseudomonas aeruginosa

variants increase airway epithelial mICAM-1 expression and enhance neutrophilic lung

inflammation. PLoS Pathog 17:e1009375.

32. Skopelja-Gardner S, Theprungsirikul J, Lewis KA, Hammond JH, Carlson KM, Hazlett HF,

Nymon A, Nguyen D, Berwin BL, Hogan DA, Rigby WFC. 2019. Regulation of

Pseudomonas aeruginosa-Mediated Neutrophil Extracellular Traps. Front Immunol

10:1670.

33. Bastaert F, Kheir S, Saint-Criq V, Villeret B, Dang PM-C, El-Benna J, Sirard J-C,

Voulhoux R, Sallenave J-M. 2018. Pseudomonas aeruginosa LasB Subverts Alveolar

Macrophage Activity by Interfering With Bacterial Killing Through Downregulation of

Innate Immune Defense, Reactive Oxygen Species Generation, and Complement

Activation. Front Immunol 9.

34. Bruscia EM, Bonfield TL. 2016. Cystic Fibrosis Lung Immunity: The Role of the

Macrophage. J Innate Immun 8:550–563.

35. Hazlett HF, Hampton TH, Aridgides DS, Armstrong DA, Dessaint JA, Mellinger DL,

Nymon AB, Ashare A. 2020. Altered iron metabolism in cystic fibrosis macrophages: the

impact of CFTR modulators and implications for Pseudomonas aeruginosa survival. Sci

Rep 10:10935.

36. Barnaby R, Koeppen K, Nymon A, Hampton TH, Berwin B, Ashare A, Stanton B. 2017.

Lumacaftor (VX-809) restores the ability of CF macrophages to phagocytose and kill

Pseudomonas aeruginosa. Am J Physiol Lung Cell Mol Physiol ajplung004612017.

37.   Aridgides DS, Mellinger DL, Gwilt LL, Hampton TH, Mould DL, Hogan DA, Ashare A.

596   2023. Comparative effects of CFTR modulators on phagocytic, metabolic and inflammatory

597   profiles of CF and nonCF macrophages. Sci Rep 13:11995.

598   38.   2018. The 32nd Annual North American Cystic Fibrosis Conference, Colorado Convention

599   Center, Denver, Colorado October 18-20, 2018. Pediatric Pulmonology 53:S148–S456.

600   39.   2019. The 33rd Annual North American Cystic Fibrosis Conference, October 31-

601   November 2, 2019. Pediatric Pulmonology 54:S155–S480.

602   40.   2020. The 34th Annual North American Cystic Fibrosis Conference, October 7-23, 2020.

603   Pediatr Pulmonol 55 Suppl 2:S38–S361.

604   41.   Aridgides D, Mellinger D, Ashare A. 2022. 553 Determinants of LasR mutant

605   Pseudomonas aeruginosa phagocytosis resistance. Journal of Cystic Fibrosis 21:S307–

606   S308.

607   42.   Aridgides D, Mellinger D, Dessaint J, Hogan D, Ashare A. 2021. 432: Pseudomonas

608   aeruginosa LasR loss-of-function mutants evade phagocytosis by CF macrophages despite

609   CFTR triple modulator therapy. Journal of Cystic Fibrosis 20:S203.

610   43.   Lund ME, To J, O'Brien BA, Donnelly S. 2016. The choice of phorbol 12-myristate 13-

611   acetate differentiation protocol influences the response of THP-1 macrophages to a pro-

612   inflammatory stimulus. J Immunol Methods 430:64–70.

44. Cruz RL, Asfahl KL, Van den Bossche S, Coenye T, Crabbé A, Dandekar AA. 2020. RhlR-

Regulated Acyl-Homoserine Lactone Quorum Sensing in a Cystic Fibrosis Isolate of

Pseudomonas aeruginosa. mBio 11:e00532-20.

45. Jacobi CA, Schiffner F, Henkel M, Waibel M, Stork B, Daubrawa M, Eberl L, Gregor M,

Wesselborg S. 2009. Effects of bacterial N-acyl homoserine lactones on human Jurkat T

lymphocytes-OdDHL induces apoptosis via the mitochondrial pathway. International

Journal of Medical Microbiology 299:509–519.

46. Josephson H, Ntzouni M, Skoglund C, Linder S, Turkina MV, Vikström E. 2020.

Pseudomonas aeruginosa N-3-Oxo-Dodecanoyl-Homoserine Lactone Impacts

Mitochondrial Networks Morphology, Energetics, and Proteome in Host Cells. Front

Microbiol 11:1069.

47. Kravchenko VV, Kaufmann GF, Mathison JC, Scott DA, Katz AZ, Wood MR, Brogan AP,

Lehmann M, Mee JM, Iwata K, Pan Q, Fearns C, Knaus UG, Meijler MM, Janda KD,

Ulevitch RJ. 2006. N-(3-oxo-acyl)homoserine lactones signal cell activation through a

mechanism distinct from the canonical pathogen-associated molecular pattern recognition

receptor pathways. J Biol Chem 281:28822–28830.

48. Neely AM, Zhao G, Schwarzer C, Stivers NS, Whitt AG, Meng S, Burlison JA, Machen

TE, Li C. N-(3-Oxo-acyl)-homoserine lactone induces apoptosis primarily through a

mitochondrial pathway in fibroblasts. Cellular Microbiology 20:e12787.

49.   Schwarzer C, Fu Z, Shuai S, Babbar S, Zhao G, Li C, Machen TE. 2014. Pseudomonas

633   aeruginosa Homoserine Lactone Triggers Apoptosis and Bak/Bax-Independent Release of

634   Mitochondrial Cytochrome C in Fibroblasts. Cell Microbiol 16:1094–1104.

635   50.   Singh PK, Yadav VK, Kalia M, Sharma D, Pandey D, Agarwal V. 2019. Pseudomonas

636   aeruginosa quorum-sensing molecule N-(3-oxo-dodecanoyl)-L-homoserine lactone triggers

637   mitochondrial dysfunction and apoptosis in neutrophils through calcium signaling. Med

638   Microbiol Immunol 208:855–868.

639   51.   Tao S, Niu L, Cai L, Geng Y, Hua C, Ni Y, Zhao R. 2018. N-(3-oxododecanoyl)-l-

640   homoserine lactone modulates mitochondrial function and suppresses proliferation in

641   intestinal goblet cells. Life Sciences 201:81–88.

642   52.   Tateda K, Ishii Y, Horikawa M, Matsumoto T, Miyairi S, Pechere JC, Standiford TJ,

643   Ishiguro M, Yamaguchi K. 2003. The Pseudomonas aeruginosa Autoinducer N-3-

644   Oxododecanoyl Homoserine Lactone Accelerates Apoptosis in Macrophages and

645   Neutrophils. Infection and Immunity 71:5785–5793.

646   53.   Telford G, Wheeler D, Williams P, Tomkins PT, Appleby P, Sewell H, Stewart GSAB,

647   Bycroft BW, Pritchard DI. 1998. The Pseudomonas aeruginosaQuorum-Sensing Signal

648   MoleculeN-(3-Oxododecanoyl)-l-Homoserine Lactone Has Immunomodulatory Activity.

649   Infection and Immunity 66:36–42.

650   54.   Yadav VK, Singh PK, Sharma D, Singh SK, Agarwal V. 2019. Mechanism underlying N-

651   (3-oxo-dodecanoyl)-L-homoserine lactone mediated intracellular calcium mobilization in

652   human platelets. Blood Cells Mol Dis 79:102340.

55. Wagner C, Zimmermann S, Brenner-Weiss G, Hug F, Prior B, Obst U, Hänsch GM. 2007.

The quorum-sensing molecule N-3-oxododecanoyl homoserine lactone (3OC12-HSL)

enhances the host defence by activating human polymorphonuclear neutrophils (PMN).

Anal Bioanal Chem 387:481–487.

56. Vikström E, Magnusson K-E, Pivoriūnas A. 2005. The Pseudomonas aeruginosa quorum-

sensing molecule N-(3-oxododecanoyl)-L-homoserine lactone stimulates phagocytic

activity in human macrophages through the p38 MAPK pathway. Microbes Infect 7:1512–

1518.

57. Cavinato L, Luly FR, Pastore V, Chiappetta D, Sangiorgi G, Ferrara E, Baiocchi P,

Mandarello G, Cimino G, Del Porto P, Ascenzioni F. 2022. Elexacaftor-Tezacaftor-

Ivacaftor corrects monocyte microbicidal deficiency in cystic fibrosis. Eur Respir J

2200725.

58. Assani K, Shrestha CL, Rinehardt H, Zhang S, Robledo-Avila F, Wellmerling J, Partida-

Sanchez S, Cormet-Boyaka E, Reynolds SD, Schlesinger LS, Kopp BT. 2018. AR-13

reduces antibiotic-resistant bacterial burden in cystic fibrosis phagocytes and improves

cystic fibrosis transmembrane conductance regulator function. J Cyst Fibros

https://doi.org/10.1016/j.jcf.2018.10.010.

59. Kopp BT, Nicholson L, Paul G, Tobias J, Ramanathan C, Hayes D. 2015. Geographic

variations in cystic fibrosis: An analysis of the U.S. CF Foundation Registry. Pediatr

Pulmonol 50:754–762.

60. Sheikh S, Britt RD, Ryan-Wenger NA, Khan AQ, Lewis BW, Gushue C, Ozuna H,

Jaganathan D, McCoy K, Kopp BT. 2023. Impact of elexacaftor-tezacaftor-ivacaftor on

bacterial colonization and inflammatory responses in cystic fibrosis. Pediatr Pulmonol

58:825–833.

61. Shrestha CL, Zhang S, Wisniewski B, Häfner S, Elie J, Meijer L, Kopp BT. 2020. (R)-

Roscovitine and CFTR modulators enhance killing of multi-drug resistant Burkholderia

cenocepacia by cystic fibrosis macrophages. Sci Rep 10:21700.

62. Bitto NJ, Baker PJ, Dowling JK, Wray-McCann G, De Paoli A, Tran LS, Leung PL, Stacey

KJ, Mansell A, Masters SL, Ferrero RL. 2018. Membrane vesicles from Pseudomonas

aeruginosa activate the noncanonical inflammasome through caspase-5 in human

monocytes. Immunology & Cell Biology 96:1120–1130.

63. Gabillard-Lefort C, Casey M, Glasgow AMA, Boland F, Kerr O, Marron E, Lyons A-M,

Gunaratnam C, McElvaney NG, Reeves EP. 2022. Trikafta Rescues CFTR and Lowers

Monocyte P2X7R-Induced Inflammasome Activation in Cystic Fibrosis. Am J Respir Crit

Care Med https://doi.org/10.1164/rccm.202106-1426OC.

64. Van den Bossche J, O'Neill LA, Menon D. 2017. Macrophage Immunometabolism: Where

Are We (Going)? Trends in Immunology 38:395–406.

65. Hamilton K, Krause K, Badr A, Daily K, Estfanous S, Eltobgy M, Khweek AA, Anne

MNK, Carafice C, Baetzhold D, Tonniges JR, Zhang X, Gavrilin MA, Parinandi NL, Amer

AO. 2020. Defective immunometabolism pathways in cystic fibrosis macrophages. Journal

of Cystic Fibrosis 0.

66.   Zhang S, Shrestha CL, Robledo-Avila F, Jaganathan D, Wisniewski BL, Brown N, Pham

695   H, Carey K, Amer AO, Hall-Stoodley L, McCoy KS, Bai S, Partida-Sanchez S, Kopp BT.

696   2022. Cystic fibrosis macrophage function and clinical outcomes after

697   elexacaftor/tezacaftor/ivacaftor. Eur Respir J 2102861.

698   67.   Hennemann LC, Nguyen D. 2021. LasR-regulated proteases in acute vs. chronic lung

699   infection: a double-edged sword. Microb Cell 8:161–163.

700   68.   Maurice NM, Bedi B, Yuan Z, Goldberg JB, Koval M, Hart CM, Sadikot RT. 2019.

701   Pseudomonas aeruginosa Induced Host Epithelial Cell Mitochondrial Dysfunction. Sci Rep

702   9:11929.

703   69.   Everett MJ, Davies DT, Leiris S, Sprynski N, Llanos A, Castandet JM, Lozano C, LaRock

704   CN, LaRock DL, Corsica G, Docquier J-D, Pallin TD, Cridland A, Blench T, Zalacain M,

705   Lemonnier M. 2023. Chemical Optimization of Selective Pseudomonas aeruginosa LasB

706   Elastase Inhibitors and Their Impact on LasB-Mediated Activation of IL-1β in Cellular and

707   Animal Infection Models. ACS Infect Dis 9:270–282.

708   70.   Sun J, LaRock DL, Skowronski EA, Kimmey JM, Olson J, Jiang Z, O'Donoghue AJ, Nizet

709   V, LaRock CN. 2020. The Pseudomonas aeruginosa protease LasB directly activates IL-1β.

710   EBioMedicine 60:102984.

711   71.   Phuong MS, Hernandez RE, Wolter DJ, Hoffman LR, Sad S. 2021. Impairment in

712   inflammasome signaling by the chronic Pseudomonas aeruginosa isolates from cystic

713   fibrosis patients results in an increase in inflammatory response. Cell Death Dis 12:241.

714   72.   LaFayette SL, Houle D, Beaudoin T, Wojewodka G, Radzioch D, Hoffman LR, Burns JL,

715   Dandekar AA, Smalley NE, Chandler JR, Zlosnik JE, Speert DP, Bernier J, Matouk E,

716   Brochiero E, Rousseau S, Nguyen D. 2015. Cystic fibrosis-adapted Pseudomonas

717   aeruginosa quorum sensing lasR mutants cause hyperinflammatory responses. Sci Adv 1.

718   73.   Pahari S, Arnett E, Simper J, Azad A, Guerrero-Arguero I, Ye C, Zhang H, Cai H, Wang Y,

719   Lai Z, Jarvis N, Lumbreras M, Maselli DJ, Peters J, Torrelles JB, Martinez-Sobrido L,

720   Schlesinger LS. 2023. A new tractable method for generating human alveolar macrophage-

721   like cells in vitro to study lung inflammatory processes and diseases. mBio 14:e0083423.

722   74.   Aridgides D, Dessaint J, Atkins G, Carroll J, Ashare A. 2021. Safety of research

723   bronchoscopy with BAL in stable adult patients with cystic fibrosis. PLoS One

724   16:e0245696.

725   75.   Jarosz-Griffiths HH, Scambler T, Wong CH, Lara-Reyna S, Holbrook J, Martinon F, Savic

726   S, Whitaker P, Etherington C, Spoletini G, Clifton I, Mehta A, McDermott MF, Peckham

727   D. 2020. Different CFTR modulator combinations downregulate inflammation differently

728   in cystic fibrosis. Elife 9:e54556.

729   76.   Casey M, Gabillard-Lefort C, McElvaney OF, McElvaney OJ, Carroll T, Heeney RC,

730   Gunaratnam C, Reeves EP, Murphy MP, McElvaney NG. 2023. Effect of

731   elexacaftor/tezacaftor/ivacaftor on airway and systemic inflammation in cystic fibrosis.

732   Thorax 78:835–839.

733   77.   Jarosz-Griffiths HH, Gillgrass L, Caley LR, Spoletini G, Clifton IJ, Etherington C, Savic S,

734   McDermott MF, Peckham D. 2024. Anti-inflammatory effects of

elexacaftor/tezacaftor/ivacaftor in adults with cystic fibrosis heterozygous for F508del.

PLoS One 19:e0304555.

78.   De Vuyst RC, Bennard E, Kam CW, McKinzie CJ, Esther CR. 2023.

Elexacaftor/tezacaftor/ivacaftor treatment reduces airway inflammation in cystic fibrosis.

Pediatr Pulmonol 58:1592–1594.

79.   Rahme LG, Stevens EJ, Wolfort SF, Shao J, Tompkins RG, Ausubel FM. 1995. Common

virulence factors for bacterial pathogenicity in plants and animals. Science 268:1899–1902.

80.   Hogan DA, Vik A, Kolter R. 2004. A Pseudomonas aeruginosa quorum-sensing molecule

influences Candida albicans morphology. Mol Microbiol 54:1212–1223.

81.   Mould DL, Botelho NJ, Hogan DA. 2020. Intraspecies Signaling between Common

Variants of Pseudomonas aeruginosa Increases Production of Quorum-Sensing-Controlled

Virulence Factors. mBio 11:e01865-20.

82.   Smith EE, Buckley DG, Wu Z, Saenphimmachak C, Hoffman LR, D'Argenio DA, Miller

SI, Ramsey BW, Speert DP, Moskowitz SM, Burns JL, Kaul R, Olson MV. 2006. Genetic

adaptation by Pseudomonas aeruginosa to the airways of cystic fibrosis patients. Proc Natl

Acad Sci U S A 103:8487–8492.

Figure 1. *lasR* mutation confers phagocytosis resistance independent of parental strain.

A) THP-1 cells were differentiated into macrophages by addition of 50 nM PMA for 48 hours.

Phagocytosis was then measured by a gentamicin protection assay at 20 minutes of bacterial

uptake. After removal of extracellular bacteria by washing and addition of gentamicin,

intracellular CFU were quantified by plating onto LB agar. Phagcoytosis of PA14 wild type and

PA14 Δ*lasR* (LasR-) was assessed in eight independent assays, log-transformed means of

three technical replicates per assay are displayed. ** p<0.01 for PA14 WT vs LasR- by mixed

model linear regression. **B)** Identical assays to **A** were run using naturally occurring paired

LasR+/- clinical isolates of *P. aeruginosa* derived from the same parental strain DH1137. **

p<0.01 for PA14WT vs LasR- by mixed model linear regression. **C)** Identical assays to **A** were

run using naturally occurring paired LasR+/- clinical isolates of *P. aeruginosa* derived from the

same parental strain DH2417. ** p<0.01 for PA14WT vs LasR- by mixed model linear

regression.

Figure 2. LasR-regulated autoinducer inhibits phagocytosis in a dose-dependent manner.

A) THP-1 macrophages were incubated with the indicated concentration of 3OC12HSL for 20

minutes prior to performing a gentamicin phagocytosis assay with PA14 WT and PA14Δ*lasR*

(LasR-) in parallel. Four separate experiments were performed with three technical replicates

each. The means of technical replicates from each individual experiment are shown in the light

lines, with darker lines at the means of the four experiments. *$p<0.05$, **** $p<0.0001$ for 50 and

250 μM 3OC12HSL relative to DMSO respectively, and LasR+ vs LasR- by mixed model linear

regression. **B)** Identical assays to **A** were performed with C4HSL in place of 3OC12HSL. ****

$p<0.0001$ by mixed model linear regression.

Figure 3. *lasR* mutation confers phagocytosis resistance in mixed cultures. A) A

PA14Δ*lasR* strain engineered to express a LacZ reporter was mixed with PA14 wild type at 0%,

10%, 30%, 50%, 70%, 90% and 100% with a fixed total MOI of 10 for the two strains and a

phagocytosis assay was performed as previously. After lysis, bacteria were plated onto X-gal

plates to allow for differentiation of colony genotype. Means of three technical replicates from 6

independent experiments are graphed, with a dark line at the mean of the 6 experiments. **B)**

The same data as in **A** are graphed relative to the percent of each strain input, rather than

percent LasR- to better visualize differences. **** p<0.0001 by mixed model linear regression.

Figure 4. *lasR* mutants exhibit reduced phagocytosis by primary MDMs irrespective of CFTR genotype or modulators. A) CF MDMs were treated for 48 hours with CFTR modulators as indicated, then infected with PA14 WT or PA14 Δ*lasR* (LasR-)*.* Lines connecting dots represent data from a single subject (n=7). **B)** As in **A** except the clinical isolate DH1137 and its derivative *lasR* mutant were used. **C)** NonCF MDMs (n=5) were infected with PA14 WT or LasR- derivative as in **A**. D) NonCF MDMs were infected with DH1137 or its *lasR* mutant derivative as in **B**. ** p<0.01, ***p<0.001, **** p<0.0001 for indicated comparisons by mixed model linear regression. Data were combined with previously published data on LasR+ strains only (37).

Figure 5. Effects of CFTR modulators and LasR+/- secreted products on macrophage

metabolism. CF and nonCF MDM were differentiated with M-CSF then treated for 48 hours

with CFTR modulators. A Seahorse mitostress assay with acute injection of *P. aeruginosa*

conditioned medium was then performed. **A-C**) Representative oxygen consumption rate

tracings from a CF subject MDMs pretreated with DMSO, TI, or ETI. **D-F**) Representative

extracellular acidification rates from the same subject. **G-L**) Calculated parameters as indicated

based on the mitostress assay of CF MDMs (n=10). Each point with connecting lines represents

the mean of technical replicates from an individual subject. Bars represent means of all the

subjects within the group. **M-R**) Calculated parameters as indicated for nonCF MDMs (n=11).

*p<0.05, ** p<0.01, ***p<0.001, **** p<0.0001 for indicated comparisons by mixed model linear

regression. Data were combined with previously published data on LasR+ strains only (37).

Figure 6. 3OC12HSL inhibits mitochondrial respiration in THP-1 macrophages. A-F)

Seahorse mitostress assay was run with acute injection of various concentrations of 3OC12HSL

or C4HSL. Parameters were calculated as per standard protocol. Points represent averages of

6-8 technical replicates from 6 individual experiments, with solid lines at the global means.

*p<0.05, ** p<0.01, ***p<0.001, **** p<0.0001 for indicated comparisons relative to DMSO

treated wells by mixed model linear regression.

Figure 7. *lasR* mutation alters inflammatory cytokine expression profile by CF and nonCF

MDMs. A) CF MDMs (n=11 subjects) were pre-treated with CFTR modulators and then infected

with DH1137 (LasR+) or DH1136 (LasR-) for two hours, then cells were washed and media

replaced. Supernatants were collected 2 hours after washing and analyzed by cytokine

multiplex assay. Means of log-transformed data are graphed. **B)** nonCF MDMs (n=4) were

infected in the same way and cytokines analyzed by multiplex. **C)** Volcano plot of fold difference

between LasR+ and LasR- cytokine results, DMSO treated group only.

831