Title

A self-limiting mechanotransduction feedback loop ensures robust organ

formation

Authors

Yusuke Mori[1,2], Paul Robin[3], Anne Belle Briggs[1], Daniel S. Levic[1], Jiacheng Wang[1], Kira

L. Heikes[1], Michel Bagnat[1], Edouard Hannezo[3], Akankshi Munjal[1,4]

[1] Department of Cell Biology, Duke University School of Medicine, Durham, NC 27710,

USA

[2] Duke Regeneration Center, Duke University School of Medicine, Durham, NC 27710,

USA

[3] Institute of Science and Technology, Am Campus 1, 3400 Klosterneuburg, Austria

[4] Correspondence: akankshi.munjal@duke.edu

Abstract

Organ development requires integration of mechanical forces, biochemical signals, and
transcriptional programs to achieve proper size and architecture. Mechanotransduction
feedback loops convert mechanical forces into gene-expression changes that, in turn,
regulate cell behaviors such as growth and extracellular matrix (ECM) production. How
these loops are coupled to developmental programs to ensure robust morphogenetic
outcomes remains poorly understood. Here, we show that Yap-dependent
mechanotransduction establishes a self-limiting positive feedback loop that drives the
successful formation of semicircular canals in zebrafish. These canals form through
reproducible steps of bud initiation, extension, and fusion within the otic epithelium. Local
swelling of a hyaluronan-rich ECM initiates bud formation. We demonstrate that this ECM
expansion activates Yap in a spatially patterned manner within the bud. Activated Yap
induces its target, *ccn1l1,* which drives further ECM production, establishing a positive
feedback loop. The ensuing rapid ECM expansion sustains bud extension. Graded
perturbations to the mechanotransduction loop reduced extension, with bud fusion
remaining robust at lower levels of disruption but blocked at the highest dose. Critically,
the loop contains its own termination mechanism: when buds fuse, the PKA-CREB
signaling is activated by the adhesion GPCR Gpr126 to suppress *ccn1l1*, preventing
overgrowth. These findings reveal how mechanotransduction-driven positive feedback
loops can be coupled to their own termination, providing developmental control through
the integration of mechanical forces with transcriptional responses and morphogenetic
outcomes.

Introduction

Signaling pathways employ sophisticated feedback mechanisms to control tissue morphogenesis– the process by which tissues remodel into complex and functional organs[1,2]. Such feedback can take the form of temporal switches or self-limiting circuits, for example, the morphogen Nodal induces its own inhibitor, Lefty, to establish early embryonic patterning[3,4]. Beyond biochemical signals, developing tissues also experience and respond to physical forces. Mechanotransduction pathways sense forces such as compression, stretching, or shear and convert them into transcriptional responses that remodel the physical environment, creating feedback loops between mechanics and gene expression during morphogenesis[5,6].

Yes-associated protein 1 (Yap), a transcriptional co-activator of the Hippo pathway, is a central mechanotransducer[7]. In response to force, Yap and its co-activator Taz translocate to the nucleus, where they bind to the transcription factor Tead to activate genes that regulate proliferation, differentiation, and extracellular matrix (ECM) production[7]. By linking mechanical inputs to transcriptional programs that reshape tissue architecture, Yap provides a molecular basis for feedback between mechanics and gene expression[8]. Yet how such mechanotransduction feedback loops are spatially and temporally organized during development—and how they are ultimately terminated to avoid unchecked amplification or premature growth restriction—remains poorly understood. Understanding these principles is central to explaining how complex organ shapes emerge from the interplay of signals, forces, and transcriptional programs.

The zebrafish semicircular canals (SCC) enable addressing these questions because they allow for the simultaneous measurement of transcriptional, biochemical, and mechanical dynamics during morphogenesi[9]. SCCs arise from the otic vesicle (OV) epithelium through three discrete steps: bud initiation, extension, and fusion. Bud initiation is driven by localized synthesis of a hyaluronan- and Versican-rich ECM that osmotically swells to push the overlying epithelium[9,10]. A key challenge is to understand how this process, initiated in a small group of cells, achieves the systems-level coordination necessary to ensure the buds extend to the right size and fuse successfully. We

hypothesized that mechanotransduction could provide this coordinating mechanism,
linking local ECM expansion to transcriptional responses.

Our work reveals that SCC morphogenesis is driven by a self-limiting positive feedback
loop operating through Yap. Using a combination of quantitative imaging, theoretical
modeling, and a live knock-in reporter of endogenous Yap, we observe that ECM
expansion mechanically activates Yap in a spatially patterned manner, with peak activity
at the bud border, recruiting these cells into the growing bud. Yap in turn induces *cellular*
communication network factor 1-like 1 (*ccn1l1,* also known as *cyr61l1*), a matricellular
protein that promotes HA and Versican synthesis, ensuing rapid ECM expansion.
Progressive loss of the Yap-Ccn1l1-ECM loop led to a dose-dependent reduction in
extension rates, yet fusion remained robust at lower levels of disruption and was
abolished only at the highest dose.

Termination of this loop occurs during bud fusion when *ccn1l1* expression declines.
Analysis of mutants in a conserved adhesion GPCR, *gpr126*, previously shown to block
bud fusion, revealed that *ccn1l1* expression persists, resulting in uncontrolled growth.
Mechanistically, bud fusion activates the transcription factor CREB via the Gpr126-cAMP-
PKA pathway, which suppresses *ccn1l1*. Thus, the morphological outcome of positive
feedback–bud fusion–serves as a stop signal that terminates the very process it drives–
bud extension.

Results

ECM swelling-mediated Yap activation is required for bud extension

The SCC in zebrafish develops through the remodeling of the lumenized single-layered
embryonic inner ear epithelium, the basement membrane (BM)-enclosed OV, at six
"canal-genesis zones" (Fig. 1a)[9,11]. Morphogenesis begins at 45 hours post fertilization
(hpf) with the initiation of buds at antero-lateral, postero-lateral, and anterior zones, where
cells extend into the lumen with reduced BM contact[9,12]. By 50 hpf, these buds extend,
and a posterior bud begins to form. By 60 hpf, antero-lateral and anterior buds fuse, while
postero-lateral and posterior buds extend, and ventral and ventro-lateral budding begins.

By 72 hpf, all bud pairs fuse to form the anterior, posterior, and ventral pillars, which
demarcate the three semicircular canals. We focused our analysis on the anterior buds,
which provide a clear example of all morphogenetic steps: prebudding (40 hpf), initiation
(45-48 hpf), extension (48-55 hpf), and fusion (55 hpf).

We recently demonstrated that the transcription factor Lmx1b promotes the expression
of *has3 (hyaluronan synthase 3)* and *vcanb (versican b),* which encode enzymes for HA
and Versican synthesis, respectively, at canal-genesis zones[10]. HA polymers and the
proteoglycan Versican that binds HA are secreted basally into the ECM, and osmotic
swelling of this HA-rich ECM generates pressure that pushes the overlying epithelium into
buds[10]. Longitudinal bud extension is maintained by tangentially distributed actomyosin
tethers, known as cytocinches[9]. Because ECM swelling imposes mechanical stress on
budding cells[9] (Extended Fig. 1a-c), we asked whether this stress engages the Yap
mechanotransduction pathway.

To characterize the dynamic localization of endogenous Yap protein *in vivo*, we
established a *TgKI(Yap1-mScarlet)* reporter line using a knock-in tagging method[13] (Fig.
1b-b'). Yap activity was quantified by calculating the nuclear-cytoplasmic ratio in budding
cells in embryos co-injected with H2B-GFP mRNA to visualize the nuclei. Prior to budding
(40 hpf), Yap was inactive, with a nuclear-cytoplasmic ratio around 1. During anterior bud
extension at 50 hpf, the ratio increased above 1, indicating active Yap (Fig 1c). We also
observed a few cells in the dorsal non-budding region of the OV with a nuclear-
cytoplasmic ratio greater than 1 at various stages of canal morphogenesis (Fig. 1b and
Extended Fig. 1d).

To test whether ECM-generated mechanical forces activate Yap in budding cells, we
injected hyaluronidase (HAase), an enzyme that degrades HA polymers, into the region
surrounding the OV (periotic space). As previously reported, HAase treatment caused
shrinkage of the bud-ECM, reduced cell stretching, and inhibited bud extension[9,14]
(Extended Fig. 1e-h). Significantly, nuclear localization of Yap was decreased in HAase-
treated embryos, with the nuclear-cytoplasmic ratio less than 1 (Fig. 1d-e). HAase
treatment had no effect on the nuclear-cytoplasmic ratio of non-budding cells in the dorsal

OV (Extended Fig. 1i). These findings were further confirmed by Yap immunostaining
experiments, in which the nuclear-cytoplasmic ratio in budding cells was greater than 1
in controls but reduced to around 1 in HAase-treated embryos (Extended Fig. 1j-m).
Together, these results demonstrate that Yap activation in budding cells is induced by
hyaluronate pressure during bud extension.

To examine the functional role of Yap, we performed knockdown of *yap1* using a
previously characterized morpholino (MO)[15]. Notably, *yap1* knockdown significantly
inhibited bud elongation (Fig. 1f-g), resulting in shorter lateral, anterior, and posterior buds
at 50 hpf. By 72 hpf, all buds had formed, but not all had fused into pillars, indicating a
delay in morphogenesis (Extended Fig. 1n). These data demonstrate that Yap drives bud
extension. To uncover the molecular mechanisms underlying this effect, we next
examined Yap's downstream transcriptional targets.

Yap-target *ccn1l1* expression is Yap-independent at initiation and Yap-dependent
during bud extension

To identify downstream targets of Yap, we used available single-cell transcriptomic
atlases[16-18]. A zebrafish inner-ear-specific atlas noted expression of a candidate gene,
ccn1l1, in the SCC-forming cell cluster of the OV, and this expression pattern was
validated at the canal-genesis zone during pre-budding and bud extension stages[16]. A
similar enrichment was observed in the embryo-wide Daniocell atlas[17,18], where
annotation of canal-forming cells using *vcanb* (Versican core protein) revealed *ccn1l1*
among the top 10 differentially expressed genes (Fig. 2a–b, Extended Fig. 2a). Ccn1l1 is
a paralog of Ccn1 and a member of the CCN family of matricellular proteins[19–21], which
are known downstream targets of Yap[22–25]. Importantly, the 5' upstream region of *ccn1l1*
contains three conserved TEAD-binding motifs recognized by Yap/Taz
(GGAATG/CATTCC)[22,26–28] (Extended Fig. 2b).

To quantitatively analyze *ccn1l1* expression dynamics at various developmental stages,
we used multiplexed hybridization chain reaction (HCR) fluorescence *in situ* hybridization
(FISH)[29]. Consistent with the single-cell atlases[16,18], *ccn1l1* expression was restricted to

the canal-genesis zones (Fig. 2c-d), and was detectable before bud formation at 40 hpf.

Its expression increased significantly during bud extension at 50 hpf, and was

subsequently downregulated to background levels upon bud fusion at 60 hpf (Fig. 2e).

Importantly, MO-mediated *yap1* knockdown led to a 50% reduction in *ccn1l1* expression

during bud extension (Fig. 2f-g), making it comparable to levels at the pre-budding stage,

confirming that *ccn1l1* is a Yap target. In contrast, *yap1* knockdown did not affect the

onset of *ccn1l1* expression at the pre-budding stage (Extended Fig. 2c-d). These findings

demonstrate that Yap is essential for increasing *ccn1l1* expression during bud extension,

but not for its initial onset.

To understand the initial regulation of *ccn1l1*, we examined its relationship with *lmx1bb*,

the transcription factor that, together with its paralog *lmx1ba,* we previously demonstrated

patterns HA and Versican synthesis during the pre-budding stage[10]. In the *lmx1bb* null

mutants (*lmx1b* *jj410/jj410*)[29], which carry a premature stop codon, we observed a 30.1%

reduction of *ccn1l1* expression at the pre-budding zone compared to sibling controls

(Extended Fig. 2e-f). Pharmacological inhibition of Notch signaling with RO4929097,

which expands *lmx1bb* expression[10], similarly expanded *ccn1l1* expression compared to

controls (Extended Fig. 2g).

Altogether, these findings reveal a two-stage regulatory mechanism: Lmx1bb in part

initiates *ccn1l1* expression during pre-budding, while Yap amplifies *ccn1l1* expression

during extension. Given that Yap sustains bud extension, we hypothesized that *ccn1l1*

might promote the synthesis of HA and Versican, which drive bud extension. We next

investigated the function of *ccn1l1* to test this possibility.

Ccn1l1 promotes HA and Versican synthesis to create a positive feedback loop

To investigate the function of *ccn1l1*, we performed CRISPR-Cas9 or MO-mediated

ccn1l1 knockdown. Consistent with its pre-budding expression pattern, the knockdown of

ccn1l1 blocked the onset of budding, resulting in an absence of buds at 50 hpf and no

SCC at 72 hpf (Fig. 3a-b). We next asked how Ccn1l1 drives epithelial budding. CCN

proteins regulate cell behaviors such as adhesion, proliferation, and ECM production

through their conserved domains[30–32]. Zebrafish Ccn1l1 exhibits high similarity to human

CCN1, supporting a conserved function (Extended Fig. 3a-b). We then hypothesized that

bud formation failure in *ccn1l1* knockdown embryos results from loss of HA and Versican

synthesis. Consistent with this, *has3* and *vcanb* colocalized with *ccn1l1* at canal-genesis

zones (Extended Fig. 3c) and exhibited nearly identical temporal expression patterns—

both showing low expression during bud initiation, peaking during bud extension, and

returning to baseline during bud fusion, as previously reported[9,10,14] (Extended Fig. 3d).

Indeed, *ccn1l1* knockdown reduced *has3* and *vcanb* expression to background levels at

the pre-budding stage and by 50.1% and 73.5%, respectively, at bud extension (Fig. 3c-

d and Extended Fig. 3e-f). We conclude that Ccn1l1 drives bud initiation and extension

by promoting *has3* and *vcanb* transcription.

Given the HA-rich bud-ECM's role in Yap activation and *ccn1l1*'s role in HA production,

we hypothesized the existence of a positive feedback loop: *ccn1l1* expression triggers HA

production, generating ECM pressure that activates Yap, which in turn enhances *ccn1l1*

expression. This predicts that disrupting any component of this loop will affect the others.

First, we tested whether disrupting the mechanical force from ECM expansion would

affect gene expression. Consistent with our prediction, HAase treatment during bud

extension suppressed *ccn1l1* expression to pre-budding levels (Fig. 3e-f) and similarly

reduced *has3* expression (Fig. 3g-h). Second, we disrupted transcriptional regulation

using a heat-shock inducible dominant-negative (DN) Yap mutant lacking the TEAD-

binding domain[33], preventing the conversion of mechanical cues into transcriptional

outputs. These mutants showed significantly reduced bud extension and bud-ECM

volume (Fig. 3i-k), confirming that Yap's transcriptional activity is required for sustained

ECM production and morphogenesis.

Together, these complementary approaches—disrupting the mechanical arm with HAase

and blocking the transcriptional arm with dominant-negative Yap—demonstrate a positive

feedback loop where Ccn1l1 promotes ECM synthesis, ECM expansion activates Yap,

and Yap enhances *ccn1l1* expression (Fig. 3l). This creates a self-amplifying system that

can drive ECM expansion and bud extension from initially small signals. We next

examined how this feedback loop is spatially patterned.

Mechanical patterning of cells at the bud border sustains extension for timely bud fusion

While cell proliferation and convergence-extension through cell rearrangement are typical behaviors underlying tissue growth during development, our previous study showed that neither is necessary for bud extension[9]. To determine the source of cells contributing to this process, we performed time-lapse imaging and cell tracking. We observed a linear increase in bud cell number (Fig. 4a), driven by bud-adjacent cells moving toward the initiated bud, turning at its border, and becoming incorporated into the extending structure (Fig. 4b). Disruption of ECM expansion with HAase treatment or mechanotransduction with *yap1* MO inhibited this increase in cell number (Fig. 4c-d). This suggests that ECM-expansion through mechanotransduction exerts pulling forces that recruit adjacent cells into the growing bud.

Because recruitment occurs specifically at the bud border, we hypothesized that mechanotransduction cues are spatially patterned rather than uniform. To explore this, we developed a theoretical vertex model of a 2D epithelial section incorporating experimentally established parameters[9]: apical, lateral, and basal cellular tension; BM adhesion at the bud border; HA-rich bud-ECM pressure; and anisotropic cytocinch tension orthogonal to the bud's long axis (Fig. 4e and Extended Fig. 4a-d; see Supplementary Note). The model defines four mechanically distinct regions: bud-adjacent cells (BM-attached), bud border cells (partially BM-attached), budding cells (newly incorporated, BM-detached), and tip cells (leading-edge, BM-detached) (Fig. 4e).

The model indicated that cell recruitment was accompanied by a transient increase in cell aspect ratio (Fig. 4f), with stronger overshoots in border and budding cells compared to tip cells. This spatial pattern was lost when BM attachment or cytocinches were removed (Extended Fig. 4a-d), suggesting that mechanical stress is concentrated at the border cells attached to the BM. To test this, we measured nuclear aspect ratios as a marker of cell deformation[34,35] and found that border cells were indeed more deformed than tip cells (Fig. 4g). Supporting the higher mechanical stress prediction, the Yap reporter showed higher activity in border and budding cells, with notably lower variability in border cells

indicating Yap activation during recruitment (Fig. 2b-b' and Fig. 4h). Notably, *ccn1l1*
expression was slightly lower in border cells than in budding cells, consistent with recent
Yap activation (Fig. 2d). Together, these results demonstrate that Yap activity is spatially
patterned at the bud borders rather than occurring throughout the bud. Consistent with
this,

To test the functional significance of spatially patterned mechanotransduction, we
modeled bud growth dynamics in which ECM production comprised two components: a
constant baseline rate from pre-patterned tip cells and a mechanotransductive
contribution proportional to the number of recruited cells. When mechanotransduction
strength is high, the model predicts accelerated, non-linear ECM volume growth (Fig. 4i).
In contrast, disabling the mechanotransductive contribution, resulted in a slower, more
linear increase in ECM volume consistent with constant ECM production from tip cells
alone (Fig. 4i). Because bud extension in the model depends on ECM volume growth,
reduced ECM production without mechanotransductive contribution led to slower
extension rates (Fig. 4k). As predicted by the model, control embryos exhibited rapid ECM
growth and faster extension compared to *yap1* MO-injected embryos that showed slower
ECM growth and reduced bud extension (Fig. 4j and 4l).

Extending buds eventually meet and fuse to form pillars, completing otic vesicle
remodeling and canal demarcation. We next examined how varying extension rates, set
by ECM produced by tip cells and by the mechanotransductive recruitment of neighboring
cells, determine whether buds achieve fusion. To address this, we generated a phase
diagram from ~500 simulations varying both parameters, with fusion defined as buds
reaching 30 μm in length (Fig. 4m), which corresponds to the average distance required
for opposing buds to make contact across the otic vesicle lumen in wild-type embryos at
60 hpf (Extended Fig. 5a). The simulations revealed a trade-off: when
mechanotransduction strength is low, fusion was delayed and required high tip cell
contribution, whereas with high mechanotransduction, even minimal tip cell production
was sufficient for bud fusion (Fig. 4m and Extended Fig. 4e). To test these predictions
experimentally, we performed dose-dependent MO-knockdowns of *ccn1l1* (Fig. 4n-o and
Extended Fig. 4f), which affects both tip cell and mechanotransductive ECM production.

At lower doses, extension slowed significantly, but buds still fused successfully by 72 hpf
(Fig. 4n-o). At the highest dose, extension dropped below ~95% of normal rates (Fig. 4n)
and buds failed to fuse (Fig. 4o). Thus, mechanotransduction ensures that buds can still
achieve fusion across a wide range of tip cell output, with failure occurring only under the
most severe loss of ECM production.

Our theoretical model and experimental data demonstrate that mechanical deformation
at bud borders activates Yap to drive cell recruitment and ECM expansion, sustaining bud
extension rates that ensure successful fusion even when tip cell ECM production varies.

CREB activation during fusion terminates the positive feedback loop

Intriguingly, *ccn1l1* expression dropped sharply once opposing buds fused (Fig. 2c and
2e). Consistent with this downregulation, both cell recruitment and bud extension
plateaued when bud pairs made contact (Fig. 4a and Extended Fig. 5a), indicating a self-
termination mechanism that stops further bud growth. A likely candidate for this stop
signal is Gpr126 (Adgrg6), a mechanosensitive adhesion-type G-protein-coupled
receptor (aGPCR) essential for bud fusion[14,36,37] and expressed in budding cells
throughout morphogenesis[14] (Extended Fig. 5b). Consistent with previous findings[14], in
gpr126^{st49/st49} mutants, which carry a premature stop codon[38], buds fail to fuse and
undergo excessive elongation, accompanied by persistent *has3* expression (Fig. 5a-c and
Extended Movie 2-3). We found that *ccn1l1* expression remained persistently elevated in
these mutants at both 60 hpf and 72 hpf, contrasting with the normal sharp
downregulation upon fusion (Fig. 5b-c and Extended Fig. 5c) . This persistent expression
drove continued growth: mutant buds were 78.7% longer and contained 76.5% more cells
than wild type at 60 hpf (Fig. 5d-e). Notably, bud-ECM volume continued to grow
unchecked, with a 464.6% increase in volume (Fig. 5f and Extended Fig. 5d), consistent
with the ongoing recruitment of HA-producing cells. This behavior is inconsistent with a
genetically predetermined bud cell number, which would have produced a plateau in
extension. Instead, this provides further evidence that mechanotransduction recruits new
cells to the bud. The 2D vertex model recapitulated this behavior: when fusion was
blocked and the mechanotransductive contribution proportional to recruited cells allowed

to persist, ECM volume expanded unchecked (Extended Fig. 5e). These results
demonstrate that bud fusion provides the essential termination signal for the positive
feedback loop; in its absence, mechanotransductive recruitment drives runaway growth.

We next investigated how Gpr126 terminates *ccn1l1* expression. Gpr126 activates the
cyclic adenosine monophosphate (cAMP) pathway[39,40]. Pharmacological cAMP elevation
with forskolin[41] strongly reduced both *ccn1l1* and *has3* expression (Extended Fig. 5f-g)
and inhibited bud elongation (Extended Fig. 5h-i). Since cAMP signals via Protein Kinase
A (PKA), leading to the phosphorylation of the transcription factor CREB[42], we examined
CREB activation during fusion. Immunostaining revealed striking CREB phosphorylation
at bud-bud contact sites, spreading into budding cells that form the pillar (Fig. 5g). This
CREB phosphorylation was absent in *gpr126* mutants (Fig. 5h), demonstrating that
Gpr126 signaling mediates the phosphorylation of CREB.

CREB has been reported to downregulate CCN gene transcription in cultured cells[43,44].
To confirm the role of CREB in the inhibition of *ccn1l1*, we expressed DN-PKA lacking the
cAMP-binding site[45]. This attenuated CREB phosphorylation and prevented both bud
fusion and *ccn1l1* downregulation (Fig. 5i-j and Extended Fig. 5j-k), demonstrating that
PKA-CREB signaling is necessary for feedback termination.

Together, these results demonstrate that the Gpr126-cAMP-PKA-CREB pathway
suppresses *ccn1l1* and terminates ECM synthesis during bud fusion. This creates a
morphological transition-triggered, self-limiting mechanism where the successful
completion of morphogenesis (fusion) terminates the positive feedback loop.

Discussion

Our findings reveal how robust SCC morphogenesis emerges from a self-limiting positive
feedback mechanism where morphological success triggers its own termination. We
demonstrate that the mechanotransducer Yap, activated by a HA-rich ECM, drives the
expression of the matricellular protein Ccn1l1, which further promotes HA and Versican
synthesis to sustain ECM expansion and bud extension. This positive feedback loop is
spatially restricted, with Yap activity highest at the bud border. Critically, the Gpr126-

cAMP-PKA-CREB pathway is activated during bud fusion and terminates *ccn1l1*
expression, preventing runaway growth. Thus, the completion of morphogenesis serves
as the signal to terminate the very process that drives it.

Most studies of Yap/Taz mechanotransduction have emphasized their role in negative
feedback loops that maintain tissue homeostasis, where mechanical forces activate
Yap/Taz to promote behaviors that ultimately dissipate those same forces[26,46–48]. More
recently, however, it has become clear that Yap/Taz can also participate in positive
feedback loops[49] to amplify mechanical cues rather than inhibiting them. Here we show
that Yap activation promotes ECM synthesis, which generates mechanical forces that
further sustain Yap activtity to generate self-reinforcing growth dynamics. We propose
that this positive feedback mechanism may be important during developmental windows
where weak initial signals must be robustly amplified to drive rapid morphogenetic
outcomes.

The identification of Ccn1l1 as a key mediator linking Yap activation to ECM production
represents a significant advance in understanding how CCN proteins contribute to
mechanotransduction networks. While CCN family proteins have been implicated in
developmental processes, wound healing, and inflammation[19–21], their specific role in
creating positive feedback loops between mechanical forces and matrix synthesis has not
been previously demonstrated. The two-stage regulation of Ccn1l1—initial Lmx1b-
dependent expression followed by Yap-dependent amplification—suggests an elegant
mechanism for transitioning from developmental patterning to mechanically-driven growth
amplification. Notably, *ccn1l1* is also expressed in other organs, including somite and
notochord[19], raising the possibility that similar feedback-driven mechanisms operate more
broadly during vertebrate development.

CCN proteins are known to interact with integrins to regulate ECM production[30–32], and
among them, β1 integrin is a well-documented receptor for CCN family members[24,31,50].
β1 integrin expression has indeed been reported in the OV of both zebrafish and mice[51,52],
suggesting that *ccn1l1* may act via integrin to promote ECM production during SCC
formation. Beyond ECM regulation, CCN proteins can also remodel the actomyosin

cytoskeleton through integrin-Rho GTPase signaling, thereby promoting cellular

protrusions and generating mechanical tension[24]. We previously reported that

cytocinches progressively increase during bud extension and are essential for longitudinal

growth[9], coinciding with the rise in *ccn1l1* expression. These observations raise the

possibility that *ccn1l1* not only promotes ECM expansion, but may also contribute to bud

extension by modulating cytoskeletal tension.

Our theoretical model predicts that BM attachment is crucial for spatially restricting Yap

activation to border cells, yet this prediction remains to be investigated. Integrins,

particularly those that bind laminin components of the BM[53], are prime candidates for

transducing mechanical forces at these sites. Focal adhesion kinase (FAK) and related

signaling molecules, such as vinculin, likely serve as mechanosensitive intermediates that

link integrin engagement to Yap activation[7]. The precise mechanical threshold required

for Yap activation, and how BM composition and stiffness modulate this threshold,

represents an important avenue for future investigation. Understanding these molecular

details could reveal how tissues fine-tune their mechanosensitivity during morphogenesis.

Adhesion GPCRs (aGPCRs) perform diverse developmental roles in many organs like

heart, sciatic nerve, cartilage, and vascular systems, and their dysfunction can lead to

organ malformation or embryonic lethality[36,54–57]. Previous work implicated Gpr126 in

downregulating ECM production during SCC formation[14], but the molecular mechanism

was unknown. We find that Gpr126 signaling is activated during fusion and terminates

ECM production by downregulating *ccn1l1*. Such termination mechanisms may represent

a conserved principle by which aGPCRs couple tissue remodeling to signaling and

transcriptional outcomes during morphogenesis. Although the precise mode of Gpr126

activation remains unclear, it likely involves ligand binding and/or mechanical forces that

expose a tethered agonist (Stachel sequence) to trigger downstream signaling[58–61].

The integration of mechanotransduction feedback loops to morphological outcomes

provides a framework for understanding how complex organ shapes emerge through self-

organizing principles. Overall, we demonstrate how robust SCC morphogenesis emerges

from an interplay of two evolutionarily conserved pathways. The mechano-transduction

feedback loop, Yap-Ccn1l1-HA, drives bud extension, and the Gpr126-cAMP-PKA-CREB

pathway couples bud fusion to extension termination. This principle likely applies broadly

to developmental systems where patterned growth-promoting signals must first be

amplified and then precisely shut down.

Materials and Methods

Animals

Zebrafish (*Danio rerio*) EK and AB wild-type strains were used in this study. Adult fish

were kept on a 14-hour light/10-hour dark cycle. Embryos were collected by crossing

female and male adults (3-18 months old) and raised in 28.5 °C egg water. In this study,

the following transgenic and mutant lines were used: *Tg(actb2:membrane-neongreen-*

neongreen)[9]*, lmx1bb* [ji410/ji410] [38], *Tg(hsp70I: RFP-DN-Yap1)* (this study), *Tg(hsp70I: DN-*

PKA-GFP) (this study), *gpr126* [st49/st49] [38], *TgKI(Yap1-mScarlet)* [pd1280] (this study), and

Tg(βActin: utrophin-mCherry)[62]. *gpr126* [st49/st49] mutant fish were maintained as an AB

strain background, whereas all other transgenic and mutant lines were maintained as an

EK strain background.

Morpholino injection

Standard control MOs and MOs targeting *ccn1l1* (5′-

CTCAGAGGAGACATGGTGTTATCCA-3′, translational blocking), *yap1* (5′-

AGCAACATTAACAACTCACTTTAGG-3′, splice-modifying between intron 2 and exon 2)

[15], and *p53* (5′-GCGCCATTGCTTTGCAAGAATTG-3′, translational blocking)[63] were

purchased from Gene Tools. The *ccn1l1* MO *target sequence* was designed based on

the Danio rerio *ccn1l1* mRNA sequence (ENSDART00000051336), retrieved from the

Ensembl database, using the Gene Tools Oligo design service. For the *ccn1l1* knockdown,

embryos at the single-cell stage were injected with 2 ng of *ccn1l1* MO or control MO,

together with 1 ng of *p53* MO. For the *yap1* knockdown, 5 ng of *yap1* MO was injected

with 2.5 ng of *p53* MO into embryos at the single-cell stage. MO-injected embryos were

screened for inner ear phenotype at 40-50 hpf and used for confocal live imaging, HCR-

FISH, or immunostaining at 40-72 hpf. To test the dose-dependent effect of *ccn1l1* MO,

embryos at the single-cell stage were injected with either standard control MO (2 ng) or
ccn1l1 MO (0.5, 1, or 2 ng), together with *p53* MO. Injected embryos were screened for
inner ear phenotype at 50 hpf, followed by confocal live imaging at 50, 56, and 72 hpf.
p53 MO was co-injected with control, *ccn1l1* or *yap1* MO to reduce MO-injection induced
apoptosis and non-specific toxity.

Genome editing by CRISPR-Cas9

ccn1l1 CRISPANT embryo

Single-guide RNAs (sgRNAs) were synthesized as previously reported[10]. Briefly, sgRNAs
targeting multiple coding regions of *ccn1l1* were synthesized using MEGAscript SP6
Transcription Kit (Invitrogen, AM1330) and purified with RNA Clean & Concentrator kit
(Zymo Research, R1018). For the knockdown of *ccn1l1*, three sgRNAs were pooled
together and injected into embryos as previously described[10]. Briefly, embryos at the
single-cell stage were injected with 2 nL of injection solution containing pooled sgRNAs
(200 ng/μL: 66.6 ng/μL of each sgRNA) and Cas9 protein (PNA Bio, CP-03, 500 ng/μL).
F0 crispant embryos were screened for inner ear phenotype at 50-72 hpf and then used
for HCR-FISH or confocal live imaging. The following target sites were used to generate
sgRNAs: *ccn1l1* sgRNA1: 5'-CGGCGGGCACAGCGCCAGTCGGG-3'; *ccn1l1* sgRNA2:
5'-GTGTGTCCTCAGGTGCAGGGAGG-3';   *ccn1l1*   sgRNA3:   5'-
AGGTGGCTGGAGCGATAGGGCGG-3'

tyrosinase CRISPANT embryo

A sgRNA targeting the *tyrosinase* gene[64] was synthesized as described above. sgRNA
(200 ng/μL) and Cas9 protein (500 ng/μL) were injected into embryos at the one-cell stage
as described above. The *tyrosinase* crispant embryos were screened for pigmentation
loss at 50-72 hpf and used for HCR-FISH or confocal live imaging. *tyrosinase* crispant
embryos were used as an sgRNA injection control, resulting in loss of pigmentation
without affecting SCC formation[10]. The following target site was used to generate the
sgRNA: 5'-GGACTGGAGGACTTCTGGGG-3'.

Generation of *TgKI(Yap1-mScarlet)* line

The *TgKI(Yap1-mScarlet)* [pd1280] line was generated using a previously reported knock-in

tagging method[13]. As the targeting cassette, a gene fragment spanning from the middle

of intron 7 to the last codon of exon 8 was cloned into pUC19-TgKI-MCS-mScarlet-MCS-

polyA (Addgene plasmid #174023) using In-Fusion seamless cloning (Takara Bio) with

the   following   primers: *yap1*   Fw: 5′-

TTCGCTAGCCCGCGGCTATCAGTGCTTTACAGATCATTGTC-3′; *yap1* Rev: 5′-

CACCTCGAGTCTAGATAGCCAGGTTAGAAAGTTCTCC-3′. The sgRNA target sites for

genomic editing were mutated in the donor plasmid using Q5 site-directed mutagenesis

(New England Biolabs, NEB) with the following primers: *yap1* SDM Fw: 5′-

ACATCCTCAACGACATGGAGTCAGTGCTGGC-3′; *yap1* SDM Rev: 5′-

CGGAGCTCAAAGCTTCCTGCAGGCTGGGCA-3′. A sgRNA targeting *yap1* (5′-

GTTGAGGATGTCGGAGCTTAGGG-3′) was synthesized using the HiScribe T7 Quick

High Yield RNA Synthesis Kit (NEB) and purified with the Monarch RNA Cleanup Kit

(NEB). The injection mix contained PCR donor (20 nM), gRNA (4 µM), Cas9 protein (3

µM; PNA Bio CP-02), and DMSO (1%). The genomic integration site was confirmed by

sequencing at the F2 generation, and the line was designated allele number pd1280.

Construct the generation and injection of plasmid and mRNA

actb2:H2B-EGFP plasmid

H2B-EGFP was amplified by PCR using Phusion Flash High-Fidelity PCR Master Mix

(Thermo Fisher Scientific, F548L) and appropriate primers from a construct harboring its

coding sequence. pMTB *actb2:membrane-neongreen-neongreen* construct[9], which

contains the SP6 promoter for mRNA synthesis, was used as the plasmid backbone. The

plasmid backbone was digested with BamHI and AgeI. Then, the amplified H2B-EGFP

fragment was cloned downstream of the actb2 promoter and the SP6 promoter in the

pMTB backbone using In-Fusion Snap Assembly Master Mix (Takara, 638948). The H2B-

EGFP mRNA was synthesized using the mMESSAGE mMACHINE SP6 transcription kit

(Invitrogen, AM1340) from the linearized *actb2:H2B-EGFP* plasmid. 2nL of mRNA (100

ng/µL) was injected into single-cell-stage embryos from *TgKI(Yap1-mScarlet)* parents.

Live embryos were observed at 40-56 hpf by confocal microscopy. The following primers

were used for PCR: Fw: 5'-

CTTGTTCTTTTTGCAGGATCCCGCCACCATGCCAGAGCCAGCGA-3'; Rev: 5'-

TTATCATGTCTGGATCACCGGTTTACTTGTACAGCTCGTCCATGC-3'.

hsp70l: RFP-dominant negative Yap1 plasmid

hsp70l: RFP-DN-Yap1[33] construct harboring the I-Sce I site for the transgene was a gift

from Kenneth Poss (Morgridge Institute, University of Wisconsin-Madison, USA) and was

used as a template for cloning. RFP-DN-Yap1 was amplified by PCR using appropriate

primers. The plasmid backbone, which contains a heat-shock-inducible promoter (hsp70l)

and Tol2 sites, was digested with BamH1 and AgeI and then gel-purified. The amplified

DNA fragment was cloned downstream of the hsp70l promoter in the plasmid backbone

using the In-Fusion system. To generate the *Tg(hsp70l: RFP-DN-Yap1)* line, 100 pg of

Tol2 mRNA and 50 pg of hsp70l: RFP-dnYap1 plasmid were injected into embryos at the

one-cell stage. The following primers were used for PCR: Fw: 5'-

TTGCAGGATCCGCCACCATGGACAACACCGAGGACG-3'; Rev: 5'-

TGGATCACCGGTTCAGATACTACTGTGGGCTGGAGGG-3'

hsp70l: dominant negative PKA-GFP plasmid

M7 pdnPKA-GFP was a gift from Randall Moon (Addgene plasmid # 16716), used as a

template for cloning. The cloning was performed, as described above. Briefly, the

amplified dnPKA-GFP fragment was cloned downstream of the hsp70l promoter in the

plasmid backbone. To generate the *Tg(hsp70l: DN-PKA-GFP)* line, 100 pg of Tol2 mRNA

and 50 pg of hsp70l: DN-PKA-GFP plasmid were injected into embryos at the one-cell

stage. The following primers were used for PCR: Fw: 5'-

TACAGTTCAGCCATCCTAGGATTTAGGTGACACTATAGAATACAAGCTACT-3'; Rev:

5'-GTTCCGGATACGTATGAATTCTTATTTGTATAGTTCATCCATGCCATGT-3'

Drug treatment

For hyaluronidase treatment, we injected hyaluronidase (HAase, Sigma Aldrich, H1136) into the periotic space of dechorionated embryos, with slight modifications to a previously reported method[9]. Briefly, HAase (500 units/mL) was injected into embryos twice at a 3-hour interval from 46 hpf, followed by HCR-FISH or confocal live imaging. For live imaging of *TgKI(Yap1-mScarlet)* embryos, HAase was injected into embryos at 50 hpf, followed by imaging at 1.5 hours post-injection. For immunostaining of Yap1 antibody, HAase was injected into embryos at 50 hpf, and embryos were fixed with 4% PFA in PBS at 3 hours post-injection, followed by immunostaining. For RO4929097 (ApexBio, A4005) treatment, dechorionated embryos at 24 hpf were soaked in egg water containing 50 μM inhibitor for 26 hours, followed by HCR-FISH. For forskolin (Cayman Chemical, 1101810) treatment, dechorionated embryos were soaked in egg water containing 50 μM of the drug at 46 hpf for 6 hours, followed by HCR-FISH or confocal live imaging.

Multiplex *in situ* hybridization chain reaction (HCR)

HCR probes for each gene were obtained from IDT (Oligo Pools, oPools). Hybridization buffer, amplification buffer, wash buffer, and amplifiers were purchased from Molecular Instruments. mRNA sequences used for HCR probe design were retrieved from the NCBI database. HCR probe sets for each mRNA were designed using a home-made Python script[29]. 40 probe oligonucleotides targeting each mRNA were generated and pooled oligonucleotides were used for HCR-FISH. HCR probe sequences are listed in Extended Table 1. HCR-FISH experiments were performed as previously described[9,10]. Dechorionated embryos at various stages were fixed with 4% paraformaldehyde (PFA) in PBS and permeabilized with prechilled acetone. The embryos were then incubated with the hybridization buffer at 37°C for 30 min, followed by incubation with the hybridization solution containing probes overnight at 37°C. After probe hybridization, the embryos were incubated overnight at room temperature in an amplification buffer with fluorescent HCR amplifiers (Alexa-488, Alexa-546, or Alexa-647). Embryos were washed with 0.1% Tween-20 in 5× SSC buffer (four times) and stored at 4°C until confocal imaging. Control and experimental samples were processed within the same tube to ensure uniform

520   treatment. This included the same incubation times, reagent concentrations, and handling

521   procedures, ensuring that any differences in fluorescence intensity were due to variations

522   in target abundance rather than experimental variability.

Immunohistochemistry

Immunohistochemistry was performed as previously described[10]. Briefly, for phospho-

CREB staining, the dechorionated embryos were fixed with 4% PFA/PBS for 2 hours at

room temperature, followed by permeabilization with pre-chilled acetone at -20 °C for 7

minutes. Embryos were then blocked with 5% bovine serum albumin (BSA; Sigma-Aldrich,

A3311) in 1× PBS containing 0.1% Tween-20 (PBT) at room temperature for 1 h, followed

by incubation in primary antibody solution containing rabbit anti-phospho-CREB (Ser133)

antibody (Cell Signaling Technology, 9198; 1:100) and mouse anti-GFP antibody

(Invitrogen, A11120; 1:500) at 4°C overnight. Embryos were washed with 1x PBT for 5

min, thrice, then incubated with a secondary antibody solution containing fluorescently

labeled anti-rabbit and anti-mouse antibodies (Thermo Fisher Scientific, A11008, A11035,

A21244, and A11001; 1:200) and phalloidin (Thermo Fisher Scientific, A22287; 1:100) at

4°C overnight. Embryos were washed with 1x PBT for 5 min, thrice, and stored at 4°C.

For Yap1 staining, embryos fixed with 4% PFA/PBS were permeabilized with 2% Triton

X-100/PBS for 2 hours. Embryos were then blocked with 2% BSA in 1× PBS containing

1% DMSO and 0.1% Triton X-100 (PBDT) for 2 hours at room temperature, followed by

incubation in a primary antibody solution containing rabbit anti-Yap1 (Proteintech, 13584-

1-AP; 1:200) at 4 °C overnight. Embryos were washed with 1× PBDT for 15 min, thrice,

then incubated with a secondary antibody solution containing fluorescently labeled anti-

rabbit antibody (1:200) and Hoechst 33342 (Thermo Fisher Scientific, 62249; 1:2000) at

4°C overnight. Following this, embryos were washed with 1× PBDT for 15 minutes, thrice.

Confocal imaging

Confocal imaging was performed with a Zeiss LSM 980 confocal microscope using a C-

Apochromat 40×1.2 NA objective for all fluorescence acquisitions as previously

described[10]. Briefly, dechorionated live embryos were anesthetized in 1xtricaine and then

mounted dorsolaterally using a canyon mount cast in 1.5% agarose dissolved in egg

water and covered with a coverslip. Images of *TgKI(Yap1-mScarlet)* embryos were

acquired using Airyscan SR mode, followed by image processing using the Zen software

(Zeiss). For imaging of embryos stained with HCR-FISH or immunohistochemistry,

embryos were mounted on 1.5% agarose dissolved in 1x PBS and then subjected to

confocal imaging. Image acquisition settings, such as laser power, gain, and exposure,

were kept constant and within the sub-saturation range during microscopy to allow for

accurate and quantitative sample comparisons. Time-lapse imaging was performed as

previously described[9]. Briefly, embryos were immobilized by injecting 500 µM α-

bungarotoxin protein in PBS (Thermo Fisher Scientific, B35450) into the heart 30 min

before imaging. The embryos were maintained in a custom-built incubator at 28.5°C

during time-lapse imaging. Z-stack images were acquired at 30-minute intervals.

FluoRender, an open-source software, was used for 3D renderings of images.

Screening and genotyping of mutants

gpr126 $^{st49/49}$ mutants, derived from *gpr126* $^{st49/+}$ parents, were screened for

immunohistochemistry based on inner ear phenotypes at 60 hpf[14]. For Fig. 5a–f and

Extended Fig. 5d, EK wild-type and *gpr126* $^{st49/49}$ mutants were obtained from EK wild-

type and *gpr126* $^{st49/49}$ parents, respectively, and subsequently subjected to HCR-FISH

and confocal live imaging at 50-72 hpf. To minimize developmental differences between

mutant and wild-type embryos, male and female fish were maintained in a mating tank

separated by a removable divider overnight, and fertilized eggs were collected within 15

minutes after the divider was removed the following morning. *lmx1bb* $^{jj410/+}$ mutants were

genotyped using the derived cleaved amplified polymorphic sequence (dCAPS) method,

as previously described[10]. The *lmx1bb* $^{jj410/jj410}$ mutant obtained from *lmx1bb* $^{jj410/+}$ parents

was screened based on inner ear phenotypes at 50 hpf, followed by HCR-FISH.

Heat-shock activation of hsp70l promoter

For experiments using DN-Yap1 mutants, *Tg(Hsp70l: RFP-DN-Yap1)* fish were crossed

with *Tg(actb2:membrane-neongreen-neongreen)* fish. Embryos obtained from this cross

were heat-shocked at 37°C for 30 min twice at 5-hour intervals at 40 hpf, followed by
confocal live imaging at 50 hpf. DN-Yap1 mutants were identified by RFP fluorescence at
5 hours post-heat shock. RFP-negative siblings were used as controls for confocal
imaging. For experiments using DN-PKA mutants, *Tg(Hsp70l: DN-PKA-GFP)* fish were
crossed with wild-type or *Tg(βActin:utrophin-mCherry)* fish. To induce acute activation of
the heat-shock promoter before bud fusion, embryos were heat-shocked at 42°C for 10
min at 48 hpf, followed by HCR-FISH or confocal live imaging at 56 hpf. DN-PKA mutants
were identified by GFP fluorescence at 3 hours post-heat shock. GFP-negative siblings
were used as controls for HCR-FISH or confocal live imaging. Heat-shock treatment at
42°C for 10 min was confirmed not to noticeably affect zebrafish embryonic development.

Modeling

See Theory Supplementary Note

Analysis of single-cell RNA sequencing data

Single-cell RNA sequencing analysis was performed using
Daniocell2023_SeuratV4.rds[17,18] available at https://daniocell.nichd.nih.gov/. The
published Seurat object was processed in R (v 4.4.0) using Seurat (v5.2.1)[65]. The R script
of this analysis can be obtained at GitHub (https://github.com/MunjalLab/ccn1l1_project).
Clusters of otic tissue cells from 36 hpf to 72 hpf were isolated from the original Daniocell
atlas. 3000 highly variable genes were selected using the Seurat function
'FindVariableFeatures' with the method 'vst'. All genes were then scaled to have a mean
expression of 0 and a variance of 1 across all selected cells. Clustering was performed
using the Louvain algorithm[66] implemented in FindClusters with a resolution of 0.6. The
canal-forming cell cluster was identified based on expression of *vcanb*, a known marker
for this cell type, supported by previous literature[10,14]. The top 10 differentially expressed
genes were identified by the 'FindAllMarkers' function in Seurat.

Comparative analysis of Ccn1l1 and Ccn1 amino acid sequences

For the analysis of the functional domains of zebrafish Ccn1l1, amino acid sequences

were retrieved from the NCBI database. Functional domains were predicted using the

SMART database[67,68], and the significance (E-value) of each predicted domain was

recorded. To compare domain conservation across species, multiple sequence

alignments were performed using Clustal Omega[69], and sequence similarity and percent

identity were assessed. Clustal alignment outputs were used to visualize conserved

residues, and the pairwise identity matrix was used to quantify percent identity between

species. Amino acid sequences of each functional domain predicted by SMART are listed

in Extended Table 2.

Analysis of the TEAD-binding domain in *ccn1l1* 5' upstream region

For this analysis, genomic DNA was extracted from EK wild-type embryos at 2 dpf, and

a 7-kb fragment of the *ccn1l1* 5′ upstream region was amplified by PCR. The sequence

of the PCR product was determined by Sanger sequencing and analyzed for the presence

of the TEAD-binding motif (GGAATG/CATTCC)[22,26–28]. The following primers were used

for   PCR: forward,   5′-   TATGTCAAACACACAAGCAGTGG-3′;   reverse,   5′-

GGTGTTATCCAAAAAATTGTTAAGAGGC-3′.

Quantification and statistical analysis

1) Bud length measurement

For quantification, images were acquired by confocal live imaging of embryos expressing

membrane-NeonGreen, with z-stacks collected at 0.5 or 1 µm intervals. 2D images

capturing the middle section of the bud were used to measure the bud length of embryos.

Bud lengths were measured manually using the "straight-line" tool in Fiji. Where budding

had not started or in perturbations where budding was affected, bud lengths correspond

to average cell height at the anterior or posterior region. For time-lapse analysis, the same

buds were tracked every 30 min, and the bud length was measured at each time point.

For the anterior bud length of *gpr126[st49/st49]* mutants or the anterior pillar of WT embryos

(Fig. 5), bud length was measured by Imaris software. Using the spot tool, spots were

placed along the centerline of the bud, and the total length was calculated by connecting
these spots.

2) Cell and nuclear aspect ratio

The cell aspect ratio of budding cells was measured using the "straight-line" tool in Fiji. Z-
stack images of the OV in embryos expressing membrane-NeonGreen were obtained by
confocal live imaging at 0.5 µm intervals. 2D images capturing the middle section of
budding cells were used to measure cell height and width, as shown in Extended Fig. 1a.
For time-lapse analysis, the same cells were tracked every 30 min, and the cell aspect
ratio was measured at each time point. To measure the nuclear aspect ratio, embryos
were injected with H2B-GFP mRNA, followed by confocal live imaging at 50 hpf. Z-stack
images of anterior buds were acquired at 0.5 µm intervals. The nuclear aspect ratio was
measured using Imaris. Using the Surface tool, masks delineating the H2B-GFP signal
were generated at 1 µm slice intervals in the 3D images and subsequently merged to
reconstruct an ellipsoidal 3D nuclear structure. The lengths of the long and short axes of
the ellipsoid were quantified, and the nuclear aspect ratio was calculated.

3) Counting cell numbers in a bud

Cell counting was performed using the Mastodon plugin (https://github.com/mastodon-
sc/mastodon) in Fiji, which enables 3D image analysis. 3D z-stack images of the otic
vesicle in embryos expressing membrane-NeonGreen were acquired by confocal live
imaging at 0.5 or 1 µm slice intervals. Using the Big Data Viewer in the Mastodon plugin,
all cells in a bud were manually marked in 3D and counted. For time-lapse analysis, the
same buds were tracked every 30 min, and the cell number was counted at each time
point.

4) Bud-ECM volume measurement

The bud-ECM region was defined as shown in Extended Fig. 1a. 3D z-stack images of
the otic vesicle in embryos expressing membrane-NeonGreen were acquired by confocal
live imaging at 1 µm slice intervals. Bud-ECM volume was measured using Imaris

software. Using the Surface tool, masks delineating the bud-ECM area were generated

at 1 μm slice intervals in the 3D images and subsequently merged to reconstruct a 3D

surface filling the bud-ECM region, and the volume was quantified. For time-lapse

analysis, the same buds were tracked every 30 min, and the volume of bud-ECM region

was measured at each time point.

5) Intensity analysis of multiplex in situ HCR (HCR-FISH)

Measurement of the fluorescence intensity of the HCR probes was performed as

previously described[10]. In brief, using Fiji, maximum intensity z-projections covering the

entire anterior or posterior bud were created. To measure the mean intensity of HCR

probes, the ROI was manually drawn along the outer cells of the bud with a segmented

line" tool. To create a histogram plot of fluorescence intensity, regions of interest (ROIs)

were manually drawn using the "segmented line" tool (10 pixels wide) along the anterior

budding region, spanning 55-65 μm, as shown in Fig. 2. An additional 20 μm on each side

of the non-budding region flanking the budding cells was also included. The "plot profile"

tool was then used to obtain an average intensity trace along the line. Fluorescence

intensities were divided into 120 and 40 bins for the budding and shoulder regions,

respectively, expressed as normalized distances, and subsequently normalized again to

the overall maximum value. The Fire heatmap was used to represent the fluorescence

intensities of individual probes. The heatmap color scale was adjusted individually for

each experiment.

6) Intensity analysis of the Yap1 nuclear-cytoplasm ratio

For measurement of the Yap1 nuclear-to-cytoplasmic ratio in immunostained samples,

nuclear and cytoplasmic regions were manually segmented based on Hoechst and

phalloidin staining, respectively, using Fiji software, and the ratio of mean nuclear to mean

cytoplasmic signal intensities was calculated. For measurement using the *TgKI(Yap1-*

mScarlet) line, nuclei were manually segmented based on the H2B-GFP signal, and the

mean nuclear Yap1 signal intensity was quantified. The mean cytoplasmic Yap1 signal

intensity was obtained using the "oval" tool in Fiji by placing four circular ROIs in the

cytoplasm and averaging their mean intensities. The Yap1 nuclear-to-cytoplasmic ratio

was then calculated as described above.

Statistical analysis

Average values were calculated from 'n,' where 'n' represents the number of buds, bud-

ECM, and cells (as labeled in each figure). Plots show individual data points, mean values,

and error bars (standard deviation of the mean, as labeled). Before performing parametric

statistical tests, the Kolmogorov-Smirnov and Shapiro-Wilk tests were used to test if the

dataset in each experiment was from a normal distribution. To compare the difference

between the two groups, unpaired or paired two-tailed Student's t-tests were performed.

For multiple comparisons, one-way ANOVA with Tukey's test for normally distributed data

and Kruskal-Wallis test with Dunn's test for non-normally distributed data were performed.

P-values less than 0.05 were considered to be significant. All statistical tests and plots

were made in GraphPad Prism 10.

Acknowledgments

We are thankful to all other members of the Principles of Tissue Morphogenesis (Munjal)

Lab for their feedback and support. We thank Brigid Hogan and Stefano Di Talia for their

critical comments on the manuscript. We thank members of the Poss and Bagnat labs for

valuable technical input during this project. We thank Sean Megason and Ian Swinburne

for their guidance and support during AM's postdoctoral training. We thank Kazunori Ando

for technical support and advice to YM. We thank Toru Kawanishi and Sean Megason for

sharing the Yap immunostaining protocol. We thank Tony Tsai's lab for sharing the *Yap*

(4xGTIIC:EGFP) reporter, which was not included in the final manuscript. We thank

Nanami Sato for illustrations. We thank the Duke Zebrafish Core Facility of the Duke

School of Medicine for their support. This work was funded by the National Institute of

Child Health and Human Development (R00HD098918 and DP2115157 to A.M.) and the

National Institute of Arthritis and Musculoskeletal and Skin Diseases (R01AR083346 to

M.B). YM is supported by JSPS Overseas Research Fellowship, Duke Regeneration

Centre Career Advancement Award, and Uehara Memorial Foundation Overseas
Research Fellowship.

Author contributions

A.M. conceived and initiated the project. Y.M., with input from A.M., subsequently
developed the project direction. Y.M. designed, performed, and analyzed most of the
experiments with the assistance of A.M., A.B.B., and K.H. P.R. developed the theoretical
model with the supervision of E.H. Y.M. generated the *Tg(hsp70l: RFP-DN-Yap1)* line,
Tg(hsp70l: DN-PKA-GFP) line, and all constructs for the project, except those required
for the *TgKI(Yap-mScarlet)* line. D.S.L. and M.B. generated the *TgKI(Yap-mScarlet)* line
and the corresponding construct. J.W. analyzed the publicly available transcriptomics
dataset. Y.M. and P.R. prepared figures with the input from E.H. and A.M. Y.M. and A.M.
interpreted the data and wrote the original manuscript with input from P.R. and E.H. All
authors reviewed and edited the manuscript. A.M. acquired funding for the project. A.M.
supervised and administered the project.

Declaration of interests

726 The authors declare no competing interests.

References

1. Hannezo, E., and Heisenberg, C.-P. (2019). Mechanochemical Feedback Loops in
Development and Disease. Cell *178*, 12–25.
https://doi.org/10.1016/j.cell.2019.05.052.

2. Collinet, C., and Lecuit, T. (2021). Programmed and self-organized flow of
information during morphogenesis. Nat. Rev. Mol. Cell Biol. *22*, 245–265.
https://doi.org/10.1038/s41580-020-00318-6.

3. Schier, A.F. (2003). Nodal signaling in vertebrate development. Annu. Rev. Cell Dev.
Biol. *19*, 589–621. https://doi.org/10.1146/annurev.cellbio.19.041603.094522.

4. Moeller, M.E., Danielsen, E.T., Herder, R., O'Connor, M.B., and Rewitz, K.F. (2013).
Dynamic feedback circuits function as a switch for shaping a maturation-inducing
steroid pulse in Drosophila. Development *140*, 4730–4739.
https://doi.org/10.1242/dev.099739.

5. Vogel, V., and Sheetz, M. (2006). Integrins and Extracellular Matrix in
Mechanotransduction Martin Alexander Schwartz 3. Nat Rev Mol Cell Biol *7*, 265–
275.

6. Di, X., Gao, X., Peng, L., Ai, J., Jin, X., Qi, S., Li, H., Wang, K., and Luo, D. (2023).
Cellular mechanotransduction in health and diseases: from molecular mechanism to
therapeutic targets. Signal Transduct. Target. Ther. *8*, 282.
https://doi.org/10.1038/s41392-023-01501-9.

7. Dupont, S., Morsut, L., Aragona, M., Enzo, E., Giulitti, S., Cordenonsi, M., Zanconato,
F., Le Digabel, J., Forcato, M., Bicciato, S., et al. (2011). Role of YAP/TAZ in
mechanotransduction. Nature *474*, 179–183. https://doi.org/10.1038/nature10137.

8. Totaro, A., Panciera, T., and Piccolo, S. (2018). YAP/TAZ upstream signals and
downstream responses. Nat. Cell Biol. *20*, 888–899. https://doi.org/10.1038/s41556-
018-0142-z.

9. Munjal, A., Hannezo, E., Tsai, T.Y.-C., Mitchison, T.J., and Megason, S.G. (2021).
Extracellular hyaluronate pressure shaped by cellular tethers drives tissue
morphogenesis. Cell *184*, 6313-6325.e18. https://doi.org/10.1016/j.cell.2021.11.025.

10. Mori, Y., Smith, S., Wang, J., Eliora, N., Heikes, K.L., and Munjal, A. (2025). Versican
controlled by Lmx1b regulates hyaluronate density and hydration for semicircular
canal morphogenesis. Development *152*, dev203003.
https://doi.org/10.1242/dev.203003.

11. Whitfield, T.T. (2015). Development of the inner ear. Curr. Opin. Genet. Dev. *32*,
112–118. https://doi.org/10.1016/j.gde.2015.02.006.

12. Haddon, C.M., and Lewis, J.H. (1991). Hyaluronan as a propellant for epithelial
movement: the development of semicircular canals in the inner ear of Xenopus.
Development *112*, 541–550. https://doi.org/10.1242/dev.112.2.541.

13. Levic, D.S., Yamaguchi, N., Wang, S., Knaut, H., and Bagnat, M. (2021). Knock-in
tagging in zebrafish facilitated by insertion into non-coding regions. Development *148*.
https://doi.org/10.1242/dev.199994.

14. Geng, F.-S., Abbas, L., Baxendale, S., Holdsworth, C.J., Swanson, A.G., Slanchev,
K., Hammerschmidt, M., Topczewski, J., and Whitfield, T.T. (2013). Semicircular
canal morphogenesis in the zebrafish inner ear requires the function of gpr126
(lauscher), an adhesion class G protein-coupled receptor gene. Development *140*,
4362–4374. https://doi.org/10.1242/dev.098061.

15. Skouloudaki, K., Puetz, M., Simons, M., Courbard, J.-R., Boehlke, C., Hartleben, B.,
Engel, C., Moeller, M.J., Englert, C., Bollig, F., et al. (2009). Scribble participates in
Hippo signaling and is required for normal zebrafish pronephros development. Proc.
Natl. Acad. Sci. U. S. A. *106*, 8579–8584. https://doi.org/10.1073/pnas.0811691106.

16. Swinburne, I. (2025). A single-cell atlas of inner ear morphogenesis in zebrafish
(Biorxiv 250924). (Zenodo). https://doi.org/10.5281/ZENODO.17188953
https://doi.org/10.5281/ZENODO.17188953.

17. Farrell, J.A., Wang, Y., Riesenfeld, S.J., Shekhar, K., Regev, A., and Schier, A.F.
(2018). Single-cell reconstruction of developmental trajectories during zebrafish
embryogenesis. Science *360*. https://doi.org/10.1126/science.aar3131.

18. Sur, A., Wang, Y., Capar, P., Margolin, G., Prochaska, M.K., and Farrell, J.A. (2023).
Single-cell analysis of shared signatures and transcriptional diversity during
zebrafish development. Dev. Cell *58*, 3028-3047.e12.
https://doi.org/10.1016/j.devcel.2023.11.001.

19. Fernando, C.A., Conrad, P.A., Bartels, C.F., Marques, T., To, M., Balow, S.A.,
Nakamura, Y., and Warman, M.L. (2010). Temporal and spatial expression of CCN
genes in zebrafish. Dev. Dyn. *239*, 1755–1767. https://doi.org/10.1002/dvdy.22279.

20. Yeger, H., and Perbal, B. (2016). CCN family of proteins: critical modulators of the
tumor cell microenvironment. J. Cell Commun. Signal. *10*, 229–240.
https://doi.org/10.1007/s12079-016-0346-6.

21. Perbal, B. (2013). CCN proteins: A centralized communication network. J. Cell
Commun. Signal. *7*, 169–177. https://doi.org/10.1007/s12079-013-0193-7.

22. Zhao, B., Ye, X., Yu, J., Li, L., Li, W., Li, S., Yu, J., Lin, J.D., Wang, C.-Y., Chinnaiyan,
A.M., et al. (2008). TEAD mediates YAP-dependent gene induction and growth
control. Genes Dev. *22*, 1962–1971. https://doi.org/10.1101/gad.1664408.

23. Lee, S., Ahad, A., Luu, M., Moon, S., Caesar, J., Cardoso, W.V., Grant, M.B., and
Chaqour, B. (2019). CCN1-Yes-Associated Protein Feedback Loop Regulates
Physiological and Pathological Angiogenesis. Mol. Cell. Biol. *39*.
https://doi.org/10.1128/MCB.00107-19.

24. Park, M.-H., Kim, A.K., Manandhar, S., Oh, S.-Y., Jang, G.-H., Kang, L., Lee, D.-W.,
Hyeon, D.Y., Lee, S.-H., Lee, H.E., et al. (2019). CCN1 interlinks integrin and hippo
pathway to autoregulate tip cell activity. Elife *8*. https://doi.org/10.7554/eLife.46012.

25. Zhang, H., Pasolli, H.A., and Fuchs, E. (2011). Yes-associated protein (YAP)
transcriptional coactivator functions in balancing growth and differentiation in skin.
Proc. Natl. Acad. Sci. U. S. A. *108*, 2270–2275.
https://doi.org/10.1073/pnas.1019603108.

26. Pocaterra, A., Romani, P., and Dupont, S. (2020). YAP/TAZ functions and their
regulation at a glance. J. Cell Sci. *133*, jcs230425. https://doi.org/10.1242/jcs.230425.

27. Zanconato, F., Forcato, M., Battilana, G., Azzolin, L., Quaranta, E., Bodega, B.,
Rosato, A., Bicciato, S., Cordenonsi, M., and Piccolo, S. (2015). Genome-wide

association between YAP/TAZ/TEAD and AP-1 at enhancers drives oncogenic
growth. Nat. Cell Biol. *17*, 1218–1227. https://doi.org/10.1038/ncb3216.

28. Benhaddou, A., Keime, C., Ye, T., Morlon, A., Michel, I., Jost, B., Mengus, G., and
Davidson, I. (2012). Transcription factor TEAD4 regulates expression of myogenin
and the unfolded protein response genes during C2C12 cell differentiation. Cell
Death Differ. *19*, 220–231. https://doi.org/10.1038/cdd.2011.87.

29. Choi, H.M.T., Schwarzkopf, M., Fornace, M.E., Acharya, A., Artavanis, G., Stegmaier,
J., Cunha, A., and Pierce, N.A. (2018). Third-generation in situ hybridization chain
reaction: multiplexed, quantitative, sensitive, versatile, robust. Development *145*.
https://doi.org/10.1242/dev.165753.

30. Kurundkar, A.R., Kurundkar, D., Rangarajan, S., Locy, M.L., Zhou, Y., Liu, R.-M.,
Zmijewski, J., and Thannickal, V.J. (2016). The matricellular protein CCN1 enhances
TGF-β1/SMAD3-dependent profibrotic signaling in fibroblasts and contributes to
fibrogenic responses to lung injury. FASEB J. *30*, 2135–2150.
https://doi.org/10.1096/fj.201500173.

31. Nishida, T., Kawaki, H., Baxter, R.M., Deyoung, R.A., Takigawa, M., and Lyons, K.M.
(2007). CCN2 (Connective Tissue Growth Factor) is essential for extracellular matrix
production and integrin signaling in chondrocytes. J. Cell Commun. Signal. *1*, 45–58.
https://doi.org/10.1007/s12079-007-0005-z.

32. Wong, M., Kireeva, M.L., Kolesnikova, T.V., and Lau, L.F. (1997). Cyr61, product of
a growth factor-inducible immediate-early gene, regulates chondrogenesis in mouse
limb bud mesenchymal cells. Dev. Biol. *192*, 492–508.
https://doi.org/10.1006/dbio.1997.8766.

33. Mateus, R., Lourenço, R., Fang, Y., Brito, G., Farinho, A., Valério, F., and Jacinto, A.
(2015). Control of tissue growth by Yap relies on cell density and F-actin in zebrafish
fin regeneration. Development *142*, 2752–2763. https://doi.org/10.1242/dev.119701.

34. Jean, R.P., Gray, D.S., Spector, A.A., and Chen, C.S. (2004). Characterization of the
nuclear deformation caused by changes in endothelial cell shape. J. Biomech. Eng.
126, 552–558. https://doi.org/10.1115/1.1800559.

35. Venturini, V., Pezzano, F., Castro, F.C., Häkkinen, H.-M., Jiménez-Delgado, S.,
Colomer-Rosell, M., Marro, M., Tolosa-Ramon, Q., Paz-López, S., Valverde, M.A.,
et al. (2020). The nucleus measures shape changes for cellular proprioception to
control dynamic cell behavior. Science *370*, eaba2644.
https://doi.org/10.1126/science.aba2644.

36. Torregrosa-Carrión, R., Piñeiro-Sabarís, R., Siguero-Álvarez, M., Grego-Bessa, J.,
Luna-Zurita, L., Fernandes, V.S., MacGrogan, D., Stainier, D.Y.R., and de la Pompa,
J.L. (2021). Adhesion G protein-coupled receptor Gpr126/Adgrg6 is essential for
placental development. Sci Adv 7, eabj5445. https://doi.org/10.1126/sciadv.abj5445.

37. Bian, F., Hansen, V., Feng, H.C., He, J., Chen, Y., Feng, K., Ebrahimi, B., Gray, R.S.,
Chai, Y., Wu, C.-L., et al. (2024). The G protein-coupled receptor ADGRG6 maintains
mouse growth plate homeostasis through IHH signaling. J. Bone Miner. Res. *39*,
1644–1658. https://doi.org/10.1093/jbmr/zjae144.

38. Pogoda, H.-M., Sternheim, N., Lyons, D.A., Diamond, B., Hawkins, T.A., Woods, I.G.,
Bhatt, D.H., Franzini-Armstrong, C., Dominguez, C., Arana, N., et al. (2006). A
genetic screen identifies genes essential for development of myelinated axons in
zebrafish. Dev. Biol. *298*, 118–131. https://doi.org/10.1016/j.ydbio.2006.06.021.

39. Mogha, A., Benesh, A.E., Patra, C., Engel, F.B., Schöneberg, T., Liebscher, I., and
Monk, K.R. (2013). Gpr126 functions in Schwann cells to control differentiation and
myelination via G-protein activation. J. Neurosci. *33*, 17976–17985.
https://doi.org/10.1523/JNEUROSCI.1809-13.2013.

40. Petersen, S.C., Luo, R., Liebscher, I., Giera, S., Jeong, S.-J., Mogha, A., Ghidinelli,
M., Feltri, M.L., Schöneberg, T., Piao, X., et al. (2015). The adhesion GPCR GPR126
has distinct, domain-dependent functions in Schwann cell development mediated by
interaction with laminin-211. Neuron *85*, 755–769.
https://doi.org/10.1016/j.neuron.2014.12.057.

41. Seamon, K., and Daly, J.W. (1981). Activation of adenylate cyclase by the diterpene
forskolin does not require the guanine nucleotide regulatory protein. J. Biol. Chem.
256, 9799–9801. https://doi.org/10.1016/s0021-9258(19)68694-1.

42. Gonzalez, G.A., and Montminy, M.R. (1989). Cyclic AMP stimulates somatostatin
gene transcription by phosphorylation of CREB at serine 133. Cell *59*, 675–680.
https://doi.org/10.1016/0092-8674(89)90013-5.

43. Dobroff, A.S., Wang, H., Melnikova, V.O., Villares, G.J., Zigler, M., Huang, L., and
Bar-Eli, M. (2009). Silencing cAMP-response element-binding protein (CREB)
identifies CYR61 as a tumor suppressor gene in melanoma. J. Biol. Chem. *284*,
26194–26206. https://doi.org/10.1074/jbc.M109.019836.

44. Yu, Z., Kong, Q., and Kone, B.C. (2010). CREB trans-activation of disruptor of
telomeric silencing-1 mediates forskolin inhibition of CTGF transcription in mesangial
cells. Am. J. Physiol. Renal Physiol. *298*, F617-24.
https://doi.org/10.1152/ajprenal.00636.2009.

45. Inhibition Protein Kinase Phenocopies Ectopic Expression ofhedgehogin CNS Wild-
Type andcyclopsMutant Embryos.

46. Aragona, M., Panciera, T., Manfrin, A., Giulitti, S., Michielin, F., Elvassore, N., Dupont,
S., and Piccolo, S. (2013). A mechanical checkpoint controls multicellular growth
through YAP/TAZ regulation by actin-processing factors. Cell *154*, 1047–1059.
https://doi.org/10.1016/j.cell.2013.07.042.

47. Mason, D.E., Collins, J.M., Dawahare, J.H., Nguyen, T.D., Lin, Y., Voytik-Harbin, S.L., Zorlutuna, P., Yoder, M.C., and Boerckel, J.D. (2019). YAP and TAZ limit cytoskeletal and focal adhesion maturation to enable persistent cell motility. J. Cell Biol. *218*, 1369–1389. https://doi.org/10.1083/jcb.201806065.

48. Mason, D.E., Goeckel, M., Vega, S.L., Wu, P.-H., Johnson, D., Heo, S.-J., Wirtz, D., Burdick, J.A., Wood, L., Chow, B., et al. (2023). Mechanotransductive feedback control of endothelial cell motility and vascular morphogenesis. https://doi.org/10.7554/elife.86668.1.

49. Panciera, T., Azzolin, L., Cordenonsi, M., and Piccolo, S. (2017). Mechanobiology of YAP and TAZ in physiology and disease. Nat. Rev. Mol. Cell Biol. *18*, 758–770. https://doi.org/10.1038/nrm.2017.87.

50. Lau, L.F. (2016). Cell surface receptors for CCN proteins. J. Cell Commun. Signal. *10*, 121–127. https://doi.org/10.1007/s12079-016-0324-z.

51. Wang, X., Li, L., and Liu, D. (2014). Expression analysis of integrin β1 isoforms during zebrafish embryonic development. Gene Expr. Patterns *16*, 86–92. https://doi.org/10.1016/j.gep.2014.10.001.

52. Littlewood Evans, A., and Müller, U. (2000). Stereocilia defects in the sensory hair cells of the inner ear in mice deficient in integrin alpha8beta1. Nat. Genet. *24*, 424–428. https://doi.org/10.1038/74286.

53. Kechagia, J.Z., Ivaska, J., and Roca-Cusachs, P. (2019). Integrins as biomechanical sensors of the microenvironment. Nat. Rev. Mol. Cell Biol. *20*, 457–473. https://doi.org/10.1038/s41580-019-0134-2.

54. Monk, K.R., Oshima, K., Jörs, S., Heller, S., and Talbot, W.S. (2011). Gpr126 is essential for peripheral nerve development and myelination in mammals. Development *138*, 2673–2680. https://doi.org/10.1242/dev.062224.

55. Patra, C., van Amerongen, M.J., Ghosh, S., Ricciardi, F., Sajjad, A., Novoyatleva, T., Mogha, A., Monk, K.R., Mühlfeld, C., and Engel, F.B. (2013). Organ-specific function of adhesion G protein-coupled receptor GPR126 is domain-dependent. Proc. Natl. Acad. Sci. U. S. A. *110*, 16898–16903. https://doi.org/10.1073/pnas.1304837110.

56. Lu, S., Liu, S., Wietelmann, A., Kojonazarov, B., Atzberger, A., Tang, C., Schermuly, R.T., Gröne, H.-J., and Offermanns, S. (2017). Developmental vascular remodeling defects and postnatal kidney failure in mice lacking Gpr116 (Adgrf5) and Eltd1 (Adgrl4). PLoS One *12*, e0183166. https://doi.org/10.1371/journal.pone.0183166.

57. Lala, T., and Hall, R.A. (2022). Adhesion G protein-coupled receptors: structure, signaling, physiology, and pathophysiology. Physiol. Rev. *102*, 1587–1624. https://doi.org/10.1152/physrev.00027.2021.

58. Paavola, K.J., Sidik, H., Zuchero, J.B., Eckart, M., and Talbot, W.S. (2014). Type IV
collagen is an activating ligand for the adhesion G protein-coupled receptor GPR126.
Sci. Signal. *7*, ra76. https://doi.org/10.1126/scisignal.2005347.

59. Mitgau, J., Franke, J., Schinner, C., Stephan, G., Berndt, S., Placantonakis, D.G.,
Kalwa, H., Spindler, V., Wilde, C., and Liebscher, I. (2022). The N Terminus of
Adhesion G Protein-Coupled Receptor GPR126/ADGRG6 as Allosteric Force
Integrator. Front Cell Dev Biol *10*, 873278. https://doi.org/10.3389/fcell.2022.873278.

60. Li, Q., Huo, A., Li, M., Wang, J., Yin, Q., Chen, L., Chu, X., Qin, Y., Qi, Y., Li, Y., et
al. (2024). Structure, ligands, and roles of GPR126/ADGRG6 in the development and
diseases. Genes Dis *11*, 294–305. https://doi.org/10.1016/j.gendis.2023.02.016.

61. Wilde, C., Mitgau, J., Suchý, T., Schöneberg, T., and Liebscher, I. (2022). Translating
the force-mechano-sensing GPCRs. Am. J. Physiol. Cell Physiol. *322*, C1047–
C1060. https://doi.org/10.1152/ajpcell.00465.2021.

62. Behrndt, M., Salbreux, G., Campinho, P., Hauschild, R., Oswald, F., Roensch, J.,
Grill, S.W., and Heisenberg, C.-P. (2012). Forces driving epithelial spreading in
zebrafish gastrulation. Science *338*, 257–260.
https://doi.org/10.1126/science.1224143.

63. Gerety, S.S., and Wilkinson, D.G. (2011). Morpholino artifacts provide pitfalls and
reveal a novel role for pro-apoptotic genes in hindbrain boundary development. Dev.
Biol. *350*, 279–289. https://doi.org/10.1016/j.ydbio.2010.11.030.

64. Sorlien, E.L., Witucki, M.A., and Ogas, J. (2018). Efficient Production and
Identification of CRISPR/Cas9-generated Gene Knockouts in the Model System
Danio rerio. J. Vis. Exp. https://doi.org/10.3791/56969.

65. Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., III,
Hao, Y., Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive integration
of single-cell data. Cell *177*, 1888-1902.e21.
https://doi.org/10.1016/j.cell.2019.05.031.

66. Blondel, V.D., Guillaume, J.-L., Lambiotte, R., and Lefebvre, E. (2008). Fast
unfolding of communities in large networks. J. Stat. Mech. *2008*, P10008.
https://doi.org/10.1088/1742-5468/2008/10/p10008.

67. Schultz, J., Milpetz, F., Bork, P., and Ponting, C.P. (1998). SMART, a simple modular
architecture research tool: identification of signaling domains. Proc. Natl. Acad. Sci.
U. S. A. *95*, 5857–5864. https://doi.org/10.1073/pnas.95.11.5857.

68. Letunic, I., Khedkar, S., and Bork, P. (2021). SMART: recent updates, new
developments and status in 2020. Nucleic Acids Res. *49*, D458–D460.
https://doi.org/10.1093/nar/gkaa937.

69. Sievers, F., and Higgins, D.G. (2014). Clustal Omega, accurate alignment of very
large   numbers   of   sequences.   Methods   Mol.   Biol.   *1079*,   105–116.
https://doi.org/10.1007/978-1-62703-646-7_6.

Figure Legends

Figure 1: ECM swelling-mediated Yap activation is required for bud extension

(a) Three-dimensional (3D) rendered OVs imaged at selected time points using *Tg(actb2:membrane-neongreen-neongreen)* embryos. A, anterior bud; AL, anterior-lateral bud; P, posterior bud; PL, posterior-lateral bud; V, ventral bud. Developmental stages shown in hours post fertilization (hpf). Scale bar: 50 μm.

(b-b') 2D sections of OVs imaged at selected time points using H2B-GFP mRNA injected *TgKI(Yap1-mScarlet)* embryos. Insets (b') show enlarged images of the anterior pre-budding or budding region corresponding to the white boxes in (b). Dorsal epithelial cells showing nuclear-Yap localization are indicated by asterisks. Scale bar: 10 or 50 μm. Representative images from three independent experiments are shown.

(c) Quantification of the Yap nuclear to cytoplasmic ratio in the cells at the anterior pre-budding or budding region at selected time points. Data are mean ± SD. *n* denotes the number of cells measured per condition from seven individual embryos from three independent experiments. *P*-value as labeled (Mann-Whitney test).

(d-e) Effect of hyaluronidase injection on Yap nuclear translocation. (d) 2D sections of the anterior bud in *TgKI(Yap1-mScarlet)*, injected with H2B-GFP mRNA, imaged before and at 1.5 h post-injection (hpi) of PBS or HAase. Pre-injection and post-injection images were acquired from the same individual embryos. Arrows indicate Yap1 translocated to the nucleus. Scale bar: 10 μm. Representative images from two independent experiments are shown. (d) Quantification of the Yap nuclear to cytoplasmic ratio in the anterior bud before and at 1.5 hpi of PBS or HAase. Data are mean ± SD. *n* denotes the number of cells measured per condition from ten individual embryos from two independent experiments. *P*-values as labeled (Kruskal–Wallis test with Dunn's test).

(f-g) Effect of *yap1* knockdown on bud growth. (f) 3D-rendered OVs from from *Tg(actb2:membrane-neongreen-neongreen)* embryos, injected with control or *yap1* MO, imaged at 50 hpf. Scale bar: 50 μm. Representative images from two independent experiments are shown. (g) Quantification of bud length in embryos injected with control or *yap1* MO at 50 hpf. Data are mean ± SD. *n* denotes the number of buds from individual embryos measured per condition from two independent experiments. *P*-values as labeled (unpaired two-tailed Student's t-test)

Figure 2: *ccn1l1*, a Yap target gene, is enriched in the canal genesis zone during SCC formation

(a) UMAP of single-cell RNA of the inner ear cell clusters from the DanioCell dataset. Single cells in the canal-forming cell cluster are colored blue.

(b) Dot plot showing the expression levels of the top 10 DEGs in the canal-forming cell cluster during SCC formation: pre-budding (36-40 hpf), bud extension (45-50 hpf), and bud fusion (60-72 hpf) stages. Color encodes the normalized gene expression level, and the dot size encodes the percentage of positive cells within a cluster.

(c) 3D-rendered OVs at select time points stained with multiplex *in situ* probes against *ccn1l1* (green) and *col2a1a* (white). Scale bar: 50 µm. Representative images from three independent experiments are shown.

(d) Normalized fluorescence probe intensities of *ccn1l1* or *col2a1a* genes across the illustrated region of interest (ROI) at 50 hpf indicated by the red dashed line along the anterior bud. Data are mean ± SEM. *n* denotes the number of buds from individual embryos measured per condition from three independent experiments.

(e) Comparison of normalized fluorescence probe intensity of *ccn1l1* in the anterior pre-budding, budding, or pillar region at each stage. Data are mean ± SD. *n* denotes the number of buds or pillars from individual embryos measured from two independent experiments. *P*-values as labeled (one-way ANOVA with Tukey's test).

(f-g) Effect of *yap1* knockdown on *ccn1l1* expression. (f) 3D-rendered OVs from *Tg(actb2:membrane-neongreen-neongreen)* embryos at 50 hpf stained with multiplex *in situ* probes against *ccn1l1* (green). Heatmap represents fluorescence intensity with the same contrast for the *ccn1l1* probe of embryos injected with control and *yap1* MO. Scale bar: 50 µm. Representative images from two independent experiments are shown. (g) Quantification of probe fluorescence intensity of *ccn1l1* probe in anterior buds. Data are mean ± SD. *n* denotes the number of buds from individual embryos measured from two independent experiments. *P*-value as labeled (unpaired two-tailed Student's t-test).

Figure 3: Ccn1l1 induces ECM production and drives positive feedback loop via Yap mechanotransduction

(a-b) Effect of *ccn1l1* knockdown on bud growth. (a) 3D-rendered OVs from *Tg(actb2:membrane-neongreen-neongreen)* embryos, injected with *tyrosinase* gRNA, *ccn1l1* gRNA, control MO or *ccn1l1* MO, imaged at 50 hpf. Scale bar: 50 µm. Representative images from three independent experiments are shown. *tyrosinase* gRNA is used as a gRNA injection control because it causes loss of pigmentation but does not affect SCC formation. (b) Quantification of bud length in control or *ccn1l1* knockdown embryos at 50 hpf. Data are mean ± SD. In the absence of buds, bud

lengths correspond to cell height. *n* denotes the number of buds from individual embryos measured per condition from three independent experiments. *P*-values as labeled (unpaired two-tailed Student's t-test).

(c-d) Effect of *ccn1l1* knockdown on *has3* and *vcanb* expression levels in buds. (c) 3D-rendered OVs at 50 hpf stained with multiplex *in situ* probes against *has3* (green), *vcanb* (magenta), and *col2a1a* (white). Heatmaps represent fluorescence intensity with the same contrast for each probe of embryos injected with control or *ccn1l1* MO. Scale bar: 50 μm. Representative images from two independent experiments are shown. (d) Quantification of probe fluorescence intensity of *has3* or *vcanb* probe in anterior buds. Data are mean ± SD. *n* denotes the number of buds from individual embryos measured per condition from two independent experiments. *P*-values as labeled (unpaired two-tailed Student's t-test).

(e-h) Effect of hyaluronidase injection on *ccn1l1* and *has3* expression levels in buds. 3D-rendered OVs and quantification of probe fluorescence intensity of *ccn1l1* (e-f) or *has3* (g-h) at 0 or 6 hpi of PBS or HAase. Heatmaps represent fluorescence intensity with the same contrast for each probe of embryos injected with PBS or HAase. Scale bar: 50 μm. Representative images from two independent experiments are shown. Data are mean ± SD. *n* denotes the number of buds from individual embryos measured per condition from two independent experiments. *P*-values as labeled (one-way ANOVA with Tukey's test).

(i-k) Effect of DN Yap1 expression on bud extension and bud-ECM volume. (i) 2D sections of anterior buds from *Tg(hsp: RFP-DN Yap1)* and sibling control embryos, both *Tg(actb2:membrane-neongreen-neongreen)*, imaged at 50 hpf following heat-shock induction. Regions marked in light blue indicate the bud-ECM region measured in quantifications. Scale bar: 20 μm. Representative images from two independent experiments are shown. Quantification of bud length (j) and bud-ECM volume (k) in sibling control or DN Yap1-expressing embryos at 50 hpf. Data are mean ± SD. *n* denotes the number of buds from individual embryos measured per condition from two independent experiments. *P*-values as labeled (unpaired two-tailed Student's t-test).

(l) A schematic model of the positive mechanotransduction feedback loop during bud growth.

Figure 4: Mechanical patterning by the positive feedback loop recruits adjacent cells to sustain bud extension and ensure timely

(a) Quantification of cell number in the anterior bud across individual time points obtained from time-lapse imaging of *Tg(actb2:membrane-neongreen-neongreen)* embryos. Data are mean ± SD. Dashed lines indicate the range of SD at each time

point. *n* denotes the number of buds from individual embryos measured from three independent experiments.

(b) 3D-rendered images of an anterior bud of the *Tg(actb2:membrane-neongreen-neongreen)* embryo at selected time points with individual cells tracked. Scale bar: 50 µm. Representative images from two independent experiments are shown.

(c) Quantification of cell number in anterior buds of embryos injected with PBS or HAase at 0 or 6 hpi. Data are mean ± SD. *n* denotes the number of buds from individual embryos measured per condition from two independent experiments. *P*-values as labeled (one-way ANOVA with Tukey's test).

(d) Quantification of cell number in anterior buds of embryos injected with control or *yap1* MO at 50 hpf. Data are mean ± SD. *n* denotes the number of buds from individual embryos measured per condition from two independent experiments. *P*-value as labeled (unpaired two-tailed Student's t-test).

(e) Schematics of the 2D vertex model simulations, showing tip, budding and border cells, as well as the relevant physical parameters. All values of physical parameters are given in the supplementary information.

(f) Trajectories of the aspect ratio of a tip cell and a recruited cell obtained with 2D vertex model simulations, exhibiting a sharp increase in aspect ratio as cell position shifts from the bud border to the budding region. Illustrations show trajectories of two individual cells from bud initiation to the end of bud extension (defined by the moment where the bud height reaches 30 µm).

(g) Quantification of 3D nuclear aspect ratio. The nuclear aspect ratio was calculated as the ratio of the long to short axes (long/short) of the ellipsoidal nucleus. Data are mean ± SD. *n* denotes the number of nuclei from individual cells measured per condition from one experiment. *P*-value as labeled (unpaired two-tailed Student's t-test).

(h) Spatial difference of Yap nuclear translocation within the anterior bud at 50 hpf. Data are mean ± SD. coefficient of variation (CV) of each condition is shown. *n* denotes the number of cells per condition from two independent experiments. *P*-values as labeled (one-way ANOVA with Tukey's test).

(i) Simulation of bud-ECM area over time, with and without mechanotransduction-driven ECM production. Tip cells are set to 100% ECM production rate.

(j) Quantification of bud-ECM volume of anterior buds in embryos injected with control or *yap1* MO at selected time points. Data are mean ± SD. *n* denotes the number of cells per condition from two independent experiments. *P*-values at selected time pointes as labeled (unpaired two-tailed Student's t-test).

(k) Simulation of bud length over time, with and without mechanotransduction-driven ECM production. Tip cells are set to 100% ECM production rate.

(l) Quantification of bud length of the anterior buds in embryos injected with control or *yap1* MO at selected time points. Data are mean ± SD. *n* denotes the number of cells per condition from two independent experiments. *P*-values at selected time points as labeled (unpaired two-tailed Student's t-test).

(m) Phase diagram showing the time needed for the bud to reach 30 μm and fuse, as a function of bud ECM production by budding cells (x axis) and tip cells (y axis). The dotted level line indicates the region where fusion happens in less than 15 hours after budding. Grey areas indicate regions where the bud failed to fuse (did not reach 30 μm within 22 hours after budding).

(n) Quantification of bud growth rate (μm/hour) of the anterior bud in wild-type embryos or embryos injected with *ccn1l1* MO. Data are mean ± SD. *n* denotes the number of cells per condition from one experiment. *P*-values as labeled (one-way ANOVA with Tukey's test). Growth rates for each bud were calculated over the following time ranges during bud extension stage: control (48-50 hpf); *ccn1l1* MO 0.5, 1, and 2 ng (50-56 hpf).

(o) 3D-rendered OVs from *Tg(actb2:membrane-neongreen-neongreen)* embryos, injected with control or *ccn1l1* MO, imaged at selected time points. Scale bar: 50 μm. Representative images from one experiment per condition are shown.

Figure 5: Gpr126-cAMP-PKA-CREB signaling during bud fusion suppresses *ccn1l1* expression to terminate bud extension

(a) 3D-rendered OVs from wild-type or *gpr126* [st49/st49] mutant embryos, both *Tg(actb2:membrane-neongreen-neongreen),* imaged at selected time points. Blue and yellow dashed lines indicate the anterior bud/pillar or the anterior-lateral portion of the pillar, respectively. Scale bar: 50 μm. Representative images from two independent experiments are shown.

(b-c) *ccn1l1* and *has3* expression levels in *gpr126* [st49/st49] mutant embryos. (b) 3D-rendered OVs of wild-type or *gpr126* [st49/st49] mutant embryos at 60 hpf stained with multiplex *in situ* probes against *ccn1l1* (green), *has3* (magenta), and *col2a1a* (white). Heatmaps represent fluorescence intensity with the same contrast for each probe of embryos. Scale bar: 50 μm. Representative images from three independent experiments are shown. (c) Quantification of probe fluorescence intensities of *ccn1l1* or *has3* probes in anterior buds or pillars at 50 hpf or 60 hpf. Data are mean ± SD. *n* denotes the number of buds or pillars from individual embryos measured from three independent experiments. *P*-values as labeled (Kruskal-Wallis test with Dunn's test).

(d-f) Effect of *gpr126* $^{st49/st49}$ mutation on bud growth. Quantification of bud length (d), cell number in the anterior bud or pillar (e), and bud-ECM volume (f) at selected time points. Data are mean ± SD. *n* denotes the number of buds or pillars from individual embryos measured from two independent experiments. *P*-values as labeled (unpaired two-tailed Student's t-test).

(g) 2D sections of OVs showing phospho-CREB and F-actin, stained using anti-phospho-CREB^{ser133} antibody and phalloidin, respectively, at selected time points. Insets show enlarged images of the anterior buds corresponding to the dashed white boxes. Scale bar: 50 μm. Representative images from three independent experiments are shown.

(h) 2D sections OVs showing phospho-CREB and F-actin, stained using anti-phospho-CREB^{ser133} antibody and phalloidin, respectively, at 60 hpf in sibling control or *gpr126* $^{st49/st49}$ mutant embryos. Scale bar: 50 μm. Representative images from two independent experiments are shown.

(i-j) Effect of DN PKA on *ccn1l1* expression in buds. (i) 3D-rendered OVs from *Tg(hsp:DN PKA-GFP)* embryos and sibling controls after heat-shock induction, stained with multiplex *in situ* probes against *ccn1l1* (magenta) and *col2a1a* (white). Heatmaps represent fluorescence intensity with the same contrast for the *ccn1l1* probe of each. Scale bar: 50 μm. Representative images from three independent experiments are shown. (j) Quantification of probe fluorescence intensity of *ccn1l1* in anterior buds or pillars at 56 hpf. Data are mean ± SD. *n* denotes the number of buds or pillars from individual embryos measured from three independent experiments. *P*-value as labeled (unpaired two-tailed Student's t-test).

Extended Figure Legends

Extended Figure 1: ECM swelling-mediated Yap activation in budding cells, related to Figure 1

(a) Illustration of budding epithelium and bud-ECM region filled with hyaluronan.

(b) 3D-rendered and 2D section of the anterior bud imaged at budding (45 hpf) and bud extension (50 hpf) stages using *Tg(actb2:membrane-neongreen-neongreen)* embryos. Yellow asterisk indicates the opposing anterior-lateral bud. Regions marked in light blue indicate the bud-ECM region. Scale bar: 20 μm. Representative images from two independent experiments are shown.

(c) Quantification of cell aspect ratio in budding cells in the anterior bud at budding (45 hpf) and bud extension (50 hpf) stages. Cell height and width in a budding cell are defined as described in Extended Fig. 1a. Data are mean ± SD. *n* denotes the number of buds from individual embryos measured per condition from two independent experiments. *P*-value as labeled (unpaired two-tailed Student's t-test).

(d) Quantification of the Yap nuclear to cytoplasmic ratio in the dorsal epithelial cells at pre-budding (40 hpf) or bud extension (50 hpf) stages. Data are mean ± SD. *n* denotes the number of cells measured per condition from nine individual embryos of two independent experiments. *P*-value as labeled (unpaired two-tailed Student's t-test).

(e) 2D sections of the anterior bud in *Tg(actb2:membrane-neongreen-neongreen)* embryos imaged at 0 and 6 hpi of PBS and HAase. 0 hpi and 6 hpi images were acquired from the same embryo. Scale bar: 20 μm. Representative images from two independent experiments are shown.

(f) Quantification of bud-ECM volume of PBS or HAase-treated anterior buds at 6 hpi (52 hpf). Data are mean ± SD. *n* denotes the number of buds measured per condition from two independent experiments. *P*-value as labeled (unpaired two-tailed Student's t-test).

(g) Quantification of cell aspect ratio of budding cells in anterior buds before (46 hpf) and at 6 hpi (52 hpf) of PBS and HAase. Data are mean ± SD. *n* denotes the number of buds from individual embryos measured per condition from two independent experiments. *P*-values as labeled (Kruskal-Wallis test with Dunn's test).

(h) Quantification of bud length of anterior buds before (46 hpf) and at 6 hpi (52 hpf) of PBS and HAase. Data are mean ± SD. *n* denotes the number of buds from individual embryos measured per condition from two independent experiments. *P*-values as labeled (one-way ANOVA with Tukey's test).

(i) Quantification of the Yap nuclear-cytoplasmic ratio in the dorsal epithelial cells before (50 hpf) and after (51.5 hpf) injection of HAase. Data are mean ± SD. *n* denotes the number of cells measured per condition from nine individual embryos of two independent experiments. *P*-value as labeled (Mann-Whitney test).

(j-k) Immunostaining of Yap1 in budding cells. (j) 2D sections of anterior buds showing Yap1 and nuclear staining using anti-Yap1 antibody and Hoechst stain respectively, at selected time points. Scale bar: 10 μm. Arrows indicate Yap1 translocated to the nucleus. Representative images from two independent experiments are shown. (k) Quantification of the Yap nuclear-cytoplasmic ratio in the epithelial cells at the canal genesis zone at selected time points. Data are mean ± SD. *n* denotes the number of cells measured per condition from two independent experiments. *P*-values as labeled (Kruskal-Wallis test with Dunn's test).

(l-m) Effect of HAase on nuclear Yap translocation. (l) 2D sections of anterior buds showing Yap1 and nuclear staining using anti-Yap1 antibody, and Hoechst stain, respectively, at 3 hpi of PBS or HAase. Scale bar: 10 µm. Arrows indicate Yap1 translocated to the nucleus. Representative images from two independent experiments are shown. (m) Quantification of the Yap nuclear to cytoplasmic ratio in the budding cells in the anterior buds at 3 hpi of PBS or HAase. Data are mean ± SD. *n* denotes the number of cells measured per condition from two independent experiments. *P*-value as labeled (Mann-Whitney test).

(n) 3D-rendered OVs from *Tg(actb2:membrane-neongreen-neongreen)* embryos, injected with control or *yap1* MO, imaged at 72 hpf. Yellow arrows indicate unfused posterior and ventral buds. Scale bar: 50 µm. Representative images from three independent experiments are shown.

Extended Figure 2: Lmx1bb regulates *ccn1l1* expression at the canal genesis zone, related to Figure 2

(a) UMAP of single-cell RNA of *vcanb*-expressing cell cluster from the DanioCell dataset. Heatmap color of each dot represents *vcanb* expression levels.

(b) Schematic image and sequence of the TEAD binding motifs in the genomic sequence upstream of the *ccn1l1* locus.

(c-d) Effect of *yap1* knockdown on the onset of *ccn1l1* expression at the pre-budding stage. (c) 3D-rendered OVs, injected with control or *yap1* MO, imaged at 40 hpf stained with multiplex *in situ* probes against *ccn1l1* (green) and *col2a1a* (white). Scale bar: 50 µm. Representative images from two independent experiments are shown. (d) Quantification of probe fluorescence intensity of *ccn1l1* probe in the anterior budding region at the pre-budding stage (40 hpf). Data are mean ± SD. *n* denotes the number of budding regions from individual embryos measured from two independent experiments. *P*-value as labeled (unpaired two-tailed Student's t-test).

(e-f) Effect of *lmx1bb* mutation on *ccn1l1* expression level. (e) 3D-rendered OVs of *lmx1bb* $^{jj410/jj410}$ or sibling control embryos at 50 hpf stained with multiplex *in situ* probes against *ccn1l1* (green) and *col2a1a* (white). Heatmap represents fluorescence intensity with the same contrast for the *ccn1l1* probe of each. Scale bar: 50 µm. Representative images from three independent experiments are shown. (f) Quantification of probe fluorescence intensity of *ccn1l1* probe in the posterior pre-budding region. Data are mean ± SD. *n* denotes the number of budding regions from individual embryos measured from two independent experiments. *P*-value as labeled (unpaired two-tailed Student's t-test).

(g) Maximum intensity projections of OVs treated with Notch receptor inhibitor RO4929097 at 50 hpf stained with multiplex *in situ* probes against *lmx1bb* (magenta), *ccn1l1* (green), and *col2a1a* (white). Scale bar: 50 µm. Representative images from three independent experiments are shown.

Extended Figure 3: Ccn1l1 induces *has3* and *vcanb* expression in budding cells, related to Figure 3

(a) E-values of predicted functional domains of zebrafish Ccn1l1 obtained with SMART.

(b) Multiple sequence alignment of amino acids of each functional domain from zebrafish (*Danio rerio*) Ccn1l1 (Dr_Ccn1l1), zebrafish Ccn1 (Dr_Ccn1), and human (*Homo sapiens*) CCN1 (Hs_CCN1) performed with Clustal Omega. Black (*), dark grey (:), and grey (.) indicate fully conserved residues, strong similarity of residues, and weak similarity of residues, respectively, across species. The percentage sequence identity of Dr_Ccn1 and Dr_Ccn1l1 relative to Hs_CCN1 is shown.

(c) 3D-rendered OVs at pre-budding (40 hpf) or bud extension (50 hpf) stages stained with multiplex *in situ* probes against *ccn1l1* (cyan), *has3* (green), and *vcanb* (green). A, anterior bud; AL, anterior-lateral bud; P, posterior bud; PL, posterior-lateral bud; V, ventral bud. Scale bar: 50 µm. Representative images from three independent experiments are shown.

(d) Quantification of probe fluorescence intensities of *has3 and vcanb* probe measured at the pre-budding stage (anterior budding region), extension stage (anterior bud region), or fusion stage (anterior pillar region). Data are mean ± SD. *n* denotes the number of buds from individual embryos measured from two independent experiments.

(e-f) Effect of *ccn1l1* knockdown on *has3* and *vcanb* expression at the pre-budding stage. (e) 3D-rendered OVs injected with control or *ccn1l1* MO at 40 hpf stained with multiplex *in situ* probes against *has3* (green), *vcanb* (magenta), and *col2a1a* (white). Heatmaps represent fluorescence intensity with the same contrast for each probe of embryos. Scale bar: 50 µm. Representative images from two independent experiments are shown. (f) Quantification of probe fluorescence intensites of *has3* and *vcanb* probes in the anterior pre-budding region. Data are mean ± SD. *n* denotes the number of buds from individual embryos measured from two independent experiments. *P*-values as labeled (unpaired two-tailed Student's t-test).

Extended Figure 4: Mechanical patterning of budding cells, related to Figure 4

(a) Simulation snapshot of a bud with low adhesion to the basement membrane obtained with the 2D vertex model, showing the trajectory of two cells during bud extension. Upper and lower panels show the bud shape after and before budding, respectively.

(b) Simulated aspect ratio of a tip cell and a recruited cell during bud extension, in the case of low adhesion to the basement membrane. The dashed black line indicates the trajectory of the aspect ratio of a recruited cell in the control case.

(c) Simulation snapshot of a bud in the absence of cytocinches obtained with the 2D vertex model, showing the trajectory of two cells during bud extension. Cells appear to be pushed away from the bud center, rather than flowing towards it. Upper and lower panels show the bud shape after and before budding, respectively.

(d) Simulated Aspect ratio of a tip cell and a recruited cell during bud extension obtained with the 2D vertex model, in the absence of cytocinches. The dashed black line indicates the evolution of the aspect ratio of a recruited cell in the control case.

(e) Simulation snapshots of bud morphology at 5 hours post-budding for different ECM production rate parameter values. The red heat map shows the ECM production rate in tip cells. Mech. t. indicates the percent of ECM production driven by mechanotransduction contribution from recruited cells.

(f) 2D sections of anterior buds from *Tg(actb2:membrane-neongreen-neongreen)* embryos injected with control or *ccn1l1* MO at the indicated concentrations, imaged at 50 hpf. Scale bar: 20 μm. Representative images from two independent experiments are shown.

Extended Figure 5: Gpr126-cAMP-PKA-CREB signaling terminates bud extension, related to Figure 5

(a) Quantification of anterior bud length across individual time points obtained from time-lapse imaging of *Tg(actb2:membrane-neongreen-neongreen)* embryos. Data are mean ± SD. Dashed lines indicate the range of SD at each time point. *n* denotes the number of buds from individual embryos measured from three independent experiments.

(b) 3D-rendered OVs at selected time points stained with multiplex *in situ* probes against *gpr126* (green) and *col2a1a* (white). Heatmap represents fluorescence intensity with the same contrast for the *gpr126* probe of each embryo. Scale bar: 50 μm. Representative images from two independent experiments are shown.

(c) 3D-rendered OVs of wild-type and *gpr126* $^{st49/st49}$ mutant embryos at 72 hpf stained with multiplex *in situ* probes against *ccn1l1* (green), *has3* (magenta), and *col2a1a* (white). Heatmaps represent fluorescence intensity with the same contrast for each

probe of wild-type or *gpr126* *st49/st49* mutant embryos. Scale bar: 50 μm. Representative images from three independent experiments are shown.

(d) Quantification of bud-ECM volume in anterior buds across individual time points obtained from time-lapse imaging of wild-type or *gpr126* *st49/st49* embryos expressing membrane-NeonGreen. Data are mean ± SD. *n* denotes the number of buds from individual embryos measured from three independent experiments. *P*-values as labeled (unpaired two-tailed Student's t-test)

(e) Simulated bud ECM area (equivalent of bud ECM volume in our 2D simulation) as function of time, with (fusion: blue line) or in the absence (no fusion: red line) of the terminating event of bud fusion. Tip and budding cell productions are set to 100%.

(f-g) Effect of forskolin treatment on *ccn1l1* and *has3* expression. (f) 3D-rendered OVs of embryos treated with forskolin at 52 hpf stained with multiplex *in situ* probes against the combination of *ccn1l1* (green) and *col2a1a* (white) or *has3* (magenta) and *col2a1a* (white). Scale bar: 50 μm. Representative images from two independent experiments are shown. (g) Quantification of probe fluorescence intensities of the *ccn1l1* and *has3* probes in the anterior bud. Data are mean ± SD. *n* denotes the number of buds from individual embryos measured from two independent experiments. *P*-values as labeled (unpaired two-tailed Student's t-test).

(h-i) Effect of forskolin treatment on bud growth. (h) 3D-rendered OVs of *Tg(actb2:membrane-neongreen-neongreen)* embryos, treated with forskolin, imaged at 52 hpf. Scale bar: 50 μm. Representative images from two independent experiments are shown. (i) Quantification of bud length in embryos treated with forskolin at 52 hpf. Data are mean ± SD. *n* denotes the number of buds from individual embryos measured per condition from two independent experiments. *P*-value as labeled (unpaired two-tailed Student's t-test).

(j) 2D sections of OVs showing phospho-CREB, DN PKA-GFP and F-actin staining using anti-phospho-CREB[ser133] antibody, anti-GFP antibody, and phalloidin, respectively, at 56 hpf in sibling control or in DN PKA-GFP-expressing embryos following heat-shock induction. Scale bar: 50 μm. Representative images from two independent experiments are shown.

(k) 3D-rendered OVs from *Tg(hsp: DN PKA-GFP)* and sibling control embryos, both *Tg(βActin: utrophin-mCherry*, imaged at 56 hpf following heat-shock induction. Insets show 2D sections of the pillar or unfused buds from the regions in the white dashed boxes. Blue and yellow dashed lines indicate the anterior bud/pillar and the anterior-lateral bud/pillar, respectively. Scale bar: 50 μm. Representative images from two independent experiments are shown.

Extended movie 1: Mechanical epithelial cell recruitment during bud extension, related to Figure 4.

3D-rendered time-lapse images of the anterior bud obtained from *a Tg(actb2:membrane-neongreen-neongreen)* embryo. Each cell is marked by uniquely colored dot. Images were captured every 30 min. Time stamps indicate developmental time (hours post fertilization). Scale bar: 20 μm. Fig. 4b shows a cropped frame cropped from this movie.

Extended movie 2: Semicircular canal morphogenesis of wild-type embryo, related to Figure 5.

3D-rendered time-lapse images of the OV showing SCC formation of a wild-type embryo expressing membrane-NeonGreen. Images were captured every 30 min. Time stamps indicate developmental time (hours post fertilization). Scale bar: 50 μm.

Extended movie 3: Semicircular canal morphogenesis of *gpr126* $^{st49/st49}$ mutant embryo, related to Figure 5.

3D-rendered time-lapse images of the OV showing SCC formation of a *gpr126* $^{st49/st49}$ mutant embryo expressing membrane-NeonGreen. Images were captured every 30 min. Time stamps indicate developmental time (hours post fertilization). Scale bar: 50 μm.

Fig 3, Mori et al.,

a **Cell number in a bud** vs hpf; n=4 buds; Bud fusion (55–60 hpf)

b 45 hpf: Budding, 46 hpf, 47 hpf, 48 hpf: Extension, 49 hpf, 50 hpf, 51 hpf, 52 hpf; 20 μm

c Cell number in a bud; ○ 0 hpi, ○ 6 hpi; PBS p=0.005, HAase p=0.521, p=0.042; n=8 buds

d Cell number in a bud; ○ cont MO, ○ yap1 MO; p<0.001; n=10 buds

e cellular tensions: apical, lateral, basal; bud-adjacent; BM; Tip cell; Budding cell; Border cell; ECM pressure; cytocinches; adhesion

f **2D vertex model** — Budding cell trajectory, Tip cell; Cell aspect ratio vs Time (A.U.); Neighboring, Border, Budding, Border; a b

g Nuclear aspect ratio; Border, Tip; p<0.001; n=15 cells

h Yap Nuc/cyto ratio; Border, Budding, Tip; p=0.529, p<0.001, p<0.001; CV= 0.20, 0.29, 0.19; n= 23, 58, 25

i **2D vertex model** — Bud-ECM area (2D, ×10² μm²) vs Time (A.U.); with feedback, without feedback

j Bud-ECM volume (×10³ μm³) vs hpf; ○ cont MO, ○ yap1 MO; n=5 buds; p=0.215, p<0.001

k **2D vertex model** — Bud length (μm) vs Time (A.U.); with feedback, without feedback

l Bud length (μm) vs hpf; ○ cont MO, ○ yap1 MO; n=6 buds; p<0.001, p<0.001

m **2D vertex model** — Bud fusion time (h); Tip cell ECM production (%) vs Mechanosensitive ECM production (%); (yap1 KD); Robust bud fusion; (ccn1l1 moderate KD); 15h; No fusion (ccn1l1 strong KD)

n Bud growth rate (μm/h); control, 0.5 ng, 1 ng, 2 ng; ccn1l1 MO conc.; p<0.001, p<0.001, p=0.004, p<0.001, p<0.001, p<0.001; n=4, 8, 8, 8

o control MO; ccn1l1 MO conc. 0.5 ng, 1 ng, 2 ng; 50 hpf, 56 hpf, 72 hpf; 50 μm

Fig 4, Mori et al.,

a

Bud extension | Bud fusion | SCC formation

Wild type

gpr126^st49/st49

50 hpf | 60 hpf | 72 hpf | 50 μm

b

ccn1l1 | *has3* | *ccn1l1*, *has3*, *col2a1a*

Wild type

gpr126^st49/st49

low ▬ high | low ▬ high | 60 hpf | 50 μm

c

□ Bud extension: 50 hpf   ▨ Bud fusion: 60 hpf

ccn1l1
(n=7 buds)

Normalized intensity (A.U.)

p=0.010
p>0.999
p>0.999

WT   *gpr126^st49*

has3
(n=6 buds)

Normalized intensity (A.U.)

p=0.001
p>0.999
p>0.999

WT   *gpr126^st49*

d

● WT   ● *gpr126^st49/st49*

Bud length (μm)

p=0.992
p<0.001

50 hpf   60 hpf   n=5 buds

e

● WT   ● *gpr126^st49/st49*

Cell number in a bud

p=0.982
p<0.001

50 hpf   60 hpf   n=6 buds

f

● WT   ● *gpr126^st49/st49*

Bud-ECM volume at 60 hpf (×10³ μm³)

p<0.001

WT   *gpr126^st49*   n=6 buds

g

Bud extension | Bud interaction | Bud fusion

p-CREB^Ser133

p-CREB^Ser133, F-actin

50 hpf | 52 hpf | 60 hpf | 50 μm

h

control | *gpr126^st49/st49*

p-CREB^Ser133

p-CREB^Ser133, F-actin

60 hpf | 50 μm

i

ccn1l1 | *ccn1l1*, *col2a1a*

control

DN PKA

low ▬ high | 56 hpf | 50 μm

j

ccn1l1
(n=7 buds)

Normalized intensity (A.U.)

p<0.001

control   dnPKA

Fig 5, Mori et al.,

a Budding epithelium — Width / Height / HA-rich Bud-ECM

b Budding / Bud extension — 3D-rendered view (45 hpf / 50 hpf) / 2D view / Bud-ECM / 20 μm

c Cell aspect ratio (width/height) — Budding / Bud extension / p=0.028 / n=4 buds

d dorsal cells — Yap Nuc/cyto ratio / Pre-budding / Bud extension / p=0.095 / n=28 cells

e PBS injection / HAase injection — 0 hpi / 6 hpi / Bud-ECM / 46 hpf / 52 hpf / 20 μm

f Bud-ECM volume (x10³ μm³) — PBS / HAase / p<0.001 / n=7 buds

g 0 hpi / 6 hpi — Cell aspect ratio / PBS / HAase / p=0.024 / p=0.010 / p>0.999 / n=8 buds

h 0 hpi / 6 hpi — Bud length (μm) / PBS / HAase / p<0.001 / p<0.001 / p=0.126 / n=8 buds

i dorsal cells — Yap Nuc/cyto ratio / Before inj. / After inj. / p=0.412 / n=28 cells

j Pre-budding / Bud extension — anti-Yap1 / anti-Yap1,Nuclei / 40 hpf / 50 hpf / 10 μm

k Yap Nuc/cyto ratio — Pre-budding / Budding / non-budding / Bud extension / p<0.001 / p<0.001 / n=45 / 45 / 20

l PBS inj. / HAase inj. — anti-Yap1 / anti-Yap1,Nuclei / 3 hpi (53 hpf) / 10 μm

m Yap Nuc/cyto ratio — PBS inj. / HAase inj. / p<0.001 / n=32 / 32

n control MO / yap1 MO — Membrane / 72hpf / 50 μm

Extended Fig. 1, Mori et al.,

a Inner ear cell clusters

vcanb expression

b Tead-binding sites   start site

ccn1l1

-1.35   -0.6   0 kb

-1350 GCATCATCACAGGAATGTCATTCCAGCTGCTCTT-------
-600 AAAGAGCAAATGACCACCAGTTTCCACCATTCCTCAAAATA

c control MO   *yap1* MO

ccn1l1

ccn1l1, col2a1a   40 hpf   50 μm

d *ccn1l1*
at pre-budding stage
(n=5 buds)

p=0.734

Normalized Intensity (A.U.)

cont MO   *yap1* MO

e sibling control   *lmx1bb^{jj410/jj410}* mutant

ccn1l1

low   high

ccn1l1, col2a1a   50 hpf   50 μm

f Budding zone
(posterior bud)

ccn1l1
(n=5 buds)

p=0.002

Normalized Intensity (A.U.)

sibling cont   *lmx1bb^{jj410/jj410}*

g *lmx1bb*   *ccn1l1*   *lmx1bb, ccn1l1, col2a1a*

DMSO

RO4929097   50 hpf   50 μm

Extended Fig. 2, Mori et al.,

a

E-value:

| 2.4e-17 | 2.93e-17 | 2.71e-6 | 1.75e-5 |

zebrafish Ccn1l1   IGFBP — VWC — TSP-1 — CT   383 aa

b

IGFBP domain

	aa		% of Identity to Hs_CCN1
Hs_CCN1	24	SSCPAVCSCAPDAPQCPPGVGLQRDACGCCLVCARQLNEDCSRSEPCDHSKGLECNYGASSHGATRGICRA	
Dr_Ccn1	19	STCPAACHCPLEAPKCAPGVGLVRDGCGCCKVCAKQLNEDCSKTQPCDHTKGLECNFGASSTALKGICR-	69.57
Dr_Ccn1l1	28	GGCPLKCSCPSAPPSCPPGISSVLDHCGCRVCAROFNQDCSPTEPCDHIKGLCHLGAEGDPDRGICRA	53.62

VWC domain

	aa		% of Identity to Hs_CCN1
Hs_CCN1	100	CEYNSRIYQNGESFQPNCKHQCTCIDGAVGCIPLCPQELSLPNLGCPNPRLVKVTGQCCEEWVC	
Dr_Ccn1	95	CEYSSRIYQNGESFQPNCKHQCTCMDGAVGCVPLCPQELSLPALGCANPRLVKVPGRCCEEWVC	89.06
Dr_Ccn1l1	104	CELDGRIYQHGEDFQPSCEHQCTCVDGVVGCLPLCPHHISLPDWRCARRRLIRKPERCCPEWVC	59.38

TSP-1 domain

	aa		% of Identity to Hs_CCN1
Hs_CCN1	231	-VQTTGWSQCSRSCGTGISTRITNNNAECKLMKESRICEVRPCS	
Dr_Ccn1	219	-VQTTSWSQCSKTCGTGISTRVTNDNPECRLVKETRICEVRPCG	76.74
Dr_Ccn1l1	228	LLQITDWSPCSCGMGVSSRVTNSNPECRLVSETRLCQIRECG	67.44

CT domain

	aa		% of Identity to Hs_CCN1
Hs_CCN1	291	KSVTPVKFSYAGCSSLKKYRPRFCGSCVDGRCCSPQNTRTIRVKFRCQDGE--TFNKNVMMIESCKCTKHCP-	
Dr_Ccn1	279	KSPEPVRFTYAGCLSVKKYRPKYCGSCVDGRCCTPQLTRTVKMRFRCEDGE--TFSKNVMMIQSCKCNYNCP-	71.43
Dr_Ccn1l1	289	RPQKPVQISFAGCSTTRRYRLRSCGFCSDGRLCVPSATRTVRLHFHCPEAERHDFTRSLMWIQRCSCRHTHSR	40.00

c

Pre-budding / Bud extension: *ccn1l1*, *has3*, *vcanb*, *ccn1l1,has3,vcanb* — 40 hpf, 50 hpf, A P L, 50 µm

d

has3 (n=6 buds), *vcanb* (n=4 buds) — Normalized intensity (A.U.) — Pre-budding, Extension, Fusion

e

control Morpholino / *ccn1l1* Morpholino: *has3*, *vcanb*, *has3,vcanb, col2a1a* — low high — 40 hpf — 50 µm

f

has3 (n=5 buds) p=0.001, *vcanb* (n=5 buds) p<0.001 — Normalized intensity (A.U.) — cont MO, *ccn1l1* MO

Extended Fig. 3, Mori et al.,

a

Low BM adhesion

b

Low BM adhesion

Cell aspect ratio

····· Border cell (control)
— Border cell (Low BM)
— Tip cell (Low BM)

Time (A.U.)

c

Without cytocinches

d

Withoout cytocinches

Cell aspect ratio

····· Border cell (control)
— Border cell (without cytocinches)
— Tip cell (without cytocinches)

Time (A.U.)

e

Bud at 5h post budding

Tip 100%, Mech. t. 80%

Tip 80%, Mech. t. 100%

Tip 100%, Mech. t. 30%

Tip 30%, Mech. t. 100%

ECM production: Tip 0%, Mech. t. 100%

ECM production of Tip cell

high ⟶ low

f

ccn1l1 MO conc.

Cont MO   0.5 ng   1 ng   2 ng

50 hpf   20 μm

Extended Fig. 4, Mori et al.,

Supplementary Information – Theory note

In these notes, we detail the vertex models used to account for the bud morphogenesis, as well as analytical models to describe the role of the mechanotranduction feedback. These notes are organized as follows. In Section 1, we present the implementation of the vertex models. In Section 2, we focus on the elongation of budding cells during the recruitment process, in relation with YAP activation. In Section 3, we investigate the growth dynamics of the bud in presence of a feedback between YAP mechanotransduction and gel production.

1 Methods: Vertex model

1.1 Bud mechanics

As the bud is axisymmetric, we use here a 2D vertex model, representing a transversal cut across the bud in the xy plane. Because the radius of the bud is much smaller than the total size of the otic vesicle, we model the later as a planar surface at $y = 0$, from which the bud is growing. The system consists in N cells (corresponding to cells already recruited into the bud, and several rows of cells neighbouring the bud). The energy of a given cell i is taken to be:

$$e_i = (\gamma_a \ell_a + \gamma_b \ell_b + \gamma_l \ell_l) + K(a - a_0)^2 + \mathcal{U}_{\text{adh}} + \mathcal{U}_{\text{cytocinches}}, \tag{1}$$

where ℓ_a, ℓ_b and ℓ_l are the apical, basal and lateral lengths of the cell, respectively, and the γ's the respective mechanical tensions from the acto-myosin cytoskeleton. The second term imposes volume conservation, with a the 2D area of the cell, a_0 its homeostatis area and K a constant energy penalty for breaking volume conservation. To account for the corner-like shapes cells go through in vivo during the recruitment process, each cell is modelled by an hexagon, with 4 vertices corresponding to contact points with adjacent cells (2 on the apical side, and 2 on the basal side), and one additional vertex on both the apical and the basal sides, to allow cells to bend. The third term corresponds to an adhesion energy, for non-budded cells still in contact with the otic vesicle:

$$\mathcal{U}_{\text{adh}} = -G \times d_{\text{contact}} e^{-y/y_{\text{contact}}}, \tag{2}$$

where G is an adhesion energy per unit of contact length, d_{contact} is the portion of the cell's basal length that lies below some fixed threshold y_{contact}, and y is the average height of the cell's basal side. The exponential term allows for a smooth delamination of cells from the substrate, rather than a sharp threshold effect.

The last term in e_i accounts for cytocinches, actin-rich cables that were previously determined to constrict the bud and make it anisotropic, we applied a constricting force on all vertices that are part of the bud at a given timepoint (that is, that are not in contact with the horizontal substrate):

$$\mathcal{U}_{\text{cytocinches}} = \sum_{\text{basal vertex } j} k_c x_j^2, \tag{3}$$

where the sum runs other all the basal vertices of the cell, with x_j their horizontal position.

The total energy of the bud is then:

$$\mathcal{E} = K_{\text{ECM}}(A_{\text{ECM}} - A_0(t))^2 + \sum_{\text{cell } i=1}^{N} e_i, \tag{4}$$

where the first term imposes the volume of the ECM below the bud, with its equilibrium 2D area $A_0(t)$ being updated at each timestep to account for ECM production in the bud.

We then simulate the bud's mechanics through a Monte Carlo algorithm. At each timestep, we try optimizing the position and shape of each cell by randomly moving their vertices, and checking if the proposed change leads to a decrease in the total energy of the bud. In the initial state, cells on the horiizontal line are placed such that the epithelium is tension-free: each cell has a width $w = \sqrt{a \frac{\gamma_l}{\gamma_a + \gamma_b - G}}$ and height $h = a_0/w$, where a_0 is the homeostatic 2D area of a cell. The bud is assumed to have an axis of symmetry at its center, which in our simulation corresponds to the right-most vertices (the other half of the bud being obtained by axial symmetry). The position of the left-most vertex is kept constant throughout the simulation. All the codes were implemented in Julia.

1.2 Values of parameters

Cytoskeletal tensions can be estimated by considering cellular aspect ratios. At the start of the bud extension phase, when pressure from the ECM is still low, cells are likely close to their prefered aspect ratio, given by $\epsilon_0 = w_0/h_0 = \gamma_l/(\gamma_a + \gamma_b)$, with w_0 and h_0 the width and thickness of cells in their stress-free state. We measured that their thickness was approximately $h = 9\,\mu m$ and their width $w = 5\,\mu m$, resulting in an aspect ratio $\epsilon = w/h \simeq 0.55$. We chose γ_l, γ_a and $\gamma_b = 1$ to be all equal as this choice would yield $\epsilon_0 = 0.5$. In practice, experimental observations show that myosin is preferentially localized on the basal side; however, it is difficult to discriminate how much this basal myosin contributes directly to contraction, rather than simply to cell stiffness. In addition, cytocinches also contribute to contraction on the basal side. Therefore, we decided to keep $\gamma_b = \gamma_a = 1$ rather than considering more complex scenarios, and tuned basal contraction only through cytocinches.

We chose to work with length units so that the cell cross-section is $a_0 = 1$. Experimentally, the cell cross-section at the start of the bud extension phase was roughly $a_0 = 50\,\mu m^2$, so a unit length in our simulation corresponds to a distance $\sqrt{a_0} = 7\,\mu m$ in experiments.

The last two parameters, G and k_c, representing substrate adhesion and cytocinches contractility, respectively, cannot be directly determined experimentally. Intuitively, the corresponding forces must be of the same order as the in-plane tissue tension $\gamma_a + \gamma_b$, and their precise values could be obtained by fitting the length and radius of the bud for different parameter values on experimental data. However, the 2D and highly simplified nature of our model makes such a fitting procedure rather questionable. Instead, we chose to set $G = 0.5$ and $k_c = 0.2$ in most cases, and we tested the impact of changing these values on the bud morphology.

In the following, we considered two separate cases for the dynamics of the ECM production, defined by $A_0(t)$. As a first step, we externally prescribed A_0 to slowly increase at a constant rate, to focus on cell recruitment alone. Then, we considered the role of the mechanotransduction feedback, and assume that the growth rate $\dot{A}_0$ was proportional to the number of recruited cells.

When testing the effects of a low adhesion, we set $G = 0.3$. In simulations without cytocinches, we set $k_c = 0.0$.

2 Cell elongation during the recruitment process

2.1 Link between cellular aspect ratio and mechanical forces

At the lowest possible degree of description, apical, basal and lateral cytoskeletal tensions create a spring-like restoring force, causing each cell to spontaneously return to a prefered aspect ratio. For a rectangular 2D cell, this prefered value can be determined as:

$$\epsilon_0 = \frac{w_0}{h_0} = \frac{w_0^2}{a_0} = \frac{\gamma_l}{\gamma_a + \gamma_b}, \tag{5}$$

where ϵ_0 is the prefered aspect ratio, w_0 the prefered width, h_0 the height of the cell, a_0 its 2D cross-section, and γ_l, γ_a, γ_b are the lateral, apical and basal tensions. As a result, a deformed cell, with aspect ratio $\epsilon > \epsilon_0$ will exert a pulling force on its environment, given by:

$$\zeta = \gamma_a + \gamma_b - \gamma_l \epsilon^{-1} \simeq (\gamma_a + \gamma_b)\frac{w - w_0}{w_0}, \tag{6}$$

so that this pulling force is proportional to the relative elongation of the cell. Therefore, in what follows, we monitor the aspect ratio $\epsilon = \frac{w^2}{a}$ as a readout of mechanical forces felt by the cells. When $\epsilon > \epsilon_0$, the cell is under stretch, and this can typically result in nuclear relocalization of Yap.

2.2 Cell aspect ratio during recruitment

Based on this result, we monitor the cellular aspect ratio of all cells in our simulations. First, we assume that bud ECM production is prescribed independently from cell recruitment, and we impose that $A_0 = rt$, with r a constant growth rate. As cells are not rectangular, we define numerically w as the average of the apical and basal sides of the cell, and compute the aspect ratio through $\epsilon = w^2/a_0$. While other definitions of the aspect ratio are possible, this one directly relates to mechanical forces as it results from the balance between apical, basal, and lateral tensions with external forces from the surrounding tissue.

Numerically, we find that cells undergo a transient stretching as they are recruited into the bud: border cells, that are partially recruited and partially still attached to the basement membrane exhibit an overshoot in their aspect ratio, and similarly for budding cells (bud cells that are adjacent to a border cell). On the other hand, tip cells showed a lower aspect ratio. Looking at trajectories of individual cells, we find that they "flow" towards the bud, and start stretching as they turn the corner. Bud adjacent cells (cells neighboring border cells, but still attached to the basement membrane) displayed a slight compression. Intuitively, we would expect that the flow of cells converging around the bud would result in an in-plane tension that decreases with distance from the bud, resulting instead to a slight stretching of bud adjacent

cells. However, our 2D simulation does not model the fluidity (through in-plane rearrangements and friction with the substrate) of the epithelium around the bud. Therefore, we assume that this slight compression of bud adjacent cell stems from a small simulation artefact.

We hypothesized that the increase in aspect ratio of border cells was the result of a tug-of-war between adhesion forces, which pull the epithelium towards the basement membrane, and tugging forces from the cytocinches. To test this idea, we considered the effects of either reducing focal adhesion, or removing forces from cytocinches. The former led to over-recruitment of cells. The latter, on the other hand, fully suppressed the converging flow of cells towards the bud. Instead, the bud appeared to radially "push" the epithelium away from the center. In both cases, the aspect ratio of cells remained low and no marked difference was found between border cells and tip cells.

3   Role of the feedback loop

Next, we consider the impact of this mechanical patterning on the bud extension dynamics. Rather than prescribing the volume increase of the bud ECM, we now assume its rate depends on the number of recruited cells. We consider the following model:

$$\frac{\mathrm{d}A_{\mathrm{ECM}}}{\mathrm{d}t} = k(aN_{\mathrm{tip}} + bN_{\mathrm{recruited}}), \tag{7}$$

where bud ECM production is the sum of two contributions: a constant rate from tip cells, and a variable contribution from recruited cells that grows as the bud recruits more and more cells. Setting $b = 0$ allows us to mimick Yap morpholino conditions, while $b = a = 1$ would be closer to wild type conditions. Here, the quantity k is a unit of ECM volume production per unit of time, setting the unit of time of our simulation. Here we choose $k = 13\,\mu\mathrm{m}^3/\mathrm{h}$, with one hour corresponding to 30000 simulation steps.

Numerically, we find that the mechanical feedback increases the bud extension rate, as well as the cell recruitment rate. The tell-sign of the feedback is the growth dynamics of the bud-ECM volume A_{ECM}: in absence of any feedback, it grows linearly and slowly, while in its presence it grows supra-linearly. This observation is fully in line with experimental findings. In particular, in the gpr126 mutant, the bud ECM volume displays a clear non linear growth as the feedback continues unimpeded in absence of bud fusion.

To study the impact of the feedback loop on bud fusion, we define bud fusion as the moment when the bud reaches 30 μm in length. The start of our simulation corresponds roughly to a timepoint of 50 hpf. If the bud does not reach this value after 22 hours (72 hpf), we assume that the bud fails to fuse.

Extended Table 1

Target gene	amplifier type	Probe Name	Sequence
ccn1l1	B3	ccn1l1_B3_1	TTTTTTTTTTTTTTTTTTTTTTTTTTTTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_2	GTCCCTGCCTCTATATCTTTGGAAATTGTAACACATTTTTATTTG
ccn1l1	B3	ccn1l1_B3_3	ATTCACCTCAAAAAATTGAGGTTTGTTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_4	GTCCCTGCCTCTATATCTTTTATGACTCTTCACAAGAACTGAAGA
ccn1l1	B3	ccn1l1_B3_5	ATATTTACAGCGATATTTGTACAAATTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_6	GTCCCTGCCTCTATATCTTTTACACTATATACAGTAAATCTGGAT
ccn1l1	B3	ccn1l1_B3_7	GTGTGTGTAAATTATATAAACCAAATTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_8	GTCCCTGCCTCTATATCTTTAGCTTTGCAGTCATACCGTAATAAA
ccn1l1	B3	ccn1l1_B3_9	TCTTCCACCTGTGTCATTACCACATTTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_10	GTCCCTGCCTCTATATCTTTATAATAAAACGGTAATATTTAAAAC
ccn1l1	B3	ccn1l1_B3_11	TATGCTGGGTTTAAATGTTTTGTAGTTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_12	GTCCCTGCCTCTATATCTTTTGAGGGCAGTGTGGATCTGTTGCTG
ccn1l1	B3	ccn1l1_B3_13	TCAGCTGGTGTGTTCAGGCCTCGTCTTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_14	GTCCCTGCCTCTATATCTTTCTGTGTGTGTGTCTGCAGCTGCAGC
ccn1l1	B3	ccn1l1_B3_15	CTCTGGACAGTGGAAATGCAGACGGTTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_16	GTCCCTGCCTCTATATCTTTTGTCCGAGTTGCAGATGGGACACAC
ccn1l1	B3	ccn1l1_B3_17	GGCGAGTAGTGGAACATCCAGCAAATTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_18	GTCCCTGCCTCTATATCTTTTGATCTGCACAGGTTTCTGTGGCCG
ccn1l1	B3	ccn1l1_B3_19	GTGTTGACGCCACACTCTCGGATCTTTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_20	GTCCCTGCCTCTATATCTTTCAGAGTCTGGTTTCGCTTACTAGCC
ccn1l1	B3	ccn1l1_B3_21	TCCACAGGTGGCAGAGCATGGCGACTTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_22	GTCCCTGCCTCTATATCTTTATCGGTGATCTGCAGCAAGCAGGTG
ccn1l1	B3	ccn1l1_B3_23	GATAAGGGGCTCCTGCACTGCTGTCTTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_24	GTCCCTGCCTCTATATCTTTAGGAAGCAGATGGAGTTACCAGTAG
ccn1l1	B3	ccn1l1_B3_25	GGCACTGGATTCACCTCAGCAATGCTTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_26	GTCCCTGCCTCTATATCTTTTTGTCGTCATCACAGACCCACTCCG
ccn1l1	B3	ccn1l1_B3_27	GCGCCAGTCGGGCAGAGAGATGTGGTTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_28	GTCCCTGCCTCTATATCTTTTGGGCACAGTGGCAGACAGCCGACC
ccn1l1	B3	ccn1l1_B3_29	AGTCCTCTCCGTGCTGGTAAATCCGTTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_30	GTCCCTGCCTCTATATCTTTCATCCAGCTCACACGGGCGACCCTG
ccn1l1	B3	ccn1l1_B3_31	CCTAGGTGGCAGCGCAGTCCTTTAATTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_32	GTCCCTGCCTCTATATCTTTTGGTCGCAGGGTTCAGTAGGGCTGC
ccn1l1	B3	ccn1l1_B3_33	GTGGTCCAAAACCGAGCTGATACCTTTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_34	GTCCCTGCCTCTATATCTTTAGGACACGAGGGGGGAGCGGAAGGA
ccn1l1	B3	ccn1l1_B3_35	CCCATGACAGCAGCACCAGTGTGAATTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_36	GTCCCTGCCTCTATATCTTTAATGTCCTTGCCTTAATATGATCCT

ccn1l1	B3	ccn1l1_B3_37	CAGAAAGTGTCTTCTGGTTTAGACCTTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_38	GTCCCTGCCTCTATATCTTTGTTCTGGATCTGTCAATTCACAAGG
ccn1l1	B3	ccn1l1_B3_39	TGGCTAAATGCAAAAAACTGCAAAGTTCCACTCAACTTTAACCCG
ccn1l1	B3	ccn1l1_B3_40	GTCCCTGCCTCTATATCTTTGTGTGCTGCTGGATTCCCTCAGTGA
has3	B2	has3_B2_1	TCAGATAAGCCCCGACAGCATTTAGAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_2	CCTCGTAAATCCTCATCAAAAGCATATAAACCATGCTAGGCTTCG
has3	B2	has3_B2_3	ACGTTTGGCCACTATGGCCAGGTAGAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_4	CCTCGTAAATCCTCATCAAACACCAGGAGTATAAGCCAGTAAGAG
has3	B2	has3_B2_5	AGTCTCGGTAAAAGGTTCTTGAACCAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_6	CCTCGTAAATCCTCATCAAAGCAGTAAATGGTGTAAACCACTCCA
has3	B2	has3_B2_7	TCCAATAAGGTTTACTACCATCTTTAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_8	CCTCGTAAATCCTCATCAAACCTGCCTGATGTGCCCCATCCTGCT
has3	B2	has3_B2_9	CAGGCTTGACATGTAGAGCAGTGAGAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_10	CCTCGTAAATCCTCATCAAAGAGTGACATGAAGATCATGACAAGG
has3	B2	has3_B2_11	ACCCACCAGCTGGACCGTCAACAAGAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_12	CCTCGTAAATCCTCATCAAAGAGTAAGATGTTCCACAACCGGCCT
has3	B2	has3_B2_13	GGGGAAGAACCCAGTAACCACAGATAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_14	CCTCGTAAATCCTCATCAAAATAGGTCATCCACAGGCTGTGCTTG
has3	B2	has3_B2_15	AGACTTGCTCCAGCGGGTCTGCTGAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_16	CCTCGTAAATCCTCATCAAAGAGCCAACGCAGGTATTGTGTTGGG
has3	B2	has3_B2_17	TTTGAAGCCAAAGCTAAGGACGCGGAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_18	CCTCGTAAATCCTCATCAAAGGTGAGATGTCGATCATCACCAAAA
has3	B2	has3_B2_19	CTCTAAGAACTGCTGAAGGAGAGAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_20	CCTCGTAAATCCTCATCAAAACGGTACATTCCTAGAGGTCCACTG
has3	B2	has3_B2_21	CTCCACATTGAAGGCCATCCAATAGAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_22	CCTCGTAAATCCTCATCAAAGACACTGCTCAGGAAGGAGATCCAG
has3	B2	has3_B2_23	TCCTCCCACATCAGGATCCTCCTCTAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_24	CCTCGTAAATCCTCATCAAAAATCTTCAACATCTCAATAGTGCAT
has3	B2	has3_B2_25	GTAATCCACTGTATCTCCAAGTGCTAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_26	CCTCGTAAATCCTCATCAAAGAAGGCTGTATACATCACTTCCCTC
has3	B2	has3_B2_27	GCTCCGCACCACTCTGGCCACTCGTAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_28	CCTCGTAAATCCTCATCAAAAGCCTCTTCGGCATGCACCGAGCCT
has3	B2	has3_B2_29	ATTCCCTTTCCACACCCACACAGCCTAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_30	CCTCGTAAATCCTCATCAAACTGCTCCACTCCCCCCATCACCTCC
has3	B2	has3_B2_31	GCCATCCACCACCAACACCACTTTCAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_32	CCTCGTAAATCCTCATCAAACCCGGGAAAAGAAATGCGGCGAATG
has3	B2	has3_B2_33	ATAGGCTGCAATGCAGAGGGCCACAAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_34	CCTCGTAAATCCTCATCAAAGCGGCGCAGGTGCTGCGGTCTGGAG
has3	B2	has3_B2_35	ACTCTGCAGGAGAAGGTGGAGGGATAAATCATCCAGTAAACCGCC

has3	B2	has3_B2_36	CCTCGTAAATCCTCATCAAAAAATGCACCGTACAAGCCAAAAGAC
has3	B2	has3_B2_37	CACATAGGCTAGTAGGATTGCGAAAAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_38	CCTCGTAAATCCTCATCAAACACCACTGCTGCAAATAAGGTGGTG
has3	B2	has3_B2_39	CTGAAGAAGGAAAAAAACATGACTTAAATCATCCAGTAAACCGCC
has3	B2	has3_B2_40	CCTCGTAAATCCTCATCAAAGTGTAGAAAGCAAAATCATTACTGC
lmx1bb	B2	lmx1bb_B2_1	TAGGCTCAGTGGCTAGCATTGTCGCAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_2	CCTCGTAAATCCTCATCAAACACTGCAGCTATATATATTTATTTA
lmx1bb	B2	lmx1bb_B2_3	TAATTTTGGTTTCTTCATGTAGCAAAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_4	CCTCGTAAATCCTCATCAAAGTTTAAGTTTTCCTAATGCATGTTC
lmx1bb	B2	lmx1bb_B2_5	CTTTATTGCATGCAGATTCATTTCAAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_6	CCTCGTAAATCCTCATCAAATTTGTTTCCGTGGATGCCAGGCAAA
lmx1bb	B2	lmx1bb_B2_7	ACGGAAGTTACTTAAGTTTATTGAAAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_8	CCTCGTAAATCCTCATCAAATTATCTACTTAATATTCTAATCATC
lmx1bb	B2	lmx1bb_B2_9	TCAAATGCTTCCATCTGACAAGCACAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_10	CCTCGTAAATCCTCATCAAAGTCTATACAGTCGAAGCTCTCCAAA
lmx1bb	B2	lmx1bb_B2_11	CCATGTGCCACCAACAACCATGCCAAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_12	CCTCGTAAATCCTCATCAAATTCAAAGTTACTTAAATCACCTTTA
lmx1bb	B2	lmx1bb_B2_13	TGTTTGGCATTTTAGTGGCGCTCAAAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_14	CCTCGTAAATCCTCATCAAAGGCGAAGCAAATTCGATTTCTGCTC
lmx1bb	B2	lmx1bb_B2_15	ATCACTTTGCTTTTACTATATGTGTAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_16	CCTCGTAAATCCTCATCAAACTTTCCTAGCTTTCGGGTCTAGTCT
lmx1bb	B2	lmx1bb_B2_17	ACAGAACTGAACAGATTGAAACAAAAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_18	CCTCGTAAATCCTCATCAAAAAACAGAACAGAATCGATTGAAACA
lmx1bb	B2	lmx1bb_B2_19	GCATTTTTCCAATAAAATCTCTCCAAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_20	CCTCGTAAATCCTCATCAAATCCTTCAGATTCTCTAGTCCATGTT
lmx1bb	B2	lmx1bb_B2_21	TTTGTTTTTCCTTTTAAATAGCATGAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_22	CCTCGTAAATCCTCATCAAAGGTTATAAAATAGCATTTTAGGTTA
lmx1bb	B2	lmx1bb_B2_23	ATCCCGCTTTTGCATATTTGCAGAGAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_24	CCTCGTAAATCCTCATCAAACTCGCTTCAGAACACCTTATATTAC
lmx1bb	B2	lmx1bb_B2_25	GGTGTGACCAGCTGTCGCCATTTAGAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_26	CCTCGTAAATCCTCATCAAACGCACGCCATCACGCCGACCAGGGG
lmx1bb	B2	lmx1bb_B2_27	TCAGTGTATTCAGATTAGTTGTGTTAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_28	CCTCGTAAATCCTCATCAAAAAGTTTAATCAGTCACTCTACCTAG
lmx1bb	B2	lmx1bb_B2_29	GTCTGACCTTTAAAAAAAGAATAAAAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_30	CCTCGTAAATCCTCATCAAAATCATCTCAGTGAATCTCTTCACAT
lmx1bb	B2	lmx1bb_B2_31	TCCAGCCGCAGCCCATTTCTGGGAAAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_32	CCTCGTAAATCCTCATCAAAATCCACACACACATTCACACACACA
lmx1bb	B2	lmx1bb_B2_33	TCATATAATTTAGTACCATTTGCTCAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_34	CCTCGTAAATCCTCATCAAACACCCAATGACGGTTGGGTTTAGGG

lmx1bb	B2	lmx1bb_B2_35	ATCTGTTATAAACGCACCACTGAACAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_36	CCTCGTAAATCCTCATCAAATGGGAAGTTTTTTGTTTGTTATGAG
lmx1bb	B2	lmx1bb_B2_37	TTGGTTGCCAATAAGCTTGTTCCTCAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_38	CCTCGTAAATCCTCATCAAACATTTAAATACCTCAGTGTTGTAGC
lmx1bb	B2	lmx1bb_B2_39	TTGCAATTTAATCAATAAATAATTTAAATCATCCAGTAAACCGCC
lmx1bb	B2	lmx1bb_B2_40	CCTCGTAAATCCTCATCAAATTATTTATTAATTAGTCTGTTACTT
vcanb	B1	vcanb_B1_1	TTTTTTTTTTTTTTTTTTTTTTTTTTTAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_2	GAGGAGGGCAGCAAACGGAATTAATATTCTAATAGCGCTTTATTA
vcanb	B1	vcanb_B1_3	TATATTTTGATGAGCATATGATTGTTAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_4	GAGGAGGGCAGCAAACGGAAGAGTGTACAATGGCGTTCTCTGAAG
vcanb	B1	vcanb_B1_5	TCACAGCTGCGTTGACTTGATCCTCTAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_6	GAGGAGGGCAGCAAACGGAATTCTACTATATACATTTAAGCCCCG
vcanb	B1	vcanb_B1_7	TTCTGCTGTTTTTGACCAGCGATGGTAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_8	GAGGAGGGCAGCAAACGGAAGTACTGAGATTCACTGAAGCGTCTC
vcanb	B1	vcanb_B1_9	CAGGACACTGTGCCCTTCTTACAGGTAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_10	GAGGAGGGCAGCAAACGGAAAAGGTCAGGTGGTAGTTGCAGGGCA
vcanb	B1	vcanb_B1_11	GACGACACTCTCTTTCTGCTGCCTCTAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_12	GAGGAGGGCAGCAAACGGAAACGTCCGCCGGTGGGTGAAGTACTT
vcanb	B1	vcanb_B1_13	GCCGTTTTCACACACACTGACCGAGTAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_14	GAGGAGGGCAGCAAACGGAAAAGTGCGTGATCTTGCAAATCCACC
vcanb	B1	vcanb_B1_15	GCAATAGTGATGGGGCCACTGGAATTAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_16	GAGGAGGGCAGCAAACGGAAACAGACGTTGAGATGAGAGGCTCCT
vcanb	B1	vcanb_B1_17	CAAATGCTTTAGTTGGTTCATTTGCTAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_18	GAGGAGGGCAGCAAACGGAACAACACTAGTTTCAACAGTTTGTTG
vcanb	B1	vcanb_B1_19	ACTAGCAGGTTCCCTTGGGTCTTTGTAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_20	GAGGAGGGCAGCAAACGGAAATGAGTACTTGTTGAGGAAGTTGTA
vcanb	B1	vcanb_B1_21	CCACCTATTTCAGTCTCGTCGGTAGTAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_22	GAGGAGGGCAGCAAACGGAAACCATATCTGGGTCAGTAGAAGCTG
vcanb	B1	vcanb_B1_23	CAGTTTTGTAGGAAGATGTAAGACCTAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_24	GAGGAGGGCAGCAAACGGAACAGCTTCGGCCAGAACTATTACTAT
vcanb	B1	vcanb_B1_25	TTCATCTGTTTTAGGGGAAGATACATAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_26	GAGGAGGGCAGCAAACGGAAACCTTCAAATTCTGACATAAGAATT
vcanb	B1	vcanb_B1_27	AAATCCAGAACACTTTCAACTGATTTAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_28	GAGGAGGGCAGCAAACGGAATTTGTATCCTTGATGTCGACTATGA
vcanb	B1	vcanb_B1_29	CGCTGATGCTTCCCTCAGCACTACTTAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_30	GAGGAGGGCAGCAAACGGAAATTCCCCATGAGGTGTATCCAAAGG
vcanb	B1	vcanb_B1_31	TGGTGCCATGCTTGCAGAGAACATTTAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_32	GAGGAGGGCAGCAAACGGAAAGGTGGTTCTTCTATGGTGTCATCG
vcanb	B1	vcanb_B1_33	CCGTCCGACAACCAGCCGAAATCGCTAGAAGAGTCTTCCTTTACG

vcanb	B1	vcanb_B1_34	GAGGAGGGCAGCAAACGGAATGGTCCATGCCGTCTCTCCAGGCGG
vcanb	B1	vcanb_B1_35	CGGCCATCCATCCGGCGTCACACTGTAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_36	GAGGAGGGCAGCAAACGGAACGAATCCGTCCTCGTATGCTGAATA
vcanb	B1	vcanb_B1_37	GCTGGGCACTGACACCCGGCCCATATAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_38	GAGGAGGGCAGCAAACGGAACTCTTGGCCCATTTTGACGATGCCG
vcanb	B1	vcanb_B1_39	ATGTGCTTCACGTCCAACAACATCATAGAAGAGTCTTCCTTTACG
vcanb	B1	vcanb_B1_40	GAGGAGGGCAGCAAACGGAATCAAACTTATTTTTCCAGTTCCTGA
gpr126	B4	gpr126_B4_1	TCTGAATGTCCAGACGCAATACAGAATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_2	CCTCAACCTACCTCCAACAACTTCATTAGGAAATTATCTTACTGT
gpr126	B4	gpr126_B4_3	ATATGAAGGTCTGAATGTGTTAGATATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_4	CCTCAACCTACCTCCAACAAGGTGTATAGCTGTTGCTTTCCGTTT
gpr126	B4	gpr126_B4_5	GGGAAAAATGCAAGCTCCCAACCATATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_6	CCTCAACCTACCTCCAACAATTGTTCTTTGAGCCCCCAACAGAGG
gpr126	B4	gpr126_B4_7	ATATTTACAGATCACATAGTCAGTAATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_8	CCTCAACCTACCTCCAACAAAGCTGAAAATCACAGTTTGTAAAAT
gpr126	B4	gpr126_B4_9	GACCTGGTGCACAGGAAGAATGGAGATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_10	CCTCAACCTACCTCCAACAAGGAAGGCTCTCCGTCAGAAGGGCTG
gpr126	B4	gpr126_B4_11	TTAGTGGCAGTCTTGCTCCAGTCTGATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_12	CCTCAACCTACCTCCAACAATTATCTGCCAAGCGTAACTTTCCAC
gpr126	B4	gpr126_B4_13	TATCCTCTCGCAGTGTCCGGTTGCTATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_14	CCTCAACCTACCTCCAACAAGCTTGCCGTTACGTCCACAGATTTG
gpr126	B4	gpr126_B4_15	CCATCCCACGATGCAAAACTTGAGGATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_16	CCTCAACCTACCTCCAACAAGTATCTCCGAATGTATGTGTTGAAG
gpr126	B4	gpr126_B4_17	CGCCGCAGCTTCTCAAAAGCGATGTATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_18	CCTCAACCTACCTCCAACAAGTGAGCAGTGTTGCCGCTGAAAAAA
gpr126	B4	gpr126_B4_19	TGAAATCCCAAAACACACACTGAGGATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_20	CCTCAACCTACCTCCAACAATCGGATCTTTCTGATACTCCAAATG
gpr126	B4	gpr126_B4_21	GACCACAGCCAATGCATTATGTGCCATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_22	CCTCAACCTACCTCCAACAACGAGTCAAAATCAATCTGAGGGTCT
gpr126	B4	gpr126_B4_23	GAAATGATGGTGACCACAGTGCTGGATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_24	CCTCAACCTACCTCCAACAAAGGGTGGCGTTGATTTTGGCTATGT
gpr126	B4	gpr126_B4_25	CGTTTCGATCTTTGTAAGGGAAACAATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_26	CCTCAACCTACCTCCAACAAGGACATACTGAGTGACTGTGGGTCT
gpr126	B4	gpr126_B4_27	CCTTGAGAGAACAGTGATAAGCTGAATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_28	CCTCAACCTACCTCCAACAAACTGATGCCATCAACATAAACTCTA
gpr126	B4	gpr126_B4_29	CAACCTGCGGTTCCTGGTGAAAGAGATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_30	CCTCAACCTACCTCCAACAAGTTATGGCAGTACTGCAGCTGAGCG
gpr126	B4	gpr126_B4_31	ATCCGCCACTCTGCAGTGTGTAAATATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_32	CCTCAACCTACCTCCAACAAGAGAGGCCGAGCAGATGCTGGAGAA

gpr126	B4	gpr126_B4_33	GGTCTCTGTGGCCTTCTGTGCAGTGATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_34	CCTCAACCTACCTCCAACAAGGCGATCTCAAAGCACACCGTGAAG
gpr126	B4	gpr126_B4_35	AAGCCGCAAAATTTGGCATCGCTGGATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_36	CCTCAACCTACCTCCAACAACCGGTCTTGACGACAACGCGGTCAT
gpr126	B4	gpr126_B4_37	AAAGGCAGATCAGCAGCAGGAAAACATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_38	CCTCAACCTACCTCCAACAACAAGAGTATTCTGGAACTTCCATCT
gpr126	B4	gpr126_B4_39	TAAATGTCGGTTTTTTGACAGATGAATTCTCACCATATTCGCTTC
gpr126	B4	gpr126_B4_40	CCTCAACCTACCTCCAACAACTGCGCTCCAGGCAGACAGATGCTG
col2a1a	B4	col2a1a_B4_1	GATGACATCAAGCCATGTTTCTGTCATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_2	CCTCAACCTACCTCCAACAATCATTGGTAGTGCTTGCATGTTCGG
col2a1a	B4	col2a1a_B4_3	AGTGTTTTGCATCCTCACAGTTATTATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_4	CCTCAACCTACCTCCAACAATCGTCATTGGTATTTTTCCATGTCT
col2a1a	B4	col2a1a_B4_5	GTACGCCATATGTAAAGCGGCTGTTATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_6	CCTCAACCTACCTCCAACAACCTCTGCTCTGATCTCCACATCATT
col2a1a	B4	col2a1a_B4_7	GAACCACACGTGCTTCTGGGCCTTGATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_8	CCTCAACCTACCTCCAACAACTTGCTTGTCCACCAGTTCTTGCGT
col2a1a	B4	col2a1a_B4_9	TGCCCGTTGATGGACTTCAGAGTGGATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_10	CCTCAACCTACCTCCAACAATCCACCTCTACATCGTGTTGTCTGA
col2a1a	B4	col2a1a_B4_11	GCTGACCATTAGAACCATCCTTACCATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_12	CCTCAACCTACCTCCAACAACAGGTCCAACAGGTCCTGAAGGTCC
col2a1a	B4	col2a1a_B4_13	AGGTCCTTGTGGGCCATTCTCTCCTATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_14	CCTCAACCTACCTCCAACAAGTCTCCCTGTTTGCCAATTGGACCA
col2a1a	B4	col2a1a_B4_15	CCTCCTGGAGCTCCCTGTTTGCCAGATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_16	CCTCAACCTACCTCCAACAATCACCAGAAGGTCCTGGAAGTCCAG
col2a1a	B4	col2a1a_B4_17	GACCATCCTTACCAGAAGGACCTGCATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_18	CCTCAACCTACCTCCAACAAGGCCGGCAGCACCAGGGTTACCGTT
col2a1a	B4	col2a1a_B4_19	GCCATCAGCACCAGGGGGTCCAGCAATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_20	CCTCAACCTACCTCCAACAAGCCAGCAGGTCCAGGTGCACCAATC
col2a1a	B4	col2a1a_B4_21	CCAACAGCTCCTCTCTCACCAGGCAATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_22	CCTCAACCTACCTCCAACAACCTTGAAGACCGGGAGGACCTTGAG
col2a1a	B4	col2a1a_B4_23	CGGGGGGCTCCAGTAGGTCCAGGCAGATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_24	CCTCAACCTACCTCCAACAACCTGGAAGCCAGATGGACCAGGTTG
col2a1a	B4	col2a1a_B4_25	AGGTGGTCCAGGGCGGCCATCTTCAATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_26	CCTCAACCTACCTCCAACAAAGGGGCACCAGTCGCTCCAACTTTT
col2a1a	B4	col2a1a_B4_27	GGAGGTCCAAGAGGTCCGGCAGAACATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_28	CCTCAACCTACCTCCAACAAGGCTCTCCTCTTGGTCCTCTCTTGC
col2a1a	B4	col2a1a_B4_29	CAGATCCTTTAGCTCCAGGAATACCATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_30	CCTCAACCTACCTCCAACAACAGTACCAGGGTTACCGGATGCACC
col2a1a	B4	col2a1a_B4_31	TGCACCAGGGCGACCTCTCTCACCGATTCTCACCATATTCGCTTC

col2a1a	B4	col2a1a_B4_32	CCTCAACCTACCTCCAACAACAGACCACGAGGACCCATTGGTCCA
col2a1a	B4	col2a1a_B4_33	CTTCCGGGTTTTCCAGGGGGGCCGGATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_34	CCTCAACCTACCTCCAACAAGGGCCACGGGGTCCAAGGGCACCAG
col2a1a	B4	col2a1a_B4_35	GTCCGGGTGGTCCAGGGGGTCCTGGATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_36	CCTCAACCTACCTCCAACAATTCCAGGTGTACCAGGCTCGCCATC
col2a1a	B4	col2a1a_B4_37	CTCTCCAGGTGAAATGATGGGATTGATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_38	CCTCAACCTACCTCCAACAAACAGTCATTCAGATCCTCACAGATG
col2a1a	B4	col2a1a_B4_39	AATCTGAACATGTCTAAAGATTAGAATTCTCACCATATTCGCTTC
col2a1a	B4	col2a1a_B4_40	CCTCAACCTACCTCCAACAATGCAGGTCCTAAGGGGTGAAAGTCG
col2a1a	B2	col2a1a_B2_1	GATGACATCAAGCCATGTTTCTGTCAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_2	CCTCGTAAATCCTCATCAAATCATTGGTAGTGCTTGCATGTTCGG
col2a1a	B2	col2a1a_B2_3	AGTGTTTTGCATCCTCACAGTTATTAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_4	CCTCGTAAATCCTCATCAAATCGTCATTGGTATTTTTCCATGTCT
col2a1a	B2	col2a1a_B2_5	GTACGCCATATGTAAAGCGGCTGTTAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_6	CCTCGTAAATCCTCATCAAACCTCTGCTCTGATCTCCACATCATT
col2a1a	B2	col2a1a_B2_7	GAACCACACGTGCTTCTGGGCCTTGAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_8	CCTCGTAAATCCTCATCAAACTTGCTTGTCCACCAGTTCTTGCGT
col2a1a	B2	col2a1a_B2_9	TGCCCGTTGATGGACTTCAGAGTGGAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_10	CCTCGTAAATCCTCATCAAATCCACCTCTACATCGTGTTGTCTGA
col2a1a	B2	col2a1a_B2_11	GCTGACCATTAGAACCATCCTTACCAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_12	CCTCGTAAATCCTCATCAAACAGGTCCAACAGGTCCTGAAGGTCC
col2a1a	B2	col2a1a_B2_13	AGGTCCTTGTGGGCCATTCTCTCCTAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_14	CCTCGTAAATCCTCATCAAAGTCTCCCTGTTTGCCAATTGGACCA
col2a1a	B2	col2a1a_B2_15	CCTCCTGGAGCTCCCTGTTTGCCAGAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_16	CCTCGTAAATCCTCATCAAATCACCAGAAGGTCCTGGAAGTCCAG
col2a1a	B2	col2a1a_B2_17	GACCATCCTTACCAGAAGGACCTGCAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_18	CCTCGTAAATCCTCATCAAAGGCCGGCAGCACCAGGGTTACCGTT
col2a1a	B2	col2a1a_B2_19	GCCATCAGCACCAGGGGGTCCAGCAAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_20	CCTCGTAAATCCTCATCAAAGCCAGCAGGTCCAGGTGCACCAATC
col2a1a	B2	col2a1a_B2_21	CCAACAGCTCCTCTCTCACCAGGCAAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_22	CCTCGTAAATCCTCATCAAACCTTGAAGACCGGGAGGACCTTGAG
col2a1a	B2	col2a1a_B2_23	CGGGGGCTCCAGTAGGTCCAGGCAGAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_24	CCTCGTAAATCCTCATCAAACCTGGAAGCCAGATGGACCAGGTTG
col2a1a	B2	col2a1a_B2_25	AGGTGGTCCAGGGCGGCCATCTTCAAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_26	CCTCGTAAATCCTCATCAAAAGGGGCACCAGTCGCTCCAACTTTT
col2a1a	B2	col2a1a_B2_27	GGAGGTCCAAGAGGTCCGGCAGAACAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_28	CCTCGTAAATCCTCATCAAAGGCTCTCCTCTTGGTCCTCTCTTGC
col2a1a	B2	col2a1a_B2_29	CAGATCCTTTAGCTCCAGGAATACCAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_30	CCTCGTAAATCCTCATCAAACAGTACCAGGGGTTACCGGATGCACC

col2a1a	B2	col2a1a_B2_31	TGCACCAGGGCGACCTCTCTCACCGAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_32	CCTCGTAAATCCTCATCAAACAGACCACGAGGACCCATTGGTCCA
col2a1a	B2	col2a1a_B2_33	CTTCCGGGTTTTCCAGGGGGGCCGGAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_34	CCTCGTAAATCCTCATCAAAGGGCCACGGGGTCCAAGGGCACCAG
col2a1a	B2	col2a1a_B2_35	GTCCGGGTGGTCCAGGGGGTCCTGGAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_36	CCTCGTAAATCCTCATCAAATTCCAGGTGTACCAGGCTCGCCATC
col2a1a	B2	col2a1a_B2_37	CTCTCCAGGTGAAATGATGGGATTGAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_38	CCTCGTAAATCCTCATCAAAACAGTCATTCAGATCCTCACAGATG
col2a1a	B2	col2a1a_B2_39	AATCTGAACATGTCTAAAGATTAGAAAATCATCCAGTAAACCGCC
col2a1a	B2	col2a1a_B2_40	CCTCGTAAATCCTCATCAAATGCAGGTCCTAAGGGGTGAAAGTCG

Extended Table 2

Species	Protein	Domain	Amino Acid sequence
Homo sapiese	CCN1	IGFBP	STCPAACHCPLEAPKCAPGVGLVRDGCGCCKVCAKQLNEDCSKTQPCDHTKGLECNFGASSTALKGICRA
Homo sapiens	CCN1	VWC	CEYNSRIYQNGESFQPNCKHQCTCIDGAVGCIPLCPQELSLPNLGCPNPRLVKVTGQCCEEWVC
Homo sapiens	CCN1	TSP-1	-VQTTSWSQCSKTCGTGISTRVTNDNPECRLVKETRICEVRPCG
Homo sapiens	CCN1	CT	KSPEPVRFTYAGCLSVKKYRPKYCGSCVDGRCCTPQLTRTVKMRFRCEDGE--TFSKNVMMIQSCKCNYNCP
Danio rerio	Ccn1	IGFBP	SSCPAVCSCAPDAPQCPPGVGLQRDACGCCLVCARQLNEDCSRSEPCDHSKGLECNYGASHGATRGICR-
Danio rerio	Ccn1	VWC	CEYSSRIYQNGESFQPNCKHQCTCMDGAVGCVPLCPQELSLPALGCANPRLVKVPGRCCEEWVC
Danio rerio	Ccn1	TSP-1	-VQTTGWSQCSRSCGTGISTRITNNNAECKLMKESRICEVRPCS
Danio rerio	Ccn1	CT	KSVTPVKFSYAGCSSLKKYRPRFCGSCVDGRCCSPQNTRTIRVKFRCQDGE--TFNKNVMMIESCKCTKHCP-
Danio rerio	Ccn1l1	IGFBP	GCPLKCSCPSAPPSCPPGISSVLDHCGCCRVCARQFNQDCGSPTEPCDHIKGLRCHLGAEGDPDRGLCRA
Danio rerio	Ccn1l1	VWC	CELDGRIYQHGEDFQPSCEHQCTCVDGVVGCLPLCPHHISLPDWRCARRRLIRKPERCCPEWVC
Danio rerio	Ccn1l1	TSP-1	LLQITDWSPCSATCGMGVSSRVTNSNPECRLVSETRLCQIRECG
Danio rerio	Ccn1l1	CT	RPQKPVQISFAGCSTTRRYRLRSCGFCSDGRLCVPSATRTVRLHFHCPEAERHDFTRSLMWIQRCSCRHTHSR