A Field-Based Study of Phyllosphere Mycobiomes in Apple Orchards Under Varying Agricultural Practices

Sophie Boutin[1,2*], Jonathan Rondeau-Leclaire[1], Alexis Roy[1], Isabelle Laforest-Lapointe[1,2*]

[1]Département de biologie, Université de Sherbrooke, Sherbrooke, QC J1K 2R1, Canada

[2]Centre SÈVE, Université de Sherbrooke, Sherbrooke, QC J1K 2R1, Canada

*Corresponding authors:

Sophie Boutin

e-mail: sophie.boutin3@usherbrooke.ca; sophieboutin13@hotmail.com

ORCID : https://orcid.org/0000-0001-6675-2748

Isabelle Laforest-Lapointe

e-mail: isabelle.laforest-lapointe@usherbrooke.ca; isabelle.laforest.lapointe@gmail.com

ORCID : https://orcid.org/0000-0003-3630-9475

ORCID

Jonathan Rondeau-Leclaire   (0000-0002-4433-3566)

Alexis Roy   (0009-0000-0763-1049)

Key words: mycobiome, apple tree, phyllosphere, agricultural practices, microbial ecology, ITS amplicon sequencing, on-farm study, metagenomics.

Abstract

Microbial communities in the phyllosphere are key players in plant health and disease resistance, yet their response to agricultural practices remains poorly understood under field conditions. Here, we compare fungal community composition and diversity across conventional and organic apple orchards using ITS amplicon sequencing. Leaf samples were collected from six sites at three distinct time points during the 2023 growing season: once in May, July, and August, corresponding approximately to monthly intervals throughout the summer, while flower samples were collected from the same trees in May. Our analyses reveal that agricultural practices are significantly associated with fungal community structure, with effects intensifying from May to July. While conventional practices are associated with increased absolute abundance of commensal and potentially beneficial taxa, higher abundance of several genera which include known pathogens is also observed at organic sites. Alpha diversity declined over time at conventional orchards but remained stable at organic sites. These patterns suggest that shared interventions and distinct inputs impose contrasting selective pressures on the phyllosphere mycobiome, thus affecting not only community composition but also diversity. Our findings underscore the ecological relevance of field-based microbiome research and offer insights to support the development of sustainable orchard management strategies informed by fungal community dynamics.

Introduction

In 2024, the world's total population reached 8.2 billion humans, with an expected increase of 1.5 billion by 2050 [1]. As the world population keeps growing, agriculture faces increasing pressure not only to meet rising food demands, but also to ensure efficient, stable, and sustainable production systems that can support long-term productivity under variable environmental conditions. In parallel, pressures on ecosystems biodiversity and functions have never been so strong, and global change is deeply impacting biomes' temperature and precipitation regimes [2]. Now more than ever, crop producers face the heightened challenges of optimizing land use and limiting the spread of plant bacterial and fungal diseases, while navigating social expectations to minimize local environment disruptions. On the one hand, conventional agricultural practices most often rely on the use of chemicals such as antibiotics, fungicides, and insecticides to maintain crop productivity. On the other hand, these practices can be damageable to the environment in the long-term as they can lead to water pollution [3] and to the emergence of resistances in microbial pathogens [4, 5]. While the use of strong chemicals in agriculture provided crucial benefits to human society in terms of food security during the Green Revolution of the 1950s, the book "Silent Spring" by Rachel Carson [6] revealed how these practices, when used without restrictions, are detrimental to ecosystems [7]. Thus, sustainable agricultural practices (e.g., organic), even if they are less effective in terms of productivity per land area in the short term, are considered to be an important alternative to preserve ecosystem biodiversity and functions [8]. Yet, organic practices also include some treatments (e.g., biocontrol products such as "Blossom Protect" [9], "Bt" [10]), which can impact the microbial communities associated with crops. In this context, research on the effects of agricultural practices (conventional and organic) on plant-microbe interactions is crucial to advance our knowledge of these ecosystems and support crop producers.

Plant flowers, leaves, and roots host diverse microbial communities (i.e., assemblies of microorganisms including bacteria and fungi) that can impact plant host growth and general health [11, 12]. For instance, members of the *Arabidopsis* microbiota can antagonize the widely distributed foliar pathogen *Pseudomonas syringae* [13], and plant growth promoting rhizobacteria (PGPR) are foreseen to be key players in plant growth [14, 15]. As the impact of agricultural practices on rhizosphere (i.e., roots)

microbial communities has been extensively researched [16, 17], the current study focuses

on the aerial parts of plants (i.e., flowers and leaves), commonly named the phyllosphere

[11], an environment that has gained scientific interest in the last two decades due to its

known impacts on general plant health.

Plant-microbe studies are often performed in controlled settings, such as tissue

culture plates filled with agar [18], growth chambers on a variety of substrates such as

calcined clay [19], or greenhouses with controlled lighting, temperature, and humidity [20].

Although these controlled growth experiments provide instrumental understanding of

molecular mechanisms and simplified interactions, they offer little extrapolation potential

to *in situ* dynamics, which is what crop producers truly experiment and deal with.

Scientific initiatives are often criticized to be too far from the real world or to not make the

effort to connect with users [21]. In view of this, initiatives from our group [22, 23] but

also from others (see **Table S1**) have addressed plant-microbe systems directly in the field.

This comes with various challenges linked to access to the sites, access to information on

practices, and study design constraints [24]. Nonetheless, a few studies have investigated

the influence of crop management practices on either phyllosphere bacterial or fungal alpha

and beta diversity (**Table S1**). However, because of the difficulties mentioned above, these

studies often yield different and sometimes contradictory results, thus reinforcing the need

to investigate how agricultural practices impact the structure of microbial phyllosphere

communities in the field to reach scientific consensus.

Fungi, although expected to be less abundant than bacteria [11] in the phyllosphere,

are known to affect host plant and ecosystems through drought tolerance improvement

[25], pathogen regulation [9], and leaf litter decomposition [26]. In a previous study, we

identified *Aureobasidium*, *Sporobolomyces*, and *Mycosphaerella* as the three main fungal

genera observed in the apple tree (*Malus domestica*, Borkh.) phyllosphere [22]. Our work

also showed that site and time are strong drivers of microbial (both bacteria and fungi)

community composition and diversity in field conditions [22, 23]. Yet, in Boutin *et al.*

(2024) [22], when stratifying our data by agricultural practices, the organic site showed no

significant difference in alpha diversity across the three time points, in sharp contrast with

the observable trend at the two conventional sites. Therefore, to broaden the impact of our

research and investigate this discrepancy, we extended our sampling design of the apple
tree phyllosphere in orchards to include more sites of varying agricultural practices.

Here, we aim to compare fungal community composition and diversity across
conventional and organic agricultural practices on both apple tree flowers and leaves.
More precisely, our main objectives are to (1) assess the associations between agricultural
practices and fungal community composition, (2) characterize how alpha diversity varies
across time, sites, and agricultural practices, and (3) identify differentially abundant taxa
within conventional and organic sites. These analyses are performed on a large dataset
including four to seven cultivars at each site, as well as a smaller subset of two cultivars
present at every sites. Building on previous results from our group [22, 23] and others
(**Table S1**), we expect that fungal community assemblies will diverge between agricultural
practices, and that these differences will be reflected in the differential abundance of
specific taxa. Yet, alpha diversity should be affected to a lesser extent. We expect selective
pressure from agricultural chemicals usage to be higher at conventional sites, leading to
stronger alterations in the community composition of dominant fungal taxa, including
apple leaf pathogens such as *Venturia inequalis* (the causal agent of apple scab). Despite
this selective pressure, we anticipate that alpha diversity (Shannon index) will be reduced
at conventional sites. Finally, we expect to observe spatial (site specific) and temporal
variation within fungal community composition and diversity, linked to fungal taxa
responding to climatic cues and treatments, as well as dispersing from local reservoirs
(e.g., soil, land use).

To achieve our objectives, we sampled apple flowers and leaves at six orchards
(three conventional and three organic) at three time points during the summer of 2023
Specifically, flowers were sampled in May, while leaves were sampled in May, July, and
August. Targeting the Internal Transcribed Spacer (ITS) region of fungal ribosomal DNA,
we sequenced fungal DNA extracted from phyllosphere samples to assess community
composition and diversity. Our results highlight a strong association between agricultural
practices and phyllosphere (leaves and flowers) fungal community composition and
diversity, explaining part of the variation that was attributed to site identity in previous
works.

Materials and methods

Sampling

Six orchards (A-B1-B2-C-D1-D2) from the Eastern Townships region of Québec (Canada)

were sampled in 2023. These orchards (sites) were selected based on available cultivars,

agricultural practices (conventional or organic), and spatial proximity. Some orchards were

selected as pairs of contrasting practices with spatial proximity. For example, B1 and B2

are respectively conventional and organic orchards situated less than a kilometer apart.

Sites D1 and D2 are conventional and organic plots within the same orchard. These two

pairs of sites were chosen to test for differences in agricultural practices while controlling

for the distance. More details about the sites and the sampled cultivars are available in

Table S2. At each site, three trees from each cultivar were randomly selected. In May, leaf

and flower samples were collected. In July and August, after the end of the blooming

period, only leaf samples were collected. In the same day of leaf and flower collection,

samples were stored at -20ºC in the laboratory until cell collection and DNA extraction.

Fungal cells collection and DNA extraction

Using the same method as previously described by Boutin *et al.* [22], samples were thawed

at room temperature approximatively one hour before proceeding to cell collection.

Then, 100 ml of 1X Redford Buffer [27] was added to each sampling bag. The bags were

shaken during five minutes, then the buffer was decanted equally into two 50 ml tubes.

These tubes were centrifuged at 4000 g for 20 minutes and the supernatants were discarded.

The pellet of the first tube was resuspended in 800 µl of solution CD1 from the DNeasy®

PowerSoil® Pro Kit (QIAGEN). The content of the first tube was transferred to the second

tube to resuspend its pellet. Up to 1 ml was transferred to a bead beating tube, which was

kept at -20ºC until DNA extraction. Before DNA extraction, bead beating tubes containing

the microbial pellets and CD1 were thawed at room temperature. Tubes were placed in a

Bead Mill Homogenizer (Fisher) set to the following parameters (speed: 3.25 m/s; time:

2 minutes; cycles: 5; breaks: 30 seconds). The rest of the extraction followed the quick start

protocol and DNA was eluted in 75 µl of solution C6. Extracted DNA was stored at -20ºC

until it was sent to the sequencing platform.

Sequencing

The ITS1 region was amplified using the ITS1F (CTTGGTCATTTAGAGGAAGTAA) and ITS2 (GCTGCGTTCTTCATCGATGC) primers. Samples were sequenced at the CERMO genomic platform (Université du Québec à Montréal) on the Illumina MiSeq platform at 2x300 bp.

Amplicon sequence variants processing

A total of 9,333,318 sequences from 364 samples were processed using the DADA2 pipeline v1.34 [28] in R v4.4.0 [29] using default parameters unless otherwise stated. Briefly, low complexity sequences and those containing ambiguous nucleotides were discarded using the *filterAndTrim* function with *rm.lowcomplex = TRUE, maxN = 0*. Primers were removed using cutadapt v2.10 [30] with parameters *-n 2 -m 21 -M 300*, then sequences were quality-filtered using *filterAndTrim*, allowing a maximum expected error *maxEE = c(4, 4)* and a minimum length *minLen = 100*. After learning the error model, sample compositions were inferred using *learnErrors(pool = "pseudo")*. Forward and reverse sequences were merged, then bimeras removed using *method = "consensus"*. The taxonomy of the resulting amplicon sequence variants (ASVs) was assigned using the UNITE ITS database v10.0, 04-04-2024 [31]. ASVs with fewer than a total of 10 sequences across all samples or with no taxonomic classification at the Phylum level were removed. Then, samples with fewer than a total of 2,000 sequences were discarded, resulting in a total of 329 samples kept for analysis (the 35 discarded samples are listed in **Table S3**. This resulted in a total of 5,022,964 total sequences (mean 15,267 ± 8,189 per sample) and 2,812 ASVs (mean 216 ± 144 per sample). On average, each ASV was found in 25 ± 39 samples. Sample metadata, sequence and taxonomy tables were built into a phyloseq object (phyloseq v1.46.0 [32]) for further analyses. For alpha and beta diversity analyses (see below), samples were rarefied to 2,500 sequences to control for uneven sequencing depth using the *rarefy_even_depth* function from the vegan package v2.7-1 [33]. This resulted in a total 822,500 sequences and 2,795 ASVs (mean 146 ± 112 per sample). On average, each ASV was found in 17 ± 31 samples after rarefaction (**Table S4**).

Statistical analyses

Statistical analyses were performed using base R functions and the rstatix v0.7.2 and vegan packages. Plots were generated using the ggplot2 v3.5.2, ggpubr v0.6.1, ggrain v0.0.4, ggh4x v0.3.1, see v0.11.0, and patchwork v1.3.1 packages.

For each dataset, we analyzed alpha diversity based on the Shannon index and beta diversity based on the Bray-Curtis index. Shannon diversity indices were calculated using the *estimate_richness* function from phyloseq. Beta diversity was estimated by first variance-stabilizing the rarefied count tables using the *varianceStabilizingTransformation* function from DESeq2 v1.48.1, then computing the Bray-Curtis dissimilarity index using the *vegdist* function from vegan. Permutational multivariate analyses of variance (PERMANOVA) were applied to the resulting dissimilarity matrices using the *adonis2* function from vegan with 999 permutations. Homogeneity of group dispersions was verified using the *betadisper* function (**Table S5**). The dissimilarity matrices were visualized using principal coordinates analysis (PCoA). Aggregate sample composition bar charts are based on the mean relative abundance of each taxon, showing the top 20 taxa that were most relatively abundant on average.

Flower fungal beta and alpha diversity

To characterize the fungal communities associated with apple tree flowers, we analyzed a subset of 54 flower samples (35 from conventional and 19 from organic orchards), collected across four sites (A, B1, B2, and C). Sites D1 and D2 were excluded due to the absence of flowers at the time of sampling.

Beta diversity dynamics were tested using the following model: [*Community ~ (Practice/Site) × Cultivar*] with 'Site' nested in 'Practice' to reflect the sampling design. A Kruskal-Wallis test revealed that alpha diversity was not normally distributed; therefore, the non-parametric Wilcoxon–Mann–Whitney test was used to confirm whether organic and conventional sample diversity originated from the same distribution.

Leaf fungal beta diversity

To assess the influence of agricultural practices on the fungal communities associated with apple tree leaves, we analyzed all 275 leaf samples, encompassing all sites and cultivars.

Beta diversity dynamics were tested using the following model: [*Community ~ (Practice/Site) × Time + (Practice/Site) × Cultivar*] with 'Site' nested in 'Practice' to reflect the sampling design. Permutations were constrained within individual trees (TreeID) to account for repeated measures. Separate analyses were also conducted for the Honeycrisp (n = 51) and Spartan (n = 53) cultivars to confirm that our results were robust despite uneven cultivar representation across sites. To control for temporal effects, we repeated the PERMANOVA and PCoA independently for each sampling time point (May, July, August), using the model: [*Community ~ (Practice/Site) × Cultivar*], allowing to estimate the variance explained by agricultural practices at each time point. To further highlight site-specific patterns, ordinations were visualized to highlight pairs of sites that were spatially closed to one another: (1) for sites B1 and B2 as well as (2) for sites D1 and D2. To visualize the variation in fungal class composition across the PCoA ordination, we performed an *envfit* analysis from the vegan package to test for correlations between the relative abundance of fungal classes and the ordination axes. Significance was assessed using permutation tests, and a Bonferroni correction was applied to account for multiple comparisons.

Leaf fungal alpha diversity

Diversity was compared using the Kruskal-Wallis non-parametric test, followed by post hoc pairwise Dunn's tests with Bonferroni correction. These analyses were performed to assess variation in diversity across sampling time points (May, July, August) within each site, across time points within each agricultural practice, and across agricultural practices at each time point.

All analyses (for composition and alpha diversity) were also performed separately for the Honeycrisp ($n_{Flowers}$ = 8; n_{Leaves} = 51) and Spartan ($n_{Flowers}$ = 9; n_{Leaves} = 53) cultivars to confirm that our results were robust to uneven cultivar representation across sites.

Differential absolute abundance analyses

Differences in absolute abundance between organic and conventional sites were estimated using the log-linear model implemented in ANCOM-BC2 [34] and are expressed as natural log fold differences (LFD) in absolute abundances. The following procedure was

done separately for flower and leaf samples. First, non-rarefied sequence counts were aggregated at the genus level, discarding any ASV with no taxonomic assignment at that level. Genera found fewer than 10% of samples (6 flower and 28 leaf) were not considered. The model controlled for cultivar, as well as sampling month in the case of the leaf samples. *P*-values were manually corrected using the Holm procedure, because ANCOM-BC2's *p*-value correction only accounts for the number of tests done in a single run, while the command was executed once for leaf and once for flower samples. Genera with an adjusted p-value < 0.01 and having passed the pseudo-count sensitivity check were considered as differentially abundant (enriched) in either group. Only genera with an absolute LFD > 1 were reported.

Results

Flower samples across agricultural practices

We first explored the impact of agricultural practices on flower fungal community composition and diversity. From the principal coordinates analysis (PCoA), we observed that samples from conventional and organic practices clustered separately (**Figure 1A**). The permutational variance analysis (PERMANOVA) revealed that agricultural practices explained up to $R^2 = 20\%$ ($P = 0.001$) of the variation observed within those samples (**Table 1**). Site identity was also a driver of fungal community structure, explaining $R^2 = 13\%$ of the variation (**Table 1**). In terms of alpha diversity, flower samples from organic sites had a lower mean Shannon index than those from conventional sites (conventional = 4.3, organic = 3.7; $P < 0.0001$) (**Figure 1B**). The three main fungal genera found in samples from conventional sites were *Cladosporium*, *Aureobasidium*, and *Fomes* which respectively represented 8%, 8%, and 7% of the total number of sequences. In organic sites, they were *Ramularia, Didymella, Gyoerffyella*, representing 17%, 17%, and 9% of sequences (**Figure 1C**).

We then sought to identify genera enriched in samples from either organic or conventional sites. To that end, we estimated fold differences in absolute abundances for each genus using a log-linear model (ANCOM-BC2), controlling for cultivar type (**Figure 2**). In flower samples, eight and twenty-six genera were found to be differentially enriched (adjusted p < 0.01) in conventional and organic sites, respectively.

Lecanoromycetes (n = 2 genera) and Eurotiomycetes (n = 1 genus) were only enriched in

conventional sites, while Agaricomycetes (n = 11 genera), Phaeococcomyces and

Cystobasidiomycetes (n = 1 genus each) were only enriched in organic sites. The most

differentially enriched genus, *Exidia*, was estimated to be almost 37 times more abundant

(3.6 ± 0.2 LFD) in organic sites compared to conventional sites.

Overall, we obtained similar results when looking at flower samples from

Honeycrisp and Spartan cultivars only. Indeed, samples clustered separately by practice

type (**Figure S1A**). Of note, practices explained an even greater amount of the variation

observed within samples (**Table S6**), and the difference in alpha diversity was not

statistically significant (conventional = 4.4, organic = 3.9; *P* = 0.07; **Figure S1B**). The two

most relatively abundant genera (*Ramularia* and *Aureobasidium*) (**Figure S1C**) were the

same as for the whole dataset.

Leaf samples across agricultural practices

The impact of agricultural practices was then investigated for leaf fungal communities.

In the PCoAs, samples from conventional and organic sites did cluster independently

across all sampling time-points (**Figure 3**). When all leaf samples were analyzed together,

agricultural practices explained 13% of the variation among samples (**Table 2A**). When the

three time-points were evaluated independently, a greater proportion of the variation was

explained by the agricultural practices in July (32%) and August (27%) than in May (13%)

(**Table 2B-D**). Site identity explained from 15 to 31% of the variation across sampling

time-points (**Table 2**). With samples from Honeycrisp and Spartan only, fungal

communities from different agricultural practices also clustered separately (**Figure S2**), as

agricultural practices explained 14% of the variation (**Table S7**).

For leaves, fungal community composition was also explored across sampling time-

points (May, July, and August) for both conventional and organic sites. The most relatively

abundant fungal genera were *Aureobasidium*, *Sporobolomyces*, and *Cladosporium*

(**Figure 4**). Similar community profiles were observed when analyzing the data subset

including only Honeycrisp and Spartan cultivars. (**Figure S3**).

For both practices, our results show that there is a shift in fungal communities' composition between May and July (**Figure 4**), which is confirmed by the change in the absolute abundance of certain taxa (**Figure 5**) as well as by general changes in the relative abundance of specific fungal classes (**Figure S5**). In leaf samples, 10 and 28 genera were found to be differentially enriched in conventional and organic sites, respectively (**Figure 5**). Lecanoromycetes genera (n = 2) identified as significantly enriched were again all associated with conventional sites, and both Tremellomycetes (n = 5) and Cystobasidiomycetes (n = 3) genera were only found to be enriched in organic sites. The most differentially enriched genus was an unnamed Dothidotthiaceae, estimated to be 75 times more abundant (4.32 ± 0.13 LFD) in organic sites compared to conventional.

In terms of alpha diversity, a general decrease in Shannon index through time is observed at conventional sites (**Figure 6A-B**). Oppositely, alpha diversity remains stable from May to August at organic sites (**Figure 6A-B**). When comparing both practices through sampling month, alpha diversity was higher at conventional sites in May compared to organic sites, while the opposite was observed in July and August (**Figure 6C**).

Discussion

Understanding the dynamics of microbial communities in agroecosystems is essential for developing sustainable agricultural practices. While many observational microbiome studies have focused on spatiotemporal variation in microbial assemblages [35, 36], few have examined these dynamics under *in situ* conditions with direct relevance to producers. In this study, we extended the design of our previous works to assess how fungal communities in the apple phyllosphere respond to agricultural practices (conventional vs. organic) across multiple sites (six orchards) and time points (three sampling months). Our results demonstrate that agricultural practices are associated with alterations in the composition and diversity of fungal communities of the apple tree flowers and leaves, with effects intensifying from May to July before stabilizing in August (**Tables 2-3**; **Figures 1-3**). Our work also confirms that both site and month of sampling are strongly correlated with changes in fungal community structure (**Table 1; Table 2A**), reinforcing the idea that orchard-specific conditions, such as microclimate and management practices, contribute to shaping the local phyllosphere mycobiome.

Although practices were broadly categorized as conventional or organic due to limited access to detailed treatment records, this binary classification still revealed clear and consistent differences in fungal assemblages between orchard types. Working *in-field* inevitably introduces constraints, such as incomplete knowledge of specific inputs or treatment schedules, yet it also ensures ecological relevance and direct applicability of findings for growers. Despite these limitations, we observed distinct fungal community profiles between conventional and organic orchards (**Figures 1-4**), suggesting that the use of synthetic versus organic inputs could exert differential selective pressures on the phyllosphere mycobiome, for example resulting in shift in community composition.

Flower mycobiome of organic orchards host more relatively abundant plant pathogens

In flower samples, fungal communities from conventional and organic orchards formed distinct clusters in PCoA ordinations (**Figure 1A**), a pattern supported by PERMANOVA results showing that agricultural practice explains a significant proportion of the variation in fungal community dissimilarities. Notably, genera such as *Alternaria*, *Didymella*, and *Ramularia*, all of which contain potential plant pathogens species (e.g., *Alternaria mali*, *Ramularia collo-cygni*, *Didymella pomorum*) [37–39], were more relatively abundant in organic orchards (**Figure 1C**). Differential abundance analysis confirmed that *Didymella* and *Ramularia* were significantly enriched at organic sites (**Figure 2**). This dominance of a few potentially pathogenic taxa may contribute to the lower alpha diversity observed in organic flower samples (**Figure 1C**), consistent with previous findings that pathogen proliferation can reduce overall community evenness and richness [40, 41]. Interestingly, Agaricomycetes such as the *Exidia* genus (known as saprotroph jelly fungi) were only enriched at organic sites. This suggests either that products used in conventional orchards may reduce the abundance of members of this fungal genus, or that organic cultivation methods favor them.

The effect of agricultural practices on leaf mycobiome beta diversity increases with time
Similar patterns were observed in leaf samples, where agricultural practices were linked to fungal composition and diversity. Importantly, the proportion of variance in fungal community composition explained by practice increased over the growing season, from 13% of community variation in May to 32% in July and 27% in August (**Table 2**).

This temporal trend likely reflects the timing of orchard treatments, which may have been

applied shortly before or after the May sampling, thus exerting stronger effects later in the

season. In conventional orchards, the mid-season shift in community structure was

characterized by an increased relative abundance of *Cladosporium*, *Sporobolomyces*, and

Aureobasidium, genera commonly associated with beneficial or commensal roles in the

phyllosphere [42–44]. Notably, the red pigmented *Sporobolomyces* produce carotenoids,

thus have antimicrobial properties and the potential to control for pathogens [45, 46].

Different species of *Cladosporium* also show biocontrol potential against plant pathogens

[42, 47]. Moreover, *Aureobasidium* is the active ingredient in the biocontrol product

Blossom Protect [9], which is widely applied in both organic and conventional orchards to

suppress fire blight, potentially explaining its consistent presence across all sites

(**Figure 4**). In contrast, organic sites showed a higher abundance of potential pathogens

such as members of the *Dothidotthiaceae*, known to cause cankers and leaf spots [48].

These findings highlight how overlapping and distinct management interventions

across organic and conventional regimes impose unique selection pressures on

phyllosphere fungal communities, leading to both shared dominant taxa, such as

Aureobasidium, and divergent overall community structures. Conventional practices

appear be associated with dominance by potentially beneficial or commensal taxa, while

organic practices showed potentially pathogenic genera to be enriched in leaf mycobiomes.

Yet, it is important to note that these interpretations are based on relative abundance data

derived from metagenomic sequencing, which reflects DNA presence rather than active

fungal biomass or viability. In terms of diversity, alpha diversity declined over time in

conventional orchards but remained stable in organic sites, a pattern consistent with our

previous observations [22]. Interestingly, alpha diversity was higher at conventional sites

in May, while organic sites showed greater diversity in July and August. This contrasts with

patterns observed in bacterial communities, where diversity was lower in organic sites in

May but showed no significant differences between practices in July and August [23].

These divergent trends suggest that agricultural practices may have distinct and taxon-

specific impacts on fungal versus bacterial communities. Taken together, our results

provide *in-field* information about flower and leaf mycobiome dynamics highlighting that

agricultural practices are associated with shifts in fungal community assembly and

diversity across the season. By linking fungal community dynamics to orchard practices,

we provide actionable insights for growers seeking to optimize plant health and disease

resistance through microbiome-informed strategies.

Apple tree cultivar is a weak but significant driver of fungal phyllosphere communities

Consistent with earlier observations by Boutin et al. (2024) [22], cultivar identity explained

a small but statistically significant portion of the variation in leaf-associated fungal

community structure (**Table 2**). This modest effect aligns with previous studies showing

that host species identity tends to exert a stronger influence on microbial communities than

cultivar-level differences [35, 49, 50]. Nonetheless, the reproducibility of this pattern

across independent datasets underscores that, although this effect is weaker than month of

sampling, site identity or agricultural practices, accounting for cultivar identity is relevant

to understand the dynamics of the apple phyllosphere mycobiome.

Limitations

While our study provides ecologically relevant insights into the influence of agricultural

practices on apple phyllosphere fungal communities, several limitations should be

acknowledged. First, working under field conditions meant that we could not fully control

environmental variables or access detailed treatment records for each orchard (e.g., absence

of flowers at sites D1 and D2, not all cultivars present at all sites). As a result, agricultural

practices were broadly categorized as conventional or organic, which may mask important

nuances in management strategies and input types. Second, our sequencing approach

targeted the ITS1 region, which, while widely used for fungal community profiling, has

known biases in taxonomic resolution and amplification efficiency across fungal taxa.

These methodological constraints may have influenced the detection and relative

abundance of certain groups, which should be held into account when interpreting or

comparing the results with other studies. Finally, although our sampling design captured

temporal and spatial variation, it remains observational in nature and cannot establish

causal relationships between specific treatments and microbial outcomes. Future studies

incorporating controlled experiments, metagenomic approaches, and detailed agronomic

metadata will be essential to refine our understanding of how orchard management shapes
fungal community structure and function.

Conclusion

This study provides novel information about how agricultural practices are associated with
fungal community composition and diversity of the apple tree phyllosphere in field
conditions. By leveraging a robust sampling design across multiple sites, cultivars, and
months of sampling, we were able to quantify the relative influence of agricultural
practices, time, and site on fungal assemblages, track alpha diversity dynamics throughout
the growing season, and identify differentially abundant taxa associated with each practice.
Our findings offer new insights into how agricultural practices used for pathogen control
may differentially affect the structure of fungal communities, including shifts in relative
abundance of both pathogenic and beneficial taxa. Beyond documenting these patterns, our
work contributes to a growing body of research emphasizing the ecological relevance of
phyllosphere fungi, which play key roles in plant health, disease resistance, and microbial
interactions. Studying these communities *in situ* allows us to generate actionable
knowledge for growers, highlighting how site-level decisions can influence microbial
composition and, ultimately, crop resilience and productivity.

While our results reveal clear differences between conventional and organic
systems, determining which practice is more sustainable remains an open question. Future
studies are required to develop integrative sustainability indices and link microbial
community structure to disease incidence, yield, and broader environmental and socio-
economic outcomes. We therefore advocate for future research that bridges microbial
ecology with on-farms agronomic performance and sustainability science to guide the
development of microbiome-informed orchard management strategies.

TABLES

Table 1. Factors explaining the variation within all flower samples (PERMANOVA on
Bray-Curtis dissimilarities)

	R^2 (%)	F	Pr(>F)	
practice	20.2	16.9	0.001	***
cultivar	12.4	1.5	0.008	**
practice:site[*1]	12.7	5.3	0.001	***
practice:cultivar	8.9	1.2	0.058	ns
practice:site:cultivar	7.7	1.3	0.050	*
Residual	38.1			
Total explained	61.9			

457   [*1] Site is nested within practice.

458   **Table 2.** Factors explaining the variation within all leaf samples (PERMANOVA on Bray-
459   Curtis dissimilarities). All leaf samples from all time-points (**A**) and separated for May (**B**),
460   July (**C**) and August (**D**).

A

	R^2(%)	F	Pr(>F)	
practice	13.8	117.4	0.001	***
time	19.1	81.1	0.001	***
cultivar	4.4	5.3	0.001	***
practice:site[*1]	15.4	32.7	0.001	***
practice:time	6.0	25.7	0.001	***
practice:cultivar	2.1	2.9	0.001	***
practice:site:time	7.6	8.0	0.001	***
practice:site:cultivar	4.5	2.8	0.001	***
Residual	27.1			
Total explained	72.9			

B

	R^2(%)	F	Pr(>F)	
practice	13.2	20.3	0.001	***
cultivar	10.2	2.2	0.001	***
practice:site[*1]	24.6	9.5	0.001	***
practice:cultivar	5.6	1.4	0.011	*
practice:site:cultivar	12.2	1.4	0.001	***
Residual	34.3			
Total explained	65.7			

C

	R^2(%)	F	Pr(>F)	
practice	32.6	108.4	0.001	***
cultivar	8.3	3.9	0.001	***
practice:site[*1]	26.6	22.1	0.001	***
practice:cultivar	4.3	2.4	0.001	***
practice:site:cultivar	10.4	2.5	0.001	***
Residual	17.8			
Total explained	82.24			

D

	R^2(%)	F	Pr(>F)	
practice	27.5	88.5	0.001	***
cultivar	8.5	3.9	0.001	***
practice:site[*1]	31.1	25.0	0.001	***
practice:cultivar	4.1	2.2	0.003	**
practice:site:cultivar	8.6	2.0	0.001	***
Residual	20.2			
Total explained	79.8			

461
[*1]Site is nested within practice.

Figure 1. Fungal composition and diversity of flower samples. **(A)** Principal Coordinates
Analysis (PCoA) based on Bray-Curtis dissimilarities among flower samples. Point shapes
and colors indicate sampling sites and agricultural practices. Ellipses represent 95%
confidence intervals. **(B)** Shannon alpha diversity index across conventional (n =35) and
organic (n = 19) sites. Asterisks indicate significance levels from Wilcoxon-Mann-Whitney
test (**** P ≤ 0.0001). **(C)** Average relative abundance of the 20 most abundant fungal
genera across flower samples. Taxa not among the top 20 are grouped under "Others." Bar
charts are separated by agricultural practice. *Asterisks denote taxa identified at the family*
level but with unknown genus.

Figure 2: Differential abundance analysis of flower communities comparing differences in absolute abundance of genera across agricultural practice, controlling for cultivar. A negative natural log fold difference (LFD) corresponds to an enrichment in the conventional sites (i.e., the reference group in the log-linear model), whereas a positive LFD corresponds to an enrichment in the organic group. Gen. I. s.: *Genus Incertae sedis*.

Figure 3. PCoAs on Bray-Curtis dissimilarities performed on leaf samples. **(A)** All leaf samples across sites and sampling months. **(B–D)** Subset of samples collected in May **(B)**, July **(C)**, and August **(D)**, with conventional sites shown in orange and organic sites in green. **(E–G)** Highlighted comparison of sites B1 (conventional) and B2 (organic for May **(E)**, July **(F)**, and August **(G)**; other sites are shown in gray. **(H–J)** Highlighted comparison of sites D1 (conventional) and D2 (organic) for May **(H)**, July **(I)**, and August **(J)**; other sites are shown in gray. Ellipses represent 95% confidence intervals.

Figure 4. Fungal community composition across all leaf samples (all cultivars). Average
relative abundance of the 20 most abundant fungal genera. Taxa not among the top 20 are
grouped under "Others." Bar charts are organized by agricultural practice and sampling
month (Conventional: n = 45, 48, 54; Organic: n = 40, 44, 44). Asterisks indicate taxa
identified at the family level but with unknown genus.

 Figure 5: Differential abundance analysis of leaf communities comparing differences in
 absolute abundance of genera across agricultural practice, controlling for cultivar. A
 negative natural log fold difference (LFD) corresponds to an enrichment in the
 conventional sites (i.e., the reference group in the log-linear model), whereas a positive
 LFD corresponds to an enrichment in the organic group. Gen. I. s.: *Genus Incertae sedis*.

Figure 6. Shannon alpha diversity of leaf samples (all cultivars). (A) Alpha diversity across sampling times for each individual site. (B) Alpha diversity across sampling times grouped by agricultural practice. (C) Comparison of alpha diversity between conventional and organic practices at each sampling time point. Asterisks indicate significance levels from post-hoc Dunn tests following a Kruskal-Wallis test ($P \leq 0.05$; $P \leq 0.01$; $P \leq 0.001$; $P \leq 0.0001$).

Acknowledgments

We want to acknowledge the help of the Laforest-Lapointe lab members who participated to field work (E. Lussier, G. Lajeunesse, M. Enea, J. Fraysse, A. Werz, R. Perreault) and revised the manuscript (E. Lussier). We also want to thank orchards managers who allowed us to go sampling on their land.

Authors contribution

Conceptualization: S.B. & I.L-L.
Data curation: S.B., J.R-L. & A.R.
Formal analysis: A.R., S.B, J.R-L & I.L-L.
Funding acquisition: I.L-L., S.B. & J.R-L.
Investigation: S.B., J.R-L. & A.R.
Methodology: S.B., J.R-L. & A.R.
Project administration: I.L-L.
Resources: I.L-L.
Software: S.B., J.R-L. & A.R.
Supervision: I.L-L.
Validation: S.B., J.R-L., A.R. & I.L-L.
Visualization: S.B., J.R-L., A.R. & I.L-L.
Writing – original draft: S.B., J.R-L., A.R. & I.L-L.
Writing – review & editing: I.L-L., S.B., J.R-L., A.R.

Supplementary materials

Supplementary materials are available at the following GitHub repository: https://github.com/alexis-roy5/orchard_phyllosphere

Conflicts of interest

The authors declare no conflict of interest.

Funding

This project was funded by the Canada Research Chairs Program (I.L-L.: CRC-2023-00177), Fonds de Recherche du Québec—Science et Technologies (S.B.: https://doi.org/10.69777/359740 & https://doi.org/10.69777/330306), and National Sciences and Engineering Research Council of Canada (S.B. & J.R-L.).

Data availability

The raw sequences are available on the European Nucleotide Archive under the following accession number: PRJEB96463

References

1. UNITED NATIONS DEPARTMENT FOR ECONOMIC AND SOCIAL AFFAIRS. WORLD POPULATION PROSPECTS 2024: summary of results. S.l.: UNITED NATIONS, 2025.

2. Calvin K et al. IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland. Intergovernmental Panel on Climate Change (IPCC), 2023.

3. Evans AE et al. Agricultural water pollution: key knowledge gaps and research needs. *Curr Opin Environ Sustain* 2019;**36**:20–27. https://doi.org/10.1016/j.cosust.2018.10.003

4. Mann A et al. Antibiotic resistance in agriculture: Perspectives on upcoming strategies to overcome upsurge in resistance. *Curr Res Microb Sci* 2021;**2**:100030. https://doi.org/10.1016/j.crmicr.2021.100030

5. McManus PS. Does a drop in the bucket make a splash? Assessing the impact of antibiotic use on plants. *Curr Opin Microbiol* 2014;**19**:76–82. https://doi.org/10.1016/j.mib.2014.05.013

6. Stein KF. Silent Spring (1962). In: Stein KF (ed.), Rachel Carson: Challenging Authors. Rotterdam: SensePublishers, 1962, 61–106.

7. Halpern BS et al. The environmental footprint of global food production. *Nat Sustain* 2022;**5**:1027–1039. https://doi.org/10.1038/s41893-022-00965-x

8. Reganold JP, Wachter JM. Organic agriculture in the twenty-first century. *Nat Plants* 2016;**2**:15221. https://doi.org/10.1038/nplants.2015.221

9. Celetti M. Blossom Protect: A New Biopesticide for Fire Blight Control. http://omaf.gov.on.ca/english/crops/hort/news/orchnews/2013/on-0113a5.htm. (2013, date last accessed).

10. Whalon ME, Wingerd BA. Bt: Mode of action and use. *Arch Insect Biochem Physiol* 2003;**54**:200–211. https://doi.org/10.1002/arch.10117

11. Vorholt JA. Microbial life in the phyllosphere. *Nat Rev Microbiol* 2012;**10**:828–840. https://doi.org/10.1038/nrmicro2910

12. Berendsen RL, Pieterse CMJ, Bakker PAHM. The rhizosphere microbiome and plant health. *Trends Plant Sci* 2012;**17**:478–486. https://doi.org/10.1016/j.tplants.2012.04.001

13. Vogel CM et al. Protective role of the Arabidopsis leaf microbiota against a bacterial pathogen. *Nat Microbiol* 2021;**6**:1537–1548. https://doi.org/10.1038/s41564-021-00997-7

14. Backer R et al. Plant growth-promoting rhizobacteria: Context, mechanisms of action, and roadmap to commercialization of biostimulants for sustainable agriculture. *Front Plant Sci* 2018;**9**.

15. Shah A et al. PGPR in Agriculture: A Sustainable Approach to Increasing Climate
Change Resilience. *Front Sustain Food Syst* 2021;**5**.

16. Moneda APC et al. Sugarcane cultivation practices modulate rhizosphere microbial
community composition and structure. *Sci Rep* 2022;**12**:19174.
https://doi.org/10.1038/s41598-022-23562-6

17. Katsoula A et al. The response of soil and phyllosphere microbial communities to
repeated application of the fungicide iprodione: accelerated biodegradation or
toxicity? *FEMS Microbiol Ecol* 2020;**96**:fiaa056.
https://doi.org/10.1093/femsec/fiaa056

18. Vogel C et al. Forward Genetic In Planta Screen for Identification of Plant-Protective
Traits of Sphingomonas sp. Strain Fr1 against Pseudomonas syringae DC3000. *Appl*
Environ Microbiol 2012. https://doi.org/10.1128/AEM.00639-12

19. Keppler A et al. Plant microbiota feedbacks through dose-responsive expression of
general non-self response genes. *Nat Plants* 2025;**11**:74–89.
https://doi.org/10.1038/s41477-024-01856-z

20. Dong C-J et al. Bacterial communities in the rhizosphere, phyllosphere and
endosphere of tomato plants. *PLOS ONE* 2019;**14**:e0223847.
https://doi.org/10.1371/journal.pone.0223847

21. Bergmann M et al. Transdisciplinary sustainability research in real-world labs:
success factors and methods for change. *Sustain Sci* 2021;**16**:541–564.
https://doi.org/10.1007/s11625-020-00886-8

22. Boutin S, Lussier E, Laforest-Lapointe I. Investigating the spatiotemporal dynamics
of apple tree phyllosphere bacterial and fungal communities across cultivars in
orchards. *Can J Microbiol* 2024;**70**:238–251. https://doi.org/10.1139/cjm-2023-0215

23. Boutin S, Laforest-Lapointe I. Reproducing plant microbiome research reveals site
and time as key drivers of apple tree phyllosphere bacterial communities. *Sci Rep*
2025;**15**:1–15. https://doi.org/10.1038/s41598-025-10729-0

24. OECD. Challenges for Agricultural Research. OECD, 2011.

25. Cambon MC et al. Drought Tolerance Traits in Neotropical Trees Correlate with the
Composition of Phyllosphere Fungal Communities. *Phytobiomes J* 2023;**7**:244–258.
https://doi.org/10.1094/PBIOMES-04-22-0023-R

26. Phyllosphere fungi on leaf litter of Fagus crenata: occurrence, colonization, and
succession. https://cdnsciencepub.com/doi/epdf/10.1139/b02-028. .

27. Kembel SW et al. Relationships between phyllosphere bacterial communities and
plant functional traits in a neotropical forest. *Proc Natl Acad Sci* 2014;**111**:13715–
13720. https://doi.org/10.1073/pnas.1216057111

28. Callahan BJ et al. DADA2: High-resolution sample inference from Illumina amplicon
data. *Nat Methods* 2016;**13**:581–583. https://doi.org/10.1038/nmeth.3869

29. R Core Team. R: A Language and Environment for Statistical   Computing. 2023.
Vienna, Austria: R Foundation for Statistical Computing, 2023.

30.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing
625   reads. *EMBnet.journal* 2011;**17**:10–12. https://doi.org/10.14806/ej.17.1.200

626   31.   Abarenkov K et al. UNITE general FASTA release for Fungi 2. 2024. UNITE
627   Community, 2024.

628   32.   McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive
629   Analysis and Graphics of Microbiome Census Data. *PLoS ONE* 2013;**8**:e61217.
630   https://doi.org/10.1371/journal.pone.0061217

631   33.   Oksanen J et al. vegan: Community Ecology Package. 2025. 2025.

632   34.   Lin H, Peddada SD. Multigroup analysis of compositions of microbiomes with
633   covariate adjustments and repeated measures. *Nat Methods* 2024;**21**:83–91.
634   https://doi.org/10.1038/s41592-023-02092-7

635   35.   Laforest-Lapointe I, Messier C, Kembel SW. Host species identity, site and time drive
636   temperate tree phyllosphere bacterial community structure. *Microbiome* 2016;**4**:27.
637   https://doi.org/10.1186/s40168-016-0174-1

638   36.   Li M et al. Phyllosphere bacterial and fungal communities vary with host species
639   identity, plant traits and seasonality in a subtropical forest. *Environ Microbiome*
640   2022;**17**:29. https://doi.org/10.1186/s40793-022-00423-3

641   37.   Havis ND et al. Ramularia collo-cygni—An Emerging Pathogen of Barley Crops.
642   *Phytopathology®* 2015;**105**:895–904. https://doi.org/10.1094/PHYTO-11-14-0337-
643   FI

644   38.   Prechsl UE et al. Fungicide-free management of Alternaria leaf blotch and fruit spot
645   on apple indicates Alternaria spp. as secondary colonizer. *Sci Rep* 2023;**13**:8431.
646   https://doi.org/10.1038/s41598-023-35448-2

647   39.   Havenga M et al. Canker and Wood Rot Pathogens Present in Young Apple Trees and
648   Propagation Material in the Western Cape of South Africa. *Plant Dis* 2019;**103**:3129–
649   3141. https://doi.org/10.1094/PDIS-04-19-0867-RE

650   40.   Li P-D et al. The phyllosphere microbiome shifts toward combating melanose
651   pathogen. *Microbiome* 2022;**10**:56. https://doi.org/10.1186/s40168-022-01234-x

652   41.   Kim S-H et al. Comparison of Bacterial Community of Healthy and Erwinia
653   amylovora Infected Apples. *Plant Pathol J* 2021;**37**:396–403.
654   https://doi.org/10.5423/PPJ.NT.04.2021.0062

655   42.   Köhl J et al. Toward an Integrated Use of Biological Control by Cladosporium
656   cladosporioides H39 in Apple Scab (Venturia inaequalis) Management. *Plant Dis*
657   2015;**99**:535–543. https://doi.org/10.1094/PDIS-08-14-0836-RE

658   43.   (PDF) Control of Storage Decay of Apples with Sporobolomyces roseus.
659   https://www.researchgate.net/publication/275070481_Control_of_Storage_Decay_o
660   f_Apples_with_Sporobolomyces_roseus. .

661   44.   Zeng Q et al. Aureobasidium pullulans from the Fire Blight Biocontrol Product,
662   Blossom Protect, Induces Host Resistance in Apple Flowers. *Phytopathology®*
663   2023;**113**:1192–1201. https://doi.org/10.1094/PHYTO-12-22-0452-R

664   45. Fatemi S et al. Sporobolomyces lactucae sp. nov. (Pucciniomycotina,
665   Microbotryomycetes, Sporidiobolales): An Abundant Component of Romaine
666   Lettuce Phylloplanes. *J Fungi* 2022;**8**:302. https://doi.org/10.3390/jof8030302

667   46. Janisiewicz. Control of Storage Decay of Apples with Sporobolomyces roseus.

668   47. Zhu (朱墨) M et al. Biocontrol Potential of Cladosporium sphaerospermum Against
669   the Wheat Powdery Mildew Fungus Blumeria graminis f. sp. tritici. *Plant Dis*
670   2024;**108**:2983–2988. https://doi.org/10.1094/PDIS-02-24-0433-SC

671   48. Senwanna C. Towards a natural classification of Dothidotthia and Thyrostroma in
672   Dothidotthiaceae (Pleosporineae, Pleosporales). *Mycosphere* 2019;**10**:701–738.
673   https://doi.org/10.5943/mycosphere/10/1/15

674   49. Li J et al. Plant identity shapes phyllosphere microbiome structure and abundance of
675   genes involved in nutrient cycling. *Sci Total Environ* 2023;**865**:161245.
676   https://doi.org/10.1016/j.scitotenv.2022.161245

677   50. Wang Y et al. Community Structure of Phyllosphere Bacteria in Different Cultivars
678   of Fingered Citron (Citrus medica 'Fingered') and Their Correlations With Fragrance.
679   *Front Plant Sci* 2022;**13**. https://doi.org/10.3389/fpls.2022.936252