Vision dominates sound in mediating classic cue-induced microsaccadic eye movement modulations in rhesus macaque monkeys

Tatiana Malevich[1, 2], Matthias P. Baumann[1, 2], Yue Yu[1, 2], Ziad M. Hafed[1, 2]

1 Werner Reichardt Centre for Integrative Neuroscience, University of Tübingen, Tübingen, Germany

2 Hertie Institute for Clinical Brain Research, University of Tübingen, Tübingen, Germany

Correspondence:
Ziad M. Hafed
ziad.m.hafed@cin.uni-tuebingen.de

Abstract

Visual onsets in the environment cause a short-latency reduction in saccade generation likelihood. Concomitant with that, the metrics of the saccades that do happen near the time of saccadic inhibition reflect the landscape of stimulus-driven neural responses in topographically organized brain maps. While certain aspects of saccadic inhibition have already been explored from a neurophysiological perspective, several questions remain, including which sensory-motor pathways propagate visual input information to the final oculomotor control circuitry with such short latencies. Here, motivated by evidence of a dissociation between temporal and spatial aspects of saccadic inhibition, we devised behavioral paradigms experimentally manipulating either aspect. Rhesus macaque monkeys maintained gaze fixation, and we characterized visually-driven microsaccadic inhibition with and without paired auditory pulses. In one condition, the sound pulses were spatially-unbiased, thus only affecting the temporal processes associated with microsaccadic inhibition; in another, the pulses were spatially-biased, affecting both the temporal and spatial components. We found that sound pulses alone barely affected microsaccade rates. However, when paired with visual stimuli, they accelerated the timing of visually-driven microsaccadic inhibition by ~14-18 ms. In terms of microsaccade directions, spatially-unbiased sounds interfered with, but did not eliminate, visually-driven direction modulations. Surprisingly, spatially-biased sounds had a much weaker effect on microsaccade direction modulations, even when the sound source was in the opposite hemifield as the appearing visual stimulus. We conclude that vision dominates sound in mediating classic microsaccadic inhibition, and we argue that our paradigms are ideal for neurophysiological investigations of the parallel sensory-motor pathways contributing to such inhibition.

Significance

Coordination of internal oculomotor eye movement plans with exogenous sensory events entails rapid resetting of the saccadic eye movement system, but the pathways mediating the resetting are unknown. We devised novel behavioral paradigms in primates, involving multisensory stimulation, which allow neurophysiological experiments to isolate the contributions of individual sensory, motor, and sensory-motor brain areas to saccadic resetting. Such paradigms underscore that resetting involves dissociable temporal and spatial processes that are dominated by the visual modality.

Introduction

Sensory transients in the environment have a powerful, unavoidable impact on the oculomotor system, and one that precedes even reflexive orienting responses towards such transients (1). This sensory-driven impact manifests itself as a robust, short-latency modulation not only of saccades (2-5) and microsaccades (6-9), but also of slow ocular position drifts during fixation, which are on the scale of approximately 1 min arc eye movements (10, 11), as well as smooth pursuit (12-14).

In saccades and microsaccades, our focus here, sensory transients cause an almost complete cessation of eye movement generation, in a phenomenon called (micro)saccadic inhibition (1-4, 6). While parts of the underlying neural mechanisms for (micro)saccadic inhibition have been explored (15, 16), much remains to be learned (1, 17), especially given the plethora of ways in which sensory information from the periphery can find its way to the final oculomotor control pathways. Prior computational accounts have supported the idea of a dissociation between temporal and spatial factors mediating (micro)saccadic inhibition (18, 19), and this is consistent with behavioral and neurophysiological evidence (16-18, 20-23). However, experimental tests of prior predictions (1, 17, 18) on the relative contributions of cortical and subcortical brain circuits to (micro)saccadic inhibition require behavioral paradigms that can experimentally manipulate and modify the inhibition's temporal and spatial factors. Here, we explored and characterized such paradigms.

In the visual modality, manipulations of the spatial extent of a visual stimulus, as well as the spatial configuration of multiple simultaneously appearing ones, have been particularly informative about the spatial component of (micro)saccadic inhibition (18). These results were also significant because they added to growing evidence (8, 9, 15, 16, 24-28) that (micro)saccadic inhibition can be effectively studied in an ideal animal model for neurophysiological mechanisms, the rhesus macaque monkey. Having said that, there is a need for additional ways to explore the temporal and spatial components of (micro)saccadic inhibition. For example, a visual stimulus alone not only manipulates the spatial aspect (via biasing of spatial maps representing the sensory environment), but the transient visual onset itself simultaneously jumpstarts the temporal processes. Here, we used multisensory stimulation to allow a further experimental differentiation between the spatial and temporal components. We paired a visual stimulus onset with either a spatially-unbiased sound pulse (thus only affecting the temporal processes of the inhibition) or a spatially-biased one (thus affecting both the spatial and temporal processes). Importantly, we performed our experiments in monkeys, as an obligatory prerequisite for us continuing further neurophysiological experiments in the sensory-motor pathways of the very same animals.

Our use of audio-visual stimulation was additionally necessary for delineating the potential brain circuits underlying (micro)saccadic inhibition. For example, one of our ongoing hypotheses (1, 17, 18, 20) is that the temporal processes associated with inhibition can critically depend on omnipause neurons in the brainstem's premotor oculomotor control network (29-32). These neurons gate saccade generation, but past work (primarily in cats) has shown that they also exhibit sensory responses (33-35), which functionally raise the threshold for saccade generation. Importantly, the visual responses of these neurons are

feature-tuned in monkeys (36, 37), in a manner that is directly consistent with the feature
tuning properties of microsaccadic inhibition in these animals (24). Similarly, it has been
suggested that the superior colliculus contributes to the spatial component of visually-driven
(micro)saccadic inhibition (16, 17). Since both the superior colliculus and omnipause
neurons are expected to be multisensory (35, 38-46), paradigms like ours will prove
extremely insightful about the pathways for inhibition. These paradigms also enable
investigating other structures, such as the inferior colliculus (47-50), which could (in the end)
prove to be as critical as omnipause neurons and the superior colliculus.
117
118   In what follows, we describe the relative contributions of sound and vision to microsaccadic
119   inhibition in rhesus macaque monkeys. We demonstrate that our experimental paradigms
120   and analyses are a tremendously useful tool for neurophysiological investigations of multiple
121   sensory-motor pathways mediating a highly robust oculomotor phenomenon.
122
123
124

Methods

Experimental animals and ethical approvals

We collected eye movement data from two adult, male rhesus macaque monkeys (F and A, aged 14 and 12 years, and weighing 13 and 9.5 kg, respectively). The monkeys were prepared for behavioral and neurophysiological experiments previously (e.g. (51, 52)). Briefly, each monkey was implanted with a head-holder for stabilizing head position during data collection, as well as with a scleral search coil in one eye, allowing accurate and precise eye tracking using the magnetic induction technique (53, 54).

All experiments were approved by ethical committees at the regional governmental offices of the city of Tübingen (Regierungspräsidium Tübingen).

Laboratory setup

The monkeys sat in a dark room in front of a CRT display having a gray background of 26.11 cd/m^2 luminance. The display was centered at eye height, and it was 72 cm in front of the animals. The display subtended ~30 deg horizontally and ~23 deg vertically.

For trials with a sound pulse (see experimental procedures below for more details), we used a pair of audio speakers that were placed in two possible configurations. For the spatially-unbiased configuration, the goal was to have a diffuse sound source that would not be spatially biased to either the right or left visual hemifield of the monkey (when it was looking directly at the display center). Thus, each speaker was placed on the same table holding the display, but with one speaker being behind the bottom right corner of the display (behind by 22 cm from the front of the display and 21 cm below the center of the display) and the other speaker being behind the bottom left corner of the display (again, behind by 22 cm from the front and 21 cm below the center). Each speaker was 9 cm wide and 15 cm high, and it was placed upright. The right speaker was aimed towards and to the right of the monkey (that is, diagonally), and the left speaker was again aimed diagonally, but now towards and to the left of the monkey. This way, the sound surrounded the display in a balanced manner and was not spatially biased to either the right or left visual hemifield.

For the spatially-biased sound configuration, we again placed two speakers (each 9.5 cm wide and 24.5 cm high), but this time, they were placed directly next to each other in either the right or left visual hemifield of the monkey. Specifically, the two speakers were laid down on their side on a platform directly under the table holding the display. Thus, the two speakers formed a long, continuous, and horizontal line of sound source (49 cm long in total). This horizontally-lying sound source was aimed diagonally towards and to one side of the monkey, in order to bias the sound waves towards one visual hemifield. For example, if the speakers were placed under the right half of the display (under by 22 cm from the display's bottom), then they were oriented such that they emitted sound diagonally towards and to the right of the monkey. If they were placed under the left half of the display, they were oriented such that they emitted sound diagonally towards and to the left side of the

monkey. Thus, the sound pulse was biased to one visual hemifield when the monkey was fixating the display center, but it could, of course, reach both ears since the speakers were in the environment and not directly placed inside each ear. This was fine for our purposes because it was sufficient to introduce a spatial bias in the sound source (55).

We measured the sound intensity at the position of the monkeys' heads using a sound meter (NL-52A, RION, Tokyo, Japan). For monkey F, the sound intensity at the head was 69-79 db in the spatially-unbiased configuration and 76 db in the spatially-biased configuration. For monkey A, the sound intensity at the head was 89 db in the spatially-unbiased configuration, and it was 76 db in the spatially-biased configuration. The background noise level was 51-54 db. In all cases, the sound pulse that we used was a pure tone (1KHz in both monkeys, except for two sessions in monkey A, in which we used 2KHz) of 50 ms duration. Thus, our sound pulses were clearly audible (suprathreshold), but they were not so loud as to cause startle reflexes by the monkeys (like blinking); in any case, at least in humans, microsaccade rates were shown to not be affected by the emotional arousal level of sound stimuli (56).

Other aspects of the experimental setup (such as visual display control) were the same as those described previously (51, 52, 57, 58). Briefly, we controlled the experiments using a custom-made computer system based on PLDAPS (59) and the Psychophysics Toolbox (60-62). We used Psychophysics Toolbox functions for triggering sound pulses with a temporal precision of a few milliseconds. This temporal precision (on trials in which we paired the sound pulse with a visual stimulus onset) was sufficient for remaining within the expected temporal window in which auditory-visual multisensory integration could happen in the brain (41, 43, 63). We recorded all data (at 1 KHz sampling resolution) using the OmniPlex data acquisition system from Plexon, and our display's refresh rate was 85 Hz.

Experimental procedures

The behavioral task for the monkeys always consisted of maintained gaze fixation on a central, white (79.9 cd/m^2) fixation spot of 10.9 by 10.9 min arc dimensions (on an otherwise gray background).

Spatially-unbiased sound task. In one set of blocks and sessions, we pseudorandomly interleaved trials of three types: control trials with only the onset of a spatially-unbiased sound pulse (20% of all trials within a session), visual-only trials in which only a visual onset occurred without a sound pulse (40% of all trials), and visual plus auditory trials in which the visual onset was paired with a simultaneous spatially-unbiased sound pulse (40% of all trials within a session). In the visual onset trials (whether with or without the spatially-unbiased sound pulse), the stimulus consisted of a small black spot (100% Weber contrast) of 0.36 deg diameter, and it remained on the display for 100 ms. Trials ended 500-800 ms after the offset of the visual stimulus (or 500-800 ms after the onset of the sound pulse on the control trials), and the monkeys were rewarded for maintaining fixation on the central fixation spot until trial end. The visual stimulus was placed at an eccentricity of 0.5-12 deg from the central spot, and in the lower visual field (1-8.5 deg direction from the horizontal). This placement, which was constant within a block of trials, was dictated by where we might

encounter response fields in areas like the primary visual cortex and superior colliculus in subsequent neurophysiological investigations with our paradigm. Moreover, lower visual field stimuli were closer to the sound sources, allowing us to amplify the potential impacts of either spatially biased or unbiased sounds on visually-driven microsaccadic modulations. We collected 171-1489 trials per session in each monkey, and we included data in this study from 10 sessions in monkey F (7180 trials in total) and 40 sessions in monkey A (22403 trials in total).

In the data analyses for this task, we also included visual-only trials from an additional set of blocks and trials from a different behavioral paradigm. In this paradigm, we were varying the visual properties of the stimuli across trials, such as the Weber contrast and luminance polarity, for purposes not directly related to the current study. However, the 100% contrast, black condition in that other paradigm was identical to our visual-only trials here. Thus, we took these particular trials (32%) and collapsed them with the visual-only trials described above. By doing so, we obtained an additional 2370 trials over 10 sessions in monkey F and 4378 trials over 20 sessions in monkey A.

Spatially-biased sound task. We ran separate blocks of trials in which the sound pulses were spatially-biased. In these blocks and in monkey A, we interleaved two trial types within a given session: control trials with only the onset of a spatially-biased sound pulse (~33.3% of all trials within a session), and vision plus sound trials in which the visual onset was paired with a spatially-biased sound pulse (~66.6% of all trials). In monkey F, these types of trials were also interleaved with visual-only trials, resulting in the following trial ratio within a given session: control trials with only the onset of a spatially-biased sound pulse (20% of all trials), vision plus sound trials (40% of all trials), and visual-only trials (40% of all trials). For the later analyses, we collapsed the visual-only trials from this task in monkey F (5648 trials) with the visual-only trials from the spatially-unbiased sound task described above.

We typically ran one block with the sound source on the right, and one block with the sound source on the left on each day (the order of the right- and left-sound blocks was counterbalanced across sessions), and we collected 223-1022 trials per sound location session per monkey. In the analyses, we included data from 11 sessions in monkey F and 18 sessions in monkey A, giving a total of 14126 and 18001 analyzed trials for each monkey, respectively.

In most days, we ran the spatially-unbiased sound blocks described above first, and then two spatially-biased sound blocks (one for a right and one for a left sound source). However, for some sessions, we sometimes only ran the spatially-unbiased sound blocks during each session, and sometimes only the spatially-biased sound blocks. This approach was not a problem for our analyses because microsaccadic inhibition is a stationary process across many trials, blocks, and sessions. For example, it is not altered across thousands of trials in individual monkeys (8) (also see Figs. 1, 5 in Results), as well as in humans (64). Also, in neurophysiological experiments, one is very likely to block the presentation of stimuli like this, and experimental constraints can sometimes force a reordering of which tasks to run first, and so on.

Note that our aim in this study was not to perfectly match the location of the visual stimulus
with the location of the spatially-biased sound source. Rather, we wanted to investigate how
hemifield biases in the sounds influenced visually-driven microsaccadic modulations when
the visual stimulus was in a congruent or incongruent hemifield. Since our visual stimulus
was just a single spot at one location, our goal was achieved with our audio speaker
placements. Specifically, because it is expected that the microsaccades would be directed
towards the visual field stimulus location shortly after its onset (18), having the sound in the
same or opposite hemifield was sufficient to explore the impacts of the sounds on these
modulations (there was no other competing visual stimulus in the hemifield, other than the
experimentally-presented spot). Moreover, auditory receptive fields in relevant multisensory
structures for our task, such as the superior colliculus, are quite large, spanning from 45 to
100 deg in diameter in the periphery (41), and multisensory integration was shown to be
tolerant to small spatial disparities between unisensory stimuli (65). Finally, there are
successful examples in the literature, in which the effects of multisensory integration on
neuronal activity (42) or on saccadic behavior (66) were achieved with stimuli matched by
hemifields.
281
282
283 *Data analysis*
284
285 Data analysis was performed in Matlab 2020b (MathWorks, USA) using custom scripts.
286
287 We detected all microsaccades using our previously established methods (67, 68). We
288 included all trials in which the monkeys were rewarded in our analyses, and we
289 characterized the properties of microsaccadic inhibition. For any condition, we pooled data
290 from all sessions in which the condition was run, again because microsaccadic inhibition in
291 monkeys is a highly stationary process across thousands of trials (8).
292
To compute microsaccade rate, we used a moving window approach. For every trial, we
stepped a window of 50 ms width in steps of 2 ms, covering a range of times around
stimulus (visual, auditory, or both) onset. We then counted the number of microsaccade
onsets that occurred within this time window and divided by 50 ms to get a rate estimate
(movements per second). We then averaged all rate curves from all trials of a given type to
get an average microsaccade rate curve, and we estimated the standard error of the mean
for each time sample across trials.
300
To measure microsaccade direction modulations, we first classified a movement's direction
according to the hemifield towards which it was directed. Because microsaccades in
monkeys orient themselves towards the direction of the appearing stimulus with a relatively
high angular fidelity (8, 18, 69), and because we had only one single visual stimulus location
in any given trial, such hemifield classification was sufficient to reveal microsaccade direction
modulations associated with the visual onset events. A similar situation was true when there
was only a spatially-biased sound pulse in a given trial. In fact, even in the vision plus sound
trials, we had the visual stimulus appear in the lower visual field, consistent with the
location of the sound pulses (thus, meaning that the hemifield direction classification was all
that we needed). After classifying individual microsaccade directions, we used two methods
for assessing direction modulations after stimulus onset.

In the first method, we repeated what we had done previously in monkeys (e.g. (9)).

Specifically, for each time window (of 50 ms width; stepped in steps of 2 ms), we calculated

the fraction of all occurring microsaccades within the time window that were directed

towards the stimulus hemifield (or towards the right or left hemifield in some analyses).

Thus, we measured microsaccade direction biases independent of rate modulation (18). If

the calculated fraction was 0.5, then microsaccades were equally likely to be in one or the

other direction; if it was higher, then microsaccades were more likely to be towards than

opposite the stimulus hemifield.

In the second method, we created a metric called the baseline-subtracted rate difference.

Here, we were motivated by the observation that individual subjects can have intrinsic

idiosyncratic biases in microsaccade directions before any stimulus onset. Thus, for one

hemifield or the other, the "fraction towards" method might show a value deviating from 0.5

even in prestimulus intervals. This can complicate looking for stimulus-induced changes in

such biases. Thus, we performed baseline subtraction. Specifically, for a given stimulus, we

calculated microsaccade rate curves exactly as above, but now separately for only

microsaccades towards or opposite the visual hemifield of a stimulus (or rightward and

leftward microsaccades, independent of stimulus location, in some analyses). Then, for each

such curve, we measured average microsaccade rate in a prestimulus interval (-50 ms to 0

ms from stimulus onset), and we subtracted this prestimulus value from the rate calculations

in each time sample of the rate curve. This centered the rate curve for towards

microsaccades around zero movements per second, and it centered the rate curve for

opposite microsaccades again around zero movements per second. We then took the

difference between the baseline-subtracted towards and opposite rate curves to identify

stimulus driven direction modulations. In this case, the metric starts at around zero before

stimulus onset, and it is positive if stimulus onset caused an increased bias towards the

stimulus location (independent of whether the monkey already had an initial bias in this

direction or in the opposite direction).

We also quantified the timing of microsaccadic inhibition. To do so, we created a metric that

we called L_{B25}, which measured the time at which the microsaccade rate curve after stimulus

onset dropped by 25% relative to the prestimulus rate. This metric can help characterize

whether microsaccadic inhibition is accelerated or delayed by multisensory stimulation. To

calculate it, we took the average microsaccade rate curve of a condition. Then, we measured

the average prestimulus microsaccade rate in an interval from -50 ms to -1 ms from stimulus

onset. Then, we calculated the 75% value of this baseline rate. We marched forward in time

from stimulus onset until microsaccade rate first dropped to this 75% point. This time was

defined as the latency to the 25% modulation in microsaccade rate (L_{B25}). Note that we

picked 25% because it gave us a robust timing value; we were less interested in the absolute

onset time of microsaccadic inhibition than in actually comparing this timing across multi-

and unisensory stimulation conditions. As a result, we picked a level for which we were

certain that microsaccadic inhibition had indeed occurred.

In all conditions, we had thousands of repetitions per monkey (see above and Figs. 1, 5 for

examples). Thus, we were highly confident that our reported means were very

representative of true population means, and we used descriptive statistics (mean and SEM

ranges) throughout the figures of the study. We also used inferential statistics to assess the
effects of our manipulations on microsaccadic inhibition latencies. Specifically, we
performed nonparametric permutation tests on the differences in the obtained L_{B25} metrics.
This approach enabled statistical inference despite having only a single L_{B25} value per
condition. Namely, for each comparison, we pooled trials from the two conditions under
consideration into a single dataset and randomly reassigned their labels while preserving the
original ratio of trial counts across conditions. We repeated this procedure 10000 times. On
each iteration, we computed the average microsaccade rate for each condition, derived the
corresponding L_{B25} value, and calculated the difference in L_{B25} between the two conditions.
The experimental L_{B25} difference was then added to the resulting permuted distribution, and
statistical significance was assessed by computing Monte Carlo p-values. These were defined
as the proportion of permuted differences (under the null hypothesis) that were at least as
extreme as the observed effect. A critical α level of 0.05 was used to determine significance.
Multiple comparisons were corrected, when necessary, using the step-down Bonferroni-
Holm method (70); the corresponding p-values are reported as p_{BH} values. The provided 95%
confidence intervals (*CI*s) were computed as the 2.5th and 97.5th percentiles of the permuted
(i.e. null) distribution (that is, observations outside these intervals were considered
statistically significant).
377
378   To assess the effect of multisensory stimulation on the magnitude of microsaccadic
379   inhibition, we repeated the procedure described above, but this time generating permuted
380   distributions of the differences between the minimum microsaccade rates in the
381   poststimulus interval (from +1 to +150 ms relative to stimulus onset). All other steps of the
382   permutation tests remained the same.
383
384
385
386

Results

Our goal was to document the effects of task-irrelevant sound pulses on visually-driven microsaccadic modulations in rhesus macaque monkeys. This enables using these monkeys as an ideal model system for mechanistically exploring the multiple parallel sensory-motor pathways mediating reflexive oculomotor orienting in the brain.

Throughout all trials in all experiments, our monkeys fixed their gaze on a central fixation spot presented over a uniform gray background (Methods). On every regular trial, a visual stimulus appeared at an eccentricity of 0.5-12 deg to the bottom right or bottom left of the fixation spot (typically at a direction in the range of -1 to -8.5 deg from the horizontal meridian). These locations were chosen because they matched ones that we used in subsequent neurophysiological experiments in the visual cortex and midbrain, using the same animals, for investigating the differential contributions of cortical and subcortical visual pathways to microsaccadic inhibition. On trials with sound pulses, the visual onset events were accompanied by a brief, suprathreshold tone of a single frequency (Methods). And, on some control trials, only the sound pulse was played with no visual stimulus onset event on the display.

When it was played, the source of the sound pulse was either spatially-unbiased, coming from a pair of speakers placed symmetrically behind the display with each speaker at an equal distance to the right or left of the central fixation spot (Methods), or it was spatially-biased, coming from a pair of speakers placed either to the right or left of the central fixation spot (and again slightly behind and below the display; Methods). In all cases, the speakers were placed on the same table holding the display; thus, the speakers were positioned near the lower half of the visual display. Note that we did not use exactly matched sound and visual stimulus locations, as might be sometimes aimed for in neurophysiological studies of audio-visual integration in multisensory structures like the superior and inferior colliculi (41, 43, 48, 65, 71, 72). Rather, we were primarily interested in the short-latency visually-driven microsaccadic modulations, and our specific scientific question was: to what extent could unbiased or biased sounds interfere with or otherwise modify these modulations? Also, since we used a single visual stimulus in either the right or left hemifield, our spatially-biased sounds achieved the goal of causing differential across-hemifield auditory stimulation.

Spatially-unbiased sound alone barely influences microsaccade rate

Our first result was that sound alone had a surprisingly mild, and almost non-existent, impact on microsaccade rate in our monkeys. This can be seen in Fig. 1. When we played the spatially-unbiased sound alone (Fig. 1A, B), there was no apparent short-latency change in the rate of microsaccades occurring after auditory stimulus onset in both monkeys. This was very different from what happened when the visual stimulus appeared alone. In this case, and consistent with the prior literature on microsaccadic inhibition in monkeys (1, 8, 9, 17, 18, 20, 24, 25), there was a robust reduction in microsaccade rate within less than 100 ms from visual stimulus onset (Fig. 1C, D). This robust visually-driven microsaccadic inhibition

was also clearly evident when both the visual stimulus onset and the auditory pulse
occurred together (Fig. 1E, F). Thus, vision dominated the occurrence of stimulus-induced
microsaccadic inhibition in our monkeys. This observation is consistent with previous human
investigations of (micro)saccadic inhibition, in which the inhibition caused by sound alone
was generally weak, absent, or inconsistent across subjects and paradigms (56, 73-76). More
importantly, this observation is scientifically valuable: it makes the use of sound pulses in
monkeys an ideal experimental tool for modifying visually-driven microsaccadic inhibition,
without itself being a major driver for such inhibition. In what follows, we provide further
evidence for this idea, again reinforcing the experimental utility of such spatially-unbiased
sound pulses in invasive pathway studies of sensory-motor transformations in primate brains
(also see Discussion).

Figure 1 Spatially-unbiased sound alone does not cause clear microsaccadic inhibition in rhesus macaque monkeys. (A, B) For each of monkey F (**A**) and monkey A (**B**), we plotted a raster of microsaccade onset times relative to the onset of a brief, spatially-unbiased sound pulse (Methods). The visual display did not contain any transient stimuli during these trials, and the monkeys were instructed to simply fixate the stable, central fixation spot. Each row is an individual trial, and each dot indicates a microsaccade onset time. There were barely any changes in microsaccade likelihood after the occurrence of the sound tone (also see Fig. 2 for more detailed quantification). **(C, D)** In sharp contrast, a visual stimulus onset (Methods) caused robust microsaccadic inhibition (<100 ms after visual stimulus onset) in both monkeys, as expected in these animals (1, 8, 9, 17, 18, 20, 24-26, 28). **(E, F)** Such robust microsaccadic inhibition also occurred when the visual stimulus onset was paired with a simultaneous sound pulse identical to that used in **A**, **B**. Note that the inhibition in this case started slightly earlier than in **C**, **D**, as quantified more clearly in Fig. 2. Also note that we collected more trials in each monkey for all subsequent analyses (Methods), meaning that the shown trials here are from example subsets of individual trials (the shown numbers of trials can be inferred from the vertical scale bars).

Multisensory stimulation with spatially-unbiased sound accelerates visually- or auditory-driven microsaccadic rate inhibition

Closer inspection of Fig. 1E, F should reveal that microsaccadic inhibition started earlier in the case of multisensory stimulation (Fig. 1E, F) than in the case of visual stimulus onsets alone (Fig. 1C, D). To better illustrate this, we converted our measured microsaccade onset

times into microsaccade rate curves (Methods). Such rate curves are plotted in Fig. 2, and
they illustrate multiple key observations. First, with the visual stimulus onset alone, we
observed classic microsaccadic inhibition followed by a rebound in microsaccade rate (Fig.
2A, D; blue curves). Second with the spatially-unbiased sound alone, there was barely any
microsaccadic inhibition in monkey F (Fig. 2A; gray curve), and very mild inhibition in
monkey A (Fig. 2D; gray curve), consistent with Fig. 1. Importantly, multisensory stimulation
accelerated whatever inhibition was observed with unisensory stimulation alone. For
example, in both monkeys, the onset of microsaccadic inhibition with the paired visual and
auditory onsets was earlier than the onset of the mild microsaccadic inhibition with the
auditory event alone (black arrows in Fig. 2B, E). Similarly, microsaccadic inhibition was
earlier on the multisensory trials than in the visual-only trials (Fig. 2C, F). Thus, we observed
evidence of multisensory integration (40, 72, 77, 78), in the sense that the results with
multisensory stimulation did not merely reflect a superposition of the individual unisensory
findings; rather, there was a nonlinear interaction in the timing of microsaccadic inhibition in
the case of multisensory stimulation. Moreover, the multisensory result was more similar to
the visual-only result, as evidenced by the strong inhibition that was absent with the
auditory sound pulses alone. In other words, vision dominated sound in the cue-driven
microsaccadic inhibition effect.

To quantitatively document the acceleration of visually-driven microsaccadic inhibition
timing by spatially-unbiased sound pulses in our monkeys, we calculated a latency measure
of microsaccadic inhibition (L_{B25}). This measure indicates the time at which the microsaccade
rate after stimulus onset dropped by 25% relative to its prestimulus level (Methods). For
monkey F, this latency was 68 ms in the visual-only condition, but it was 50 ms in the
multisensory condition (L_{B25} difference = 18 ms, $p < 0.0001$, 95% CI [-6,8]). Similarly, in
monkey A, L_{B25} was 62 ms in the visual-only condition and 48 ms in the case with a paired
spatially-unbiased sound (L_{B25} difference = 14 ms, $p < 0.0001$, 95% CI [-6,6]). Thus, the
spatially-unbiased sound accelerated the visually-driven microsaccadic inhibition.

Interestingly, even though we could not reliably estimate the L_{B25} parameter for the sound-
only condition of our task (due to the very weak inhibition seen in Figs. 1, 2), visual
inspection suggested that multisensory integration additionally also accelerated the timing
of microsaccadic inhibition relative to the sound-only trials. Specifically, while there was no
apparent latency difference between sound-only trials and visual-only trials for monkey F
(Fig. 2A), the multisensory trials had a shorter latency of inhibition than the sound-only trials
for this monkey (Fig. 2B). In monkey A's case, the sound-only trials exhibited earlier
inhibition than the visual-only trials (Fig. 2D), and the multisensory trials still showed a hint
of a latency benefit compared to the sound-only trials (Fig. 2E), in agreement with monkey
F's results. Thus, the spatially-unbiased sound also accelerated microsaccadic inhibition
relative to sound-only conditions.

530
531
Figure 2 Multisensory stimulation accelerates the timing of unisensory-driven microsaccadic inhibition, with
vision dominating the strength of the inhibition. (A) In monkey F, we estimated microsaccade rate from
individual trials like those shown in Fig. 1 (Methods). Each curve shows average microsaccade rate across all trials
of a given type (blue for visual stimulus alone, and gray for spatially-unbiased sound alone); error bars indicate
across-trial SEM. In the visual modality, classic microsaccadic inhibition, followed by a rate rebound, was
observed. However, with the sound alone, there was only a very mild reduction in microsaccade rate, at
approximately the same time as the visually-driven inhibition. **(B)** When we paired the visual stimulus with a
simultaneous sound pulse (Methods), the microsaccadic rate signature (red) looked very similar to the visually-
driven signature in **A**. However, the inhibition was clearly accelerated relative to the small rate modulation seen
in the case of the sound pulse alone (small black arrow). **(C)** Consistent with this, plotting the rate signatures for
vision in isolation (blue) and multisensory stimulation (red) together revealed a clear acceleration of
microsaccadic inhibition with multisensory stimulation. Thus, multisensory stimulation does not cause a
superposition of effects expected from each sensory modality individually, but it accelerates microsaccadic
inhibition. **(D-F)** Similar results for monkey A. Note how the mild sound-only inhibition in this monkey started
earlier than in monkey F (**D**, **E**). Despite that, multisensory stimulation still accelerated the inhibition relative to
sound alone (**E**; compare red and gray curves) or relative to vision alone (**F**; compare red and blue curves). The
numbers of trials in each condition are indicated in each panel.

549
550
551
552

Finally, and as mentioned above, the strength of microsaccadic inhibition was vastly different
between the sound-only trials and the multisensory trials. In the latter trials, peak inhibition
(i.e. the minimum value reached by the microsaccade rate curve shortly after stimulus onset,
relative to the prestimulus rate value) was similar to the peak inhibition observed in the
visual-only trials, suggesting that the visual modality dominated the stimulus-driven
modulations in microsaccade rate that we observed. We statistically confirmed this
observation. The difference in peak inhibition between auditory-only and visual-only trials
was 1.8831 microsaccades/s in monkey F (p_{BH} = 0.0003, 95% CI [-0.2959,0.1997]) and 0.8016
microsaccades/s in monkey A (p_{BH} = 0.0003, 95% CI [-0.1663,0.1491]). Similarly, between the
auditory-only and multisensory trials, the differences were 2.2345 microsaccades/s (p_{BH} =

0.0003, 95% CI [-0.4798,0.4240]) in monkey F and 0.8821 microsaccades/s (p_{BH} = 0.0003,
95% CI [-0.1668,0.1620]) in monkey A. In contrast, the differences in maximal inhibition
between the visual-only and multisensory trials were tiny and non-significant in both
monkeys: 0.1853 microsaccades/s (p_{BH} = 0.0526, 95% CI [-0.1731,0.1989]) in monkey F, and
0.0804 microsaccades/s (p_{BH} = 0.1725, 95% CI [-0.1146,0.1178]) in monkey A. Importantly,
these differences between the visual-only and multisensory trials were markedly smaller
than those expected from the peak inhibition of the auditory-only trials; in those trials, peak
inhibition relative to prestimulus baselines (i.e. the absolute value between prestimulus and
minimum microsaccade rate) was 0.3571 microsaccades/s in monkey F and 0.7772
microsaccades/s in monkey A. Thus, the multisensory effects were not a superposition of the
individual unisensory ones. This could be because our visual and auditory stimuli were not
spatially aligned (Methods) like in other studies (73), and also because we did not use low-
contrast visual stimuli and/or low-intensity sound pulses, which are known to amplify
multisensory integration effects (72, 73, 79, 80). It is also interesting to note here that we
also saw no clear differences in the post-inhibition microsaccadic rate rebound between
visual-only and multisensory trials (Fig. 2C, F; the first microsaccade rate peak occurring after
the inhibition phase).
580
581
582
583
584 *Spatially-unbiased sound modulates, but does not eliminate, visually-driven*
585 *microsaccade direction modulations*
586
Prior research in humans (19, 81-85) and monkeys (8, 9, 16-18) has revealed the occurrence
of consistent modulations in microsaccade directions after extrafoveal visual stimulus
onsets. If vision really did dominate the cue-driven effects of multisensory stimulation on
microsaccades in our monkeys, then analyzing the monkeys' microsaccade directions in the
experiments of Figs. 1, 2 should still reveal such modulations, despite the spatially-unbiased
sound pulses. This is what we found.
593
With visual stimulation alone, we replicated the previous classic results. For example, for
monkey F, the visually-driven inhibition of microsaccades congruent to the visual stimulus
hemifield (Fig. 3A; congruent) was delayed relative to the inhibition of microsaccades
towards the opposite hemifield (Fig. 3A; incongruent). This means that microsaccades in the
direction of the appearing visual stimulus were more likely to survive microsaccadic
inhibition than microsaccades in the opposite direction; the net result is a biasing of
microsaccade directions towards the appearing visual stimulus shortly after its onset. This
biasing can also be seen for the same monkey in Fig. 3C (blue curve). In this analysis, we
plotted the likelihood for an occurring microsaccade within a given time bin to be in the
direction of the visual stimulus hemifield (Methods). During the short-latency period of
microsaccadic rate inhibition, almost all microsaccades that did occur in this monkey were
congruent with the visual stimulus hemifield (black arrow), and later microsaccades were in
the opposite direction (Fig. 3C; blue curve). Similar effects were also observed in monkey A
(Fig. 3D and Fig. 3F; blue curve), even though this monkey had weaker cueing effects than
monkey F, in general.
609

With the spatially-unbiased sound pulses, the visually-driven microsaccade direction modulations persisted, but they were weakened. For example, in monkey F, L_{B25} for congruent microsaccades with vision alone was 84 ms, and it was 46 ms for incongruent microsaccades (Fig. 3A); giving a time difference of 38 ms ($p < 0.0001$, 95% CI [-4,4]). With the spatially-unbiased sound pulse occurring in synchrony with the visual stimulus onset, the L_{B25} values were 62 ms and 40 ms for the congruent and incongruent microsaccades, respectively; giving a time difference of only 22 ms ($p = 0.0016$, 95% CI [-12,12]) (Fig. 3B). For monkey A, the L_{B25} difference was 10 ms with vision alone (68 ms and 58 ms for congruent and incongruent microsaccades, respectively; $p = 0.046$, 95% CI [-8,8]; Fig. 3D), and it was 8 ms for vision plus the spatially-unbiased sound pulses (52 ms and 44 ms for congruent and incongruent microsaccades, respectively; $p = 0.2651$, 95% CI [-12,12]; Fig. 3E). These effects were also evident in the proportion plots of Fig. 3C, F (red curves). As can be seen, in both monkeys, the cue-driven direction modulations of microsaccades reflected the visual landscape of sensory stimulation, but they were weaker with multisensory stimuli than with visual onsets alone.

Thus, the spatially-unbiased sounds in our experiments interfered with, but did not completely eliminate, the visually-driven microsaccade direction modulations shortly after stimulus onset. This indicates that spatially-unbiased sounds can have multiple divergent effects on reflexive cue-driven oculomotor behavior: on the one hand, they can enhance the salience of the sensory transients (say, via increased alertness) and accelerate the timing of microsaccadic inhibition (Fig. 2); on the other hand, their spatially-diffuse nature can increase the noise in the spatial location representation induced by the visual stimulus onset (putatively in spatially-organized visual-motor maps of the brain, like the superior colliculus; Fig. 3) (40, 41, 71, 72, 86).

The results of Fig. 3D-F for monkey A are particularly convincing that visual cueing effects, despite being initially weak in this monkey, still showed the same trends with the spatially-unbiased sound pulses. However, we further analyzed the data from this and the other monkey, and we could additionally identify a potential reason for this weaker visual-only cueing effect in this particular animal. We believe that this reason can become especially important in neurophysiological experiments involving lateralized brain structures, like the primary visual cortex or the superior colliculus. For example, in causal perturbation neurophysiological experiments, one might experimentally activate or inactivate only one side of the primary visual cortex or superior colliculus, requiring analysis of trials with visual stimuli in only one hemifield (rather than by combining hemifields like we did in Fig. 3). Thus, we asked whether the generally weaker cueing effects for monkey A (with or without the spatially-unbiased sounds) could have been related to potential hemifield asymmetries in this monkey's microsaccades.

Figure 3 Spatially-unbiased sound modulates, but does not abolish, visually-driven microsaccade direction modulations in monkeys. (A) With only visual onsets, the inhibition of microsaccades congruent to the hemifield of the appearing visual stimulus was later than the inhibition for opposite microsaccades (black arrow). This means that near the time of microsaccadic inhibition, the microsaccades that did occur were more likely to be in the direction of the appearing visual stimulus (9, 17, 18). **(B)** When adding a simultaneous spatially-unbiased sound with the visual onset, this effect remained, although it was markedly weakened. **(C)** For the same monkey, we plotted on the x-axis the time from visual stimulus onset, and on the y-axis the proportion of all occurring microsaccades within a given time bin that were directed towards the visual field of the appearing stimulus (Methods). We confirmed the results of **A**, **B**: at the time of microsaccadic inhibition, most microsaccades that did occur were directed towards the appearing visual stimulus, and the spatially-unbiased sound interfered with, but did not completely eliminate, this phenomenon. **(D-F)** Similar observations in monkey A. Note that this monkey had a weaker cueing effect on microsaccade directions (**D**), but this effect was still visible with the interfering sound (**E**, **F**). The generally weaker effect was explained by stronger prestimulus biases in microsaccade directions in this monkey than in monkey F (see Fig. 4). Each curve in **A**, **B**, **D**, **E** is the mean across trials, and the error bars indicate SEM. The total number of trials in each panel was the same as in Fig. 2.

We found that prestimulus microsaccade directions could indeed exhibit idiosyncratic biases in individual monkeys, like in humans (87), and which likely reflect biases in ocular drift directions (10, 11). To demonstrate this, we split trials according to the visual field location of the visual stimulus, and we also split the observed microsaccades within each group of trials according to whether they were rightward or leftward in direction. We found that monkey F tended to make more rightward than leftward microsaccades before stimulus onset (Fig. 4A, B; vertical bidirectional black arrows). In contrast, monkey A had an even stronger bias, but in the opposite direction: there were almost twice as many leftward than rightward microsaccades in this monkey before stimulus onset (Fig. 4D, E; vertical bidirectional black arrows), and the prestimulus rightward microsaccade rate was uncharacteristically low for macaque monkeys. Thus, for any given monkey, a leftward or rightward visual stimulus onset could be in or out of line with the monkey's intrinsic

687 microsaccade direction bias. The next question became, then, whether this could cancel
688 visually-driven cueing effects within specific hemifields or not.
689
The individual-hemifield cueing effects on microsaccade directions remained consistent with
the global results of Fig. 3, despite individual monkey differences in prestimulus
microsaccade direction biases. For example, in monkey F, when the visual stimulus onset was
in the left visual field (Fig. 4A), leftward microsaccades were inhibited later than rightward
microsaccades (L_{B25} of 86 ms versus 46 ms, respectively; L_{B25} difference = 40 ms, $p < 0.0001$,
95% CI [-8,8]), even though leftward microsaccades were less frequent than rightward
microsaccade before stimulus onset in this monkey. Similarly, in the same monkey, when the
visual stimulus onset was in the right visual field (Fig. 4B), now the rightward microsaccades
were inhibited later than the leftward movements (L_{B25} of 82 ms versus 44 ms, respectively;
L_{B25} difference = 38 ms, $p < 0.0001$, 95% CI [-8,6]). In monkey A, with the much bigger
prestimulus bias in microsaccade directions, the results were similar. For the left visual field
stimulus (Fig. 4D), L_{B25} was 62 ms versus 54 ms for leftward and rightward microsaccades,
respectively (L_{B25} difference = 8 ms, $p = 0.1232$, 95% CI [-10,10]); note that the result of the
significance test could reflect the different dynamics of rate inhibition between the two
curves. For the right visual field stimulus (Fig. 4E), the results were 78 ms versus 58 ms for
rightward and leftward microsaccades, respectively (L_{B25} difference = 20 ms, $p < 0.0009$, 95%
CI [-12,12]). Thus, even when analyzing individual hemifields, as might be the case in
neurophysiological experiments in lateralized brain structures like the visual cortex or the
superior colliculus, visually-driven cueing effects will still be observed despite sometimes
strong prestimulus microsaccade biases; congruent microsaccade rate (with the visual
stimulus location) is inhibited later than incongruent microsaccade rate, and for each visual
stimulus hemifield individually (Fig. 4A, B, D, E).
712
The above results, besides supporting the conclusions from Fig. 3, allowed us to next devise
a measure of cue-driven microsaccade direction oscillations that is independent of
prestimulus direction biases. Specifically, we developed a metric called the baseline-
subtracted rate difference (Methods). With this metric, we could measure the rate of
congruent microsaccades and the rate of incongruent microsaccades as a function of time
from stimulus onset. Then, before comparing these two rates to identify cueing effects on
microsaccade directions, we first subtracted the prestimulus microsaccade rate from each
curve individually (Methods). Only after such baseline subtraction did we calculate the
difference in microsaccade rates between congruent and incongruent movements. As seen
in Fig. 4C, F for both monkeys, this results in a metric that is close to zero in the prestimulus
interval and thus isolates the cue-driven modulatory direction effects despite idiosyncratic
prestimulus biases. The results from both monkeys (Fig. 4C, F) allow consistent
interpretations with those from all of our other analyses above. Specifically, for the vision-
only trials (blue curves), there was a biphasic oscillation in microsaccade directions, first
towards and then opposite the recently appearing stimuli. And, with the visual onsets being
paired with spatially-unbiased sound pulses, the direction oscillations persisted but were
weaker. We think that such a metric for measuring cue-driven microsaccade direction
modulations, despite prestimulus microsaccade direction biases, can be a very handy tool,
especially when analyzing individual hemifield data in subsequent neurophysiological
experiments in lateralized sensory-motor brain structures.
733

Figure 4 Persistence of the visual dominance over microsaccade direction modulations, even in the face of potentially strong prestimulus biases in microsaccade directions. (A, B) Similar data to those in Fig. 3A (visual stimulus onset alone), but after separating trials based on the visual field location of the appearing stimulus: left visual field in **A** and right visual field in **B**. In the former case, leftward microsaccades were congruent with the stimulus location (labeled "congruent"), and were inhibited later than rightward microsaccades (**A**); in the latter case (right visual field stimulus), the later inhibition was for the rightward movements (**B**), consistent with Fig. 3. This way of plotting the data was particularly useful for revealing prestimulus microsaccade direction biases (more leftward movements in this monkey). **(C)** A measure of stimulus-induced direction oscillations (like in Fig. 3C), but that can be applied in each hemifield individually despite prestimulus biases. In this measure, we calculated the baseline-subtracted difference in microsaccade rates between different microsaccade directions (here, between congruent and incongruent microsaccades across all visual stimulus locations). Note that we applied this measure for both hemifields combined in this case, for easier comparison to Fig. 3C; however, the measure is equally useful within each hemifield individually. **(D-F)** Similar results for monkey A. There was an even larger prestimulus bias in microsaccade directions (**D**, **E**; approximately twice as many leftward than rightward movements). Despite that, there was still later inhibition for the rate curve of the congruent microsaccades to the visual stimulus (leftward in **D** and rightward in **E**). Consistent with this (and Fig. 3F), the baseline-subtracted rate difference between congruent and incongruent microsaccades (**F**) shifted towards the positive direction during microsaccadic inhibition, even with the spatially-unbiased sound pulses (note that the y-axis range is smaller than that in **C**, reflecting this monkey's weaker cueing effects in general). Error bars denote SEM. All other conventions are similar to earlier figures.

Spatially-biased sound similarly influences microsaccade rate as spatially-
unbiased sound, with and without visual stimulation

Our results so far indicate that spatially-unbiased sound pulses alone had very mild impacts
on stimulus-driven microsaccade rate modulations, but that they still accelerated visually-
driven microsaccadic inhibition. Moreover, such sound pulses interfered with, but did not
completely eliminate, visually-driven microsaccade direction modulations. We interpret
these results as suggesting that vision dominates sound in transient cue-induced
microsaccadic eye movement modulations in rhesus macaque monkeys. Next, we show that
this conclusion holds even when using spatially-biased sound pulses that were either
congruent or incongruent with the hemifield location of the appearing visual stimuli.

In separate blocks of trials, we repeated the same experiments as above, but with the sound
pulse on every trial coming from either the right or left side of the display placed in front of
the animals (Methods). Again, in the absence of a visual onset, such spatially-biased sound
pulses barely affected ongoing microsaccade rates in both monkeys. This can be seen in Fig.
5A, B, in which no microsaccadic inhibition occurred (also see Fig. 7C, F later for a more
quantitative microsaccade rate version of the same data). In contrast, trials with a visual
onset paired with a spatially-biased sound pulse showed clear microsaccadic inhibition,
regardless of whether the sound source was in a hemifield congruent (Fig. 5C, D) or
incongruent (Fig. 5E, F) with the hemifield of the appearing visual stimulus. Thus, we
replicated the same results of Fig. 1, but now with a spatially-biased sound source.

792
793
794

Figure 5 Spatially-biased sound alone again does not cause clear microsaccadic inhibition in monkeys. (A, B) Similar to Fig. 1A, B but this time when the sound source was placed on only one side (either right or left) of the visual display. In the shown trials, the sound pulse occurred alone, and there were no visual transients on the display. There were again barely any changes in microsaccade rate caused by the sound pulse alone (also see Fig. 7). **(C, D)** Microsaccadic inhibition was robust when a visual stimulus onset was paired with the spatially-biased sound pulse. In this case, the visual stimulus appeared in the same hemifield as the spatially-biased sound pulse (Methods). **(E, F)** Microsaccadic inhibition was also robust when the visual onset was in the opposite hemifield from the spatially-biased sound pulse. All other conventions similar to Fig. 1.

We also replicated the observation that multisensory stimulation accelerated the onset of visually-driven microsaccadic inhibition in both monkeys. Consider, for example, Fig. 6A for monkey F and Fig. 6C for monkey A. Here, we plotted microsaccade rates in the visual-only condition in blue (these data are the same as those in Fig. 2 for the same condition). We also plotted microsaccade rates when the visual onset was paired with a spatially-biased sound pulse that was either congruent (magenta) or incongruent (purple) with the hemifield of the appearing visual stimulus. In both cases, microsaccadic inhibition occurred earlier than with a visual stimulus alone. Specifically, for monkey F (Fig. 6A), L_{B25} for the visual-only condition was 68 ms, whereas it was 58 ms (L_{B25} difference = 10 ms, p_{BH} = 0.0356, 95% CI [-6,8]) and 56 ms (L_{B25} difference = 12 ms, p_{BH} = 0.0050, 95% CI [-6,6]) for the congruent and incongruent sound trials, respectively. Similarly, for monkey A (Fig. 6C), L_{B25} for the visual-only condition

819 was 62 ms, whereas it was 50 ms (L_{B25} difference = 12 ms, p_{BH} = 0.0070, 95% CI [-6,6]) and 50
820 ms (L_{B25} difference = 12 ms, p_{BH} = 0.0072, 95% CI [-6,6]) for the congruent and incongruent
821 sound trials, respectively. Thus, sound, whether spatially-unbiased (Fig. 2) or spatially-biased
822 (Fig. 6), accelerated visually-driven microsaccadic inhibition in our macaque monkeys.
823
The acceleration by sound of visually-driven microsaccadic inhibition (Figs. 2, 6) was also
independent of the spatial biases in the sound pulses themselves. Specifically, when we
plotted microsaccade rates from the vision plus spatially-unbiased sound trials (like in Fig. 2)
along with the vision plus spatially-biased sound trials, we found almost perfect overlap in
the timing of microsaccadic inhibition in both monkeys (Fig. 6B, D). Quantitatively, for
monkey F, the difference between L_{B25} in the vision plus spatially-unbiased sound trials and
L_{B25} in the congruent sound trials was -8 ms (p_{BH} = 0.1932, 95% CI [-8,8]), and it was -6 ms
(p_{BH} = 0.3972, 95% CI [-8,8]) between the vision plus spatially-unbiased sound trials and the
incongruent sound trials. Similarly, no differences were observed for monkey A (L_{B25}
difference between unbiased vs congruent sound = -2 ms, p_{BH} = 1, 95% CI [-8,10]; L_{B25}
difference between unbiased vs incongruent sound = -2 ms, p_{BH} = 1, 95% CI [-8,8]).
Moreover, even in the vision plus spatially-biased sound condition itself, the latency of
microsaccadic inhibition did not differ between the trials in which the sound pulse was
congruent or incongruent with the visual stimulus location (for monkey F, L_{B25} difference = 2
ms, p_{BH} = 0.7226, 95% CI [-6,6]; for monkey A, L_{B25} difference = -2 ms, p_{BH} = 1, 95% CI [-8,8]).
Thus, with suprathreshold sounds, it did not matter whether they were spatially biased or
spatially unbiased for their impact on the timing of visually-driven microsaccadic inhibition.
841
In terms of the strength of microsaccadic inhibition, it was again dominated by the visual
condition and not by the sound direction. However, we did notice that peak inhibition was
slightly stronger in the spatially-biased than spatially-unbiased trials (Fig. 6B, D). For monkey
F, the difference between the peak inhibition in the spatially-unbiased trials and the peak
inhibition in the congruent sound trials was 0.2405 microsaccades/s (p_{BH} = 0.0522, 95% CI [-
0.2071,0.2123]); the difference between the peak inhibition in the spatially-unbiased trials
and the peak inhibition in the incongruent sound trials was 0.3603 microsaccades/s (p_{BH} =
0.0012, 95% CI [-0.1944,0.1969]). For monkey A, these values were 0.1687 microsaccades/s
(p_{BH} = 0.0898, 95% CI [-0.1313,0.1320]) and 0.1383 microsaccades/s (p_{BH} = 0.0012, 95% CI [-
0.1328,0.1366]), respectively. Though some p-values did not withstand correction for
multiple comparisons, the trend was consistent across the monkeys. This is unlikely to be
due to the small differences in sound intensity that existed between the experiments
(Methods), because these differences could be of opposite signs across the two monkeys
(Methods) despite both monkeys showing the same effect of stronger inhibition for the
spatially-biased sound configuration.
857
858
859
860

Figure 6 Spatially-biased sound accelerates the timing of visually-driven microsaccadic inhibition similarly to spatially-unbiased sound. (A) In blue, we plotted the microsaccade rate of monkey F on the trials with only a visual onset (same data as in Fig. 2A). In magenta and purple, we plotted microsaccade rate when the visual onset was paired with a spatially-biased sound (congruent with the visual stimulus hemifield in magenta, and incongruent in dark purple). The rate inhibition occurred earlier with multisensory than visual-only stimulation, consistent with the case of a spatially-unbiased sound (Fig. 2). **(B)** The acceleration of the timing of microsaccade rate inhibition by sound was the same whether the sound was spatially-biased or unbiased. The magenta and dark purple curves are the same as those in **A**, and the red curve shows microsaccade rate in monkey F when the visual stimulus onset was paired with a spatially-unbiased sound (same data as in Fig. 2B). Microsaccadic inhibition timing was indistinguishable across all three conditions **(C, D)** The same conclusions were reached for monkey A. Error bars denote SEM, and the numbers of trials for the blue and red curves are the same as those reported earlier in Fig. 2.

Moreover, the rebound in microsaccade rate was also different, but monkey-specific (Fig. 6B, D). As we discuss in more detail below, this latter effect might reflect the individual monkey's rebound in microsaccade rate after sound-only trials, as well as other top-down strategies in recovering from microsaccadic inhibition (9, 15, 17, 27, 88). For example, monkey F had a slight increase in microsaccade rate approximately 100-200 ms after spatially-biased sound onset alone (can be seen in Fig. 7C below), whereas monkey A had a slight decrease at such a time (can be seen in Fig. 7F below). Thus, irrespective of spatial bias in the sound pulses, multisensory stimulation resulted in microsaccade rate inhibition that was largely dominated by the inhibition caused by visual stimulation alone; any small differences in the timing or strength of peak inhibition did not reflect a superposition of unisensory effects, but the sound acted to modulate the properties of the visually-driven outcomes.

Visually-driven microsaccade direction modulations do not change with the hemifield congruency of spatially-biased sound

The strongest evidence that we had for the dominance of vision over sound in cue-driven microsaccadic modulations in rhesus macaque monkeys emerged when we looked at the microsaccade direction modulations shortly after stimulus onset. We repeated the analyses of Fig. 4A, B, D, E, but this time for multisensory stimulation with either a spatially-biased sound pulse in the right hemifield or a spatially-biased sound pulse in the left hemifield. From the earlier analyses (Fig. 4A, B, D, E), we already established that there was a visual cueing effect despite prestimulus biases in microsaccade directions in the individual monkeys (which were particularly strong in monkey A). Now, we found that the visual cueing effect in each hemifield of each monkey (that is, the timing of microsaccadic inhibition for microsaccades congruent and incongruent with visual stimulus location) did not depend on the location of the spatially-biased sound during the critical microsaccadic inhibition periods. This conclusion can be reached from the data shown in Fig. 7A, B for monkey F and those shown in Fig. 7D, E for monkey A. In these data, we plotted the microsaccade rate curves exactly as we did earlier in Fig. 4. When the visual stimulus was in the left hemifield (Fig. 7A, D), microsaccadic inhibition was later for leftward than rightward microsaccades whether the paired sound pulse with stimulus onset was in the right (yellowish curves) or the left (greenish curves) hemifield; in each panel, the corresponding yellow and green curves had almost perfect overlap during the microsaccadic inhibition phase (black oblique arrows). Similarly, when the visual stimulus was on the right (Fig. 7B, E), microsaccadic inhibition was later for rightward than leftward microsaccades (the expected visually-driven cueing effect), and the microsaccadic inhibition timing for each of the microsaccade direction curves was almost identical regardless of whether the sound pulse was in the right or left hemifield (compare the yellowish and greenish curves in each panel; black oblique arrows). Therefore, the transient cue-driven modulation in microsaccade rate (microsaccadic inhibition) and its associated microsaccade direction biasing was clearly dominated by the visual stimulus location, and this happened even despite each monkey's individual prestimulus microsaccade direction biases.

Quantitatively, with a left sound, L_{B25} in monkey F was 44 ms and 76 ms for rightward and leftward microsaccades and a leftward visual stimulus, and it was 48 ms and 70 ms with a right sound (Fig. 7A). For the rightward visual field stimulus location (Fig. 7B), L_{B25} in the same monkey was 72 ms and 34 ms for rightward and leftward microsaccades and a left sound, and it was 76 ms and 32 ms with a right sound. These numbers, when comparing right and left sound groups to each other, were more similar to each other than the actual cueing effects expected from the visual stimulus location alone (Fig. 4), suggesting that the biased sounds barely influenced the visually-driven modulations. Statistically, with the visual stimulus on the left, the effect of the sound being on the left or right was not significant for leftward microsaccades (L_{B25} difference = 6 ms, p_{BH} = 1, 95% CI [-16,16]), nor was it for rightward microsaccades (L_{B25} difference = -4 ms, p_{BH} = 1, 95% CI [-12,12]). Similarly, with the rightward visual stimulus, the L_{B25} difference between the right and left sounds was not significant for either rightward (L_{B25} difference = 4 ms, p_{BH} = 0.9385, 95% CI [-10,10]) or leftward (L_{B25} difference = -2 ms, p_{BH} = 0.9698, 95% CI [-18,18]) microsaccades. That is, the timing difference of L_{B25} was clearly bigger when considering the congruency of

microsaccade directions with the visual stimulus location than the congruency of microsaccade directions with the sound pulse location.

Figure 7 Spatially-biased sound location does not matter for the impacts of visual stimulus location on microsaccade direction modulations during microsaccadic inhibition. (A) Similar to Fig. 4A, but now exploring the effect of the hemifield location of the spatially-biased sound. For a left visual field stimulus, green shows microsaccade rates for rightward or leftward microsaccades and a left sound source; yellow shows the same curves but now with a right sound source. At the time of microsaccadic inhibition, it did not matter whether the sound was on the right or left for each curve. That is, the congruency to the visual stimulus location was what dictated the timing of rate inhibition (black oblique arrows highlight the similarity of inhibition timing between corresponding pairs of yellowish and greenish curves). **(B)** Similarly, when the visual stimulus was in the right hemifield, microsaccadic inhibition reflected the congruency of the microsaccades with the visual stimulus, regardless of whether the sound was congruent (yellow) or incongruent (green) with the visual stimulus location (black oblique arrows). **(C)** With a spatially-biased sound alone, this monkey did not show any appreciable microsaccadic inhibition, regardless of sound or movement direction, consistent with Fig. 5. Critically, the green and yellow curves were overlapping at the time of expected microsaccadic inhibition in **A**, **B**. At later times, rate rebounds could depend on sound location, explaining some of the differentials between the green and yellow curves in the post-inhibition rebound phases in **A**, **B**. **(D-F)** Similar results for monkey A. This monkey was more sensitive to rightward sounds (slightly inhibited leftward microsaccades with the rightward sound alone in **F**). This caused a slightly earlier onset of inhibition in the yellow curves of **D**, **E**, but the overall timing of visually-driven microsaccadic inhibition was unaffected by sound location (oblique black arrows). Error bars denote SEM, and the total numbers of trials for the vision+spatially-biased sound trials were the same as those in Fig. 6.

For monkey A and a left visual field visual stimulus (Fig. 7D), the L_{B25} difference between the leftward and rightward sounds was not significant for either leftward or rightward microsaccades (L_{B25} difference = 4 ms, p_{BH} = 1, 95% CI [-18,18] and L_{B25} difference = -10 ms, p_{BH} = 1, 95% CI [-30,32], respectively). Similarly, for the right visual field visual stimulus (Fig. 7E), the L_{B25} difference between the rightward and leftward sound conditions was not significant for either rightward or leftward microsaccades (L_{B25} difference = -12 ms, p_{BH} = 0.8633, 95% CI [-20,18] and L_{B25} difference = -12 ms, p_{BH} = 0.8633, 95% CI [-20,18], respectively). Thus, even with spatially-biased sounds, vision dominated the early cue-driven modulations of microsaccades in our monkeys.

There were other interesting features of the data in Fig. 7 that are worth documenting here. First, a spatially-biased sound alone did not cause strong microsaccadic inhibition, consistent with all of our prior observations (with spatially biased or unbiased sounds). This can be seen in Fig. 7C, F for each monkey (compare to the strong microsaccadic inhibition in the vision plus sound trials, and also in the vision only trials of Fig. 4). Second, with a rightward sound alone, monkey A showed an early reduction in rightward microsaccades and a slightly smaller increase in leftward microsaccades at approximately 50 ms after sound onset (Fig. 7F; yellowish curves). This appeared as an early modulation in the microsaccade rate curves with a paired visual onset, especially in the right visual field visual stimulus condition (see Fig. 7E; yellowish curves). However, this early modulation effect in the vision plus sound trials was not enough to alter the fact that the visually-driven microsaccadic inhibition remained unaffected by the sound location (black oblique arrows in Fig. 7E illustrate this). Finally, after the microsaccadic inhibition phase (e.g. after more than 100 ms from sound pulse onset), monkey F showed a subtle increase in microsaccade rate on the sound-only trials (Fig. 7C). With a rightward sound, the increase happened for both rightward and leftward microsaccades equally (yellowish curves); with a leftward sound, the increase was stronger for leftward microsaccades (bright green curve). These so-called post-inhibition effects likely reflect top-down control over microsaccades (17), when recovering from the reflexive stimulus-induced inhibition (9, 17, 18, 27, 88), and they can explain why the yellowish and greenish curves in this monkey diverged from each other after the end of microsaccadic inhibition in the vision plus sound conditions of Fig. 7A, B.

In summary, all of our analyses revealed that vision dominated sound in driving the classic cue-driven modulations in microsaccade rate and directions in our rhesus macaque monkeys.

Discussion

We were motivated by the question of how (micro)saccadic inhibition, and related ocular drift phenomena (10, 11), might be mediated in the brain. Clearly, sensory information needs to arrive at the final oculomotor control pathways with very short latencies, since measurable changes in eye movements can occur as early as 50-100 ms after sensory stimulus onset. However, this alone is not sufficient knowledge to pinpoint which sensory-motor pathways are involved in the inhibition and related phenomena, or which sensory-motor pathways are more prominent than others in these phenomena in the healthy brain.

Our starting point in the current study was the dissociation between the temporal and spatial aspects of visually-driven (micro)saccadic inhibition (17, 18, 20). That is, there seems to be a process that implements a phase-resetting of the oculomotor system by visual stimulus onsets (the temporal factor influencing saccade rates), and one in which the spatial readout of maps experiencing visual responses to stimulus onsets modulates the directions of the movements that do occur (the spatial factor influencing saccade directions). We aimed to ask how multisensory stimulation can allow further experimental control over these two types of processes.

In our first set of experiments, we experimentally manipulated the temporal aspect of microsaccadic inhibition by the use of spatially-unbiased sound pulses. Here, we basically injected an additional "phase-resetting" command to visually-driven microsaccadic inhibition, without otherwise differentially influencing the spatial landscape of visual stimulus configurations. We found a result that seems perfectly ideal from a neurophysiological experimentalist's perspective: spatially-unbiased sounds alone did not cause substantial microsaccadic inhibition (Figs. 1, 2), but they still accelerated visually-driven inhibition (Fig. 2). Thus, we succeeded in modifying the temporal component of microsaccadic inhibition. Experimentally, this observation is ideal for us, especially because we can now use spatially-unbiased sounds as a way to only "boost" the temporal aspect of microsaccadic inhibition in the presence of visual stimuli, without the sounds alone themselves being a direct driver for the observed eye movement modulations. When combined with, say, causal perturbation experiments of predominantly sensory (e.g. visual cortex) or predominantly motor (e.g. superior colliculus) brain areas, we can begin to identify, or rule out, which parts of the multiple parallel pathways from the retina to the eye muscles may be most, or least, important for the temporal aspects of the phenomenon of microsaccadic inhibition. For example, if one neurophysiological manipulation were to impair microsaccadic inhibition with only visual stimuli alone but not with multisensory stimulation, then this can inspire investigations of how multisensory structures (even other than the superior colliculus) can contribute to microsaccadic inhibition. In other words, our spatially-unbiased sound results really do widen the space of possible brain targets to explore for getting a better grip over the neural mechanisms of (micro)saccadic inhibition and related oculomotor phenomena.

In our second set of experiments, we experimentally manipulated the spatial component of microsaccadic inhibition. Here, we placed the sound and visual stimulation sources either in spatial congruence or in spatial conflict with each other. Remarkably, we observed that the visual modality was the primary determinant of microsaccade direction modulations. This, in

itself, is a very worthy observation to document, and it would be interesting to ask how this
dominance can be affected in brain perturbation scenarios. For example, along the lines of
the hypothetical perturbation experiments mentioned above for the temporal aspects, if a
causal neurophysiological manipulation were to now severely impair microsaccade direction
modulations with only visual stimuli alone, then does that mean that spatially-biased sounds
can now "kick in" and dominate the spatial aspects of microsaccadic inhibition? This, and
other questions, have the potential to allow substantial progress towards finally
understanding much more of the neural substrates for (micro)saccadic inhibition, despite
the phenomenon itself having been behaviorally documented for over two decades ago
now.

Another intriguing observation that we made in our experiments is that there was a
relatively large difference between how spatially-biased and spatially-unbiased sounds
influenced visually-driven microsaccade direction modulations. With spatially-unbiased
sounds, there was an interference effect with the visually-driven direction modulations,
especially in monkey F (Figs. 3, 4). On the one hand, this might be expected: a spatially-
diffuse sound can introduce noise in the spatial representations encoding the appearing
visual stimulus in the environment (40, 41, 71, 72, 86). On the other hand, with spatially-
biased sounds, there was barely any alteration in the visually-driven direction modulations
with a sound congruent or incongruent with the hemifield of the visual stimulus (Fig. 7). This
is a somewhat surprising observation when compared to the spatially-unbiased sound
situation, but it could also be a perfect opportunity for neurophysiological investigations, as
mentioned above (especially with causal perturbation experiments that might weaken the
visual-only modulations). What we can say, for sure, is that this lack of influence of sound
congruence with the visual stimulus in Fig. 7 was not a failure of the experimental
manipulation itself. Indeed, we did see a clear and consistent impact of spatially-biased
sounds on the temporal aspect of microsaccadic inhibition (Fig. 6), suggesting that the
experimental manipulation itself was present and at least as effective in its impacts on
inhibition timing as the spatially-unbiased case. Moreover, in humans, it was suggested that
spatial cueing by sounds did not cause systematic microsaccadic direction modulations (89).

One possibility for explaining the above observations associated with spatially-biased sounds
is that the auditory effect occurred earlier than the visual effect, resulting in the visual effect
for spatially-biased sounds to be unaffected by spatial congruence with the visual stimulus.
For example, in monkey A, especially for a rightward sound source, there was a slight and
early biasing of microsaccades in the sound-only trials (Fig. 7F; yellowish curves). However,
while this early biasing did cause an early biasing of the visually-driven rate curves in this
monkey (Fig. 7D, E; yellowish curves), it does not really explain why the visually-driven
inhibition timing was unaffected by the congruence of the sound location relative to the
visual stimulus location. For example, the temporal aspects of microsaccadic inhibition (Fig.
6) were clearly influenced by the spatially-biased sounds (and not just in a very early phase
after stimulus onset). Alternatively, a more likely possibility for the lack of change in visually-
driven modulations in Fig. 7 by the spatial congruence of the sound with the visual stimulus
location could be that the oculomotor system is much better equipped to deal with
competing lateralized representations of sensory stimuli (e.g. vision on one side and sound
on another) than with a globally introduced noise injection into the system, like was the case
with our spatially-unbiased sounds. Moreover, we believe that there was still some amount

of interference by the spatially-biased sound trials when compared to the visual-only trials.
For example, the inhibition times of the congruent and incongruent microsaccade rate
curves were more different from each other in Fig. 4A (visual-only trials) than in Fig. 7A
(vision plus spatially-biased sound trials). Thus, there was still likely some interference by the
spatially-biased sound, but the key result remains that microsaccade direction modulations
did not depend on whether the sound was congruent or incongruent with the visual
stimulus location.
1118
Regardless of the true explanation, which we hope to further investigate with
neurophysiological versions of the spatially-biased and spatially-unbiased sound paradigms,
our results revealed that vision really does dominate sound in classic cue-driven
modulations in microsaccadic eye movements in macaque monkeys. This does not mean
that there cannot be any (micro)saccadic inhibition with auditory stimuli. In contrast, there is
plenty of evidence for this in humans. For example, single auditory pulses, whether
unilateral or bilateral, reliably induce microsaccadic inhibition, although its strength and
temporal dynamics tend to differ from those elicited by visual stimuli (7, 56, 73, 74, 85).
Microsaccadic inhibition was also observed in selective attention paradigms, even in
response to distractor stimuli, although it was weaker in that case (90). Furthermore,
microsaccadic inhibition was observed in oddball sound paradigms, specifically in response
to rare, oddball sound transients (88, 91-93). Rather, our results indicate that when a visual
stimulus is paired with sound, the visual modality does dominate the observed effects on
the oculomotor system, and we think that this provides an important foundation, and a
tremendously useful tool, for further neurophysiological investigations exploring the neural
substrates of (micro)saccadic inhibition and related phenomena.

1140

Competing interests statement

1143 The authors declare no competing interests.

1144
1145
1146

Acknowledgements

We were funded by the Deutsche Forschungsgemeinschaft (DFG; German Research
Foundation) through the following projects: SPP 2411 Sensing LOOPS: Cortico-subcortical
Interactions for Adaptive Sensing (project numbers: 520617944 and 520283985, HA
6749/11-1); BU4031/1-1; SPP2205 Evolutionary optimization of neuronal processing (project
numbers: 430158665 and 430157666, HA6749/3-2); SFB 1233 Robust Vision (project
number: 276693517); BO5681/1-1.

1155
1156
1157

Author contributions

1160 TM, MPB, and ZMH designed the experiments. All authors collected the data. TM and ZMH
1161 analyzed the data, and subsequently wrote the manuscript. All authors edited the
1162 manuscript.

1163
1164
1165

References

1.   **Buonocore A, and Hafed ZM**. The inevitability of visual interruption. *J Neurophysiol* 130: 225-237, 2023.

2.   **Buonocore A, and McIntosh RD**. Saccadic inhibition underlies the remote distractor effect. *Exp Brain Res* 191: 117-122, 2008.

3.   **Edelman JA, and Xu KZ**. Inhibition of voluntary saccadic eye movement commands by abrupt visual onsets. *J Neurophysiol* 101: 1222-1234, 2009.

4.   **Reingold EM, and Stampe DM**. Saccadic inhibition in voluntary and reflexive saccades. *J Cogn Neurosci* 14: 371-388, 2002.

5.   **Bompas A, and Sumner P**. Saccadic inhibition reveals the timing of automatic and voluntary signals in the human brain. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 31: 12501-12512, 2011.

6.   **Engbert R**. Microsaccades: A microcosm for research on oculomotor control, attention, and visual perception. *Prog Brain Res* 154: 177-192, 2006.

7.   **Rolfs M, Kliegl R, and Engbert R**. Toward a model of microsaccade generation: the case of microsaccadic inhibition. *J Vis* 8: 5 1-23, 2008.

8.   **Hafed ZM, Lovejoy LP, and Krauzlis RJ**. Modulation of microsaccades in monkey during a covert visual attention task. *Journal of Neuroscience* 31: 15219-15230, 2011.

9.   **Malevich T, Buonocore A, and Hafed ZM**. Dependence of the stimulus-driven microsaccade rate signature in rhesus macaque monkeys on visual stimulus size and polarity. *J Neurophysiol* 125: 282-295, 2021.

10.   **Malevich T, Buonocore A, and Hafed ZM**. Rapid stimulus-driven modulation of slow ocular position drifts. *eLife* 9: 2020.

11.   **Khademi F, Zhang T, Baumann MP, Malevich T, Yu Y, and Hafed ZM**. Visual Feature Tuning Properties of Short-Latency Stimulus-Driven Ocular Position Drift Responses during Gaze Fixation. *J Neurosci* 44: 2024.

12.   **Buonocore A, Skinner J, and Hafed ZM**. Eye Position Error Influence over "Open-Loop" Smooth Pursuit Initiation. *J Neurosci* 39: 2709-2721, 2019.

13.   **Kreyenmeier P, Bhuiyan I, Gian M, Chow HM, and Spering M**. Smooth pursuit inhibition reveals audiovisual enhancement of fast movement control. *J Vis* 24: 3, 2024.

14.   **Kerzel D, Born S, and Souto D**. Inhibition of steady-state smooth pursuit and catch-up saccades by abrupt visual and auditory onsets. *J Neurophysiol* 104: 2573-2585, 2010.

15.   **Peel TR, Hafed ZM, Dash S, Lomber SG, and Corneil BD**. A Causal Role for the Cortical Frontal Eye Fields in Microsaccade Deployment. *PLoS Biol* 14: e1002531, 2016.

16.   **Hafed ZM, Lovejoy LP, and Krauzlis RJ**. Superior colliculus inactivation alters the relationship between covert visual attention and microsaccades. *Eur J Neurosci* 37: 1169-1181, 2013.

17.   **Hafed ZM, Yoshida M, Tian X, Buonocore A, and Malevich T**. Dissociable Cortical and Subcortical Mechanisms for Mediating the Influences of Visual Cues on Microsaccadic Eye Movements. *Front Neural Circuits* 15: 638429, 2021.

18.   **Hafed ZM, and Ignashchenkova A**. On the dissociation between microsaccade rate and direction after peripheral cues: microsaccadic inhibition revisited. *J Neurosci* 33: 16220-16235, 2013.

19. **Tian X, Yoshida M, and Hafed ZM**. A Microsaccadic Account of Attentional Capture
and Inhibition of Return in Posner Cueing. *Frontiers in systems neuroscience* 10: 23, 2016.
20. **Buonocore A, Chen CY, Tian X, Idrees S, Munch TA, and Hafed ZM**. Alteration of the
microsaccadic velocity-amplitude main sequence relationship after visual transients:
implications for models of saccade control. *J Neurophysiol* 117: 1894-1910, 2017.
21. **Buonocore A, McIntosh RD, and Melcher D**. Beyond the point of no return: effects of
visual distractors on saccade amplitude and velocity. *J Neurophysiol* 115: 752-762, 2016.
22. **Buonocore A, Tian X, Khademi F, and Hafed ZM**. Instantaneous movement-unrelated
midbrain activity modifies ongoing eye movements. *eLife* 10: 2021.
23. **Tian X, Yoshida M, and Hafed ZM**. Dynamics of fixational eye position and
microsaccades during spatial cueing: the case of express microsaccades. *J Neurophysiol* 119:
1962-1980, 2018.
24. **Khademi F, Zhang T, Baumann MP, Buonocore A, Malevich T, Yu Y, and Hafed ZM**.
Visual feature tuning properties of stimulus-driven saccadic inhibition in macaque monkeys.
J Neurophysiol 130: 1282-1302, 2023.
25. **Xue L, Huang D, Wang T, Hu Q, Chai X, Li L, and Chen Y**. Dynamic modulation of the
perceptual load on microsaccades during a selective spatial attention task. *Scientific reports*
7: 16496, 2017.
26. **Orczyk JJ, Barczak A, O'Connell MN, and Kajikawa Y**. Saccadic inhibition during free
viewing in macaque monkeys. *J Neurophysiol* 129: 356-367, 2023.
27. **Cui J, Wilke M, Logothetis NK, Leopold DA, and Liang H**. Visibility states modulate
microsaccade rate and direction. *Vision Res* 49: 228-236, 2009.
28. **Yu G, Herman JP, Katz LN, and Krauzlis RJ**. Microsaccades as a marker not a cause for
attention-related modulation. *eLife* 11: 2022.
29. **Buttner-Ennever JA, Cohen B, Pause M, and Fries W**. Raphe nucleus of the pons
containing omnipause neurons of the oculomotor system in the monkey, and its homologue
in man. *J Comp Neurol* 267: 307-321, 1988.
30. **Keller EL**. Participation of medial pontine reticular formation in eye movement
generation in monkey. *J Neurophysiol* 37: 316-332, 1974.
31. **Luschei ES, and Fuchs AF**. Activity of brain stem neurons during eye movements of
alert monkeys. *J Neurophysiol* 35: 445-461, 1972.
32. **Cohen B, and Henn V**. Unit activity in the pontine reticular formation associated with
eye movements. *Brain Res* 46: 403-410, 1972.
33. **Busettini C, and Mays LE**. Pontine omnipause activity during conjugate and
disconjugate eye movements in macaques. *J Neurophysiol* 90: 3838-3853, 2003.
34. **Missal M, and Keller EL**. Common inhibitory mechanism for saccades and smooth-
pursuit eye movements. *J Neurophysiol* 88: 1880-1892, 2002.
35. **Evinger C, Kaneko CR, and Fuchs AF**. Activity of omnipause neurons in alert cats
during saccadic eye movements and visual stimuli. *J Neurophysiol* 47: 827-844, 1982.
36. **Buonocore A, and Hafed ZM**. A sensory race between oculomotor control areas for
coordinating motor timing (Abstract). *Journal of Vision* 21: 2420, 2021.
37. **Buonocore A, Baumann MP, and Hafed ZM**. Visual pattern analysis by motor
neurons (Abstract). *Computational and Systems Neuroscience (Cosyne) 2020 Conference* 147,
2020.
38. **Middlebrooks JC, and Knudsen EI**. A neural code for auditory space in the cat's
superior colliculus. *J Neurosci* 4: 2621-2634, 1984.

39.   **Stein BE, Magalhaes-Castro B, and Kruger L**. Superior colliculus: visuotopic-
somatotopic overlap. *Science* 189: 224-226, 1975.
40.   **Meredith MA, and Stein BE**. Visual, auditory, and somatosensory convergence on
cells in superior colliculus results in multisensory integration. *J Neurophysiol* 56: 640-662,
1986.
41.   **Wallace MT, Wilkinson LK, and Stein BE**. Representation and integration of multiple
sensory inputs in primate superior colliculus. *J Neurophysiol* 76: 1246-1266, 1996.
42.   **Bell AH, Corneil BD, Meredith MA, and Munoz DP**. The influence of stimulus
properties on multisensory processing in the awake primate superior colliculus. *Can J Exp*
Psychol 55: 123-132, 2001.
43.   **Frens MA, and Van Opstal AJ**. Visual-auditory interactions modulate saccade-related
activity in monkey superior colliculus. *Brain Res Bull* 46: 211-224, 1998.
44.   **Jay MF, and Sparks DL**. Sensorimotor integration in the primate superior colliculus. I.
Motor convergence. *J Neurophysiol* 57: 22-34, 1987.
45.   **Jay MF, and Sparks DL**. Sensorimotor integration in the primate superior colliculus. II.
Coordinates of auditory signals. *J Neurophysiol* 57: 35-55, 1987.
46.   **Sparks DL, and Hartwich-Young R**. The deep layers of the superior colliculus. *Rev*
Oculomot Res 3: 213-255, 1989.
47.   **Bulkin DA, and Groh JM**. Distribution of visual and saccade related information in the
monkey inferior colliculus. *Front Neural Circuits* 6: 61, 2012.
48.   **Schmehl MN, Herche JL, and Groh JM**. Visually evoked activity and variable
modulation of auditory responses in the macaque inferior colliculus. *J Neurophysiol* 133:
1456-1467, 2025.
49.   **Gruters KG, and Groh JM**. Sounds and beyond: multisensory and other non-auditory
signals in the inferior colliculus. *Front Neural Circuits* 6: 96, 2012.
50.   **Porter KK, Metzger RR, and Groh JM**. Visual- and saccade-related signals in the
primate inferior colliculus. *Proc Natl Acad Sci U S A* 104: 17855-17860, 2007.
51.   **Malevich T, Zhang T, Baumann MP, Bogadhi AR, and Hafed ZM**. Faster Detection of
"Darks" than "Brights" by Monkey Superior Colliculus Neurons. *J Neurosci* 42: 9356-9371,
2022.
52.   **Trottenberg C, Yu Y, Zhang T, Baumann MP, Malevich T, Prasad S, and Hafed ZM**.
Much higher covariation with foveation timing by superior colliculus than primary visual
cortical neuronal activity. *BioRxiv* 2025.
53.   **Judge SJ, Richmond BJ, and Chu FC**. Implantation of magnetic search coils for
measurement of eye position: an improved method. *Vision Res* 20: 535-538, 1980.
54.   **Fuchs AF, and Robinson DA**. A method for measuring horizontal and vertical eye
movement chronically in the monkey. *J Appl Physiol* 21: 1068-1070, 1966.
55.   **Takahashi TT, Keller CH, Nelson BS, Spitzer MW, Bala AD, and Whitchurch EA**.
Object localization in cluttered acoustical environments. *Biol Cybern* 98: 579-586, 2008.
56.   **Chen JT, Yep R, Hsu YF, Cherng YG, and Wang CA**. Investigating Arousal, Saccade
Preparation, and Global Luminance Effects on Microsaccade Behavior. *Front Hum Neurosci*
15: 602835, 2021.
57.   **Baumann MP, Bogadhi AR, Denninger AF, and Hafed ZM**. Sensory tuning in neuronal
movement commands. *Proc Natl Acad Sci U S A* 120: e2305759120, 2023.
58.   **Bogadhi AR, and Hafed ZM**. Express detection of visual objects by primate superior
colliculus neurons. *Scientific reports* 13: 21730, 2023.

59.   **Eastman KM, and Huk AC**. PLDAPS: A Hardware Architecture and Software Toolbox
for Neurophysiology Requiring Complex Visual Stimuli and Online Behavioral Control. *Front*
Neuroinform 6: 1, 2012.
60.   **Brainard DH**. The Psychophysics Toolbox. *Spatial vision* 10: 433-436, 1997.
61.   **Pelli DG**. The VideoToolbox software for visual psychophysics: transforming numbers
into movies. *Spatial vision* 10: 437-442, 1997.
62.   **Kleiner M, Brainard D, and Pelli DG**. What's new in Psychtoolbox-3? (Abstract).
Perception 36: 2007.
63.   **Meredith MA, Nemitz JW, and Stein BE**. Determinants of multisensory integration in
superior colliculus neurons. I. Temporal factors. *J Neurosci* 7: 3215-3229, 1987.
64.   **Bonetti F, Valsecchi M, and Turatto M**. Microsaccades inhibition triggered by a
repetitive visual distractor is not subject to habituation: Implications for the programming of
reflexive saccades. *Cortex* 131: 251-264, 2020.
65.   **Kadunce DC, Vaughan JW, Wallace MT, and Stein BE**. The influence of visual and
auditory receptive field organization on multisensory integration in the superior colliculus.
Exp Brain Res 139: 303-310, 2001.
66.   **Corneil BD, and Munoz DP**. The influence of auditory and visual distractors on
human orienting gaze shifts. *J Neurosci* 16: 8193-8207, 1996.
67.   **Chen CY, and Hafed ZM**. Postmicrosaccadic enhancement of slow eye movements.
The Journal of neuroscience : the official journal of the Society for Neuroscience 33: 5375-
5386, 2013.
68.   **Bellet ME, Bellet J, Nienborg H, Hafed ZM, and Berens P**. Human-level saccade
detection performance using deep neural networks. *J Neurophysiol* 121: 646-661, 2019.
69.   **Chen CY, Ignashchenkova A, Thier P, and Hafed ZM**. Neuronal Response Gain
Enhancement prior to Microsaccades. *Curr Biol* 25: 2065-2074, 2015.
70.   **Holm S**. A Simple Sequentially Rejective Multiple Test Procedure. *Scandinavian*
Journal of Statistics 6: 65-70, 1979.
71.   **Meredith MA, and Stein BE**. Spatial factors determine the activity of multisensory
neurons in cat superior colliculus. *Brain Res* 365: 350-354, 1986.
72.   **Stanford TR, Quessy S, and Stein BE**. Evaluating the operations underlying
multisensory integration in the cat superior colliculus. *J Neurosci* 25: 6499-6508, 2005.
73.   **Wang CA, Blohm G, Huang J, Boehnke SE, and Munoz DP**. Multisensory integration
in orienting behavior: Pupil size, microsaccades, and saccades. *Biological psychology* 129: 36-
44, 2017.
74.   **Zhao S, Yum NW, Benjamin L, Benhamou E, Yoneya M, Furukawa S, Dick F, Slaney
M, and Chait M**. Rapid Ocular Responses Are Modulated by Bottom-up-Driven Auditory
Salience. *J Neurosci* 39: 7703-7714, 2019.
75.   **Pannasch S, Dornhoefer SM, Unema PJ, and Velichkovsky BM**. The omnipresent
prolongation of visual fixations: saccades are inhibited by changes in situation and in
subject's activity. *Vision Res* 41: 3345-3351, 2001.
76.   **Graupner ST, Velichkovsky BM, Pannasch S, and Marx J**. Surprise, surprise: two
distinct components in the visually evoked distractor effect. *Psychophysiology* 44: 251-261,
2007.
77.   **Meredith MA, and Stein BE**. Interactions among converging sensory inputs in the
superior colliculus. *Science* 221: 389-391, 1983.
78.   **Rowland BA, Quessy S, Stanford TR, and Stein BE**. Multisensory integration shortens
physiological response latencies. *J Neurosci* 27: 5879-5884, 2007.

79. **Populin LC, and Yin TC**. Bimodal interactions in the superior colliculus of the behaving cat. *J Neurosci* 22: 2826-2834, 2002.

80. **Fetsch CR, DeAngelis GC, and Angelaki DE**. Bridging the gap between theories of sensory cue integration and the physiology of multisensory neurons. *Nat Rev Neurosci* 14: 429-442, 2013.

81. **Hafed ZM, and Clark JJ**. Microsaccades as an overt measure of covert attention shifts. *Vision Res* 42: 2533-2545, 2002.

82. **Engbert R, and Kliegl R**. Microsaccades uncover the orientation of covert attention. *Vision Res* 43: 1035-1045, 2003.

83. **Laubrock J, Engbert R, and Kliegl R**. Microsaccade dynamics during covert attention. *Vision Res* 45: 721-730, 2005.

84. **Hafed ZM**. Alteration of visual perception prior to microsaccades. *Neuron* 77: 775-786, 2013.

85. **Rolfs M, Engbert R, and Kliegl R**. Crossmodal coupling of oculomotor control and spatial attention in vision and audition. *Exp Brain Res* 166: 427-439, 2005.

86. **Rowland B, Stanford T, and Stein B**. A Bayesian model unifies multisensory spatial localization with the physiological properties of the superior colliculus. *Exp Brain Res* 180: 153-161, 2007.

87. **Nachmias J**. Two-dimensional motion of the retinal image during monocular fixation. *J Opt Soc Am* 49: 901-908, 1959.

88. **Zhao S, Contadini-Wright C, and Chait M**. Cross-Modal Interactions Between Auditory Attention and Oculomotor Control. *J Neurosci* 44: 2024.

89. **Laubrock J, Kliegl R, Rolfs M, and Engbert R**. When do microsaccades follow spatial attention? *Atten Percept Psychophys* 72: 683-694, 2010.

90. **Liu X, and Chait M**. Dissociable Pupil and Oculomotor Markers of Attention Allocation and Distractor Suppression during listening. *BioRxiv* 2025.

91. **Valsecchi M, and Turatto M**. Microsaccadic responses in a bimodal oddball task. *Psychol Res* 73: 23-33, 2009.

92. **Kadosh O, and Bonneh YS**. Involuntary oculomotor inhibition markers of saliency and deviance in response to auditory sequences. *J Vis* 22: 8, 2022.

93. **Widmann A, Engbert R, and Schroger E**. Microsaccadic responses indicate fast categorization of sounds: a novel approach to study auditory cognition. *J Neurosci* 34: 11152-11158, 2014.