Lifestyle-associated variation in type IV secretion systems between phytopathogenic and environmental *Ralstonia*

Author names

Tabitha C. Cowell[1] (0009-0002-6055-6978)

Matthew L. Cope-Arguello[1] (0000-0002-2860-8179)

Gi Yoon Shin[1] (0000-0001-8168-5870)

Adam J. Bogdanove[2] (0000-0003-1683-4117)

Sara C. D. Carpenter[2] (0000-0002-3485-4042)

Kamrun Nahar[3] (0009-0007-8785-8243)

Apurba Saha[3]

Tiffany M. Lowe-Power[1] (0000-0003-2681-3563)

Affiliations

[1]Department of Plant Pathology, University of California, Davis, Davis, CA, USA 95616

[2]Plant Pathology and Plant-Microbe Biology Section, School of Integrative Plant Science, Cornell University, Ithaca, NY, USA 14850

[3]Biotechnology Division, Bangladesh Agricultural Research Institute, Gazipur 1701, Bangladesh

Corresponding author and email address

Tiffany M. Lowe-Power, tlowepower@ucdavis.edu

Keywords

Ralstonia, type IV secretion system, conjugation, accessory plasmids, integrative and conjugative elements

Abstract

Type IV secretion systems (T4SSs) are versatile machines with variable functions including DNA uptake and release, protein translocation, and DNA conjugation. However, the diversity, distribution, and functional roles of the T4SS in the *Ralstonia* genus remain poorly understood. The *Ralstonia solanacearum* species complex (RSSC) comprises three species of plant-pathogenic bacteria that cause bacterial wilt disease. The *Ralstonia* genus also includes non-RSSC species that are primarily environmental bacteria and rare opportunistic human pathogens. This study compared the diversity and phylogenetic distribution of T4SSs in the RSSC phytopathogens vs. non-RSSC environmentals. Phylogenetic analysis of VirB4 sequences and synteny analysis revealed 16 distinct T4SS clusters in *Ralstonia*, with ten clusters found in RSSC phytopathogen genomes, twelve in non-RSSC environmental genomes, and six clusters in both groups. Collectively, these gene clusters were more prevalent in non-RSSC environmental genomes. The presence of type IV coupling protein and relaxase genes suggests that at least 14 of these T4SS gene clusters could be putative DNA conjugation systems. The clusters were encoded on accessory plasmids of various sizes or as integrative and conjugative elements (ICEs) on the chromosome or megaplasmid. The putative regions of transfer for T4SS gene clusters in the RSSC phytopathogen genomes often contained type III effectors, type VI secretion toxin/antitoxin clusters, and hemagglutinin gene clusters. In contrast, the non-RSSC environmentals were enriched in heavy metal metabolism and resistance genes. One of the 16 T4SS clusters, cluster i, exhibited evidence of specialization for the RSSC phytopathogens. These findings shed light on the eco-evolutionary differences in the *Ralstonia* genus.

Impact statement

The *Ralstonia* genus contains the *Ralstonia solanacearum* species complex (RSSC), a group of globally important plant pathogens that impact food security. These pathogens are known to have an expansive, open pangenome with high levels of gene flow. To shed light on the eco-evolutionary differences between RSSC phytopathogens and closely related environmental species, we explored the potential role of type IV secretion systems (T4SSs) in horizontal gene flow within the *Ralstonia* genus. Surprisingly, these mobile genetic elements were less common in the phytopathogens than the environmentals. Nevertheless, we identified a particular T4SS gene cluster encoded on an accessory plasmid that appears to be specialized to the pathogenic lifestyle of the RSSC. This specialized cluster harbors genes that drive RSSC host range (type III effectors) and others that likely allow these pathogens to antagonize the plant microbiota (type VI secretion toxins and hemagglutinin two partner secretion systems). Moreover, this study sheds light on the cryptic lifestyles of the poorly studied environmental *Ralstonia* species, revealing that their conjugative T4SSs frequently carry heavy metal metabolism and resistance genes. Overall, this study provides a foundation to investigate the functional roles of T4SS clusters and their cargo genes in the fitness of phytopathogenic and environmental *Ralstonia*.

Data summary

Supplemental Table S1 lists details of the RSSC phytopathogen, non-RSSC environmental, and Burkholderiaceae family genomes used in this study. Supplemental Table S2A is the final list of 503 VirB4 sequences collectively identified in 644 RSSC phytopathogen genomes and 143 non-RSSC environmental genomes. Supplemental Table S2B includes the annotations and NCBI accessions for every gene and protein sequence in reference clusters a-p. Supplemental Table S2C is the list of putative T4SS cargo genes identified in complete *Ralstonia* genomes. Supplemental Table S2D is the list of putative T4SS cargo genes identified in cluster i regions from draft and complete *Ralstonia* genomes.

Supplemental File S1 is the multiple sequence alignment (MSA) input for HMMER, made from the 753 VirB4 protein sequences from the Burkholderiaceae family genomes. Supplemental Files S2 and S3 are the species trees displaying the presence of T4SS gene clusters in RSSC phytopathogens and non-RSSC environmentals, respectively. Supplemental File S4 is the species tree displaying the presence of RSp0179 and RSp1521 in the *Ralstonia* genus. Supplemental File S5 is the MSA input for FastTree 2, made from the VirB4 protein sequences from *Ralstonia* genomes. Supplemental Files S6-S9 are various formats of the VirB4 protein tree in Figure 1. Supplemental Files S10-S25 are the GBK files for reference clusters a-p. Supplemental File S26 is the clinker output comparing the reference clusters. Supplemental Files S27-S36 are the clinker outputs comparing the putative regions of transfer of T4SSs in complete *Ralstonia* genomes. Supplemental File S37 is the clustered ANI matrix for the putative regions of transfer of cluster i T4SSs in draft and complete *Ralstonia* genomes. Supplemental File S38 is the clinker output comparing the cluster i regions in draft and complete *Ralstonia* genomes. All Supplemental Tables and Files are available through Zenodo (DOI: 10.5281/zenodo.14873519).

This study's newly sequenced genomes, Gazipur 4 and Gazipur 5, are available on NCBI with the following accessions: GCF_049860735.1 and GCF_049860725.1, respectively. The two contigs containing the cluster j T4SS were submitted to NCBI through the Third Party Annotation (TPA) section of the DDBJ/ENA/GenBank databases with the following accessions: BK072068-BK072069. These contigs were assembled from SRA reads under the accession: SRR18649448.

Introduction

Each environment presents specific opportunities and challenges for bacteria. For example, soil-dwelling bacteria must survive unpredictable nutrient and terminal electron acceptor availability, variations in moisture, and intermicrobial competition. Host-colonizing bacteria must survive many of those challenges as well as host immune factors, such as bursts of reactive oxygen species and antimicrobial molecules. To seize the opportunities and overcome the challenges of their lifestyles, bacteria adapt through evolution where selection acts on pre-existing genetic variation derived from *de novo* mutations or horizontal gene transfer (HGT). Cases have been identified in which certain strains of an environmental bacterial species become pathogenic due to the presence of horizontally acquired DNA (1–5). Horizontally transferred DNA can include conjugative plasmids, integrative and conjugative elements (ICEs), non-conjugative plasmids, transposons, and phages (6,7).

By definition, conjugative plasmids and ICEs are self-transmissible because they encode conjugative molecular machines called type IV secretion systems (T4SSs) (8–10). Additionally, these mobile genetic elements often carry non-T4SS genes that confer new functions that may enable niche adaptation. A common marker of T4SSs is the VirB4 protein sequence because VirB4 family proteins are highly conserved ATPases that are associated with all known T4SSs (10–13). This family includes VirB4, TrbE, CagE, TraC, TraU, IcmB, DotO, and Tfc16. Determining that a given T4SS gene cluster encodes a conjugation machine requires experimental validation because in addition to DNA conjugation, T4SSs may carry out DNA uptake, environmental release of DNA, or protein translocation (8,10,14). Nevertheless, the presence of genes encoding the type IV coupling protein (T4CP) and relaxase can provide circumstantial evidence that the T4SS is conjugative. The T4CP recruits the DNA and/or protein substrate(s) to the membrane-bound T4SS complex (8,15). A T4CP is required for DNA conjugation (15,16) and DNA release systems (10), and is usually, but not always, associated with protein translocation systems (10,13). Additionally, a relaxase is required for DNA conjugation (10,13,15,17) and DNA release systems (10,13), although the relaxase is not considered a true T4SS component. Therefore, the presence of a T4CP gene in a T4SS gene cluster and a relaxase gene in the vicinity is an indicator that the T4SS in question may function as a DNA conjugation or DNA release system. However, the absence of an identifiable T4CP or relaxase gene cannot be used to rule out these potential functions.

A model genus to explore the question of niche adaptation of pathogenic versus environmental lifestyles is *Ralstonia*. The *Ralstonia* genus is composed of two major clades: (i) the aggressive plant pathogens in the *Ralstonia solanacearum* species complex (RSSC) and (ii) the environmental bacteria isolated from diverse habitats. Hereafter, we call these groups the RSSC phytopathogens and the non-RSSC environmentals, respectively. The RSSC phytopathogens are the primary causative agent of bacterial wilt disease and include three species: *Ralstonia solanacearum*, *R. pseudosolanacearum*, and *R. syzygii* (18). Most RSSC phytopathogens are environmentally transmitted through soil and surface water between infection cycles where they colonize the rhizosphere, xylem, and apoplastic spaces (19). The non-RSSC environmentals include at least seventeen species: *R. chuxiongensis*, *R. condita*, *R. edaphi*, *R. flaminis*, *R. holmesii*, *R. insidiosa*, *R. mannitolilytica*, *R. mojiangensis*, *R. pickettii*, *R. psammae*, *R. thomasii*, *R. wenshanensis*, and five unnamed *Ralstonia* spp. Less is known about the ecological niches of non-RSSC environmentals, but they have been isolated from more diverse natural and human-made habitats: plant rhizosphere, soil, surface water, built water infrastructure, filter-sterilized pharmaceuticals, and opportunistic infections of physiologically and immunocompromised patients. Although the environmentals can be opportunistic pathogens, these are rare outcomes, largely resulting from colonization of indoor plumbing and contamination of pharmaceuticals (20).

T4SSs are widespread and can be found in many bacterial and archaeal taxa (8,10,12,14), and are often studied for their contribution to behavior and evolution of pathogens (1,4,21). Few studies have compared T4SS biology between well-established pathogens and their environmental relatives. Here, we elucidate the diversity and phylogenetic distribution of T4SSs within RSSC phytopathogens and non-RSSC environmentals. We used VirB4 protein sequences to identify T4SS gene clusters in the *Ralstonia* pangenome. We characterized the gene clusters by synteny and sequence diversity, compared their phylogenetic distributions within the genus, and inferred their adaptive potential.

Methods

Genomes and accessions

Supplemental Table S1 lists details of the RSSC phytopathogen, non-RSSC environmental,
and Burkholderiaceae family genomes used in this study. The quality of all genomes was
assessed using CheckM (22) in KBase (23) and only genomes with less than 1%
contamination were included. Bioinformatic analysis used sets of genomes: 394 RSSC
phytopathogen genomes available as NCBI assemblies in 2023, 644 RSSC phytopathogen
genomes available as NCBI assemblies or as raw reads on NCBI SRA in 2024, 143 non-
RSSC environmental genomes from 17 genomospecies available as NCBI assemblies in
2023, and 923 Burkholderiaceae family genomes that include the 394 RSSC phytopathogen
genomes and 529 genomes from at least 252 genomospecies. Most genomes were
downloaded from the NCBI assembly database, but 145 RSSC phytopathogen genomes
mostly from Greenrod *et al.* (2022) (24) were assembled from raw reads acquired from NCBI
SRA. Genome assembly was performed on KBase; reads were trimmed with Trimmomatic
(Illumina) (25) or Filtlong (Nanopore) (26), assembled with SPAdes (Illumina) (27) or
Unicycler (hybrid Nanopore and Illumina) (28), and annotated with Prokka (29). These
assemblies can be accessed on a public KBase narrative
(https://narrative.kbase.us/narrative/189849). The two contigs containing the cluster j T4SS
were submitted to NCBI through the Third Party Annotation (TPA) section of the
DDBJ/ENA/GenBank databases with the following accessions: BK072068-BK072069. These
contigs were assembled from SRA reads under the accession: SRR18649448.

We isolated two *Ralstonia pseudosolanacearum* strains from *Solanum melongena* stems in
Gazipur, Bangladesh. Each strain was grown on SMSA agar plates (1 g/L casamino acids,
10 g/L peptone, 5 mL/L glycerol, 15 g/L agar, 50 mg/L 2,3,5-triphenyltetrazolium chloride, 5
mg/L crystal violet, 100 mg/L polymyxin B sulphate, 25 mg/L bacitracin A, 5 mg/L
chloramphenicol, and 0.5 mg/L penicillin G) (30) and single colonies were inoculated into 6
mL of CPG broth (1 g/L casamino acids, 10 g/L peptone, 5 g/L glucose, and 1 g/L yeast
extract) (31). Liquid cultures were grown overnight in a shaking incubator at 250 rpm and
28°C. We extracted gDNA using 2 mL of culture and the MasterPure Gram Positive DNA
Purification Kit (Biosearch Technologies, Cat. No. MGP04100) using the manufacturer's
protocol with doubled reagent volumes. The genomes of these strains were sequenced by
Plasmidsaurus using Oxford Nanopore Technology and assembled with Flye v2.9.1 (32).
The genome sequences for Gazipur 4 and Gazipur 5 are available on NCBI with the
following accessions: GCF_049860735.1 and GCF_049860725.1, respectively.

Synteny analysis

We used clinker (33) to visualize synteny of gene neighborhoods with a threshold of 30%
global amino acid identity.

Identification of VirB4 protein sequences using a custom hidden Markov model

Initial BLASTp (34) searches for VirB4 sequences in RSSC phytopathogen genomes revealed there were at least six VirB4 variants, so we used a hidden Markov model (HMM) approach in the KBase platform to robustly identify divergent VirB4 protein sequences. Briefly, we BLASTp searched the six VirB4 proteins against 923 Burkholderiaceae family genomes, used the hits to build a preliminary HMM, searched the preliminary HMM against the 923 Burkholderiaceae family genomes, built a final HMM from the hits, and searched the final HMM against 644 RSSC phytopathogen and 143 non-RSSC environmental genomes. We queried six divergent VirB4 protein sequences against the 923 Burkholderiaceae family genomes using BLASTp with 20% identity and 20% alignment overlap thresholds (WP_011002496.1, WP_003274595.1, WP_042568051.1, WP_247391909.1, WP_230646236.1, and WP_230645232.1). The 617 combined hits were aligned in a multiple sequence alignment (MSA) using MUSCLE (35) with a maximum of 16 iterations. With HMMER (36), this MSA was queried against the 923 Burkholderiaceae family genomes with thresholds of 20% model coverage and a bit score of 20, yielding 753 VirB4 sequences. Using the previously described parameters, the 753 VirB4 protein sequences were aligned using MUSCLE and the resulting MSA (Supplemental File S1) was queried against 644 RSSC phytopathogen and 143 non-RSSC environmental genomes using HMMER. The final list of 498 VirB4 sequences identified in those genomes is included in Supplemental Table S2A. We used clinker to analyze the gene neighborhoods of a subset of the identified *virB4* genes to verify that they were near other T4SS genes.

Typically, pseudogenes are missed in protein sequence-based searches. So, in addition to the 498 VirB4 sequences identified by the HMM search, we manually identified five T4SS gene clusters with *virB4* genes that were pseudogenized. These are included in the list in Supplemental Table S2A.

Construction of species trees and protein trees

All trees were constructed on the KBase platform. Species trees were built using the KBase "Insert Set of Genomes Into SpeciesTree" pipeline, which uses FastTree 2 (37) to create an approximately-maximum-likelihood tree based on 49 conserved genes (Supplemental Files S2-S4). Protein trees were built by constructing MSAs with MUSCLE with a maximum of 16 iterations before building approximately-maximum-likelihood protein trees with FastTree 2 using default settings in KBase (Supplemental Files S5-S9). Newick files were visualized in iTOL (38).

Gene annotation of T4SS gene clusters

To annotate T4SS genes in the T4SS reference gene clusters, we used three annotation sources: NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (39), NCBI Conserved Domain Database (CDD) (40), and MacSyFinder v2 with CONJscan models (41,42). When annotations conflicted, priority was given to MacSyFinder/CONJscan, then NCBI RefSeq/GenBank, then NCBI Conserved Domain Database. Operon-mapper (43) was used to predict operons in genomes used to build the reference T4SS gene clusters. Any predicted operons that contained an annotated T4SS gene were included in the reference

cluster. NCBI accessions for genomes, genes, and proteins for every gene in reference
clusters a-p can be found in Supplemental Table S2B and Supplemental Files S10-S26.

To broadly classify the T4SS gene clusters, we used the nomenclature established by the
F/P/I/GI-type classification system (10,11). However, the CONJscan models are based on
the different mating pair formation (MPF) classification system and include MPF_F, MPF_T,
MPF_I, MPF_G, MPF_B, MPF_C, MPF_{FA}, and MPF_{FATA} models (13,41,44). Between these two
classification systems, MPF_F corresponds to F-type, MPF_T to P-type, MPF_I to I-type, and
MPF_G to GI-type. The remaining MPF classes include T4SSs that are not represented by the
F/P/I/GI-type classes.

Determining the phylogenetic distribution of T4SS gene clusters using the VirB4 protein sequences

To determine the phylogenetic distribution of the T4SS gene clusters in *Ralstonia* (clusters
a-p as described in the Results), we used BLASTp with 40% sequence identity and 50%
alignment overlap thresholds to search 644 RSSC phytopathogen genomes and 143 non-
RSSC environmental genomes for each of the 16 identified VirB4 protein sequences, one
from each of the reference gene clusters. The NCBI accessions for the VirB4 protein or gene
sequences of each reference cluster and the genome it came from are as follows:
WP_011002496.1, GMI1000 (a); WP_247391909.1, T25_UW811 (b); WP_214372649.1,
SY1 (c); WP_123203755.1, Tg03 (d); WP_230645232.1, LMG10661 (e); WP_003274595.1,
MolK2 (f); WP_165858801.1, SL1931 (g); WP_230646236.1, LMG10661 (h);
WP_042568051.1, UW163 (i); YO100T327_00165, YO100_T327 (j); WP_103518556.1,
ICMP-8657 (k); WP_009242089.1, WCHRI065437 (l); WP_182594753.1, s11 (m);
WP_004630343.1, WCHRI065162 (n); WP_354534808.1, 1138 (o); and WP_024979717.1,
SSH4 (p). There was one exception for cluster d, which has two members with 24.97%
identity and 92% coverage by BLASTp comparison of their VirB4 protein sequences.
Therefore, the 40% sequence identity threshold was too stringent for this particular case. We
used the iTOL annotation editor for Google Sheets (38) and the results from BLASTp to
annotate the RSSC phytopathogen and non-RSSC environmental species trees. Please
note that the genomes SY1 and LMG10661 are suppressed by NCBI. However, the protein
sequence accessions provided are still accessible through NCBI.

Identifying putative region of transfer and putative cargo genes

The analysis of putative cargo genes was only performed for T4SS gene clusters from
complete genomes, so clusters b, h, j, k, o, and p were not analyzed due to a lack of
complete genomes. For the RSSC phytopathogens, 33 putative regions of transfer from 25
complete genomes were analyzed. For the non-RSSC environmentals, 33 putative regions
of transfer from 14 complete genomes were analyzed. For T4SS gene clusters encoded on
accessory plasmids, the putative region of transfer was considered to be the entire plasmid,
except for NZ_CP146093.1, which was an accessory plasmid with two T4SS clusters. For
NZ_CP146093.1 and for T4SS gene clusters encoded on the chromosome or megaplasmid,
the putative region of transfer was determined by manual inspection using synteny analysis
with clinker. During manual inspection, regions were determined by identifying genes that
consistently acted as a boundary, such as an integrase gene for cluster a, and/or by

identifying a consistent insertion location characteristic, such as a tRNA gene for cluster c.
See Supplemental Files S27-S36 for clinker output files with putative regions of transfer from
each cluster.

Within each putative region of transfer, we searched for type III secretion system (T3SS)
effector genes from the Ralsto T3E database (45) using BLAST+ with a threshold of 40%
amino acid identity (34) followed by manual verification of each hit. We also searched for
type VI secretion system (T6SS) *vgrG*-linked auxiliary clusters. We have previously curated
25 *vgrG*-linked auxiliary clusters from RSSC phytopathogen genomes (46) and we curated
19 additional *vgrG*-linked auxiliary clusters from non-RSSC environmental genomes using
the method described in Aoun and Georgoulis *et al.* (2024). We first used clinker to search
the putative regions of transfer for *vgrG*-linked auxiliary clusters based on a global amino
acid identity threshold of 30%. As a second approach, we performed BLASTp searches with
seven representative VgrG alleles with 20% local identity and 20% coverage thresholds. The
representatives were chosen using clinker to identify VgrG alleles with less than 30% global
amino acid identity between each other. We searched for hemagglutinin two-partner
secretion (TPS) clusters using BLASTp followed by manual verification of each hit,
regardless of percent identity and percent coverage (query sequences: WP_240442336.1,
WP_118891273.1, and WP_118891272.1). Additionally, we searched for metabolism-related
genes using DRAM with a bit score threshold of 60 (47), dbCAN2 with an e-value threshold
of 1e-15 (48), and ModelSEED v2 with sequential gapfilling (49) in KBase and METGeneDb
v2 (50) using BLAST+ with a 40% identity threshold. We combined the results of the four
different metabolism gene searches, then divided them into two categories: (i) heavy metal
metabolism and resistance genes and (ii) other metabolism genes. Finally, we used
MOBscan (51) to search for relaxase genes. MOBscan uses a protein sequence-based
HMM profile for each of the following MOB families: MOB_{P1}, MOB_{P2}, MOB_{P3}, MOB_Q, MOB_H,
MOB_C, MOB_V, MOB_B, MOB_F, MOB_T, and MOB_M.

Average nucleotide identity of putative regions of transfer

313　Average nucleotide identity (ANI) of putative regions of transfer was calculated using pyANI-
314　plus v0.0.1 (52) with the ANIb method, which uses BLAST+. Hierarchical clustering of the
315　ANI matrix was performed using Morpheus (53) with the Euclidean distance metric.

Statistical analysis

318　All graphing and statistical analyses were performed in GraphPad Prism v10.4.1 for
319　Windows.

Results

There are sixteen distinct T4SS gene clusters in the *Ralstonia* genus

To broadly identify T4SSs in *Ralstonia*, we used VirB4 as a marker. We built a hidden
Markov model (HMM) from 753 VirB4 protein sequences identified by preliminary BLASTp
searches against 923 Burkholderiaceae genomes. Querying the HMM against 644 RSSC

phytopathogen and 143 non-RSSC environmental genomes identified 498 VirB4 sequences.
A protein tree of the 498 sequences revealed at least 13 clades. To test the hypothesis that
there were multiple, distinct T4SS gene clusters, we analyzed the synteny and global amino
acid sequence identity of the *virB4* gene neighborhoods using clinker (33). In most cases,
gene organization patterns reinforced the phylogenetic groupings, but in a few cases,
synteny analysis allowed us to further divide certain groups. For example, two T4SS loci with
long VirB4 branch lengths were shown to belong to the same cluster based on their
consistent gene organization (cluster d). In two other cases, synteny conservation within a
cluster and differences in synteny between the clusters allowed us to differentiate and divide
the clusters (b vs. k and h vs. p). Through this combined phylogenetic and syntenic
approach, we ultimately identified 16 distinct T4SS gene clusters in the *Ralstonia* genus,
which we referred to as clusters a-p (**Figure 1**).

The clusters varied in their prevalence in the *Ralstonia* genus as well as in their prevalence
in the RSSC phytopathogens vs. the non-RSSC environmentals. The most commonly
identified T4SS clusters were cluster a (n=160), cluster i (n=75), and cluster f (n=74), while
the least commonly identified were cluster j (n=1), cluster o (n=1), and cluster d (n=2)
(**Figure 1**). Six T4SS gene clusters were found in both RSSC phytopathogen and non-RSSC
environmental genomes (clusters a, b, c, e, f, and i). Four T4SS clusters were exclusively
found in RSSC phytopathogen genomes (clusters d, g, h, and j) while six T4SS clusters
were exclusively found in the non-RSSC environmental genomes (clusters k, l, m, n, o, and
p).

Figure 1. Sequence diversity of the T4SS marker VirB4 in the *Ralstonia* genus. VirB4 protein sequences (n=498) were identified by a custom hidden Markov model (HMM) search, and an approximately-maximum-likelihood tree was built with FastTree 2 from a MUSCLE multiple sequence alignment (MSA) of the VirB4 sequences. The unrooted tree was visualized in iTOL and labeled using Affinity Designer. The 16 distinct T4SS gene clusters (a-p) are outlined and labeled. Circles denote clusters that are present in the RSSC phytopathogens and triangles denote clusters that are present in the non-RSSC environmentals. Colored clusters were found in RSSC phytopathogen genomes, and clusters with dashed outlines were not found in the RSSC phytopathogens. Supplemental files contain the *virB4* gene list (Supplemental Table S2A), the VirB4 protein sequence MSA (Supplemental File S5), the Newick file for the tree (File S6), a PDF of the unrooted tree with branch labels (File S7), and PDFs of the rectangular tree with branch labels, without and with colors (Files S8 and S9).

We selected one member of each of the 16 *Ralstonia* T4SS gene clusters to be reference clusters (**Figure 2**, Supplemental Table S2B). Reference clusters a-j are from RSSC phytopathogen genomes, and reference clusters k-p are from non-RSSC environmental genomes. The references include T4SS genes predicted by multiple tools including CONJscan/MacSyFinder, NCBI CDD, and NCBI PGAP. The T4SS reference clusters contain 14-28 genes, with a median of 20.5 genes. The *virB4* genes range in size from 2,375 to 3,065 bp. To assign *vir/tra/trb/tfc* gene names to the reference clusters, we used CONJscan to classify the T4SS gene clusters into the established F/P/I/GI-type classification system. P-type systems use the *virB*/*D4* naming convention, GI-type systems use *tfc*, and F-type and I-type systems both use *tra*/*trb*. Notably, the F-type and I-type systems reuse certain gene names for non-homologous genes. Twelve clusters were classified as P-type T4SSs (clusters a, b, e, f, g, h, i, k, m, n, o, and p), two as F-type (clusters d and l), one as I-type (cluster j), and one as GI-type (cluster c) (**Figure 2**, Supplemental Figure S1).

There are certain recurring gene organization patterns among the twelve P-type systems, despite the overall variability across the 16 T4SS gene clusters. These patterns in gene organization can be used to infer that a given T4SS gene cluster is a P-type T4SS. This is especially useful for more distantly related gene clusters where certain homologous genes have less than the 30% global identity, which is a default parameter for clinker. In all twelve cases, *virB4* is preceded by two small 225-417 bp genes on the same strand. In six cases, the two small genes are *virB2* and *virB3*, in that order. In the other six cases, only *virB3* is predicted by sequence alignment. The genes *virB8*, *virB9*, and *virB10* occur in succession, but in two clusters the *virB9* and *virB10* genes are separated by an additional gene. *virB11* either precedes *virB2*-*virB3*-*virB4* (8 clusters) or follows *virB8*-*virB9*-*virB10* (4 clusters). A second *virB9* paralog immediately follows *virB8*-*virB9*-*virB10* in four cases, only when that set of genes is not followed by *virB11*. To compare T4SSs in future genome sequences to the reference clusters, the GenBank files are available on Zenodo (DOI: 10.5281/zenodo.14873519; Supplemental Files S10-S26).

Figure 2. The T4SS gene clusters in *Ralstonia* have variable gene organization and composition.
Clusters are ordered based on their phylogenetic distribution: found only in RSSC phytopathogens,
found in both, or found only in non-RSSC environmentals. Genes in each cluster are labeled based
on F/P/I/GI-type classification conventions: *tra*/*trb* (F-type), *virB*/*D4* (P-type), an independent *tra*/*trb*

scheme (I-type), and *tfc* (GI-type). The *virB4* homologs have a bolded name (*traC*, *virB4*, *traU*, and
tfc16) and a bolded gene arrow border. The genes annotated as the type IV coupling protein are
labeled "CP". "LT" indicates genes annotated as lytic transglycosylase genes. Genes marked with an
asterisk (*) are T4SS genes that were not able to be annotated within the indicated F/P/I/GI-type
convention. Gene clusters were defined using Operon-mapper, MacSyFinder v2 with CONJscan
models, NCBI RefSeq annotations, and NCBI Conserved Domain Database. Gene clusters were
visualized with clinker and labeled with Affinity Designer. Gaps are displayed where the
representative cluster had regions without predicted T4SS genes; patterns of non-T4SS genes are
not always consistent among all members of each cluster. The representative cluster j is found on two
separate contigs in the only *Ralstonia* genome with cluster j. For additional details regarding genome,
gene, and protein NCBI accessions, see Supplemental Table S2B and Supplemental Files S10-S26.

T4SS gene clusters are more abundant in the non-RSSC environmentals than in the RSSC phytopathogens

We hypothesized that T4SS gene clusters would be more abundant in the RSSC
phytopathogens than in the non-RSSC environmentals because clinical pathogens are
known to have abundant mobile genetic elements (1,2,6,54). We first looked at T4SS
abundance qualitatively by visualizing presence and absence patterns on species trees of
644 RSSC phytopathogen genomes and 143 non-RSSC environmental genomes (**Figure
3A-B**). We queried the genomes for the VirB4 sequences from each reference cluster using
multiple BLASTp searches, identifying 498 VirB4 sequences. The BLASTp results were
consistent with the previous HMM search that led to Figure 1. Queries with VirB3 and VirB8
yielded a similar number of hits as VirB4 and protein trees with similar topologies (data not
shown), but VirB3 and VirB8 searches only identified the P-type clusters. Nevertheless,
during data curation, we identified and added five clusters with a pseudogenized *virB4* gene
to the tree (four cluster a and one cluster f). Overall, the T4SS gene clusters had a sporadic
phylogenetic distribution across both the RSSC phytopathogens and the non-RSSC
environmentals.

Because the non-RSSC environmental genomes appeared to harbor more T4SS clusters,
we quantified the abundance of T4SSs in each *Ralstonia* clade. We first quantified the
incidence of any T4SS cluster, revealing the non-RSSC environmentals encoded more
T4SS gene clusters (Chi-Square test, α = 0.001) (**Figure 3C**). Specifically, at least one
T4SS gene cluster was found in 77.6% (111/143) of non-RSSC environmental genomes and
26.6% (171/644) of RSSC phytopathogen genomes. As a second metric, we quantified the
number of T4SS gene clusters per genome. Once again, the non-RSSC environmental
genomes contained more T4SS gene clusters (Kolmogorov-Smirnov test, P < 0.0001)
(**Figure 3C**). The median number of T4SS gene clusters was 0 (IQR 0-1) in RSSC
phytopathogen genomes and 2 (IQR 1-3) in non-RSSC environmental genomes. Of the
genomes searched, no RSSC phytopathogen genomes had more than three T4SS gene
clusters, while the non-RSSC environmental genomes contained up to six T4SS gene
clusters.

A) *virB4* Gene Presence in RSSC

Cluster
a b c d e f g h i j

Absent Clusters: k, l, m, n, o, p

B) *virB4* Gene Presence in Non-RSSC *Ralstonia*

Cluster
a b c e f i k l m n o p

Absent Clusters: d, g, h, j

C) *virB4* Gene Presence in RSSC and Non-RSSC

Figure 3. RSSC phytopathogen and non-RSSC environmentals had different abundance and phylogenetic distributions of sixteen T4SS clusters. Phylogenetic distribution of **A)** T4SS clusters a-j across 644 RSSC phytopathogen genomes and **B)** clusters a-c, e-f, i, and k-p across 143 non-RSSC environmental genomes. The species trees are approximately-maximum-likelihood trees built with FastTree 2 using 49 conserved genes. The presence of the T4SS gene clusters was determined by BLASTp queries with each cluster's representative VirB4 sequence, and synteny was manually checked for multiple hits per cluster. Each column represents one T4SS gene cluster, with the dark rectangles indicating presence of the cluster and the light background indicating absence of the cluster in each genome. For the non-RSSC environmental species tree, species are designated by alternating gray shading, with names labeled for large groups. The missing labels are, from top to bottom: *Ralstonia sp.*, *R. condita*, *Ralstonia sp.*, *R. flaminis*, *Ralstonia sp.*, *R. psammae*, *R. holmesii*, *R. chuxiongensis*, and *R. edaphis.* The trees with presence/absence data were visualized in iTOL. Supplemental Table S2A contains a list of *virB4* genes, and Supplemental Files S2 and S3 contain PDFs of the phylogenetic trees with labeled branches. **C)** Top: the proportion of RSSC phytopathogen and non-RSSC environmental genomes containing at least one T4SS marker gene (*virB4*). Fewer RSSC phytopathogen genomes contained *virB4* genes (Chi-Square test, * corresponds to α = 0.001).

Bottom: the number of *virB4* copies per genome. Each point represents one genome. RSSC phytopathogen genomes had fewer T4SS gene clusters per genome than non-RSSC environmental genomes (Kolmogorov-Smirnov test, * corresponds to *P* < 0.0001).

We investigated whether the specific T4SS clusters varied in prevalence among RSSC phytopathogens and non-RSSC environmentals. When focusing only on the six T4SS gene clusters that were found in both the RSSC phytopathogens and non-RSSC environmentals (clusters a, b, c, e, f, and i), cluster a were consistently found in high prevalence and cluster b was consistently found in low prevalence. Cluster i was more common in the RSSC phytopathogens while clusters c, e, and f were more common in the non-RSSC environmentals.

The RSSC phytopathogen genes RSp0179 and RSp1521 are unlikely to be T4SS-associated

In RSSC phytopathogen genomes, two genes (RSp0179/RS_RS18010 and RSp1521/RS_RS24455) have been described in the literature as T4SS-associated because they are annotated as *virB10* and *virJ* by the NCBI CDD and other tools (55). However, we were skeptical that these genes were related to T4SSs because they were not present in any of the T4SS gene clusters we identified. In *Agrobacterium*, VirJ is a periplasmic protein that associates with the T4SS complex and the T4CP to aid in DNA transfer (56), and VirB10 is a component of the transmembrane channel of the T4SS complex (10,11,15,16,57). If RSp0179 and RSp1521 are associated with *Ralstonia* T4SSs, we would expect lines of evidence beyond annotation, such as being co-located with T4SS genes or sharing phylogenetic distributions with T4SS clusters. We inspected the neighborhoods of RSp0179 and RSp1521 in complete genomes with and without T4SS gene clusters (Figure S2). Both genes were located in consistent neighborhoods in separate regions of the megaplasmid. Importantly, neither gene was located near T4SS gene clusters. To test whether RSp0179 and RSp1521 have sporadic phylogenetic distributions similar to any of the 16 T4SS clusters, we conducted BLASTp searches against the *Ralstonia* genus using thresholds of 40% identity and 50% coverage. In contrast with the sporadic distribution of T4SS gene

clusters, RSp0179 and RSp1521 are broadly conserved in the phytopathogenic lineage or in
the genus as a whole, respectively. RSp0179 was present in 95.7% (377/394) of RSSC
phytopathogen genomes and none of the 143 non-RSSC environmental genomes, and
RSp1521 was present in 95.9% (378/394) of RSSC phytopathogen genomes and 100%
(143/143) of non-RSSC environmental genomes. Overall, these genes have different
evolutionary histories than T4SS gene clusters, so they are unlikely to be T4SS-associated,
despite their annotations.

The T4SS gene clusters vary in prevalence and co-occurrence patterns in *Ralstonia*
genomes

Since one genome can contain more than one T4SS gene cluster, we investigated which
specific clusters co-occurred and in what quantities (**Figure 4A**). Out of the genomes that
encoded at least one T4SS gene cluster, co-occurrence existed in 21.1% (36/171) of RSSC
phytopathogen genomes and 71.2% (79/111) of non-RSSC environmental genomes. We
found that co-occurrence was generally more common for more prevalent T4SS clusters
(RSSC: clusters a, i, f; non-RSSC: clusters a, f, c). However, there were exceptions where
low prevalence clusters co-occurred. Although RSSC phytopathogen cluster e occurred in
only seven genomes, cluster e co-occurred twice each with other rare clusters g and h,
which were only found in five and four genomes, respectively. Additionally, there were
exceptions where two reasonably abundant clusters were exclusively found in different
genomes; the non-RSSC environmentals contained 72 cluster a and 17 cluster m T4SSs,
but no genome contained both. Both of these clusters commonly co-occurred with other
T4SS clusters.

In several cases, two members of the same cluster exhibited co-occurrence (**Figure 4A**).
Five RSSC phytopathogen genomes and 24 non-RSSC environmental genomes contained
two or more cluster a T4SSs. Additionally, five non-RSSC environmental genomes contained
two cluster c T4SS. We ruled out a putative co-occurrence of cluster i in the draft genome of
RSSC phytopathogen strain CIP221_UW464, because both occurrences were on opposite
ends of the same contig. Those two putative cluster i copies had 99.95% nucleotide identity,
so the dual presence likely reflects error assembling a ~125 kb plasmid. Other than these
instances, no other T4SS gene clusters were observed to co-occur with themselves in the
same genome.

The T4SS gene clusters are variably located on the chromosome, megaplasmid, and
accessory plasmids

We investigated whether cluster a-p are predominantly maintained as accessory plasmids or
ICEs on the bipartite genomes, which are composed of a ~3.5 Mb chromosome and a ~2.0
Mb megaplasmid. In two cases, the co-occurrence analysis provided indirect evidence that
clusters were ICEs. In 29 genomes, there were multiple cluster a T4SSs, and in five
genomes there were two cluster c T4SSs (**Figure 4A**). Assuming that members of the same
cluster are likely to be part of the same plasmid incompatibility group, the co-occurrence
suggests that cluster a and c are not replicating plasmids. To confidently determine the
genomic location of the T4SS gene clusters, we narrowed our analysis to complete
genomes: 33 T4SS gene clusters from 25 RSSC phytopathogen genomes and 33 T4SS

gene clusters from 14 non-RSSC environmental genomes. With this dataset, we were able
to infer locations of clusters a, c, d, e, f, g, i, l, m, and n, but there were no complete
genomes containing clusters b, h, j, k, o, and p.

Analysis of complete genomes provided strong evidence that clusters a and c are often
ICEs. We analyzed 26 cluster a T4SSs, which were encoded on the chromosome (n=24) or
megaplasmid (n=2) (**Figure 4B**). For cluster c, we analyzed twelve T4SSs, revealing nine
chromosomal clusters, two megaplasmid clusters, and one accessory plasmid cluster
(**Figure 4B**). The cluster c on the accessory plasmid of non-RSSC environmental strain RRA
could be an ICE that integrated into an existing plasmid because this plasmid contained a
second T4SS (cluster f). Gonçalves *et al.* (2020) described six types of chromosomally-
encoded ICEs that align with this paper's cluster a (Tn*4371*, ICE*Rps1*, ICE*Rps2*, ICE*Rps3*,
and ICE*Rsy1*) and cluster c (ICE*Rm1*) (58). Thus, we support the inference that clusters a
and c are both generally maintained as ICEs.

The other clusters were predominantly present on accessory plasmids. For clusters d, e, g, i,
l, m, and n, we analyzed one, six, two, seven, one, one, and four T4SS gene clusters,
respectively. All of these clusters were identified on accessory plasmids (**Figure 4B**).
Moreover, the clusters did not confidently co-occur with variants of their own cluster type,
further supporting their plasmid nature. For cluster f, we analyzed six T4SSs. We found one
on a chromosome and five clusters on accessory plasmids, including the plasmid described
above that also contained cluster c (**Figure 4B**). Since clusters f and c were identified on
plasmids and as ICEs, it is possible that a cluster f ICE integrated into an existing cluster c
plasmid. However, the high prevalence of cluster c in ICE format and cluster f in plasmid
format suggests cluster c integrated into a cluster f plasmid. Analyzing a larger sample size
of *Ralstonia* T4SSs can shed light on which are commonly maintained as accessory
plasmids.

Figure 4. Genomic context sheds light on the ICE vs. accessory plasmid status of T4SS gene
clusters. **A)** Pairwise comparison of T4SS cluster co-occurrence in genomes with two or more
clusters. Numbers show how many genomes contain the indicated pair of T4SS gene clusters;
genomes with more than two clusters are counted for each pairwise co-occurrence. Colored circles
indicate clusters that were found in RSSC phytopathogen genomes, and black rings indicate clusters
that were not found in the RSSC phytopathogens. **B)** T4SS gene clusters are encoded on accessory
plasmids and ICEs. Rings indicate an accessory plasmid and rectangles indicate an ICE. T4SS
cluster locations were determined using only complete genomes. Supplemental Table S2C shows the
genomes used and the locus information for each T4SS region.

567 **At least fourteen T4SS gene clusters are predicted to be DNA conjugation systems**

568 We hypothesized that the T4SS gene clusters in *Ralstonia* may encode self-transmissible
569 DNA conjugation systems due to their sporadic distribution across the genus. Generally, the
570 T4SSs that function as DNA conjugation systems require the proteins that facilitate the
571 mobilization of the DNA cargo, the T4CP and relaxase. We searched for these mobilization
572 genes in complete genomes containing T4SS gene clusters and also draft genomes if they
573 contained a reference cluster from Figure 2, for a total of 38 T4SS gene clusters from 29
574 RSSC phytopathogen genomes and 39 T4SS gene clusters from 20 non-RSSC
575 environmental genomes.

576 To identify T4CP genes, we first searched the 16 reference T4SS gene clusters using
577 CONJscan/MacSyFinder. Then we used the synteny visualization tool clinker to confirm that
578 the T4CPs were conserved in all 77 T4SS gene clusters searched. To search for relaxase
579 genes in the vicinity of T4SS gene clusters, we used MOBscan. We identified a relaxase
580 gene near 84.4% (65/77) of T4SS gene clusters searched. The T4SS gene clusters without
581 an identifiable relaxase gene were cluster d (1/2), cluster e (7/7), and cluster m (2/2). Eight
582 of these are clearly encoded on accessory plasmids, and the other two are reference
583 clusters from draft genomes. Note that the absence of an identifiable relaxase gene is not
584 sufficient to exclude DNA conjugation as a possible function. Overall, all 16 T4SS clusters
585 appear to contain a T4CP (**Figure 2**) and 14 out of 16 are near a relaxase gene. Therefore,
586 we predict that at least 14 gene clusters are DNA conjugation systems.

587 Because we used MOBscan to identify relaxase genes, we were able to determine the MOB
588 class of each relaxase identified. MOBscan uses HMM profiles to classify MOB families. We
589 found three of the twelve MOBscan MOB families in our dataset: P1, H, and F. A MOB_{P1}
590 gene was associated with clusters a, b, h, i, j, and k; MOB_H with clusters c, f, n, and o; and
591 MOB_F with clusters d, g, l, and p. In general, the MOB class of the relaxase gene near a
592 T4SS gene cluster was consistent between members of the same cluster (Supplemental
593 Table S2A). This consistency further supports the differentiation of clusters, such as in the
594 case of the closely related clusters h (MOB_{P1}) and p (MOB_F).

595 Since we predicted that the majority of these T4SS gene clusters encode DNA conjugation
596 systems, we investigated their putative regions of transfer. To do this, we analyzed 33
597 putative regions of transfer from 25 complete RSSC phytopathogen genomes and 33
598 putative regions of transfer from 14 complete non-RSSC environmental genomes. For
599 accessory plasmids with a single T4SS cluster, the entire plasmid was considered the
600 putative region of transfer. Otherwise, the putative region of transfer was determined by
601 clinker synteny analysis, where the putative boundaries of the regions of transfer were
602 inferred by comparing closely related genomes with and without T4SS clusters. On the
603 plasmid with two T4SS clusters, synteny comparisons to single-T4SS plasmids and ICEs
604 revealed the likely boundaries of clusters f and c, respectively. Collectively, the sizes of the
605 putative regions of transfer were 32.6-389.8 kb (Figure S3A). Usually, members of the same
606 cluster had similarly sized putative regions of transfer. For example, the regions for cluster a
607 were 37.3-57.0 kb. In contrast, some regions had variable lengths, such as those from
608 cluster f members where size ranged from 146.4 to 379.1 kb.

609 Additionally, we compared the GC contents of the putative regions of transfer and their
610 parental genomes because differences in GC content could result from horizontal gene
611 transfer (Figure S3B) (59). The putative regions in RSSC phytopathogen genomes had 57.2-

64.3% GC, which was lower than the 66.5-67.0% GC of the whole genomes. For the non-
RSSC environmentals, the regions had 58.2-67.0% GC compared to the 63.0-66.0% GC of
the whole genomes. There was a significant difference in GC content between the regions
and whole genomes for T4SS gene clusters in the RSSC phytopathogens (Wilcoxon test, P
< 0.0001), but there was no significant difference in GC content for clusters in the non-RSSC
environmentals (Wilcoxon test, $P = 0.11$). When considering GC content of the region vs.
parental genome per cluster, the difference in GC content was significant for clusters a
(Wilcoxon test, $P < 0.0001$), e ($P = 0.031$), f ($P = 0.031$), and i ($P = 0.016$), but not significant
for clusters c ($P = 0.30$) and n ($P = 0.13$). Clusters d, g, l, and m had sample sizes too small
for statistical comparison.

RSSC phytopathogens and non-RSSC environmentals differ in cargo gene
composition of T4SSs

Because the RSSC phytopathogens and non-RSSC environmentals have different lifestyles,
we investigated if there was a difference in the cargo gene composition of their T4SSs'
putative regions of transfer. We focused on complete *Ralstonia* genomes to infer cargo gene
functions.

The RSSC phytopathogens manipulate plants by using a type III secretion system (T3SS) to
inject an arsenal of effector proteins into host cells (60), effector composition between RSSC
phytopathogen genomes is highly variable (45), and effectors have been previously
identified on an RSSC accessory plasmid (61). Thus, we hypothesized that RSSC
phytopathogen T4SSs might contribute to the dynamic patterns by carrying T3 effectors and
that T3 effectors would be more common in regions of transfer in RSSC phytopathogen
T4SSs. Five of 33 RSSC phytopathogen genomes and zero of 33 non-RSSC environmental
genomes had at least one T3 effector gene (**Figure 5**, Supplemental Table S2C), but there
was not a significant difference in frequency (Mann-Whitney test, $P = 0.053$). The cluster i
accessory plasmid in HA4-1 encoded one *ripP1* gene (also known as *ripBR* (61)) and two
ripAC paralogs (also known as *ripBS* (61)). The cluster i plasmids in 068_2 and 077-080_1
each encoded a single copy of *ripBH*. The cluster i plasmid in RS24 and the cluster f plasmid
in MolK2 each encoded a single copy of *ripBA*.

Recently, we showed that diverse classes of mobile genetic elements in RSSC
phytopathogen genomes are often associated with microbial warfare-associated type VI
secretion system (T6SS) genes, specifically the *vgrG*-linked auxiliary clusters with paired
toxins and antitoxins (46). Several of the *vgrG*-linked toxin clusters were located on T4SS-
containing plasmids, so we decided to investigate further. Here, we investigated the
Ralstonia T4SS regions of transfer for the 25 previously classified *vgrG*-linked auxiliary
clusters (46) and 19 additional *vgrG*-linked clusters from non-RSSC environmental genomes
(Lowe-Power, unpublished). Consistent with the known enrichment of *vgrG*-linked auxiliary
clusters in RSSC phytopathogen genomes (46), the RSSC phytopathogens had significantly
more T6 *vgrG*-linked auxiliary clusters per region of transfer than the non-RSSC
environmentals (Mann-Whitney test, $P = 0.011$) (**Figure 5**); seven of 33 RSSC
phytopathogen genomes and zero of 33 non-RSSC environmental genomes had a single
vgrG-linked auxiliary cluster (Supplemental Table S2C). The *vgrG*-linked auxiliary clusters
were encoded on cluster i and d accessory plasmids of RSSC phytopathogens. In addition to
the presence of *vgrG*-linked auxiliary clusters *aux6a* and *aux11* on four previously noted

cluster i containing accessory plasmids (46), *aux11* was encoded on the cluster i plasmids in
068_2 and 077-080_1, and *aux6a* was present on the cluster d accessory plasmid of this
study's newly sequenced genome of RSSC strain Gazipur 4.

Filamentous hemagglutinin-associated two-partner secretion (TPS) system gene clusters
have been previously identified in the RSSC phytopathogens (62) and hypothesized to
function as contact-dependent inhibition systems participating in kin-recognition and
competition (63,64). TPS systems are class b type V secretion systems (T5bSSs) known in
other bacteria to be involved in diverse functions like surface adhesion, bacterial
aggregation, contact-dependent inhibition, cytolysis/hemolysis, and iron acquisition (8,65–
71). Hemagglutinin TPS clusters include a transporter gene (*fhaC*) and at least one
hemagglutinin-repeat containing gene (*fhaB*/FHA) (65–71). The RSSC phytopathogens had
significantly more hemagglutinin TPS clusters per region of transfer than the non-RSSC
environmentals (Mann-Whitney test, P = 0.027) (**Figure 5**); eight of 33 RSSC phytopathogen
genomes and one of 33 non-RSSC environmental genomes had a single hemagglutinin TPS
cluster (Supplemental Table S2C). These clusters were encoded on seven cluster i
accessory plasmids and one cluster f accessory plasmid in RSSC phytopathogen genomes.
In the non-RSSC environmentals, there was one cluster n accessory plasmid in strain 12D
that encoded a hemagglutinin TPS cluster.

Because the non-RSSC environmentals occupy more diverse environments than the RSSC
phytopathogens, we reasoned that the non-RSSC environmentals might benefit from T4SS-
associated metabolism genes. To identify metabolism genes in the complete genome T4SS
regions of transfer, we used four tools: ModelSEED v2 (49), METGeneDb v2 (50), DRAM
(47), and dbCAN2 (48). We separated the aggregated results into two categories: (i) heavy
metal metabolism and resistance genes and (ii) other metabolism genes. Here, the term
"heavy metal" refers to transition (d-block) metals, post-transition (p-block) metals, and
metalloids. Other metabolism genes were defined as any metabolism genes other than
those identified as heavy metal metabolism and resistance genes.

Heavy metal metabolism and resistance genes were found in 22/33 non-RSSC
environmental regions of transfer but only 2/33 RSSC phytopathogen regions of transfer
(Supplemental Table S2C). The non-RSSC environmentals had significantly more heavy
metal metabolism and resistance genes per region of transfer than the RSSC
phytopathogens (Mann-Whitney test, P < 0.0001) (**Figure 5**). Within the non-RSSC
environmentals, the number of heavy metal metabolism and resistance genes per region of
transfer was 1-43 in the 22 regions identified. These genes were identified in regions of
transfer of diverse clusters: cluster a (n=2), c (n=9), e (n=1), f (n=5), l (n=1), m (n=1), and n
(n=3). Gene annotations suggest that these genes are related to mercury, nickel, zinc,
cadmium, lead, cobalt, arsenic, chromium, iron, copper, silver, antimony, and tellurium
interactions. Within the RSSC phytopathogens, six heavy metal metabolism and resistance
genes were identified within a cluster a region of transfer in strain UW386, and two genes
were identified on a cluster d accessory plasmid in strain Gazipur 4. Gene annotations
suggest that these genes are related to mercury and cobalt interactions, respectively.

Genes for other metabolic functions were found in 32/33 RSSC phytopathogen regions of
transfer and 33/33 non-RSSC environmental regions of transfer (Supplemental Table S2C).
Within the RSSC phytopathogens, the number of other metabolism genes per region of
transfer was 1-16 in the 32 regions identified. The single RSSC phytopathogen region for

which we did not identify other metabolism genes was the cluster g accessory plasmid in
strain SL1931. Within the non-RSSC environmentals, the number of other metabolism genes
per region of transfer was 1-34. Despite the trend, there was no significant difference in the
number of other metabolism genes per region of transfer between the RSSC
phytopathogens and the non-RSSC environmentals (Mann-Whitney test, *P* = 0.089) (**Figure**
5). An annotated clinker visualization of the cluster a putative regions of transfer is available
in Supplemental Figure S4. Clinker files for putative regions of transfer from each of the
other clusters with complete genomes are available in Supplemental Files S27-S36.

Figure 5. Cargo gene content of T4SS clusters differs between RSSC phytopathogens and non-
RSSC environmentals. The graphs show the number of type III secretion system (T3SS) effector
genes, type VI secretion system (T6SS) *vgrG*-linked auxiliary clusters, hemagglutinin two-partner
secretion (TPS) clusters, heavy metal metabolism and resistance genes, and other metabolism genes
present in each putative region of transfer. "Heavy metal" refers to transition (d-block) metals, post-
transition (p-block) metals, and metalloids. Each region of transfer is represented by a circle (RSSC
phytopathogen) or triangle (non-RSSC environmental) and is colored based on cluster identity.
Significance was determined using the Mann-Whitney test: * corresponds to *P* < 0.05, ** corresponds
to *P* < 0.0001, and ns corresponds to *P* > 0.05. Data was graphed and analyzed using GraphPad
Prism v10.4.1. See Supplemental Table S2C for additional details about the specific genes identified

in the putative regions of transfer. Supplemental Figure S3 conveys the GC content and sizes of the
putative regions of transfer.

──────────────────────────────

Specialization of a T4SS cluster to the RSSC phytopathogens

We hypothesized that cluster i might have specialized to the RSSC phytopathogens based
on the previous results. As described, cluster i T4SSs are more prevalent in RSSC
phytopathogens (n=71) as compared to non-RSSC environmentals (n=4). This is in contrast
to the overall trend of greater T4SS prevalence in the non-RSSC environmentals (**Figure 3**).

Closer analysis of the VirB4 phylogeny (**Figure 1**) revealed that the cluster i T4SSs in the
RSSC phytopathogens were almost entirely grouped by the species of *Ralstonia* they came
from (**Figure 6**). However, the branching pattern is not perfectly congruent with the species
tree made from the genomes that the cluster i T4SSs were found in. In contrast, the VirB4
phylogenies for other prevalent T4SS gene clusters, such as cluster a, were not grouped by
the *Ralstonia* species in which the VirB4 sequences were identified. The cluster i VirB4
phylogeny has four major clades (**Figure 6**). The first clade is the outgroup, which includes
two VirB4 sequences from one phylotype II *R. solanacearum* genome (Rs545_UW752) and
one *R. chuxiongensis* genome. The second clade includes six VirB4 sequences from three
phylotype I *R. pseudosolanacearum* genomes, two *R. chuxiongensis* genomes, and one *R.
psammae* genome. The third clade includes 16 VirB4 sequences from 16 phylotype I *R.
pseudosolanacearum* genomes. The fourth clade includes 51 VirB4 sequences from 49
phylotype II *R. solanacearum* genomes and one phylotype I *R. pseudosolanacearum*
genome (RALF-MA). The RSSC phytopathogen phylotypes exhibit different geographical
distributions, with phylotype I *R. pseudosolanacearum* originating in Asia and phylotype II *R.
solanacearum* originating in the Americas. The phylotype I *R. pseudosolanacearum* strain
RALF-MA was isolated from Costa Rica, most likely having been introduced to the Americas
in the last few hundred years. Therefore, the origin of the cluster i T4SS in RALF-MA could
be a phylotype II *R. solanacearum* strain.

We computed an ANI matrix for cluster i putative regions of transfer, which includes T4SS
genes. Seven cluster i VirB4 sequences were found in complete genomes and 68 in draft
genomes (n=75 total). For clusters in draft genomes, we treated the putative region of
transfer as the whole contig that encoded the *virB4* gene. Two of the 75 cluster i T4SSs
were found in nearly identical copies on either end of the same contig and were treated as
one cluster. Additionally, two clusters were found in early genomes that had hundreds of
contigs concatenated randomly into one sequence before submission to NCBI; these were
excluded from the analysis because we could not confidently delineate contig boundaries.
The ANI matrix was computed with ANIb using pyANI-plus for the 72 remaining regions.
Hierarchical clustering using Morpheus resulted in a dendrogram with the same four major
clades as the VirB4 phylogeny (Supplemental File S37).

Our cargo gene analysis revealed that the T3 effectors, T6 *vgrG*-linked auxiliary clusters,
and hemagglutinin TPS clusters were common within cluster i regions (**Figure 5**). To
discover if the trend was consistent, we searched for these putatively adaptive genes in 72
cluster i putative regions of transfer from draft and complete *Ralstonia* genomes
(Supplemental Table S2D, Supplemental File S38). Four regions (5.6%) had at least one T3

effector gene and these regions were all previously described accessory plasmids from
complete genomes (**Figures 5** and **6**). These regions were grouped within clades 3 and 4 of
the cluster i VirB4 phylogeny. We identified a single T6 *vgrG* gene in 50 regions (69.4%), five
from *aux6a*, five from *aux6b*, 29 from *aux11*, and a unique *aux9*-like cluster with divergent
gene content from the previously classified *aux9a* and *aux9b* (**Figure 6**) (46). Ten of the
identified *vgrG* genes were not able to be matched with a particular auxiliary cluster because
they were truncated at the ends of contigs. The different T6 auxiliary clusters corresponded
with the clades of the VirB4 phylogeny: *aux6b* and *aux9a/b*-like in clade 2, *aux6a* in clade 3,
and *aux11* in clade 4. The identified T6 auxiliary clusters were generally in similar locations a
few genes downstream of the T4SS cluster. No *vgrG* genes were identified in the remaining
22 regions, however, at least 16 of those regions had contigs that ended before the
expected location of a T6 auxiliary cluster. We identified hemagglutinin TPS clusters in 35
regions of transfer (48.6%) (**Figure 6**). Of these clusters, 17 were whole clusters, meaning
they included both *fhaC* and *fhaB* genes. The 18 partial clusters had at least one component
of a TPS cluster, with the other part truncated by the ends of contigs. These TPS clusters
were found in clades 2, 3, and 4 of the VirB4 phylogeny.

Figure 6. Phylogenetic analysis suggests cluster i T4SSs evolved a specialized relationship with RSSC phytopathogens. The left cladogram is a VirB4 protein tree made from an MSA of 75 cluster i sequences. The right tree is a species tree made from the 74 *Ralstonia* genomes that contain a cluster i T4SS. The columns show the presence of specific T3 effector genes, T6 *vgrG*-linked auxiliary clusters, and hemagglutinin TPS clusters in each cluster i T4SS. The sizes of the putative regions of transfer are noted, bold indicates the cluster is encoded in a complete genome, and grayed "N/A" labels are listed for T4SSs encoded in old genomes with contigs randomly concatenated into one scaffold. Branches of the phylogenetic trees were rotated manually in iTOL, and the tanglegram linkages and labels were added in Affinity Designer. See Supplemental File S37 for a cluster i region average nucleotide identity (ANI) matrix and Supplemental File S38 for a cluster i region clinker visualization.

Discussion

Contrary to our initial hypothesis, T4SS were less prevalent in RSSC phytopathogen genomes than in non-RSSC environmental genomes. Notably, there were functional differences in the genetic cargo associated with the T4SSs. What does this mean about the eco-evolutionary differences between the three RSSC phytopathogen species and the 17 non-RSSC environmental species?

Assuming that this study's T4SS clusters are *bona fide* conjugative systems, the non-RSSC environmental species may inhabit bacterial communities with a greater diversity of conjugative donors, have fewer physiological barriers to acquiring and maintaining conjugated DNA, or more often benefit from the functions conferred by the conjugative elements. The prevalence of heavy metal metabolism and resistance genes in non-RSSC environmental putative regions of transfer may reflect their broader environmental distribution, in which exposure to toxic heavy metals may be more frequent. Correspondingly, many bacteria have evolved various mechanisms to resist the toxicity (72,73) and/or metabolize these metals. Therefore, the enrichment of heavy metal resistance genes in the putative T4SS regions of transfer is consistent with horizontal acquisition of adaptive traits.

The three monophyletic RSSC phytopathogen species have a more tailored and predictable life cycle of plant infection and environmental survival in soil and surface water. The phytopathogens' lower prevalence of T4SSs is surprising because they are known to be naturally transformable (74,75). Additionally, T4SSs benefit other phytopathogens by conferring virulence (76,77). Thus, the RSSC phytopathogens might encounter barriers to acquiring conjugated DNA. For example, they may encounter a limited pool of non-kin cells that could donate conjugative elements during their explosive replication in plants. Nevertheless, we and others have identified ecologically beneficial genetic cargo on some of the T4SS regions of transfer in RSSC phytopathogen genomes: T3 effectors, and auxiliary T6 toxin clusters and hemagglutinin TPS clusters (45,46,58,61–64). These three classes of genes mediate or putatively mediate interactions with RSSC phytopathogens' ecological targets: plants (T3 effectors) and microbial competitors (T6 toxins and hemagglutinin TPS toxins). At the genome level, it has been previously established that T6 auxiliary clusters are enriched in the RSSC phytopathogens (46), and we speculate that systematic comparison

will also show that phytopathogen genomes are also enriched in T3 effectors and
hemagglutinin TPS clusters.

Interestingly, cluster i appears to be relatively specialized to RSSC phytopathogens. Their
regions of transfer often encoded T6 *vgrG*-linked auxiliary clusters and hemagglutinin TPS
clusters, while they occasionally encoded T3 effector genes. Cluster i T4SSs were enriched
in the RSSC phytopathogens as compared to the non-RSSC environmentals. Additionally, in
the VirB4 phylogeny, the cluster i T4SSs in the RSSC phytopathogens were strongly
clustered by the species of *Ralstonia* they came from. Based on these results, we speculate
that cluster i T4SSs are specialized for RSSC phytopathogens.

Several of the T4SS gene clusters that we named have been previously identified in
comparative genomics papers: cluster a ICEs (55,58,61,62,78–81), cluster c ICEs (58),
cluster g plasmids (55,61,78,82), and cluster i plasmids (61,79). Additionally, Morales and
Sequeira (1985) identified and estimated the sizes of several accessory plasmids in RSSC
phytopathogen strains, including a 120 MDa (~197 kb) plasmid in UW256, which likely
encodes the cluster i T4SS identified in the UW256 genome (83). Although multiple T4SS
clusters are present on accessory plasmids, there are also accessory plasmids in *Ralstonia*
genomes that lack T4SS gene clusters (data not shown). One of these T4SS-null accessory
plasmids co-occurs with a cluster i plasmid in UW163, so it is possible that it is mobilizable
by the cluster i T4SS.

This study has several limitations. Like any survey of public genomic data, a limitation of this
study is that there is sampling bias in publicly available data. Although more diverse
Ralstonia genomes have become available in recent years, our dataset is not completely
representative of the total biological variation of this genus. Additionally, it is likely that our
T4SS search approach yielded occasional false negatives against the analyzed genomes
because the HMM and BLASTp searches queried translated amino acid sequences. For
instance, we manually identified five additional T4SS gene clusters with *virB4* genes that
were pseudogenized after identifying the initial 498 VirB4 sequences. Another limitation lies
in the identification of metabolism-related genetic cargo. Although we used four databases to
identify metabolic genes, there are certainly metabolic proteins that are unrepresented in
available databases.

We used the identification of a T4CP gene and a relaxase gene to infer that most of the
identified T4SSs could be DNA conjugation systems. This inferential approach cannot
differentiate between DNA conjugation and DNA release systems because experimental
testing is needed. However, DNA release systems have only rarely been identified; DNA
release systems are limited to the T4SS encoded on the gonococcal genetic island (GGI) in
Neisseria gonorrhoeae (84,85) and a T4SS in nontypeable *Haemophilus influenzae* (86).
Therefore, it is more likely that a T4SS with a T4CP and an associated relaxase functions as
a DNA conjugation system rather than a DNA release system. Although, the possibility that
these T4SSs could function as DNA release systems introduces an alternative hypothesis
for their mechanism of transmission to *Ralstonia* species. It has been determined that RSSC
phytopathogens are naturally transformable in multiple conditions including *in vitro* and *in*
planta, and are able to take up exogenous DNA ranging in size from 30 to 90 kb (74,75).
Therefore, natural transformation has the potential to contribute to T4SS acquisition. DNA
release via a T4SS could provide the exogenous DNA for natural transformation. This is the
mechanism of horizontal transfer of the GGI of *Neisseria gonorrhoeae* (84,85). The GGI is

released by its T4SS, then the naturally transformable *Neisseria* spp. in the vicinity
recognize a specific sequence in the released DNA that triggers uptake by natural
transformation (84,87). Nevertheless, the circumstantial evidence and Occam's razor
suggest most of the T4SSs in this study are part of self-transmissible mobile genetic
elements.

Overall, the differences in T4SS prevalence and cargo gene composition between the RSSC
phytopathogens and non-RSSC environmentals likely reflect differences in their ecological
niches.

Author statements

Author contributions

TCC: Conceptualization, Data curation, Analysis, Experiments / Investigation, Supervision,
Visualization, Writing – original draft, MLCA: Conceptualization, Project administration,
Supervision, Writing – review & editing, GYS: Conceptualization, Experiments /
Investigation, Writing – review & editing, AJB: Resources, Supervision, Writing – review &
editing, SCDC: Experiments / Investigation, Resources, Writing – review & editing, KN:
Resources, Writing – review & editing, AS: Resources, Writing – review & editing, TMLP:
Conceptualization, Project administration, Resources, Supervision, Writing – review &
editing.

Conflicts of interest

894 The authors declare that there are no conflicts of interest.

Funding information

897 This work is supported by the joint NSF / USDA NIFA Plant Biotic Interactions program (NSF
898 award # 2336557 and NIFA award # 2024-67013-43303), USDA NIFA Pests and Beneficial
899 Species in Agricultural Production Systems (A1112) program (award # 2024-67013-42781),
900 the USDA Hatch Program (Project #1023861), USDA NIFA predoctoral fellowship #2024-
901 67011-42914 to M. Cope-Arguello, and a Provost's Undergraduate Fellowship to T. Cowell.

Acknowledgements

904 We would like to acknowledge Titus Brown and Clay Fuqua for their valuable discussions,
905 and Jonathan Jacobs and Taylor Klass for sharing unpublished genomes.

References

1.   Hazen TH, Pan L, Gu JD, Sobecky PA. The contribution of mobile genetic elements to
the evolution and ecology of *Vibrios. FEMS Microbiology Ecology*. 2010;74(3): 485–
499. https://doi.org/10.1111/j.1574-6941.2010.00937.x.

2.　Tenaillon O, Skurnik D, Picard B, Denamur E. The population genetics of commensal
Escherichia coli. *Nature Reviews Microbiology*. 2010;8(3): 207–217.
https://doi.org/10.1038/nrmicro2298.

3.　Savory EA, Fuller SL, Weisberg AJ, Thomas WJ, Gordon MI, Stevens DM, et al.
Evolutionary transitions between beneficial and phytopathogenic *Rhodococcus*
challenge disease management. *eLife*. 2017;6: e30925.
https://doi.org/10.7554/eLife.30925.

4.　Martínez JL. Ecology and Evolution of Chromosomal Gene Transfer between
Environmental Microorganisms and Pathogens. Baquero F, Bouza E, Gutiérrez-
Fuentes JA, Coque TM (eds) *Microbiology Spectrum*. 2018;6(1).
https://doi.org/10.1128/microbiolspec.MTBP-0006-2016.

5.　Melnyk RA, Hossain SS, Haney CH. Convergent gain and loss of genomic islands drive
lifestyle changes in plant-associated *Pseudomonas*. *The ISME Journal*. 2019;13(6):
1575–1588. https://doi.org/10.1038/s41396-019-0372-5.

6.　Gyles C, Boerlin P. Horizontally Transferred Genetic Elements and Their Role in
Pathogenesis of Bacterial Disease. *Veterinary Pathology*. 2014;51(2): 328–340.
https://doi.org/10.1177/0300985813511131.

7.　Fernandes AS, Gonçalves OS, De Lima LMO, Santana MF. Does the mobilome of
Ralstonia solanacearum influence the evolution and virulence of this pathogen?
Tropical Plant Pathology. 2025;50(1): 29. https://doi.org/10.1007/s40858-025-00718-z.

8.　Chang JH, Desveaux D, Creason AL. The ABCs and 123s of Bacterial Secretion
Systems in Plant Pathogenesis. *Annual Review of Phytopathology*. 2014;52(1): 317–
345. https://doi.org/10.1146/annurev-phyto-011014-015624.

9.　Johnson CM, Grossman AD. Integrative and Conjugative Elements (ICEs): What They
Do and How They Work. *Annual Review of Genetics*. 2015;49(1): 577–601.
https://doi.org/10.1146/annurev-genet-112414-055018.

10.　Alvarez-Martinez CE, Christie PJ. Biological Diversity of Prokaryotic Type IV Secretion
Systems. *Microbiology and Molecular Biology Reviews*. 2009;73(4): 775–808.
https://doi.org/10.1128/MMBR.00023-09.

11.　Juhas M, Crook DW, Hood DW. Type IV secretion systems: tools of bacterial horizontal
gene transfer and virulence. *Cellular Microbiology*. 2008;10(12): 2377–2386.
https://doi.org/10.1111/j.1462-5822.2008.01187.x.

12.　Fronzes R, Christie PJ, Waksman G. The structural biology of type IV secretion
systems. *Nature Reviews Microbiology*. 2009;7(10): 703–714.
https://doi.org/10.1038/nrmicro2218.

13.　Guglielmini J, De La Cruz F, Rocha EPC. Evolution of Conjugation and Type IV
Secretion Systems. *Molecular Biology and Evolution*. 2013;30(2): 315–331.
https://doi.org/10.1093/molbev/mss221.

14.　Costa TRD, Patkowski JB, Macé K, Christie PJ, Waksman G. Structural and functional
diversity of type IV secretion systems. *Nature Reviews Microbiology*. 2024;22(3): 170–
185. https://doi.org/10.1038/s41579-023-00974-3.

15.　Christie PJ. The Mosaic Type IV Secretion Systems. Lovett ST, Bernstein HD (eds)

EcoSal Plus. 2016;7(1). https://doi.org/10.1128/ecosalplus.esp-0020-2015.

16.   Cascales E, Christie PJ. The versatile bacterial type IV secretion systems. *Nature*
Reviews Microbiology. 2003;1(2): 137–149. https://doi.org/10.1038/nrmicro753.

17.   Garcillán-Barcia MP, Francia MV, De La Cruz F. The diversity of conjugative relaxases
and its application in plasmid classification. *FEMS Microbiology Reviews*. 2009;33(3):
657–687. https://doi.org/10.1111/j.1574-6976.2009.00168.x.

18.   Prior P, Ailloud F, Dalsing BL, Remenant B, Sanchez B, Allen C. Genomic and
proteomic evidence supporting the division of the plant pathogen *Ralstonia*
solanacearum into three species. *BMC Genomics*. 2016;17(1): 90.
https://doi.org/10.1186/s12864-016-2413-z.

19.   Safni I, Subandiyah S, Fegan M. Ecology, Epidemiology and Disease Management of
Ralstonia syzygii in Indonesia. *Frontiers in Microbiology*. 2018;9: 419.
https://doi.org/10.3389/fmicb.2018.00419.

20.   Ryan MP, Adley CC. *Ralstonia* spp.: emerging global opportunistic pathogens.
European Journal of Clinical Microbiology & Infectious Diseases. 2014;33(3): 291–304.
https://doi.org/10.1007/s10096-013-1975-9.

21.   Bartoli C, Roux F, Lamichhane JR. Molecular mechanisms underlying the emergence
of bacterial pathogens: an ecological perspective. *Molecular Plant Pathology*.
2016;17(2): 303–310. https://doi.org/10.1111/mpp.12284.

22.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing
the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Research*. 2015;25: 1043–1055.
https://doi.org/10.1101/gr.186072.114.

23.   Arkin AP, Cottingham RW, Henry CS, Harris NL, Stevens RL, Maslov S, et al. KBase:
The United States Department of Energy Systems Biology Knowledgebase. *Nature*
Biotechnology. 2018;36(7): 566–569. https://doi.org/10.1038/nbt.4163.

24.   Greenrod STE, Stoycheva M, Elphinstone J, Friman VP. Global diversity and
distribution of prophages are lineage-specific within the *Ralstonia solanacearum*
species complex. *BMC Genomics*. 2022;23: 689. https://doi.org/10.1186/s12864-022-
08909-7.

25.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
data. Birol I (ed.) *Bioinformatics*. 2014;30(14): 2114–2120.
https://doi.org/10.1093/bioinformatics/btu170.

26.   Wick RR. *Filtlong*. GitHub. https://github.com/rrwick/Filtlong

27.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: A
New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing.
Journal of Computational Biology. 2012;19(5): 455–477.
https://doi.org/10.1089/cmb.2012.0021.

28.   Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome
assemblies from short and long sequencing reads. *PLOS Computational Biology*.
2017;13(6): e1005595. https://doi.org/10.1371/journal.pcbi.1005595.

994 29. Seemann T. Prokka: rapid prokaryotic genome annotation. Valencia A (ed.)
995   *Bioinformatics*. 2014;30(14): 2068–2069. https://doi.org/10.1093/bioinformatics/btu153.

996 30. Elphinstone JG, Hennessy J, Wilson JK, Stead DE. Sensitivity of different methods for
997   the detection of *Ralstonia solanacearum* in potato tuber extracts. *EPPO Bulletin*.
998   1996;26(3–4): 663–678. https://doi.org/10.1111/j.1365-2338.1996.tb01511.x.

999 31. Kelman A. The relationship of pathogenicity of *Pseudomonas solanacearum* to colony
1000   appearance in a tetrazolium medium. *Phytopathology*. 1954;44(12): 693–695.

1001 32. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using
1002   repeat graphs. *Nature Biotechnology*. 2019;37(5): 540–546.
1003   https://doi.org/10.1038/s41587-019-0072-8.

1004 33. Gilchrist CLM, Chooi YH. clinker & clustermap.js: automatic generation of gene cluster
1005   comparison figures. Robinson P (ed.) *Bioinformatics*. 2021;37(16): 2473–2475.
1006   https://doi.org/10.1093/bioinformatics/btab007.

1007 34. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:
1008   architecture and applications. *BMC Bioinformatics*. 2009;10(1): 421.
1009   https://doi.org/10.1186/1471-2105-10-421.

1010 35. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high
1011   throughput. *Nucleic Acids Research*. 2004;32(5): 1792–1797.
1012   https://doi.org/10.1093/nar/gkh340.

1013 36. Eddy SR. Accelerated Profile HMM Searches. Pearson WR (ed.) *PLoS Computational*
1014   *Biology*. 2011;7(10): e1002195. https://doi.org/10.1371/journal.pcbi.1002195.

1015 37. Price MN, Dehal PS, Arkin AP. FastTree 2 – Approximately Maximum-Likelihood Trees
1016   for Large Alignments. Poon AFY (ed.) *PLoS ONE*. 2010;5(3): e9490.
1017   https://doi.org/10.1371/journal.pone.0009490.

1018 38. Letunic I, Bork P. Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic
1019   tree display and annotation tool. *Nucleic Acids Research*. 2024;52(W1): W78–W82.
1020   https://doi.org/10.1093/nar/gkae268.

1021 39. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, et al.
1022   NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Research*. 2016;44(14):
1023   6614–6624. https://doi.org/10.1093/nar/gkw569.

1024 40. Wang J, Chitsaz F, Derbyshire MK, Gonzales NR, Gwadz M, Lu S, et al. The
1025   conserved domain database in 2023. *Nucleic Acids Research*. 2023;51(D1): D384–
1026   D388. https://doi.org/10.1093/nar/gkac1096.

1027 41. Cury J, Abby SS, Doppelt-Azeroual O, Néron B, Rocha EPC. Identifying Conjugative
1028   Plasmids and Integrative Conjugative Elements with CONJscan. In: De La Cruz F (ed.)
1029   *Horizontal Gene Transfer*. New York, NY: Springer US; 2020. p. 265–283.
1030   https://doi.org/10.1007/978-1-4939-9877-7_19.

1031 42. Néron B, Denise R, Coluzzi C, Touchon M, Rocha EPC, Abby SS. MacSyFinder v2:
1032   Improved modelling and search engine to identify molecular systems in genomes. *Peer*
1033   *Community Journal*. 2023;3: e28. https://doi.org/10.24072/pcjournal.250.

1034 43. Taboada B, Estrada K, Ciria R, Merino E. Operon-mapper: a web server for precise

operon identification in bacterial and archaeal genomes. Hancock J (ed.)
Bioinformatics. 2018;34(23): 4118–4120. https://doi.org/10.1093/bioinformatics/bty496.

44. Guglielmini J, Néron B, Abby SS, Garcillán-Barcia MP, La Cruz FD, Rocha EPC. Key
components of the eight classes of type IV secretion systems involved in bacterial
conjugation or protein secretion. *Nucleic Acids Research*. 2014;42(9): 5715–5727.
https://doi.org/10.1093/nar/gku194.

45. Peeters N, Carrère S, Anisimova M, Plener L, Cazalé AC, Genin S. Repertoire, unified
nomenclature and evolution of the Type III effector gene set in the *Ralstonia*
solanacearum species complex. *BMC Genomics*. 2013;14(1): 859.
https://doi.org/10.1186/1471-2164-14-859.

46. Aoun N, Georgoulis SJ, Avalos JK, Grulla KJ, Miqueo K, Tom C, et al. A pangenomic
atlas reveals eco-evolutionary dynamics that shape type VI secretion systems in plant-
pathogenic *Ralstonia*. Ruby EG, Allen C (eds) *mBio*. 2024;15(10): e00323-24.
https://doi.org/10.1128/mbio.00323-24.

47. Shaffer M, Borton MA, Bolduc B, Faria JP, Flynn RM, Ghadermazi P, et al. kb_DRAM:
annotation and metabolic profiling of genomes with DRAM in KBase. Marschall T (ed.)
Bioinformatics. 2023;39(4): btad110. https://doi.org/10.1093/bioinformatics/btad110.

48. Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, et al. dbCAN2: a meta server
for automated carbohydrate-active enzyme annotation. *Nucleic Acids Research*.
2018;46(W1): W95–W101. https://doi.org/10.1093/nar/gky418.

49. Faria JP, Liu F, Edirisinghe JN, Gupta N, Seaver SMD, Freiburger AP, et al.
ModelSEED v2: High-throughput genome-scale metabolic model reconstruction with
enhanced energy biosynthesis pathway prediction. *bioRxiv*. 2023;
https://doi.org/10.1101/2023.10.04.556561.

50. Dziurzynski M, Gorecki A, Decewicz P, Ciuchcinski K, Dabrowska M, Dziewit L.
Development of the LCPDb-MET database facilitating selection of PCR primers for the
detection of metal metabolism and resistance genes in bacteria. *Ecological Indicators*.
2022;145: 109606. https://doi.org/10.1016/j.ecolind.2022.109606.

51. Garcillán-Barcia MP, Redondo-Salvo S, Vielva L, De La Cruz F. MOBscan: Automated
Annotation of MOB Relaxases. In: De La Cruz F (ed.) *Horizontal Gene Transfer*. New
York, NY: Springer US; 2020. p. 295–308. https://doi.org/10.1007/978-1-4939-9877-
7_21.

52. Pritchard L, Glover RH, Humphris S, Elphinstone JG, Toth IK. Genomics and taxonomy
in diagnostics for food security: soft-rotting enterobacterial plant pathogens. *Analytical*
Methods. 2016;8: 12–24. https://doi.org/10.1039/C5AY02550H.

53. *Morpheus*. https://software.broadinstitute.org/morpheus

54. Weisberg AJ, Chang JH. Mobile Genetic Element Flexibility as an Underlying Principle
to Bacterial Evolution. *Annual Review of Microbiology*. 2023;77(1): 603–624.
https://doi.org/10.1146/annurev-micro-032521-022006.

55. Li P, Wang D, Yan J, Zhou J, Deng Y, Jiang Z, et al. Genomic Analysis of Phylotype I
Strain EP1 Reveals Substantial Divergence from Other Strains in the *Ralstonia*
solanacearum Species Complex. *Frontiers in Microbiology*. 2016;7: 1719.
https://doi.org/10.3389/fmicb.2016.01719.

56. Pantoja M, Chen L, Chen Y, Nester EW. *Agrobacterium* type IV secretion is a two-step
process in which export substrates associate with the virulence protein VirJ in the
periplasm. *Molecular Microbiology*. 2002;45(5): 1325–1335.
https://doi.org/10.1046/j.1365-2958.2002.03098.x.

57. Das A, Xie YH. The *Agrobacterium* T-DNA Transport Pore Proteins VirB8, VirB9, and
VirB10 Interact with One Another. *Journal of Bacteriology*. 2000;182(3): 758–763.
https://doi.org/10.1128/JB.182.3.758-763.2000.

58. Gonçalves OS, De Queiroz MV, Santana MF. Potential evolutionary impact of
integrative and conjugative elements (ICEs) and genomic islands in the *Ralstonia
solanacearum* species complex. *Scientific Reports*. 2020;10(1): 12498.
https://doi.org/10.1038/s41598-020-69490-1.

59. Ravenhall M, Škunca N, Lassalle F, Dessimoz C. Inferring Horizontal Gene Transfer.
Wodak S (ed.) *PLOS Computational Biology*. 2015;11(5): e1004095.
https://doi.org/10.1371/journal.pcbi.1004095.

60. Macho AP. Subversion of plant cellular functions by bacterial type-III effectors: beyond
suppression of immunity. *New Phytologist*. 2016;210(1): 51–57.
https://doi.org/10.1111/nph.13605.

61. Tan X, Qiu H, Li F, Cheng D, Zheng X, Wang B, et al. Complete Genome Sequence of
Sequevar 14M *Ralstonia solanacearum* Strain HA4-1 Reveals Novel Type III Effectors
Acquired Through Horizontal Gene Transfer. *Frontiers in Microbiology*. 2019;10: 1893.
https://doi.org/10.3389/fmicb.2019.01893.

62. Genin S, Boucher C. Lessons Learned from the Genome Analysis of *Ralstonia
solanacearum*. *Annual Review of Phytopathology*. 2004;42(1): 107–134.
https://doi.org/10.1146/annurev.phyto.42.011204.104301.

63. Evseeva D, Pecrix Y, Kucka M, Weiler C, Franzl C, Vlková-Žlebková M, et al.
Emergence and host range expansion of an epidemic lineage of *Ralstonia
solanacearum*. *bioRxiv*. 2024; https://doi.org/10.1101/2024.10.16.618685.

64. Prokchorchik M, Pandey A, Moon H, Kim W, Jeon H, Jung G, et al. Host adaptation
and microbial competition drive *Ralstonia solanacearum* phylotype I evolution in the
Republic of Korea. *Microbial Genomics*. 2020;6(11).
https://doi.org/10.1099/mgen.0.000461.

65. Nash ZM, Cotter PA. *Bordetella* Filamentous Hemagglutinin, a Model for the Two-
Partner Secretion Pathway. Sandkvist M, Cascales E, Christie PJ (eds) *Microbiology
Spectrum*. 2019;7(2): PSIB-0024-2018. https://doi.org/10.1128/microbiolspec.psib-
0024-2018.

66. Hodak H, Clantin B, Willery E, Villeret V, Locht C, Jacob-Dubuisson F. Secretion signal
of the filamentous haemagglutinin, a model two-partner secretion substrate. *Molecular
Microbiology*. 2006;61(2): 368–382. https://doi.org/10.1111/j.1365-2958.2006.05242.x.

67. Pérez A, Merino M, Rumbo-Feal S, Álvarez-Fraga L, Vallejo JA, Beceiro A, et al. The
FhaB/FhaC two-partner secretion system is involved in adhesion of *Acinetobacter
baumannii* AbH12O-A2 strain. *Virulence*. 2017;8(6): 959–974.
https://doi.org/10.1080/21505594.2016.1262313.

68. Trouillon J, Attrée I, Elsen S. The regulation of bacterial two-partner secretion systems.

Molecular Microbiology. 2023;120(2): 159–177. https://doi.org/10.1111/mmi.15112.

69. Jacob-Dubuisson F, Guérin J, Baelen S, Clantin B. Two-partner secretion: as simple as it sounds? *Research in Microbiology.* 2013;164(6): 583–595. https://doi.org/10.1016/j.resmic.2013.03.009.

70. Guérin J, Bigot S, Schneider R, Buchanan SK, Jacob-Dubuisson F. Two-Partner Secretion: Combining Efficiency and Simplicity in the Secretion of Large Proteins for Bacteria-Host and Bacteria-Bacteria Interactions. *Frontiers in Cellular and Infection Microbiology.* 2017;7: 148. https://doi.org/10.3389/fcimb.2017.00148.

71. Leo JC, Grin I, Linke D. Type V secretion: mechanism(s) of autotransport through the bacterial outer membrane. *Philosophical Transactions of the Royal Society B: Biological Sciences.* 2012;367: 1088–1101. https://doi.org/10.1098/rstb.2011.0208.

72. Mathivanan K, Chandirika JU, Vinothkanna A, Yin H, Liu X, Meng D. Bacterial adaptive strategies to cope with metal toxicity in the contaminated environment – A review. *Ecotoxicology and Environmental Safety.* 2021;226: 112863. https://doi.org/10.1016/j.ecoenv.2021.112863.

73. Pal A, Bhattacharjee S, Saha J, Sarkar M, Mandal P. Bacterial survival strategies and responses under heavy metal stress: a comprehensive overview. *Critical Reviews in Microbiology.* 2022;48(3): 327–355. https://doi.org/10.1080/1040841X.2021.1970512.

74. Coupat B, Chaumeille-Dole F, Fall S, Prior P, Simonet P, Nesme X, et al. Natural transformation in the *Ralstonia solanacearum* species complex: number and size of DNA that can be transferred. *FEMS Microbiology Ecology.* 2008;66(1): 14–24. https://doi.org/10.1111/j.1574-6941.2008.00552.x.

75. Guidot A, Coupat B, Fall S, Prior P, Bertolla F. Horizontal gene transfer between *Ralstonia solanacearum* strains detected by comparative genomic hybridization on microarrays. *The ISME Journal.* 2009;3(5): 549–562. https://doi.org/10.1038/ismej.2009.14.

76. Weisberg AJ, Wu Y, Chang JH, Lai EM, Kuo CH. Virulence and Ecology of Agrobacteria in the Context of Evolutionary Genomics. *Annual Review of Phytopathology.* 2023;61(1): 1–23. https://doi.org/10.1146/annurev-phyto-021622-125009.

77. Shin GY, Asselin JA, Smith A, Aegerter B, Coutinho T, Zhao M, et al. Plasmids encode and can mobilize onion pathogenicity in *Pantoea agglomerans. The ISME Journal.* 2025;19(1): wraf019. https://doi.org/10.1093/ismejo/wraf019.

78. Banerjee G, Chatterjee S, Banerjee P, Chattopadhyay P. Diversity of Secretion System Apparatus in Tomato Wilt Causing *Ralstonia solanacearum* Strains: a Comparative Analysis Using *in-silico* Approach. *bioRxiv.* 2022; https://doi.org/10.1101/2022.04.21.489029.

79. Ding S, Yu L, Lan G, Tang Y, Li Z, He Z, et al. Identification and genomic characterization of *Ralstonia pseudosolanacearum* strains isolated from pepino melon in China. *Physiological and Molecular Plant Pathology.* 2023;125: 101977. https://doi.org/10.1016/j.pmpp.2023.101977.

80. Ding S, Ma Z, Yu L, Lan G, Tang Y, Li Z, et al. Comparative genomics and host range analysis of four *Ralstonia pseudosolanacearum* strains isolated from sunflower reveals

genomic and phenotypic differences. *BMC Genomics*. 2024;25(1): 191.
https://doi.org/10.1186/s12864-024-10087-7.

81.   Salanoubat M, Genin S, Artiguenave F, Gouzy J, Mangenot S, Arlat M, et al. Genome
sequence of the plant pathogen *Ralstonia solanacearum*. *Nature*. 2002;415: 497–502.
https://doi.org/10.1038/415497a.

82.   Remenant B, Coupat-Goutaland B, Guidot A, Cellier G, Wicker E, Allen C, et al.
Genomes of three tomato pathogens within the *Ralstonia solanacearum* species
complex reveal significant evolutionary divergence. *BMC Genomics*. 2010;11: 379.
https://doi.org/10.1186/1471-2164-11-379.

83.   Morales VM, Sequeira L. Indigenous Plasmids in *Pseudomonas solanacearum*.
Phytopathology. 1985;75(7): 767–771. https://doi.org/10.1094/Phyto-75-767.

84.   Hamilton HL, Domínguez NM, Schwartz KJ, Hackett KT, Dillard JP. *Neisseria*
gonorrhoeae secretes chromosomal DNA via a novel type IV secretion system.
Molecular Microbiology. 2005;55(6): 1704–1721. https://doi.org/10.1111/j.1365-
2958.2005.04521.x.

85.   Callaghan MM, Heilers JH, Van Der Does C, Dillard JP. Secretion of Chromosomal
DNA by the *Neisseria gonorrhoeae* Type IV Secretion System. In: Backert S,
Grohmann E (eds) *Type IV Secretion in Gram-Negative and Gram-Positive Bacteria*.
Springer International Publishing; 2017. p. 323–345. https://doi.org/10.1007/978-3-319-
75241-9_13.

86.   Jurcisek JA, Brockman KL, Novotny LA, Goodman SD, Bakaletz LO. Nontypeable
Haemophilus influenzae releases DNA and DNABII proteins via a T4SS-like complex
and ComE of the type IV pilus machinery. *Proceedings of the National Academy of*
Sciences. 2017;114(32): E6632–E6641. https://doi.org/10.1073/pnas.1705508114.

87.   Goodman SD, Scocca JJ. Identification and arrangement of the DNA sequence
recognized in specific transformation of *Neisseria gonorrhoeae. Proceedings of the*
National Academy of Sciences. 1988;85(18): 6982–6986.
https://doi.org/10.1073/pnas.85.18.6982.

1192

1193 **Supplemental Figures**
1194 In addition to the four supplemental figures, a Zenodo repository contains supplemental
1195 tables and files like multiple sequence alignments, phylogenetic trees, and GenBank files
1196 that can be used for clinker synteny analyses (DOI: 10.5281/zenodo.14873519).
1197

1198
Figure S1. Distribution of T4SS gene clusters by T4SS class using F/P/I/GI-type classification
nomenclature. The majority of the T4SS gene clusters found in *Ralstonia* genomes are P-type.
Numbers indicate the number of T4SS gene clusters of that cluster/class in the RSSC pathogens vs.
non-RSSC environments. The counts of T4SS gene clusters are based on the number of *virB4* genes
in each category, including the 498 identified by the HMMER and BLASTp searches as well as the
five manually identified sequences.
1205

A) RSp0179 Synteny

B) RSp1521 Synteny

1206

Figure S2. Synteny analysis of the gene neighborhoods of two genes that are unlikely to be T4SS-
associated: **A)** RSp0179 and **B)** RSp1521. Both genes were consistently found on the megaplasmid.
Gene arrow colors are assigned by clinker and genes of the same exact color are homologous, other
than gray genes, which do not have homologs in this depiction. The linkages between gene arrows
are shaded based on global amino acid identity. The synteny analyses were performed using clinker.
Visualization adjustments were made using Affinity Designer.

1213

1214

Figure S3. Properties of the regions of transfer. **A)** Sizes of putative regions of transfer. The size in
kilobases is shown on the y-axis and the cluster is on the x-axis. Each point represents one region of
transfer. Circles indicate RSSC phytopathogens and triangles indicate non-RSSC environmentals. **B)**
GC content of putative regions of transfer. Left: the GC content of the whole genome on the x-axis
and the GC content of only the putative region of transfer on the y-axis. The dashed line represents a
one to one relationship between the GC content of the region and the GC content of the whole
genome. Each point represents one region of transfer in one genome. Circles indicate RSSC
phytopathogens and triangles indicate non-RSSC environmentals. Right: the GC content of the
putative region of transfer on the y-axis and the cluster on the x-axis. The dashed lines labeled
"Maximum (67%)" and "Minimum (63%)" represent the maximum and minimum GC content of the
whole genomes used in this analysis. Each point represents one region of transfer. Circles indicate
RSSC phytopathogens and triangles indicate non-RSSC environmentals.
1227

Figure S4. Example of cargo gene analysis using cluster a. The genes in T4SS cluster a are red, the heavy metal metabolism and resistance genes are green, and the other metabolism genes are brown. Genes that do not fall into the selected cargo categories are gray or white. Gray genes are those that have homologs among the genes analyzed. White genes are those that do not have homologs present in this analysis. The synteny analysis was performed using clinker. Gene colors were edited

1234   using Affinity Designer. Interactive HTML files for other putative region of transfer visualizations like
1235   this for the other clusters with complete genomes are available in Supplemental Files S27-S36.

Identified in:
● RSSC genomes
▲ Non-RSSC genomes

● ▲ Cluster i
n=75

▲ Cluster p
n=5

● Cluster h
n=4

● Cluster g
n=5

● ▲ Cluster c
n=49

▲ Cluster l
n=18

● Cluster d
n=2

● Cluster j
n=1

● ▲ Cluster b
n=7

▲ Cluster k
n=5

● ▲ Cluster a
n=160

Cluster f ● ▲
n=74

Cluster n ▲
n=34

Cluster m ▲
n=17

Cluster o ▲
n=1

Cluster e ● ▲
n=41

A) *virB4* Gene Presence in RSSC

B) *virB4* Gene Presence in Non-RSSC *Ralstonia*

C) *virB4* Gene Presence in RSSC and Non-RSSC

virB4 Gene Presence

virB4 Genes per Genome

A) Pairwise Cluster Co-Occurrence

RSSC Cluster

Cluster	a	i	f	b	c	d	h	g	e	j
a	5	10	8	1	1	1	1	0	0	0
i		1	8	0	0	0	0	0	0	0
f			0	0	0	0	0	0	0	0
b				0	0	0	0	0	0	0
c					0	0	0	0	0	0
d						0	0	0	0	0
h							0	0	2	0
g								0	2	0
e									0	0
j										0

Non-RSSC Cluster

Cluster	a	f	n	e	c	l	m	p	k	b	i	o
a	24	25	19	15	11	11		0	2	2	1	0	0
f		20	21	20	13	9	3	2	1	1	0	0
n			15	7	12	2	1	3	0	0	0	0
e				5	13	4	0	3	1	0	0	0
c					1	1	1	0	1	0	0	0
l						1	0	0	0	0	0	0
m							0	1	1	0	0	0
p								0	0	0	0	0
k									0	0	0	0
b										0	0	0
i											0	0
o												0

B) T4SS Cluster Locations in Complete Genomes

Chromosome
Megaplasmid

Accessory Plasmid

Cluster c/f n=1
Cluster d n=1
Cluster e n=6
Cluster f n=4
Cluster g n=2
Cluster i n=7
Cluster l n=1
Cluster m n=1
Cluster n n=4

Integrative and Conjugative Element (ICE)

Chromosome:
Cluster a n=24
Cluster c n=9
Cluster f n=1

Megaplasmid:
Cluster a n=2
Cluster c n=2