Time in the city: Exploration and problem-solving behaviours reveal long-term urban

adaptation in wild red foxes (*Vulpes vulpes*)

F. Blake Morton*[1], Dylan Thompson-Jones[1], Kristy A. Adaway[1], Koko M. Sutter[1], Catia Matos[2],

Georgia B. Freer[3], Georgie Fletcher[3], Carl D. Soulsbury[3]

[1] Department of Psychology, University of Hull, Hull, UK

[2] School of Environmental Sciences, University of Hull, Hull, UK

[3] School of Natural Sciences, University of Lincoln, Lincoln, UK

Author Note

F. Blake Morton 🆔 https://orcid.org/0000-0002-8644-1557

Dylan Thompson-Jones 🆔 https://orcid.org/0009-0005-0460-4033

Kristy A. Adaway 🆔 https://orcid.org/0009-0006-1041-8592

Koko M. Sutter 🆔 https://orcid.org/0009-0003-7565-3663

Catia Matos 🆔 https://orcid.org/0000-0001-8452-5629

Georgia B. Freer 🆔 https://orcid.org/0009-0009-5896-0956

Carl D. Soulsbury 🆔 https://orcid.org/0000-0001-8808-5210

*Author for correspondence: Dr Blake Morton, Department of Psychology, University of Hull,

Hull, HU6 7RX, United Kingdom. E-mail: b.morton@hull.ac.uk

Abstract

Urbanisation is one of the most important forms of human-driven landscape change,

creating novel situations in which some species thrive. Animals' willingness to touch, explore,

and solve novel challenges, such as exploiting unfamiliar resources, are thought to be key

behavioural characteristics for urban adaptation. Little is known about the impact of

spatiotemporal patterns of urbanisation, particularly historical patterns of change, on these

behaviours. We tested this in the world's most urbanised carnivore, the red fox, using novel

tasks deployed across 284 sites in the United Kingdom. We compared tactile and problem-

solving behaviours in rural populations, recently colonised urban populations, and long-

established urban populations (>40 years). Similar to previous work, urban foxes were more

likely to touch the tasks compared to rural populations; however, in our new extended dataset,

we show for the first time that older urban populations – particularly from London – display

greater exploratory diversity and higher problem-solving success, despite more recent urban

populations being equally likely to touch them. Our findings suggest that certain population

characteristics, such as the likelihood of touching/engaging with novelty, emerge early in

urbanisation while other characteristics, such as exploratory diversity and behavioural

innovation, may emerge after long-term urban exposure across many decades.

Key words: rapid environmental changes, behavioural innovation, boldness, biodiversity

crisis, urbanisation, problem-solving ability

Introduction

Earth is presently facing a catastrophic decline in biodiversity, often termed the sixth

mass extinction or biodiversity crisis (1, 2). This decline is largely the result of human-driven

environmental changes (hereafter "environmental changes") (3). One of the most important

forms of environmental change on the planet is urbanisation (4), defined as a dynamic process

characterised by expanding human settlements, such as increased infrastructure (e.g.,

housing and building densities), human population growth, and reduced habitat heterogeneity

(5, 6). By 2050, over 68% of the world human population is expected to live within cities,

compared to only 30% in 1950 (4). Such rapid changes expose wildlife to a wide variety of

novel situations, such as new or modified habitats (7), anthropogenic resources (8),

competitive interactions (9), and potential predators and diseases (10, 11). Most species are

experiencing these changes for the first time in their evolutionary history; meaning, evolution

has not necessarily equipped them to withstand such unprecedented conditions (12-14). While

many species are at risk of decline or extinction due to slow adaptation, others are

unexpectedly thriving for reasons that are still unclear (12, 15). Thus, there is a pressing need

to better understand what factors help wildlife respond and adapt to environmental changes,

particularly those related to urbanisation.

Animals' initial responses to environmental changes are often the result of behavioural

plasticity, not rapid genetic evolution (16). Several behavioural phenotypes have been

proposed as being important for coping with novel situations arising from environmental

changes. One is bold behaviour, broadly defined as an animal's willingness to engage with

novelty (17, 18). Another behaviour is exploration, which reflects an animal's tendency to

actively investigate situations within its environments (19). Unlike boldness, exploration does

not always necessarily involve risk-taking and, instead, can be driven by other factors, such

as curiosity or persistence (20, 21). Finally, a third behaviour is innovation, defined as using a

new or modified behaviour to solve a new or familiar task (22, 23). Bolder animals are more

likely to touch and engage with unfamiliar, and therefore potentially risky, resources compared

to other organisms (24-26). Animals with greater exploratory and innovative tendencies are

better equipped to gain information and discover solutions to novel challenges (19, 21, 23,

27). Thus, greater environmental changes are often hypothesised to favour bolder, more

exploratory and innovative behaviours (14, 28), potentially shaping the likelihood of these

behaviours being observed within a population (29). However, the likelihood of animals

displaying such behaviours is not always related to the magnitude of environmental change

(14, 26, 28, 29). In terms of urban-rural gradients, for example, many studies find that urban

populations are more likely to display bolder and more innovative behaviour than rural

populations, while other studies show that rural populations can be more, or equally, as bold

and innovative as their urban counterparts (18, 26, 28-30). Thus, the relationship between

environmental changes and how animals respond to novelty is complex and warrants further

research.

One underexplored factor is the long-term historical patterning of environmental

changes, such as the rate at which such changes occur over decades. Most studies examining

the impact of environmental changes on animals' behavioural responses to novelty are based

on static, cross-sectional ecological gradients from the present day (14, 26, 28, 29). While

such analyses are important for understanding how different degrees or "magnitudes" of

change might impact animal populations (e.g., a steeper gradient suggests greater impact),

behavioural characteristics of present-day populations are often a reflection of their ecological

past, and so cross-sectional analyses cannot account for historical patterning. If urbanisation,

for example, follows a gradual and contiguous pattern, populations may have longer to adapt

to human pressures, leading to incremental behavioural changes across many generations.

By contrast, if urbanisation is abrupt and rapid, such as sudden expansion or land-use

changes, it may trigger an abrupt shift in population-level behaviour, favouring animals that

quickly learn or already possess these behavioural characteristics (12, 26, 31). Thus, to better

understand how wildlife might adapt to environmental changes, there is a pressing need to

understand whether (and how) historical patterns of environmental change predict the

likelihood of populations displaying bold, exploratory, and innovative behaviours.

Red foxes, *Vulpes vulpes*, offer a powerful model for understanding how historical

patterns of urbanisation influence present-day behavioural adaptation. As the most globally

widespread terrestrial carnivore on the planet (32), foxes were among the first mammals on

record to have expanded their ecological niche to urban areas (33). Reports of urban foxes

have existed in British cities at least since the 1930s (33). In more recent decades, foxes have

established themselves in other cities throughout Europe, Asia, Australia, and North America

(8, 34-36). Although urbanisation is a relatively recent phenomenon in this species'

evolutionary history, notable behavioural and morphological differences are already being

documented in urban fox populations compared to the countryside. Compared to their rural

counterparts (29, 36), for instance, urban foxes often have smaller dietary repertoires (37),

consume anthropogenic food more frequently (8, 37, 38), occupy smaller home ranges (39,

40) and display bolder behaviours. There is also some evidence that urban foxes are evolving

to have smaller skulls (41) which, taken together, suggests that urbanisation may lead to

microevolution of behaviours and other phenotypes, including morphology. The colonization

of urban areas by foxes, however, has been an uneven process (42). If microevolution or other

long-term processes are occurring, we should expect there to be localized variation in the

speed and timing of behavioural changes within the species – a possibility that has not yet

been formally tested.

In the current study, we investigated whether long-term historical patterns of

urbanisation predict population differences in bold, exploratory, and innovative behaviours in

present-day foxes from the United Kingdom (UK). Specifically, we tested whether present-day

urban fox populations differ from rural populations in these behaviours, and whether such

differences are predicted by historical spatio-temporal patterns of urbanisation over the last

several decades. We also tested whether populations from relatively older, more established

urban areas (before 1986) are more likely to display bolder, more exploratory and innovative

behaviours compared to more recently urbanized areas. By integrating behavioural data with

historical patterns of urban development, we aim to clarify how past environmental changes

potentially shape present-day behavioural strategies, offering new insights into the adaptive

potential of wildlife's exposure to urbanisation.

Methods

Study Sites and Subjects

We studied 284 locations throughout Great Britain (Figure 1a), including areas within

and around many towns and cities (e.g. London, Glasgow, Edinburgh, Bristol, Southampton,

and York). Landscapes for these locations were diverse, including public parks, residential

gardens, monostand pine/spruce plantations, open fields (including farms), mixed woodland,

and coastal and mountainous scrubland. Foxes were not tagged and their participation in our

behavioural tests was entirely voluntary. We acquired permission to visit 242 of these locations

by contacting city councils and other organizations that owned the land. The remaining 42

locations were accessed by advertising the study through social media and regional wildlife

groups; these locations were primarily comprised of golf courses and residential gardens. All

locations were spaced at least 3.5 km away from any of our other study locations, which

reduced the likelihood of acquiring data on the same foxes visiting multiple locations since $\geq$

3.5 km is larger than the typical dispersal distance and home range diameter of foxes within

the UK (43, 44).

Measuring population-level differences in bold, exploratory, and innovative behaviours in

foxes

We used methods already developed and validated to assess population-level

differences in foxes' behavioural responses to novel tasks, the details of which can be found

in Morton et al. (29). Data from 200 of our locations come from our original Morton et al. (29)

study, collected between August 2021 and November 2022. Data from an additional 84

locations were collected specifically for the current analysis, between March 2023 and May

2024.

Briefly, a single novel food-related task was selected from a battery of eight designs

(Figure S1) and deployed at a given location for 15.6 ± 1.8 days before it was retrieved by

researchers. Each object varied in terms of design and was made from basic household

materials (e.g., PVC piping, metal screws and wooden rods). Tasks were counterbalanced

across locations to ensure that our findings were not unique to any particular type of task and,

instead, were more generalisable. Our previous work found that, for these specific task

designs, population differences in fox behaviour were not significantly influenced by the type

of design deployed at a location (29). Some tasks were sprayed with scent deodorizer to

understand whether the scent of objects (e.g. human odour) influenced fox behaviour which,

in our previous study (29), was also shown to have no significant effect.

All tasks had bait scattered around them to attract foxes to those locations. The objects

also contained a food reward inside them, which could be obtained by foxes performing natural

behaviours (e.g., lifting, biting, pushing, or pulling materials). Examples of these behaviours

can be found here: https://youtu.be/SYHyjXPdcZs (29). The tasks were designed to

understand animals' *willingness* to engage with novelty, and so in this context, solving a task

meant that animals needed to be bold (and otherwise motivated) enough to physically touch

and explore it during the two-week period in which it was available. Stemming from this, many

factors can potentially underpin animals' willingness to display bold, exploratory, and

innovative behaviours (23, 45-47). For example, animals that are too afraid, not hungry, or not

persistent may be less likely to exploit novel food-related objects. Importantly, however, the

purpose of our study was to determine *whether*, not why, our study species would display

bold, exploratory, and innovative behaviours, and so the only way they could do this was by

physically engaging with the tasks and displaying those behaviours.

Fox visitations occurred when researchers were not present, and 'no glow' (940 nm)

infrared motion-sensor cameras (Apeman H45) were used to record fox behaviours. Across

all locations, we measured whether a fox was detected on camera (1=yes, 0=no) as well as

the total number of visits made by foxes during the period in which a task was deployed at that

location. To reduce temporal autocorrelation between successful visits, we considered a "visit"

to have ended if a fox was absent from the camera's field of view for $\geq$10 minutes. The use of

this time interval to measure independent red fox detections has been validated in a previous

trail camera study (48). Because foxes were free ranging and their participation was entirely

voluntary, we based our analysis on foxes that were at least able and willing to visit the

locations, which still enabled us to assess urban-rural differences in foxes' behavioural

responses to the novel tasks (29).

We only coded behaviours from locations where foxes clearly acknowledged the task

during their visit, which we defined in terms of the fox turning its head to look/smell in the

object's direction. As in our previous study (29), we measured the likelihood of behavioural

innovation occurring within fox populations in terms of the number of locations in which we

saw (1=yes, 0=no) a fox displaying a behaviour at any point that led it to operate and

successfully gain access to the food inside the novel task within the timeframe in which the

task was available at that location. Also, we were not interested in trait boldness in the current

study, which describes stable individual differences in behaviour across a wide variety of

contexts (17). Instead, as with Morton et al. (29) and other studies of free-ranging animals (18,

20, 47), we were interested in the likelihood of fox populations displaying outwardly bold

behaviour, which we defined in terms of a fox physically touching (yes or no) the object at any

point within the timeframe in which the task was available at that location (e.g., pushing,

pulling, licking, or biting it).

Of the foxes that were bold enough to make physical contact with the novel tasks, we

also quantified four additional behavioural measures. First, latency to first touch was defined

as the time from the first acknowledgement of a task by any fox at a location until the first

instance of physical contact by a fox from that location. Second, we measured the total time

foxes spent engaged in tactile exploration of the task at a location, which captured foxes'

sustained interactions with the task within a given area. Third, we recorded exploration

diversity (21), defined here as the total number of distinct exploratory behaviours directed at

the task at any point by foxes during the period in which it was deployed at a location (Table

1). Finally, we measured latency to solve, which we defined in terms of the amount of time

elapsed between the first time a fox touched the object until the time when a fox ultimately

solved it by gaining access to the food inside.

Table 1. Ethogram of behaviours used by foxes to physically explore novel food-related tasks.

Behaviour	Definition
Nip	Quick contact with object using teeth, with a fast recoil. Quickly moving body or head away from object after making contact.
Bite	Longer contact with teeth compared to a 'nip'. Not moving position of the mouth, staying stationary. Each close of the jaw counts as a new bite.
Teeth pull	Grabbing the object with teeth and trying to pull it towards their body. Each close of the jaw counts as new pull.
Rostrum touch	Contact with the front of face/nose, which does not include teeth, such as pushing snout into or towards the object, or licking it.
Left paw contact	Using left paw to make contact with the object other than standing on the object.
Right paw contact	Using right paw to make contact with the object other than standing on the object.
Combined paw scratch	Quick contact with the object using both paws at the same time. One paw may make contact first, but both must be touching at some point during the behaviour for combined contact.

Measuring rates of environmental change due to urbanisation

We obtained spatial data on landscape changes typically associated with urbanisation

at each of our study locations from 1994 to 2020, which was the maximum consistent

timeframe available. These landscape changes were based on predefined categories from

ESA Land Cover Time Series with a resolution of 300 metres (49). Buffers of 3,500 metres

were created around each location to analyse the proportion of urban land cover present; this

buffer distance corresponded to the typical annual home range of male and female foxes in

the UK (43, 44). For each buffer, we calculated the proportion of urban habitat, allowing us to

quantify urbanisation in 1994, 2020, and the amount of change between these two time points.

Raster data processing was performed with ArcGIS Pro (ESRI), and landscape cover and

change analysis were performed with R version 4.4.2 (50).

Measuring length of exposure to urbanisation

The exact timing of urban foxes colonising different cities is uncertain. We do, however,

have a snapshot from a national survey of 378 local UK authorities from 1986 which

documented urban fox population status as being either present, absent, or uncertain (51).

We therefore cross-referenced data on fox presence/absence from our current study with

records from 1986; two locations from 1986 were uncertain, both in Hamilton, Scotland, and

were excluded, resulting in 84 locations for our analysis. Locations where urban foxes were

already present in 1986 were defined as relatively "older" urban populations, whereas

locations where foxes were absent in 1986 (but present in the current study) were defined as

relatively "newer" urban populations.

Statistical Analyses

We used Cohen's kappa tests and intra-class correlation coefficients (ICC 3,1) to

evaluate interobserver agreement for all fox behaviours coded from videos. There was

excellent agreement across all coders (Tables S1-S8).

We first identified clusters of high or low values and spatial outliers of foxes touching

and then solving the novel tasks using the Local Moran's I statistic, i.e., a local indicator of

spatial association (LISA), using the R package *spdep* (52). To ensure coverage of the UK,

we created a neighbour list using a distance threshold of 35 km to identify spatial relationships

between points. The results were visualized using Moran's scatterplots and maps, highlighting

areas of significant spatial autocorrelation, with permutation tests providing p-values to assess

the statistical significance of the observed spatial patterns either as areas of high-high (hot

spots) or low-low (cold spots) of behaviours.

We then tested whether foxes were more likely to display our target behaviours

depending on the length of time in which the trail camera was operational (29); however, we

found no significant effect (Table S9), and so we excluded camera operation time as a variable

in our main analysis to avoid model oversaturation. We then tested whether (a) touching

(yes/no) or (b) solving the task (yes/no) was associated with current (2020) urbanisation data

using binomial general linear models (GLMs). We also used poisson GLMs to examine the

effects of urbanisation on the number of visits and exploratory diversity of foxes at each

location, followed by GLMs to assess how urbanisation predicted the time taken to touch or

solve the task.

We tested whether change in the environment was a key driver of behaviour using

binomial GLMs, comparing the (a) probability of touching and (b) solving the tasks in relation

to the increase in urbanisation between 1990 (earliest available data) and 2020. Finally, we

compared the likelihood of older versus newer urban fox populations (a) touching and (b)

solving the tasks using binomial GLMs, as well as their (c) latency to solve the tasks using

GLMs and (d) number of exploratory behaviours using poisson GLMs.

Results

Spatial patterns of fox behaviour

Foxes touched the novel tasks at 27.4% of locations (34/124; Figure 1b) and solved

them at 12.4% of locations (14/122; Figure 1c). Foxes mainly solved tasks in urban areas,

including Leeds, Edinburgh, Sheffield and London (Figure 1c). The local indicators of spatial

association (LISA) identified multiple hotspots for foxes acknowledging the tasks (Figure S2),

but only indicated a cluster of 13 significant hot spots for both touching and solving them – all

in and around London (Figure 1d). Similar to our previous work, the likelihood of touching a

task was positively associated with urbanisation (Likelihood Ratio (LR) χ^2=11.05, p<0.001;

Figure 2a). However, in our extended dataset, there was also a significant positive association

between urbanisation and the likelihood of foxes solving the task (LR χ^2=15.6-, p<0.001;

Figure 2b).

Behavioural differences between urban and rural fox populations

Number of fox visits to tasks was positively related to urbanisation (LR χ^2=1205.5,

p<0.001; Fig S3). For locations where foxes touched a task, there was no effect of urbanisation

on the latency to touch (LR χ^2=2.03, p=0.153) or solve (LR χ^2=0.47, p=0.491) the tasks. Urban

foxes did not display more exploratory behaviours when interacting with the tasks (LR χ^2=0.03,

p=0.866), nor did they spend any longer exploring the tasks (LR χ^2=0.12, p=0.732), compared

to rural populations.

Temporal drivers of fox behaviour

The majority of locations (80.2%, 228/284) showed some increase in urbanisation

between 1994 and 2020. However, most of these locations had relatively small changes in the

amount of urbanisation across this timeframe (mean±SE=0.008±0.004), with only 12 locations

showing an increase in urbanisation by >2%. Areas with the greatest increases in urbanisation

between 1994 to 2020 did not have a higher likelihood of a fox touching (LR χ^2=2.82, p=0.093)

or solving (LR χ^2=0.01, p=0.976) the tasks.

When comparing relatively older versus newer urban fox populations, there was no

difference in the likelihood of foxes touching the tasks (LR χ^2=0.90, p=0.341), but older

population were significantly more likely to solve them (LR χ^2=8.11, p=0.004). In total, 26.1%

(12/46) of foxes from older urban populations solved the tasks compared with 0% (0/16) of

foxes from newer populations. Older urban populations also displayed significantly more

exploratory behaviours when interacting with tasks (LR χ^2=9.08, p=0.002; Figure 3). There

was only a tendency for older urban populations to touch the tasks faster (LR χ^2=2.06,

p=0.154) and spend longer exploring them (LR χ^2=3.03, p=0.081). Removing locations in

London (identified as a major hotspot) yielded similar results across all analyses (touch: LR

χ^2=0.27, p=0.605; solve: LR χ^2=4.22, p=0.039), indicating that behavioural differences

between old versus new urban populations are not solely driven by factors unique to London

and, instead, more likely reflect broader patterns of urbanisation among our study sites.

Discussion

Consistent with our previous findings (29), the current study found that urban foxes

were more likely to touch novel food-related tasks. We also extend this work by showing, for

the first time, that certain urban fox populations are more likely to solve novel tasks using

innovative behaviour, and that older, more established urban populations display greater

exploratory diversity and problem-solving success compared to more recent urban

populations. Collectively, these new findings suggest that historical patterns of urbanisation –

specifically, where and how long a city has been inhabited by foxes – may contribute to

population differences in bold, exploratory, and innovative behaviours.

Although behavioural innovation was observed in multiple urban locations, London

emerged as a particular hotspot, likely reflecting its status as the oldest urban fox population

in the UK (34). This population has previously shown distinct phenotypic changes in skull

morphology (41), suggesting that behavioural and morphological phenotypes may be

influenced by long-term exposure to urban environments. By contrast, foxes from Bristol,

which are another relatively old urban population, did not display bold, exploratory, or

innovative behaviours. Historical declines from sarcoptic mange in Bristol during the 1990's

(53), followed by recolonisation from surrounding rural areas, may have introduced a genetic

pool that is less-shaped by urban selective pressures, reducing the likelihood of such

behaviours (54). If so, then local population history could play a potential role in shaping urban

adaptation, including the age of colonisation, demographic events, and recolonisation

dynamics. Such ecological factors might also help explain why urban-rural differences in

behaviour are not always consistent between studies (26, 28).

Urban foxes were first reported in UK cities in the 1930s (34), with 33.1% (125/378) of

surveyed urban areas hosting populations by the mid-1980s (51). Colonisation has continued,

meaning that some of our study populations are older than others (51, 55). A key finding from

our study is that the rate of urbanisation between 1994 and 2020 did not appear to influence

urban-rural differences in our target behaviours, which may reflect the modest changes to

urban areas in the UK during this period. Alternatively, this result could reflect the spatial

resolution of available landscape datasets, which may not have detected more subtle

environmental changes, such as fine-scale shifts in human activity. Regardless, exploratory

diversity and problem-solving success were more common in older, more established urban

fox populations, even though newer populations (established after 1986 or <40 years ago)

were equally likely to touch the tasks. Since foxes first colonised UK cities ~90 years ago (34),

these findings suggest that certain behaviours, such as the likelihood of touching/engaging

with novelty, may emerge relatively early (<40 years), whereas other behaviours, including

exploratory diversity and innovative behaviour, may require 40-90 years of urban exposure. In

other words, the behavioural characteristics of older urban fox populations suggest a more

gradual pattern of adaptation rather than more recent rates of urbanisation, potentially via

individual learning, cultural transmission, dispersal patterns, or a combination of these and

other mechanisms across many generations. Newer urban populations may not yet exhibit

these behaviours to the same extent, being in earlier stages of adjustment or colonisation.

Adaptation to urban environments, including the micro-evolutionary or other long-term

processes that drive it, may take time to emerge in at least some, but not necessarily all,

contexts. Indeed, evidence from our study supports this notion. In other species, older urban

coot populations (*Fulica atra*) are bolder (56) while older commensal house mice populations

are better problem-solvers (57) compared to younger urban populations. By contrast,

Johnson-Ulrich et al. (58) found that rural hyenas (*Crocuta Crocuta*) were more innovative

than relatively newer and older urban populations. Whether such differences are explained by

genetic or non-genetic factors is not clear (59), highlighting the need for longitudinal research

to better understand urban-rural divergence in behaviour.

Building on our proposed temporal framework for behavioural adaptation in foxes,

studies in other species show that bold and innovative behaviours sometimes co-occur within

the same population, with novelty avoidance potentially limiting innovation (19, 60). In our

study, innovation required foxes to display sufficient boldness to interact with the tasks, but

not all bold fox populations displayed innovative behaviour (e.g., 59% of tasks that were

touched were not solved). Although further research is needed, such patterns could be

explained by the notion of a temporal sequence, where innovation emerges over time after

urban fox populations are sufficiently bold to engage with novelty (61). Because our study

focused on population-level behavioural patterns, future research should test whether bold

and innovative behaviours co-occur within the same individuals, as suggested in some other

species (60).

Urban areas typically have higher fox densities than rural settings (34), and in the

current study, urban foxes were also more likely to visit the tasks. This raises the possibility

that differences in density or visitation could influence the likelihood of observing our target

behaviours. However, several lines of evidence suggest that these factors alone cannot

account for our results. First, among locations where foxes interacted with tasks, urban and

rural populations did not differ in their latency to touch or solve the tasks, nor did they differ in

terms of the number of exploratory behaviours or the time spent engaging with tasks. Second,

spatial hotspots for touching and solving were concentrated in specific areas (e.g., London),

rather than being more uniform across all high-density urban fox sites (e.g., Glasgow). Third,

the overall likelihood of problem-solving was relatively low across all urban and rural fox

populations (12.4% of locations), suggesting that sheer fox numbers or visits cannot fully

explain the observed behavioural differences. This suggests that other factors associated with

long-term urban exposure may provide a more parsimonious explanation for these findings.

For instance, urban environments, particularly older cities, likely expose foxes to greater

diversity of human-made objects, such as discarded litter (62), providing repeated

opportunities to discover, explore, and learn how to exploit novel resources (63). Alternatively,

novel food-related objects, including those used in our study, may represent relatively easy-

to-access resources in urban habitats, where high fox densities and smaller home ranges

intensify competition and increase the attractiveness of such resources. Further research is

needed to explore these and other possibilities.

Conclusions

Overall, our study demonstrates that historical patterns of urbanisation, particularly the

length of time populations have been established in urban environments, are an important

predictor of population differences in bold, exploratory, and innovative behaviours in wild red

foxes. While initial engagement with novel food-related opportunities, such as our novel tasks,

may emerge relatively quickly in newly established urban populations, more complex

behaviours – such as diverse exploration and problem-solving – were predominantly observed

in older urban populations (particularly London), highlighting the potential role of local variation

and long-term exposure to urbanisation in shaping population behaviour. By linking long-term

urbanisation history to present-day behavioural traits, our findings provide new insights into

the temporal and spatial processes underlying wildlife adaptation to human-altered

environments.

Ethical Note

This research was granted ethical approval by the Animal Welfare Ethics Board of the

University of Hull (FHS356) and followed all ASAB/ABS guidelines. Foxes were not handled,

trail cameras were placed away from footpaths to reduce public disturbance, and all food items

used to attract foxes were safe for ingestion by other species.

Data Availability

All data are provided in data set S1 in the Supplementary material.

Declaration of Interest

The authors declare no conflict of interest.

Acknowledgments

We thank everyone involved with the *British Carnivore Project*, including the researchers, students, and citizen scientists. We are grateful to the many organizations that helped us acquire landowner permissions for the study, particularly staff from the Wildlife Trusts, Scottish Wildlife Trust, The Land Trust, the National Trust, Forestry England (particularly C. Bashforth), Forestry & Land Scotland, Yorkshire Water, and numerous city and regional councils (particularly Hull, East Riding of Yorkshire, and Glasgow). We thank Yacob Haddou (University of Glasgow) for helping us obtain the landscape measures for our study sites. K.A.A. and D.T-J. were both supported at the time of writing by scholarships from the Doctoral College at the University of Hull. K.M.S. was supported at the time of writing by the Leverhulme Doctoral Scholarships Centre for Water Cultures. G.B.F. was supported by a research grant (awarded to C.D.S. and F.B.M.) from the Wild Animal Initiative. F.B.M. also thanks the University of Hull and UKRI Natural Environment Research Council (NERC; Grant No. NE/X018342/1) for funding, and the EU Social Plus programme for providing funding to the University of Hull, which supported three of our research assistants.

References

1.   Ceballos G, Ehrlich PR. Biological annihilation via the ongoing sixth mass extinction signaled by vertebrate population losses and declines Proceedings of the National Academy of Sciences of the United States of America. 2017;114:E6089-E96.
2.   Ceballos G, Ehrlich PR, Barnosky AD, Garcia A, Pringle RM, Palmer TM. Accelerated modern human-induced species losses: Entering the sixth mass extinction. Science Advances. 2015;1:e1400253.
3.   Young HS, McCauley DJ, Galetti M, Dirzo R. Patterns, causes, and consequences of anthropocene defaunation Annual Review of Ecology, Evolution, and Systematics. 2016;47:333-58.
4.   United_Nations_Population_Division. World urbanization prospects 2018: Highlights (ST/ESA/SER.A/421). United Nations, Department of Economic and Social Affairs, Population Division; 2019.
5.   Grimm NB, Faeth SH, Golubiewski NE, Redman CL, Wu J, Bai X, et al. Global change and the ecology of cities. Science 2008;319:756-60.
6.   Angel S, Parent J, Civco DL, Blei AM. Making room for a planet of cities: Lincoln Institute of Land Policy. ; 2011.
7.   Šálek M, Drahníková L, Tkadlec E. Changes in home range sizes and population densities of carnivore species along the natural to urban habitat gradient. Mammal Review. 2015;45:1-14.

8. Contesse P, Hegglin D, Gloor S, Bontadina F, Deplazes P. The diet of urban foxes
(Vulpes vulpes) and the availability of anthropogenic food in the city of Zurich, Switzerland.
Mammalian Biology. 2004;2:81-95.
9. Martin PR, Bonier F. Species interactions limit the occurrence of urban-adapted birds
in cities. Proceedings of the National Academy of Sciences of the United States of America.
2018;115:E11495-E504.
10. Guiden P, Bartel SL, Byer NW, Shipley AA, Orrock JL. Predator-prey interactions in
the anthropocene: Reconciling multiple aspects of novelty. Ecology and Evolution.
2019;34:P616-27.
11. Pedroso-Santos F, Costa-Campos CE. Novel predator-prey interactions of Rhinella
major (Anura: Bufonidae) from an urban area in northern Brazil. Phyllomedusa: Journal of
Herpetology. 2020;19:287-92.
12. Sih A, Ferrari MCO, Harris DJ. Evolution and behavioural responses to human-
induced rapid environmental change. Evolutionary Applications. 2011;4:367-87.
13. Garant D. Natural and human-induced environmental changes and their effects on
adaptive potential of wild animal populations. Evolutionary Applications. 2020;13:1117-27.
14. Gunn RL, Hartley IR, Algar AC, Niemela PT, Keith SA. Understanding behavioural
responses to human-induced rapid environmental change: a meta-analysis. Oikos.
2022;2022:e08366.
15. Finn C, Grattarola F, Pincheira-Donoso D. More losers than winners: investigating
anthropocene defaunation through the diversity of population trends Biological Reviews.
2023;98:1732-48.
16. Hendry AP, Farrugia TJ, Kinnison MT. Human influences on rates of phenotypic
change in wild animal populations. Molecular Ecology. 2008;17:20-9.
17. Bergvall UA, Schapers A, Kjellander P, Weiss A. Personality and foraging decisions
in fallow deer, *Dama dama*. Animal Behaviour. 2011;81(1):101-12.
18. Breck SW, Poessel SA, Mahoney P, Young JK. The intrepid urban coyote: a
comparison of bold and exploratory behavior in coyotes from urban and rural environments.
Scientific Reports. 2019;9:2104.
19. Sol D, Griffin AS, Bartomeus I, Boyce H. Exploring or avoiding novel food resources?
The novelty conflict in an invasive bird. PLoS ONE. 2011;6(5):e19535.
20. Adaway KA, Snider MH, Vinciguerra C, Kupferman CA, Beasley JC, Morton FB. Wild
raccoons are more innovative but not bolder than another generalist, the Virginia opossum,
on a pull-string task. Animal Behaviour. 2025;227:123279.
21. Daniels SE, Fanelli RE, Gilbert A, Benson-Amram S. Behavioral flexibility of a
generalist carnivore. Animal Cognition. 2019;22:387-96.
22. Lee. Adaptations to environmental change: an evolutionary perspective. In: Box HO,
editor. Primate responses to environmental change. Dordrecht: Springer; 1991. p. 39-56.
23. Reader SM, Laland KN. Animal innovation. Oxford ; New York: Oxford University
Press; 2003. x, 344 p. p.
24. Forss SIF, Schuppli C, Haiden D, Zweifel N, van Schaik CP. Contrasting responses
to novelty by wild and captive orangutans. American Journal of Primatology.
2015;77(10):1109-21.
25. Webster MM, Ward AJW, Hart PJB. Individual boldness affects interspecific
interactions in sticklebacks. Behavioral Ecology and Sociobiology. 2009;63:511-20.
26. Griffin AS, Netto K, Peneaux C. Neophilia, innovation and learning in an urbanised
world: a critical evaluation of mixed findings. Current Opinion in Behavoral Sciences.
2017;16:15-22.
27. Wild S, Krutzen M, Rankin RW, Hoppitt WJE, Gerber L, Allen SJ. Long-term decline
in survival and reproduction of dolphins following a marine heatwave. Current Biology.
2019;29:R239-R40.
28. Vincze E, Kovacs B. Urbanisation's effects on problem solving abilities: A meta-
analysis. Frontiers in Ecology and Evolution. 2022;10:834436.

29.   Morton FB, Gartner M, Norrie E-M, Haddou Y, Soulsbury CD, Adaway KA. Urban
foxes are bolder but not more innovative than their rural conspecifics Animal Behaviour.
2023;203:101-13.
30.   Lazzaroni M, Brogi R, Napolitano V, Apollonio M, Range F, Marshall-Pescini S.
Urbanisation does not affect red foxes' interest in anthropogenic food, but increases their
initial cautiousness. Current Zoology. 2024;70:394-405.
31.   Uchida K, Suzuki K, Shimamoto T, Yanagawa H, Koizumi I. Seasonal variation of
flight initiation distance in Eurasian red squirrels in urban versus rural habitat. Journal of
Zoology. 2016;298:225-31.
32.   Marsh CJ, Sica YV, Burgin CJ, Dorman WA, Anderson RC, Del Toro Mijares I, et al.
Expert range maps of global mammal distributions harmonised to three taxonomic
authorities. J Biogeogr. 2022;49(5):979-92.
33.   Harris S, Baker P. Urban foxes: New Edition; 2001.
34.   Soulsbury CD, Baker PJ, Iossa G, Harris S. Red foxes (Vulpes vulpes). In: Gehrt SD,
Riley SPD, Cypher BL, editors. Urban carnivores: Ecology, conflict, and conservation: John
Hopkins University Press; 2010. p. 63-75.
35.   Plumer L, Davison J, DSaarma U. Rapid urbanization of red foxes in Estonia:
distribution, behaviour, attacks on domestic animals, and health-risks related to zoonotic
diseases. PLoS ONE. 2014;9:e115124.
36.   Gil-Fernandez M, Harcourt R, Newsome T, Towerton A, Carthey A. Adaptations of
the red fox (Vulpes vulpes) to urban environments in Sydney, Australia. Journal of Urban
Ecology. 2020;6:1-9.
37.   Scholz C, Firozpoor J, Kramer-Schadt S, Gras P, Schulze C, Kimmig SE, et al.
Individual dietary specialisation in a generalist predator: A stable isotope analysis of urban
and rural red foxes. Ecology and Evolution. 2020;10:8855-70.
38.   Fletcher JWJ, Tollington S, Cox R, Tolhurst BA, Newton J, McGill RAR, et al.
Utilisation of anthropogenic food by red foxes (Vulpes vulpes) in Britain as determined by
stable isotope analysis Ecology and Evolution. 2025;15:e70844.
39.   Kobryn HT, Swinhoe EJ, Bateman PW, Adams PJ, Shephard JM, Fleming PA. Foxes
at your front door? Habitat selection and home range estimation of suburban red foxes
(Vulpes vulpes). Urban Ecosystems 2022.
40.   Main MT, Davis RA, Blake D, Mills H, Doherty TS. Human impact overrides
bioclimatic drivers of red fox home range size globally. Diversity and Distributions.
2020;26:1083-92.
41.   Parsons KJ, Rigg A, Conith AJ, Kitchener AC, Harris S, Zhu H. Skull morphology
diverges between urban and rural populations of red foxes mirroring patterns of
domestication and macroevolution Proceedings of the Royal Society B: Biological Sciences.
2020;287:20200763.
42.   Scott DM, Berg MJ, Tolhurst BA, Chauvenet ALM, Smith GC, Neaves K, et al.
Changes in the distribution of red foxes (Vulpes vulpes) in urban areas in Great Britain:
Findings and limitations of a media-driven nationwide survey. PLoS ONE. 2014;9:e99059.
43.   Trewhella WJ, Harris S, McAllister FE. Dispersal distance, home-range size and
population density in the red fox (Vulpes vulpes): A quantitative analysis. Journal of Applied
Ecology. 1988;25:423-34.
44.   Soulsbury CD, Iossa G, Baker PJ, White PCL, Harris S. Behavioral and spatial
analysis of extraterritorial movements in red foxes (Vulpes vulpes). Journal of Mammology.
2011;92(1):190-9.
45.   Griffin AS, Diquelou M, Perea M. Innovative problem solving in birds: a key role of
motor diversity. Animal Behaviour. 2014;92:221-7.
46.   Lee PC, Moura A. Necessity, unpredictability and opportunity: An exploration of
ecological and social drivers of behavioral innovation. In: Kaufman AB, Kaufman JC, editors.
Animal Creativity and Innovation: Academic Press; 2015. p. 317-30.
47.   Morton FB. Do wild raccoons (*Procyon lotor*) use tools? Animal Cognition.
2021;24:433-41.

48. Kolowski JM, Forrester TD. Camera trap placement and the potential for bias due to trails and other features. PLoS ONE. 2017;12:e0186679.

49. ESA. Land cover CCI Product User Guide Version 2. 2017.

50. Team RC. R: A language and environment for statistical computing. Vienna, Austria R Foundation for Statistical Computing 2021.

51. Harris S, Rayner JMV. A discriminant analysis of the current distribution of urban foxes (Vulpes vulpes) in Britain. Journal of Animal Ecology. 1986;55:605-11.

52. Pebesma E, Bivand R. Sp: Classes and methods for spatial data. 2023.

53. Soulsbury CD, Iossa G, Baker PJ, Cole NC, Funk, S. M., Harris S. The impact of sarcoptic mange Sarcoptes scabiei on the British fox Vulpes vulpes population. Mammal Review. 2007;37(4):278-96.

54. Baker P, Newman T, Harris S. Bristol's foxes - 40 years of change. British Wildlife. 2001;12:411-7.

55. Wilkinson D, Smith GC. A preliminary survey for changes in urban fox (Vulpes vulpes) densities in England and Wales, and implications for rabies control. Mammal Review. 2001;31:107-10.

56. Minias P, Jedlikowskil J, Wlodarczyk R. Development of urban behaviour is associated with time since urbanization in a reed-nesting waterbird. Urban Ecosystems. 2018;21:1021-8.

57. Vrbanec L, Matijevic V, Guenther A. Enhanced problem-solving ability as an adaptation to urban environments in house mice. Proceedings of the Royal Society B: Biological Sciences. 2021;288:20202504.

58. Johnson-Ulrich L, Yirga G, Strong RL, Holekamp KE. The effect of urbanization on innovation in spotted hyenas. Animal Cognition. 2021;24:1027-38.

59. Miranda AC, Schielzeth H, Sonntag T, Partecke J. Urbanization and its effects on personality traits: a result of microevolution or phenotypic plasticity? Global Change Biology. 2013;19:2634-44.

60. Mazza V, Guenther A. City mice and country mice: innovative problem solving in rural and urban noncommensal rodents. Animal Behaviour. 2021;172:197-210.

61. Griffin AS. Innovativeness as an emergent property: A new alignment of comparative and experimental research on animal innovation. Philosophical Transactions of the Royal Society B: Biological Sciences. 2016;371:20150544.

62. DEFRA. ENV23 - UK statistics on waste 2022 [Available from: https://www.gov.uk/government/statistical-data-sets/env23-uk-waste-data-and-management.

63. Greggor AL, Clayton NS, Fulford AJC, AThornton A. Street smart: faster approach towards litter in urban areas by highly neophobic corvids and less fearful birds Animal Behaviour. 2016;117:123-33.

599 **Figure 1.** (a) The spatial locations of all sampling sites across the UK (dark red points), (b) the recorded locations where foxes touched the novel

600 tasks (light blue points), (c) the recorded locations where foxes solved the tasks (light green points), and (d) significant (p<0.05) local clustering

601 of locations where foxes touched then solved the tasks (red points).

Figure 2. The relationship between urbanisation and whether a fox from a given location (a)

does not touch (red) or touches (green) a task, and (b) does not solve (red) or solves (green)

a task.

607 **Figure 3.** Exploratory diversity (based on the number of unique tactile behaviours from Table

608 1) displayed by foxes physically interacting with the novel tasks from relatively newer (post-

609 1986) versus older (pre-1986) urban fox populations.