**Chilling out or heating up? Thermal plasticity of seminal fluid proteins in *Drosophila***

***melanogaster***

Claudia Londoño-Nieto[1], Roberto García-Roa[1,2], Stuart Wigby[3], Irem Sepil[4] and Pau Carazo[1]

[1]Ethology Lab, Cavanilles Institute of Biodiversity and Evolutionary Biology, University of

Valencia, Valencia, Spain.

[2] Department of Biology, Lund University, Lund, Sweden.

[3] Department of Evolution, Ecology, and Behaviour, Institute of Infection, Veterinary &

Ecological Sciences, University of Liverpool, Liverpool, United Kingdom.

[4] Department of Biology, University of Oxford, Oxford, United Kingdom.

**Authors for correspondence:** Claudia Patricia Londoño Nieto (Claudia.londono@uv.es) +37

674572407.

**Abstract**

Seminal fluid proteins (SFPs) are key mediators of post-mating interactions, shaping fertility,

sperm use and female physiology with major consequences for both sexes. Because they often

enhance male fitness while imposing costs on females, SFPs are central to sexual selection and

sexual conflict. Their composition and allocation strategies are influenced by both environmental

conditions and male state. However, how abiotic ecological factors regulate SFP expression, and

how this interacts with sperm competition risk, remains poorly understood. Here, we use adult flies

from a wild *Drosophila melanogaster* population to test how exposure to natural temperature

variation (20ºC, 24ºC and 28ºC) across two-time scales (48h and 13d), and under either no or high

sperm competition risk, interact to shape SFP production and transfer to females. SFP production

and transfer profiles were significantly affected by temperature, both consistently reduced after

long-term (13d) exposure to low (20ºC) and high (28ºC) temperatures, compared to 24ºC. Proteins

from the sex peptide and ovulin networks, which mediate female post-copulatory responses and

are key to male fitness, followed this same pattern. Our results show that natural temperature

fluctuations, well within the optimal reproductive range for flies in the wild, can have a strong

effect on SFP production and transfer, and thus on post-copulatory sexual selection dynamics.

**Introduction**

The last couple of decades have witnessed a revolution in our understanding of the complexity and function of male ejaculates, much of which has been driven by an increasing realization of the importance of non-sperm components (Gillott, 2003; Perry et al., 2013; Wedell et al., 2002). Sperm is not transferred in isolation within the ejaculate, but alongside a wide variety of molecules, including proteins (seminal fluid proteins: SFPs) that are crucial for reproduction (Hopkins et al., 2017). Produced in the male reproductive organs and transferred to the female during mating, SFPs play essential roles in fertility, sperm storage and usage, and strongly impact the fitness of both sexes by affecting female physiology, behaviour, immunity, and life history, as well as biasing the outcome of sperm competition (Avila et al., 2011; Gillott, 2003; Hopkins & Perry, 2022; Perry et al., 2013; Poiani, 2006). SFPs are thus fundamental to male and female reproductive success and are subject to strong natural and sexual selection (Perry et al., 2013; Sirot et al., 2015). For example, SFPs can alter female physiology and behaviour in ways that are costly to the female, yet beneficial to the male, making them a case study in sexual conflict (Sirot et al., 2015).

Sexual conflict arises because, in sexually reproducing organisms, anisogamy often leads to distinct sex roles (Janicke et al., 2016), which in turn result in divergent reproductive interests between the sexes (Arnqvist & Rowe, 2005; Pizzari & Snook, 2003). Sexual conflict is particularly intense in polygynous species, where it frequently results in male adaptations that increase male fitness at the expense of harming females (i.e. male harm; Arnqvist & Rowe, 2005). Male harming of females is taxonomically widespread and can significantly depress net female productivity, with potentially substantial demographic consequences for populations (Flintham et al., 2023; Gómez-Llano et al., 2024; Martinez-Ruiz & Knell, 2017). An important avenue of male harm are male adaptations driven by strong sperm competition, such as traumatic insemination and "harmful"

SFPs (Mueller et al., 2007; Perry et al., 2013; Reinhardt et al., 2015). Work on *Drosophila*

*melanogaster* has been at the forefront of this field (Wigby et al., 2020), providing compelling

evidence that individual SFPs are sufficient to alter female post-mating physiology and behaviour

(Hopkins & Perry, 2022). For example, SFPs such as sex peptide (SP) and ovulin regulate key

female post-mating responses. SP stimulates oogenesis, reduces female receptivity to remating,

and facilitates sperm storage and release (Chapman, 2001; Chapman et al., 2003; Liu & Kubli,

2003; Soller et al., 1999), while ovulin increases the rate of release of existing eggs (ovulation) by

the ovaries. Furthermore, there are proteins that are crucial for SP and ovulin to function: *antr,*

*aqrs, intr, lectin-46Ca, lectin-46Cb, Sems, CG17575* and *CG9997*, contribute to the long-term

persistence of reduced female receptivity to remating, effective sperm release, and fecundity

stimulation (Findlay et al., 2014; Ravi Ram & Wolfner, 2007). *Semp1*, in turn, is necessary to

process the ovulin in mated females (Ravi Ram et al., 2006). Together, SP, ovulin and their

associated network proteins increase male reproductive success, and SP and its network proteins

are believed to be under sexual conflict, as they can substantially reduce female lifespan and long-

term reproductive output (Wigby & Chapman, 2005).

A male's state, environment and condition affect both the composition and allocation

strategies of his seminal fluid proteome (Wigby et al., 2009). *Drosophila* males, for instance, can

tailor the quantity of SFPs they transfer in response to perceived sperm competition risk (Hopkins,

Sepil, Thézénas, et al., 2019), and may even exploit SFPs from rival ejaculates to maximize their

own reproductive success (Sirot et al., 2011). There is also increasing evidence that SFPs' post-

mating effects on females can be highly plastic themselves (Hopkins & Perry, 2022). However,

our understanding of how abiotic ecological factors influence the production and transfer of SFPs,

and how these interact with the processes of sexual selection and conflict remain limited. This gap

persists despite mounting evidence suggesting that the environment is a crucial determinant of

both fertility and competitive fertilisation success (Sales et al., 2018; Vasudeva et al., 2014; Yun et

al., 2017). In particular, responses of SFPs to environmental variables are prime candidates for

mediating these effects. For example, studies show that the larval environment can shape SFP

allocation patterns later in adult life (Von Hellfeld et al., 2025; Wigby et al., 2016).

Temperature is a particularly salient ecological factor in this respect, due to its potentially

far-reaching effects on sexual reproduction and sexual selection/conflict processes (Dougherty et

al., 2024; García-Roa et al., 2020). In nature, temperature is subject to substantial spatiotemporal

variability at different scales (e.g. circadian, seasonal), potentially imposing strong selective

pressures on sperm and ejaculate traits related to sexual selection and reproductive performance,

especially in ectothermic species. For instance, high temperatures can impair gonad development,

reduce sperm production, viability and function, (Peña Jr. et al., 2019; Vasudeva et al., 2014; Wang

& Gunderson, 2022), and compromise protein stability and function (Berger et al., 2021). Indeed,

*D. melanogaster* SFPs are highly sensitive to heat stress (Canal Domenech & Fricke, 2022), and

recent evidence shows that, in this species, high temperatures affect SFP-induced female post-

mating responses and ensuing male harm levels (Londoño-Nieto et al., 2023). A small temperature

increase well within the optimal reproductive range of a natural population (i.e.from 24ºC to 28ºC,

where female lifetime reproductive success is equal across temperatures), is sufficient to reduce

both the impact of male ejaculates on female receptivity and overall male harm to females

(Londoño-Nieto et al., 2023). This finding is especially relevant, as it suggests that even mild,

naturally occurring temperature fluctuations may influence SFP production and transfer. Hence,

understanding how temperature influences SFP dynamics is crucial to understand how

environmental variation shapes male ejaculate composition and sexual selection dynamics, and

may also offer important insight into the thermal limits of fertility (Dougherty et al., 2024).

Here, we used *D. melanogaster* flies to investigate how typical temperature variation,

within the optimal reproductive range for a wild population, affects the production and transfer of

SFPs, and how these effects change with male response to perceived sperm competition risk

(hereafter SCR). We assessed these effects exposing adult males to temperature (i.e. 20ºC, 24ºC

and 28ºC) and SCR treatments (i.e. no SCR vs. high SCR) across two different time scales

(experiment 1: short-term exposure - 48 h; experiment 2: long-term exposure - 13 d).

We predicted that:

1) The seminal fluid proteome would be thermally plastic, such that SFP production and/or

transfer would be reduced at high and low temperatures (20ºC and 28ºC). This is based on

the hypothesis that male harm is mediated by SFPs, and harm was previously found to be

reduced at 20ºC and 28ºC relative to 24ºC (Londoño-Nieto et al., 2023).

2) Long-term exposure to high temperatures (28ºC) would alter the seminal fluid proteome,

especially the abundance of functionally important SFPs, (i.e. the sex peptide, ovulin and

their network proteins), such that they would be produced and/or transferred in lower

quantities. This is based on the finding that high temperatures reduce the impact of male

ejaculates on female receptivity compared with lower temperatures (Londoño-Nieto et al.,

2023).

**Materials and Methods**

<u>Stock Maintenance</u>

We carried out all experiments between November 2020 and March 2021, using individuals from

our field-collected population, "Vegalibre", established in October 2018 from three wineries in

Requena (Spain), and re-sampled annually to maintain natural genetic variation (Londoño-Nieto

et al., 2023). We keep Vegalibre in the laboratory with overlapping generations at an average

temperature of 24°C with daily pre-programmed fluctuations (±4°C) mimicking natural daily

temperature conditions during the reproductively active season, at ~60% humidity and on a

12:12 hr light:dark cycle. We used maize-malt medium (7 g of agar, 72 g of malt, 72 g of maize,

9 g of soya, 15 g of live yeast, 20 g of molasses, and 33 ml of Nipagin mix: 3 g of methyl 4-

hydroxy-benzoate and 28 ml of ethanol 90% per 1000 ml of water) as a food source throughout

maintenance and experiments. To collect experimental flies, we introduced yeasted grape juice

agar plates into stock populations to induce female oviposition. We then collected eggs and

placed them in bottles containing ~75 ml of medium to be incubated at $24 \pm 4$°C at a controlled

density  (Clancy & Kennington, 2001). We collected virgin flies within 6 hr of eclosion, under

ice anesthesia, and then sexed and kept them in vials with food until their use (see below). All

females used in our experiments were 4 days old, while males were 4 days old for experiment 1

and 16 days old for experiment 2, at the time when samples were collected.

<u>Proteomics experiments</u>

*Experimental overview and detailed methods*

To study the effect of temperature and its interaction with the sociosexual environment (i.e.

perceived SCR) in shaping the production and transfer of SFPs, we exposed adult male flies to

either no or high precopulatory levels of SCR (Hopkins, Sepil, Thézénas, et al., 2019), across

three stable temperature treatments of 20°C, 24°C, and 28°C, for two different treatment

durations/scales.

In experiment 1, we simulated short-term temperature effects (i.e. intragenerational effects) on

the production and transfer of SFPs. Upon eclosion we randomly placed virgin males either

individually (no-SCR) or in a group of 8 males (high-SCR) in plastic vials with food for 2 days.

Next, we randomly divided them into three groups that we allocated to one of the three stable

temperature treatments (20°C, 24°C, and 28°C) for 48-hours (Fig. S1).

In experiment 2, we examined longer-term temperature effects on the production and transfer of

newly synthesized SFPs that would be typical of seasonal effects. We took 2 days-old males and

first depleted their ejaculates by housing them individually with four virgin females for 24 hours,

since *D. melanogaster* males can become sperm and SFP depleted when mated sequentially over

a short period (Macartney et al., 2021; Sanghvi et al., 2025; Sirot et al., 2009). Then, we placed

the 3 days-old depleted males for 13 days in the same temperature-SCR treatments of the

experiment 1 (Fig. S2). Because males take at least 3 days to fully replenish their SFPs after

depletion (Sirot et al., 2009), this design ensured that males synthesized new SFPs under our

temperature treatments.

Immediately after treating males, we collected all samples at a common garden

temperature of 24ºC. To explore SFP production patterns, we measured the abundance of SFPs in

virgin (experiment 1) or unmated (for 13d, experiment 2) males across all temperature-SCR

treatment combinations. To explore SFP transfer patterns, we compared the abundance of SFPs

in virgin or unmated males with the abundance of SFPs in recently mated males across the same

treatment combinations (Hopkins, Sepil, Thézénas, et al., 2019; Sepil et al., 2019, 2020).

*Experimental matings*

Four days before sample collection, we collected females as virgins and held them in same sex

groups of 15-20 flies at $24 \pm 4°C$. On the day of sample collection, we isolated 240 females in

vials with food, after which we immediately introduced one focal male either into a vial

containing a female or into an empty vial, to be retained as virgin or unmated. We flash froze

recently mated males in liquid nitrogen 25 min after the start of mating, freezing a virgin or

unmated male from the same temperature and SCR treatment combination at the same time

(Figs. S1-2). Freezing males at 25 min after the start of mating ensures a complete mating and is

consistent with the protocol used previously for proteomics experiments in this species (Hopkins,

Sepil, Bonham, et al., 2019; Hopkins, Sepil, Thézénas, et al., 2019; Sepil et al., 2019). To obtain

three independent biological replicates, we repeated this procedure over three different cohorts

from the same stock population on different days (i.e., blocks) for each experiment, with all SCR

and temperature treatments equally represented within each block.

We stored all frozen samples at -80°C until dissection, for which we thawed flash frozen

males and dissected their accessory glands, severing the ejaculatory duct and removing the

seminal vesicles and testes on ice in phosphate-buffered saline (PBS) under a binocular scope

(Hopkins, Sepil, Thézénas, et al., 2019). Each biological replicate (i.e. sample) consisted of a

pool of 20 reproductive glands in 25-µL PBS buffer. Samples were analysed with label-free

quantitative proteomics, using the sample preparation and quantification protocol (SWATH-MS;

Gillet et al., 2012) at the SCSIE proteomics service at the University of Valencia. Samples were

centrifuged and protein pellets resuspended in Laemmli buffer. Proteins were separated by SDS-

PAGE, full gel separation was performed for spectral library construction, while a short gel run

was used to clean and concentrate individual samples, followed by in-gel trypsin digestion as

described by Shevchenko et al. (1996). Peptides were analysed using a nanoESI qQTOF 6600+

TripleTOF mass spectrometer (AB SCIEX) coupled to a nanoflow LC system. Protein

identification and quantification were carried out using ProteinPilot and PeakView software with

a 1% false discovery rate threshold. Full methodological details are available in the

Supplementary Information. We analysed six samples per each temperature treatment-SCR

combination (three from virgin or unmated males and three from recently mated males). In total,

we had 72 samples (36 from experiment 1 and 36 from experiment 2).

Data analysis

We conducted all proteomics analysis on normalized abundances. We normalized the protein

areas calculated by the total sum of the areas of all the quantified proteins. We excluded proteins

with fewer than 2 unique peptides to restrict analysis to proteins with high-confidence

quantitation (Carr et al., 2004). We identified all proteins described as SFPs, based on a high-

confidence SFPs reference list from Wigby et al. (2020). We used an Elastic net penalized

logistic regression model and a partial least-squares discriminant analysis (PLS-DA) to analyse

our data sets, using *glmnet* (Friedman et al., 2010), *caret* (Kuhn, 2008) and *mixOmics* (Rohart et

al., 2017) packages in RStudio. Elastic net regression is a hybrid method that combines features

of Lasso and Ridge regularization (Zou & Hastie, 2005). It is particularly suited for situations

where the number of predictors far exceeds the number of observations, as it performs both

regularization and variable selection. The method applies a combined L1 (Lasso) and L2 (Ridge)

penalty that shrinks regression coefficients: weakly associated or irrelevant predictors are

penalized to exactly zero, effectively excluding them from the model, whereas variables with

stronger effects retain non-zero coefficients. Two parameters control the model: $\alpha$, which

balances the contribution of Lasso ($\alpha=1$) and Ridge ($\alpha=0$), and $\lambda$, which determines the overall

penalty strength. Both parameters were optimized using cross-validation (Zou & Hastie, 2005).

Under this framework, proteins with non-zero coefficients represent variables with the strongest

effects on proteome quantification.

To avoid overparameterization and maximize statistical power, we broke down analyses for both

experiments using the following scheme. To analyse how temperature affects SFP production

patterns (i.e. the abundance of SFPs in the accessory glands of virgin or unmated males), we

generated two datasets (one per SCR level), including all virgin (experiment 1) or unmated

(experiment 2) male samples across all temperature treatments. To analyse how temperature

influences SFP allocation, we first inferred the relative abundances of SFPs transferred by

dividing the abundance of each SFP in virgin or unmated males by its abundance in recently

mated males within the same temperature treatment, SCR level, and replicate. Based on these

ratios, we then generated two datasets (one per SCR level) including SFP transfer estimates

across all temperature treatments. To visualize the abundance of SFPs produced and transferred,

we took the average across the three biological replicates for each SFP within each temperature

treatment and SCR level. Heatmaps were generated from log2-transformed protein abundances,

using a Euclidean correlation distance metric and plotted with the aheatmap function from the

*NMF* package (Gaujoux & Seoighe, 2010) in RStudio (version R 4.2.2; R Core Team, 2022).

**Results**

We found a total of 1492 proteins, of which we retained 1328 for having 2 or more unique peptides.

Among these we focused on the 146 which were previously identified as SFPs.

*Temperature effects on the overall plasticity of SFP production and transfer*

Our results show that SFP production and transfer are highly plastic in response to temperature

variation across both treatment durations, but with important differences. In males treated for 48-

hour (short-term -experiment 1), elastic net regression identified 74 (under no-SCR) and 3 (under

high-SCR) SFPs as predictors contributing most strongly to temperature variation in protein

production (non-zero coefficients indicate proteins retained by the model; see Tables S2 and S3).

Similarly, the transfer of 79 (under no-SCR) and 6 (under high-SCR) SFPs was most strongly

associated with temperature variation (Tables S4 and S5). Heatmaps revealed that both production

and transfer of SFPs changed with temperature, with clearer effects under no-SCR. Partial least-

squares discriminant analysis (PLS-DA) supported these findings (Fig. 1), suggesting that SFP

production and transfer were more strongly affected by temperature in males exposed to no-SCR

than in those under high-SCR. Both the number of SFPs showing temperature-mediated changes

in abundance and the magnitude of these changes were markedly reduced under high-SCR (Fig.

1), suggesting that under high SCR male ejaculates are less thermally plastic after a short-term

exposure. In males treated for 13-days (long-term -experiment 2), elastic net regression selected

39 (under no-SCR) and 68 (under high-SCR) SFPs for production (Tables S6 and S7), and 49

(under no-SCR) and 46 (under high-SCR) SFPs for transfer (Tables S8 and S9) as predictors

contributing most strongly to differential response to temperature. Heatmap inspection showed

clear temperature-driven variation in SFP production and transfer, confirmed by the PLS-DA

analysis (Fig. 2). Overall, we found clear evidence that temperature effects after long-term

exposure were more uniform across SCR conditions, consistently reducing SPF production and

transfer at both low (20ºC) and hot (28ºC) temperatures (Fig. 2).

*Temperature effects on the production and transfer of SFPs involved in female post-mating*

*responses.*

For proteins involved in female post-mating responses, we identified 8 members of the SP network

(*antr, aqrs, intr, lectin-46Ca, lectin-46Cb, Sems, CG17575* and *CG9997*) as well as SP itself, ovulin

and its precursor, *Semp1*. Our results show that at 20ºC and 28°C, the production and transfer of

these proteins, and specifically those involved in the SP network, frequently diverged from that

observed at 24ºC, with clearer effects after the long-term treatment duration (experiment 2).

In experiment 1 (short-term exposure), SFP production revealed a consistent pattern:

protein abundance at 24ºC was markedly reduced compared to 20ºC and 28ºC, but only under no-

SCR (Fig. 1A and 3A). Among the proteins selected as important, having the strongest contribution

to temperature variation responses, under no-SCR, *Sems, lectin-46Cb, CG17575,* all belonging to

the sex peptide (SP) network, followed this general pattern. *SP* itself, however, and *Semp1*, a

member of the ovulin network (Ravi Ram et al., 2006), were also selected as important proteins

but displayed different patterns, with SP decreasing and *Semp1* increasing at 28ºC relative to lower

temperatures (Figs. 1A and S3). In contrast, under high-SCR we did not identify a consistent

pattern. However *Mst57Dc,* a protein related to the regulation of female receptivity (Findlay et al.,

2008, 2009; Simmerl et al., 1995) was selected as important under both SCR treatments, but with

different patterns: it showed lower abundance at 28ºC under no-SCR, whereas under high-SCR its

lowest abundance was observed at 24ºC (Figs. 1A and S3), suggesting a complex interaction

between thermal and social cues. Consistent with differences in production, SFP transfer under no-

SCR was generally reduced at 24ºC compared to 20ºC and 28ºC, while we did not find a clear

temperature-dependent pattern under high-SCR (Fig. 1D and 3B). Moreover, *ovulin, SP* and

proteins associated with SP network (i.e. *lectin-46Cb, CG9997*) were among those selected as

important only under no-SCR, following the general pattern (Figs. 1D and S4). Under high-SCR,

*lectin-46Ca*, a SP-network protein, was the only SPF known to regulate female post-mating

physiology selected as important contributing to temperature variation responses, with higher

transfer at 28ºC compared to lower temperatures (Figs. 1D and S4).

In experiment 2 (long-term exposure), SFP production showed a clear trend of increased

production at 24ºC compared to 20ºC and 28ºC, regardless of SCR (Fig. 2A and 3C). SP and ovulin

network proteins (i.e., *Semp1*, *antr*, *SP*, *CG17575)* were only selected as important predictors in

males under high-SCR (Fig 2A), showing the general pattern of increased production at 24ºC (Figs

2A and S5). *Acp26Ab, Acp53Ea* and *Mst57Dc* were also selected as important proteins only in

males under high-SCR but with increased production at 28ºC compared with at lower temperatures

(Figs 2A and S5). Patterns of SFP transfer mirrored those observed for production. In both SCR

conditions, 24ºC yielded the highest transfer levels (Figs. 2D and 3D). Among proteins showing

temperature-dependent transfer, elastic net identified *Semp1*, *antr*, *CG17575, lectin-46Cb, Ovulin*

and *Mst57Dc* in males under no-SCR, and *Semp1*, *CG17575, lectin-46Cb, lectin-46Ca, Acp53Ea*

and *Mst57Dc* in males under high-SCR as the strongest contributors to temperature variation.

These proteins exhibited markedly higher transfer at 24ºC compared to 20ºC and 28º (Figs. 2D and

S4). Despite the general similarity across SCR levels, more proteins from the SP and ovulin

networks were identified by elastic net as strongly temperature-sensitive in terms of production

under high-SCR, while transfer patterns showed similar thermal response across SCR levels (Figs.

2A and 2D).

**Discussion**

In this study, we show that temperature variation, within the optimal reproductive range for a wild

*D. melanogaster* population, affects both the production and transfer of SFPs, with effects shaped

by SCR and treatment duration. First, our results show that SFP production and transfer are very

plastic in response to temperature. In particular, long-term (13d) exposure to both low (20ºC) and

high (28ºC) temperatures resulted in lower production and transfer of SFPs. Second, we show that

these effects carry over to proteins with known important functions in female post-mating

respondes, whereby long-term exposure to high and low temperatures reduced the production and transfer of SFPs involved in the SP and ovulin networks. Overall, these findings highlight the role of natural temperature fluctuations in shaping ejaculate composition, with potential implications for the dynamics of sexual selection, sexual conflict, and temperature effects on reproduction.

*SFP production and transfer are thermally plastic*

In line with our predictions, we found a pronounced plastic response in SFP production and transfer after long-term exposure to both high and low temperatures, even though both temperatures fall well within the optimal reproductive range for this population in the wild (Londoño-Nieto et al., 2023). Males kept at 20°C or 28°C produced and transferred lower abundances of SFPs than at 24°C, with temperature effects on both the production and transfer being consistent across SCR levels. In the long-term exposure experiment, sexually mature males were exposed to different stable temperatures for 13-days following the depletion of accessory-glands via repeated matings (Macartney et al., 2021). While we cannot exclude the possibility that the number of females used to deplete male ejaculates (four) resulted in only a partial rather than full depletion (males can mate consecutively with up to 10 females; Sanghvi et al., 2025), males take at least 3 days to fully replenish their SFPs after depletion (Sirot et al., 2009). Thus, our design ensures that males were forced to synthesize new SFPs under our temperature treatments, fulfilling our aim of capturing effects on the production and transfer of new SFPs that would be typical of seasonal effects. Our results thus support the idea that seasonal changes in temperature may impose physiological constraints on the synthesis and/or transfer of SFPs, and potentially on associated trade-offs, that warrant further investigation. Furthermore, these effects fit well with the finding that male-induced harm to females is significantly reduced at 20ºC and 28ºC, as compared to 24ºC, in this same population of flies (Londoño-Nieto et al., 2023).

In contrast, our results following short-term exposure to high and low temperatures were

less clear and somewhat paradoxical. We again found clear temperature effects on both ejaculate

production and transfer, but thermal plasticity was strongly modulated by sociosexual context in a

way that is not clearcut. Under no SCR, we found higher production and transfer at both low and

high temperatures. Furthermore, these differences disappeared under high SCR. Our short-term

experiment was designed to capture effects on a fine-grained ecological scale (i.e. intragenerational

effects), given that sexually mature males were exposed to different stable temperatures for 48h

and flies typically live considerably longer lifespans in the wild. Compared to our long-term

treatment, we may expect temperature effects on the production of SFPs to be relatively limited in

this treatment, as males experienced only a brief thermal exposure. One possibility is that the

pattern observed under no-SCR in the short-term experiment may reflect compensatory plasticity,

in which males transiently increase SFP production and transfer to counteract negative long-term

thermal effects on SFPs at low and high temperatures (see above). Under high-SCR, previous

studies indicate that males invest maximally in the production and transfer of SFP at 25ºC

(Hopkins, Sepil, Thézénas, et al., 2019). This could explain why we find no differences across

temperature treatments under high-SCR, if males invested maximally in SFP, making short-term

thermal exposure to lower/higher temperatures insufficient to produce detectable changes.

However, it is important to stress that this is, at this point, speculative and predicated on the

assumption that short-term temperature effects on SFPs are adaptive, which may not be the case.

Broadly, our findings align with previous work showing that high temperatures disrupts

several stages of protein synthesis, including transcription, translation, folding, and post-

translational modification, which collectively affect protein functionality (Zheng et al., 2024).

These disruptions are particularly detrimental to protein and sperm phenotype and function across

animal taxa (Dougherty et al., 2024; Reinhardt et al., 2015; Sales et al., 2018). For instance, heat

stress commonly reduces sperm production, motility, viability, and longevity (Wang & Gunderson,

2022). It also increases molecular entropy, which can interfere with correct protein folding and

decreases the proportion of functional proteins (Berger et al., 2021). Recent work has extended

these findings to seminal fluid components, suggesting that SFPs may also be thermally sensitive

(Canal Domenech & Fricke, 2022; Martinet et al., 2023). Our findings build on this work by

showing that non-stressful thermal variation within the optimal reproductive range can

significantly alter SFP production and transfer. The finding that even subtle and relatively fine-

grained temperature variation can have clear effects on SFP production and transfer opens up a

new avenue to understand how ecology affects sexual selection traits/dynamics, and may also shed

light on expected effects of climate change on populations. There is growing recognition that

fertility limits, rather than lethal temperature limits, may be more accurate predictors of extinction

vulnerability (Dougherty et al., 2024). For instance, some *Drosophila* species can become sterile

at temperatures well below their lethal limit (Parratt et al., 2021; Van Heerwaarden & Sgrò, 2021).

Our results suggest that even below these fertility thresholds, natural temperature fluctuations may

still influence sexual selection dynamics, potentially increasing vulnerability to climate change in

ways not yet fully appreciated (García-Roa et al., 2020).

*Long-term exposure to low and high temperatures reduces the production and transfer of SFP that*

*mediate key post-mating processes*

Beyond the general effects of temperature on SFPs plasticity, we predicted that prolonged exposure

to 28ºC would particularly impact SFPs involved in the SP and ovulin networks based on the

evidence that the effect of male ejaculates on female remating behaviour is modulated by the

thermal environment males experienced prior to mating (Londoño-Nieto et al., 2023). Specifically,

ejaculates from males exposed during 13 days to 28°C induced a markedly weaker reduction in

female receptivity (i.e. shorter remating latency) compared to males kept at 20ºC and, especially,

at 24 °C (Londoño-Nieto et al., 2023). Our results support this prediction. After 13 days of

exposure, we observed consistently lower production and transfer of SP- and ovulin-associated

proteins at both 20ºC and 28ºC compared to 24ºC (Fig 2B).

Ejaculate composition is shaped by trade-offs between ejaculate components that mediate post-

mating sexual interactions and sperm competition (Perry et al., 2013). This, along with our results

rise an interesting question of whether temperature may affect males´ ability to strategically adjust

the production and transfer of SFPs in response to perceived SCR, where higher investment is

typically observed under high-SCR vs. no-SCR conditions (Hopkins, Sepil, Thézénas, et al., 2019;

Wigby et al., 2009). Although our design was not optimized to test this interaction directly, we

explored visual comparisons across temperature treatments to explore potential trends in

production and transfer of SFPs by males kept under high-SCR vs. no -SCR. We found a trend in

that high temperatures seem to disrupt the expected SCR-related patterns of SFP production and

transfer in both experiments, including proteins with key roles in female post-mating responses,

specifically those in the SP and ovulin networks (Figs. 4 and 5). At 28ºC, after short-term exposure,

the production of every member of both SP and ovulin networks (including ovulin but excluding

SP itself), was lower by males under high-SCR compared to males under no-SCR (Fig 4A).

Similarly, males under high-SCR transferred less *Semp1, lectin-46Cb, aqrs, ovulin, Sems, antr,*

*CG17575, CG9997,* and *SP* than those under no-SCR (Fig. 4B). After prolonged thermal exposure

at the same temperature (28ºC), males under high-SCR produced less *Semp1, lectin-46Cb, lectin-*

*46Ca, intr, Sems, antr, CG17575* and *SP* (Fig. 5A), and transferred less *intr, Sems, CG17575, lectin-*

*46Cb* and *Semp1* compared to males under no-SCR (Fig. 5B). Needless to say, this is just a trend

and thus must be taken as preliminary, but it suggests that temperature, and particularly high

temperatures, may limit males' ability to strategically adjust their ejaculate composition in

response to SCR. These results point to a potentially important interaction that warrants further

investigation.

*The ecological relevance of thermal effects on SFPs*

Taken together, our results underscore the broader implications of thermal sensitivity in ejaculate

composition. Due to the importance of SFPs that regulate key post-mating processes, their

plasticity (or lack thereof) under thermal variation could mediate individual fitness and shape

population responses to environmental change. Our results from experiment 1 suggest that short

term fine-scale temperature fluctuations (e.g. circadian) are unlikely to have major effects on

females and populations, especially under high SCR. However, our results form experiment

2capture the cumulative impact of thermal variation on ejaculate production and transfer over a

timeframe that spans both intragenerational and intergenerational seasonal effects. Such thermal

fluctuations are extremely common in the field, particularly in temperate climates, underscoring

the ecological relevance of our findings.

At the intragenerational scale, sustained temperature effects on continuous SFP production

and transfer are bound to alter male reproductive dynamics and overall reproductive success across

an individual's lifetime, potentially making plasticity itself a target of selection. Supporting this

idea, there is evidence showing that seminal fluid composition can vary among genotypes in

response to environmental conditions, with direct consequences for reproductive success (Patlar

& Ramm, 2020). Consistent with this, in the same wild *D. melanogaster* population used here, we

have detected strong GxE interactions in male reproductive success under thermal variation, with

individual genotypes exhibiting divergent, and sometimes opposing, quadratic reaction norms

(Londoño-Nieto et al., 2025). Interestingly, these GxE effects appear to be sexually dimorphic

(Panyam et al., unpublished results), indicating that males and females may experience, and

potentially adapt to, thermal variation in different ways. Moreover, experimental evolution under

sustained high-temperature regimes in this population revealed shifts in the abundance of several

SP- and ovulin-network proteins, suggesting that thermal environments can drive evolutionary

changes in SFPs composition (Londoño-Nieto et al., 2025).

At the intergenerational scale, prolonged periods of heat or cold may shift the targets of

post-copulatory sexual selection across seasons, potentially driving seasonal adaptive tracking

and/or balancing selection (Bergland et al., 2014; Hoffmann et al., 2005; Machado et al., 2021;

Rudman et al., 2022). For example, if higher temperatures consistently reduce the abundance or

effectiveness of key proteins like SP, this could weaken male ability to manipulate female post-

mating behaviour, potentially shifting the relative importance of pre- vs. post-copulatory sexual

selection towards the former. Moreover, as climate change increases the frequency and magnitude

of temperature fluctuations, such effects may be increasingly important for temperate populations

of ectothermic species.

**Conclusion**.

Our results show that natural temperature fluctuations can have a strong effect on SFP production

and transfer, and thus on post-copulatory sexual selection dynamics. We show that SFP production

and transfer are thermally plastic and modulated by sociosexual context and exposure duration.

Notably, long-term exposure to low and high temperatures consistently reduced the production and

transfer of SFPs, including key proteins involved in female post-mating responses. These findings

highlight how seasonal environmental variation can shape ejaculate traits under realistic ecological

conditions, and provide a mechanistic link for effects shown in recent studies linking habitat

complexity, reproductive plasticity, and eco-evolutionary processes shaping sexual selection (Berger & Liljestrand-Rönn, 2024; Gomez-Llano et al., 2018; Londoño-Nieto et al., 2023, 2025; Malek & Long, 2019; Yun et al., 2017, 2018). Importantly, these results may have important implications for understanding species vulnerability to climate change. We suggest a focus of future studies should be to examine how temperature influences the male´s ability to tailor SPF production and transfer in response to SCR, as well as the plasticity in female reproductive traits (i.e. their responses to male ejaculates), and how these female traits co-evolve with male ejaculates in response to temperature variation. Finally, these findings may also have applied relevance, as the thermal sensitivity of SFPs could improve strategies in pest control and assisted reproduction.

**References**

Arnqvist, G., & Rowe, L. (2005). *Sexual Conflict*. Princeton University Press, Princeton, NJ.

Avila, F. W., Sirot, L. K., LaFlamme, B. A., Rubinstein, C. D., & Wolfner, M. F. (2011). Insect Seminal Fluid Proteins: Identification and Function. *Annual Review of Entomology*, *56*(1), 21–40. https://doi.org/10.1146/annurev-ento-120709-144823

Berger, D., & Liljestrand-Rönn, J. (2024). Environmental complexity mitigates the demographic impact of sexual selection. *Ecology Letters*, *27*(1), e14355. https://doi.org/10.1111/ele.14355

Berger, D., Stångberg, J., Baur, J., & Walters, R. (2021). Elevated temperature increases genome-wide selection on de novo mutations. *Proceedings of the Royal Society B: Biological Sciences*, *287*. https://doi.org/10.1098/rspb.2020.3094

Bergland, A. O., Behrman, E. L., Brien, K. R. O., Schmidt, P. S., & Petrov, D. A. (2014). Genomic Evidence of Rapid and Stable Adaptive Oscillations over Seasonal Time Scales in Drosophila. *PLoS Genetics*, *10*(11). https://doi.org/10.1371/journal.pgen.1004775

Canal Domenech, B., & Fricke, C. (2022). Recovery from heat-induced infertility—A study of

reproductive tissue responses and fitness consequences in male *Drosophila melanogaster*.

*Ecology and Evolution*, *12*(12), e9563. https://doi.org/10.1002/ece3.9563

Carr, S., Aebersold, R., Baldwin, M., Burlingame, A., Clauser, K., & Nesvizhskii, A. (2004). The

Need for Guidelines in Publication of Peptide and Protein Identification Data. *Molecular*

*& Cellular Proteomics*, *3*(6), 531–533. https://doi.org/10.1074/mcp.T400006-MCP200

Chapman, T. (2001). Seminal fluid-mediated fitness traits in Drosophila. *Genetics*, *87*.

Chapman, T., Bangham, J., Vinti, G., Seifried, B., Lung, O., Wolfner, M. F., Smith, H. K., &

Partridge, L. (2003). The sex peptide of Drosophila melanogaster: Female post-mating

responses analyzed by using RNA interference. *Proceedings of the National Academy of*

*Sciences of the United States of America*, *100*(17), 9923–9928.

https://doi.org/10.1073/pnas.1631635100

Clancy, D. J., & Kennington, J. (2001). A simple method to achieve consistent larval density in

bottle cultures. *Drosophila Information Service*, *84*(January), 168–169.

Dougherty, L. R., Frost, F., Maenpaa, M. I., Rowe, M., Cole, B. J., Vasudeva, R., Pottier, P.,

Schultner, E., Macartney, E. L., Lindenbaum, I., Smith, J. L., Carazo, P., Graziano, M.,

Weaving, H., Domenech, B. C., Berger, D., Meena, A., Bishop, T. R., Noble, D. W. A., …

Price, T. A. R. (2024). A systematic map of studies testing the relationship between

temperature and animal reproduction. *Ecological Solutions and Evidence*, *5*(1).

https://doi.org/DOI:%252010.1002/2688-8319.12303

Findlay, G. D., MacCoss, M. J., & Swanson, W. J. (2009). Proteomic discovery of previously

unannotated, rapidly evolving seminal fluid genes in *Drosophila*. *Genome Research*,

*19*(5), 886–896. https://doi.org/10.1101/gr.089391.108

484 Findlay, G. D., Sitnik, J. L., Wang, W., Aquadro, C. F., Clark, N. L., & Wolfner, M. F. (2014).

485   Evolutionary Rate Covariation Identifies New Members of a Protein Network Required

486   for Drosophila melanogaster Female Post-Mating Responses. *PLoS Genetics*, *10*(1),

487   e1004108. https://doi.org/10.1371/journal.pgen.1004108

488 Findlay, G. D., Yi, X., MacCoss, M. J., & Swanson, W. J. (2008). Proteomics Reveals Novel

489   Drosophila Seminal Fluid Proteins Transferred at Mating. *PLoS Biology*, *6*(7), e178.

490   https://doi.org/10.1371/journal.pbio.0060178

491 Flintham, E. O., Savolainen, V., & Mullon, C. (2023). Male harm offsets the demographic

492   benefits of good genes. *Proceedings of the National Academy of Sciences of the United*

493   *States of America*, *120*(10), 1–9. https://doi.org/10.1073/pnas.2211668120

494 Friedman, J., Hastie, T., & Tibshirani, R. (2010). Regularization Paths for Generalized Linear

495   Models via Coordinate Descent. *Journal of Statistical Software*, *33*(1).

496   https://doi.org/10.18637/jss.v033.i01

497 García-Roa, R., Garcia-Gonzalez, F., Noble, D. W. A., & Carazo, P. (2020). Temperature as a

498   modulator of sexual selection. *Biological Reviews*, *95*(6), 1607–1629.

499   https://doi.org/10.1111/brv.12632

500 Gaujoux, R., & Seoighe, C. (2010). A flexible R package for nonnegative matrix factorization.

501   *BMC Bioinformatics*, *11*. https://doi.org/10.1186/1471-2105-11-367

502 Gillet, L. C., Navarro, P., Tate, S., Röst, H., Selevsek, N., Reiter, L., Bonner, R., & Aebersold, R.

503   (2012). Targeted data extraction of the MS/MS spectra generated by data-independent

504   acquisition: A new concept for consistent and accurate proteome analysis. *Molecular and*

505   *Cellular Proteomics*, *11*(6), 1–17. https://doi.org/10.1074/mcp.O111.016717

Gillott, C. (2003). Male Accessory Gland Secretions: Modulators of Female Reproductive

    Physiology and Behavior. *Annual Review of Entomology*, *48*(1), 163–184.

    https://doi.org/10.1146/annurev.ento.48.091801.112657

Gomez-Llano, M. A., Bensch, H. M., & Svensson, E. I. (2018). Sexual conflict and ecology:

    Species composition and male density interact to reduce male mating harassment and

    increase female survival. *Evolution*, *72*(4), 906–915. https://doi.org/10.1111/evo.13457

Gómez-Llano, M., Faria, G. S., García-Roa, R., Noble, D. W. A., & Carazo, P. (2024). Male

    harm suppresses female fitness, affecting the dynamics of adaptation and evolutionary

    rescue. *Evolution Letters*, *8*(1), 149–160. https://doi.org/10.1093/evlett/qrac002

Hoffmann, A. A., Hallas, R., Anderson, A., & Telonis-Scott, M. (2005). *Evidence for a robust*

    *sex-specific trade-off between cold resistance and starvation resistance in Drosophila*

    *melanogaster*. *18*, 804–810. https://doi.org/10.1111/j.1420-9101.2004.00871.x

Hopkins, B. R., & Perry, J. C. (2022). The evolution of sex peptide: Sexual conflict, cooperation,

    and coevolution. *Biological Reviews*, *97*(4), 1426–1448.

    https://doi.org/10.1111/brv.12849

Hopkins, B. R., Sepil, I., Bonham, S., Miller, T., Charles, P. D., Fischer, R., Kessler, B. M.,

    Wilson, C., & Wigby, S. (2019). BMP signaling inhibition in Drosophila secondary cells

    remodels the seminal proteome and self and rival ejaculate functions. *Proceedings of the*

    *National Academy of Sciences of the United States of America*, *116*(49), 24719–24728.

    https://doi.org/10.1073/pnas.1914491116

Hopkins, B. R., Sepil, I., Thézénas, M. L., Craig, J. F., Miller, T., Charles, P. D., Fischer, R.,

    Kessler, B. M., Bretman, A., Pizzari, T., & Wigby, S. (2019). Divergent allocation of

    sperm and the seminal proteome along a competition gradient in Drosophila

melanogaster. *Proceedings of the National Academy of Sciences of the United States of America*, *116*(36), 17925–17933. https://doi.org/10.1073/pnas.1906149116

Hopkins, B. R., Sepil, I., & Wigby, S. (2017). Seminal fluid. *Current Biology*, *27*(11), R404–R405. https://doi.org/10.1016/j.cub.2017.03.063

Janicke, T., Häderer, I. K., Lajeunesse, M. J., & Anthes, N. (2016). Evolutionary Biology: Darwinian sex roles confirmed across the animal kingdom. *Science Advances*, *2*(2), 1–11. https://doi.org/10.1126/sciadv.1500983

Kuhn, M. (2008). Building Predictive Models in *R* Using the **caret** Package. *Journal of Statistical Software*, *28*(5). https://doi.org/10.18637/jss.v028.i05

LaFlamme, B. A., Avila, F. W., Michalski, K., & Wolfner, M. F. (2014). A *Drosophila* Protease Cascade Member, Seminal Metalloprotease-1, Is Activated Stepwise by Male Factors and Requires Female Factors for Full Activity. *Genetics*, *196*(4), 1117–1129. https://doi.org/10.1534/genetics.113.160101

Liu, H., & Kubli, E. (2003). Sex-peptide is the molecular basis of the sperm effect in Drosophila melanogaster. *Proceedings of the National Academy of Sciences of the United States of America*, *100*(17), 9929–9933. https://doi.org/10.1073/pnas.1631700100

Londoño-Nieto, C., Butler-Margalef, M., García-Roa, R., & Carazo, P. (2025). Temperature Drives the Evolutionary Diversification of Male Harm in *Drosophila melanogaster* Flies. *Ecology Letters*, *28*(3), e70102. https://doi.org/10.1111/ele.70102

Londoño-Nieto, C., García-Roa, R., Garcia-Co, C., González, P., & Carazo, P. (2023). Thermal phenotypic plasticity of pre- and post-copulatory male harm buffers sexual conflict in wild Drosophila melanogaster. *eLife*, *12*, e84759. https://doi.org/10.7554/eLife.84759

Macartney, E. L., Zeender, V., Meena, A., De Nardo, A. N., Bonduriansky, R., & Lüpold, S.
    (2021). Sperm depletion in relation to developmental nutrition and genotype in
    Drosophila melanogaster. *Evolution*, *75*(11), 2830–2841.
    https://doi.org/10.1111/evo.14373

Machado, H. E., Bergland, A. O., Taylor, R., Tilk, S., Behrman, E., Dyer, K., Fabian, D. K., Flatt,
    T., González, J., Karasov, T. L., Kim, B., Kozeretska, I., Lazzaro, B. P., Merritt, T. J.,
    Pool, J. E., O'Brien, K., Rajpurohit, S., Roy, P. R., Schaeffer, S. W., … Petrov, D. A.
    (2021). Broad geographic sampling reveals the shared basis and environmental correlates
    of seasonal adaptation in Drosophila. *eLife*, *10*, e67577.
    https://doi.org/10.7554/eLife.67577

Malek, H., & Long, T. (2019). Spatial environmental complexity mediates sexual conflict and
    sexual selection in Drosophila melanogaster. *Ecology and Evolution*, *9*(5).
    https://doi.org/10.1002/ece3.4932

Martinet, B., Przybyla, K., Decroo, C., Wattiez, R., & Aron, S. (2023). Proteomic differences in
    seminal fluid of social insects whose sperm differ in heat tolerance. *Royal Society Open*
    *Science*, *10*(11), 231389. https://doi.org/10.1098/rsos.231389

Martinez-Ruiz, C., & Knell, R. J. (2017). Sexual selection can both increase and decrease
    extinction probability: Reconciling demographic and evolutionary factors. *Journal of*
    *Animal Ecology*, *86*, 117–127. https://doi.org/doi:%252010.1111/1365-2656.12601

Mueller, J. L., Page, J. L., & Wolfner, M. F. (2007). An Ectopic Expression Screen Reveals the
    Protective and Toxic Effects of Drosophila Seminal Fluid Proteins. *Genetics*, *175*(2),
    777–783. https://doi.org/10.1534/genetics.106.065318

Parratt, S. R., Walsh, B. S., Metelmann, S., White, N., Manser, A., Bretman, A. J., Hoffmann, A.

A., Snook, R. R., & Price, T. A. R. (2021). Temperatures that sterilize males better match

global species distributions than lethal temperatures. *Nature Climate Change*, *11*(6), 481–

484. https://doi.org/10.1038/s41558-021-01047-0

Patlar, B., & Ramm, S. A. (2020). Genotype-by-environment interactions for seminal fluid

expression and sperm competitive ability. *Journal of Evolutionary Biology*, *33*(2), 225–

236. https://doi.org/10.1111/jeb.13568

Peña Jr., S. T., Stone, F., Gummow, B., Parker, A. J., & Paris, D. B. B. P. (2019). Tropical

summer induces DNA fragmentation in boar spermatozoa: Implications for evaluating

seasonal infertility. *Reproduction, Fertility and Development*, *31*(3), 590.

https://doi.org/10.1071/RD18159

Perry, J. C., Sirot, L., & Wigby, S. (2013). The seminal symphony: How to compose an ejaculate.

*Trends in Ecology and Evolution*, *28*(7), 414–422.

https://doi.org/10.1016/j.tree.2013.03.005

Pizzari, T., & Snook, R. R. (2003). Perspective: Sexual conflict and sexual selection: Chasing

away paradigm shifts. *Evolution*, *57*(6), 1223–1236. https://doi.org/10.1111/j.0014-

3820.2003.tb00331.x

Poiani, A. (2006). Complexity of seminal fluid: A review. *Behavioral Ecology and Sociobiology*,

*60*(3), 289–310. https://doi.org/10.1007/s00265-006-0178-0

R Core Team. (2022). *R: A language and environment for statistical computing.* [Computer

software]. R   Foundation for Statistical Computing, Vienna, Austria. https://www.R-

project.org/.

Ravi Ram, K., Sirot, L. K., & Wolfner, M. F. (2006). Predicted seminal astacin-like protease is

      required for processing of reproductive proteins in *Drosophila melanogaster*.

      *Proceedings of the National Academy of Sciences*, *103*(49), 18674–18679.

      https://doi.org/10.1073/pnas.0606228103

Ravi Ram, K., & Wolfner, M. F. (2007). Seminal influences: Drosophila Acps and the molecular

      interplay between males and females during reproduction. *Integrative and Comparative*

      *Biology*, *47*(3), 427–445. https://doi.org/10.1093/icb/icm046

Reinhardt, K., Anthes, N., & Lange, R. (2015). Copulatory wounding and traumatic

      insemination. *Cold Spring Harbor Perspectives in Biology*, *7*(5), 1–26.

      https://doi.org/10.1101/cshperspect.a017582

Rohart, F., Gautier, B., Singh, A., & Lê Cao, K.-A. (2017). mixOmics: An R package for 'omics

      feature selection and multiple data integration. *PLOS Computational Biology*, *13*(11),

      e1005752. https://doi.org/10.1371/journal.pcbi.1005752

Rudman, S. M., Greenblum, S. I., Rajpurohit, S., Betancourt, N. J., Hanna, J., Tilk, S.,

      Yokoyama, T., Petrov, D. A., & Schmidt, P. (2022). Direct observation of adaptive

      tracking on ecological time scales in *Drosophila*. *Science*, *375*(6586), eabj7484.

      https://doi.org/10.1126/science.abj7484

Sales, K., Vasudeva, R., Dickinson, M. E., Godwin, J. L., Lumley, A. J., Michalczyk, Ł.,

      Hebberecht, L., Thomas, P., Franco, A., & Gage, M. J. G. (2018). Experimental

      heatwaves compromise sperm function and cause transgenerational damage in a model

      insect. *Nature Communications*, *9*(1), 4771. https://doi.org/10.1038/s41467-018-07273-z

Sanghvi, K., Shandilya, S., Brown, A., Todorova, B., Jahn, M., Gascoigne, S. J. L., Camilleri, T.-

      L., Pizzari, T., & Sepil, I. (2025). Reproductive output of old males is limited by seminal

fluid, not sperm number. *Evolution Letters*, *9*(2), 282–291.

https://doi.org/10.1093/evlett/qrae071

Sepil, I., Hopkins, B. R., Dean, R., Bath, E., Friedman, S., Swanson, B., Ostridge, H. J., Harper,

L., Buehner, N. A., Wolfner, M. F., Konietzny, R., Thézénas, M.-L., Sandham, E.,

Charles, P. D., Fischer, R., Steinhauer, J., Kessler, B. M., & Wigby, S. (2020). Male

reproductive aging arises via multifaceted mating-dependent sperm and seminal

proteome declines, but is postponable in *Drosophila*. *Proceedings of the National*

*Academy of Sciences*, *117*(29), 17094–17103. https://doi.org/10.1073/pnas.2009053117

Sepil, I., Hopkins, B. R., Dean, R., Thézénas, M.-L., Charles, P. D., Konietzny, R., Fischer, R.,

Kessler, B. M., & Wigby, S. (2019). Quantitative Proteomics Identification of Seminal

Fluid Proteins in Male Drosophila melanogaster. *Molecular & Cellular Proteomics*, *18*,

S46–S58. https://doi.org/10.1074/mcp.RA118.000831

Simmerl, E., Schäfer, M., & Schäfer, U. (1995). Structure and regulation of a gene cluster for

male accessory gland transcripts in Drosophila melanogaster. *Insect Biochemistry and*

*Molecular Biology*, *25*(1), 127–137. https://doi.org/10.1016/0965-1748(94)00034-F

Sirot, L. K., Buehner, N. A., Fiumera, A. C., & Wolfner, M. F. (2009). Seminal fluid protein

depletion and replenishment in the fruit fly, Drosophila melanogaster: An ELISA-based

method for tracking individual ejaculates. *Behavioral Ecology and Sociobiology*, *63*(10),

1505–1513. https://doi.org/10.1007/s00265-009-0806-6

Sirot, L. K., Wolfner, M. F., & Wigby, S. (2011). Protein-specific manipulation of ejaculate

composition in response to female mating status in Drosophila melanogaster. *Proceedings*

*of the National Academy of Sciences of the United States of America*, *108*(24), 9922–

9926. https://doi.org/10.1073/pnas.1100905108

Sirot, L. K., Wong, A., Chapman, T., & Wolfner, M. F. (2015). Sexual Conflict and Seminal Fluid

Proteins: A Dynamic Landscape of Sexual Interactions. *Cold Spring Harbor Perspectives*

*in Biology*, *7*(2), a017533. https://doi.org/10.1101/cshperspect.a017533

Soller, M., Bownes, M., & Kubli, E. (1999). Control of Oocyte Maturation in Sexually

MatureDrosophilaFemales. *Developmental Biology*, *208*(2), 337–351.

https://doi.org/10.1006/dbio.1999.9210

Van Heerwaarden, B., & Sgrò, C. M. (2021). Male fertility thermal limits predict vulnerability to

climate warming. *Nature Communications*, *12*(1), 2214. https://doi.org/10.1038/s41467-

021-22546-w

Vasudeva, R., Deeming, D. C., & Eady, P. E. (2014). Developmental temperature affects the

expression of ejaculatory traits and the outcome of sperm competition in *Callosobruchus*

*maculatus*. *Journal of Evolutionary Biology*, *27*(9), 1811–1818.

https://doi.org/10.1111/jeb.12431

Von Hellfeld, R., Konietzny, R., Charles, P. D., Fischer, R., Kessler, B. M., Wigby, S., Sepil, I., &

Morimoto, J. (2025). *The developmental environment mediates adult seminal proteome*

*allocation in male* Drosophila melanogaster. Evolutionary Biology.

https://doi.org/10.1101/2025.01.09.632097

Wang, W. W.-Y., & Gunderson, A. R. (2022). The Physiological and Evolutionary Ecology of

Sperm Thermal Performance. *Frontiers in Physiology*, *13*, 754830.

https://doi.org/10.3389/fphys.2022.754830

Wedell, N., Gage, M. J. G., & Parker, G. A. (2002). Sperm competition, male prudence and

sperm-limited females. *TREE*, *17*(7), 313–320.

Wigby, S., Brown, N. C., Allen, S. E., Misra, S., Sitnik, J. L., Sepil, I., Clark, A. G., & Wolfner,

M. F. (2020). The Drosophila seminal proteome and its role in postcopulatory sexual

selection. *Philosophical Transactions of the Royal Society of London B Biological*

*Sciences*, 20200072.

Wigby, S., & Chapman, T. (2005). Sex peptide causes mating costs in female Drosophila

melanogaster. *Current Biology*, *15*(4), 316–321.

Wigby, S., Perry, J. C., Kim, Y., & Sirot, L. K. (2016). Developmental environment mediates

male seminal protein investment in *D rosophila melanogaster* . *Functional Ecology*,

*30*(3), 410–419. https://doi.org/10.1111/1365-2435.12515

Wigby, S., Sirot, L. K., Linklater, J. R., Buehner, N., Calboli, F. C. F., Bretman, A., Wolfner, M.

F., & Chapman, T. (2009). Seminal Fluid Protein Allocation and Male Reproductive

Success. *Current Biology*, *19*(9), 751–757. https://doi.org/10.1016/j.cub.2009.03.036

Yun, L., Chen, P. J., Kwok, K. E., Angell, C. S., Rundle, H. D., & Agrawal, A. F. (2018).

Competition for mates and the improvement of nonsexual fitness. *Proceedings of the*

*National Academy of Sciences of the United States of America*, *115*(26), 6762–6767.

https://doi.org/10.1073/pnas.1805435115

Yun, L., Chen, P. J., Singh, A., Agrawal, A. F., & Rundle, H. D. (2017). The physical

environment mediates male harm and its effect on selection in females. *Proceedings of*

*the Royal Society B: Biological Sciences*. https://doi.org/10.1098/rspb.2017.0424

Zheng, J., Guo, N., Huang, Y., Guo, X., & Wagner, A. (2024). High temperature delays and low

temperature accelerates evolution of a new protein phenotype. *Nature Communications*,

*15*(1), 2495. https://doi.org/10.1038/s41467-024-46332-6

Zou, H., & Hastie, T. (2005). Regularization and Variable Selection Via the Elastic Net. *Journal*

*of the Royal Statistical Society Series B: Statistical Methodology*, *67*(2), 301–320.

https://doi.org/10.1111/j.1467-9868.2005.00503.x

**Figure 1 | Effect of temperature on SFPs production and transfer (short-term exposure / experiment 1)**. A) Heatmap showing the abundance of 145 SFPs detected in accessory gland samples. Elastic net regression selected 74 (under no-SCR) and 3 (under High-SCR) out of 145 SFPs as predictors with a strong effect differentiating proteomic responses across temperature treatments. Each cell gives the across-biological replicate mean for that protein, at each temperature treatment. Row annotations indicate SFPs with strong response to temperature in each SCR level and include functional information relating to protein functions as part of the sex peptide or ovulin networks, or other known roles in sperm competition. B) Partial least-squares discriminant analysis (PLS-DA) plot of proteins production for no-SCR treatment and C) PLS-DA plot of proteins production for high-SCR treatment. Points represent biological replicates. Ellipses denote variability among replicates. D) Heatmap showing the inferred relative abundance of 145 SFPs transferred to females during mating. Elastic net regression selected 79 (no SCR treatment) and 6 (High SCR treatment) out of 145 SFPs as predictors with strong differential transfer response to temperature. Each cell shows the mean ratio of protein abundance in virgin males relative to recently mated males, across biological replicates, for each temperature treatment, reflecting the relative loss of protein upon mating. Values closer to 1 indicate proteins reduced after mating (i.e., transferred). Heatmaps were generated from log2-transformed protein abundances for better visualization. E) PLS- DA plot of proteins transfer for no-SCR treatment and F) PLS-DA plot of proteins transfer for high-SCR treatment. Points in each PLS-DA plot represent biological replicates. Ellipses denote variability among replicates.

**Figure 2 | Effect of temperature on SFPs production and transfer (long-term exposure / experiment 2)**. A) Heatmap showing the abundance of the 145 SFPs detected in the accessory gland samples. Elastic net regression selected 39 (no SCR treatment) and 68 (High SCR treatment) out of 145 SFPs as predictors with a strong effect differentiating proteomic responses across temperature treatments. Each cell gives the across-biological replicate mean for that protein, at each temperature treatment. Row annotations indicate SFPs with strong response to temperature in each SCR level and include functional information relating to protein functions as part of the sex peptide or ovulin networks, or other known roles in sperm competition. B) PLS-DA plot of proteins production for no-SCR treatment and C) PLS- DA plot of proteins production for high-SCR treatment. Points represent biological replicates. Ellipses denote variability among replicates. D) Heatmap showing the inferred relative abundance of 145 SFPs transferred to females during mating. Elastic net regression selected 49 (no SCR treatment) and 46 (High SCR treatment) out of 145 SFPs as predictors with strong differential transfer response to temperature. Each cell shows the mean ratio of protein abundance in unmated males relative to recently mated males, across biological replicates, for each temperature treatment, reflecting the relative loss of protein upon mating. Values closer to 1 indicate proteins reduced after mating (i.e., transferred). Heatmaps were generated from log2-transformed protein abundances for better visualization. E) PLS- DA plot of proteins transfer for no-SCR treatment and F) PLS-DA plot of proteins transfer for high-SCR treatment. Points in each PLS-DA plot represent biological replicates. Ellipses denote variability among replicates.

**Figure 3 | Effect of temperature on production and transfer of SFPs involved in Ovulin and SP networks**. A – B) Relative abundance production and transfer profiles of SPFs associated with the SP and ovulin networks across different temperatures and SCR levels for short-term exposure (experiment 1). C – D) Relative abundance production and transfer profiles of SPFs associated with the SP and ovulin networks across different temperatures and SCR levels for long-term exposure (experiment 2). Each faint line represents the mean across the 3 replicates for an individual SFP within each treatment combination. Abundance values were normalized by mean-centering and averaged across replicates. Blue points represent the average (± s.e.) of SPFs in the SP network, across temperature and SCR conditions. Red points represent the average (± s.e.) of SFPs in the ovulin network, across temperature and SCR conditions.

**Figure 4 | Comparison of SFPs production and transfer under high-SCR vs. no-SCR across temperature treatments (short-term exposure / experiment 1).** A) Comparison for production. B) Comparison for transfer. Each cell shows the mean ratio of protein abundance under high-SCR relative to no-SCR across biological replicates for each temperature. Values closer to 1 indicate higher production/transfer under high-SCR conditions. Row annotations indicate functional information relating to protein functions as part of the sex peptide or ovulin networks, or other known roles in sperm competition.

**Figure 5 | Comparison of SFPs production and transfer under high-SCR vs. no-SCR across temperature treatments (long-term exposure / experiment 2).** A) Comparison for production. B) Comparison for transfer. Each cell shows the mean ratio of protein abundance under high-SCR relative to no-SCR across biological replicates for each temperature. Values closer to 1 indicate higher production/transfer under high-SCR conditions. Row annotations indicate functional information relating to protein functions as part of the sex peptide or ovulin networks, or other known roles in sperm competition.

SFPs' PRODUCTION

SFPs' TRANSFER

B  PLS-DA No-SCR

C  PLS-DA High-SCR

E  PLS-DA No-SCR

F  PLS-DA High-SCR

20ºC    24ºC    28ºC    20ºC    24ºC    28ºC
No-SCR              High-SCR

Important under No-SCR
Important under High-SCR
Others
SPnetwork
OvulinNetwork
PostMating Bahaviour / SpermCompetition
Other

SFPs' PRODUCTION

48-hour treatment duration (Exp. 1)

A

SFPs' TRANSFER

B

13 days treatment duration (Exp. 2)

C

D

Network
Ovulin
SP

A **SFPs' PRODUCTION**

B **SFPs' TRANSFER**

SPnetwork
OvulinNetwork
PostMating Baheviour / SpermCompetition
Other