Chilling out or heating up? Thermal plasticity of seminal fluid proteins in *Drosophila*

melanogaster

Claudia Londoño-Nieto[1], Roberto García-Roa[1,2], Stuart Wigby[3], Irem Sepil [4] and Pau Carazo[1]

[1]Ethology Lab, Cavanilles Institute of Biodiversity and Evolutionary Biology, University of

Valencia, Valencia, Spain.

[2] Department of Biology, Lund University, Lund, Sweden.

[3] Department of Evolution, Ecology, and Behaviour, Institute of Infection, Veterinary &

Ecological Sciences, University of Liverpool, Liverpool, United Kingdom.

[4] Department of Biology, University of Oxford, Oxford, United Kingdom.

Author for correspondence: Claudia Patricia Londoño Nieto (Claudia.londono@uv.es) +37

674572407.

Abstract

13   1. Seminal fluid proteins (SFPs) play crucial roles in reproduction by shaping female post-mating physiology and behaviour, male sperm competition, and sexual conflict between the sexes.

16   2. We mostly ignore how abiotic ecological factors regulate SFP expression and transfer.

17   3. Here, we use quantitative proteomics in adult flies from a wild *Drosophila melanogaster* population to test how exposure to natural temperature variation (20ºC, 24ºC and 28ºC) across two-time scales (48h and 13d), and under either no or high sperm competition risk, interact to shape SFP production and transfer.

21   4. We show that both production and transfer of SFPs was reduced after long-term (13d) exposure to low (20ºC) and high (28ºC) temperatures, including in key proteins (sex peptide and ovulin networks) mediating female post-copulatory responses, sperm competition and sexual conflict. Our results show that natural temperature fluctuations can have a strong effect on SFP production and transfer, and thus on post-copulatory sexual selection dynamics.

Keywords: Temperature, ecology, sexual selection, SFPs, reproduction performance, Drosophila, plasticity.

Introduction

The last couple of decades have witnessed a revolution in our understanding of the complexity and function of male ejaculates, much of which has been driven by an increasing realization of the importance of non-sperm components (Gillott, 2003; Perry et al., 2013; Wedell et al., 2002). Sperm is not transferred in isolation within the ejaculate, but alongside a wide variety of molecules, including proteins (seminal fluid proteins: SFPs) that are crucial for reproduction (Hopkins et al., 2017). Produced in the male reproductive organs and transferred to the female during mating, SFPs play essential roles in fertility, sperm storage and usage, and strongly impact the fitness of both sexes by affecting female physiology, behaviour, immunity, and life history, as well as biasing the outcome of sperm competition (Avila et al., 2011; Gillott, 2003; Hopkins & Perry, 2022; Perry et al., 2013; Poiani, 2006). SFPs are thus fundamental to male and female reproductive success and are subject to strong natural and sexual selection (Perry et al., 2013; Sirot et al., 2015). For example, SFPs can alter female physiology and behaviour in ways that are costly to the female, yet beneficial to the male, making them a case study in sexual conflict (Sirot et al., 2015).

Sexual conflict arises because, in sexually reproducing organisms, anisogamy often leads to distinct sex roles (Janicke et al., 2016), which in turn result in divergent reproductive interests between the sexes (Arnqvist & Rowe, 2005; Pizzari & Snook, 2003). Sexual conflict is particularly intense in polygynous species, where it frequently results in male adaptations that increase male fitness at the expense of harming females (i.e. male harm; Arnqvist & Rowe, 2005). Male harming of females is taxonomically widespread and can significantly depress net female productivity, with potentially substantial demographic consequences for populations (Flintham et al., 2023; Gómez-Llano et al., 2024; Martinez-Ruiz & Knell, 2017). An important avenue of male harm are male adaptations driven by strong sperm competition, such as traumatic insemination and "harmful"

SFPs (Mueller et al., 2007; Perry et al., 2013; Reinhardt et al., 2015). Work on *Drosophila*

melanogaster has been at the forefront of this field (Wigby et al., 2020), providing compelling

evidence that individual SFPs are sufficient to alter female post-mating physiology and behaviour

(Hopkins & Perry, 2022). For example, SFPs such as sex peptide (SP) and ovulin regulate key

female post-mating responses. SP stimulates oogenesis, reduces female receptivity to remating,

and facilitates sperm storage and release (Chapman, 2001; Chapman et al., 2003; Liu & Kubli,

2003; Soller et al., 1999), while ovulin increases the rate of release of existing eggs (ovulation) by

the ovaries. Furthermore, other proteins in the SP/ovulin network (i.e., *antr, aqrs, intr, lectin-46Ca,*

lectin-46Cb, Sems, CG17575 and *CG9997*) contribute to the long-term persistence of reduced

female receptivity to remating, effective sperm release, and fecundity stimulation (Findlay et al.,

2014; Ravi Ram & Wolfner, 2007), while *Semp1* is necessary to process ovulin in mated females

(Ravi Ram et al., 2006). Together, SP, ovulin and their associated network proteins increase male

reproductive success, and SP and its network proteins are believed to be under sexual conflict, as

they increase male fitness but can substantially reduce female lifespan and long-term reproductive

output (Wigby & Chapman, 2005).

There is increasing evidence that SFPs production and transfer is highly plastic. A male's

state, socio-sexual environment and condition can affect both the composition and allocation

strategies of his seminal fluid proteome (Wigby et al., 2009). *Drosophila* males, for instance, can

tailor the quantity of SFPs they transfer in response to perceived sperm competition risk (Hopkins

et al., 2019b), and may even exploit SFPs from rival ejaculates to maximize their own reproductive

success (Sirot et al., 2011). There is also evidence that SFPs' post-mating effects on females can

be highly plastic themselves (Hopkins & Perry, 2022). However, our understanding of how abiotic

ecological factors influence the production and transfer of SFPs, and how these interact with sexual

selection and conflict, remain limited. This gap persists despite mounting evidence suggesting that

the environment is a crucial determinant of both fertility and competitive fertilisation success

(Sales et al., 2018; Vasudeva et al., 2014; Yun et al., 2017). In particular, responses of SFPs to

environmental variables are prime candidates for mediating these effects. For example, studies

show that the larval environment can shape SFP allocation patterns later in adult life (Von Hellfeld

et al., 2025; Wigby et al., 2016).

Temperature is a particularly salient ecological factor in this respect, due to its potentially

far-reaching effects on sexual reproduction and sexual selection/conflict processes (Dougherty et

al., 2024; García-Roa et al., 2020). In nature, temperature is subject to substantial spatiotemporal

variability at different scales (e.g. circadian, seasonal), potentially imposing strong selective

pressures on sperm and ejaculate traits related to sexual selection and reproductive performance,

especially in ectothermic species. For instance, high temperatures can impair gonad development,

reduce sperm production, viability and function, (Peña et al., 2019; Vasudeva et al., 2014; Wang

& Gunderson, 2022), and compromise protein stability and function (Berger et al., 2021). Indeed,

D. melanogaster SFPs are highly sensitive to heat stress (Canal Domenech & Fricke, 2022), and

recent evidence shows that even a small temperature increase, well within the wild females'

optimal reproductive range under no sexual conflict (i.e. from 24ºC to 28ºC), is sufficient to reduce

both the impact of male ejaculates on female receptivity and overall male harm to females under

high sexual conflict (Londoño-Nieto et al., 2023). This finding is especially relevant, as it suggests

that even mild, naturally occurring temperature fluctuations may influence SFP production and

transfer. Hence, understanding how temperature influences SFP dynamics is crucial to understand

how environmental variation shapes male ejaculate composition and sexual selection dynamics,

and may also offer important insight into the thermal limits of fertility (Dougherty et al., 2024).

Here, we used *D. melanogaster* flies to investigate how natural temperature variation,

within the optimal reproductive range for a wild population, affects male production and transfer

of SFPs, and how these effects change with perceived sperm competition risk (hereafter SCR). We

assessed these effects exposing adult males to temperature (i.e. 20ºC, 24ºC and 28ºC) and SCR

treatments (i.e. no SCR vs. high SCR) across two different time scales (experiment 1: short-term

exposure-48h; experiment 2: long-term exposure-13d). We predicted that the seminal fluid

proteome would be thermally plastic, with 28ºC impacting the production and/or transfer of SFP,

particularly after long-term exposure (Berger et al., 2021; Zheng et al., 2024). We also predicted

that production/transfer of SFPs under sexual conflict (i.e. the sex peptide, ovulin and their network

proteins) would be generally reduced at 28ºC and 20ºC after long-term exposure, based on the fact

that male harm is reduced after long-term exposure to 20ºC and 28ºC relative to 24ºC in *Drosophila*

melanogaster (García-Roa et al., 2019; Londoño-Nieto et al., 2023). More specifically, we

predicted that long-term exposure to 28ºC would reduce the abundance of functionally important

SFPs involved in mediating female receptivity (i.e. *SP, antr, aqrs, intr, lectin-46Ca, lectin-46Cb,*

Sems, CG17575 and *CG9997*), given the reduced impact of male ejaculates on female receptivity

after long-term exposure to 28ºC (Londoño-Nieto et al., 2023).

Materials and Methods

Stock Maintenance

We carried out all experiments using individuals from our field-collected population, "Vegalibre",

established in October 2018 from three wineries in Requena (Spain), and re-sampled annually to

maintain natural genetic variation (Londoño-Nieto et al., 2023). We keep Vegalibre in the

laboratory with overlapping generations at an average temperature of 24°C with daily pre-

programmed fluctuations (±4°C) mimicking natural daily temperature conditions during the

reproductively active season, at ~60% humidity and on a 12:12 hr light:dark cycle. We used maize-

malt medium (see Londoño-Nieto et al., 2023) as a food source throughout maintenance and

experiments. To collect experimental flies, we introduced yeasted grape juice agar plates into stock

populations to induce female oviposition. We then collected eggs and placed them in bottles

containing ~75 ml of medium to be incubated at 24 ± 4°C at a controlled density (Clancy &

Kennington, 2001). We collected virgin flies within 6 hr of eclosion, under ice anesthesia, and then

sexed and kept them in vials with food until their use (see below). All females used in our

experiments were 4 days old, while males were 4 days old for experiment 1 and 16 days old for

experiment 2, at the time when samples were collected.

<u>Proteomics experiments</u>

Experimental overview and detailed methods

To study the effect of temperature, and its interaction with perceived SCR, in the production and

transfer of SFPs, we exposed adult male flies to either no or high precopulatory levels of SCR

(Hopkins et al., 2019b), across three stable temperature treatments of 20°C, 24°C, and 28°C, for

two different treatment durations/scales.

In experiment 1, we simulated short-term temperature effects (intragenerational effects) on

the production and transfer of SFPs. Upon eclosion, we randomly placed virgin males either

individually (no-SCR) or in a group of 8 males (high-SCR) in plastic vials with food for 2 days.

Next, we randomly divided them into three groups that we allocated to one of the three stable

temperature treatments (20°C, 24°C, and 28°C) for 48-hours (Fig. S1).

In experiment 2, we examined longer-term temperature effects on the production and

transfer of newly synthesized SFPs that would be typical of seasonal (intra- and inter-generational)

effects. We took 2 days-old males and first depleted their ejaculates by housing them individually

with four virgin females for 24 hours, since *D. melanogaster* males can become sperm and SFP

depleted when mated sequentially over a short period (Sirot et al., 2009). Then, we placed the 3

days-old depleted males for 13 days in the same temperature-SCR treatments of the experiment 1

(Fig. S2). Because males take at least 3 days to fully replenish their SFPs after depletion (Sirot et

al., 2009), this design ensured that males synthesized new SFPs under our temperature treatments.

Quantitative proteomics

Immediately after treating males, we explored SFP production patterns at a common garden

temperature of 24ºC, by measuring the abundance of SFPs in virgin (experiment 1) or recently

unmated (for 13d, experiment 2) males across all temperature-SCR treatment combinations. To

explore SFP transfer patterns, we compared the abundance of SFPs in virgin/recently unmated

males vs. recently mated males across the same treatment combinations (Hopkins et al., 2019b;

Sepil et al., 2019, 2020).

Four days before sample collection, we collected females as virgins and held them in same

sex groups of 15-20 flies at 24±4°C. On the day of sample collection, we isolated 240 females in

vials with food, after which we immediately introduced one focal male either into a vial containing

a female or into an empty vial. We flash froze recently mated males in liquid nitrogen 25 min after

the start of mating, freezing a virgin/recently unmated male from the same temperature-SCR

treatment combination at the same time (Figs. S1-2). Freezing males at 25 min after the start of

mating ensures a complete mating (see Hopkins et al., 2019a,b; Sepil et al., 2019). To obtain three

independent biological replicates, we repeated this procedure over three different cohorts from the

same stock population on different days (i.e. blocks) for each experiment, with all SCR and

temperature treatments equally represented within each block.

We stored all frozen samples at -80°C until dissection, for which we thawed flash frozen

males and dissected their accessory glands, severing the ejaculatory duct and removing the seminal

vesicles and testes on ice in phosphate-buffered saline (PBS) under a binocular scope (Hopkins et

al., 2019b). Each biological replicate (i.e. sample) consisted of a pool of 20 reproductive glands in

25-µL PBS buffer. Samples were analysed with label-free quantitative proteomics, using the

sample preparation and quantification protocol (SWATH-MS; Gillet et al., 2012) at the SCISIE

proteomics service at the University of Valencia. Samples were centrifuged and protein pellets

resuspended in Laemmli buffer. Proteins were separated by SDS-PAGE, full gel separation was

performed for spectral library construction, while a short gel run was used to clean and concentrate

individual samples, followed by in-gel trypsin digestion as described by Shevchenko et al. (1996).

Peptides were analysed using a nanoESI-qQTOF-6600+ TripleTOF mass spectrometer (AB

SCIEX) coupled to a nanoflow LC system. Protein identification and quantification were carried

out using ProteinPilot and PeakView software with a 1% false discovery rate threshold. Full

methodological details are available in SI. We analysed six samples per each temperature

treatment-SCR combination (three from virgin or recently unmated males and three from recently

mated males). In total, we had 72 samples (36 from experiment 1 and 36 from experiment 2) from

1440 males.

Data analysis

Scale of inference	Scale at which the factor of interest is applied	Number of replicates at the appropriate scale
Individuals	Sample - consisted of a pool of 20 reproductive glands	6 per each temperature treatment-SCR combination

We conducted all proteomics analysis on normalized abundances. We normalized protein areas

calculated by the total sum of the areas of all quantified proteins. We excluded proteins with fewer

than 2 unique peptides to restrict analysis to proteins with high-confidence quantification (Carr et

al., 2004). We identified SFPs based on the high-confidence SFPs reference list from Wigby et al.

(2020). We used an Elastic net penalized logistic regression model and a partial least-squares

discriminant analysis (PLS-DA) to analyse our data sets, using *glmnet* (Friedman et al., 2010),

caret (Kuhn, 2008) and *mixOmics* (Rohart et al., 2017) packages in RStudio. Elastic net regression

is a hybrid method that combines features of Lasso and Ridge regularization (Zou & Hastie, 2005).

It is particularly suited for situations where the number of predictors far exceeds the number of

observations, as it performs both regularization and variable selection. The method applies a

combined Lasso and Ridge penalty that shrinks regression coefficients: weakly associated or

irrelevant predictors are penalized to exactly zero, effectively excluding them from the model,

whereas variables with stronger effects retain non-zero coefficients. Two parameters control the

model: α, which balances the contribution of Lasso ($\alpha=1$) and Ridge ($\alpha=0$), and λ, which

determines the overall penalty strength. Both parameters were optimized using cross-validation

(Zou & Hastie, 2005). Under this framework, proteins with non-zero coefficients represent

variables with the strongest effects on proteome quantification.

To avoid overparameterization and maximize statistical power, we broke down analyses

for both experiments using the following scheme. To analyse how temperature affects SFP

production patterns (i.e. the abundance of SFPs in the accessory glands of virgin –in experiment

1– or recently unmated –in experiment 2– males), we generated two datasets (one per SCR level),

including all virgin (experiment 1) or recently unmated (experiment 2) male samples across all

temperature treatments. To analyse how temperature influences SFP allocation, we first inferred

the relative abundances of SFPs transferred by dividing the abundance of each SFP in virgin or

recently unmated males by its abundance in recently mated males within the same temperature

treatment, SCR level, and replicate. Based on these ratios, we then generated two datasets (one per

SCR level) including SFP transfer estimates across all temperature treatments. Note, however, that

we also analysed absolute abundances of SFPs transferred, with qualitatively very similar results

(see SI). To visualize the abundance of SFPs produced and transferred, we took the average across

the three biological replicates for each SFP within each temperature treatment and SCR level, and

then generated heatmaps using a Euclidean correlation distance metric and plotted with the

aheatmap function from the *NMF* package (Gaujoux & Seoighe, 2010) in RStudio (version R 4.2.2;

R Core Team, 2022).

Results

We found a total of 1492 proteins, of which we retained 1328 with 2 or more unique peptides.

Among these, we focused on the 146 previously identified as SFPs.

Temperature effects on the overall plasticity of SFP production and transfer

Our results show that SFP production and transfer are highly plastic in response to temperature

variation across both treatment durations, but with important differences across experiments. In

males treated for 48-hour (short-term -experiment 1), elastic net regression identified 74 (under

no-SCR) and 3 (under high-SCR) SFPs as predictors contributing most strongly to temperature

variation in protein production (see Tables S2 and S3), and 79 (under no-SCR) and 6 (under high-

SCR) in the case of SFP transfer (Tables S4 and S5). Heatmaps revealed that both production and

transfer of SFPs changed with temperature, with clearer effects under no-SCR. Partial least-squares

discriminant analysis (PLS-DA) supported these findings (Fig. 1), suggesting that SFP production

and transfer were more strongly affected by temperature in males exposed to no-SCR than in those

under high-SCR. Both the number of SFPs showing temperature-mediated changes in abundance

and the magnitude of these changes were markedly reduced under high-SCR (Fig. 1), suggesting

that under high SCR male ejaculates are less thermally plastic after a short-term exposure. In males

treated for 13-days (long-term -experiment 2), elastic net regression selected 39 (under no-SCR)

and 68 (under high-SCR) SFPs for production (Tables S6 and S7), and 49 (under no-SCR) and 46

(under high-SCR) SFPs for transfer (Tables S8 and S9) as predictors contributing most strongly to

differential response to temperature. Heatmap inspection showed clear temperature-driven

variation in SFP production and transfer, confirmed by the PLS-DA analysis (Fig. 2). Overall, we

found clear evidence that temperature effects after long-term exposure were more uniform across

SCR conditions, consistently reducing SPF production and transfer at both low (20ºC) and hot

(28ºC) temperatures (Fig. 2). Estimates of SFP transfer derived from absolute abundances

produced comparable heatmap patterns, supporting the observed temperature effects (Figs. S3-4).

Temperature effects on the production and transfer of SFPs involved in female post-mating

responses.

For proteins involved in female post-mating responses, we identified 8 members of the SP network

(*antr, aqrs, intr, lectin-46Ca, lectin-46Cb, Sems, CG17575* and *CG9997*) as well as SP itself,

ovulin, and its precursor *Semp1*. Our results show that, at 20ºC and 28°C, the production and

transfer of these proteins, and specifically those involved in the SP network, frequently diverged

from that observed at 24ºC, with clearer effects after the long-term treatment duration (experiment

2).

In experiment 1 (short-term exposure), SFP production revealed a consistent pattern:

protein abundance at 24ºC was markedly reduced compared to 20ºC and 28ºC, but only under no-

SCR (Fig. 1A and 3A). Among the selected proteins under no-SCR, SP network proteins *Sems,*

lectin-46Cb and *CG17575* followed this pattern. In contrast, *SP* decreased and *Semp1* increased at

28ºC (Figs. 1A and S5). Under high-SCR we did not identify a consistent pattern. Consistent with

differences in production, SFP transfer under no-SCR was generally reduced at 24ºC compared to

20ºC and 28ºC, while we did not find a clear temperature-dependent pattern under high-SCR (Fig.

1D, and 3B). Moreover, *ovulin, SP* and SP network proteins *lectin-46Cb* and *CG9997* were among

those selected as important only under no-SCR (Figs. 1D and S6). Note, however, that the absolute

transfer of *SP* was higher at 24ºC than at 20º and 28ºC (Fig. S3). Under high-SCR, *lectin-46Ca*, a

SP-network protein, was the only SPF known to regulate female post-mating physiology selected

as important, with higher relative and absolute transfer at 28ºC compared to lower temperatures

(Figs. 1D. S3 and S6).

In experiment 2 (long-term exposure), SFP production showed a clear trend of increased

production at 24ºC compared to 20ºC and 28ºC, regardless of SCR (Fig. 2A and 3C). SP and ovulin

network proteins (i.e., *Semp1*, *antr*, *SP*, *CG17575)* were only selected as important predictors in

males under high-SCR (Fig 2A), showing the general pattern of increased production at 24ºC (Figs

2A and S7). *Acp26Ab, Acp53Ea* and *Mst57Dc* were also selected as important proteins only in

males under high-SCR but with increased production at 28ºC compared with at lower temperatures

(Figs 2A and S7). Patterns of SFP transfer mirrored those observed for production. In both SCR

conditions, 24ºC yielded the highest relative and absolute transfer levels (Figs. 2D, 3D and S4).

Among proteins showing temperature-dependent transfer, elastic net identified *Semp1*, *antr*,

CG17575, lectin-46Cb, Ovulin and *Mst57Dc* in males under no-SCR, and *Semp1, CG17575,*

lectin-46Cb, lectin-46Ca, Acp53Ea and *Mst57Dc* in males under high-SCR as those most strongly

associated with temperature variation. These proteins exhibited markedly higher transfer at 24ºC

compared to 20ºC and 28ºC (Figs. 2D, S4 and S8). Despite the general similarity across SCR

levels, more proteins from the SP and ovulin networks were identified by elastic net as strongly

temperature-sensitive in terms of production under high-SCR, while transfer patterns showed

similar thermal response across SCR levels (Figs. 2A and 2D).

Discussion

In this study, we show that temperature variation, within the optimal reproductive range for a wild

D. melanogaster population, affects both the production and transfer of SFPs, with effects shaped

by SCR and treatment duration. First, our results show that SFP production and transfer are very

plastic in response to temperature. In particular, long-term (13d) exposure to both low (20ºC) and

high (28ºC) temperatures resulted in lower production and transfer of SFPs. Second, we show that

these effects carry over to proteins with known important functions in female post-mating

responses, whereby long-term exposure to high and low temperatures reduced the production and

transfer of SFPs involved in the SP and ovulin networks. Overall, these findings highlight the role

of natural temperature fluctuations in shaping ejaculate composition, with potential implications

for the dynamics of sexual selection, sexual conflict, and temperature effects on reproduction at

large.

SFP production and transfer are thermally plastic

In line with our predictions, we found a pronounced plastic response in SFP production and transfer

after long-term exposure to high but also to low temperatures, even though both temperatures fall

well within the optimal reproductive range for this population in the wild (Londoño-Nieto et al.,

2023). Males kept at 20°C or 28°C produced and transferred lower abundances of SFPs than at

24°C across SCR levels. In the long-term exposure experiment, sexually mature males were

exposed to different stable temperatures for 13-days following partial depletion (males can mate

consecutively with up to 10 females; Sanghvi et al., 2025) of accessory-glands via repeated

matings (Sirot et al., 2009). Thus, our design ensures that males were forced to synthesize new

SFPs under our temperature treatments, fulfilling our aim of capturing effects on the production

and transfer of new SFPs that would be typical of seasonal effects. Wild *D. melanogaster* flies are

likely to live long enough to experience seasonal thermal fluctuations, given that they can survive

up to 90 days in controlled conditions (Brischigliaro et al., 2023). Our results thus support the idea

that seasonal changes in temperature may impose physiological constraints on the synthesis and/or

transfer of SFPs, and fit well with the finding that male-induced harm to females is significantly

reduced at 20ºC and 28ºC (vs. 24ºC) in this same population of flies (Londoño-Nieto et al., 2023).

In contrast, our results following short-term exposure to high and low temperatures were

less clear and somewhat paradoxical. We again found clear temperature effects on both ejaculate

production and transfer, but thermal plasticity was strongly modulated by sociosexual context in a

complex way. Under no SCR, we found higher production and transfer at both low and high

temperatures, but these differences disappeared under high SCR. Our short-term experiment was

designed to capture effects on a fine-grained ecological scale (i.e. short intragenerational effects),

given that sexually mature males were exposed to different stable temperatures for 48h and flies

likely live considerably longer lifespans in the wild. Compared to our long-term treatment, we may

thus expect temperature effects on the production of SFPs to be relatively limited in this treatment.

Broadly, our findings align with previous work showing that high and low temperatures

influence several stages of protein synthesis, including transcription, translation, folding, and post-

translational modification, which collectively affect protein functionality (Feller & Gerday, 2003;

Zheng et al., 2024). In particular, high temperatures are known to be especially detrimental to

protein and sperm phenotype and function across animal taxa (Dougherty et al., 2024; Reinhardt et al., 2015; Sales et al., 2018). For instance, heat stress commonly reduces sperm production, motility, viability, and longevity (Wang & Gunderson, 2022). It also increases molecular entropy, which can interfere with correct protein folding and decreases the proportion of functional proteins (Berger et al., 2021). Recent work has extended these findings to seminal fluid components, suggesting that SFPs may also be thermally sensitive (Canal Domenech & Fricke, 2022; Martinet et al., 2023). Our findings build on this work by showing that non-stressful thermal variation within the optimal reproductive range can significantly alter SFP production and transfer. The finding that even subtle and relatively fine-grained temperature variation can have clear effects on SFP production and transfer opens up a new avenue to understand how ecology affects sexual selection, and may also shed light on expected effects of climate change on populations. There is growing recognition that fertility limits, rather than lethal temperature limits, may be more accurate predictors of extinction vulnerability (Dougherty et al., 2024). For instance, some *Drosophila* species can become sterile at temperatures well below their lethal limit (Parratt et al., 2021; Van Heerwaarden & Sgrò, 2021). Our results suggest that, even below these fertility thresholds, natural temperature fluctuations may still influence sexual selection dynamics, potentially increasing vulnerability to climate change in ways not yet fully appreciated (García-Roa et al., 2020).

Long-term exposure to low and high temperatures reduces the production and transfer of SFP that mediate key post-mating processes

Beyond the general effects of temperature on SFPs plasticity, we predicted that prolonged exposure to 28ºC would particularly impact functionally important SFPs involved in mediating female receptivity (Londoño-Nieto et al., 2023). Specifically, Londoño-Nieto et al. (2023) showed that ejaculates from males exposed during 13 days to 28°C induced a markedly weaker reduction in

female receptivity (i.e. shorter remating latency) compared to males kept at 20ºC and, especially,

at 24 °C. Our results fit well with this finding, as, after 13 days of exposure, we observed

consistently lower production and transfer of SP -and ovulin- associated proteins at both 28ºC and

20ºC compared to 24ºC (Fig 2B).

Ejaculate composition is shaped by trade-offs between ejaculate components that mediate

post-mating sexual interactions and sperm competition (Perry et al., 2013). Our results thus beg

the question of whether temperature may affect males´ ability to strategically adjust the production

and transfer of SFPs in response to perceived SCR, where higher investment is typically observed

under high-SCR vs. no-SCR conditions (Hopkins et al., 2019b; Wigby et al., 2009). Although our

design was not optimized to test this interaction directly, we explored visual comparisons across

temperature treatments to explore potential trends in production and transfer of SFPs by males

kept under high-SCR vs. no -SCR. We found a trend in that high temperatures seem to disrupt the

expected SCR-related patterns of SFP production and transfer in both experiments, especially in

SP and ovulin network proteins (Figs. 4 and 5). At 28ºC, after short-term exposure, males under

high-SCR produced relatively less of the proteins belonging to the SP and ovulin networks

(including ovulin but excluding SP itself) compared to males under no-SCR, than they did at 24ºC

(Fig 4A), and transferred less *lectin-46Ca, lectin-46Cb, CG9997,* and *SP* than those under no-SCR

(Fig. 4B). After prolonged thermal exposure at 28ºC, males under high-SCR produced less *Semp1,*

lectin-46Cb, lectin-46Ca, intr, Sems, antr, CG17575 and *SP* (Fig. 5A), and transferred less *Sems,*

lectin-46Ca, *antr*, *Semp1, ovulin* and *SP* than males under no-SCR, than they did at 24ºC (Fig.

5B). This trend must be taken as preliminary, but suggests that high temperatures may limit males'

ability to strategically adjust their ejaculate composition in response to SCR.

The ecological relevance of thermal effects on SFPs

Taken together, our results underscore the broader implications of thermal sensitivity in ejaculate

composition. Due to the importance of SFPs, their plasticity (or lack thereof) under thermal

variation could mediate individual fitness and shape population responses to environmental

change. Our results from experiment 1 suggest that short term fine-scale temperature fluctuations

(e.g. circadian) are unlikely to have major effects on females and populations, especially under

high SCR. However, our results from experiment 2 capture the cumulative impact of thermal

variation on ejaculate production and transfer over a timeframe that spans both intra- and inter-

generational seasonal effects. Such thermal fluctuations are extremely common in the field

(particularly in temperate climates), and thus are of clear ecological relevance.

At the intragenerational scale, sustained temperature effects on continuous SFP production

and transfer are bound to alter male reproductive dynamics and overall reproductive success across

an individual's lifetime, potentially making plasticity itself a target of selection. Supporting this

idea, there is evidence showing that seminal fluid composition can vary among genotypes in

response to environmental conditions, with direct consequences for reproductive success (Patlar

& Ramm, 2020). In addition, *D. melanogaster* from the same population used in this study show

strong thermal GxE interactions in male reproductive success (Londoño-Nieto et al., 2025).

Interestingly, these GxE effects appear to be sexually dimorphic (Panyam et al., unpublished

results), indicating that males and females may experience, and potentially adapt to, thermal

variation in different ways. Moreover, a recent experimental evolution experiment shows the

thermal environment can drive quick evolutionary changes in SFPs composition (Londoño-Nieto

et al., 2025).

At the intergenerational scale, prolonged periods of heat or cold may shift the targets of

post-copulatory sexual selection across seasons, potentially driving seasonal adaptive tracking

and/or balancing selection (Bergland et al., 2014; Hoffmann et al., 2005; Machado et al., 2021;

Rudman et al., 2022). For example, if higher temperatures consistently reduce the abundance or

effectiveness of key proteins like SP, this could weaken male ability to manipulate female post-

mating behaviour, potentially shifting the relative importance of pre- vs. post-copulatory sexual

selection towards the former. As climate change increases the frequency and magnitude of

temperature fluctuations, such effects may be increasingly important for temperate populations of

ectothermic species.

Conclusions

Our results show that natural temperature fluctuations can have a strong effect on SFP production

and transfer, and thus on post-copulatory sexual selection dynamics. Notably, long-term exposure

to low and high temperatures consistently reduced the production and transfer of SFPs, including

key proteins involved in female post-mating responses. These findings highlight how seasonal

environmental variation can shape ejaculate traits under realistic ecological conditions, and

provide a mechanistic link for effects shown in recent studies linking habitat complexity,

reproductive plasticity, and eco-evolutionary processes shaping sexual selection (Berger &

Liljestrand-Rönn, 2024; Gomez-Llano et al., 2018; Londoño-Nieto et al., 2023, 2025; Malek &

Long, 2019; Yun et al., 2017, 2018). Importantly, these results may have important implications

for understanding species vulnerability to climate change. We suggest a focus of future studies

should be to examine how temperature influences the male´s ability to tailor SPF production and

transfer in response to SCR, as well as the plasticity in female reproductive traits (i.e. their

responses to male ejaculates), and how these female traits co-evolve with male ejaculates in

response to temperature variation. Finally, these findings may also have applied relevance, as the

thermal sensitivity of SFPs could improve strategies in pest control and assisted reproduction.

References

Arnqvist, G., & Rowe, L. (2005). *Sexual Conflict*. Princeton University Press, Princeton, NJ.

Avila, F. W., Sirot, L. K., LaFlamme, B. A., Rubinstein, C. D., & Wolfner, M. F. (2011). Insect Seminal Fluid Proteins: Identification and Function. *Annual Review of Entomology*, *56*(1), 21–40. https://doi.org/10.1146/annurev-ento-120709-144823

Berger, D., & Liljestrand-Rönn, J. (2024). Environmental complexity mitigates the demographic impact of sexual selection. *Ecology Letters*, *27*(1), e14355. https://doi.org/10.1111/ele.14355

Berger, D., Stångberg, J., Baur, J., & Walters, R. (2021). Elevated temperature increases genome-wide selection on de novo mutations. *Proceedings of the Royal Society B: Biological Sciences*, *287*. https://doi.org/10.1098/rspb.2020.3094

Bergland, A. O., Behrman, E. L., Brien, K. R. O., Schmidt, P. S., & Petrov, D. A. (2014). Genomic Evidence of Rapid and Stable Adaptive Oscillations over Seasonal Time Scales in Drosophila. *PLoS Genetics*, *10*(11). https://doi.org/10.1371/journal.pgen.1004775

Brischigliaro, M., Fernandez-Vizarra, E., & Viscomi, C. (2023). Mitochondrial Neurodegeneration: Lessons from Drosophila melanogaster Models. *Biomolecules*, *13*(2), 378. https://doi.org/10.3390/biom13020378

Canal Domenech, B., & Fricke, C. (2022). Recovery from heat-induced infertility—A study of reproductive tissue responses and fitness consequences in male *Drosophila melanogaster*. *Ecology and Evolution*, *12*(12), e9563. https://doi.org/10.1002/ece3.9563

Carr, S., Aebersold, R., Baldwin, M., Burlingame, A., Clauser, K., & Nesvizhskii, A. (2004). The Need for Guidelines in Publication of Peptide and Protein Identification Data. *Molecular & Cellular Proteomics*, *3*(6), 531–533. https://doi.org/10.1074/mcp.T400006-MCP200

Chapman, T. (2001). Seminal fluid-mediated fitness traits in Drosophila. *Genetics*, *87*.

Chapman, T., Bangham, J., Vinti, G., Seifried, B., Lung, O., Wolfner, M. F., Smith, H. K., &

Partridge, L. (2003). The sex peptide of Drosophila melanogaster: Female post-mating

responses analyzed by using RNA interference. *Proceedings of the National Academy of*

Sciences of the United States of America, *100*(17), 9923–9928.

https://doi.org/10.1073/pnas.1631635100

Clancy, D. J., & Kennington, J. (2001). A simple method to achieve consistent larval density in

bottle cultures. *Drosophila Information Service*, *84*(January), 168–169.

Dougherty, L. R., Frost, F., Maenpaa, M. I., Rowe, M., Cole, B. J., Vasudeva, R., Pottier, P.,

Schultner, E., Macartney, E. L., Lindenbaum, I., Smith, J. L., Carazo, P., Graziano, M.,

Weaving, H., Domenech, B. C., Berger, D., Meena, A., Bishop, T. R., Noble, D. W. A., …

Price, T. A. R. (2024). A systematic map of studies testing the relationship between

temperature and animal reproduction. *Ecological Solutions and Evidence*, *5*(1).

https://doi.org/DOI:%252010.1002/2688-8319.12303

Feller, G., & Gerday, C. (2003). Psychrophilic enzymes: Hot topics in cold adaptation. *Nature*

Reviews Microbiology, *1*(3), 200–208. https://doi.org/10.1038/nrmicro773

Findlay, G. D., Sitnik, J. L., Wang, W., Aquadro, C. F., Clark, N. L., & Wolfner, M. F. (2014).

Evolutionary Rate Covariation Identifies New Members of a Protein Network Required

for Drosophila melanogaster Female Post-Mating Responses. *PLoS Genetics*, *10*(1),

e1004108. https://doi.org/10.1371/journal.pgen.1004108

Flintham, E. O., Savolainen, V., & Mullon, C. (2023). Male harm offsets the demographic

benefits of good genes. *Proceedings of the National Academy of Sciences of the United*

States of America, *120*(10), 1–9. https://doi.org/10.1073/pnas.2211668120

Friedman, J., Hastie, T., & Tibshirani, R. (2010). Regularization Paths for Generalized Linear

Models via Coordinate Descent. *Journal of Statistical Software*, *33*(1).

https://doi.org/10.18637/jss.v033.i01

García-Roa, R., Chirinos, V., & Carazo, P. (2019). The ecology of sexual conflict: Temperature

variation in the social environment can drastically modulate male harm to females.

Functional Ecology, *33*(4), 681–692. https://doi.org/10.1111/1365-2435.13275

García-Roa, R., Garcia-Gonzalez, F., Noble, D. W. A., & Carazo, P. (2020). Temperature as a

modulator of sexual selection. *Biological Reviews*, *95*(6), 1607–1629.

https://doi.org/10.1111/brv.12632

Gaujoux, R., & Seoighe, C. (2010). A flexible R package for nonnegative matrix factorization.

BMC Bioinformatics, *11*. https://doi.org/10.1186/1471-2105-11-367

Gillet, L. C., Navarro, P., Tate, S., Röst, H., Selevsek, N., Reiter, L., Bonner, R., & Aebersold, R.

(2012). Targeted data extraction of the MS/MS spectra generated by data-independent

acquisition: A new concept for consistent and accurate proteome analysis. *Molecular and*

Cellular Proteomics, *11*(6), 1–17. https://doi.org/10.1074/mcp.O111.016717

Gillott, C. (2003). Male Accessory Gland Secretions: Modulators of Female Reproductive

Physiology and Behavior. *Annual Review of Entomology*, *48*(1), 163–184.

https://doi.org/10.1146/annurev.ento.48.091801.112657

Gomez-Llano, M. A., Bensch, H. M., & Svensson, E. I. (2018). Sexual conflict and ecology:

Species composition and male density interact to reduce male mating harassment and

increase female survival. *Evolution*, *72*(4), 906–915. https://doi.org/10.1111/evo.13457

Gómez-Llano, M., Faria, G. S., García-Roa, R., Noble, D. W. A., & Carazo, P. (2024). Male

harm suppresses female fitness, affecting the dynamics of adaptation and evolutionary

rescue. *Evolution Letters*, *8*(1), 149–160. https://doi.org/10.1093/evlett/qrac002

Hoffmann, A. A., Hallas, R., Anderson, A., & Telonis-Scott, M. (2005). *Evidence for a robust*

sex-specific trade-off between cold resistance and starvation resistance in Drosophila

melanogaster. *18*, 804–810. https://doi.org/10.1111/j.1420-9101.2004.00871.x

Hopkins, B. R., & Perry, J. C. (2022). The evolution of sex peptide: Sexual conflict, cooperation,

and coevolution. *Biological Reviews*, *97*(4), 1426–1448.

https://doi.org/10.1111/brv.12849

Hopkins, B. R., Sepil, I., Bonham, S., Miller, T., Charles, P. D., Fischer, R., Kessler, B. M.,

Wilson, C., & Wigby, S. (2019a). BMP signaling inhibition in Drosophila secondary cells

remodels the seminal proteome and self and rival ejaculate functions. *Proceedings of the*

National Academy of Sciences of the United States of America, *116*(49), 24719–24728.

https://doi.org/10.1073/pnas.1914491116

Hopkins, B. R., Sepil, I., Thézénas, M. L., Craig, J. F., Miller, T., Charles, P. D., Fischer, R.,

Kessler, B. M., Bretman, A., Pizzari, T., & Wigby, S. (2019b). Divergent allocation of

sperm and the seminal proteome along a competition gradient in Drosophila

melanogaster. *Proceedings of the National Academy of Sciences of the United States of*

America, *116*(36), 17925–17933. https://doi.org/10.1073/pnas.1906149116

Hopkins, B. R., Sepil, I., & Wigby, S. (2017). Seminal fluid. *Current Biology*, *27*(11), R404–

R405. https://doi.org/10.1016/j.cub.2017.03.063

Janicke, T., Häderer, I. K., Lajeunesse, M. J., & Anthes, N. (2016). Evolutionary Biology:

Darwinian sex roles confirmed across the animal kingdom. *Science Advances*, *2*(2), 1–11.

https://doi.org/10.1126/sciadv.1500983

Kuhn, M. (2008). Building Predictive Models in *R* Using the **caret** Package. *Journal of*

Statistical Software, *28*(5). https://doi.org/10.18637/jss.v028.i05

LaFlamme, B. A., Avila, F. W., Michalski, K., & Wolfner, M. F. (2014). A *Drosophila* Protease

Cascade Member, Seminal Metalloprotease-1, Is Activated Stepwise by Male Factors and

Requires Female Factors for Full Activity. *Genetics*, *196*(4), 1117–1129.

https://doi.org/10.1534/genetics.113.160101

Liu, H., & Kubli, E. (2003). Sex-peptide is the molecular basis of the sperm effect in Drosophila

melanogaster. *Proceedings of the National Academy of Sciences of the United States of*

America, *100*(17), 9929–9933. https://doi.org/10.1073/pnas.1631700100

Londoño-Nieto, C., Butler-Margalef, M., García-Roa, R., & Carazo, P. (2025). Temperature

Drives the Evolutionary Diversification of Male Harm in *Drosophila melanogaster* Flies.

Ecology Letters, *28*(3), e70102. https://doi.org/10.1111/ele.70102

Londoño-Nieto, C., García-Roa, R., Garcia-Co, C., González, P., & Carazo, P. (2023). Thermal

phenotypic plasticity of pre- and post-copulatory male harm buffers sexual conflict in

wild Drosophila melanogaster. *eLife*, *12*, e84759. https://doi.org/10.7554/eLife.84759

Machado, H. E., Bergland, A. O., Taylor, R., Tilk, S., Behrman, E., Dyer, K., Fabian, D. K., Flatt,

T., González, J., Karasov, T. L., Kim, B., Kozeretska, I., Lazzaro, B. P., Merritt, T. J.,

Pool, J. E., O'Brien, K., Rajpurohit, S., Roy, P. R., Schaeffer, S. W., … Petrov, D. A.

(2021). Broad geographic sampling reveals the shared basis and environmental correlates

of seasonal adaptation in Drosophila. *eLife*, *10*, e67577.

https://doi.org/10.7554/eLife.67577

Malek, H., & Long, T. (2019). Spatial environmental complexity mediates sexual conflict and

sexual selection in Drosophila melanogaster. *Ecology and Evolution*, *9*(5).

https://doi.org/10.1002/ece3.4932

Martinet, B., Przybyla, K., Decroo, C., Wattiez, R., & Aron, S. (2023). Proteomic differences in

seminal fluid of social insects whose sperm differ in heat tolerance. *Royal Society Open*

Science, *10*(11), 231389. https://doi.org/10.1098/rsos.231389

Martinez-Ruiz, C., & Knell, R. J. (2017). Sexual selection can both increase and decrease

extinction probability: Reconciling demographic and evolutionary factors. *Journal of*

Animal Ecology, *86*, 117–127. https://doi.org/doi:%252010.1111/1365-2656.12601

Mueller, J. L., Page, J. L., & Wolfner, M. F. (2007). An Ectopic Expression Screen Reveals the

Protective and Toxic Effects of Drosophila Seminal Fluid Proteins. *Genetics*, *175*(2),

777–783. https://doi.org/10.1534/genetics.106.065318

Parratt, S. R., Walsh, B. S., Metelmann, S., White, N., Manser, A., Bretman, A. J., Hoffmann, A.

A., Snook, R. R., & Price, T. A. R. (2021). Temperatures that sterilize males better match

global species distributions than lethal temperatures. *Nature Climate Change*, *11*(6), 481–

484. https://doi.org/10.1038/s41558-021-01047-0

Patlar, B., & Ramm, S. A. (2020). Genotype-by-environment interactions for seminal fluid

expression and sperm competitive ability. *Journal of Evolutionary Biology*, *33*(2), 225–

236. https://doi.org/10.1111/jeb.13568

Peña Jr., S. T., Stone, F., Gummow, B., Parker, A. J., & Paris, D. B. B. P. (2019). Tropical

summer induces DNA fragmentation in boar spermatozoa: Implications for evaluating

seasonal infertility. *Reproduction, Fertility and Development*, *31*(3), 590.

https://doi.org/10.1071/RD18159

Perry, J. C., Sirot, L., & Wigby, S. (2013). The seminal symphony: How to compose an ejaculate.

Trends in Ecology and Evolution, *28*(7), 414–422.

https://doi.org/10.1016/j.tree.2013.03.005

Pizzari, T., & Snook, R. R. (2003). Perspective: Sexual conflict and sexual selection: Chasing

away paradigm shifts. *Evolution*, *57*(6), 1223–1236. https://doi.org/10.1111/j.0014-

3820.2003.tb00331.x

Poiani, A. (2006). Complexity of seminal fluid: A review. *Behavioral Ecology and Sociobiology*,

60(3), 289–310. https://doi.org/10.1007/s00265-006-0178-0

R Core Team. (2022). *R: A language and environment for statistical computing*. [Computer

software]. R   Foundation for Statistical Computing, Vienna, Austria. https://www.R-

project.org/.

Ravi Ram, K., Sirot, L. K., & Wolfner, M. F. (2006). Predicted seminal astacin-like protease is

required for processing of reproductive proteins in *Drosophila melanogaster*.

Proceedings of the National Academy of Sciences, *103*(49), 18674–18679.

https://doi.org/10.1073/pnas.0606228103

Ravi Ram, K., & Wolfner, M. F. (2007). Seminal influences: Drosophila Acps and the molecular

interplay between males and females during reproduction. *Integrative and Comparative*

Biology, *47*(3), 427–445. https://doi.org/10.1093/icb/icm046

Reinhardt, K., Anthes, N., & Lange, R. (2015). Copulatory wounding and traumatic

insemination. *Cold Spring Harbor Perspectives in Biology*, *7*(5), 1–26.

https://doi.org/10.1101/cshperspect.a017582

Rohart, F., Gautier, B., Singh, A., & Lê Cao, K.-A. (2017). mixOmics: An R package for 'omics feature selection and multiple data integration. *PLOS Computational Biology*, *13*(11), e1005752. https://doi.org/10.1371/journal.pcbi.1005752

Rudman, S. M., Greenblum, S. I., Rajpurohit, S., Betancourt, N. J., Hanna, J., Tilk, S., Yokoyama, T., Petrov, D. A., & Schmidt, P. (2022). Direct observation of adaptive tracking on ecological time scales in *Drosophila*. *Science*, *375*(6586), eabj7484. https://doi.org/10.1126/science.abj7484

Sales, K., Vasudeva, R., Dickinson, M. E., Godwin, J. L., Lumley, A. J., Michalczyk, Ł., Hebberecht, L., Thomas, P., Franco, A., & Gage, M. J. G. (2018). Experimental heatwaves compromise sperm function and cause transgenerational damage in a model insect. *Nature Communications*, *9*(1), 4771. https://doi.org/10.1038/s41467-018-07273-z

Sanghvi, K., Shandilya, S., Brown, A., Todorova, B., Jahn, M., Gascoigne, S. J. L., Camilleri, T.-L., Pizzari, T., & Sepil, I. (2025). Reproductive output of old males is limited by seminal fluid, not sperm number. *Evolution Letters*, *9*(2), 282–291. https://doi.org/10.1093/evlett/qrae071

Sepil, I., Hopkins, B. R., Dean, R., Bath, E., Friedman, S., Swanson, B., Ostridge, H. J., Harper, L., Buehner, N. A., Wolfner, M. F., Konietzny, R., Thézénas, M.-L., Sandham, E., Charles, P. D., Fischer, R., Steinhauer, J., Kessler, B. M., & Wigby, S. (2020). Male reproductive aging arises via multifaceted mating-dependent sperm and seminal proteome declines, but is postponable in *Drosophila*. *Proceedings of the National Academy of Sciences*, *117*(29), 17094–17103. https://doi.org/10.1073/pnas.2009053117

Sepil, I., Hopkins, B. R., Dean, R., Thézénas, M.-L., Charles, P. D., Konietzny, R., Fischer, R., Kessler, B. M., & Wigby, S. (2019). Quantitative Proteomics Identification of Seminal

Fluid Proteins in Male Drosophila melanogaster. *Molecular & Cellular Proteomics*, *18*, S46–S58. https://doi.org/10.1074/mcp.RA118.000831

Shevchenko, A., Jensen, O. N., Podtelejnikov, A. V., Sagliocco, F., Wilm, M., Vorm, O., Mortensen, P., Shevchenko, A., Boucherie, H., & Mann, M. (1996). Linking genome and proteome by mass spectrometry: Large-scale identification of yeast proteins from two dimensional gels. *Proceedings of the National Academy of Sciences of the United States of America*, *93*(25), 14440–14445. https://doi.org/10.1073/pnas.93.25.14440

Sirot, L. K., Buehner, N. A., Fiumera, A. C., & Wolfner, M. F. (2009). Seminal fluid protein depletion and replenishment in the fruit fly, Drosophila melanogaster: An ELISA-based method for tracking individual ejaculates. *Behavioral Ecology and Sociobiology*, *63*(10), 1505–1513. https://doi.org/10.1007/s00265-009-0806-6

Sirot, L. K., Wolfner, M. F., & Wigby, S. (2011). Protein-specific manipulation of ejaculate composition in response to female mating status in Drosophila melanogaster. *Proceedings of the National Academy of Sciences of the United States of America*, *108*(24), 9922–9926. https://doi.org/10.1073/pnas.1100905108

Sirot, L. K., Wong, A., Chapman, T., & Wolfner, M. F. (2015). Sexual Conflict and Seminal Fluid Proteins: A Dynamic Landscape of Sexual Interactions. *Cold Spring Harbor Perspectives in Biology*, *7*(2), a017533. https://doi.org/10.1101/cshperspect.a017533

Soller, M., Bownes, M., & Kubli, E. (1999). Control of Oocyte Maturation in Sexually MatureDrosophilaFemales. *Developmental Biology*, *208*(2), 337–351. https://doi.org/10.1006/dbio.1999.9210

Van Heerwaarden, B., & Sgrò, C. M. (2021). Male fertility thermal limits predict vulnerability to

climate warming. *Nature Communications*, *12*(1), 2214. https://doi.org/10.1038/s41467-

021-22546-w

Vasudeva, R., Deeming, D. C., & Eady, P. E. (2014). Developmental temperature affects the

expression of ejaculatory traits and the outcome of sperm competition in *Callosobruchus*

maculatus. *Journal of Evolutionary Biology*, *27*(9), 1811–1818.

https://doi.org/10.1111/jeb.12431

Von Hellfeld, R., Konietzny, R., Charles, P. D., Fischer, R., Kessler, B. M., Wigby, S., Sepil, I., &

Morimoto, J. (2025). *The developmental environment mediates adult seminal proteome*

allocation in male Drosophila melanogaster. Evolutionary Biology.

https://doi.org/10.1101/2025.01.09.632097

Wang, W. W.-Y., & Gunderson, A. R. (2022). The Physiological and Evolutionary Ecology of

Sperm Thermal Performance. *Frontiers in Physiology*, *13*, 754830.

https://doi.org/10.3389/fphys.2022.754830

Wedell, N., Gage, M. J. G., & Parker, G. A. (2002). Sperm competition, male prudence and

sperm-limited females. *TREE*, *17*(7), 313–320.

Wigby, S., Brown, N. C., Allen, S. E., Misra, S., Sitnik, J. L., Sepil, I., Clark, A. G., & Wolfner,

M. F. (2020). The Drosophila seminal proteome and its role in postcopulatory sexual

selection. *Philosophical Transactions of the Royal Society of London B Biological*

Sciences, 20200072.

Wigby, S., & Chapman, T. (2005). Sex peptide causes mating costs in female Drosophila

melanogaster. *Current Biology*, *15*(4), 316–321.

Wigby, S., Perry, J. C., Kim, Y., & Sirot, L. K. (2016). Developmental environment mediates

637   male seminal protein investment in *D rosophila melanogaster* . *Functional Ecology*,

638   *30*(3), 410–419. https://doi.org/10.1111/1365-2435.12515

639   Wigby, S., Sirot, L. K., Linklater, J. R., Buehner, N., Calboli, F. C. F., Bretman, A., Wolfner, M.

640   F., & Chapman, T. (2009). Seminal Fluid Protein Allocation and Male Reproductive

641   Success. *Current Biology*, *19*(9), 751–757. https://doi.org/10.1016/j.cub.2009.03.036

642   Yun, L., Chen, P. J., Kwok, K. E., Angell, C. S., Rundle, H. D., & Agrawal, A. F. (2018).

643   Competition for mates and the improvement of nonsexual fitness. *Proceedings of the*

644   *National Academy of Sciences of the United States of America*, *115*(26), 6762–6767.

645   https://doi.org/10.1073/pnas.1805435115

646   Yun, L., Chen, P. J., Singh, A., Agrawal, A. F., & Rundle, H. D. (2017). The physical

647   environment mediates male harm and its effect on selection in females. *Proceedings of*

648   *the Royal Society B: Biological Sciences*. https://doi.org/10.1098/rspb.2017.0424

649   Zheng, J., Guo, N., Huang, Y., Guo, X., & Wagner, A. (2024). High temperature delays and low

650   temperature accelerates evolution of a new protein phenotype. *Nature Communications*,

651   *15*(1), 2495. https://doi.org/10.1038/s41467-024-46332-6

652   Zou, H., & Hastie, T. (2005). Regularization and Variable Selection Via the Elastic Net. *Journal*

653   *of the Royal Statistical Society Series B: Statistical Methodology*, *67*(2), 301–320.

654   https://doi.org/10.1111/j.1467-9868.2005.00503.x

Figure 1 | Effect of temperature on SFPs production and transfer (short-term exposure / experiment 1). A) Heatmap showing the abundance of 145 SFPs detected in accessory gland samples. Elastic net regression selected 74 (under no-SCR) and 3 (under High-SCR) out of 145 SFPs as predictors with a strong effect differentiating proteomic responses across temperature treatments. Each cell gives the across-biological replicate mean for that protein, at each temperature treatment. Row annotations indicate SFPs with strong response to temperature in each SCR level and include functional information relating to protein functions as part of the sex peptide or ovulin networks, or other known roles in sperm competition. B) Partial least-squares discriminant analysis (PLS-DA) plot of proteins production for no-SCR treatment and C) PLS-DA plot of proteins production for high-SCR treatment. Points represent biological replicates. Ellipses denote variability among replicates. D) Heatmap showing the inferred relative abundance of 145 SFPs transferred to females during mating. Elastic net regression selected 79 (no SCR treatment) and 6 (High SCR treatment) out of 145 SFPs as predictors with strong differential transfer response to temperature. Each cell shows the mean ratio of protein abundance in virgin males relative to recently mated males, across biological replicates, for each temperature treatment, reflecting the relative loss of protein upon mating. Values closer to 1 indicate proteins reduced after mating (i.e., transferred). Heatmaps were generated from log2-transformed protein abundances for better visualization. E) PLS- DA plot of proteins transfer for no-SCR treatment and F) PLS-DA plot of proteins transfer for high-SCR treatment. Points in each PLS-DA plot represent biological replicates. Ellipses denote variability among replicates.

Figure 2 | Effect of temperature on SFPs production and transfer (long-term exposure / experiment 2). A) Heatmap showing the abundance of the 145 SFPs detected in the accessory gland samples. Elastic net regression selected 39 (no SCR treatment) and 68 (High SCR treatment) out of 145 SFPs as predictors with a strong effect differentiating proteomic responses across temperature treatments. Each cell gives the across-biological replicate mean for that protein, at each temperature treatment. Row annotations indicate SFPs with strong response to temperature in each SCR level and include functional information relating to protein functions as part of the sex peptide or ovulin networks, or other known roles in sperm competition. B) PLS-DA plot of proteins production for no-SCR treatment and C) PLS- DA plot of proteins production for high-SCR treatment. Points represent biological replicates. Ellipses denote variability among replicates. D) Heatmap showing the inferred relative abundance of 145 SFPs transferred to females during mating. Elastic net regression selected 49 (no SCR treatment) and 46 (High SCR treatment) out of 145 SFPs as predictors with strong differential transfer response to temperature. Each cell shows the mean ratio of protein abundance in unmated males relative to recently mated males, across biological replicates, for each temperature treatment, reflecting the relative loss of protein upon mating. Values closer to 1 indicate proteins reduced after mating (i.e., transferred). Heatmaps were generated from log2-transformed protein abundances for better visualization. E) PLS- DA plot of proteins transfer for no-SCR treatment and F) PLS-DA plot of proteins transfer for high-SCR treatment. Points in each PLS-DA plot represent biological replicates. Ellipses denote variability among replicates.

Figure 3 | Effect of temperature on production and transfer of SFPs involved in Ovulin and
SP networks. A – B) Relative abundance production and transfer profiles of SPFs associated with
the SP and ovulin networks across different temperatures and SCR levels for short-term exposure
(experiment 1). C – D) Relative abundance production and transfer profiles of SPFs associated
with the SP and ovulin networks across different temperatures and SCR levels for long-term
exposure (experiment 2). Each faint line represents the mean across the 3 replicates for an
individual SFP within each treatment combination. Abundance values were normalized by mean-
centering and averaged across replicates. Blue points represent the average (± s.e.) of SPFs in the
SP network, across temperature and SCR conditions. Red points represent the average (± s.e.) of
SFPs in the ovulin network, across temperature and SCR conditions.

Figure 4 | Comparison of SFPs production and transfer under high-SCR vs. no-SCR across
temperature treatments (short-term exposure / experiment 1). A) Comparison for production.
B) Comparison for transfer. Each cell shows the mean ratio of protein absolute abundance under
high-SCR relative to no-SCR across biological replicates for each temperature. Values closer to 1
indicate higher production/transfer under high-SCR conditions. Row annotations indicate
functional information relating to protein functions as part of the sex peptide or ovulin networks,
or other known roles in sperm competition.

Figure 5 | **Comparison of SFPs production and transfer under high-SCR vs. no-SCR across temperature treatments (long-term exposure / experiment 2).** A) Comparison for production. B) Comparison for transfer. Each cell shows the mean ratio of protein absolute abundance under high-SCR relative to no-SCR across biological replicates for each temperature. Values closer to 1 indicate higher production/transfer under high-SCR conditions. Row annotations indicate functional information relating to protein functions as part of the sex peptide or ovulin networks, or other known roles in sperm competition.

SFPs' PRODUCTION

SFPs' TRANSFER

48-hour treatment duration (Exp. 1)

13 days treatment duration (Exp. 2)

Network
- Ovulin
- SP

A   SFPs' PRODUCTION

B   SFPs' TRANSFER

SPnetwork
OvulinNetwork
PostMating Baheviour / SpermCompetition
Other