Beta bursts mediate amygdala gating of hippocampal emotional encoding

Yajun Zhou[1†], Yuqing Huang[1†], Giana Pittaro[1], Behzad Zareian[1], Ayman Aljishi[1], Alondra P. Toro[1], Zion Zibly[1], Sami Obaid[1], Adithya Sivaraju[2], Ilan Harpaz-Rotem[3,4], Alfred Kaye[3,4,7], John Krystal[3,4,5,7,8], Kevin Sheth[2], Xiaosi Gu[3], Christopher Pittenger[3,4,6,8], Eyiyemisi C. Damisah[1,4,5,6*]

[1]Department of Neurosurgery, Yale University, New Haven, CT, USA.
[2]Department of Neurology, Yale University, New Haven, CT, USA.
[3]Department of Psychiatry, Yale University, New Haven, CT, USA.
[4]Wu Tsai Institute, Yale University, New Haven, CT, USA.
[5]Department of Neuroscience, Yale University, New Haven, CT, USA.
[6]Center for Brain and Mind Health, Yale University, CT, USA.
[7]Clinical Neurosciences Division, VA National Center for PTSD, USA.
[8]Department of Psychology, Yale University, New Haven, CT, USA.

*Corresponding author(s). E-mail(s): eyiyemisi.damisah@yale.edu;
[†]These authors contributed equally to this work.

Abstract

The amygdala and hippocampus are central to emotional processing, yet the transient neural dynamics coordinating these regions remain unclear. We simultaneously recorded single-neuron activity and local field potentials from both regions in epilepsy patients during an emotional image-rating task. Neurons in both regions responded to images with firing rate changes that predicted subjective ratings of extreme pleasantness or unpleasantness. To examine the underlying oscillatory mechanisms, we analyzed beta bursts (13–30 Hz)—transient, high-power events—since conventional spectral analyses revealed no valence-specific patterns. Beta bursts were associated with increased gamma amplitude and enhanced phase coherence in both structures, with beta-gamma phase-amplitude coupling capturing emotion-related dynamics. Critically, amygdala beta bursts strongly suppressed hippocampal firing through interneuron activation during negative valence processing, whereas hippocampal bursts showed no reciprocal influence. These findings suggest that beta bursts provide a temporal code for

emotion and represent a candidate mechanism for targeted neuromodulation in mood disorders.

Keywords: Emotional Processing, Beta Bursts, Amygdala-Hippocampus Circuit, Single-Neuron Activity, LFP

1 Introduction

Emotion assessment and modulation are critical for human cognitive and executive functions. The amygdala, a key region for evaluating emotionally salient stimuli, communicates extensively with the hippocampus, which provides contextual information to support the encoding and modulation of emotional experiences[1–4]. Through reciprocal connections, these structures prioritize emotionally salient information and shape future responses to affective stimuli [1, 5–8]. Specifically, the bidirectional circuit between the basolateral amygdala (BLA) and the ventral hippocampus (vHPC) has been highlighted in various emotion-related tasks, especially for negative emotion-related behaviors [9–12]. Disruption of the amygdala-hippocampal circuit is seen in various neuro-psychiatric conditions, including major depression, posttraumatic stress disorder, and pathological anxiety—affecting millions of individuals worldwide [9, 10, 13–22]. However, the temporal dynamics that coordinate rapid amygdala–hippocampus interactions at the single-neuron level in humans during emotional processing remain poorly understood.

Recent studies have highlighted the importance of oscillations in modulating neuronal activity [23–25]. In particular, beta oscillations (13–30 Hz) have been found to mediate long-range, state-dependent signaling for movement control, sensorimotor integration, working memory, and decision making [26–29]. However, there are contradictory findings regarding the role of beta oscillations in emotion processing. Although some human EEG, MEG, and fMRI studies report that beta power decreases during emotional processing [30–35], others find that beta power increases, particularly during states of negative affect or anxiety [36–38]. These inconsistencies may be due to differences in species, methods, and behavioral paradigms. It is also possible that spectral analyses that average neural activity over extended time windows could obscure millisecond-scale dynamics relevant for emotional processing. [39, 40].

Beta oscillations occur not as sustained rhythms but as bursts—transient, high-power events that coordinate neural activity with millisecond precision [40–43]. In sensorimotor cortices, human EEG and non-human primate intracranial studies showed that individual beta bursts predict behavior better than averaged beta power [44–46]. In prefrontal networks, intracranial studies in non-human primates indicate that burst timing shapes information routing across regions [39, 47]. Most importantly, in rodent anxiety models, 25 millisecond shifts in beta coherence timing in the amygdala-hippocampus circuit can transform anxious behavior into calm exploration [18]. These findings suggest that emotion processing may depend on the precise timing of transient events, which may be obscured by analyzing the broadband oscillatory power.

We hypothesized that transient beta bursts coordinate amygdala-hippocampus communication during human emotional processing. To test this, we simultaneously recorded single neuron activity and local field potentials from the amygdala and hippocampus in human subjects performing an affective image rating task. Simultaneous recording allowed us to examine how population-level oscillatory events relate to the timing of neuronal spiking, providing insight into the circuit and neuron-level mechanisms of emotional processing in humans. We found that beta bursts modulate high gamma (HG) amplitude and single-neuron firing rates differentially across emotional conditions with amygdala bursts gating hippocampal spiking activity during negative emotions. These findings suggest a temporal code for emotion appraisal, which may serve as a neuronal biomarker and potential therapeutic targets for neuromodulation.

2 Results

2.1 Experimental design and behavioral responses

To investigate the neural mechanisms underpinning the coding of emotional valence in humans, we recorded LFPs and single neuron activity from the hippocampus and amygdala of 15 epilepsy patients. Patient demographics including age sex, handedness, anti-seizure medications, Beck Anxiety Inventory (BAI) and Beck Depression Inventory (BDI) are listed. Of note the average BAI score was 15.25 ± 3.42 (mean $\pm$ SEM), suggestive of mild anxiety, and average BDI score was 14.63 ± 3.04 (mean $\pm$ SEM), indicating mild depression (see Methods for inventory scales, Fig. 1a, Table S1 and Table S2).

Each electrode consisted of macro and microcontacts for parallel recordings of the field and unit action potentials throughout the amygdala (n = 29) and hippocampus (n = 46). During the recordings, participants rated images [48][49] on a 5-point valence scale from 1 (very pleasant, VP) to 5 (very unpleasant, VU) following a 3-second presentations (Fig. 1b). Across participants, scores were distributed across all valence categories, with neutral responses being the most common (30.4% $\pm$ 12.1%) and extreme scores least frequent (very pleasant: 20.3% $\pm$ 10.0%; very unpleasant: 16.3% $\pm$ 10.8%; Fig. 1c). Semantic categories influenced these distributions, as objects and humans elicited primarily neutral ratings, whereas animals and scenes evoked more positive appraisals (Fig. S1a). Stimulus categories exhibited different nonuniform rating distribution (aligned ranks transformation (ART) ANOVA, animal: $F(4, 90) = 6.76, p = 8.34 \times 10^{-5}$, human: $F(4, 90) = 6.29, p = 1.64 \times 10^{-4}$, scene: $F(4, 90) = 15.89, p = 7.11 \times 10^{-10}$, object: $F(4, 95) = 36.62, p = 2.22 \times 10^{-16}$). Permutation test with multiple comparisons correction by FDR revealed that participants rated objects ($p < 0.01$ for 3 vs. all others) and humans ($p < 0.05$ for 3 vs. all others) more frequently as neutral, and rated animals ($p = 0.005$ for 1 vs. 4 and $p < 0.001$ for 1 vs. 5) and scenes ($p < 0.001$ for 1 vs. all others) more frequetly as very pleasant.

Response times (RT) revealed a behavioral correlate of emotional processing. We observed an inverted-U pattern ($F_{(4,1958)} = 10.01$, $p = 5.15 \times 10^{-8}$), with significantly faster responses for emotional extremes compared to moderate ratings (very pleasant vs. moderate: all $p < 0.001$; very unpleasant vs. moderate: all $p < 0.01$;

Fig. 1d). In contrast, we found no significant difference in RT across semantic categories ($F(3, 1970) = 0.97, p = 0.41$, Fig S1b). This inverted U pattern suggests faster processing of high-valence stimuli. This effect appears to be independent of arousal, as pupil responses did not differ across conditions (Fig. 1e, ART ANOVA, $p > 0.05$), and across semantic categories (Fig. S1c, ART ANOVA, $p > 0.05$). These findings suggest that emotional extremes receive preferential processing, providing a behavioral benchmark for neural analyzes.

2.2 Amygdala and hippocampal neurons encode emotional valence with graded selectivity

To investigate how individual neurons in the amygdala and hippocampus process emotional information, we computed trial-wise peristimulus time histograms (PSTH) for each subjective emotional rating. We identified 218 neurons across amygdala and hippocampus with valence-specific firing patterns (Fig. 2a, Table S3). 65.1% (54/82) of neurons in the amygdala and 59.6% (81/136) in the hippocampus significantly modulated their firing rates to emotional stimuli (Fig. 2b). Many emotion-responsive neurons in both regions exhibited elevated firing for specific emotional ratings compared to others, with firing rates diverging significantly by 500-800 ms post-stimulus (cluster-based permutation t-test, $p < 0.05$, Fig. 2c-d). While most responsive neurons were selective for single emotional categories, we observed a subset (38.89% in the amygdala and 36.59% in the hippocampus) that responded to adjacent ratings, suggesting a graded rather than categorical representation of valence (Fig. 2b). This partial overlap was more pronounced for moderate emotional ratings than for extreme ratings.

To evaluate how valence modulated neuronal firing, we fitted a population-level GLM to the firing of valence-responsive neurons. In amygdala very unpleasant-responsive neurons (Fig. 2e), baseline firing rate was a strong positive predictor of the firing rate in the task period ($\beta = 0.401$, $p = 2.44 \times 10^{-24}$). The firing rates of these neurons in the very unpleasant condition were higher than those for all other emotional conditions(very pleasant: $\beta = -0.090$, $p = 5.91 \times 10^{-6}$; pleasant: $\beta = -0.115$, $p = 8.54 \times 10^{-9}$; neutral: $\beta = -0.099$, $p = 2.97 \times 10^{-7}$; unpleasant: $\beta = -0.074$, $p = 6.31 \times 10^{-4}$). In hippocampal very pleasant-responsive neurons (Fig. 2f), baseline firing rate again positively predicted the firing rate in the task period ($\beta = 0.381$, $p = 7.73 \times 10^{-39}$). The firing rates of these neurons in the very pleasant condition were higher than those for all other emotional ratings (pleasant: $\beta = -0.049$, $p = 2.96 \times 10^{-4}$; neutral: $\beta = -0.040$, $p = 1.85 \times 10^{-3}$; unpleasant: $\beta = -0.038$, $p = 0.012$; very unpleasant: $\beta = -0.034$, $p = 0.024$). These GLM analyses indicate that baseline firing rate is a strong positive predictor of task-period activity, and support the conclusion that neuronal responses within each region are modulated by emotional valence.

2.3 From sustained power to transient bursts-uncovering emotion-specific dynamics

Having established the existence of valence-specific neuronal populations, we examined whether neural oscillatory dynamics coordinated these responses. Time-frequency decomposition showed stimulus-evoked changes—beta power (13-30 Hz) decreased while gamma power (> 40 Hz) increased following image onset (cluster-based permutation, $p < 0.05$; Fig. 3a-b, top). However, these power modulations were identical across emotional conditions in both amygdala or hippocampus (ART ANOVA, all $p > 0.05$; Fig. 3a-b, bottom), indicating that sustained beta and gamma power alone do not distinguish subjective emotional appraisal.

We therefore turned to transient oscillatory beta burst dynamics [40, 50, 51]. First, we extracted brief high-power beta bursts (see example burst in Fig. 3c) from macroelectrode LFP recordings in both regions. Across all participants, we identified 13,260 bursts (5,097 in amygdala, 8,163 in hippocampus). In the hippocampus, burst duration, rate, and peak frequency showed no significant main effect of valence (all $p > 0.05$). In the amygdala, burst duration and rate were also not significantly modulated by valence (all $p > 0.05$). However, peak frequency differed significantly across emotional ratings ($F(4, 180) = 4.21, p = 0.0028$). Stimuli rated as neutral (20.83 ± 1.36 Hz) were associated with higher peak frequencies than unpleasant ones (unpleasant: 19.83 ± 1.52 Hz, $p = 0.037$, very unpleasant: 19.65 ± 2.22 Hz, $p = 0.009$), with no other pairwise differences surviving after multiple comparisons correction using FDR. These results suggest that while the occurrence and duration of beta bursts were insensitive to valence, spectral characteristics in the amygdala may encode specific aspects of emotional appraisal.

2.4 Amygdala beta bursts gate hippocampal activity during negative emotions

To better understand the function of beta bursts, we examined their impact on local and cross-regional neural dynamics by analyzing HG and single neuron spiking activity during bursts. Beta bursts triggered robust high-gamma increases (80-140 Hz) compared to surrogate events in both regions ($p < 0.05$; Fig. 4a-b, top). We then asked whether HG amplitude was associated with valence ratings. In the amygdala, we found no significant change in burst associated HG amplitude across emotional conditions (Fig. 4a, bottom). However, the very pleasant condition exhibited a higher HG compared to moderate conditions following bursts (ART ANOVA followed by cluster based permutation testing, $p < 0.05$). This finding is concordant with increased firing rates of the very pleasant responsive neurons in the hippocampus. Together, these results suggest that beta bursts may play a role in the coordination of local neuronal populations.

This increase in gamma during bursts coincided with spiking activity changes in the hippocampus but not in the amygdala. Specifically, the amygdala neurons did not show significant changes in firing rates after the onset of their own bursts ($p < 0.05$; Fig. 4e, top left; $p > 0.05$), while the hippocampal neurons exhibited suppression of the firing rate after the onset of beta bursts. (Fig. 4f, top left; $p < 0.05$). However, the burst-firing

relationship was emotion-specific. During very unpleasant ratings, amygdala neurons showed enhanced preburst activity followed by suppression during bursts (Fig. 4e, top right; $p < 0.05$). Hippocampal neurons also exhibited earlier, deeper inhibition to very unpleasant ratings around bursts(Fig. 4f, top right; $p < 0.05$). This pattern may reflect selective modulation of different neuronal subpopulations in each region.

Furthermore, we observed an asymmetric valence-dependent manner of cross-regional interaction during bursts. Amygdala beta bursts potently suppressed hippocampal firing by 80.96%, with maximal effect during very unpleasant ratings. (Fig. 4e, bottom; $p < 0.05$). However, hippocampal bursts exerted no reciprocal influence on amygdala single neuron activity (Fig. 4f, bottom; $p > 0.05$).

To better understand this relationship at the cellular level, we analyzed cell-type contributions to firing rate changes during bursts in each region. In the hippocampus, pyramidal neurons drove the firing rate decrease observed during very unpleasant ratings while narrow and wide hippocampal interneurons did not exhibit any firing rate changes across valence conditions (Fig. 5b; $p < 0.05$). Further, amyygdala beta bursts suppressed hippocampal interneuron activity (narrow and wide) during very unpleasant trials, but not the activity of pyramidal cells (Fig. 5c; $p < 0.05$). We did not have enough amygdala interneurons (n=4) to conduct the same analysis. These results suggest that this asymmetric relationship may reflect a hierarchical mechanism in emotional processing: the amygdala may gate hippocampal processing specifically during very negative emotional states.

2.5 Beta-gamma coupling distinguishes emotional valence through phase-specific modulation

Despite the absence of differences in spectral power between emotional ratings, we found that discrete time-locked beta bursts modulate neuronal activity within and across regions during emotional processing. We then asked if phase-specific neural modulation might account for differences in emotional appraisal at the LFP level. To do this, we first implemented inter trial phase coherence during bursts. We found that the timing and magnitude of burst-locked dynamics varied with emotional valence. Very unpleasant stimuli elicited stronger beta phase-locking in both the hippocampus and the amygdala, indicated by the increase in intertrial phase coherence (ITPC) (ART ANOVA test with multiple comparisons FDR correction $p < 0.05$; pairwise cluster-based permutation t test, $p < 0.05$; Fig. 4c-d).

Next, we quantified phase-amplitude coupling (PAC) between beta-phase and gamma-amplitude using a modulation index (MI) approach [52, 53]. While sustained beta power showed no differences across emotional conditions, PAC revealed robust emotion-specific modulation (Fig. 6a). The beta phase at which gamma amplitude peaked differed across emotional ratings (Pairwise Mardia-Watson-Wheeler tests with multiple comparisons FDR correction, $p < 0.01$ for all conditions, Fig. 6b) with each emotional condition exhibiting a preferred beta phase (Rayleigh test, $p < 0.01$ for all conditions) in both regions. However, PAC strength, as measured by MI, was highest during very unpleasant trials in both amygdala and hippocampus (Fig. 6c). This emotion-specific coupling decreased systematically across the beta band (ART ANOVA test with FDR correction $p < 0.05$; pairwise cluster-based permutation t test,

$p < 0.05$; Fig. 6d). This suggests that emotional valence may not only be encoded by precise temporal coordination between oscillatory rhythms (PAC) but critically by coupling strength (MI)).

3 Discussion

Our results suggest a new mechanism for human emotional processing: transient beta bursts that coordinate amygdala-hippocampus communication through millisecond-precision temporal dynamics. A subset of neurons were found in both the amygdala and hippocampus that displayed responsiveness to emotional stimuli, with firing rate differed across subjective emotional valence ratings. While the average spectral power in the beta and gamma bands did not discriminate among emotional conditions, beta burst dynamics revealed robust emotion-specific modulation across scales—from single-neuron firing to cross-regional communication. Furthermore, emotional valence may be encoded through systematic phase relationships within these bursts rather than through power changes, suggesting a temporal code for emotion.

We believe that our findings enhance the understanding of the neural mechanisms underlying emotional processing by providing converging evidence from single-unit and oscillatory activity in the amygdala and hippocampus. Prior works suggest the amygdala rapidly encodes emotional valence [54–56]. Consistent with this, we identified amygdala neurons sensitive to pleasant and unpleasant stimuli, replicating prior intracranial studies (e.g.[57, 58]). We also found that hippocampal neurons, mostly from the anterior hippocampus, showed similar valence selectivity, aligning with recent evidence that the ventral hippocampus contributes to emotion in rodents [18, 59]. Beyond spiking activity, oscillatory dynamics also play a role: prior studies have linked beta [17, 60] and gamma [61, 62] oscillations to emotional evaluation and salience encoding. Our time-frequency decomposition revealed similar spectral features, however, these were insufficient to distinguish emotional valence appraisals, echoing concerns that sustained oscillations may overlook key dynamic features of emotional processing [63, 64]. Instead, transient beta bursts offered a more precise neural marker, aligning with recent paradigm in a rodent study [18], suggesting that transient beta bursts may reflect a key neurocomputational process on emotion.

Building on this, our findings offer new insight into the functional role of beta bursts in affective encoding within the human MTL. Emerging evidence suggests oscillatory bursts—rather than sustained rhythms—play a key role in modulating cognitive and emotional functions [65, 66]. Beta bursts, in particular, have been implicated in routing and gating information during cognitive tasks [40, 67]. Rodent and nonhuman primate studies show that beta bursts within hippocampus and amygdala circuits regulate attention states [41] and memory encoding [18]. Extending this, we found that beta bursts in the amygdala and hippocampus were modulated by emotional stimuli and linked to transient changes in spiking activity and gamma amplitudes. These bursts provided brief windows in which affective information was selectively enhanced, especially for highly pleasant or unpleasant stimuli. Neuronal responsiveness was stronger during burst windows, suggesting that beta bursts may coordinate ensemble-level encoding of emotional valence. Moreover, amygdala bursts exhibited

a different peak frequency for neutral versus negative valence ratings, while the hippocampus lacked such valence-specific burst modulation (Fig. 3d). This suggests that there might be differences in regional specialization across the hippocampus and amygdala. Beta bursts also showed robust cross-frequency coupling (CFC) with gamma, supporting hierarchical organization across frequency bands during emotion-related tasks [18, 40, 42, 65].

The amygdala and hippocampus form a tightly interconnected circuit critical for emotional processing and affective memory. Anatomically, these regions share reciprocal projections: the basolateral amygdala projects to dorsal and ventral hippocampus [17, 54], while the hippocampus sends input back via the subiculum [43]. Functionally, this circuitry is crucial for integrating emotional information with contextual memory, enabling behavioral responses and long-term memory encoding. For example, fMRI studies show increased amygdala-hippocampus activation during emotional memory and aversive recall [68–70]. Meanwhile, human electrophysiology confirms dynamic coordination between these regions during emotion processing. Zheng et al. [64] demonstrated theta–alpha multiplexing between amygdala and hippocampus, revealing frequency-specific communication for affective salience. The beta and gamma rhythms in both regions also play key roles. For instance, Costa et al. [61] and Qasim et al. [20] reported enhanced synchrony and ripple coactivity during aversive memory tasks. Our results align with this view, showing that beta burst coordination can temporally constrain amygdala-hippocampus communication, reflecting emotional processing rather than general arousal. We observed that beta bursts not only modulated local gamma amplitude and neuronal firing within the amygdala and hippocampus but also exerted cross-structural effects. Specifically, beta bursts of the amygdala may influence the firing rates in the ipsilateral hippocampus, but not vice versa. Cell-type analysis further supported the circuit understanding: that the suppressed firing rate seen in the hippocampus during the amygdala burst was driven by hippocampal interneurons and not by pyramidal cells. (Fig. 5c), This finding aligns with rodent studies showing that somatostatin interneurons modulate sychronization between these two regions [18]. The asymmetric, burst-triggered coordination supports a dynamic, state-dependent model of emotional coding, where transient oscillatory events shape communication between amygdala and hippocampus [17, 71].

Our ITPC and PAC analyses suggest that phase relationships may be better than oscillatory power in distinguishing emotional states. Both the amygdala and hippocampus exhibited enhanced inter trial phase coherence (Fig. 4c-d) specifically for very negative valence ratings. Furthermore, while the preferred phase for maximal gamma amplitude varied across valence conditions, the coupling strength was remarkably specific with very negative valence ratings showing increased coupling strength. Together, these results suggest that phase modulation may underlie selective enhancement and temporal coordination of affective information processing, especially for strong negative emotions.

Aberrant amygdala-hippocampus dynamics has been implicated in psychiatric conditions such as major depressive disorder, PTSD, and generalized anxiety [55, 56, 72, 73]. Disruptions in oscillatory coordination—particularly in beta and gamma bands—and cross-regional communication may contribute to maladaptive emotional

encoding and recall. Our findings suggest a potential mechanism for emotional processing and offer translational relevance for neuropsychiatric interventions. Specifically, beta bursts in amygdala and hippocampus, by modulating local spiking and gamma activity, may serve as real-time physiological markers of emotional states. This temporally precise neural motif is especially relevant for closed-loop neuromodulation strategies, including deep brain stimulation, transcranial alternating current stimulation, and transcranial magnetic stimulation [74].

Rodent studies have shown that manipulating somatostatin interneurons during beta coherence bursts can influence anxiety-related behavior [18]. In humans, high-frequency activity (HFA)—a proxy for local spiking [75] in MTL—has been used to modulate emotional recall [20] and alleviate depressive symptoms via personalized closed-loop stimulation [76]. By identifying beta bursts as a modifiable and behaviorally relevant neural event, our results suggest that this transient, dynamic view of emotional processing may open new avenues for understanding and treating disorders of emotional regulation.

Several limitations merit consideration. First, recordings from epilepsy patients may not generalize to healthy populations. Our participant cohort has a heterogeneous clinical background, encompassing different psychiatric and mood disorder histories as well as treatment with anti-seizure medications. Second, while our task effectively elicited emotional responses, naturalistic emotional experiences involve additional complexity that our task may not account for. Third, the correlational nature of our findings precludes causal inference—whether beta bursts actively shape emotional processing remains unresolved without direct manipulations such as burst-inducing stimulation as mentioned above. Fourth, limited arousal measurements (pupil data from 8 participants including 6 healthy controls) constrain valence-arousal dissociation. Although we focused on beta bursts due to their alignment with emotional stimuli, transient activity in other bands and their interaction with beta, such as through cross-frequency coupling may also contribute to emotional processing. Lastly we acknowledge that our study is purely mechanistic and thus claims as to its therapeutic potential without extensive clinical trial studies to support this hypothesis. Furthermore our cohort study was only evaluated using a limited number of tests including the BDI and BAI, which may not capture the nuances of psychiatric disease. Future studies should employ causal interventions, include clinical populations, incorporate comprehensive autonomic measures, and examine multi-band burst interactions. Nevertheless, by revealing how transient beta bursts coordinate emotional processing through asymmetric control and phase-specific encoding, our findings establish a new framework for understanding and potentially treating emotional dysfunction.

4 Methods

4.1 Human participants

Fifteen participants (9 female, 6 male; age: 34.13 ± 12.15 years) with pharmacologically intractable epilepsy were enrolled at Yale New Haven Hospital. All procedures

were approved by the Yale University Institutional Review Board and participants provided with written informed consent. Recordings occurred during clinical monitoring with depth electrodes placed solely for clinical purposes. Channels with epileptiform activity (IED) were excluded using automated IED detection and manual inspection. Data within 4 hours of clinical or electrographic seizures were excluded. Clinical characteristics (Table S1), including age, handedness, sex, anti-seizure medications, Beck Anxiety Inventory (BAI, [77])and Beck Depression Inventory (BDI, [78]) scores, are provided in Supplementary Table 1. BAI and BDI scores were obtained from each participant's routine presurgical neuropsychological assessment.

4.2 Electrophysiological recording

Participants underwent stereotactic implantation of Behnke-Fried depth electrodes (Ad-Tech Medical) containing eight 40-µm microwires protruding 3-4 mm from the macroelectrode tip, enabling simultaneous recording of LFPs and single-unit activity from the amygdala and hippocampus (Fig. 1a). Microwire signals were digitized at 30 kHz (0.3-7.5 kHz bandpass) using a NeuroPort system (Blackrock Microsystems). Macroelectrode signals were sampled at 2 kHz via the clinical EEG system [79, 80].

4.3 Electrode localization

Pre-implantation T1-weighted MRI scans were co-registered with post-implantation CT scans. Electrode positions were identified using LeGUI [81] and mapped to standardized Montreal Neurological Institute (MNI) space (Table S2). Gray and white matter boundaries were automatically segmented to exclude contacts residing in white matter. Regional assignments used the Neuromorphometrics atlas [82]. For visualization, we plotted the MNI coordinates of all macroelectrodes across participants on the 152 template brain from the MNI [83, 84].

4.4 Behavioral task

Participants viewed 120 color photographs while seated in hospital beds. Visual stimuli were presented on a 24-inch monitor (2K, 120 Hz) using MATLAB and Psychtoolbox. Latencies between stimulus onsets and TTL pulses were logged and later compensated for in neural analyses. Participants were provided with a Gamepad or keyboard for ratings. Each trial consisted of fixation (1 s), image presentation (3 s), and rating period (maximum 5 s). The participants rated the images from 1 (very pleasant) to 5 (very unpleasant), with moderate values corresponding to pleasant (2), neutral (3), and unpleasant (4) (Fig. 1b). Trials with incorrect or absent responses were excluded from behavioral and neural analyzes. The stimuli consisted of emotionally evocative images selected from two validated datasets: the International Affective Picture System (IAPS) for 8 patients and the Nencki Affective Picture System (NAPS) for 8 patients, which was used to match visual properties across categories (e.g., luminance, contrast, spatial frequency). Both datasets included an equal number of images from four semantic categories: scenes, objects, animals, and humans with 30 images per category, ensuring consistent visual diversity and balanced emotional content. All images were presented at their original size and were novel to the participants at the time of

testing. Verbal instructions were given prior to the task. Eye tracking was performed in 8 participants (2 participants and 6 healthy controls) using Tobii Pro (60 Hz) to examine arousal differences across valence conditions.

4.5 Behavioral analysis

We quantified the distribution of emotional ratings across all trials and subjects (Fig. 1c). The trials assigned to each category were counted and differences between rating conditions were tested using ART ANOVA with permutation tests and FDR correction (Fig. S1a). Response times (RT) was defined as the interval between image offset and the participant's rating response by to button press (Fig. 1d and Fig. S1b). Trials with RTs shorter than 0.2 s (anticipatory responses) or longer than 5 s (inattentive responses) were excluded, resulting in the removal of 8.87% of trials. ART ANOVA test was also performed followed by post-hoc permutation with FDR correction across ratings and image categories. For pupil analysis (Fig. 1e and Fig. S1c), raw pupil diameter data were preprocessed to extract the most valid eye per sample. Artifacts (>3 SD change or confidence <0.8) were removed and missing samples interpolated. Data were smoothed with a third-order Butterworth low-pass filter (4 Hz low-pass), and z-score normalized per subject. ART ANOVA assessed differences across emotional ratings, with pairwise cluster-based permutation t-test for time-resolved analysis.

4.6 LFP preprocessing

Macroelectrode channels were selected based on their anatomical localization in the hippocampus and amygdala. LFPs were downsampled to 1000 Hz and common-average referenced across all channels within each subject. Bipolar referencing between neighboring contact pairs was used for beta burst detection [85, 86]. All prepreprocessing was conducted with the FieldTrip toolbox[87]. Data were bandpass filtered with a zero-phase lag FIR filter (1-200 Hz) and notch filtered at 60 Hz harmonics (59–61 Hz, 119–121 Hz, 179–181 Hz). Epochs were extracted from -1 to 3s relative to stimulus onset. Artifacts were rejected using automated detection ($z > 3$), followed by manual inspection to remove segments containing non-physiological artifacts (e.g., movement, clipping, or sudden transients).

4.7 Time-frequency analysis

Time-frequency decomposition was performed to characterize task-induced power changes relative to baseline activity for each channel based on FieldTrip toolbox with Morlet wavelets (4-14 cycles) over 1 to 150 Hz in 1 Hz steps [88]. Power was baseline-normalized using the -1 to 0 s pre-stimulus window. Cluster-based permutation tests were used to assess differences in spectral power between the pre-stimulus and stimulus windows (0–3 s after image onset) or across valence conditions. Statistical significance was determined by identifying clusters in time-frequency space exceeding chance levels based on the permutation distribution.

4.8 Beta burst detection and characterization

Beta bursts were identified and characterized based on established protocols from recent intracranial and MEG literature [40, 50, 51]. Signals were bandpass filtered (13–30 Hz; two-pass, fourth-order Butterworth) and transformed via the Hilbert method to extract amplitude envelopes, smoothed with a 20 ms moving average (Fig. 3c). Beta bursts detection criteria includes: (1) power spectrum > 2 SD above the trial mean, (2) ≥ 3 oscillatory cycles (i.e., three peaks and three troughs), (3) smoothened envelope > 1.5 SD above the tiral mean, and (4) duration between 80 ms to 500 ms. Note that the first two criteria were applied on the bandpass-filtered signal, while the last two criteria used the envelope. Manual verification by two independent raters was used for burst verification based on comparative assessments (10% of detected bursts, achieving above 90% agreement) of signal quality. Sensitivity analysis confirmed results were robust to detection threshold variations (1.5-2.5 SD). The characteristics (Fig. 3d), including burst rate, duration, and peak frequency, were calculated per trial. Peak frequency was calculated using a wavelet-based spectral estimate (cfg.toi = -1 all:3). Burst rate was defined as the number of bursts per 3 s stimulus period normalized by trial length. ART ANOVA with permutation t tests were used for analysis between conditions in these two regions.

4.9 PAC

The instantaneous amplitude of HG and phase of beta of the LFPs for each electrode during the 3 s following image onset of each trial were extracted using the Hilbert transform (Fig. 6). Phases were binned into 18 bins of 20 degrees. PAC was quantified by the MI, using the Kullback-Leibler distance method combined with Shannon's entropy [52, 53]. Significance was assessed against 100 surrogate MI distributions generated by trial shuffling. All comparisons of MI used ART ANOVA test followed by permutation tests with FDR correction ($p < 0.05$). Pairwise Mardia–Watson–Wheeler tests with FDR correction were used to compare the phase preference of the maximum amplitudes across emotional ratings, with the Rayleigh test used to compare the phase distribution against a uniform distribution.

4.10 Spike sorting

Microwire signals were bandpass filtered (300-3000 Hz) and spike-sorted using Wave_Clus3 (single-cell quality metrics are shown in Fig. S2) [89]. Detected spike clusters were subsequently curated manually using custom MATLAB scripts and the Wave_Clus GUI. Units were retained if: (1) $<5\%$ inter-spike intervals (ISIs) violations (ISI < 3 ms), (2) >200 spikes per session, and (3) physiological waveforms [50, 90]. Waveform stability and isolation quality were visually verified through ISI histograms, PCA projections, and consistency across time. This yielded 136 hippocampal and 82 amygdala units across participants. Neurons were classified as putative pyramidal cells or narrow interneurons based on waveform characteristics [91]. Wide interneurons were identified by trough-to-peak duration >0.425 ms, acg tau rise $> 6ms$, and firing rates <10 Hz, while narrow interneurons had short trough-to-peak duration (<0.425 ms) and higher firing rates (>10 Hz). The remaining cells with wide waveforms and low

firing rates are assigned as pyramidal cells. Notably, the waveform and firing properties of wide interneurons do not clearly separate from the clusters of pyramidal or narrow interneurons so we placed them as a separate category.

4.11 Neural response analysis

Firing rates were calculated in 2-ms non-overlapping bins and smoothened with a Gaussian kernel ($\sigma = 200$ ms). Population peristimulus time histograms (PSTHs) were computed by pooling neurons within each region for each emotion rating condition. Cluster-based permutation tests identified emotion-specific responses. Responsive neurons (Fig. 2a, b) were defined as units that showed significant rate changes during 300-1300 ms after image onset versus baseline (-1000-0 ms) (two-tailed cluster-based permutation t-test, P < 0.05) [92, 93]. This yielded multiple subsets of neurons preferentially responding to specific emotional rating levels. To assess population-level dynamics (Fig. 2c, d), emotional-responsive units from all subjects were pooled separately for the hippocampus and amygdala to form regional populations. For each brain region and each emotional rating, we computed trial-averaged population firing rates. These population PSTHs were then grouped and com- pared across different emotional conditions. Group-level statistical comparisons were conducted using a cluster-based permutation test ($p < 0.05$).

To assess for the effect of valence effects on the neuronal population while accounting for baseline activity, we fitted population-level generalized linear models (GLMs) to all trials from the valence-selective neurons in Fig. 2e-f. For very unpleasant -responsive neurons in amygdala, the model was:

$$\mathrm{FR_{task}} \sim \mathrm{FR_{baseline}} + \mathrm{Valence}$$

with very unpleasant as the reference. $\mathrm{FR_{task}}$ was the mean firing rate 0.5–1.0 s after image onset, $\mathrm{FR_{baseline}}$ was the mean firing rate from -1.0 to 0 s before onset, and *Valence* was a categorical factor. The same procedure was applied to very pleasant - responsive neurons in hippocampus, with very pleasant as the reference. β coefficients indicate adjusted differences from the reference valence, and model-predicted FRs were computed at the mean baseline FR. Significance for contrasts with the reference was taken from GLM coefficient p-values.

4.12 High-gamma and spike activity during beta bursts

To examine the physiological relevance of transient beta bursts, we analyzed both the dynamics of HG amplitude (Fig. 4a, b) and neuronal firing rates (Fig. 4e, f) aligned to beta burst onset within these two regions. HG amplitude was obtained using a zero-phase FIR filter and Morlet wavele transform with a frequency resolution of 0.5 Hz (80:0.5:140 Hz) and a temporal resolution of 20 ms (-1 to 3 s around burst time). The extracted gamma envelopes were z-score normalized within each trial relative to the baseline window (-1000-0 ms) and averaged per channel. Responses were then concatenated across subjects to form group-level representations. For neuronal spiking data, Gaussian-smoothed firing rates (200 ms kernel) were aligned to the onset each beta burst and aggregated across trials.

To assess the specificity of bursts, we created surrogate bursts by selecting random time points within 200 ms before or after each real burst, excluding any intervals with artifacts or overlapping bursts. Surrogate-aligned HG amplitudes and firing rates were calculated using the same procedure. Subsequently, HG amplitude and Gaussian-smoothed firing rates were aligned to burst onset and compared to surrogate events [40]. We also computed inter-trial phase coherence (ITPC, Fig. 4c, d) of beta-band LFPs to assess whether beta bursts were associated with phase-aligned activity. Phase estimates were extracted using Morlet wavelets, and ITPC was calculated per channel and concatenated across subjects. Finally, to evaluate cross-regional interactions, we extracted high-gamma amplitude and firing rates from ipsilateral electrodes during beta bursts detected in the other region, aligned to burst onset times. The significance of real bursts versus surrogate bursts was assessed using cluster-based permutation tests. The comparisons across conditions during bursts were evaluated by ART ANOVA with multiple comparisons FDR correction and pairwise permutation tests.

4.13 Statistical analysis

We employed non-parametric statistical methods throughout to account for violations of normality and the presence of unequal trial counts across conditions. Statistical tests were chosen based on the number of groups being compared and whether the analysis was time-resolved. Non-parametric approaches were adopted for all analyses as described below:

To compare conditions without a temporal dimension, we used two-tailed permutation tests to assess differences in trial-level distributions between the two groups. To assess differences in time-resolved signals between two conditions, we used two-tailed cluster-based non-parametric permutation tests. For the permutation, we generated 1000 random surrogate datasets by shuffling the labels of two groups with the Monte Carlo method. At each timepoint, an uncorrected p-value was calculated; timepoints with $p < 0.05$ were grouped into contiguous clusters. The empirical p-values for each cluster were computed as the proportion of permutations yielding a cluster statistic greater than or equal to the observed one [88, 94].

For comparisons involving three or more conditions without a temporal dimension, we used an ART ANOVA test [95, 96] as a non-parametric alternative to repeated-measures ANOVA. If a significant effect was detected, post-hoc pairwise comparisons were performed using permutation tests with p-values corrected for multiple comparisons using the Benjamini–Hochberg false discovery rate (FDR) procedure. For time-resolved multi-group analyses, ART ANOVA was performed at each time point, followed by pairwise comparison using cluster-based permutation test with FDR correction. All statistics were conducted in Matlab, except ART ANOVA which was finished in R.

5 Data availability

All data produced in this work are available on request from the corresponding author via email, with no conditions or restrictions.

6 Code availability

Custom MATLAB code associated with this study for data analysis are available online at https://github.com/damisahlab/AMY-HPC-Emotion.

7 Acknowledgements

We thank the members of the Yale Comprehensive Epilepsy Center for their excellent patient care, the patients who participated in this study. This study was supported by grants from the National Institutes of Health (R01MH138291 ECD; the Hypothesis Fund Seed Grant (ECD), Center for Brain and Mind Health (ECD, AK, IHR, AS), VA National Center for PTSD and Wellcome Trust Grant: 310142/Z/24/Z (ECD, AK, IHR and JK).

8 Author Contributions

Conceptualization, E.C.D.; Methodology, E.C.D, Y.Z., Y.H., G.P., B.Z., A.A. and A.P.T; Software, Y.Z., Y.H., B.Z., A.A. and A.P.T; Validation, Y.Z., Y.H.; Formal analysis, E.C.D, Y.Z., Y.H.; Investigation, E.C.D, Y.Z., Y.H.; Resources, E.C.D.; Data curation, E.C.D, Y.Z., Y.H.; Writing—original draft, Y.Z., Y.H.; Writing—review and editing, E.C.D, Y.Z., Y.H., G.P., B.Z., A.A, A.P.T, A.S., Z.Z., S.O, I.H., A.K., J.K., K.S., X.G., C.P.; Visualization, Y.Z., Y.H.; Supervision, E.C.D.; Project administration, E.C.D, Y.Z., Y.H.; Funding acquisition E.C.D.

9 Competing Interests

The authors declare no competing interests.

References

[1] Heilman, K.M., Gilmore, R.L.: Cortical influences in emotion. Journal of Clinical Neurophysiology **15**(5), 409–423 (1998)

[2] LeDoux, J.: The emotional brain, fear, and the amygdala. Cellular and molecular neurobiology **23**(4), 727–738 (2003)

[3] Phelps, E.A., LeDoux, J.E.: Contributions of the amygdala to emotion processing: from animal models to human behavior. Neuron **48**(2), 175–187 (2005)

[4] Constantinou, M., Yankouskaya, A., Burianová, H.: Valence effects on episodic memory in young and old adults following exposure to emotional stimuli. European Journal of Neuroscience **61**(5), 70041 (2025)

[5] Ramos, C.G.L., Tan, H., Yamamoto, E.A., Cleary, D.R., Mazur-Hart, D.J., Shahin, M.N., Raslan, A.M.: Stereotactic electroencephalography in epilepsy patients for mapping of neural circuits related to emotional and psychiatric behaviors: a systematic review. Neurosurgical focus **54**(2), 4 (2023)

[6] McDonald, A.J., Mott, D.D.: Functional neuroanatomy of amygdalohippocampal interconnections and their role in learning and memory. Journal of neuroscience research **95**(3), 797–820 (2017)

[7] Kwon, D., Kim, J., Yoo, S.B.M., Shim, W.M.: Coordinated representations for naturalistic memory encoding and retrieval in hippocampal neural subspaces. Nature Communications **16**(1), 641 (2025)

[8] Pirastru, A., Di Tella, S., Cazzoli, M., Esposito, F., Baselli, G., Baglio, F., Blasi, V.: The impact of emotional valence and stimulus habituation on fMRI signal reliability during emotion generation. NeuroImage **284**, 120457 (2023)

[9] Felix-Ortiz, A.C., Beyeler, A., Seo, C., Leppla, C.A., Wildes, C.P., Tye, K.M.: BLA to vHPC inputs modulate anxiety-related behaviors. Neuron **79**(4), 658–664 (2013)

[10] Pi, G., Gao, D., Wu, D., Wang, Y., Lei, H., Zeng, W., Gao, Y., Yu, H., Xiong, R., Jiang, T., *et al.*: Posterior basolateral amygdala to ventral hippocampal CA1 drives approach behaviour to exert an anxiolytic effect. Nature communications **11**(1), 183 (2020)

[11] Kim, W.B., Cho, J.-H.: Encoding of contextual fear memory in hippocampal–amygdala circuit. Nature communications **11**(1), 1382 (2020)

[12] Nguyen, R., Koukoutselos, K., Forro, T., Ciocchi, S.: Fear extinction relies on ventral hippocampal safety codes shaped by the amygdala. Science Advances **9**(22), 4881 (2023)

[13] Akiki, T.J., Jubeir, J., Bertrand, C., Tozzi, L., Williams, L.M.: Neural circuit basis of pathological anxiety. Nature Reviews Neuroscience **26**(1), 5–22 (2025)

[14] Butler, O., Willmund, G., Gleich, T., Zimmermann, P., Lindenberger, U., Gallinat, J., Kühn, S.: Cognitive reappraisal and expressive suppression of negative emotion in combat-related posttraumatic stress disorder: a functional MRI study. Cognitive therapy and research **43**, 236–246 (2019)

[15] Cruces-Solis, H., Babaev, O., Ali, H., Piletti Chatain, C., Mykytiuk, V., Balekoglu, N., Wenger, S., Krueger-Burg, D.: Altered theta and beta oscillatory synchrony in a genetic mouse model of pathological anxiety. The FASEB Journal **35**(6), 21585 (2021)

[16] Fales, C., Barch, D., Burgess, G., Schaefer, A., Mennin, D., Gray, J., Braver, T.: Anxiety and cognitive efficiency: differential modulation of transient and sustained neural activity during a working memory task. Cognitive, affective, & behavioral neuroscience **8**(3), 239–253 (2008)

[17] Kirkby, L.A., Luongo, F.J., Lee, M.B., Nahum, M., Van Vleet, T.M., Rao, V.R., Dawes, H.E., Chang, E.F., Sohal, V.S.: An amygdala-hippocampus subnetwork that encodes variation in human mood. Cell **175**(6), 1688–1700 (2018)

[18] Jackson, A.D., Cohen, J.L., Phensy, A.J., Chang, E.F., Dawes, H.E., Sohal, V.S.: Amygdala-hippocampus somatostatin interneuron beta-synchrony underlies a cross-species biomarker of emotional state. Neuron **112**(7), 1182–1195 (2024)

[19] Khodadadifar, T., Soltaninejad, Z., Ebneabbasi, A., Eickhoff, C.R., Sorg, C., Van Eimeren, T., Vogeley, K., Zarei, M., Eickhoff, S.B., Tahmasian, M.: In search of convergent regional brain abnormality in cognitive emotion regulation: A transdiagnostic neuroimaging meta-analysis. Human brain mapping **43**(4), 1309–1325 (2022)

[20] Qasim, S.E., Mohan, U.R., Stein, J.M., Jacobs, J.: Neuronal activity in the human amygdala and hippocampus enhances emotional memory encoding. Nature human behaviour **7**(5), 754–764 (2023)

[21] Sehlmeyer, C., Dannlowski, U., Schöning, S., Kugel, H., Pyka, M., Pfleiderer, B., Zwitserlood, P., Schiffbauer, H., Heindel, W., Arolt, V., *et al.*: Neural correlates of trait anxiety in fear extinction. Psychological medicine **41**(4), 789–798 (2011)

[22] Zhu, Y., Gao, H., Tong, L., Li, Z., Wang, L., Zhang, C., Yang, Q., Yan, B.: Emotion regulation of hippocampus using real-time fMRI neurofeedback in healthy human. Frontiers in human neuroscience **13**, 242 (2019)

[23] Balconi, M., Brambilla, E., Falbo, L.: Appetitive vs. defensive responses to emotional cues. Autonomic measures and brain oscillation modulation. Brain research **1296**, 72–84 (2009)

[24] Ertl, M., Hildebrandt, M., Ourina, K., Leicht, G., Mulert, C.: Emotion regulation by cognitive reappraisal—the role of frontal theta oscillations. NeuroImage **81**, 412–421 (2013)

[25] Symons, A.E., El-Deredy, W., Schwartze, M., Kotz, S.A.: The functional role of neural oscillations in non-verbal emotional communication. Frontiers in Human Neuroscience **10**, 239 (2016)

[26] Khanna, P., Carmena, J.M.: Beta band oscillations in motor cortex reflect neural population signals that delay movement onset. elife **6**, 24573 (2017)

[27] Torrecillos, F., Alayrangues, J., Kilavik, B.E., Malfait, N.: Distinct modulations in sensorimotor postmovement and foreperiod β-band activities related to error salience processing and sensorimotor adaptation. Journal of Neuroscience **35**(37), 12753–12765 (2015)

[28] Hanslmayr, S., Matuschek, J., Fellner, M.-C.: Entrainment of prefrontal beta oscillations induces an endogenous echo and impairs memory formation. Current biology **24**(8), 904–909 (2014)

[29] Zavala, B.A., Jang, A.I., Zaghloul, K.A.: Human subthalamic nucleus activity during non-motor decision making. Elife **6**, 31007 (2017)

[30] Balconi, M., Lucchiari, C.: EEG correlates (event-related desynchronization) of emotional face elaboration: a temporal analysis. Neuroscience letters **392**(1-2), 118–123 (2006)

[31] Meng, X., Liu, W., Zhang, L., Li, X., Yao, B., Ding, X., Yuan, J., Yang, J.: EEG oscillation evidences of enhanced susceptibility to emotional stimuli during adolescence. Frontiers in psychology **7**, 616 (2016)

[32] Mishra, S., Srinivasan, N., Tiwary, U.S.: Dynamic functional connectivity of emotion processing in beta band with naturalistic emotion stimuli. Brain sciences **12**(8), 1106 (2022)

[33] Schubring, D., Schupp, H.T.: Emotion and brain oscillations: high arousal is associated with decreases in alpha-and lower beta-band power. Cerebral Cortex **31**(3), 1597–1608 (2021)

[34] Schneider, T.R., Hipp, J.F., Domnick, C., Carl, C., Büchel, C., Engel, A.K.: Modulation of neuronal oscillatory activity in the beta-and gamma-band is associated with current individual anxiety levels. NeuroImage **178**, 423–434 (2018)

[35] Schutter, D.J., Putman, P., Hermans, E., Honk, J.: Parietal electroencephalogram beta asymmetry and selective attention to angry facial expressions in healthy human subjects. Neuroscience letters **314**(1-2), 13–16 (2001)

[36] Güntekin, B., Başar, E.: Event-related beta oscillations are affected by emotional eliciting stimuli. Neuroscience letters **483**(3), 173–178 (2010)

[37] Pavlenko, V., Chernyi, S., Goubkina, D.: EEG correlates of anxiety and emotional stability in adult healthy subjects. Neurophysiology **41**(5), 337–345 (2009)

[38] Sonkusare, S., Qiong, D., Zhao, Y., Liu, W., Yang, R., Mandali, A., Manssuer, L., Zhang, C., Cao, C., Sun, B., *et al.*: Frequency dependent emotion differentiation and directional coupling in amygdala, orbitofrontal and medial prefrontal cortex network with intracranial recordings. Molecular Psychiatry **28**(4), 1636–1646 (2023)

[39] Boroujeni, K.B., Womelsdorf, T.: Routing states transition during oscillatory bursts and attentional selection. Neuron **111**(18), 2929–2944 (2023)

[40] Lundqvist, M., Rose, J., Herman, P., Brincat, S.L., Buschman, T.J., Miller, E.K.: Gamma and beta bursts underlie working memory. Neuron **90**(1), 152–164 (2016)

[41] Voloh, B., Womelsdorf, T.: Cell-type specific burst firing interacts with theta and beta activity in prefrontal cortex during attention states. Cerebral Cortex **28**(12), 4348–4364 (2018)

[42] Fries, P.: Rhythms for cognition: Communication through coherence. Neuron **88**(1), 220–235 (2015)

[43] Li, J., Cao, D., Yu, S., Xiao, X., Imbach, L., Stieglitz, L., Sarnthein, J., Jiang, T.: Functional specialization and interaction in the amygdala-hippocampus circuit during working memory processing. Nature Communications **14**(1), 2921 (2023)

[44] Shin, H., Law, R., Tsutsui, S., Moore, C.I., Jones, S.R.: The rate of transient beta frequency events predicts behavior across tasks and species. elife **6**, 29086 (2017)

[45] Wessel, J.R.: Beta-bursts reveal the trial-to-trial dynamics of movement initiation and cancellation. Journal of Neuroscience **40**(2), 411–423 (2020)

[46] Feingold, J., Gibson, D.J., DePasquale, B., Graybiel, A.M.: Bursts of beta oscillation differentiate postperformance activity in the striatum and motor cortex of monkeys performing movement tasks. Proceedings of the National Academy of Sciences **112**(44), 13687–13692 (2015)

[47] Spitzer, B., Haegens, S.: Beyond the status quo: a role for beta oscillations in endogenous content (re) activation. eneuro **4**(4) (2017)

[48] Bradley, M.M., Lang, P.J.: The International Affective Picture System (IAPS) in the study of emotion and attention. In: Coan, J.A., Allen, J.J. (eds.) Handbook of Emotion Elicitation and Assessment, pp. 29–46. Oxford University Press, New York (2007)

[49] Marchewka, A., Żurawski, Ł., Jednoróg, K., Grabowska, A.: The Nencki Affective Picture System (NAPS): Introduction to a novel, standardized, wide-range, high-quality, realistic picture database. Behavior research methods **46**, 596–610 (2014)

[50] Kunz, L., Staresina, B.P., Reinacher, P.C., Brandt, A., Guth, T.A., Schulze-Bonhage, A., Jacobs, J.: Ripple-locked coactivity of stimulus-specific neurons and human associative memory. Nature neuroscience **27**(3), 587–599 (2024)

[51] Zich, C., Quinn, A.J., Bonaiuto, J.J., O'Neill, G., Mardell, L.C., Ward, N.S., Bestmann, S.: Spatiotemporal organisation of human sensorimotor beta burst activity. Elife **12**, 80160 (2023)

[52] Daume, J., Kamiński, J., Schjetnan, A.G.P., Salimpour, Y., Khan, U., Kyzar, M., Reed, C.M., Anderson, W.S., Valiante, T.A., Mamelak, A.N., Rutishauser, U.: Control of working memory by phase–amplitude coupling of human hippocampal neurons. Nature, 1–9 (2024)

[53] Saint Amour di Chanaz, L., Perez-Bellido, A., Wu, X., Lonzano-Soldevilla, D., Pacheco-Estefan, D., Lehongre, K., Conde-Blanco, E., Roldan, P., Adam, C., Lambrecq, V., Frazzini, V., Donaire, A., Carreno, M., Navarro, V., Valero-Cabre, A., Fuentemilla, L.: Gamma amplitude is coupled to opposed hippocampal theta-phase states during the encoding and retrieval of episodic memories in humans. Curr Biol (2023)

[54] Pignatelli, M., Beyeler, A.: Valence coding in amygdala circuits. Current opinion in behavioral sciences **26**, 97–106 (2019)

[55] Anderson, D.J., Adolphs, R.: A framework for studying emotions across species. Cell **157**(1), 187–200 (2014)

[56] Disner, S.G., Beevers, C.G., Haigh, E.A., Beck, A.T.: Neural mechanisms of the cognitive model of depression. Nature Reviews Neuroscience **12**(8), 467–477 (2011)

[57] Wang, S., Tudusciuc, O., Mamelak, A.N., Ross, I.B., Adolphs, R., Rutishauser, U.: Neurons in the human amygdala selective for perceived emotion. Proceedings of the National Academy of Sciences **111**(30), 3110–3119 (2014)

[58] Wang, S., Yu, R., Tyszka, J.M., Zhen, S., Kovach, C., Sun, S., Huang, Y., Hurlemann, R., Ross, I.B., Chung, J.M., *et al.*: The human amygdala parametrically encodes the intensity of specific facial emotions and their categorical ambiguity. Nature communications **8**(1), 14821 (2017)

[59] Volitaki, E., Forro, T., Li, K., Nevian, T., Ciocchi, S.: Activity of ventral hippocampal parvalbumin interneurons during anxiety. Cell reports **43**(6) (2024)

[60] Clark, D.L., Brown, E.C., Ramasubbu, R., Kiss, Z.H.: Intrinsic local beta

oscillations in the subgenual cingulate relate to depressive symptoms in treatment-resistant depression. Biol. Psychiatry **80**, 93–94 (2016)

[61] Costa, M., Lozano-Soldevilla, D., Gil-Nagel, A., Toledano, R., Oehrn, C.R., Kunz, L., Yebra, M., Mendez-Bertolo, C., Stieglitz, L., Sarnthein, J., *et al.*: Aversive memory formation in humans involves an amygdala-hippocampus phase code. Nature Communications **13**(1), 6403 (2022)

[62] Zheng, J., Anderson, K.L., Leal, S.L., Shestyuk, A., Gulsen, G., Mnatsakanyan, L., Vadera, S., Hsu, F.P., Yassa, M.A., Knight, R.T., *et al.*: Amygdala-hippocampal dynamics during salient information processing. Nature communications **8**(1), 14413 (2017)

[63] Chen, S., Tan, Z., Xia, W., Gomes, C.A., Zhang, X., Zhou, W., Liang, S., Axmacher, N., Wang, L.: Theta oscillations synchronize human medial prefrontal cortex and amygdala during fear learning. Science Advances **7**(34), 4198 (2021)

[64] Zheng, J., Stevenson, R.F., Mander, B.A., Mnatsakanyan, L., Hsu, F.P., Vadera, S., Knight, R.T., Yassa, M.A., Lin, J.J.: Multiplexing of theta and alpha rhythms in the amygdala-hippocampal circuit supports pattern separation of emotional information. Neuron **102**(4), 887–898 (2019)

[65] Lundqvist, M., Miller, E.K., Nordmark, J., Liljefors, J., Herman, P.: Beta: bursts of cognition. Trends in cognitive sciences (2024)

[66] Ede, F., Quinn, A.J., Woolrich, M.W., Nobre, A.C.: Neural oscillations: Sustained rhythms or transient burst-events? Trends in Neurosciences **41**(7), 415–417 (2018)

[67] Banaie Boroujeni, K., Womelsdorf, T.: Routing states transition during oscillatory bursts and attentional selection. Neuron **111**(18), 2929–294411 (2023)

[68] Dolcos, F., LaBar, K.S., Cabeza, R.: Interaction between the amygdala and the medial temporal lobe memory system predicts better memory for emotional events. Neuron **42**(5), 855–863 (2004)

[69] Hayes, D.J., Northoff, G.: Common brain activations for painful and non-painful aversive stimuli. BMC Neuroscience **13**(1), 60 (2012)

[70] Wen, Z., Pace-Schott, E.F., Lazar, S.W., Rosén, J., Åhs, F., Phelps, E.A., LeDoux, J.E., Milad, M.R.: Distributed neural representations of conditioned threat in the human brain. Nature Communications **15**(1), 2231 (2024)

[71] Inman, C.S., Hollearn, M.K., Augustin, L., Campbell, J.M., Olson, K.L., Wahlstrom, K.L.: Discovering how the amygdala shapes human behavior: From lesion studies to neuromodulation. Neuron **111**(24), 3906–3910 (2023)

[72] Bijanki, K.R., van Rooij, S.J.H., Ely, T.D., Stevens, J.S., Inman, C.S., Fasano, R.E., Carter, S.E., Winters, S.J., Baman, J.R., Drane, D.L., Jovanovic, T., Willie, J.T.: Case series: Unilateral amygdala ablation ameliorates post-traumatic stress disorder symptoms and biomarkers. Neurosurgery **87**(4), 796 (2020)

[73] Gill, J.L., Schneiders, J.A., Stangl, M., Aghajan, Z.M., Vallejo, M., Hiller, S., Topalovic, U., Inman, C.S., Villaroman, D., Bari, A., Adhikari, A., Rao, V.R., Fanselow, M.S., Craske, M.G., Krahl, S.E., Chen, J.W.Y., Vick, M., Hasulak, N.R., Kao, J.C., Koek, R.J., Suthana, N., Langevin, J.-P.: A pilot study of closed-loop neuromodulation for treatment-resistant post-traumatic stress disorder. Nature Communications **14**(1), 2997 (2023)

[74] Zheng, J., Lin, J.J.: Modulating amygdala–hippocampal network communication: A potential therapy for neuropsychiatric disorders. Neuropsychopharmacology **43**(1), 218 (2017)

[75] Manning, J.R., Jacobs, J., Fried, I., Kahana, M.J.: Broadband shifts in local field potential power spectra are correlated with single-neuron spiking in humans. Journal of Neuroscience **29**(43), 13613–13620 (2009)

[76] Scangos, K.W., Khambhati, A.N., Daly, P.M., Makhoul, G.S., Sugrue, L.P., Zamanian, H., Liu, T.X., Rao, V.R., Sellers, K.K., Dawes, H.E., *et al.*: Closed-loop neuromodulation in an individual with treatment-resistant depression. Nature medicine **27**(10), 1696–1700 (2021)

[77] Beck, A.T., Epstein, N., Brown, G., Steer, R.A.: An inventory for measuring clinical anxiety: psychometric properties. Journal of consulting and clinical psychology **56**(6), 893 (1988)

[78] Beck, A.T., Steer, R.A., Brown, G.: Beck depression inventory–II. Psychological assessment (1996)

[79] Fried, I., Wilson, C.L., Maidment, N.T., Engel, J., Behnke, E., Fields, T.A., Macdonald, K.A., Morrow, J.W., Ackerson, L.: Cerebral microdialysis combined with single-neuron and electroencephalographic recording in neurosurgical patients. Journal of neurosurgery **91**(4), 697–705 (1999)

[80] Rutishauser, U., Reddy, L., Mormann, F., Sarnthein, J.: The architecture of human memory: insights from human single-neuron recordings. Journal of Neuroscience **41**(5), 883–890 (2021)

[81] Davis, T.S., Caston, R.M., Philip, B., Charlebois, C.M., Anderson, D.N., Weaver, K.E., Smith, E.H., Rolston, J.D.: LeGUI: a fast and accurate graphical user interface for automated detection and anatomical localization of intracranial electrodes. Frontiers in Neuroscience **15**, 769872 (2021)

[82] Neuromorphometrics, Inc.: Neuromorphometrics Atlas. https://www.

neuromorphometrics.com/. Available: https://www.neuromorphometrics.com/ (2022)

[83] Fonov, V., Evans, A.C., Botteron, K., Almli, C.R., McKinstry, R.C., Collins, D.L., Group, B.D.C., *et al.*: Unbiased average age-appropriate atlases for pediatric studies. Neuroimage **54**(1), 313–327 (2011)

[84] Fonov, V.S., Evans, A.C., McKinstry, R.C., Almli, C.R., Collins, D.: Unbiased nonlinear average age-appropriate brain templates from birth to adulthood. NeuroImage **47**, 102 (2009)

[85] He, S., Mostofi, A., Syed, E., Torrecillos, F., Tinkhauser, G., Fischer, P., Pogosyan, A., Hasegawa, H., Li, Y., Ashkan, K., Pereira, E., Brown, P., Tan, H.: Subthalamic beta-targeted neurofeedback speeds up movement initiation but increases tremor in Parkinsonian patients. Elife **9** (2020)

[86] Vaz, A.P., Wittig Jr, J.H., Inati, S.K., Zaghloul, K.A.: Replay of cortical spiking sequences during human memory retrieval. Science **367**(6482), 1131–1134 (2020)

[87] Oostenveld, R., Fries, P., Maris, E., Schoffelen, J.-M.: FieldTrip: open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. Computational intelligence and neuroscience **2011**(1), 156869 (2011)

[88] Gill, J.L., Schneiders, J.A., Stangl, M., Aghajan, Z.M., Vallejo, M., Hiller, S., Topalovic, U., Inman, C.S., Villaroman, D., Bari, A., *et al.*: A pilot study of closed-loop neuromodulation for treatment-resistant post-traumatic stress disorder. Nature communications **14**(1), 2997 (2023)

[89] Chaure, F.J., Rey, H.G., Quian Quiroga, R.: A novel and fully automatic spike-sorting implementation with variable number of features. Journal of Neurophysiology **120**(4), 1859–1871 (2018)

[90] Axmacher, N.: Intracranial EEG: a Guide for Cognitive Neuroscientists. Springer, Cham, Switzerland (2023)

[91] Petersen, P.C., Siegle, J.H., Steinmetz, N.A., Mahallati, S., Buzsáki, G.: CellExplorer: A framework for visualizing and characterizing single neurons. Neuron **109**(22), 3594–3608 (2021)

[92] Kehl, M.S., Mackay, S., Ohla, K., Schneider, M., Borger, V., Surges, R., Spehr, M., Mormann, F.: Single-neuron representations of odours in the human brain. Nature **634**(8034), 626–634 (2024)

[93] Quian Quiroga, R., Boscaglia, M., Jonas, J., Rey, H.G., Yan, X., Maillard, L., Colnat-Coulbois, S., Koessler, L., Rossion, B.: Single neuron responses underlying face recognition in the human midfusiform face-selective cortex. Nature Communications **14**(1), 5661 (2023)

[94] Maris, E., Oostenveld, R.: Nonparametric statistical testing of EEG-and MEG-data. Journal of neuroscience methods **164**(1), 177–190 (2007)

[95] Elkin, L.A., Kay, M., Higgins, J.J., Wobbrock, J.O.: An aligned rank transform procedure for multifactor contrast tests. In: The 34th Annual ACM Symposium on User Interface Software and Technology, pp. 754–768 (2021)

[96] Ricci, L., Croce, P., Pulitano, P., Boscarino, M., Zappasodi, F., Narducci, F., Lanzone, J., Sancetta, B., Mecarelli, O., Di Lazzaro, V., *et al.*: Levetiracetam modulates EEG microstates in temporal lobe epilepsy. Brain Topography **35**(5), 680–691 (2022)

Fig. 1: **Experimental paradigm and behavioral validation of emotional processing**. **a**, Recording setup and electrode localization. Top: Behnke-Fried depth electrode with protruding microwires enabling simultaneous LFP and single-unit recordings. Bottom: Microwire locations across all participants (n=15) shown in MNI space. AMY microwire bundles (n=16, pink), hippocampus (n=23 microwire bundles orange). **b**, Emotional valence rating task. Participants viewed IAPS/NAPS images for 3 s followed by a 5-point valence rating (1=very pleasant, 5=very unpleasant). Inter-trial interval: 1s. **c**, Distribution of valence ratings across all trials (n=2,270). Neutral ratings were most frequent (30.4%), extreme ratings least frequent (very pleasant: 20.3%, very unpleasant: 16.3%). **d**, Response times showed an inverted-U pattern with faster decisions for extreme emotional rating. Violin plots show distribution with median (white dot), IQR (thick bar), and full range (thin lines). ART ANOVA test followed by permutation tests with FDR correction. $**p < 0.01$, $***p < 0.001$. **e**, Pupil diameter responses (n=8 participants) showed no differences across valence conditions, ruling out arousal confounds. Shaded areas: ±s.e.m. ART ANOVA: no main effect of condition ($p > 0.05$). AMY: amygdala, HPC: hippocampus. VP: very pleasant, P: pleasant, N: neutral, U: unpleasant, VU: very unpleasant.

Fig. 2: **Valence-specific single neuron responses in the human AMY and HPC**. **a**, Representative single neurons showing distinct emotional response patterns. Each panel shows raster plots (top) of trials sorted by valence rating and PSTHs (bottom) with baseline-corrected firing rates (± s.e.m.). Four exampler emotional rating associated neuronal responses are shown: Very pleaseant (top left, from HPC), Very unpleasant (top right from AMY), broadly responsive (bottom left, AMY), and non-responsive (bottom right, HPC) neurons. Insets show spike waveforms with amplitude scale bars. **b**, Venn diagrams revealing overlap of valence-responsive neurons in AMY (left, 54 total) and HPC (right, 81 total), with numbers indicating neurons selective for each rating combination. **c**, Population responses of valence-selective neurons in AMY. Top row: very pleasant-responsive neurons (n=13); bottom row: very unpleasant-responsive neurons (n=15). Left panels compare the corresponding trials to all other trials among the responsive neurons; right panels show direct comparison between trials for very pleasant and very unpleasant conditions. Black bars indicate significant differences (cluster-based permutation, $p < 0.05$). **d**, Same analysis for HPC neurons showing very pleasant-responsive (n=27, top) and very unpleasant-responsive (n=12, bottom) populations. **e**, GLM results for VU-responsive neurons in AMY. Left: β coefficients (mean ± 95% CI) from a population-level GLM with baseline FR (FR_B) and valence (VU as reference) as predictors of task-period FR. Right: model-predicted FRs (mean ± 95% CI) at the mean baseline FR. **f**, Same as **e**, but for VP-responsive neurons in HPC with VP as the reference. *$p < 0.05$, ***$p < 0.001$.

Fig. 3: **From sustained power to transient bursts-uncovering emotion-specific dynamics**. **a**, Time-frequency analysis in AMY. Top: spectrograms showing stimulus-evoked changes in gamma (40-140 Hz, left) and beta (13-30 Hz, right) power. Black contours indicate significant changes from baseline (cluster-based permutation, $p < 0.05$). Bottom: averaged power traces for each valence rating (1-5) show no emotion-specific differences despite clear stimulus responses. (ART ANOVA FDR corrected $p > 0.05$). Shading: $\pm$ s.e.m. **b**, Same analysis for HPC, similarly showing no valence-specific power modulation. **c**, Beta burst detection pipeline. Example trial showing (top to bottom): raw macroelectrode LFP trace, LFP bandpass-filtered in the beta band, smoothed amplitude envelope of the beta-filtered signal with detection threshold (red dashed line), time-frequency spectrogram, and aligned gamma-band activity. Green highlighted line indicates detected bursts. **d**, Beta burst characteristics across emotional ratings. Top row: AMY; bottom row: HPC. Left to right: burst duration, rate, and peak frequency. Each dot represents one LFP channel; bars show group means. Asterisks indicate significant differences unpleasant vs neutral ratings in the AMY (*$p < 0.05$, permutation test with ART ANOVA multiple comparisons FDR corrected).

Fig. 4: **Asymmetric beta burst control: amygdala gates hippocampal emotional processing. a**, AMY beta bursts are associated with local high-gamma (80-140 Hz) increase. Top: averaged HG amplitude for real bursts (purple) significantly exceeds surrogate events (black) ($p < 0.05$). Middle: single-channel heatmaps showing consistent HG enhancement across recording sites. Bottom: Comparisons of HG amplitude during bursts in the AMY between extreme and moderate emotional ratings. 0 is burst onset **b**, Same analysis for HPC beta bursts, showing similar local HG enhancement. **c,d**, Inter-trial phase coherence (ITPC) during beta bursts in AMY (c) and HPC (d). Very unpleasant stimuli (purple) show significantly stronger intertrial phase coherence than neutral and pleasant ratings ($p < 0.05$). **e**, Population neuronal response in the AMY and HPC during AMY bursts. Left panels: Comparisons of FRs in AMY neurons (n = 82 neurons) (top) and in HPC neurons (bottom, n = 136 neurons) aligned to AMY burst onset vs surrogate bursts. Only the HPC neurons show a decrease in FRs (cluster permutation test $p < 0.05$). Right panels: Increased AMY firing for very unpleasant ratings at the onset of AMY beta burst, while decreased HPC firing for very unpleasant ratings during AMY bursts (all, ART ANOVA FDR corrected $p < 0.05$, permutation test $p < 0.05$) **f**, Population neuronal response in the HPC and AMY during HPC bursts (n=136 neurons). Left panels: Comparisons of FRs in HPC neurons (top) and in AMY neurons (bottom) aligned to HPC bursts vs surrogate bursts. Only the HPC neurons show a decrease in FRs (cluster permutation test $p < 0.05$). Right panels: Decreased HPC firing for very unpleasant ratings at the onset of HPC beta burst (ART ANOVA FDR corrected $p < 0.05$, permutation test $p < 0.05$) but no effect on AMY firing pattern during HPC bursts (ART ANOVA FDR adjusted $p > 0.05$), indicating unidirectional AMY→HPC influence. Shading: ± s.e.m. Horizontal bars indicate significant periods (black: ratings VP vs VU; red: VP vs P/N/U; blue: VU vs P/N/U; all $p < 0.05$).

Fig. 5: **Hippocampal pyramidal cells (PYR) and interneurons (INT) show differing responses to beta bursts during negative emotional valence rating. a**, Cell-type classification based on waveform characteristics. Pyramidal cells (AMY: 61% N=50, HPC: 59% N=80) constitute the majority, with narrow interneurons (AMY: 5% N=4, HPC: 9% N=12), wide interneurons (AMY: 16% N=13, HPC: 15% n=21), and unclassified cells making up the remainder. **b**, Cell-type-specific burst modulation in HPC neurons. Pyramidal cells (PYR, left, N=80) show early suppression during beta bursts for very unpleasant stimuli. In contrast, narrow interneurons (INT, middle, N=12) and wide interneurons (INT, right, N=21) did not exhibit significant difference across valence. **c**, Cell-type-specific AMY burst modulation of HPC neurons. Pyramidal cells (PYR, left, N=80) show no difference across valence. However narrow interneurons (INT, middle, N=12) and wide interneurons (INT, right, N=21) show suppression for very unpleasant stimuli (rating 5). This opposing pattern suggests beta bursts may coordinate emotional processing through differential modulation of excitatory and inhibitory circuits. Black bars: $p < 0.05$, cluster-based permutation test.

Fig. 6: **Beta-gamma PAC encodes emotional valence through coupling strength relationships. a**, Example PAC analysis from a single trial. From top: raw LFP, beta-filtered (13-30 Hz), gamma-filtered (40-140 Hz), and computed PAC showing moments of strong coupling (red). **b**, Preferred coupling phase varies with emotional valence. Circular histograms showing the preferred beta phase at which gamma amplitude peaks for each emotional rating in the AMY (top) and HPC (bottom). All pairwise comparisons: $p < 0.01$, Mardia-Watson-Wheeler test with FDR correction. **c**, Emotion-specific PAC patterns. Comodulograms show coupling strength (MI) between beta phase and gamma amplitude frequencies for each emotional condition. Very unpleasant stimuli show strongest coupling at beta band in both AMY (top) and HPC (bottom). **d**, Group-averaged MI as a function of beta frequency for each emotional condition. Very unpleasant (purple) shows significantly stronger coupling than all other conditions. Shading area: $\pm$ s.e.m.

Fig. S1: **Stimulus category effects on behavior are limited to rating distributions. a**, Distribution of emotional ratings varies by stimulus category. Objects and humans elicit predominantly neutral ratings, while animals and scenes evoke more extreme ratings, particularly pleasant. Bars show mean number of trials $\pm$ s.e.m. ART ANOVA followed by permutation tests with FDR correction revealed significant category effects (all $p < 0.001$). **b**, Response times show no category specificity. Mean RTs ($\pm$ s.e.m.) are similar across all four stimulus categories (ART ANOVA: $F_{(3,2072)} = 0.73$, $p = 0.53$). **c**, Pupil diameter responses are category-independent. Time courses show no arousal differences between stimulus categories (ART ANOVA: $p > 0.05$). Shaded areas: $\pm$ s.e.m. Vertical dashed line indicates image onset.

Fig. S2: **Quality metrics to evaluate reliable single-unit isolation.** **(a)** Proportion of inter-spike intervals (ISI) below 3 ms ($1.13\% \pm 1.63\%$, mean $\pm$ s.d.). **(b)** Signal-to-noise ratio (SNR) for the mean waveform across all spikes as compared to the standard deviation of the background noise (2.01 ± 1.00, mean $\pm$ s.d.). **(c)** Average firing rate within the entire recording session for all identified single units (3.77 ± 4.83, mean $\pm$ s.d.).

Table S1: Subject information

Subject ID	Age	Handedness	Sex	BAI	BDI	ASM
1	23	R	Male	6	5	CLB, LTG
2	61	R	Female	14	11	OXC
3	28	R	Male	−	−	CLB, CNB, PER
4	21	R	Male	−	−	CLB, OXC, TPM
5	18	R	Female	−	−	CLB, LEV, OXC
6	49	R	Female	11	2	CLB, CNB, PGB
7	47	R	Female	−	−	BRV, CNB, LTG
8	27	R	Female	14	18	CLB, OXC
9	36	R	Male	16	21	CNB, PHT, TPR
10	35	L	Female	26	28	CLB, LCM, TPR
11	45	R	Female	3	13	CLB, CNB, LCM
12	41	R	Female	−	−	CLB, CNB, CBD
13	32	R	Male	32	19	CLB, CNB, LCM
14	22	R	Male	−	−	CLB, CNB, LCM
15	23	R	Female	−	−	CLB, LCM

Table S2: MNI coordinates of microwires for different brain regions.

Electrode ID	Patient ID	Location	Side	X	Y	Z
1	1	Hippocampus	Left	−25.68	−18.76	−19.12
2	1	Amygdala	Left	−20.63	−5.92	−20.86
3	2	Hippocampus	Left	−24.66	−18.37	−17.27
4	2	Hippocampus	Right	27.60	−16.62	−17.10
5	2	Amygdala	Left	−18.97	−6.17	−20.79
6	2	Amygdala	Right	24.96	−6.31	−25.30
7	3	Hippocampus	Right	37.76	−13.55	−18.30
8	3	Amygdala	Right	16.08	−2.12	−22.97
9	4	Hippocampus	Right	20.56	−15.02	−15.85
10	5	Hippocampus	Right	27.65	−15.71	−23.18
11	5	Amygdala	Right	21.79	−4.43	−21.48
12	6	Hippocampus	Right	18.68	−7.66	−20.70
13	6	Amygdala	Left	−18.22	−9.02	−25.21
14	7	Amygdala	Right	21.58	−3.52	−26.20
15	7	Amygdala	Left	−20.98	−7.06	−23.74
16	8	Hippocampus	Left	−19.09	−7.05	−25.35
17	8	Hippocampus	Right	23.33	−12.54	−20.00
18	8	Amygdala	Left	−19.86	1.47	−27.02
19	8	Amygdala	Right	19.99	0.05	−28.69
20	9	Hippocampus	Left	−26.11	−9.49	−23.71
21	9	Hippocampus	Right	24.24	−10.36	−19.10
22	10	Hippocampus	Right	25.41	−12.53	−17.62
23	10	Hippocampus	Right	31.39	−28.20	−6.98
24	10	Hippocampus	Left	−28.83	−21.17	−13.09
25	10	Amygdala	Right	20.59	−1.82	−24.19
26	10	Amygdala	Left	−20.49	−5.15	−21.68
27	11	Hippocampus	Left	−23.11	−14.84	−15.39
28	11	Hippocampus	Left	−22.35	−31.47	−4.94
29	11	Hippocampus	Right	26.51	−15.82	−15.76
30	11	Amygdala	Left	−16.84	−5.02	−18.27
31	12	Amygdala	Left	−15.93	−5.97	−24.35
32	12	Hippocampus	Left	−21.00	−16.91	−19.73
33	12	Hippocampus	Right	21.70	−23.40	−21.56
34	13	Hippocampus	Right	25.22	−9.19	−20.95
35	13	Amygdala	Left	19.27	−3.14	−18.30
36	14	Hippocampus	Left	−33.66	−33.05	−8.26
37	14	Hippocampus	Left	−31.56	−14.61	−16.81
38	15	Amygdala	Right	19.80	−1.10	−20.07
39	15	Hippocampus	Right	33.51	−14.97	−17.82

Table S3: Detected neuron/burst count per region

Subject ID	Session	Number of detected units			LFP Channel number		Number of detected bursts	
		All units	units in HPC	units in AMY	AMY	HPC	Bursts in AMY	Bursts in HPC
1	1	11	6	5	2	2	252	288
2	2	33	27	6	3	3	762	848
3	1	3	0	3	1	2	141	305
4	1	0	0	0	2	9	297	1308
5	1	1	1	0	2	3	291	514
6	1	16	14	2	2	2	258	287
7	1	10	0	10	3	2	406	291
8	1	42	29	13	1	3	129	433
9	1	3	3	0	0	2	0	270
10	1	10	5	5	2	3	248	382
11	1	2	2	0	3	4	366	678
12	1	15	4	11	3	5	448	734
13	2	36	32	4	2	2	547	564
14	2	5	5	0	2	2	599	595
15	2	31	8	23	1	2	353	666
Total	19	218	136	82	29	46	5097	8163