Integrated Human Transcriptomics Identifies Fallopian Tube Progenitors as Plausible Precursors of High-Grade Serous Ovarian Cancer

Qian Li[1,2,#], Keren Cheng[3,#,*], Lili Sun[2], Wei Yan □,□,□,*

[1]Department of Medical Genetics, China Medical University, Shenyang, Liaoning 110122, China
[2]Department of Gynecological Oncology, Cancer Hospital of China Medical University / Liaoning Cancer Hospital & Institute, Shenyang, Liaoning 110042, China
[3]Department of Obstetrics and Gynecology, Center for Reproductive Medicine, Fourth Affiliated Hospital, School of Medicine, International School of Medicine, Zhejiang University, Yiwu 322000, China
□The Lundquist Institute for Biomedical Innovation at Harbor-UCLA Medical Center, Torrance, CA 90502, USA
□Department of Medicine, David Geffen School of Medicine at UCLA, Los Angeles, CA 90095, USA
□School of Molecular Biosciences and Center for Reproductive Biology, Washington State University, Pullman, WA 99164, USA

#co-first authors

*Correspondence: Keren.Cheng@zju.edu.cn or wei.yan@wsu.edu

Lead Contact: Wei Yan (wei.yan@wsu.edu)

One sentence summary: Fallopian tube progenitor cells represent a key intermediate state linking normal epithelial hierarchy to the earliest stages of high-grade serous ovarian cancer development.

Abstract

High-grade serous ovarian cancer (HGSOC) is the most lethal gynecologic malignancy,

yet its earliest cellular precursors remain incompletely defined. Here, we integrate bulk

and single-nucleus RNA sequencing of human fallopian tubes, ovaries, and primary

HGSOC tumors to resolve the epithelial hierarchy of the fallopian tube and examine its

relationship to malignant states. We identify a rare LGR5$^+$/PGR$^+$ basal population as

putative epithelial stem cells and delineate a large pool of OVGP1$^+$/RNPC3$^+$ progenitors

that give rise to both secretory and ciliated lineages. By jointly analyzing normal and

malignant transcriptomes, we find that these progenitor populations show the strongest

transcriptional and developmental continuity with multiple HGSOC cell states,

suggesting that they represent the epithelial compartments most plausibly poised for

oncogenic divergence. Transcription factor network reconstruction highlights SPDEF as

a key regulator of progenitor identity and NR2F6 as a candidate oncogenic driver, while

inferred copy number variations and microenvironmental collagen-integrin signaling

further illuminate the genomic and stromal factors that consolidate malignant fate.

Together, these findings refine the cellular landscape of the human fallopian tube and

identify progenitor-enriched molecular features that could inform earlier detection,

improved risk stratification, and new preventive strategies for women at risk of

developing HGSOC.

INTRODUCTION

High-grade serous ovarian carcinoma (HGSOC) is the most common and lethal subtype of ovarian cancer, accounting for more than 70% of ovarian cancer deaths and associated with a 10-year survival rate below 30% (*1, 2*). For decades, HGSOC was thought to arise from the ovarian surface epithelium (OSE), but extensive molecular and pathological evidence has fundamentally reshaped this view over the past two decades (*3, 4*). HGSOC more closely resembles Müllerian duct-derived epithelia of the fallopian tube than the mesothelial-like OSE, based on its transcriptional, proteomic, and morphological features (*5, 6*). These observations have led to the now widely supported hypothesis that the fallopian tube epithelium (FTE), particularly the distal fimbrial region, represents the most plausible origin for many HGSOCs (*7, 8*).

Compelling support for this hypothesis comes from serous tubal intraepithelial carcinomas (STICs), early precursor lesions that harbor TP53 mutations identical to those in matched invasive tumors (*9, 10*). STICs are frequently found in the fimbriae of high-risk BRCA1/2 mutation carriers and in patients with sporadic HGSOC (*11-14*). While these lesions strongly link the distal fallopian tube to HGSOC pathogenesis, they do not identify which epithelial populations within the tube are most susceptible to transformation, nor how these populations depart from normal differentiation trajectories. Historically, although mature secretory cells have been proposed as likely precursors, recent single-cell studies reveal greater epithelial complexity (*15-18*), suggesting that ciliated-lineage or transitional states may also contribute to malignant evolution.

The fallopian tube consists of four anatomical segments: the infundibulum, ampulla, isthmus, and the uterotubal junction, each exposed to distinct physiological environments. The distal infundibulum and fimbria, positioned adjacent to the ovary, experience repeated bursts of inflammatory and oxidative stress during ovulation, necessitating frequent epithelial repair (*19-22*). Early histological studies described small, round "basal" or "intercalary" cells as potential stem cells capable of generating both ciliated and secretory lineages (*20, 23*), yet their molecular identity remains

uncertain. Advances in single-nucleus RNA sequencing (snRNA-seq) offer a way to

clarify such lineage relationships in human tissues, as this approach avoids

dissociation-induced artifacts, accommodates frozen clinical samples, and produces

high-resolution transcriptomic data suiData File for integration with pathology (*24-27*).

Understanding the epithelial hierarchy of the human fallopian tube and identifying the

populations capable of initiating HGSOC have remained challenging due to limited

access to healthy human tissues, a lack of definitive stem/progenitor markers, and the

absence of lineage-tracing models in women. High-resolution transcriptomic

technologies now allow for reconstruction of epithelial differentiation trajectories and

direct comparison of normal epithelia with malignant states in primary tumors. In this

study, we integrate bulk RNA sequencing of normal fallopian tubes, ovaries, and

HGSOC samples with snRNA-seq of cancer-free fallopian tubes and primary HGSOC

tissues. We define a stem-progenitor hierarchy within the FTE, identifying rare

$LGR5^+/PGR^+$ basal cells as putative stem cells that give rise to a large and widely

distributed pool of $OVGP1^+/RNPC3^+$ progenitor cells. By jointly analyzing normal and

malignant transcriptomes, we show that these progenitors exhibit the strongest

transcriptional and developmental continuity with multiple malignant trajectories, making

them the epithelial populations most poised for oncogenic deviation. We further

delineate transcription factor networks, copy-number alterations, and

microenvironmental signaling pathways that accompany progression from normal

epithelial differentiation to malignant transformation. Rather than asserting a definitive

cell of origin, our findings provide a mechanistic and spatial framework indicating that

fallopian tube progenitors are the most biologically plausible intermediates capable of

evolving into HGSOC under oncogenic, inflammatory, or microenvironmental stress.

These insights refine our understanding of epithelial organization in the fallopian tube

and illuminate the earliest cellular and molecular events that may underlie HGSOC

development, with implications for early detection, risk assessment, and targeted

prevention.

RESULTS

Dual origins of HGSOC identified by bulk RNA-seq

High-grade serous ovarian cancer (HGSOC) has been proposed to arise from either the

fallopian tube epithelium or the ovarian surface epithelium (*11, 28, 29*). To evaluate

these possibilities, we performed bulk RNA-seq to examine transcriptomic relationships

among normal fallopian tubes, normal ovaries, and HGSOC tumors (Fig. 1A). Twenty-

six human specimens were analyzed, including four cancer-free ovaries, four cancer-

free fallopian tubes dissected into three anatomical segments (fimbriae/infundibulum,

ampulla, and isthmus), and ten HGSOC tumors. Cancer-free tissues were collected

from women undergoing surgery for benign uterine conditions, whereas tumor samples

were obtained from patients pathologically diagnosed with epithelial ovarian cancer and

subsequently confirmed as HGSOC (Table S1).

Sequencing produced 38-65 million high-quality reads per sample (Fig. S1A). Principal

component analysis revealed three clearly separated clusters corresponding to ovary,

fallopian tube (FT), and HGSOC tissues (Fig. 1B). Pearson correlation analysis further

highlighted the transcriptional distances among these groups. Notably, eight HGSOC

samples clustered with fallopian tube tissues, whereas two samples (EOC-3 and EOC-4)

clustered with ovaries, suggesting distinct tissue affinities. Hierarchical clustering

confirmed this separation, allowing classification of EOC-3 and EOC-4 as ovary-like

HGSOC (EOC-O) and the remaining eight tumors as fallopian tube-derived HGSOC

(EOC-F) (Fig. 1C; Fig. S1B). Differential gene expression analysis revealed that

fallopian tube-enriched genes, including *FOXJ1, SOX17, PODXL, MUC16*, and *WFDC2,*

were markedly elevated in EOC-F tumors. In contrast, ovarian marker genes, including

FOXL2, ARX, KIT, STAR, and *WFIKKN2,* were predominantly expressed in EOC-O

tumors (Fig. S1C). Gene Ontology (GO) analysis showed enrichment of cell division,

cell adhesion, and spermatogenesis pathways in EOC-F, whereas EOC-O exhibited

signatures associated with gonadal development and WNT signaling (Fig. S1C).

Together, these results support a dual-origin model of HGSOC, with the majority of

tumors in our cohort derived from the fallopian tube.

Clustering of all differentially expressed genes across the ovary, FT, and EOC-F

samples identified seven major gene sets representing core transcriptional programs

(Fig. 1D; Data File S1). Ovarian tissues showed enrichment for genes involved in TGF-

β signaling and gonadal development (Geneset 1), whereas normal FT and ovary

shared immune regulatory and transcriptional programs (Geneset 2). Additional gene

sets reflected smooth muscle components (Geneset 3), FT-specific developmental

signatures, including *PAX2* and *HOX* family genes (Geneset 4), and genes upregulated

in both FT and EOC-F, such as *SPDEF, POU5F1, FOXJ1, HOXB2*, and *HOXB4*

(Geneset 5). Cilium assembly genes (Geneset 6) and cancer-associated pathways

involving proliferation, metabolism, and keratinization (Geneset 7) were also distinctly

represented. Pairwise comparisons among EOC-F, FT, and ovary samples (Fig. S1D–F)

highlighted pathways involved in cell division, chemokine signaling (e.g., *CXCL9/10/11-*

CXCR3, CXCL6-CXCR2), inflammatory responses (*IL1-IL1RAP, GAL-3, NR1H4*),

apoptosis (*IL24, TNFAIP8*), and antigen presentation via MHC class II. Immune-related

genes, such as *HLA-DMA* and *HLA-DRA*, which are key to T-cell recruitment and

differentiation, were strongly expressed, consistent with the inflammatory and immune-

rich microenvironment of HGSOC. These findings underscore the immunologic

complexity of fallopian tube-derived tumors. Collectively, our bulk RNA-seq analyses

establish that HGSOC comprises at least two transcriptionally distinct subtypes, one

resembling fallopian tube epithelium and the other resembling ovarian tissue. The

predominance of the FT-like subtype in our cohort further reinforces the fallopian tube

as the principal tissue of origin for most HGSOCs.

Identification of a large pool of progenitors capable of differentiating into

secretory or ciliated cells in the fallopian tube epithelium

Although several recent single-cell studies have profiled human fallopian tube

epithelium (FTE) (*21, 30-33*), the molecular identity of true FTE stem cells, the hierarchy

of secretory cell subtypes, and the developmental relationship between mature

secretory and ciliated cells remain incompletely defined. To address these uncertainties,

we collected cancer-free fallopian tubes from three perimenopausal women undergoing

hysterectomy with bilateral salpingo-oophorectomy for benign uterine conditions (Table

S2). Each tube was dissected into fimbriae/infundibulum, ampulla, and isthmus, and

single-nucleus RNA sequencing (snRNA-seq) was performed on nine samples.

Using the 10x Genomics platform, we obtained high-quality transcriptomes from 87,920

nuclei after initial quality control, with 36,000-63,000 mean reads per nucleus and

approximately 2,000 genes detected per cell (Fig. S2A). After regressing out cell cycle

effects, mitochondrial gene content, and library size, no cell cycle-driven clustering was

observed (Fig. S2B), and all segments were evenly represented on UMAP, indicating

minimal batch-related segregation (Fig. S2C). These results confirmed the robustness

of the dataset. Initial clustering of all fallopian tube samples identified five major cell

types: secretory cells, ciliated cells, smooth muscle cells, endothelial cells, and

macrophages (Fig. S2D). Marker gene (Fig. S2E) and GO enrichment (Fig. S2F; Data

File S2) analyses suggested that the secretory cell cluster contained the putative stem

cell population. This cluster showed higher expression of genes associated with adult

stem cell maintenance and epithelial progenitor states, including *SOX17, LGR5,*

TFCP2L1, and *HES1*. Endothelial, smooth muscle, ciliated, and macrophage clusters

each expressed expected lineage-specific signatures (Fig. S2F). The distribution of

major epithelial populations across segments showed an increasing proportion of

ciliated cells from the isthmus to the fimbriae, and the inverse pattern for secretory cells

(Fig. S2G). To resolve epithelial heterogeneity at higher resolution, we reclustered

34,267 epithelial cells, yielding 12 distinct epithelial subclusters (Fig. 2A). Hierarchical

clustering grouped these into five broader categories (Fig. 2B). Based on marker

expression, cluster 12 was identified as the FTE stem cell population, defined by high

expression of *LGR5, PGR, PODXL,* and *SOX17*. Clusters 2 and 6 expressed *PAX8,*

ZBTB16, and *EGR1* and were annotated as mature secretory cells. Clusters 1, 4, 5, and

8 represented ciliated cells expressing *CAPS, CFAPs*, and dynein-associated genes.

The remaining clusters, 3, 7, 9, 10, and 11, expressed *OVGP1, CRISP3, RNPC3,* and

BDP1 but lacked stem cell and mature cell markers, indicating they are likely progenitor

cells (Fig. 2C).

RNA velocity analysis demonstrated that progenitor cells occupy transitional positions

between stem cells and both mature lineages, with directional flow toward either

secretory or ciliated fates (Fig. 2D). PAGA confirmed strong connectivity: stem cells

(Cluster 12) positioned at the root, progenitors forming intermediate nodes, and mature

secretory and ciliated cells at opposite termini (Fig. 2E). PHATE visualization revealed

two branching developmental trajectories, supported by pseudotime inference (Fig. 2F–

G). Marker genes such as *LGR5, PAX8, FOXJ1,* and *CFAP54* aligned with expected

pseudotime patterns (Fig. S2H). Transcriptomic dynamics along the secretory trajectory

showed gradual downregulation of stem cell markers (*LGR5, SPDEF, NR2F1, AR*) and

transient induction of estradiol-responsive genes (*POSTN, MAP1B, PTGS2, EZH2*),

followed by increased expression of mature secretory genes, including *EGR1, PAX2,*

PAX8, CD74, and *ZBTB16* (Fig. 2H; Data File S3). Along the ciliated trajectory, early

decreases in *LGR5* and *LAMA5* were followed by transient induction of xenobiotic

response and adhesion-related genes, and terminal enrichment of CFAPs and dyneins

associated with ciliogenesis (Fig. 2I; Data File S4). Quantification of epithelial subtypes

revealed that progenitors comprise the largest epithelial population, ~38% of all

epithelial cells. Stem cells were concentrated in the isthmus and decreased toward the

fimbriae, whereas progenitors remained consistently abundant across segments (Fig.

S2I–J). Immunostaining supported these observations: TUBB4B$^+$ ciliated cells

increased from isthmus to fimbriae, while PAX8$^+$ secretory cells decreased (Fig. S3A–B),

matching snRNA-seq distributions. LGR5 and PGR co-localized in small round basal

cells, confirming their identity as FTE stem cells (Fig. 2J–K; Fig. S3C). Stem cells were

enriched in the isthmus and were rare in the fimbriae, whereas progenitors were

abundant in all segments. Together, these data define a hierarchical epithelial

organization in which rare LGR5$^+$/PGR$^+$ stem cells generate a large, widely distributed

pool of OVGP1$^+$ progenitors that can differentiate into either mature secretory or ciliated

cells. This progenitor population likely represents the key transitional state underlying

both normal epithelial turnover and vulnerability to malignant transformation.

High inter-patient variability and heterogeneity in HGSOC cell composition

High-grade serous ovarian cancer (HGSOC) is characterized by substantial cellular

complexity and inter-patient variability. To define the composition of malignant and non-

malignant cell populations within HGSOC, we analyzed 15,322 nuclei from snRNA-seq

datasets generated from four primary tumors (Fig. S3D). After removing batch effects

and regressing out cell cycle and library size using LIGER (*34*), datasets showed no

residual technical clustering (Fig. S3E), yet pronounced biological heterogeneity across

patients remained evident (Fig. S3F–G). Unsupervised Louvain clustering resolved 13

transcriptionally distinct clusters (Fig. 3A). Hierarchical clustering grouped these into ten

malignant cell clusters and three non-malignant clusters corresponding to macrophages,

cancer-associated fibroblasts (CAFs), and endothelial cells (Fig. 3B). Malignant clusters

represented five major cancer cell states: *proliferative, mesenchymal-like, ciliated-like,*

epithelial-to-mesenchymal transition (EMT), and MHC (immunoreactive) (Fig. 3C).

Cluster-specific marker analysis (Fig. 3D; Data File S5) showed that proliferative

cancer cells strongly expressed genes regulating cell cycle progression and mitosis,

consistent with a tumor stem-like or progenitor-like phenotype. Mesenchymal-like

cancer cells expressed transcription factors linked to cellular plasticity and migration

(*SOX17, MECOM, PAX8*) along with markers of matrix remodeling, suggesting a role in

invasion and stromal engagement. Ciliated-like cancer cells expressed cilium assembly

and motility genes, indicating partial retention of their fallopian tube lineage identity.

EMT-like cancer cells upregulated genes associated with cell motility and extracellular

matrix interactions, reflecting active epithelial-mesenchymal transition. MHC

(immunoreactive) cancer cells displayed robust expression of antigen processing and

MHC class II-related genes, underscoring their interaction with the immune

microenvironment.

Re-clustering of malignant cells alone revealed strong patient-specific patterns (Fig. 3E–

F) that persisted despite computational correction, indicating genuine biological diversity

among tumors. Correlation analyses confirmed substantial transcriptomic divergence

among malignant cell states (Fig. 3G). Several markers typically associated with the

FTE, including *PAX8* and *MUC16*, were broadly expressed across all malignant

subtypes (Fig. S4A). Their expression in matched fallopian tube tissues (Fig. S4B–C)

further reinforced the FT origin of most tumors. Ciliated-like malignant cells expressed

TUBB4B, a hallmark of fallopian tube ciliated epithelium (*22*), consistent with retention

of lineage features (Fig. S4A–B). EMT cells exhibited high levels of *MUC16* and

ARGLU1 (Figs. 3H, S4D), whereas mesenchymal-like cells expressed matrix-

associated genes such as *COL1A1, COL1A2,* and *VCAN* (Fig. 3H), reflecting tumor

core structural phenotypes. Immunoreactive tumor cells expressed *CLU, CD74, HLA-*

DRA, and related immune-interaction genes (Figs. 3H, S4C–D; Data File S6).

Collectively, these analyses demonstrate that HGSOC exhibits striking inter-patient

variability and comprises diverse malignant subpopulations spanning proliferative,

ciliated-like, EMT-like, mesenchymal-like, and immunoreactive states. These findings

establish a foundation for linking specific malignant phenotypes to the differentiation

hierarchy of normal fallopian tube epithelial cells in subsequent analyses.

The FTE progenitor cells as the most likely origin of HGSOC

Multiple lines of evidence, including our bulk RNA-seq findings, support the concept that

HGSOC arises from FTE (*21, 35*). Yet the specific epithelial populations that deviate

from normal lineage progression and initiate carcinogenesis have not been firmly

established. To address this question, we integrated 11,028 malignant cells from

HGSOC samples with 34,267 epithelial cells from normal fallopian tubes, enabling direct

comparison of their transcriptional relationships at single-cell resolution (Fig. 4). Using

force-directed graph embedding (*36*), we identified 15 distinct cellular clusters

encompassing both normal and malignant states (Fig. 4A). Slingshot trajectory analysis

suggested that epithelial stem cells (Cluster 11) give rise to three major lineages:

mature secretory cells (Cluster 1), ciliated cells (Cluster 3), and malignant cells

(Clusters 12 and 13). PAGA connectivity mapping supported these lineage relationships,

revealing strong transitions from stem/progenitor clusters toward both normal and

cancerous endpoints (Fig. 4B). RNA velocity confirmed that epithelial stem cells and

progenitor clusters (0, 4, 8, 11) occupy early pseudotime states and generate

trajectories that culminate in either differentiated FTE cell types or HGSOC cell states

(Fig. 4C). Global pseudotime analysis demonstrated continuity across all epithelial and

cancerous cells, with malignant clusters emerging as late-stage branches diverging

from the stem–progenitor axis (Fig. 4D).

Hierarchical clustering grouped the 15 clusters into three major compartments: ciliated,

secretory, and cancer cell populations (Fig. 4E). Consistent with developmental

positioning, progenitor clusters (4, 0, 8) grouped within the secretory compartment

alongside epithelial stem cells (Cluster 11) and mature secretory cells (Cluster 1).

Although annotated as a ciliated subtype, Cluster 14 exhibited gene expression features

and trajectory placement consistent with malignant identity (Figs. 4B–D, 4F–G).

Similarly, a subset of malignant cells co-clustered with FTE progenitors (Cluster 4),

indicating shared transcriptional programs between early progenitors and cancer-

initiated states. Among cancer clusters, we identified four major malignant phenotypes:

mesenchymal-like (Clusters 10; $VEGFA^+/NEAT1^+$), EMT-like (Cluster 13;

$SPARC^+/COL1A^+$), proliferative (Cluster 9; $TOP2A^+/ASPM^+$), and immunoreactive

(Cluster 12; $CD74^+/HLA\text{-}DRA^+$). Notably, proliferative cancer cells incorporated nuclei

from fimbriae, ampulla, and isthmus, reflecting strong lineage continuity between

progenitors across anatomical regions and the malignant proliferative state (Fig. 4F).

To determine how progenitors diverge toward malignant phenotypes, we analyzed

pseudotime progression along two tumor-forming trajectories. From epithelial stem cells

to immunoreactive cancer cells (Cluster 12), we observed early downregulation of Notch

signaling and cell fate-regulating genes, followed by upregulation of antiapoptotic

pathways, cilium assembly programs, and MHC class II antigen processing, consistent

with immune-interacting tumor states. Late pseudotime was marked by increased

expression of EMT-associated and collagen-organizing genes (Fig. 4H; Data File S7).

Along the trajectory leading to mesenchymal-like cancer cells (Cluster 13), early

pseudotime reflected loss of stemness-associated transcripts, while late pseudotime

showed activation of angiogenesis, integrin signaling, and fibroblast growth pathways,

signatures characteristic of proliferating, invasive tumor cells (Fig. 4I; Data File S8).

These integrated analyses point to $OVGP1^+/RNPC3^+$ progenitor cells, particularly those

enriched in the fimbrial region, as the epithelial population most primed for malignant

transformation. Under normal conditions, progenitors differentiate into either secretory

or ciliated epithelium. Under carcinogenic stress or chronic inflammatory stimuli,

however, these progenitors can instead divert into malignant pathways, giving rise to

immunoreactive or mesenchymal-like tumor states. This dual trajectory mirrors the

molecular heterogeneity observed across HGSOC subtypes.

Transcription factor regulatory networks driving the differentiation from FTE

progenitors into normal epithelial or HGSOC cells

To identify the transcriptional regulators that govern normal fallopian tube epithelial

(FTE) differentiation and malignant transformation, we performed single-cell regulatory

network inference and clustering (SCENIC) on combined datasets of pure FTE cells and

HGSOC malignant cells (Fig. 5A). This analysis defined five major regulon-driven cell

states corresponding to stem cells, progenitors, secretory cells, ciliated cells, and

HGSOC cells. Regulon activity patterns revealed distinct transcription factor (TF)

programs defining each cellular state (Fig. 5B). Ciliated cells were characterized by

strong activation of *NR2F1, FOXJ1,* and *HOXC4* regulons, canonical drivers of

ciliogenesis and epithelial polarization. In contrast, cancer cells exhibited unique activity

of regulons such as *TPI1, NR2F6*, and *MAFB*, consistent with metabolic rewiring,

proliferative advantage, and tumor-associated immune modulation. Stem cells,

progenitors, and mature secretory cells shared substantial overlap in regulon usage,

underscoring the fluidity of transitions among these epithelial compartments. Ranking

regulons based on specificity (AUC scores) identified *SPDEF* as a prominent regulator

in stem and progenitor cells (Fig. 5C). *SPDEF* has previously been implicated in

epithelial differentiation and tumor suppression, suggesting a central role in maintaining

epithelial identity and restraining aberrant expansion (*37-39*). In contrast, *NR2F6*

emerged as one of the most specific and highly active regulators in HGSOC cells (Fig.

5C–D). NR2F6, a member of the nuclear receptor superfamily, is known to influence

hormonal response, immune regulation, and oncogenic processes, and has been

implicated in multiple types of cancer (*40, 41*). Its high activity across malignant clusters

positions it as a potential master regulator of HGSOC cell fate.

Gene regulatory network reconstruction further highlighted SOX5 and SPDEF as master
regulators in stem and progenitor states; PAX2, TBX2, and TCF7L2 in mature secretory
cells; HOXC4 and ZNF214 in ciliated cells; and NR2F6 and TPI1 in HGSOC cells (Fig.
5E; Fig. S5A). These lineage-specific TF modules delineate the molecular pathways
that guide normal epithelial differentiation and, when dysregulated, redirect progenitors
toward malignant trajectories. To functionally validate key findings from SCENIC, we
knocked down NR2F6 in two human ovarian carcinoma cell lines (ES2 and OVCAR3).
NR2F6 suppression significantly reduced colony formation in both lines, indicating
reduced proliferative capacity (Fig. 5F–G). Transwell invasion assays revealed a
marked decrease in invasive potential following NR2F6 knockdown (Fig. 5H–I),
demonstrating that NR2F6 promotes both proliferation and invasion, key hallmarks of
HGSOC progression. Together, these data identify distinct transcriptional regulators
governing FTE differentiation and HGSOC development, spotlighting NR2F6 as a
potential driver of malignant transformation and an attractive therapeutic target.
372

DNA copy number variations in HGSOC
Genomic instability is a defining feature of high-grade serous ovarian cancer (HGSOC).
To characterize large-scale chromosomal alterations associated with malignant
transformation, we inferred DNA copy number variations (CNVs) from snRNA-seq data
using inferCNV (*42, 43*), leveraging normal fallopian tube epithelial (FTE) cells as the
reference population (Fig. 6A). This approach enabled the detection of extensive
chromosomal deletions and amplifications across malignant cells, consistent with the
well-documented genomic chaos that typifies HGSOC.
381

Using a Hidden Markov Model (HMM), we identified 138 CNV regions classified into six
states reflecting complete loss (0x), single-copy loss (0.5x), neutral state (1x), single-
copy gain (1.5x), two-copy gain (2x), and high-level gain modeled as ≥3 copies (3x) (Fig.
6B; Data File S9). Across all malignant nuclei, ~39% of genomic regions were copy-
neutral, 34% exhibited gains, and 27% displayed losses. Although CNVs were
distributed across the genome, Chromosome 1 harbored the highest number of events,
while Chromosome 6 displayed the most pronounced amplitude of alteration (Fig. 6C).

Integration of CNV profiles with gene expression revealed that gene dosage effects

directly influence many hallmark HGSOC genes. Frequently amplified genes, including

S100A6, MUC16, COL1A1, COL1A2, CD74, and *WFDC2*, showed corresponding

overexpression in malignant clusters (Fig. 6D). Conversely, reduced expression of

genes such as *CLU, PODXL,* and *PGR* aligned with copy number loss. These gene-

level dosage effects likely contribute to the biased differentiation of progenitor cells

toward malignant fates rather than terminally differentiated epithelial states. Collectively,

these analyses reveal widespread CNVs in HGSOC and suggest that copy number-

driven transcriptional reprogramming reinforces cancer-promoting pathways. These

genomic alterations may underlie the divergence of progenitors from normal epithelial

maturation trajectories, thereby fostering malignant transformation.

Pivotal role of collagen signaling in the tumor microenvironment of HGSOC

Extensive interactions between malignant cells and surrounding stromal populations

shape the tumor microenvironment (TME) of HGSOC. To delineate these interactions,

we examined ligand-receptor networks among cancer cell subtypes, cancer-associated

fibroblasts (CAFs), endothelial cells, and macrophages. CAFs emerged as the dominant

source of outgoing signals and the most influential regulators of intercellular

communication within the TME (Figs. S5B–C). Among all signaling pathways, collagen

signaling surfaced as the most prominent extracellular matrix (ECM) regulatory axis.

Collagen ligands and their cognate receptors showed extensive cross-talk between

CAFs and malignant cells, consistent with collagen's established roles in matrix

remodeling, cellular adhesion, EMT induction, and tumor progression (*44, 45*) (Figs.

S5D–E). Multiple collagen-encoding genes, including *COL1A1, COL1A2, COL4A1,*

COL4A2, COL6A1, COL6A2, and *COL6A3*, were strongly upregulated in mesenchymal-

like and ciliated-like cancer subtypes as well as in CAFs (Fig. S5F), indicating

coordinated ECM restructuring. Integrin receptors ITGAV and ITGB8, which form key

binding pairs for fibronectin and collagen, were co-expressed in both CAFs and

malignant cells. This bilateral expression suggests a feedback loop in which CAFs and

tumor cells mutually reinforce an ECM-rich, pro-invasive niche. These interactions likely

facilitate tumor cell migration, survival under mechanical stress, and the establishment

of metastatic sites. Together, these findings highlight collagen signaling as a central

orchestrator of stromal-tumor communication in HGSOC. The prominence of collagen-

integrin networks within the TME suggests potential therapeutic opportunities to target

CAF-tumor signaling circuits and disrupt ECM remodeling, immunosuppression, and

tumor progression.

DISCUSSION

High-grade serous ovarian carcinoma (HGSOC) remains the most lethal gynecologic

malignancy, mainly because its earliest cellular precursors are incompletely understood.

Although accumulating evidence implicates the fallopian tube epithelium (FTE),

particularly the distal fimbrial region, as the predominant site of origin for many

HGSOCs (*46-49*), the specific epithelial populations most vulnerable to malignant

transformation have been challenging to define. By integrating bulk RNA sequencing of

normal and malignant tissues with high-resolution single-nucleus transcriptomics of

cancer-free fallopian tubes and primary HGSOC tumors, this study reconstructs the

epithelial hierarchy of the human fallopian tube and identifies epithelial progenitors as

the populations most transcriptionally and developmentally aligned with malignant

trajectories.

Across all donors, we identify a rare LGR5$^+$/PGR$^+$ basal cell population that functions as

a putative epithelial stem cell pool. These cells give rise to a large, widely distributed

population of OVGP1$^+$/RNPC3$^+$ progenitors, which, in turn, generate both secretory and

ciliated epithelial lineages (Fig. 7). This stem-progenitor architecture refines earlier

models that primarily emphasized mature secretory cells as the likely cells of origin for

HGSOC (*21, 29, 46, 50, 51*). Our integrated analysis demonstrates that it is the

progenitor state, not terminal secretory or ciliated states, that shows the strongest

transcriptomic and developmental continuity with multiple malignant phenotypes,

including immunoreactive, proliferative, and mesenchymal-like tumor states. The

transcriptional plasticity, proliferative potential, and intermediate differentiation state of

these progenitors may collectively underlie their susceptibility to malignant deviation.

Transcription factor network inference provides additional mechanistic insight. SPDEF

emerges as a key regulator of progenitor identity, whereas NR2F6 is highlighted as a

candidate driver of malignant lineage divergence. These regulons illustrate how

deviations from the normal progenitor program may initiate transcriptional drift toward

tumor-like states. Yet transcriptional reprogramming alone does not fully explain the

sData File separation between normal epithelial lineages and malignant phenotypes.

HGSOC is characterized by pervasive genomic instability, and our inferred DNA copy

number variation (CNV) profiles reveal recurrent chromosomal gains and losses that

align with oncogenes and malignant regulons. These genomic alterations likely reinforce

the lineage deviations identified at the transcriptional level, consolidating malignant

identity as tumor trajectories diverge from their normal developmental paths.

The spatial distribution of epithelial cell types along the fallopian tube provides important

context for understanding their differential susceptibility to oncogenesis. Our snRNA-seq

analyses show that LGR5$^+$/PGR$^+$ stem cells are most enriched in the isthmus, whereas

OVGP1$^+$/RNPC3$^+$ progenitors form a much larger and more evenly distributed epithelial

population across the ampulla and fimbria. Immunofluorescence staining corroborates

this pattern, revealing that progenitors occupy the luminal surface throughout distal

regions of the tube. These spatial patterns intersect with long-recognized anatomic and

physiologic features of the fimbria, including its close proximity to the ovary and its

repeated exposure to ovulatory follicular fluid. The fimbria is subjected to cyclical

inflammatory microtrauma, bursts of reactive oxygen species, extracellular DNA, and

proteolytic enzymes, all of which are known to induce DNA damage and impair

epithelial homeostasis (*13, 52-54*). It is, therefore, noData File that the region most

frequently harboring serous tubal intraepithelial carcinoma (STIC) lesions also contains

a large pool of transcriptionally plastic progenitor cells. The combination of progenitor

abundance, ovulatory injury, and a distinct extracellular matrix architecture may create a

spatially defined niche in which progenitor states are disproportionately vulnerable to

the earliest oncogenic events. Our ligand-receptor analysis further reveals that collagen-

integrin signaling between cancer-associated fibroblasts (CAFs) and malignant cells

exerts a dominant influence on the HGSOC microenvironment. The fimbrial region's

collagen-rich architecture and dynamic immune milieu may provide additional segment-

specific cues that promote malignant divergence once early transcriptional or genomic

abnormalities arise in progenitor populations.

Our findings complement and extend existing models of HGSOC initiation. Bulk RNA-

seq analyses reveal two transcriptionally distinct tumor groups, one aligning with

ovarian surface epithelium and another with fallopian tube epithelium, consistent with a

dual-origin model. Earlier models posited that fallopian tube secretory epithelial cells

(FTSECs), particularly PAX8[+]/OVGP1[+] cells, serve as the primary cells of origin for

HGSOC (*51, 55, 56*). Still, recent single-cell studies demonstrate substantial secretory

heterogeneity and raise questions about which subtypes are most at risk (*57, 58*). Our

integrated analysis clarifies this paradigm by showing that progenitors, not mature

secretory cells, constitute the most plausible at-risk population (Fig. 7).

Transcriptomic changes along malignant trajectories reveal early tumorigenic features,

including anti-apoptotic signaling, metabolic reprogramming, and ECM remodeling,

consistent with prior models of serous carcinogenesis (*59-61*). These trajectories

converge into well-established HGSOC subtypes with distinct clinical implications (*62,*

63). The tumor microenvironment (TME) plays a central role in shaping malignant

progression. CAFs and immune cells regulate ECM dynamics, angiogenesis, and

immune suppression. Collagen signaling, driven by CAFs and mesenchymal-like tumor

cells, emerged as the dominant ECM pathway in our data, aligning with previous studies

demonstrating that collagen deposition, matrix stiffness, and integrin engagement

promote tumor invasion, EMT, and metastatic potential (*64, 65*). Interactions between

ECM remodeling and tumor-associated macrophages are also well documented (*66-68*),

reinforcing the role of the TME in supporting malignant divergence from progenitor

states.

Several limitations warrant consideration. First, despite strong developmental continuity

between progenitors and malignant states, our data do not constitute direct lineage

proof. Definitive demonstration of cellular origin would require in vivo lineage tracing or

transformation assays using purified human progenitors, approaches currently not

feasible in human tissue. Second, cancer-free fallopian tube samples were obtained

from three perimenopausal donors with benign disease; thus, they may not fully capture

epithelial diversity across ages or genetic backgrounds. Importantly, tissues from

BRCA1/2 mutation carriers, who often harbor early STIC lesions, were unavailable.

Future studies should incorporate prophylactic salpingectomy specimens and early

lesions. Finally, although our analyses suggest strong microenvironmental influences,

spatial transcriptomics and functional assays will be necessary to delineate how stromal

and immune contexts shape progenitor vulnerability.

Despite these limitations, our study provides a comprehensive, mechanistically

grounded framework linking the normal epithelial hierarchy of the fallopian tube to

HGSOC development (Fig. 7). The progenitor-associated markers identified here may

inform early detection, risk stratification, and preventive approaches. Targeting lineage-

informative regulators, such as NR2F6, or microenvironmental pathways, such as

collagen-integrin signaling, may disrupt malignant trajectories rooted in fallopian tube

biology. Together, these results support the emerging paradigm that fallopian tube

progenitors represent biologically and anatomically plausible epithelial intermediates

predisposed to early serous carcinogenesis, offering a foundation for future efforts to

intercept HGSOC before clinical manifestation.

MATERIALS AND METHODS

Study Design

This study was designed to define the epithelial hierarchy of the human fallopian tube

and to determine which epithelial populations show the strongest transcriptional

continuity with high-grade serous ovarian carcinoma (HGSOC). We combined bulk RNA

sequencing of normal fallopian tubes, ovaries, and primary HGSOC samples with

single-nucleus RNA sequencing (snRNA-seq) of cancer-free fallopian tubes and primary

HGSOC tissues. Cancer-free fallopian tubes were obtained from perimenopausal

donors undergoing gynecologic surgery for benign indications, and HGSOC samples

were collected at the time of debulking surgery. No randomization or blinding was

applicable to tissue acquisition because the samples derive from human clinical

specimens. All snRNA-seq datasets were processed using standardized pipelines,

integrated across batches, and analyzed to identify epithelial cell states, developmental

trajectories, and inferred transcription factor regulons. Findings from snRNA-seq were

validated using bulk RNA-seq comparisons, immunofluorescence staining, pathway

analyses, and ligand–receptor modeling. The sample sizes reflect the availability of

high-quality human tissues and are consistent with similar single-cell transcriptomic

studies. The study was exploratory and mechanistic in nature, aimed at establishing a

cellular framework for early serous carcinogenesis rather than testing a prespecified

hypothesis.

Single-Nucleus RNA Sequencing (snRNA-seq)

Single nuclei were isolated from frozen tissues following the **1**0x Genomics Chromium

Next GEM v3.1 protocol. Nuclei were encapsulated, barcoded, and amplified to

construct 3′ gene expression libraries, which were sequenced to approximately 200

million reads per sample on an Illumina NovaSeq platform (LC-BIO). Data were

demultiplexed using the mkfastq function in Cell Ranger v7.1.0, aligned to the hg38

reference provided by 10x Genomics, and quantified using cellranger count (*69*).

Secondary data analyses were performed in Python with Scanpy (v1.8.1) (*70*) or in R

(v.4.1.0) with the Seurat (v.4.3.0) (*71*), SeuratObject (v4.1.3), SeuratWrappers (v0.3.2)

(*71*), tidyverse (v.1.3.2) (*72*), destiny(v3.10.0) (*73*), pheatmap (v1.0.12) packages.

Briefly, outputs of cellranger were first loaded into R by using the Read10X function in

Seurat, and Seurat objects were built from each sample. Nuclei expressing fewer than

200 genes, more than 7,000 genes, or more than 15% mitochondrial reads were

excluded. Data were processed in Seurat using SCTransform for normalization (*74*).

Batch effects were corrected using Harmony and LIGER (*34*), and dimensionality

reduction was performed using the top 3,000 highly variable genes and the first 20

principal components. Clustering was performed using shared nearest-neighbor

modularity optimization, and clusters were annotated based on canonical marker gene

expression. Differential gene expression was evaluated using MAST or Seurat's ROC-

based classifier (avgLog$_2$FC > 0.25, p < 0.01). Lineage trajectories were reconstructed

using PHATE (*75*), diffusion maps (*70, 73*), dpt (*73*), and PAGA (*76*), providing a

continuous representation of epithelial and malignant differentiation.

More detailed methods are available in the supplementary material file.

Statistical Analysis

All statistical analyses were performed using R, Python, Seurat, Scanpy, and

associated packages. For snRNA-seq, quality filtering thresholds, normalization

methods, dimensionality reduction, integration strategies, and clustering parameters

were applied uniformly across samples using edgeR (*77, 78*). Differential gene

expression between clusters or conditions was calculated using the Wilcoxon rank-sum

test with Benjamini–Hochberg false discovery rate (FDR) correction; genes with

adjusted *P* < 0.05 and |log$_2$FC| ≥ defined thresholds were considered significant. Gene

set enrichment analyses used GSEA or Reactome with FDR correction. Inferred copy

number variation (CNV) analyses applied a six-state Hidden Markov Model, and RNA

velocity analyses used both stochastic and dynamical models for robustness. PAGA

connectivity, lineage trajectories, regulon activity (SCENIC), and ligand–receptor

interactions (CellChat/NATMI) were evaluated using default or recommended settings.

Sensitivity analyses comparing alternative normalization and integration strategies

(SCTransform, log-normalization, Harmony, LIGER, Seurat Integration) confirmed the

stability of clustering, velocity patterns, and PAGA structure. No statistical methods

were used to predetermine sample size due to reliance on human clinical specimens.

All statistical tests were two-sided unless otherwise specified. Quantification of

immunohistochemistry and immunofluorescence data was conducted using two-tailed

Student's t-tests. Data are presented as mean ± SEM, and p-values <0.05, <0.01, and

<0.001 indicate increasing levels of significance.

List of Supplementary Materials

Materials and Methods

Figs. S1 to S5

Data File s S1 to S2

Data files S1 to S9 (Excel files)

References 1-9 (numbers for references only cited in SM)
613

REFERENCES

1. R. J. Kurman, M. Shih Ie, The origin and pathogenesis of epithelial ovarian cancer: a proposed unifying theory. *Am J Surg Pathol* **34**, 433-443 (2010).

2. D. D. Bowtell, S. Bohm, A. A. Ahmed, P. J. Aspuria, R. C. Bast, Jr., V. Beral, J. S. Berek, M. J. Birrer, S. Blagden, M. A. Bookman, J. D. Brenton, K. B. Chiappinelli, F. C. Martins, G. Coukos, R. Drapkin, R. Edmondson, C. Fotopoulou, H. Gabra, J. Galon, C. Gourley, V. Heong, D. G. Huntsman, M. Iwanicki, B. Y. Karlan, A. Kaye, E. Lengyel, D. A. Levine, K. H. Lu, I. A. McNeish, U. Menon, S. A. Narod, B. H. Nelson, K. P. Nephew, P. Pharoah, D. J. Powell, Jr., P. Ramos, I. L. Romero, C. L. Scott, A. K. Sood, E. A. Stronach, F. R. Balkwill, Rethinking ovarian cancer II: reducing mortality from high-grade serous ovarian cancer. *Nat Rev Cancer* **15**, 668-679 (2015).

3. K. M. Feeley, M. Wells, Precursor lesions of ovarian epithelial malignancy. *Histopathology* **38**, 87-95 (2001).

4. R. J. Kurman, M. Shih Ie, Molecular pathogenesis and extraovarian origin of epithelial ovarian cancer--shifting the paradigm. *Hum Pathol* **42**, 918-931 (2011).

5. T. T. Gong, S. Guo, F. H. Liu, Y. L. Huo, M. Zhang, S. Yan, H. X. Zhou, X. Pan, X. Y. Wang, H. L. Xu, Y. Kang, Y. Z. Li, X. Qin, Q. Xiao, D. H. Huang, X. Y. Li, Y. Y. Zhao, X. X. Zhao, Y. L. Wang, X. X. Ma, S. Gao, Y. H. Zhao, S. W. Ning, Q. J. Wu, Proteomic characterization of epithelial ovarian cancer delineates molecular signatures and therapeutic targets in distinct histological subtypes. *Nat Commun* **14**, 7802 (2023).

6. F. Coscia, K. M. Watters, M. Curtis, M. A. Eckert, C. Y. Chiang, S. Tyanova, A. Montag, R. R. Lastra, E. Lengyel, M. Mann, Integrative proteomic profiling of ovarian cancer cell lines reveals precursor cell associated proteins and functional status. *Nat Commun* **7**, 12645 (2016).

7. R. Bhattacharya, A. Ghosh, S. Mukhopadhyay, High-grade serous ovarian carcinoma, the "Achiles' hill" for clinicians and molecular biologists: a molecular insight. *Mol Biol Rep* **50**, 9511-9519 (2023).

8. E. K. Colvin, V. M. Howell, Why the dual origins of high grade serous ovarian cancer matter. *Nat Commun* **11**, 1200 (2020).

9. E. Kuhn, R. J. Kurman, R. Vang, A. S. Sehdev, G. Han, R. Soslow, T. L. Wang, M. Shih Ie, TP53 mutations in serous tubal intraepithelial carcinoma and concurrent pelvic high-grade serous carcinoma--evidence supporting the clonal relationship of the two lesions. *J Pathol* **226**, 421-426 (2012).

10. J. Kraiczy, B. Yu, Human fallopian tube epithelial organoids with TP53 mutation recapitulate features of serous tubal intraepithelial carcinoma (STIC). *Gynecol Oncol* **203**, 198-208 (2025).

11. S. I. Labidi-Galy, E. Papp, D. Hallberg, N. Niknafs, V. Adleff, M. Noe, R. Bhattacharya, M. Novak, S. Jones, J. Phallen, C. A. Hruban, M. S. Hirsch, D. I. Lin, L. Schwartz, C. L. Maire, J. C. Tille, M. Bowden, A. Ayhan, L. D. Wood, R. B. Scharpf, R. Kurman, T. L. Wang, I. M. Shih, R. Karchin, R. Drapkin, V. E. Velculescu, High grade serous ovarian carcinomas originate in the fallopian tube. *Nat Commun* **8**, 1093 (2017).

12. C. Poon, S. Hyde, P. Grant, M. Newman, K. Ireland Jenkin, Incidence and Characteristics of Unsuspected Neoplasia Discovered in High-Risk Women

Undergoing Risk Reductive Bilateral Salpingooophorectomy. *Int J Gynecol Cancer* **26**, 1415-1420 (2016).

13. D. Luvero, R. Angioli, E. Notaro, F. Plotti, C. Terranova, A. M. Angioli, A. Festa, A. Stermasi, S. Manco, M. Diserio, R. Montera, Serous Tubal Intraepithelial Carcinoma (STIC): A Review of the Literature on the Incidence at the Time of Prophylactic Surgery. *Diagnostics (Basel)* **14**, (2024).

14. F. Chen, K. Gaitskell, M. J. Garcia, A. Albukhari, J. Tsaltas, A. A. Ahmed, Serous tubal intraepithelial carcinomas associated with high-grade serous ovarian carcinomas: a systematic review. *BJOG* **124**, 872-878 (2017).

15. Q. Hao, J. Li, Q. Zhang, F. Xu, B. Xie, H. Lu, X. Wu, X. Zhou, Single-cell transcriptomes reveal heterogeneity of high-grade serous ovarian carcinoma. *Clin Transl Med* **11**, e500 (2021).

16. C. Chai, L. Liang, N. S. Mikkelsen, W. Wang, W. Zhao, C. Sun, R. O. Bak, H. Li, L. Lin, F. Wang, Y. Luo, Single-cell transcriptome analysis of epithelial, immune, and stromal signatures and interactions in human ovarian cancer. *Commun Biol* **7**, 131 (2024).

17. J. Xu, Y. Fang, K. Chen, S. Li, S. Tang, Y. Ren, Y. Cen, W. Fei, B. Zhang, Y. Shen, W. Lu, Single-Cell RNA Sequencing Reveals the Tissue Architecture in Human High-Grade Serous Ovarian Cancer. *Clin Cancer Res* **28**, 3590-3602 (2022).

18. B. J. Winterhoff, M. Maile, A. K. Mitra, A. Sebe, M. Bazzaro, M. A. Geller, J. E. Abrahante, M. Klein, R. Hellweg, S. A. Mullany, K. Beckman, J. Daniel, T. K. Starr, Single cell sequencing reveals heterogeneity within ovarian cancer epithelium and cancer associated stromal cells. *Gynecol Oncol* **144**, 598-606 (2017).

19. S. M. King, T. S. Hilliard, L. Y. Wu, R. C. Jaffe, A. T. Fazleabas, J. E. Burdette, The impact of ovulation on fallopian tube epithelial cells: evaluating three hypotheses connecting ovulation and serous ovarian cancer. *Endocrine-related cancer* **18**, 627-642 (2011).

20. D. Y. Paik, D. M. Janzen, A. M. Schafenacker, V. S. Velasco, M. S. Shung, D. Cheng, J. Huang, O. N. Witte, S. Memarzadeh, Stem-like epithelial cells are concentrated in the distal end of the fallopian tube: a site for injury and serous cancer initiation. *Stem cells (Dayton, Ohio)* **30**, 2487-2497 (2012).

21. H. Q. Dinh, X. Lin, F. Abbasi, R. Nameki, M. Haro, C. E. Olingy, H. Chang, L. Hernandez, S. A. Gayther, K. N. Wright, P. J. Aspuria, B. Y. Karlan, R. I. Corona, A. Li, B. J. Rimel, M. T. Siedhoff, F. Medeiros, K. Lawrenson, Single-cell transcriptomics identifies gene expression networks driving differentiation and tumorigenesis in the human fallopian tube. *Cell reports* **35**, 108978 (2021).

22. S. Yuan, Z. Wang, H. Peng, S. M. Ward, G. W. Hennig, H. Zheng, W. Yan, Oviductal motile cilia are essential for oocyte pickup but dispensable for sperm and embryo transport. *Proceedings of the National Academy of Sciences of the United States of America* **118**, (2021).

23. M. Zhu, T. Iwano, S. Takeda, Fallopian Tube Basal Stem Cells Reproducing the Epithelial Sheets In Vitro-Stem Cell of Fallopian Epithelium. *Biomolecules* **10**, (2020).

24. M. Slyper, C. B. M. Porter, O. Ashenberg, J. Waldman, E. Drokhlyansky, I. Wakiro, C. Smillie, G. Smith-Rosario, J. Wu, D. Dionne, S. Vigneau, J. Jané-Valbuena, T. L. Tickle, S. Napolitano, M. J. Su, A. G. Patel, A. Karlstrom, S. Gritsch, M.

Nomura, A. Waghray, S. H. Gohil, A. M. Tsankov, L. Jerby-Arnon, O. Cohen, J. Klughammer, Y. Rosen, J. Gould, L. Nguyen, M. Hofree, P. J. Tramontozzi, B. Li, C. J. Wu, B. Izar, R. Haq, F. S. Hodi, C. H. Yoon, A. N. Hata, S. J. Baker, M. L. Suvà, R. Bueno, E. H. Stover, M. R. Clay, M. A. Dyer, N. B. Collins, U. A. Matulonis, N. Wagle, B. E. Johnson, A. Rotem, O. Rozenblatt-Rosen, A. Regev, A single-cell and single-nucleus RNA-Seq toolbox for fresh and frozen human tumors. *Nature medicine* **26**, 792-802 (2020).

25. H. Wu, Y. Kirita, E. L. Donnelly, B. D. Humphreys, Advantages of Single-Nucleus over Single-Cell RNA Sequencing of Adult Kidney: Rare Cell Types and Novel Cell States Revealed in Fibrosis. *Journal of the American Society of Nephrology : JASN* **30**, 23-32 (2019).

26. G. Wiegleb, S. Reinhardt, A. Dahl, N. Posnien, Tissue dissociation for single-cell and single-nuclei RNA sequencing for low amounts of input material. *Frontiers in zoology* **19**, 27 (2022).

27. J. Ding, X. Adiconis, S. K. Simmons, M. S. Kowalczyk, C. C. Hession, N. D. Marjanovic, T. K. Hughes, M. H. Wadsworth, T. Burks, L. T. Nguyen, J. Y. H. Kwon, B. Barak, W. Ge, A. J. Kedaigle, S. Carroll, S. Li, N. Hacohen, O. Rozenblatt-Rosen, A. K. Shalek, A. C. Villani, A. Regev, J. Z. Levin, Systematic comparison of single-cell and single-nucleus RNA-sequencing methods. *Nature biotechnology* **38**, 737-746 (2020).

28. J. Kim, E. Y. Park, O. Kim, J. M. Schilder, D. M. Coffey, C. H. Cho, R. C. Bast, Jr., Cell Origins of High-Grade Serous Ovarian Cancer. *Cancers (Basel)* **10**, (2018).

29. S. Zhang, I. Dolgalev, T. Zhang, H. Ran, D. A. Levine, B. G. Neel, Both fallopian tube and ovarian surface epithelium are cells-of-origin for high-grade serous ovarian carcinoma. *Nat Commun* **10**, 5367 (2019).

30. Z. Hu, M. Artibani, A. Alsaadi, N. Wietek, M. Morotti, T. Shi, Z. Zhong, L. Santana Gonzalez, S. El-Sahhar, E. M. Carrami, G. Mallett, Y. Feng, K. Masuda, Y. Zheng, K. Chong, S. Damato, S. Dhar, L. Campo, R. Garruto Campanile, H. Soleymani Majd, V. Rai, D. Maldonado-Perez, S. Jones, V. Cerundolo, T. Sauka-Spengler, C. Yau, A. A. Ahmed, The Repertoire of Serous Ovarian Cancer Non-genetic Heterogeneity Revealed by Single-Cell Sequencing of Normal Fallopian Tube Epithelial Cells. *Cancer Cell* **37**, 226-242 e227 (2020).

31. E. Lengyel, Y. Li, M. Weigert, L. Zhu, H. Eckart, M. Javellana, S. Ackroyd, J. Xiao, S. Olalekan, D. Glass, S. Iyer, R. Krishnan, A. J. Bilecz, R. Lastra, M. Chen, A. Basu, A molecular atlas of the human postmenopausal fallopian tube and ovary from single-cell RNA and ATAC sequencing. *Cell Rep* **41**, 111838 (2022).

32. N. D. Ulrich, Y. C. Shen, Q. Ma, K. Yang, D. F. Hannum, A. Jones, J. Machlin, J. F. Randolph, Jr., Y. R. Smith, S. B. Schon, A. Shikanov, E. E. Marsh, R. Lieberman, S. J. Gurczynski, B. B. Moore, J. Z. Li, S. Hammoud, Cellular heterogeneity of human fallopian tubes in normal and hydrosalpinx disease states identified using scRNA-seq. *Dev Cell* **57**, 914-929 e917 (2022).

33. J. Brand, M. Haro, X. Lin, B. J. Rimel, S. M. McGregor, K. Lawrenson, H. Q. Dinh, Fallopian tube single cell analysis reveals myeloid cell alterations in high-grade serous ovarian cancer. *iScience* **27**, 108990 (2024).

34. J. Liu, C. Gao, J. Sodicoff, V. Kozareva, E. Z. Macosko, J. D. Welch, Jointly
defining cell types from multiple single-cell datasets using LIGER. *Nat Protoc* **15**,
3632-3662 (2020).

35. Z. Hu, M. Artibani, A. Alsaadi, N. Wietek, M. Morotti, T. Shi, Z. Zhong, L. Santana
Gonzalez, S. El-Sahhar, E. M. Carrami, G. Mallett, Y. Feng, K. Masuda, Y. Zheng,
K. Chong, S. Damato, S. Dhar, L. Campo, R. Garruto Campanile, H. Soleymani
Majd, V. Rai, D. Maldonado-Perez, S. Jones, V. Cerundolo, T. Sauka-Spengler, C.
Yau, A. A. Ahmed, The Repertoire of Serous Ovarian Cancer Non-genetic
Heterogeneity Revealed by Single-Cell Sequencing of Normal Fallopian Tube
Epithelial Cells. *Cancer cell* **37**, 226-242.e227 (2020).

36. M. Jacomy, T. Venturini, S. Heymann, M. Bastian, ForceAtlas2, a continuous
graph layout algorithm for handy network visualization designed for the Gephi
software. *PLoS One* **9**, e98679 (2014).

37. X. H. Cheng, M. Black, V. Ustiyan, T. Le, L. Fulford, A. Sridharan, M. Medvedovic,
V. V. Kalinichenko, J. A. Whitsett, T. V. Kalin, SPDEF inhibits prostate
carcinogenesis by disrupting a positive feedback loop in regulation of the Foxm1
oncogene. *PLoS Genet* **10**, e1004656 (2014).

38. T. K. Noah, Y. H. Lo, A. Price, G. Chen, E. King, M. K. Washington, B. J. Aronow,
N. F. Shroyer, SPDEF functions as a colorectal tumor suppressor by inhibiting
beta-catenin activity. *Gastroenterology* **144**, 1012-1023 e1016 (2013).

39. Y. H. Lo, E. Chung, Z. Li, Y. W. Wan, M. M. Mahe, M. S. Chen, T. K. Noah, K. N.
Bell, H. K. Yalamanchili, T. J. Klisch, Z. Liu, J. S. Park, N. F. Shroyer,
Transcriptional Regulation by ATOH1 and its Target SPDEF in the Intestine. *Cell
Mol Gastroenterol Hepatol* **3**, 51-71 (2017).

40. L. Wang, H. Long, Q. Zheng, X. Bo, X. Xiao, B. Li, Circular RNA circRHOT1
promotes hepatocellular carcinoma progression by initiation of NR2F6
expression. *Mol Cancer* **18**, 119 (2019).

41. J. Liu, T. Li, X. L. Liu, DDA1 is induced by NR2F6 in ovarian cancer and predicts
poor survival outcome. *Eur Rev Med Pharmacol Sci* **21**, 1206-1213 (2017).

42. A. De Falco, F. Caruso, X. D. Su, A. Iavarone, M. Ceccarelli, A variational
algorithm to detect the clonal copy number substructure of tumors from scRNA-
seq data. *Nat Commun* **14**, 1074 (2023).

43. A. Shlien, D. Malkin, Copy number variations and cancer. *Genome Med* **1**, 62
(2009).

44. X. Cui, T. Shan, L. Qiao, Collagen type IV alpha 1 (COL4A1) silence hampers the
invasion, migration and epithelial-mesenchymal transition (EMT) of gastric
cancer cells through blocking Hedgehog signaling pathway. *Bioengineered* **13**,
8972-8981 (2022).

45. H. Huang, S. Wright, J. Zhang, R. A. Brekken, Getting a grip on adhesion:
Cadherin switching and collagen signaling. *Biochim Biophys Acta Mol Cell Res*
1866, 118472 (2019).

46. P. T. Kroeger, Jr., R. Drapkin, Pathogenesis and heterogeneity of ovarian cancer.
Curr Opin Obstet Gynecol **29**, 26-34 (2017).

47. T. M. Bergsten, J. E. Burdette, M. Dean, Fallopian tube initiation of high grade
serous ovarian cancer and ovarian metastasis: Mechanisms and therapeutic
implications. *Cancer Lett* **476**, 152-160 (2020).

48. W. G. McCluggage, L. Hirschowitz, C. B. Gilks, N. Wilkinson, N. Singh, The
Fallopian Tube Origin and Primary Site Assignment in Extrauterine High-grade
Serous Carcinoma: Findings of a Survey of Pathologists and Clinicians. *Int J*
Gynecol Pathol **36**, 230-239 (2017).
49. S. H. George, R. Garcia, B. M. Slomovitz, Ovarian Cancer: The Fallopian Tube
as the Site of Origin and Opportunities for Prevention. *Front Oncol* **6**, 108 (2016).
50. P. Galhenage, Y. Zhou, E. Perry, B. Loc, K. Fietz, S. Iyer, F. Reinhardt, T. Da Silva,
V. Botchkarev, J. Chen, C. P. Crum, R. A. Weinberg, S. Pathania, Replication
stress and defective checkpoints make fallopian tube epithelial cells putative
drivers of high-grade serous ovarian cancer. *Cell Rep* **42**, 113144 (2023).
51. N. Y. Wu, C. Fang, H. S. Huang, J. Wang, T. Y. Chu, Natural history of ovarian
high-grade serous carcinoma from time effects of ovulation inhibition and
progesterone clearance of p53-defective lesions. *Mod Pathol* **33**, 29-37 (2020).
52. D. Wloga, E. Joachimiak, A. Osinka, S. Ahmadi, S. Majhi, Motile Cilia in Female
and Male Reproductive Tracts and Fertility. *Cells* **13**, (2024).
53. Y. W. Chien, Y. Wang, P. Huang, B. C. Lawson, D. L. Kolin, M. H. Chui, R. Vang,
T. A. Numan, T. R. Soong, B. G. Wang, S. A. Smith, C. L. Chen, R. Stone, C.
Douville, T. L. Wang, I. M. Shih, Morphologic and Molecular Heterogeneity of
High-grade Serous Carcinoma Precursor Lesions. *Am J Surg Pathol* **48**, 475-486
(2024).
54. V. Popat, E. Han, From Fallopian Tube to Ovarian Cancer: Understanding the
Evaluation and Management of Serous Tubal Intraepithelial Carcinoma Lesions.
Curr Treat Options Oncol **26**, 910-919 (2025).
55. K. Nakamura, K. Nakayama, N. Ishikawa, M. Ishikawa, R. Sultana, T. Kiyono, S.
Kyo, Reconstitution of high-grade serous ovarian carcinoma from primary
fallopian tube secretory epithelial cells. *Oncotarget* **9**, 12609-12619 (2018).
56. J. Mei, H. Tian, H. S. Huang, C. F. Hsu, Y. Liou, N. Wu, W. Zhang, T. Y. Chu,
Cellular models of development of ovarian high-grade serous carcinoma: A
review of cell of origin and mechanisms of carcinogenesis. *Cell Prolif* **54**, e13029
(2021).
57. Z. Lin, S. Liang, Z. Pu, Z. Zou, L. He, C. J. Lyon, Y. Zhang, T. Y. Hu, M. Wu,
Phenotypic plasticity and secretory heterogeneity in subpopulations derived from
single cancer cell. *Acta Pharm Sin B* **15**, 2723-2735 (2025).
58. Z. Kang, J. Wang, W. Huang, J. Liu, W. Yan, Identification of Transcriptional
Heterogeneity and Construction of a Prognostic Model for Melanoma Based on
Single-Cell and Bulk Transcriptome Analysis. *Front Cell Dev Biol* **10**, 874429
(2022).
59. A. H. Lee, C. Mejia Pena, M. R. Dawson, Comparing the Secretomes of
Chemorefractory and Chemoresistant Ovarian Cancer Cell Populations. *Cancers*
(Basel) **14**, (2022).
60. C. Liu, W. Liu, J. Huang, Z. Wu, W. Li, B. Chen, Y. Yang, H. Lin, Y. Xu, Metabolic
Reprogramming Shapes the Progression and Therapeutic Landscape of Ovarian
Cancer. *Cancer Manag Res* **17**, 1707-1722 (2025).
61. E. H. Puttock, E. J. Tyler, M. Manni, E. Maniati, C. Butterworth, M. Burger Ramos,
E. Peerani, P. Hirani, V. Gauthier, Y. Liu, G. Maniscalco, V. Rajeeve, P. Cutillas, C.
Trevisan, M. Pozzobon, M. Lockley, J. Rastrick, H. Laubli, A. White, O. M. T.

Pearce, Extracellular matrix educates an immunoregulatory tumor macrophage
phenotype found in ovarian cancer metastasis. *Nat Commun* **14**, 2514 (2023).
62. W. Kassuhn, O. Klein, S. Darb-Esfahani, H. Lammert, S. Handzik, E. T. Taube, W.
D. Schmitt, C. Keunecke, D. Horst, F. Dreher, J. George, D. D. Bowtell, O. Dorigo,
M. Hummel, J. Sehouli, N. Bluthgen, H. Kulbe, E. I. Braicu, Classification of
Molecular Subtypes of High-Grade Serous Ovarian Cancer by MALDI-Imaging.
Cancers (Basel) **13**, (2021).
63. M. Khashaba, M. Fawzy, A. Abdel-Aziz, G. Eladawei, R. Nagib, Subtyping of high
grade serous ovarian carcinoma: histopathological and immunohistochemical
approach. *J Egypt Natl Canc Inst* **34**, 6 (2022).
64. S. Natarajan, K. M. Foreman, M. I. Soriano, N. S. Rossen, H. Shehade, D. R.
Fregoso, J. T. Eggold, V. Krishnan, O. Dorigo, A. J. Krieg, S. C. Heilshorn, S.
Sinha, K. C. Fuh, E. B. Rankin, Collagen Remodeling in the Hypoxic Tumor-
Mesothelial Niche Promotes Ovarian Cancer Metastasis. *Cancer Res* **79**, 2271-
2284 (2019).
65. H. Fujimoto, M. Yoshihara, R. Rodgers, S. Iyoshi, K. Mogi, E. Miyamoto, S.
Hayakawa, M. Hayashi, S. Nomura, K. Kitami, K. Uno, M. Sugiyama, Y. Koya, Y.
Yamakita, A. Nawa, A. Enomoto, C. Ricciardelli, H. Kajiyama, Tumor-associated
fibrosis: a unique mechanism promoting ovarian cancer metastasis and
peritoneal dissemination. *Cancer Metastasis Rev* **43**, 1037-1053 (2024).
66. J. Xiong, R. Xiao, J. Zhao, Q. Zhao, M. Luo, F. Li, W. Zhang, M. Wu, Matrix
stiffness affects tumor-associated macrophage functional polarization and its
potential in tumor therapy. *J Transl Med* **22**, 85 (2024).
67. K. X. Yu, W. J. Yuan, H. Z. Wang, Y. X. Li, Extracellular matrix stiffness and
tumor-associated macrophage polarization: new fields affecting immune
exclusion. *Cancer Immunol Immunother* **73**, 115 (2024).
68. W. Du, X. Xia, F. Hu, J. Yu, Extracellular matrix remodeling in the tumor immunity.
Front Immunol **14**, 1340634 (2023).
69. G. X. Zheng, J. M. Terry, P. Belgrader, P. Ryvkin, Z. W. Bent, R. Wilson, S. B.
Ziraldo, T. D. Wheeler, G. P. McDermott, J. Zhu, M. T. Gregory, J. Shuga, L.
Montesclaros, J. G. Underwood, D. A. Masquelier, S. Y. Nishimura, M. Schnall-
Levin, P. W. Wyatt, C. M. Hindson, R. Bharadwaj, A. Wong, K. D. Ness, L. W.
Beppu, H. J. Deeg, C. McFarland, K. R. Loeb, W. J. Valente, N. G. Ericson, E. A.
Stevens, J. P. Radich, T. S. Mikkelsen, B. J. Hindson, J. H. Bielas, Massively
parallel digital transcriptional profiling of single cells. *Nat Commun* **8**, 14049
(2017).
70. F. A. Wolf, P. Angerer, F. J. Theis, SCANPY: large-scale single-cell gene
expression data analysis. *Genome Biol* **19**, 15 (2018).
71. Y. Hao, S. Hao, E. Andersen-Nissen, W. M. Mauck, 3rd, S. Zheng, A. Butler, M. J.
Lee, A. J. Wilk, C. Darby, M. Zager, P. Hoffman, M. Stoeckius, E. Papalexi, E. P.
Mimitou, J. Jain, A. Srivastava, T. Stuart, L. M. Fleming, B. Yeung, A. J. Rogers, J.
M. McElrath, C. A. Blish, R. Gottardo, P. Smibert, R. Satija, Integrated analysis of
multimodal single-cell data. *Cell* **184**, 3573-3587 e3529 (2021).
72. H. Wickham, M. Averick, J. Bryan, Chang, W., , L. McGowan, R. François, G.
Grolemund, A. Hayes, L. Henry, J. Hester, , Welcome to the Tidyverse. *JOSS* **4**,
1686 (2019).

73. P. Angerer, L. Haghverdi, M. Büttner, F. J. Theis, C. Marr, F. Buettner, destiny:
diffusion maps for large-scale single-cell data in R. . *Bioinformatics* **32**, 1241–
1243 (2016).
74. C. Hafemeister, R. Satija, Normalization and variance stabilization of single-cell
RNA-seq data using regularized negative binomial regression. *Genome Biol* **20**,
296 (2019).
75. K. R. Moon, D. van Dijk, Z. Wang, S. Gigante, D. B. Burkhardt, W. S. Chen, K.
Yim, A. V. D. Elzen, M. J. Hirn, R. R. Coifman, N. B. Ivanova, G. Wolf, S.
Krishnaswamy, Visualizing structure and transitions in high-dimensional
biological data. *Nat Biotechnol* **37**, 1482-1492 (2019).
76. F. A. Wolf, F. K. Hamey, M. Plass, J. Solana, J. S. Dahlin, B. Gottgens, N.
Rajewsky, L. Simon, F. J. Theis, PAGA: graph abstraction reconciles clustering
with trajectory inference through a topology preserving map of single cells.
Genome Biol **20**, 59 (2019).
77. M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package
for differential expression analysis of digital gene expression data. *Bioinformatics*
26, 139-140 (2010).
78. Y. Chen, A. T. Lun, G. K. Smyth, From reads to genes to pathways: differential
expression analysis of RNA-Seq experiments using Rsubread and the edgeR
quasi-likelihood pipeline. *F1000Res* **5**, 1438 (2016).

ACKNOWLEDGMENTS

We thank LC-BIO Co., Ltd. (Hangzhou, China) for performing the Chromium 10✕

Genomics single-nucleus RNA-sequencing and Genedenovo Biotechnology Co., Ltd.

(Guangzhou, China) for conducting the Illumina NovaSeq 6000 bulk RNA-seq.

Funding

Startup funds from The Lundquist Institute for Biomedical Innovation at Harbor-UCLA

Medical Center (WY).

Internal research funding from China Medical University (QL).

Startup funds from the Fourth Affiliated Hospital and International School of Medicine,

Zhejiang University (KC).

Competing Interests

Authors declare that they have no competing interests.

Data and Code Availability

Bulk RNA-seq and single-nucleus RNA-seq (snRNA-seq) datasets have been deposited

in the NCBI Gene Expression Omnibus under accession number GSE223426. All

custom analysis scripts, computational workflows, and software settings used in this

study are openly accessible at https://github.com/chengkeren/fallopiantube, and no

restrictions apply to their use. Software versions, computational dependencies, and

scripts used in data processing and analysis are documented comprehensively in the

GitHub repository (https://github.com/chengkeren/fallopiantube).

FIGURE LEGENDS

Figure 1. Transcriptomic relationships among human fallopian tubes, ovaries, and high-grade serous ovarian cancer.

(A) Overview of human samples used for bulk RNA-seq, including cancer-free ovaries, cancer-free fallopian tube segments (isthmus, ampulla, fimbria), and high-grade serous ovarian cancer (HGSOC) tumors.

(B) Principal component analysis of bulk RNA-seq data showing three distinct clusters corresponding to ovary, fallopian tube, and HGSOC.

(C) Hierarchical clustering and heatmap of pairwise Pearson correlations identifying two HGSOC subgroups: fallopian tube–like (EOC_F) and ovary-like (EOC_O).

(D) Heatmap of seven major gene sets differentially enriched among ovary, fallopian tube, and EOC_F samples. Representative genes for each set are shown. See also **Data File S1**.

Figure 2. Cellular composition and differentiation hierarchy of the human fallopian tube epithelium revealed by snRNA-seq.

(A) UMAP visualization of 34,267 epithelial nuclei from human fallopian tubes, identifying 12 epithelial subclusters.

(B) Hierarchical clustering of subclusters grouped into five major epithelial categories.

(C) Dot plot showing marker genes defining stem cells, progenitors, secretory cells, and ciliated cells.

(D) RNA velocity fields projected onto UMAP indicate directional flow from stem cells toward either secretory or ciliated lineages.

(E) PAGA graph depicting connectivity among epithelial states, placing stem cells at the root.

(F–G) PHATE embeddings and pseudotime trajectories showing two major differentiation paths: secretory and ciliated.

(H–I) Heatmaps of dynamic gene expression along secretory (H) and ciliated (I) trajectories.

(J–K) Immunofluorescence localization of LGR5 and PGR in basal epithelial stem cells

of the fallopian tube. Scale bars: 50 μm.

See also **Figure S2** and **Data File S2–S4**.

Figure 3. Cellular diversity and inter-patient heterogeneity in human HGSOC.

(A) UMAP plot of 15,322 nuclei from four HGSOC tumors identifying 13 cellular clusters.

(B) Dendrogram showing three non-malignant clusters (macrophages, cancer-

associated fibroblasts (CAFs), endothelial cells) and five major HGSOC cell states.

(C) Annotation of cancer subtypes: proliferative, mesenchymal-like, epithelial-to-

mesenchymal transition (EMT), ciliated-like, and immunoreactive (MHC).

(D) Dot plot of representative marker genes across cancer subtypes.

(E–F) UMAP embeddings highlighting strong inter-patient heterogeneity among

malignant cells.

(G) Correlation matrix demonstrating transcriptional divergence among malignant

subtypes.

(H) Expression patterns of markers defining proliferative, EMT, mesenchymal-like,

ciliated-like, and immunoreactive cancer cell states.

See also **Figure S3–S4** and **Data File S5–S6**.

Figure 4. Lineage relationships linking fallopian tube epithelial progenitors to

HGSOC malignant states.

(A) Force-directed graph layout showing 45,295 combined epithelial and cancer nuclei,

identifying 15 integrated clusters.

(B) PAGA analysis reveals three major lineages from epithelial stem cells: secretory

cells, ciliated cells, and malignant cells.

(C) RNA velocity projection demonstrating early progenitors giving rise to either

differentiated epithelial states or cancer trajectories.

(D) Pseudotime analysis showing malignant clusters emerging as late branches from

the progenitor axis.

(E) Hierarchical clustering divides clusters into ciliated, secretory, and cancer cell

compartments.

(F–G) UMAP overlays illustrating that a subset of malignant cells co-cluster with

epithelial progenitors, indicating shared transcriptional programs.

(H–I) Pseudotime heatmaps depicting temporal gene expression patterns along

immunoreactive (H) and mesenchymal-like (I) malignant trajectories.

See also **Data File S7–S8**.

1003

1004

Figure 5. Transcription factor regulatory networks governing normal fallopian

tube differentiation and HGSOC malignant transformation.

(A) Integrative analysis workflow combining normal FTE and HGSOC nuclei for

SCENIC regulon inference.

(B) Heatmap of regulon activity (AUC scores) identifying stem, progenitor, secretory,

ciliated, and cancer-specific transcriptional programs.

(C–D) Ranking of regulons reveals SPDEF as a key stem/progenitor regulator and

NR2F6 as a highly specific cancer-associated regulator.

(E) Gene regulatory network map showing lineage-specific TF modules.

(F–G) NR2F6 knockdown reduces colony formation in ES2 and OVCAR3 cells.

(H–I) NR2F6 knockdown decreases invasive capacity of ES2 and OVCAR3 cells in

Transwell assays.

Data presented as mean ± SEM. Student's t test. $p < 0.05$; $p < 0.01$; $p < 0.001$.

1018

1019

Figure 6. Genome-wide DNA copy number variation (CNV) landscape in HGSOC.

(A) Overview of inferCNV workflow using FTE epithelial cells as the reference

population.

(B) Hidden Markov Model–based classification of copy number states (0x, 0.5x, 1x, 1.5x,

2x, ≥3x).

(C) Genome-wide CNV heatmap showing chromosomal losses and gains across

malignant cells.

(D) CNV-driven gene expression biases, highlighting amplified oncogenes (S100A6,

MUC16, COL1A1/2, CD74, WFDC2) and genes reduced through copy loss.

See also **Data File S9**.

1030

1031

Figure 7. Collagen signaling networks orchestrating stromal–tumor

communication in HGSOC.

(A–B) Ligand–receptor interaction networks showing CAFs as the dominant source of

outgoing signals and major regulators of the tumor microenvironment.

(C) Selected ligand–receptor interaction patterns across malignant and stromal cell

types.

(D) Collagen signaling network indicating extensive cross-talk between CAFs and

malignant cells.

(E) Violin plots of collagen-related gene expression (COL1A1, COL1A2, COL4A1/2,

COL6A1/2/3) across major tumor and non-tumor cell subtypes.

See also **Figure S5**.

1043

Figure 1

Z-score
normalized expression

−3   0   3

E

1
GO:0008584~male gonad development
NR5A1, STAR, AMHR2, KIT, GATA4, INHA
GO:0007179~transforming growth factor beta receptor signaling pathway
LEFTY1, AMHR2, BAMBI, WFIKKN2, LEFTY2
GO:0001541~ovarian follicle development
KIT, FOXL2, INHA

2
GO:0006955~immune response
IGHV2-5, IGHV1-2, IGHV3-48, IGLV3-1, IGHV4-39, IGKV3D-15, IGLV8-61, JCHAIN, IGLV2-11, IGLV2-8, IGLV3-25, IGLV1-40, IGKV1-16, IGKC,...
GO:0010628~positive regulation of gene expression
OSR2, IL33, SLC24A3, PID1, CITED2, TCF23, PDE2A, MIR27B, GATA5, FBLN1, GATA2, KLF4, ACTG2, ACTA2, CLDN5, SFRP4, ALDH1A2, HAND2, ID1, EMILIN1

3
GO:0035914~skeletal muscle cell differentiation
EGR1, NR4A1, EGR2, BTG2, FOS, ATF3
GO:0006357~regulation of transcription from RNA polymerase II promoter
NR4A2, EGR1, CSRNP1, NR4A1, EGR2, JUN, EGR3, FOSB, FOS, JUNB, ATF3

4
GO:0048704~embryonic skeletal system morphogenesis
HOXA3, HOXB6, HOXA5, HOXA4, HOXB5
GO:0048793~pronephros development
HNF1B, PAX2
GO:0007501~mesodermal cell fate specification
SFRP2, PAX2

5
GO:0007275~multicellular organism development
EHF, PAEP, HOXB2, EPHA1, SPDEF
GO:0006357~regulation of transcription from RNA polymerase II promoter
HES6, EHF, HOXD1, IRX3, RORC, HOXB4, FOXJ1, HOXB2, IRF6, POU5F1, SPDEF

6
GO:0003341~cilium movement
ODAD1, RSPH4A, ODAD3, DNAI2, CFAP73, SPEF1, SPAG6, ODAD4, CCDC103, DNAAF1, DNAH9, RSPH9, CFAP206, TTC29, CFAP100, ...
GO:0030317~flagellated sperm motility
DRC7, ODAD3, CFAP52, TCTE1, CFAP251, CATSPERD, TEKT1, TEKT2, ...

7
GO:0006508~proteolysis
ST14, MMP7, MMP1, TMPRSS3, TMPRSS4, HPN, MMP3, KLK5, PRSS22, KLK8, MMP9, KLK11, WFDC2, KLK7, MMP10, C2, PRSS21, MMP11, ..
GO:0051301~cell division
UBE2C, CDCA5, CDCA8, BUB1B, ZWINT, AURKB, AURKA, CDC20, CCNB2, ...
GO:0031424~keratinization
KRT81, KRT17, KRT8, KRT7, SFN, KRT5, KRT6A

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6