1

Unveiling The Origin of Human High-Grade Serous "Ovarian" Cancer Using Bulk and Single-Nucleus RNA Sequencing

Qian Li[1,2,#], Keren Cheng[3,#,*], Lili Sun[2], Wei Yan[4,5,6,*]

[1]Department of Medical Genetics, China Medical University, Shenyang, Liaoning, 8 110122, China.

[2]Department of Gynecological Oncology, Cancer Hospital of China Medical University/ 10 Liaoning Cancer Hospital & Institute, Shenyang, Liaoning, 110042, China.

[3]Department of Obstetrics and Gynecology, Center for Reproductive Medicine, the 12 Fourth Affiliated Hospital of School of Medicine, and International School of Medicine, 13 International Institutes of Medicine, Zhejiang University, Yiwu, China,322000

[4]The Lundquist Institute for Biomedical Innovation at Harbor-UCLA, Torrance, CA 90502, 15 USA.

[5]Department of Medicine, David Geffen School of Medicine at UCLA, Los Angeles, CA 17 90095, USA.

[6]Current Address: School of Molecular Biosciences, Washington State University, 19 Pullman, WA 99164, USA

[#]Co-first authors

*Correspondence: Keren.Cheng@zju.edu.cn or wei.yan@wsu.edu

Contact Lead: Dr. Wei Yan, wei.yan@wsu.edu

In Brief

This report describes comprehensive cell atlases of the human fallopian tube and high-

grade serous "ovarian" cancer (HGSOC), the relationships among major cell subtypes

within the human fallopian tube epithelium (FTE), and the most likely pathways by which

FTE progenitor cells are transformed into HGSOC cells. The data will serve as an

important resource and open new avenues for further investigation into diagnostic,

therapeutic, and preventive measures for FTE cell-derived HGSOC.

Highlights

1. $LGR5^+$/PGR^+ cells in human FTE represent stem cells, mostly quiescent and

capable of differentiating into $OVGP1^+$/$RNPC3^+$ progenitors. These progenitors,

in turn, differentiate into secretory and ciliated cells to maintain cellular

homeostasis in the FTE.

2. Progenitor cells, morphologically resembling secretory cells, are abundant in all

segments of the fallopian tube, and those in the fimbriae are the most likely

precursors of HGSOC.

3. Stem or progenitor cells can be transformed into cancerous cells following two

distinct trajectories, leading to the formation of immunoreactive and

mesenchymal cancer cell types common in HGSOC.

ABSTRACT

Secretory cells in the fallopian tube epithelium (FTE) have been suspected to be the

origin of high-grade serous "ovarian" cancer (HGSOC). However, due to the high

heterogeneity of the FTE cells, the true identity of the FTE cells that are transformed

into HGSOC cells and the spatiotemporal molecular events leading to carcinogenesis

remain unknown. Our integrative analyses, utilizing both bulk and single-nucleus

transcriptomics, revealed that the FTE cells are indeed highly heterogeneous,

consisting of multiple subtypes of secretory and ciliated cells in addition to a small stem

cell pool. In the adult fallopian tube, the $LGR5^+/PGR^+$ stem cells, often situated in the

basal layer of the FTE, can proliferate and differentiate into $OVGP1^+/RNPC3^+$

progenitors, constituting the predominant cell types in all segments of the fallopian tube.

These progenitors can quickly differentiate into ciliated or mature secretory cells to

replenish the lost FTE cells when needed. Exposure to certain oncogenic factors may

cause the progenitors to undergo carcinogenesis along two differentiation pathways,

forming mesenchymal-like and immunoreactive cancerous cells, commonly seen in

HGSOC. Key transcription factors and genetic changes in critical genes were also

identified during carcinogenesis. The present study provides comprehensive cell atlases

of the human fallopian tube and HGSOC and helps gain insight into the molecular

events leading to the transformation of FTE progenitors into HGSCO cells. (217 words)

KEYWORDS: *Ovarian cancer; fallopian tube; stem cell; progenitor; multiciliogenesis;*

tumorigenesis; differentiation; cancer microenvironment.

INTRODUCTION

High-grade serous carcinoma represents the most common type of ovarian cancer, accounting for ~75% of all epithelial ovarian cancer cases. It is also the most lethal histotype of ovarian cancer, with a 10-year survival rate of <30%. While histologically different from typical ovarian epithelial cells, cancerous cells in high-grade serous "ovarian" cancer (HGSOC) resemble epithelial cells of the Müllerian duct-derived internal reproductive tracts, leading to the hypothesis that the fallopian tube may be the source of HGSOC. Indeed, phenotypic, genetic, and proteomic analyses have revealed similarities between HGSOC and fallopian tube epithelial (FTE) cells, especially those from the infundibulum of the fallopian tube.[1,2] While secretory cells are considered the main source of HGSOC, signatures of ciliated cells have also been found in HGSOC, giving rise to the postulation that there exist FTE progenitors, which are capable of differentiating into ciliated cells to replenish the damaged ciliated cells or entering carcinogenesis, leading to the formation of serous intraepithelial carcinoma which eventually spread to the ovary.[3,4]

The fallopian tube can be divided into four distinct segments from the ovary to the uterus: the infundibulum, ampulla, isthmus, and uterotubal junction. The infundibulum's proximity to the ovary subjects this portion to inflammatory factors liberated from oocyte expulsion during the monthly ovulatory events,[5] leading to frequent tissue injury, repair, and remodeling within the epithelial layer.[6] The fallopian tube possesses regenerative capacities to replenish the injured epithelial cells to maintain proper cellular homeostasis.[7,8] The small round cells occasionally seen at the basal layer of FTE (also called intercalary cells) have been postulated as the FTE stem

cells, which can both self-renew and differentiate into ciliated cells or secretory cells

(also referred to as "Club" or Peg cells) to fulfill the replenishment function.[6,9] However,

the molecular signature of the FTE stem cell remains elusive.

Single-nucleus RNA sequencing (snRNA-seq) allows for in-depth transcriptomic

profiling using nuclei extracted from frozen samples, with sequencing deferred until

pathological validation.[10] It offers distinct advantages, including minimized dissociation

bias, compatibility with frozen specimens, elimination of dissociation-induced

transcriptional stress responses,[11] and a heightened efficacy in capturing intricate cell

subtypes.[12] Furthermore, snRNA-seq parallels single-cell RNA sequencing (scRNA-seq)

in terms of sensitivity and classification of cell subtypes.[13] In the present study, we

performed both bulk RNA-seq and snRNA-seq using human fallopian tubes and

HGSOC samples collected from patients. We identified the LGR5$^+$/PGR$^+$ cells as the

FTE stem cells that either self-renew or differentiate into the progenitor cells

(OVGP1$^+$/RNPC3$^+$), which can further differentiate into normal FTE cells (i.e., secretory

and ciliated cells) under physiological conditions, or get transformed into cancerous

cells leading to HGSOC.

RESULTS

Dual origins of HGSOC identified by bulk RNA-seq

It has been hypothesized that HGSOC originates either from the fallopian tube or

ovarian epithelium. To test the hypothesis, we first performed bulk RNA-seq analyses to

explore transcriptomic correlations among the HGSOC, fallopian tube and ovary (Figure

1A). With the patients' consent, twenty-six specimens, including four cancer-free ovaries,

four cancer-free fallopian tubes with each dissected into three segments

(fimbriae/infundibulum, ampulla, and isthmus), and ten HGSOC samples. The cancer-

free ovary and fallopian tube samples were collected from five patients undergoing

surgery due to hysteromyoma or endometrial polyps, whereas the cancerous tissues

were from patients diagnosed with epithelial ovarian cancer (EOC) and subsequently

confirmed to have HGSOC based on pathology (Table 1).

In our bulk RNA-seq analyses, each specimen yielded between 38 and 65 million

high-quality reads (Figure S1A). We first performed principal component analyses and

observed three distinct clusters, representing the three sample types (ovary, fallopian

tube and HGSOC) analyzed (Figure 1B). The fallopian tube segments showed a close

transcriptomic resemblance, and the biological replicates of the fallopian tube and

ovarian specimens were similar (Figures 1B and S1B). Of interest, while eight EOC

samples were clustered together, two (EOC-3 and -4) grouped with the ovaries,

suggesting their potential ovarian origin. Despite transcriptomic heterogeneity among

the eight HGSOC samples clustered together, these HGSOC samples appeared closer

to the fallopian tube than the ovarian samples, based on hierarchical correlation and

pair-wise Pearson coefficient assays (Figures 1C and S1B). We, therefore, categorized

EOC-3 and EOC-4 as ovary-derived HGSOC (EOC-O) and the remaining HGSOC

samples as fallopian tube-derived HGSOC (EOC-F) (Figure 1C).

By comparing the differentially expressed genes (DEGs) between EOC-F and

EOC-O, we identified that the fallopian tube-specific genes (e.g., *FOXJ1*, *SOX17*,

PODXL, *MUC16*, *WFDC2*) were upregulated in EOC-F, while the ovarian-specific genes

(*FOXL2*, *ARX*, *KIT*, *STAR*, *WFIKKN2*) were upregulated in EOC-O (Figure S1C). Gene

ontology (GO) enrichment analyses of these DEGs revealed pathways such as "cellular

division," "cell adhesion," and "spermatogenesis" in EOC-F, whereas "male gonad

development," "female gonad development," and the "WNT signaling pathway" were

predominant in EOC-O (Figure S1C). The fact that eight out of ten HGSOC samples

analyzed in this study are likely derived from the fallopian tubes supports the notion that

most of the HGSOC are derived from the fallopian tube.

DEGs identified among the EOC-F, FT, and ovary samples were grouped into

seven genesets based on GO term enrichment analyses (Figure 1D and Table S1).

Geneset 1, predominantly upregulated in the ovarian specimens, was enriched in male

gonad development, ovarian follicle development, and the transforming growth factor-

beta receptor signaling pathway. Geneset 2 showcased an enrichment in immune

response and positive regulation of gene expression in normal fallopian tube and ovary

samples. Notably, key transcription factors, including *HAND2, CITED2, OSR2, GATA2,*

and *ALDH1A2,* known to play pivotal roles in ovarian and fallopian tube development,

were among the upregulated genes. Geneset 3 captured the presence of smooth

muscle components within the tissues. Geneset 4 was marked by fallopian tube-specific

transcriptomic signatures, including those related to pronephros development,

mesodermal cell fate specification, and embryonic skeletal system morphogenesis, with

expression of key transcription factor *PAX2* and various *HOX* genes. Of interest was

geneset 5, which was enriched with multicellular organism development and the

regulation of transcription from the RNA polymerase II promoter. This set was

prominently upregulated in both fallopian tubes and EOC-F, with key transcription

factors like SPDEF, POU5F1, FOXJ1, HOXB2 and HOXB4 notably upregulated.

Geneset 6 is highlighted by cilium movement and flagellated sperm motility, showcasing

the expression of cilia and flagella-associated proteins, outer dynein arm docking

complexes and tektins. Geneset 7, reflecting cancer-specific metabolism, rapid cell

cycle and keratinization of epithelium, was enriched with proteolysis, cell division, and

keratinization pathways (Figures 1D &1E, Table S1).

By contrasting EOC-F, fallopian tube and ovary samples in a pairwise fashion

(Figures S1D, S1E &S1F), we discerned that genes related to cell division, chemokine-

mediated signaling pathways (CXCL9, CXCL10, CXCL11/CXCR3 axis, CXCL6/CXCR2

axis), inflammatory responses (IL1/IL1RAP axis, GAL-3 and NR1H4), apoptotic

processes (IL24, TNFAIP8), and antigen presentation via MHC class II (CTSS induced

by TNF-α/IFN-γ, CTSV9 induced by NF-κB) were augmented in EOC-F. These genes

are known to bolster inflammation and tissue fibrosis, reflecting the multifaceted

characteristics of EOC-F. In addition, immune-related genes HLA-DMA and HLA-DRA,

which induce chemotaxis and promote the differentiation of T cells by Notch signaling,

widely participate in immunocyte recruitment, impact tumor progression and modulate

the tumor microenvironment,[14,15] were conspicuous. Overall, these findings underscored

the pivotal roles of these genes in orchestrating inflammation, immune responses and

the tumor microenvironment involved in tumorigenesis and progression of HGSOC.

Taken together, our bulk RNA-seq data support the notion that HGSOCs have dual

origins, i.e., the ovary and fallopian tube, and most HGSOCs are derived from the

fallopian tube.

187 **Identification of a large pool of progenitors capable of differentiating into**

188 **secretory or ciliated cells in the fallopian tube epithelium**

189 While bulk RNA-seq offers a broad transcriptomic overview, it lacks the nuanced

190 resolution to discern the complexity inherent to individual cell subtypes and their

191 transcriptomic dynamics contributing to the origin of HGSOC. Even though several

192 previous studies have performed single-cell RNA-seq analyses of human fallopian

193 tubes,[7,16–19] the definition of FTE stem cells, the distinction between various secretory

194 cell subtypes, and the identity of mature secretory and ciliated cells remain controversial.

195 To solve these problems, we collected human cancer-free fallopian tubes from three

196 perimenopausal women undergoing hysterectomy with bilateral salpingo-oophorectomy

197 due to benign uterine diseases, sectioned them into infundibulum/fimbriae, ampulla, and

198 isthmus based on morphology, then performed single-nucleus RNA-seq (snRNA-seq)

199 analyses.

200   Employing the 10X Genomics platform for snRNA-seq, we analyzed 87,920 cells

201 (post-initial quality control) with a range of 36,000-63,000 mean reads per cell and a

202 median of ~2,000 genes per cell from 9 specimens, spanning three fallopian tube

203 segments: fimbriae/infundibulum, ampulla and isthmus (Figure S2A). After performing

204 cell cycle scoring and regressing out the effects of cell cycle, mitochondrial genes and

205 library size for subsequent analyses, our data did not reveal cell clustering predicated

206 on cell cycles, a routine quality control (QC) step in single-cell data analyses (Figure

207 S2B). Utilizing the uniform manifold approximation and projection (UMAP) for

208 embedding, the various fallopian tube segments displayed an even distribution,

precluding the necessity for clustering based on sample preparation (Figure S2C). The

QC assays demonstrated the high quality of our snRNA-seq datasets.

We first combined all fallopian tube samples and conducted initial cell clustering

and annotations, leading to the identification of five cell types, including secretory cells,

ciliated cells, smooth muscle cells, endothelial cells and macrophages (Figure S2D)

based on well-documented marker genes (Figure S2E). By analyzing the differentially

expressed genes and enriched GO terms (Figure S2F and Table S2), we discovered

that secretory cells likely contain the potential stem cells because genes known critical

for adult stem cell homeostasis in hair follicle development, metanephric epithelium

development, and regulation of transcription from RNA polymerase II promoter, were

among the most upregulated one (Figure S2F). Intriguingly, some genes are markers for

adult stem cells, for example, *SOX17, LGR5, TFCP2L1* and *HES1,* which are highly

expressed in the secretory cell cluster. Endothelial cell cluster exhibited hallmarks of

angiogenesis, vasculogenesis and cell adhesion – quintessential characteristics of this

cell lineage. The ciliated cell cluster demonstrated enrichment in cilium assembly,

epithelial cilium movement, and outer dynein arm assembly. In the context of the

fallopian tube, smooth muscle cells, vital to its function, were enriched in actin

cytoskeleton organization and smooth muscle contraction. Meanwhile, macrophages

displayed enhanced inflammatory responses, differentiation patterns, and positive

modulation of cytokine production (Figure S2F). Moreover, a trend was observed where

the number of secretory cells was the highest in the isthmus, decreasing towards the

ovary, whereas ciliated cells increased in number from the isthmus to the fimbriae

(Figure S2G).

We then collected all data points from 34,267 FTE cells to identify cell subtypes

and determine the transcriptomic dynamics in the FTE. Utilizing Louvain for re-

clustering, we segregated the FTE cells into 12 distinct clusters, as visualized using

UMAP (Figure 2A). Cells were subsequently partitioned with ward.D2, based on mean

expression across clusters, resulting in five discernible subgroups (Figure 2B). Based

on the top marker genes and insights from existing literature, we designated cell cluster

12 as FTE stem cells (*LGR5$^+$, PGR$^+$, PODXL$^+$, SOX17$^+$*). Clusters 2 and 6, marked by

elevated expression of *PAX8, ZBTB16* and *EGR1*, were annotated as mature secretory

cells, while clusters 1, 4, 5, and 8 were identified as ciliated cells (*CAPS$^+$, CFAPs$^+$*).

Significantly, we postulated that the remaining clusters (Clusters 3, 7, 9, 10 and 11)

represented progenitor cells characterized by the pronounced expression of *OVGP1,*

CRISP3, RNPC3 and *BDP1* and the absence of *LGR5, PGR, PODXL, PAX8, CAPS*

and *FOXJ1* expression (Figure 2C).

Leveraging RNA velocity analyses, we discerned that these progenitor cells were

not fully mature, while harboring some secretory traits, and retained the potential to

differentiate into ciliated cells (Figure 2D). Using PAGA to analyze developmental

trajectory and connectivity among cell clusters, we found that FTE stem cells (Cluster

12) were located at the bottom tip, and progenitor cells (Clusters 3, 7, 9, 10 and 11)

straddled between ciliated cells and secretory cells located at distinct trajectory termini,

consistent with our RNA velocity findings (Figure 2E). We then used PHATE to better

visualize cell developmental trajectories. Our findings revealed that cluster 12 remained

at a terminal tip, while the other cells branched off in two directions: one leading to

secretory cells and the other to ciliated cells (Figure 2F). Following the establishment of

these trajectories, we employed PHATE embedding to fit the principal curve and then

computed the pseudotime based on these trajectories (Figure 2G). Key marker gene

expressions were then projected to validate our trajectory analyses (Figure S2H). These

trajectory analyses revealed a dynamic continuum encompassing the fallopian tube

epithelial stem cell pool and progenitor cells with the potential to differentiate into either

mature secretory cells or ciliated cells.

Based on the cell developmental trajectories, we analyzed the transcriptomic

dynamics transitioning from stem cells to either secretory or ciliated cells (Figures 2H

and 2I). Within the trajectory delineating FTE stem cell differentiation into secretory cells,

we observed a declining expression of *LGR5, SPDEF, NR2F1* and *AR* in congruence

with pseudotime. Notably, genes responsive to estradiol (*POSTN, MAP1B, PTGS2, and*

EZH2) experienced a transient up-regulation, likely reflecting post-ovulation hormonal

modulations that propel stem cells or progenitors towards the secretory cell phenotype.

Mature secretory cells exhibited an amplified expression of *EGR1, PAX2, PAX8, CD74,*

and *ZBTB16* (Figure 2H and Table S3). Conversely, transitioning from FTE stem cells to

ciliated cells coincided with a decrease in *LGR5* and *LAMA5* expressions concurrent

with pseudotime. As cells initiate differentiation, certain genes responsive to xenobiotic

stimulus and cell adhesion were transiently up-regulated in the middle of the

differentiation trajectory. Ciliated cells, positioned at the end of the trajectory, were

characterized by elevated expressions of *CFAPs* and *Dynein*, consistent with active

cilial assembly and movement (Figure 2I and Table S4). By calculating the proportional

distribution of the annotated epithelial cell types, we found the FTE stem cells reside

predominantly in the isthmus, coexisting with progenitor cells and collectively

representing approximately 2/3 of the total epithelial cell population (Figure S2I & S2J).

In contrast, the stem cells were scarce in the fimbriae and ampulla, but progenitors

constituted roughly 1/3 (Figure S2I & S2J) of the total epithelial population in these two

segments.

Immunohistochemical staining of the ciliated cell marker TUBB4B and the

secretory cell marker PAX8 across the isthmus, ampulla and fimbria (Figure S3A)

revealed an incline in the density of TUBB4B$^+$ ciliated cells and a decline in the density

of PAX8$^+$ secretory cells from the isthmus to the fimbria (Figure S3B), aligning with our

snRNA-seq data (Figure S2G, S2I&S2J). Additionally, our immunofluorescent staining

confirmed that the basal cells, with the typical small and round morphology, showed

prominently nuclear expressions of LGR5 and PGR, consistent with their identity as the

FTE stem cells (Figures 2J, 2K and S3C). Consistent with our snRNA-seq data (Figure

S2J), our immunostaining also showed that the LGR5+/PGR+ stem cells were

predominantly localized within the isthmus, followed by the ampulla, with a conspicuous

absence in the fimbria (Figures 2J, 2K and S3C). Taken together, our snRNA-seq data

reveal that progenitor cells (*OVGP1*+) represent the most abundant cell type,

accounting for ~38% of the total FTE cells and that the progenitor cells are derived from

the FTE stem cells, which are those small round basal cells, with the potential to

develop into either ciliated (*CFAPs$^+$; DNAs$^+$*) or mature secretory (*PAX8*+) cells.

High inter-patient variability and heterogeneity in HGSOC cell composition

Given the intra-tumor heterogeneity and infiltration of non-cancer cells, we collected

15,322 cells from HGSOC snRNA-seq datasets (Figure S3D) for further analyses after

removing the batch effect and regressing out the effects of cell cycle and library size

(Figure S3E) using LIGER (Linked Inference of Genomic Experimental Relationships).[20]

As expected, marked heterogeneity was observed across patients (Figures S3F & S3G).

Within HGSOC samples, we delineated 13 cell clusters through Louvain

clustering, as visualized on UMAP (Figure 3A), including ten cancer cell clusters and

three non-cancer cell clusters (Figure 3B). The non-cancer cell clusters were annotated

as macrophage, cancer-associated fibroblasts (CAFs) and endothelial cells. Cancer cell

clusters were classified into proliferative, mesenchymal, ciliated-like, epithelial-to-

mesenchymal transition (EMT) and MHC (immunoreactive) subtypes (Figure 3C).

Analyses of the most upregulated genes in each cluster (Figure 3D and Table S5)

revealed that genes instrumental in cell division and cell cycle are enriched in

proliferative cancer cells, whereas the mesenchymal cell type exhibited overexpression

of genes pertinent to the negative regulation of cell migration alongside key transcription

factors like SOX17, MECOM and PAX8. While cilium-like cancer cells expressed genes

involved in cilium assembly and movement, EMT cancer cells predominantly expressed

genes associated with cell migration and adhesion, signifying their ongoing transition

from an epithelial to a mesenchymal phenotype. MHC (immunoreactive) cancer cells

were predominantly characterized by antigen processing and peptide or polysaccharide

antigen presentation via MHC class II (Figure 3D and Table S5). We also assessed

several HGSOC marker genes recently identified (Figure S3H).[21]

Furthermore, upon re-clustering the malignant cells (with minimal representation

from HGSOC_1) and visualizing their subtypes on UMAP (Figure 3E, 3F), a strong

patient-derived batch effect was discernible even post-LIGR-based batch effect

mitigation. Intriguingly, the mesenchymal subtype resembled the proliferative subtype

(Figure 3G), leading us to postulate that these cell subtypes might function analogously

to tumor stem cells. Cilium-like cancer cells, albeit sparse, were universally present

across all four HGSOC samples (Figure 3G). All five cancer cell subtypes express high

levels of PAX8 and MUC16 (Figure S4A), which are normally expressed in the FTE

(Figure 2C), supporting the origin of HGSOC from the FTE cells. Indeed, both PAX8 and

MUC16 are abundantly expressed in HGSOC cells (Figure S4B, S4C). MKI67,

exhibiting wide and high expression in the fallopian tube, underscores the active

proliferative state of FTE cells (Figure 2J). TUBB4B, while prevalent in HGSOC, is also

a hallmark of ciliated cells in the FTE (Figure 2C), attesting to the presence of ciliated

cell-like cancer cells (Figure S4A & S4B). MHC (immunoreactive) cancer cells

predominantly expressed Clusterin (CLU) and specific immune cell genes (Figures 3H,

S4C, S4D and Table S6). MUC16 and *ARGLU1* exhibited up-regulation in the EMT

subtype, suggesting cellular migratory characteristics (Figures 3H, S4A, S4D and Table

S6). MHC (immunoreactive) cancer cells also demonstrated CD74 expression, a trait

observed in normal FT, albeit at diminished levels (Figures 3H, S4A, S4D and Table S6).

The mesenchymal subtype was characterized by Collagens (COL1A1, COL1A2) and

Versican (VCAN) expressions, epitomizing extracellular matrix features and tissue

connectivity, likely constituting the tumor's core mass (Figures 3H, S4C, S4D and Table

S6). Taken together, our snRNA-seq data reveal that HGSOCs display a high degree of

inter-patient variability in tumor cell composition and prominent heterogeneity in the

subtypes of malignant cells.

The FTE progenitor cells as the origin of HGSOC

Both previous reports and our data support the notion that most of the HGSOC

originates from FTE cells.[7,16] However, it remains unclear which specific FTE cells

deviate from their normal developmental program and get transformed into cancerous

cells. By merging 11,028 HGSOC cells and 34,267 FTE cells, we attempted to scrutinize

the relationship between normal FTE and HGSOC cells (Figure 4). The large enough

number of cells analyzed on the same sequencing platform at single-cell resolution

allowed us to include potential rare cell types which may be crucial for connecting the

normal FTE cells and those undergoing carcinogenic transformation. Using Force-

directed embedding in ForceAtlas2,[22] we identified 15 discernible cell clusters (Figure

4A). Using Slingshot, we discovered four developmental lineages, including those from

epithelial stem cells (Cluster 11) to secretory cells (Cluster 1), ciliated cells (Cluster 3)

and cancerous cells (Clusters 12 and 13), respectively. We also confirmed these

trajectories using PAGA (Figure 4B). Supplementing RNA velocity to the Force-directed

graph embedding, we discovered that clusters 0/4/8/11 (fallopian epithelial stem cells

and progenitors) represented a potential starting state and developed into Cluster 1

(secretory cells), Cluster 3 (ciliated cells) and Cluster 12/13 (cancer cells) (Figure 4C).

Furthermore, pseudotime analyses revealed concordant results with those from RNA

velocity and PAGA trajectory assays (Figure 4D). In addition to normal FTE cell

developmental trajectories (Cluster 11 to Cluster 3 or 1), two cancer cell trajectories

(Cluster 11 to Cluster 12/13) were also identified.

Hierarchical clustering analyses categorized the 15 cell clusters into three

primary groups: ciliated, secretory, and cancer cells (Figure 4E). In addition to the

epithelial stem cells (Cluster 11), the secretory subgroup contained progenitor cells

(Clusters 4, 0 and 8) and mature secretory cells (Cluster 1), collectively classifiable as

general secretory cells. Although classified as one of the five ciliated cell subtypes,

Cluster 14 appeared to exhibit characteristics of HGSOC cells based on their position in

the developmental trajectory (Figures 4B, C, and D), appearance in the cancerous

tissue (Figure 4F) and gene expression signature (Figure 4G). Similarly, progenitors in

Cluster 4, present in all segments of the fallopian tube, also displayed cancerous traits

(Figures 4F and G), suggesting that the FTE progenitors are a potential source of

HGSOC. Of significance, identification of the ciliated cell-like cancer cells (Cluster 14) is

consistent with previous reports showing that HGSOC tumor cells retain some of the

ciliated cell attributes characteristic of their progenitor lineage.[7,16]

Among cancer cells, four major subtypes were discerned, including

mesenchymal-like subtype (Clusters 10 with upregulated *VEGFA and NEAT1*), EMT-like

subtype (Cluster 13 with upregulated *SPARC* and *COL1A*), proliferative subtype

(Cluster 9 with upregulated *TOP2A, ASPM* and *MKI67*) and immunoreactive subtype

(Cluster 12, inflammatory-like cells, labeled as MHC with upregulated expression of

CD74 and *HLA-DRA*). It is of note that while the majority of Cluster 9 consisted of

cancer cells, it also incorporated cells from the fimbriae, ampulla and isthmus, indicating

that the progenitors within each segment of the fallopian tube can all be transformed

into cancerous cells (Figure 4F).

Although increasing lines of evidence support the notion that HGSOC is derived

from FTE cells, it remains unknown how the FTE cells become cancerous cells at

molecular levels. By exploring the expression patterns of the top 5,000 variable genes

along the two cancer cell trajectories mapped to pseudotime, we attempted to gain

insight into the molecular events during FTE cell transformation into HGSOC cells

(Figure 4H and I). From epithelial stem cells (Cluster 11) to immunoreactive cancer

cells (Cluster 12), genes associated with the Notch signaling pathway and cell fate

determination were first downregulated at the beginning of pseudotime, followed by

upregulation of genes mediating anti-apoptotic processes, cilium assembly, and peptide

antigen assembly via the MHC II protein complex during the intermediate or

transformative phase of pseudotime. In the terminal stages of pseudotime, genes

associated with EMT, cellular response to tumor necrosis, and collagen fibril

organization exhibited elevated expression (Figure 4H and Table S7). During the

trajectory from epithelial stem cells (Cluster 11) to mesenchymal-like cancer cells

(Cluster 13), the early pseudotime phase marked a decline in genes modulating stem

cell division. Toward the end of this trajectory, genes promoting angiogenesis, integrin-

mediated signaling, and cellular responses to fibroblast growth became dominant,

echoing the canonical transcriptomic signatures characteristic of tumor proliferation,

invasion and angiogenesis (Figure 4I and Table S8).

Broadly construed, stem cells and progenitors tend to differentiate into ciliated

cells in physiological environments (Figure 2). During this normal differentiation, certain

cilium-related genes either transiently increase or demonstrate a brief upregulation

followed by a decline (Figure 2I and Table S4). In contrast, under carcinogenic

conditions, genes mediating cellular responses to tumor necrosis and integrin signaling

become upregulated (Figures 4H, 4I). Such aberrant cellular environments also activate

the cell cycle and metabolic shifts, favoring aerobic respiration (Figure 4I). This

metabolic reprogramming, synonymous with the Warburg effect, potentially facilitates an

environment conducive to tumor growth by optimizing nutrient acquisition, proliferation,

metastasis and drug resistance mechanisms.[23,24] Overall, our findings pinpoint FTE

progenitors located at fimbriae as the primary source of cancer cells. Under the aegis of

inflammatory stimuli or cellular injury, these progenitors undergo transformation and

become cancer stem cell-like entities *via* two distinct trajectories, eventually culminating

as either immunoreactive or mesenchymal-like HGSOC cells.

Key transcriptional factors driving the differentiation from FTE progenitors into

normal epithelial or HGSOC cells

We used pure cancer and FTE cells to conduct single-cell regulatory network inference

and clustering (SCENIC) analyses (Figure 5A). Notably, the highest-ranking

transcription factors (TFs) in ciliated cells were identified as NR2F1, FOXJ1 and HOXC4.

In contrast, TPI1, NR2F6, and MAFB manifested distinct specificity in cancer cells.

Secretory cells, progenitors, and stem cells exhibited a commonality of TFs (Figure 5B),

reinforcing the dynamic cell fate interplay among these subsets, corroborating our

preceding findings. We delineated several TFs potentially influencing cell fate during

normal and anomalous fallopian epithelium maturation, subsequently categorizing

regulon specificity based on AUC scores (Figure 5C). NR2F6, a newly identified

superfamily member of TFs in multiple types of cancer,[25,26] can potentially modulate

diverse biological events in response to hormones or nutrients. NR2F6 might represent

a novel therapeutic target for HGSOC intervention. Distinctively, SPDEF manifested

specificity in progenitors and stem cells, further emphasizing its role as a pivotal

regulator across diverse cancers (Figure 5C).[27,28] Historically, SPDEF was recognized

as an integral component of the Notch signaling cascade.[29] Given the shared motif

sequence among the SOX family, our identification of SOX5, rather than SOX17, was

unsurprising. We further corroborated these findings by analyzing overlap patterns

through the AUC score (bottom row) and gene expression levels (top row) of regulons

(Figure 5D). By this criterion, NR2F6 emerged as a central regulator in HGSOC,

whereeases SPDEF featured prominently in progenitors and stem cells. Through the

prediction of gene regulatory networks within each cell subtype, we identified SOX5 and

SPDEF as master regulators in stem cells and progenitors, PAX2, TBX2 and TCF7L2 in

mature secretory cells, HOXC4 and ZNF214 in ciliated cells, NR2F6 and TPI1 in

HGSOC cells (Figure 5E).

DNA Copy Number Variations in HGSOC

To elucidate potential genomic changes in HGSOC, we analyzed copy number

variations (CNVs) using the inferCNVs tool,[30,31] leveraging normal FTEs as a reference

benchmark (Figure 6A). We documented the presence of both gain and loss of

expression spanning extensive genomic scales. Utilizing the Hidden Markov Model

(HMM), we identified 138 CNV regions. These regions were further categorized via a

six-state CNV model delineating the following CNV levels: State1 denoting a complete

loss (0), State2 reflecting a loss of a single copy (0.5x), State3 being neutral (1x), State4

indicating the addition of one copy (1.5x), State5 highlighting the addition of two copies

(2x), and State6, which essentially serves as a placeholder for > 2x copies but is

modeled as 3x (3x) (Figure 6B and Table S9).

Aggregating the distributions across these CNV regions revealed that neutral

variants constituted ~39%, gains of copies represented ~34%, and loss of copies

comprised ~27%. Predominantly, these events were localized to Chromosome 1, with

the most pronounced alterations observed on Chromosome 6 (Figure 6C). Our analyses

further authenticated that the aberrant up-regulation of *S100A6, MUC16, COL1A1,*

COL1A2, CD74 and *WFDC2* in HGSOC could be attributed to the amplification of copy

numbers. Conversely, the diminished expression of *CLU, PODXL* and *PGR* likely results

from the deletion of copy numbers (Figure 6D). These insights could provide a rationale

for the preferential differentiation of progenitors into HGSOC cells rather than terminally

differentiated cells.

Pivotal Role of Collagen Signaling in the Tumor Microenvironment of HGSOC.

Within the tumor microenvironment (TME) of HGSOC, cancer-associated fibroblasts

(CAFs) emerged as dominant regulators of other cell subtypes (Figures S5A, S5B). A

comprehensive assessment illuminated the collagen signaling pathway as the

paramount extracellular matrix (ECM) regulator, influencing tissue architecture, cellular

adhesion, EMT, and tumorigenesis (Figures S5C, S5D).[32,33]   Further, our analyses

underscored the intricate interactions between tumor and non-tumor cell subtypes of

HGSOC, mediated by key collagen signaling pathway ligands and receptors. Notably,

ITGAV and ITGB8, encoding integrin receptors for fibronectin, were co-expressed in

both HGSOC tumor cells and CAFs. This mutual expression pattern suggests a

synergistic TME response orchestrated by these cell types. Moreover, genes encoding

specific collagen types (*COL1A1, COL1A2, COL4A1, COL4A2, COL6A1, COL6A2* and

COL6A3) displayed upregulated expression in mesenchymal-like and cilium-like cancer

cells and CAFs, highlighting their pivotal roles in modulating ECM remodeling within the

TME (Figure S5E).

DISCUSSION

It is becoming increasingly clear that HGSOC has a dual origin: ovarian surface

epithelium and FTE, with the latter being the majority, and our bulk RNA-seq data

support this notion. Given numerous DEGs identified between the two types of HGSOC

and their known physiological functions in the normal fallopian tube and ovary, it is

feasible to develop diagnostic biomarkers to distinguish the two origins. Based on

marker genes identified in our snRNA-seq analyses, it is also possible to further classify

the fallopian tube-derived HGSOCs into various cancer subtypes, which may facilitate

developing early detection, precision treatment, and tumor prevention strategies.

While increasingly accepted, the notion that HGSOC originates from FTE

secretory cells remains confusing because it implies that some of the FTE secretory

cells would undergo carcinogenesis and become cancerous cells. The secretory cells

are terminally differentiated cells, which should not develop into tumor cells easily.

Histologically, the secretory cells are heterogeneous, and many so-called secretory cell

markers, e.g., PAX8 and OVGP1, have been found to be expressed in a subset of

secretory cells but absent in other subsets of secretory cells. Recent single-cell

transciptomes of fallopian tube have clearly demonstrated that the FTE secretory cells

are indeed heterogeneous, and several subtypes of the secretory cells have been

defined.[18] However, it remains unclear which subtype of secretory cells tends to

undergo carcinogenesis, why this specific subtype is prone to carcinogenesis, and what

roles these cells play in normal FTE. To address these questions, we employed both

bulk RNA-seq and snRNA-seq to dissect the intricate cellular homeostasis in normal

FTE and to trace the specific FTE cell subtypes that undergo transformation and

become cancerous cells. The large number of single cells used in this study for

transcriptomic analyses allowed us to obtain a comprehensive human fallopian tube cell

atlas with higher resolution, compared to all previous studies.[19,34,35] The latest cell

developmental trajectory analyses adopted in this study made it possible to dissect the

intricate relationship among FTE stem cells and various subtypes of secretory and

ciliated cells.

We've discovered that the LGR5$^+$/PGR$^+$ basal small round cells are the FTE stem

cells, characterized by their relative quiescence and attenuated MKI67 expression.

Similar to mature secretory cells, the number of FTE stem cells gradually decreases

from the isthmus to the ampulla and infundibulum/fimbriae. The FTE stem cell

population is capable of differentiation into secretory and ciliated cells *via* intermediate

progenitors, substantiating a dynamic interchange amongst stem cells, progenitors and

secretory cells, thereby preserving fallopian epithelial homeostasis.

The stem cells (LGR5$^+$/PGR$^+$) are scarce in the FTE but can produce a large

pool of progenitor cells (OVGP1$^+$). Although histologically resembling secretory cells

due to a lack of cilia, these FTE progenitors are in a poised state and can differentiate

into either ciliated (CFAPs$^+$) or mature secretory (PAX8$^+$) cells when needed. It is

noteworthy that the progenitors represent the most abundant cell type in the FTE,

accounting for ~38% of the total FTE cells. It is well-established that stem cell-derived

progenitors with the potential to differentiate into one or multiple functional cell types

tend to undergo carcinogenesis under certain conditions.[36,37] It is noteworthy that

despite the very low number of stem cells, the progenitors are abundant in fimbriae,

probably reflecting more active replenishing due to the high turnover rate of FTE cells

due to damages caused by monthly ovulatory events.

Drawing from comprehensive gene expression and active chromatin datasets,

Lawrenson K. et al. show evidence that a preponderance of HGSOCs, particularly those

of the mesenchymal type, are descendants of fallopian tube secretory epithelial cells

(FTSECs).[38] Experiments by Nakamura K. et al., encompassing the inactivation of

dominant negative p53 and the upregulation of KRAS, c-myc, and PI3K-Akt,

emphatically validate that selective gene modulations can shepherd immortalized

FTSECs, suggesting that two patterns of three gene alterations (p53/KRASV12/c-Myc

and p53/KRASV12/PI3K-AKT) can develop immortalized FTSECs into early stage

HGSOC *in vitro.*[39] Here, we report that fallopian tube-derived HGSOC originates from

OVGP1[+]/RNPC3[+] (Cluster 4) progenitors in the epithelium of infundibulum/fimbriae,

which possess dual differentiation prowess towards both tumor and ciliated cells.

Historically, HGSOCs have been parsed into four main tumor subtypes:

differentiated, proliferative, mesenchymal and immunoreactive.[40] We, however, based

on the discrete functional attributes, classify HGSOC into five cellular subgroups,

including proliferative, mesenchymal-like, ciliated-like, EMT and MHC subgroups. Data

from the Australian Ovarian Cancer Study (AOCS) illustrate the ciliated subtype's

preponderance within low-grade tumors (grade 1),[41] while the EMT-High (mesenchymal)

subtype bears significantly poor prognostic implications, as gleaned from eight

independent datasets.[42–44]

The two distinct carcinogenic trajectories identified in the present study revealed

the intricate interplay of immune infiltrates and fibroblasts within the TME. The

prevalence of immuno-reactive cells potentially correlates with heightened tumor-

infiltrating lymphocyte (TIL) abundance, portending favorable prognosis and augmented

immunotherapeutic responsiveness.[45] Notably, the "immunoreactive" and

"mesenchymal" reflected the expression programs of macrophage and CAFs,

respectively, rendering them more as indicators of tumor composition than distinct tumor

cell phenotypes.[40]

Cancer-associated fibroblasts (CAFs) and immune cells constitute integral facets

of the TME. Upon activation, CAFs can enhance tumor proliferation, angiogenesis,

invasion, metastasis, extracellular matrix (ECM) reconfiguration, and even confer

chemoresistance through various molecular pathways. Many previous studies have

underscored the pivotal interplay between CAFs and neoplastic cells in tumorigenesis

and progression.[46,47] Our investigations confirmed that the trajectory of tumorigenesis is

characterized by salient transcriptomic hallmarks, encapsulating tumor proliferation,

invasion and angiogenesis. In response to hypoxic stimuli, CAFs secrete an array of

angiogenic mediators, notably VEGF, leading to the downregulation of endothelial cell

adhesion molecules, such as ICAM-1/2 and VCAM-1.[48] Tumor-associated macrophages

(TAMs), bifurcated into the M1 and M2 phenotypes, display divergent functionalities. M1

macrophages execute their antineoplastic activity in the tumor immune

microenvironment (TIME) predominantly through mediating antibody-dependent

cytotoxicity coupled with the generation of ROS and tumor necrosis factor (TNF).[49]

Conversely, M2 macrophages foster tumorigenesis by augmenting angiogenesis,

immunosuppression, tumor cell invasion, metastasis, and ECM dynamics.[50,51] The

aberrant neoplastic collagen matrix beckons TAMs and reciprocally enables TAMs to

modulate collagen deposition, culminating in escalated matrix rigidity and hastened

ECM metamorphosis.[52] Our data spotlighted the collagen signaling pathway as the

paramount cellular communication conduit, potentially elucidating the modus operandi

by which CAFs orchestrate the immunosuppression of TME, thus facilitating the

translation of CAF-centric therapeutic interventions into clinical trials.

Compared to other populations, individuals harboring BRCA mutations exhibit a

markedly elevated oncogenic predisposition. Approximately 20% of HGSOCs are

tethered to germline or somatic BRCA mutations.[53] A prospective study articulated that

by age 80, BRCA1 and BRCA2 mutation carriers bear 44% and 17% cumulative ovarian

cancer risks, respectively.[54] In fallopian tube specimens from BRCA mutation-positive

individuals undergoing Risk-Reducing Salpingo-Oophorectomy (RRSO), isolated serous

tubal intraepithelial carcinoma (STIC) lesions were discerned in 2.3% of cases, whereas

p53 signatures were present in a quarter of the cohort.[55] Clinically, STIC lesions

coexisted with HGSC in 11% to 61% of cases, with a prevalence surpassing 60%, even

in serous ovarian cancers devoid of BRCA linkage.[56] This ubiquity has ensconced STIC

as a precursor lesion to HGSC. While previous studies predominantly leveraged

fallopian tube samples from HGSOC-afflicted individuals, our approach pivoted towards

samples from oncologically unscathed women. Additionally, our findings elucidated that

the genesis of HGSOC from the fallopian tubes was a common phenomenon. Notably,

this occurrence exhibited no correlation with variables such as family history of tumors

(designated as a high-risk population), menstrual history, marital and reproductive

records, or a history of hormone therapy (a recognized high-risk factor).

In conclusion, our data provide evidence that fallopian tube-derived HGSOCs

originate from OVGP$^+$/RNPC3$^+$ (Cluster 4) progenitor cells. Intriguingly, this is the

inaugural identification of LGR5$^+$/PGR$^+$ epithelial stem cells, poised in a relative

quiescence within the fallopian tube epithelium. HGSOCs encompass a ciliated cell-like

subtype, which reciprocally interacts with MHC-like and EMT subtypes. Normal

development programs ensuring proper differentiation from FTE stem cells to

progenitors and then to mature secretory cells or ciliated cells, can be derailed by

carcinogenic factors, e.g., chronic inflammation and cell damage due to exposure to

harmful chemicals released during ovulation, etc., causing malignant transformation into

MHC-like and mesenchymal-like cancer cell subtypes, which, in turn, interact with the

TME, catalyzing tumor initiation and progression.

STAR ★ Methods

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit Anti-OVGP1 antibody	Bioss	Cat#bs-17571R
Rabbit Anti-Ki67 antibody	Bioss	Cat#bsm-60738R
Rabbit Anti-LGR5 antibody	Bioss	Cat#bs-20747R
Rabbit Anti-PGR antibody	Bioss	Cat#bs-23376R
Acetylated-Tubulin Rabbit pAb	Abcam	Cat#ab179484
Anti-PAX8 Rabbit pAb	Abcam	Cat#ab53490
Rabbit Anti-CA125 antibody	Bioss	Cat#bs-0091R
Biological Samples		
Surgical specimen from patients with ovarian cancer	The Cancer Hospital of China Medical University	N/A
Surgical specimen from benign patients	The Cancer Hospital of China Medical University	N/A
Critical Commercial Assays		
Trizol reagent kit	Invitrogen, Carlsbad, CA,USA	15596026CN
iQ™ SYBR® Green Supermix kit	Bio-Rad	1708886
UltraSensitive™ SP (Mouse/Rabbit) IHC Kit	MXB, Fujian, CA	KIT-9720
Deposited Data		
Bulk RNA seq	This paper	GSE223426
snRNA seq	This paper	GSE270250
Software and Algorithms		
Cell Ranger	10X Genomics; https://support. 10xgenomics.com/	v7.1, RRID:SCR_017344

	single-cell-gene-expression/software/ overview/welcome	
R package Seurat	https://cran.r-project.org/web/packages/Seurat/index.html	v.4.3.0,RRID:SCR_016341
R package MAST	https://bioconductor.org/packages/release/bioc/html/MAST.html	v1.30.0
R package SCENIC	https://github.com/aertslab/SCENIC	V1.1.2, RRID:SCR_017247
R package edgeR	https://bioconductor.org/packages/release/bioc/html/edgeR.html	v3.34.1, RRID:SCR_012802
RcisTarget	https://www.bioconductor.org/packages/release/bioc/html/RcisTarget.html	v1.24.0, RRID:SCR_024860
Scanpy	https://scanpy.readthedocs.io/en/stable/	v1.8.1, RRID:SCR_018139
PAGA	https://github.com/theislab/paga	
velocyto	http://velocyto.org/velocyto.py/tutorial/index.html	V0.17, RRID:SCR_018167
scVelo	https://scvelo.readthedocs.io/en/stable/	v.0.2.2, RRID:SCR_018168
PHATE	https://phate.readthedocs.io/en/latest/	V0.3.0
Cytoscape	https://cytoscape.org/	v3.9.1, RRID:SCR_003032
infercnv	https://github.com/broadinstitute/infercnv	V1.3.3, RRID:SCR_021140
DAVID	https://david.ncifcrf.gov/	v2023q4, RRID:SCR_001881
In-house analysis	https://github.com/chengkeren/fallopiantube	NA

Data and code availability

The accession numbers generated in this study are GSE223426 for bulk RNA-seq and GSE270250 for snRNA-seq analysis. The codes are available at the GitHub repository: https://github.com/chengkeren/fallopiantube). Ethics approval was obtained in all participating countries and, where required, in all participating centers. All patients provided written informed consent.

LEAD CONTACT AND MATERIALS AVAILABILITY

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Wei Yan (weiyanucla@ucla.edu). This study did not generate new unique reagents.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Human Fallopian Tube and Ovarian Cancer Samples

This study was approved by the Institutional Review Board (IRB) at Liaoning Cancer Hospital and the Cancer Hospital of China Medical University (Shenyang, P.R. China) (Protocol#: 20220245). All participating patients provided informed consent. For bulk RNA-seq analyses, HGSOC samples were collected from 10 patients (#1-10) who were diagnosed with EOC and underwent surgery. Benign ovaries and fallopian tubes came from patients (#11-15) who underwent total hysterectomy combined with bilateral salpingo-oophorectomy for benign uterine diseases, e.g., endometrial polyps, hysteromyomas, etc. (Table 1). For snRNA-seq analyses, the 13 samples were

collected from 9 patients, including 5 (#1-5) with hysteromyoma and adenomyosis, and

4 (#6-9) with EOC/HGSOC (Table 2).

METHOD DETAILS

Tissue processing

After removal from the patients, the cancer tissue was immediately rinsed in cold PBS

and dissected into several portions: one for fixation in 10% formalin overnight, followed

by preparation of paraffin blocks for immunohistochemical analyses, and the rest were

snap-frozen in LN2, followed by storage at -80°C until RNA isolation for bulk RNA-seq

or preparation of single nuclei for snRNA-seq. After removal, the human fallopian tubes

were immediately rinsed in cold PBS, followed by dissection under a dissection scope

into three portions: isthmus (proximal), ampulla (enlarged part), and fimbria (distal). The

dissected fallopian tube segments were snap-frozen in LN_2, followed by storage at -

80°C until RNA isolation for bulk RNA-seq or preparation of single nuclei for snRNA-seq.

Bulk RNA-seq library preparation and analysis

Total RNA was extracted using Trizol (Invitrogen, Carlsbad, CA, USA). RNA quality and

quantity were assessed using an Agilent 2100 Bioanalyzer (Agilent Technologies, Palo

Alto, CA, USA). mRNA was enriched by oligo(dT) beads, and rRNA was removed by

Ribo-ZeroTM Magnetic Kit (Epicentre, Madison, WI, USA). The enriched mRNA was

fragmented into short fragments, followed by reverse transcription into cDNA using

random primers. Second-strand cDNA was synthesized by DNA polymerase I, and the

cDNA fragments were purified using a QiaQuick PCR extraction kit (Qiagen, Venlo, The

Netherlands), followed by end repair, poly(A) addition, and ligation to Illumina

sequencing adapters. The ligation products were then size selected, followed by PCR

amplification. The libraries were sequenced on Illumina NovaSeq6000 (PE150) at

Genedenovo Biotechnology Co., Ltd (Guangzhou, China).

Raw fastq files were demultiplexed with bcl2fastq2 (v.2.20.0.422). Barcodes and

adapters were removed with Trimmomatic (v.0.32).[57] Fastq files were mapped to the

UCSC human reference genome (GRH38) using STAR (2.7.10a).[58] The raw gene count

table was generated with featurecounts,[59] and weakly expressed genes were filtered

with edgeR with the filterByExpr function with default parameters. Briefly, the raw counts

were normalized to library size, and then genes with counts per million above 10 were

included in downstream analysis. DEGs were analyzed with edgeR (v.3.36.0) with

log2fold change > 1, p-value < 0.05, and FDR < 0.05. Reads per kilobase per million

(RPKM) values were calculated in edgeR,[60] and the gene length was obtained from the

UCSC table browser.[61] Downstream data analyses and visualization were performed

with R (v.4.1.0). Hierarchical clustering was performed with hclust in R (v.4.1.0).

snRNA-seq analysis

Preparation of single nuclei from frozen human HGSOC and fallopian tube samples and

constructions of 3' single-cell gene expression libraries were conducted using the 10x

Genomics Chromium system (Next GEM v3.1) following the manufacturer's protocol.

The libraries were sequenced with ~200 million PE150 reads per sample on Illumina

NovaSeq at LC-BIO Co., Ltd (Hangzhou, China).

Raw data were demultiplexed with the mkfastq command in Cell Ranger(v7.1.0)

to generate Fastq files. Trimmed sequence files were mapped to the reference genome

for humans (GRCh38) provided by 10X Genomics using Cell Ranger mkfastq. Read

counts were obtained by using cellranger count[62]. Secondary data analyses were

performed in Python with Scanpy(v1.8.1)[63] or in R(v.4.1.0) with the Seurat(v.4.3.0),[64]

SeuratObject(v4.1.3), SeuratWrappers(v0.3.2),[64] tidyverse(v.1.3.2),[65] destiny(v3.10.0),[66]

pheatmap (v1.0.12), plot3D (v1.4) and Matrix (v.1.5-1) packages and Excel (Microsoft).

UMI count tables were first loaded into R by using the Read10x function in Seurat, and

Seurat objects were built from each sample. Cells with less than 200 genes, an aberrant

high gene count above 7000, or total mitochondrial genes above 15% were filtered out.

Samples were combined using Seurat merge function. The effects of mitochondrial

genes, cell cycle genes, were regressed out by SCTransform during normalization in

Seurat. Batch effect was removed using Harmony. Mitochondrial genes and cell cycles

genes were excluded during Cell Clustering, dimensional reductions and trajectory

analysis. Cells were clustered by a shared nearest neighbor (SNN) modularity

optimization-based clustering algorithm in Seurat. Clusters were annotated on the basis

of previously characterized marker gene expression by using the FeaturePlot function

and the gene expression matrix file. Then, annotated cell clusters were used for

downstream analyses. Dimensional reductions were performed with top 3000 highly

variable genes and first 20 principal components by using Seurat. Differentially

expressed genes (DEGs) in different clusters were calculated by MAST or area under

curve (AUC) integrated in Seurat findallmarkers funtion, with a threshold of avgLog2FC >

0.25, p-value < 0.01. For LogNormalize Expression (LogExp) in Seurat, genes for each

cell were divided by the total counts for that cell and multiplied by the cal factor (10,000),

then natural-log transformed using log1p.

Developmental trajectories of cells were simulated with first 30 PCs and 30

diffusion components by DiffusionMap in destiny and scanpy.[63] Trajectory principal lines

were fitted by DiffusionMap.[66] Pseudotime was calculated by dpt in destiny. Single-cell

pseudotime trajectories were constructed using PAGA,[67] and PHATE.[68]

For Pseudobulk differential expression analysis of scRNA-seq data, DEGs

between two groups in scatterplot were identified by edgeR 3.34.1 with applying quasi-

likelihood approach (QLF) and the fraction of detected genes per cell as covariate[60].

The DEGs were defined as the genes with FDR < 0.05, p-value < 0.05, and Log2-fold

change above 1. Expression value was normalized by edgeR using log2(CPM+1) (CPM

= Counts Per Million). Cell Cycle was analyzed using CellCycleScoring in Seurat. Data

was visualized by ggplot2 and pheatmap. Genes in heatmap were hierarchically

clustered by measuring euclidean distance, scaled by row, then visualized using

pheatmap. Gene ontology enrichment was analyzed by DAVID v6.8.[69]

RNA velocity

The velocyto command line interface was used to generate spliced/unspliced

expression matrices from the 10x Genomics Chromium samples.[70] RNA velocity

analysis was then performed by using scVelo v.0.2.2.[71]

Regulatory network inference using SCENIC

We used single-Cell regulatory network inference and clustering method (SCENIC) to

analyze transcription factors (TFs) regulatory networks, from co-expression of

transcription factor and candidate target genes from RcisTarget database.[72] After

defining the key TFs, the network was drew by using Cytoscape (v3.9.1).[73]

Single-cell copy number analysis

Initial CNVs (CNV) were estimated by sorting the analyzed genes by their chromosomal

location and applying a moving average to the relative expression values, with a sliding

window of 100 genes within each chromosome, as described.[74] To avoid considerable

impact of any particular gene on the moving average, we limited the relative expression

values to [-3,3] by replacing all values above 3 by a ceiling of 3, and replacing values

below -3 by a floor of -3. This was performed only in the context of CNV estimation. This

initial analysis is based on the average expression of genes in each cell relative to the

other cells and therefore does not have a proper reference to define the baseline. We

thus defined the gene expression clusters annotated as ciliated epithelial cells,

progenitor, secretory cells, FTE stem cells by gene expression as the nonmalignant

cells, and used the average CNV estimate at each gene across those cells as the

baseline. The final CNV estimate of cell i at position j was defined as: the 6 states

correspond to the following CNV events: State 1; 0x, complete loss. State 2; 0.5x, loss

of one copy. State 3; 1x, neutral. State 4; 1.5x, addition of one copy. State 5; 2x,

addition of two copies. State 6; 3x, essentially a placeholder for >2x copies but modeled

as 3x.

Immunohistochemistry and Immunofluorescence staining

Immunohistochemical staining was conducted as described.[75] Tissue samples were fixed in 10% buffered formalin (Fisher Healthcare) with gentle rocking overnight at room temperature. After dehydration, tissues were embedded in paraffin, serially sectioned at 5 μm thickness with a microtome (Leica) and placed on Superfrost Microscope glass slides. Paraffin sections were then de-paraffinized with xylene. Antigens were retrieved with 10mM Sodium Citrate for 35 min at 90° C and then for 15 min at room temperature. The staining and incubation procedure for paraffin sections was with the following modifications: the blocking solution was 5% normal donkey serum in PBST; the primary antibody incubation was performed overnight at 4° C; and slides were washed with PBS six times after each incubation. Slides were mounted in Vectashield mounting medium for imaging. The dilution rates of primary antibodies used in this study are as follows: anti-LGR5 (1:3000), anti-PGR (1:400), anti-Ki67 (1:2000), anti-OVGP1 (1:200), anti-CRISP3 (1:200).

DECLARATION OF INTERESTS

The authors declare no competing interests.

AUTHOR CONTRIBUTIONS

Visualization, methodology and statistics, K.C., W.Y.; writing-original draft, Q.L., K.C.; writing-revise, review and editing, W.Y., K.C.; conceptualization, design and supervision, W.Y.; acquisition of specimens, Q.L., C.Y.W.; acquisition of clinical data, Q.L.; histological and pathological guidance, L.L.S.

778 **ACKNOWLEDGMENTS**

779 We thank LC-BIO Co. Ltd. (Hangzhou, China) for providing the Chromium 10X

780 genomics single-nucleus transcriptome sequencing services and Genedenovo

781 Biotechnology Co. Ltd. (Guangzhou, China) for providing the Illumina NovaSeq 6000

782 sequencing services for bulk RNA-seq.

REFERENCES

1.   Gong, T.-T., Guo, S., Liu, F.-H., Huo, Y.-L., Zhang, M., Yan, S., Zhou, H.-X., Pan, X., Wang, X.-Y., Xu, H.-L., et al. (2023). Proteomic characterization of epithelial ovarian cancer delineates molecular signatures and therapeutic targets in distinct histological subtypes. Nat Commun *14*, 7802. https://doi.org/10.1038/s41467-023-43282-3.

2.   Coscia, F., Watters, K.M., Curtis, M., Eckert, M.A., Chiang, C.Y., Tyanova, S., Montag, A., Lastra, R.R., Lengyel, E., and Mann, M. (2016). Integrative proteomic profiling of ovarian cancer cell lines reveals precursor cell associated proteins and functional status. Nat Commun *7*, 12645. https://doi.org/10.1038/ncomms12645.

3.   Bhattacharya, R., Ghosh, A., and Mukhopadhyay, S. (2023). High-grade serous ovarian carcinoma, the "Achiles' hill" for clinicians and molecular biologists: a molecular insight. Mol Biol Rep *50*, 9511–9519. https://doi.org/10.1007/s11033-023-08760-3.

4.   Colvin, E.K., and Howell, V.M. (2020). Why the dual origins of high grade serous ovarian cancer matter. Nat Commun *11*, 1200. https://doi.org/10.1038/s41467-020-15089-z.

5.   King, S.M., Hilliard, T.S., Wu, L.Y., Jaffe, R.C., Fazleabas, A.T., and Burdette, J.E. (2011). The impact of ovulation on fallopian tube epithelial cells: evaluating three hypotheses connecting ovulation and serous ovarian cancer. Endocrine-Related Cancer *18*, 627–642. https://doi.org/10.1530/ERC-11-0107.

6.   Paik, D.Y., Janzen, D.M., Schafenacker, A.M., Velasco, V.S., Shung, M.S., Cheng, D., Huang, J., Witte, O.N., and Memarzadeh, S. (2012). Stem⬜Like Epithelial Cells Are Concentrated in the Distal End of the Fallopian Tube: A Site for Injury and Serous Cancer Initiation. STEM CELLS *30*, 2487–2497. https://doi.org/10.1002/stem.1207.

7.   Dinh, H.Q., Lin, X., Abbasi, F., Nameki, R., Haro, M., Olingy, C.E., Chang, H., Hernandez, L., Gayther, S.A., Wright, K.N., et al. (2021). Single-cell transcriptomics identifies gene expression networks driving differentiation and tumorigenesis in the human fallopian tube. Cell Reports *35*, 108978. https://doi.org/10.1016/j.celrep.2021.108978.

8.   Yuan, S., Wang, Z., Peng, H., Ward, S.M., Hennig, G.W., Zheng, H., and Yan, W. (2021). Oviductal motile cilia are essential for oocyte pickup but dispensable for sperm and embryo transport. Proc. Natl. Acad. Sci. U.S.A. *118*, e2102940118. https://doi.org/10.1073/pnas.2102940118.

9.   Zhu, M., Iwano, T., and Takeda, S. (2020). Fallopian Tube Basal Stem Cells Reproducing the Epithelial Sheets In Vitro—Stem Cell of Fallopian Epithelium. Biomolecules *10*, 1270. https://doi.org/10.3390/biom10091270.

10.   Slyper, M., Porter, C.B.M., Ashenberg, O., Waldman, J., Drokhlyansky, E., Wakiro,
I., Smillie, C., Smith-Rosario, G., Wu, J., Dionne, D., et al. (2020). A single-cell and
single-nucleus RNA-Seq toolbox for fresh and frozen human tumors. Nat Med *26*, 792–
802. https://doi.org/10.1038/s41591-020-0844-1.

11.   Wu, H., Kirita, Y., Donnelly, E.L., and Humphreys, B.D. (2019). Advantages of
Single-Nucleus over Single-Cell RNA Sequencing of Adult Kidney: Rare Cell Types and
Novel Cell States Revealed in Fibrosis. JASN *30*, 23–32.
https://doi.org/10.1681/ASN.2018090912.

12.   Wiegleb, G., Reinhardt, S., Dahl, A., and Posnien, N. (2022). Tissue dissociation
for single-cell and single-nuclei RNA sequencing for low amounts of input material.
Front Zool *19*, 27. https://doi.org/10.1186/s12983-022-00472-x.

13.   Ding, J., Adiconis, X., Simmons, S.K., Kowalczyk, M.S., Hession, C.C.,
Marjanovic, N.D., Hughes, T.K., Wadsworth, M.H., Burks, T., Nguyen, L.T., et al. (2020).
Systematic comparison of single-cell and single-nucleus RNA-sequencing methods. Nat
Biotechnol *38*, 737–746. https://doi.org/10.1038/s41587-020-0465-8.

14.   Tokunaga, R., Zhang, W., Naseem, M., Puccini, A., Berger, M.D., Soni, S.,
McSkane, M., Baba, H., and Lenz, H.-J. (2018). CXCL9, CXCL10, CXCL11/CXCR3 axis
for immune activation – A target for novel cancer therapy. Cancer Treatment Reviews *63*,
40–47. https://doi.org/10.1016/j.ctrv.2017.11.007.

15.   Wu, L., Awaji, M., Saxena, S., Varney, M.L., Sharma, B., and Singh, R.K. (2020).
IL-17-CXC Chemokine Receptor 2 Axis Facilitates Breast Cancer Progression by Up-
Regulating Neutrophil Recruitment. Am J Pathol *190*, 222–233.
https://doi.org/10.1016/j.ajpath.2019.09.016.

16.   Hu, Z., Artibani, M., Alsaadi, A., Wietek, N., Morotti, M., Shi, T., Zhong, Z.,
Santana Gonzalez, L., El-Sahhar, S., KaramiNejadRanjbar, M., et al. (2020). The
Repertoire of Serous Ovarian Cancer Non-genetic Heterogeneity Revealed by Single-
Cell Sequencing of Normal Fallopian Tube Epithelial Cells. Cancer Cell *37*, 226-242.e7.
https://doi.org/10.1016/j.ccell.2020.01.003.

17.   Lengyel, E., Li, Y., Weigert, M., Zhu, L., Eckart, H., Javellana, M., Ackroyd, S.,
Xiao, J., Olalekan, S., Glass, D., et al. (2022). A molecular atlas of the human
postmenopausal fallopian tube and ovary from single-cell RNA and ATAC sequencing.
Cell Reports *41*. https://doi.org/10.1016/j.celrep.2022.111838.

18.   Ulrich, N.D., Shen, Y., Ma, Q., Yang, K., Hannum, D.F., Jones, A., Machlin, J.,
Randolph, J.F., Smith, Y.R., Schon, S.B., et al. (2022). Cellular heterogeneity of human

fallopian tubes in normal and hydrosalpinx disease states identified using scRNA-seq.
Developmental Cell *57*, 914-929.e7. https://doi.org/10.1016/j.devcel.2022.02.017.

19.   Brand, J., Haro, M., Lin, X., Rimel, B.J., McGregor, S.M., Lawrenson, K., and
Dinh, H.Q. (2024). Fallopian tube single cell analysis reveals myeloid cell alterations in
high-grade serous ovarian cancer. iScience *27*, 108990.
https://doi.org/10.1016/j.isci.2024.108990.

20.   Liu, J., Gao, C., Sodicoff, J., Kozareva, V., Macosko, E.Z., and Welch, J.D.
(2020). Jointly defining cell types from multiple single-cell datasets using LIGER. Nat
Protoc *15*, 3632–3662. https://doi.org/10.1038/s41596-020-0391-8.

21.   Garsed, D.W., Pandey, A., Fereday, S., Kennedy, C.J., Takahashi, K., Alsop, K.,
Hamilton, P.T., Hendley, J., Chiew, Y.-E., Traficante, N., et al. (2022). The genomic and
immune landscape of long-term survivors of high-grade serous ovarian cancer. Nat
Genet, 1–12. https://doi.org/10.1038/s41588-022-01230-9.

22.   Jacomy, M., Venturini, T., Heymann, S., and Bastian, M. (2014). ForceAtlas2, a
Continuous Graph Layout Algorithm for Handy Network Visualization Designed for the
Gephi Software. PLOS ONE *9*, e98679. https://doi.org/10.1371/journal.pone.0098679.

23.   Jiang, P., Du, W., and Wu, M. (2014). Regulation of the pentose phosphate
pathway in cancer. Protein & Cell *5*, 592–602. https://doi.org/10.1007/s13238-014-0082-
8.

24.   Yang, T., Shu, X., Zhang, H.-W., Sun, L.-X., Yu, L., Liu, J., Sun, L.-C., Yang, Z.-H.,
and Ran, Y.-L. (2020). Enolase 1 regulates stem cell-like properties in gastric cancer
cells by stimulating glycolysis. Cell Death Dis *11*, 870. https://doi.org/10.1038/s41419-
020-03087-4.

25.   Wang, L., Long, H., Zheng, Q., Bo, X., Xiao, X., and Li, B. (2019). Circular RNA
circRHOT1 promotes hepatocellular carcinoma progression by initiation of NR2F6
expression. Mol Cancer *18*, 119. https://doi.org/10.1186/s12943-019-1046-7.

26.   Liu, J., Li, T., and Liu, X.-L. (2017). DDA1 is induced by NR2F6 in ovarian cancer
and predicts poor survival outcome. Eur Rev Med Pharmacol Sci *21*, 1206–1213.

27.   Cheng, X.-H., Black, M., Ustiyan, V., Le, T., Fulford, L., Sridharan, A.,
Medvedovic, M., Kalinichenko, V.V., Whitsett, J.A., and Kalin, T.V. (2014). SPDEF
Inhibits Prostate Carcinogenesis by Disrupting a Positive Feedback Loop in Regulation
of the Foxm1 Oncogene. PLoS Genet *10*, e1004656.
https://doi.org/10.1371/journal.pgen.1004656.

28.   Noah, T.K., Lo, Y., Price, A., Chen, G., King, E., Washington, M., Aronow, B.J.,
and Shroyer, N.F. (2013). SPDEF Functions as a Colorectal Tumor Suppressor by
Inhibiting β-Catenin Activity. Gastroenterology *144*, 1012-1023.e6.
https://doi.org/10.1053/j.gastro.2013.01.043.

29.   Lo, Y.-H., Chung, E., Li, Z., Wan, Y.-W., Mahe, M.M., Chen, M.-S., Noah, T.K.,
Bell, K.N., Yalamanchili, H.K., Klisch, T.J., et al. (2017). Transcriptional Regulation by
ATOH1 and its Target SPDEF in the Intestine. Cellular and Molecular Gastroenterology
and Hepatology *3*, 51–71. https://doi.org/10.1016/j.jcmgh.2016.10.001.

30.   De Falco, A., Caruso, F., Su, X.-D., Iavarone, A., and Ceccarelli, M. (2023). A
variational algorithm to detect the clonal copy number substructure of tumors from
scRNA-seq data. Nat Commun *14*, 1074. https://doi.org/10.1038/s41467-023-36790-9.

31.   Shlien, A., and Malkin, D. (2009). Copy number variations and cancer. Genome
Medicine *1*, 62. https://doi.org/10.1186/gm62.

32.   Cui, X., Shan, T., and Qiao, L. (2022). Collagen type IV alpha 1 (COL4A1)
silence hampers the invasion, migration and epithelial–mesenchymal transition (EMT) of
gastric cancer cells through blocking Hedgehog signaling pathway. Bioengineered *13*,
8972–8981. https://doi.org/10.1080/21655979.2022.2053799.

33.   Huang, H., Wright, S., Zhang, J., and Brekken, R.A. (2019). Getting a grip on
adhesion: Cadherin switching and collagen signaling. Biochimica et Biophysica Acta
(BBA) - Molecular Cell Research *1866*, 118472.
https://doi.org/10.1016/j.bbamcr.2019.04.002.

34.   Xu, J., Fang, Y., Chen, K., Li, S., Tang, S., Ren, Y., Cen, Y., Fei, W., Zhang, B.,
Shen, Y., et al. (2022). Single-Cell RNA Sequencing Reveals the Tissue Architecture in
Human High-Grade Serous Ovarian Cancer. Clinical Cancer Research *28*, 3590–3602.
https://doi.org/10.1158/1078-0432.CCR-22-0296.

35.   Johnson, B.K., Rhodes, M., Wegener, M., Himadewi, P., Foy, K., Schipper, J.L.,
Siwicki, R.A., Rossell, L.L., Siegwald, E.J., Chesla, D.W., et al. (2022). Single-cell Total
RNA Miniaturized sequencing (STORM-seq) reveals differentiation trajectories of
primary human fallopian tube epithelium. Preprint at bioRxiv,
https://doi.org/10.1101/2022.03.14.484332 https://doi.org/10.1101/2022.03.14.484332.

36.   Siegel, R.L., Miller, K.D., and Jemal, A. (2016). Cancer statistics, 2016. CA: A
Cancer Journal for Clinicians *66*, 7–30. https://doi.org/10.3322/caac.21332.

37.   Cook, D.P., and Vanderhyden, B.C. (2022). Transcriptional census of epithelial-
mesenchymal plasticity in cancer. Science Advances *8*, eabi7640.
https://doi.org/10.1126/sciadv.abi7640.

38.   Lawrenson, K., Fonseca, M.A.S., Liu, A.Y., Segato Dezem, F., Lee, J.M., Lin, X.,
Corona, R.I., Abbasi, F., Vavra, K.C., Dinh, H.Q., et al. (2019). A Study of High-Grade
Serous Ovarian Cancer Origins Implicates the SOX18 Transcription Factor in Tumor
Development. Cell Reports *29*, 3726-3735.e4.
https://doi.org/10.1016/j.celrep.2019.10.122.

39.   Nakamura, K., Nakayama, K., Ishikawa, N., Ishikawa, M., Sultana, R., Kiyono, T.,
and Kyo, S. (2018). Reconstitution of high-grade serous ovarian carcinoma from
primary fallopian tube secretory epithelial cells. Oncotarget *9*, 12609–12619.
https://doi.org/10.18632/oncotarget.23035.

40.   Izar, B., Tirosh, I., Stover, E.H., Wakiro, I., Cuoco, M.S., Alter, I., Rodman, C.,
Leeson, R., Su, M.-J., Shah, P., et al. (2020). A single-cell landscape of high-grade
serous ovarian cancer. Nat Med *26*, 1271–1279. https://doi.org/10.1038/s41591-020-
0926-0.

41.   Yoshihara, K., Tajima, A., Yahata, T., Kodama, S., Fujiwara, H., Suzuki, M.,
Onishi, Y., Hatae, M., Sueyoshi, K., Fujiwara, H., et al. (2010). Gene Expression Profile
for Predicting Survival in Advanced-Stage Serous Ovarian Cancer Across Two
Independent Datasets. PLoS ONE *5*, e9615.
https://doi.org/10.1371/journal.pone.0009615.

42.   Ganzfried, B.F., Riester, M., Haibe-Kains, B., Risch, T., Tyekucheva, S., Jazic, I.,
Wang, X.V., Ahmadifar, M., Birrer, M.J., Parmigiani, G., et al. (2013).
curatedOvarianData: clinically annotated data for the ovarian cancer transcriptome.
Database *2013*. https://doi.org/10.1093/database/bat013.

43.   Konecny, G.E., Wang, C., Hamidi, H., Winterhoff, B., Kalli, K.R., Dering, J.,
Ginther, C., Chen, H.-W., Dowdy, S., Cliby, W., et al. (2014). Prognostic and Therapeutic
Relevance of Molecular Subtypes in High-Grade Serous Ovarian Cancer. JNCI: Journal
of the National Cancer Institute *106*. https://doi.org/10.1093/jnci/dju249.

44.   Tan, T.Z., Miow, Q.H., Huang, R.Y., Wong, M.K., Ye, J., Lau, J.A., Wu, M.C., Bin
Abdul Hadi, L.H., Soong, R., Choolani, M., et al. (2013). Functional genomics identifies
five distinct molecular subtypes with clinical relevance and pathways for growth control
in epithelial ovarian cancer. EMBO Mol Med *5*, 1051–1066.
https://doi.org/10.1002/emmm.201201823.

45.   Axelrod, M.L., Cook, R.S., Johnson, D.B., and Balko, J.M. (2019). Biological
Consequences of MHC-II Expression by Tumor Cells in Cancer. Clin Cancer Res *25*,
2392–2402. https://doi.org/10.1158/1078-0432.CCR-18-3200.

46.   Sharma, P., Goswami, S., Raychaudhuri, D., Siddiqui, B.A., Singh, P., Nagarajan,
A., Liu, J., Subudhi, S.K., Poon, C., Gant, K.L., et al. (2023). Immune checkpoint
therapy—current perspectives and future directions. Cell *186*, 1652–1669.
https://doi.org/10.1016/j.cell.2023.03.006.

47.   Chen, Y., McAndrews, K.M., and Kalluri, R. (2021). Clinical and therapeutic
relevance of cancer-associated fibroblasts. Nat Rev Clin Oncol *18*, 792–804.
https://doi.org/10.1038/s41571-021-00546-5.

48.   De Francesco, E.M., Lappano, R., Santolla, M.F., Marsico, S., Caruso, A., and
Maggiolini, M. (2013). HIF-1α/GPER signaling mediates the expression of VEGF
induced by hypoxia in breast cancer associated fibroblasts (CAFs). Breast Cancer Res
15, R64. https://doi.org/10.1186/bcr3458.

49.   Shapouri☐Moghaddam, A., Mohammadian, S., Vazini, H., Taghadosi, M.,
Esmaeili, S., Mardani, F., Seifi, B., Mohammadi, A., Afshari, J.T., and Sahebkar, A.
(2018). Macrophage plasticity, polarization, and function in health and disease. Journal
Cellular Physiology *233*, 6425–6440. https://doi.org/10.1002/jcp.26429.

50.   Allavena, P., Sica, A., Garlanda, C., and Mantovani, A. (2008). The Yin☐Yang of
tumor☐associated macrophages in neoplastic progression and immune surveillance.
Immunological Reviews *222*, 155–161. https://doi.org/10.1111/j.1600-
065X.2008.00607.x.

51.   Mantovani, A., Allavena, P., Sica, A., and Balkwill, F. (2008). Cancer-related
inflammation. Nature *454*, 436–444. https://doi.org/10.1038/nature07205.

52.   Varol, C. (2019). Tumorigenic Interplay Between Macrophages and Collagenous
Matrix in the Tumor Microenvironment. In Collagen Methods in Molecular Biology., I.
Sagi and N. A. Afratis, eds. (Springer New York), pp. 203–220.
https://doi.org/10.1007/978-1-4939-9095-5_15.

53.   Ledermann, J.A., Drew, Y., and Kristeleit, R.S. (2016). Homologous
recombination deficiency and ovarian cancer. European Journal of Cancer *60*, 49–58.
https://doi.org/10.1016/j.ejca.2016.03.005.

54.   Kuchenbaecker, K.B., Hopper, J.L., Barnes, D.R., Phillips, K.-A., Mooij, T.M.,
Roos-Blom, M.-J., Jervis, S., Van Leeuwen, F.E., Milne, R.L., Andrieu, N., et al. (2017).
Risks of Breast, Ovarian, and Contralateral Breast Cancer for *BRCA1* and *BRCA2*
Mutation Carriers. JAMA *317*, 2402. https://doi.org/10.1001/jama.2017.7112.

55.   Stanciu, P.I., Ind, T.E.J., Barton, D.P.J., Butler, J.B., Vroobel, K.M., Attygalle, A.D.,
and Nobbenhuis, M.A.E. (2019). Development of Peritoneal Carcinoma in women
diagnosed with Serous Tubal Intraepithelial Carcinoma (STIC) following Risk-Reducing

Salpingo-Oophorectomy (RRSO). J Ovarian Res *12*, 50. https://doi.org/10.1186/s13048-
019-0525-1.

56.   Chen, F., Gaitskell, K., Garcia, M., Albukhari, A., Tsaltas, J., and Ahmed, A.
(2017). Serous tubal intraepithelial carcinomas associated with high☐grade serous
ovarian carcinomas: a systematic review. BJOG *124*, 872–878.
https://doi.org/10.1111/1471-0528.14543.

57.   Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer
for Illumina sequence data. Bioinformatics *30*, 2114–2120.
https://doi.org/10.1093/bioinformatics/btu170.

58.   Dobin, A., and Gingeras, T.R. (2015). Mapping RNA-seq Reads with STAR. Curr
Protoc Bioinformatics *51*, 11.14.1-11.14.19.
https://doi.org/10.1002/0471250953.bi1114s51.

59.   Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general
purpose program for assigning sequence reads to genomic features. Bioinformatics *30*,
923–930. https://doi.org/10.1093/bioinformatics/btt656.

60.   Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data.
Bioinformatics *26*, 139–140. https://doi.org/10.1093/bioinformatics/btp616.

61.   Chen, Y., Lun, A.T.L., and Smyth, G.K. (2016). From reads to genes to pathways:
differential expression analysis of RNA-Seq experiments using Rsubread and the
edgeR quasi-likelihood pipeline. F1000Res *5*, 1438.
https://doi.org/10.12688/f1000research.8987.2.

62.   Zheng, G.X.Y., Terry, J.M., Belgrader, P., Ryvkin, P., Bent, Z.W., Wilson, R.,
Ziraldo, S.B., Wheeler, T.D., McDermott, G.P., Zhu, J., et al. (2017). Massively parallel
digital transcriptional profiling of single cells. Nat Commun *8*, 14049.
https://doi.org/10.1038/ncomms14049.

63.   Wolf, F.A., Angerer, P., and Theis, F.J. (2018). SCANPY: large-scale single-cell
gene expression data analysis. Genome Biology *19*, 15. https://doi.org/10.1186/s13059-
017-1382-0.

64.   Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., Zheng, S., Butler, A., Lee,
M.J., Wilk, A.J., Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal
single-cell data. Cell *184*, 3573-3587.e29. https://doi.org/10.1016/j.cell.2021.04.048.

65.   Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R.,
Grolemund, G., Hayes, A., Henry, L., Hester, J., et al. (2019). Welcome to the Tidyverse.
JOSS *4*, 1686. https://doi.org/10.21105/joss.01686.

66.   Angerer, P., Haghverdi, L., Büttner, M., Theis, F.J., Marr, C., and Buettner, F.
(2016). destiny: diffusion maps for large-scale single-cell data in R. Bioinformatics *32*,
1241–1243. https://doi.org/10.1093/bioinformatics/btv715.

67.   Wolf, F.A., Hamey, F.K., Plass, M., Solana, J., Dahlin, J.S., Göttgens, B.,
Rajewsky, N., Simon, L., and Theis, F.J. (2019). PAGA: graph abstraction reconciles
clustering with trajectory inference through a topology preserving map of single cells.
Genome Biol *20*, 59. https://doi.org/10.1186/s13059-019-1663-x.

68.   Moon, K.R., van Dijk, D., Wang, Z., Gigante, S., Burkhardt, D.B., Chen, W.S.,
Yim, K., Elzen, A. van den, Hirn, M.J., Coifman, R.R., et al. (2019). Visualizing structure
and transitions in high-dimensional biological data. Nature Biotechnology *37*, 1482–
1492. https://doi.org/10.1038/s41587-019-0336-3.

69.   Sherman, B.T., Hao, M., Qiu, J., Jiao, X., Baseler, M.W., Lane, H.C., Imamichi, T.,
and Chang, W. (2022). DAVID: a web server for functional enrichment analysis and
functional annotation of gene lists (2021 update). Nucleic Acids Res *50*, W216–W221.
https://doi.org/10.1093/nar/gkac194.

70.   La Manno, G., Soldatov, R., Zeisel, A., Braun, E., Hochgerner, H., Petukhov, V.,
Lidschreiber, K., Kastriti, M.E., Lönnerberg, P., Furlan, A., et al. (2018). RNA velocity of
single cells. Nature *560*, 494–498. https://doi.org/10.1038/s41586-018-0414-6.

71.   Bergen, V., Lange, M., Peidli, S., Wolf, F.A., and Theis, F.J. (2020). Generalizing
RNA velocity to transient cell states through dynamical modeling. Nat Biotechnol *38*,
1408–1414. https://doi.org/10.1038/s41587-020-0591-3.

72.   Aibar, S., González-Blas, C.B., Moerman, T., Huynh-Thu, V.A., Imrichova, H.,
Hulselmans, G., Rambow, F., Marine, J.-C., Geurts, P., Aerts, J., et al. (2017). SCENIC:
single-cell regulatory network inference and clustering. Nature Methods *14*, 1083–1086.
https://doi.org/10.1038/nmeth.4463.

73.   Cline, M.S., Smoot, M., Cerami, E., Kuchinsky, A., Landys, N., Workman, C.,
Christmas, R., Avila-Campilo, I., Creech, M., Gross, B., et al. (2007). Integration of
biological networks and gene expression data using Cytoscape. Nat Protoc *2*, 2366–
2382. https://doi.org/10.1038/nprot.2007.324.

74.   Patel, A.P., Tirosh, I., Trombetta, J.J., Shalek, A.K., Gillespie, S.M., Wakimoto, H.,
Cahill, D.P., Nahed, B.V., Curry, W.T., Martuza, R.L., et al. (2014). Single-cell RNA-seq

highlights intratumoral heterogeneity in primary glioblastoma. Science *344*, 1396–1401.
https://doi.org/10.1126/science.1254257.

75.   Cheng, K., Seita, Y., Moriwaki, T., Noshiro, K., Sakata, Y., Hwang, Y.S., Torigoe,
T., Saitou, M., Tsuchiya, H., Iwatani, C., et al. (2022). The developmental origin and the
specification of the adrenal cortex in humans and cynomolgus monkeys. Science
Advances *8*, eabn8485. https://doi.org/10.1126/sciadv.abn8485.

1062

1063

1064

FIGURE LEGENDS

Figure 1. Bulk RNA-seq analyses unveil transcriptomic correlation among human HGSOCs, fallopian tubes and ovaries

(A) Overview of the study design and workflow.

(B) Principal component (PC) analyses of 26 human samples analyzed by bulk RNA-seq, including 10 HGSOC, 4 ovarian, and 12 segmented fallopian tube samples. EOC, epithelial ovarian cancer; FTF, fallopian tube fimbria; FTA, fallopian tube ampulla; FTI, fallopian tube isthmus; OV, ovary.

(C) Dendrogram showing transcriptomic distance among all 26 samples analyzed by bulk RNA-seq based on ward.D2 method.

(D) Heatmap showing the top 2,000 variable genes and the transcriptomic signatures of EOC-Fs (Fallopian-derived EOCs), fallopian tubes, and ovaries. Color bar showing Z-score normalized gene expression. Red is for higher expression and blue is for lower expression.

(E) Gene ontology (GO) term enrichment analyses of seven genesets representing the transcriptomic signatures of EOC-Fs (Fallopian-derived EOCs), fallopian tubes, and ovaries.

See also Figures S1, Tables S1.

Figure 2. snRNA-seq analyses reveal human FTE cell subpopulations and the development trajectory

(A) UMAP representation of 12 distinct FTE cell clusters revealed by graph-based Louvain clustering. Each color represents a cell cluster, and the color scheme remains the same in all subsequent panels (B-F).

(B) Dendrogram showing hierarchical clustering of the twelve subpopulations identified in human FTE. Cells were partitioned using ward.D2 based on the average expression of each cell cluster.

(C) Bubble plot showing average expression levels and prevalence of major markers used to annotate the FTE cell types. Color indicating averaged gene expression and the size of circle showing the percentage of cells expressing corresponding gene in the cell cluster.

(D) RNA velocity analyses of all FTE cell subpopulations to predict the developmental
trajectory. Each color represents a cell cluster. Arrows pointing the potential
developmental directions.

(E) PAGA analyses of developmental trajectory and connectivity among 12 FTE
subpopulations. Each color represents a cell cluster, and the thickness of the lines
represents correlations between FTE cell clusters.

(F) Visualization of the developmental trajectories using PHATE. Each point represents
a cell, and each color stands for a cell sub-cluster. Branches show different
differentiation directions or fates.

(G) Pseudotime of the developmental trajectories. Pseudotime was calculated based on
the principal curve by PHATE embedding regardless of sub-clusters.

(H and I) Pseudotime heat maps showing the transcriptomic dynamics from FTE stem
cells to secretory cells (H) and ciliated cells (I) in human FTE. Color bar showing the Z-
score normalized gene expression.

(J and K) Immunofluorescence showing the localization and co-expression of LGR5,
PGR and Ki67 in the isthmus and ampulla segments of human FTE. White, DAPI; Blue,
Ki67; Green, PGR; Red, LGR5. Yellow arrows point to the small round basal stem cells.
Scale bars, 10μm.

See also Figures S2, Tables S3 and S4.

1115

Figure 3. snRNA-seq analyses uncover a high degree of variation and cellular
heterogeneity of HGSOCs.

(A) UMAP representation of 13 distinct cell subclusters identified in HGSOC samples by
graph-based Louvain clustering. Each color represents a cell cluster.

(B) Dendrogram showing hierarchical clustering of the 13 subclusters identified in
HGSOC samples. The HGSOC cells were partitioned using ward.D2 based on the
average expression of each cell cluster. CAFs, cancer-associated fibroblasts; EMT,
epithelial-mesenchymal transition; MHC, major histocompatibility complexes.

(C) UMAP representation of 8 cell subgroups identified in HGSOC samples. Each color
represents a subgroup of cells in HGSOC samples.

(D) Heatmap showing the transcriptomic signatures for each of the eight HGSOC cell
subtypes. Enriched GO terms in each of the 8 subtypes of HGSCO cells are presented
next to the heatmap.

(E and F) UMAP display of cell subclusters and their identity in all malignant cells of
HGSOC samples.

(G) Calculated correlation of transcriptomic distance of pure malignant cell subgroups
identified in HGSOC samples. Color density represents correlation coefficient size.

(H) Average expression levels and prevalence of major markers used to annotate the
malignant cell types in HGSOC samples.

See also Figures S3 and Table S5.

1136

Figure 4. snRNA-seq analyses predict the trajectory leading to the transformation
from normal FTE cells to HGSOC cells

(A) Combination of cancer cells of HGSOC and all FTE cells by force-directed graph
visualization identified 15 cell clusters that were further divided into three big partitions
for predicting potential cell lineages using Slingshot.

(B) Cell trajectories built by PAGA. Each color represents a cell cluster, and the line
thickness indicates the degree of correlation among all cell clusters.

(C) RNA velocity analyses based on force-directed graph embedding, with the arrows
pointing to the developmental directions.

(D) Pseudotime analyses of all tumor and FTE cells regardless of subclusters. The
differentiation from 0 to 1 is represented by blue to yellow.

(E) Dendrogram showing hierarchical clustering of the 15 subclusters identified in
combined HGSOC tumor and FTE cells, which were partitioned using ward.D2 based
on the average expression of each cell cluster.

(F) Histogram showing the proportional composition of 15 cell subtypes in the three
segments of human fallopian tubes (Isth: isthmus; Amp: ampulla; Fimb: fimbria) and
HGSOC.

(G) Bubble plot showing expression of top marker genes in the 15 cell clusters identified.

(H and I) Pseudotime heatmaps showing the transcriptomic dynamics from FTE stem
cells to immunoreactive-type tumor cells (cluster 12) (H) and mesenchymal-type tumor
cells (cluster 13) (I).

See also Table S7 and S8.

1159

Figure 5. SCENIC analyses of snRNA-seq data.

(A) Single-cell regulatory network inference and clustering (SCENIC) analyses of the combined HGSOC tumor and FTE cells, with five cell subtypes annotated.

(B) SCENIC analyses of transcription factor (TF) regulatory networks. Heatmap was drawn using Cytoscape, and the rows (representing genes) and columns (indicating cell subtypes) were ordered using unsupervised hierarchical clustering based on Euclidian distances. Each module represents a cell.

(C) Specific regulons in the five cell subtypes are listed based on AUC score.

(D) Overlapping between gene expression (the top panels) and AUC score (the bottom panels) of the regulons.

(E) Genes regulatory networks with predicted master regulators in different cell subtypes.

Figure 6. The copy number variation of HGSOC.

(A) Significant copy number variants (CNVs) detected in HGSOC, with FTE cells as reference.

(B) CNV regions detected using the Hidden Markov Model (HMM). The blue-to-red color transition represents the CNV region's state from 0 to 6. Each row represents a region.

(C) Pie chart showing the proportional distribution of the six CNV states (top panels) and histogram indicating the frequency of the six states of CNV regions on each chromosome (bottom panel). State1 is 0x: complete loss, State2 is 0.5x: loss of one copy, State3 is 1x: neutral, State4 is 1.5x: addition of one copy, State5 is 2x: addition of two copies, State6 is 3x: essentially a placeholder for >2x copies but modeled as 3x..

(D) Heatmaps showing the gain or loss of gene copy numbers.

See also Table S9.

Figure 7. Schematic summary of the major findings of the present study.

1188 **Figure S1-S5**

1189

Figure S1. Transcriptomic correlation among human HGSOCs, fallopian tubes,
and ovaries revealed by Bulk RNA-seq.

(A) Box graph showing expression distribution, based on FPKM values, in each of the
26 samples analyzed by bulk RNA-seq.

(B) Heatmap showing pair-wise Pearson correlation coefficient among the 26 samples.

(C, D, E and F) Scatter plot of the DEGs between fallopian-derived EOC (EOC_F) and
ovarian-derived EOC (EOC_O) (C), EOC_F and fallopian tube (D), EOC_F and ovary
(E), fallopian tube and ovary (F), and top enriched GO terms.

1198

Figure S2. Proportional composition of major cell subtypes and functional
enrichment in human fallopian tubes uncovered by snRNA-seq.

(A) snRNA-seq analyses of human fallopian tubes led to 87,920 cells that passed
through the initial quality control, 36,000-63,000 mean reads per cell, and ~2,000
median genes per cell.

(B and C) UMAP representation of cell cycle scoring of all FTE cells after removing cell
cycle effects, colored by (B) cell cycle state and (C) samples.

(D) UMAP representation of five major cell types identified in human fallopian tubes.

(E) UMAP representation of the expression of selected cell subgroup marker genes.

(F) Heatmap showing top genes and enriched gene ontology (GO) terms in each of the
five major cell types in human fallopian tubes. Expression of genes is Z-score
normalized.

(G) Histogram showing the proportional distribution of five major cell types in three
segments of human fallopian tubes.

(H) Expression of seven key marker genes projected to PHATE-based trajectory
analyses.

(I) Histograms showing the proportional distribution of subtypes in three segments of
human fallopian tubes.

(J) Histogram showing proportional distribution of 12 subclusters in three segments of
the human fallopian tubes.

See also Table S2.

1220

Figure S3. Prediction and identification of cell subtypes of the fallopian tube and
HGSOC.

(A and B) Immunohistochemical analyses showing the localization (A) and semi-
quantitative results (B) of TUBB4B (red arrow) and PAX8 (yellow arrow) in three
segments of human fallopian tubes. Scale bars, 100µm. The statistical results are
presented as means ± SEM. Student's t-test. ns ≥ 0.05; *p < 0.05; **p < 0.01; ***p <
0.001. Isth: isthmus, Amp: ampulla; Fim: fimbria.

(C) Immunofluorescence analysis showing the localization and co-expression of LGR5,
PGR, and Ki67 in the FTE of fimbria. White, DAPI; Blue, Ki67; Green, PGR; Red, LGR5.
Scale bars, 50µm.

(D) Quality control data of 4 HGSOC samples in scRNA-seq analyses.

(E and F) UMAP representation of cell cycle scoring of all HGSOC cells after removing
cell cycle effects, colored by (E) cell cycle state and (F) samples.

(G) Pearson correlation among four HGSOC samples.

(H) UMAP representation of the expression of selected cell subgroup marker genes.

1236

Figure S4. Specificity and heterogeneity of pure HGSOC tumor cells in snRNA-
seq.

(A and B) Immunohistochemical analyses showing localization (A) and relative
abundance (B) of PAX8, TUBB4B, MUC16, and Ki67 in HGSOC and isthmus of the
fallopian tube. Scale bars, 20µm. The statistical results are presented as means ± SEM.
Student's t-test; ns ≥ 0.05; **p< 0.01; ****p< 0.0001. Isth: isthmus; HGSOC: High-grade
serous ovarian cancer.

(C) Heatmap showing top genes and enriched gene ontology (GO) terms in each of the
five major HGSOC tumor cell subtypes. Expression of genes is Z-score normalized.

(D) UMAP representation of the expression of selected HGSOC tumor cell subgroup
marker genes.

See also Table S6.

1249

Figure S5. Intercellular connections analysis in HGSOC.

(A and B) Ligand-receptor connections between each cancer and non-cancer cell
subgroup of HGSOC, including number of interactions (A) and weights/strength of
interactions (B). Lines indicate intercellular connections between ligands and receptors,
and the line thickness is proportional to the number of ligand-receptor pairs.

(C) Overview of selected interactions of ligand-receptor pairs in outgoing and incoming
signaling patterns**.**

(D) Ligand-receptor connections between each cancer and non-cancer cell subgroup of
HGSOC in the collagen signaling pathway network.

(E) Violin plot showing marker genes of collagen signaling pathways in HGSOC tumor
and non-tumor cell subgroups.

1261

SUPPLEMENTARY DATASETS: see separate Excel documents

Table S1. Top 1000 variable genes among different samples, related with Figure 1DE.

Table S2. Markers identified among different cell types in human fallopian tube snRNA-seq, related with Figure S2F.

Table S2. Top 2000 variable genes along the secretory trajectory, related with Figure 2H.

Table S4. Top 2000 variable genes along the ciliated trajectory, related with Figure 2I.

Table S5. Markers identified in HGSOC snRNA-seq, related with Figure 3D.

Table S6. Markers identified among subtypes of HGSOC cancer, related with Figure S4C.

Table S7. Top 200 variable genes along the trajectory from cluster 11 to cluster 12, related with Figure 4H.

Table S8. Top 200 variable genes along the trajectory from cluster 11 to cluster13, related with Figure 4I.

Table S9. HMM predicted CNVs regions, related with Figure 6B.

Figure 1

Figure 1

A Healthy fallopian tube / Ovarian cancer — Sample collection — Bulk RNA-seq + 10X snRNA-seq

B PCA plot (PC1, PC2, PC3)

C 18579 genes ward.D2

D Z-score normalized expression (-3 to 3)

E

GO:0008584~male gonad development
NR5A1, STAR, AMHR2, KIT, GATA4, INHA
1 GO:0007179~transforming growth factor beta receptor signaling pathway
LEFTY1, AMHR2, BAMBI, WFIKKN2, LEFTY2
GO:0001541~ovarian follicle development
KIT, FOXL2, INHA

GO:0006955~immune response
IGHV2-5, IGHV1-2, IGHV3-48, IGLV3-1, IGHV4-39, IGKV3D-15, IGLV8-61,
JCHAIN, IGLV2-11, IGLV2-8, IGLV3-25, IGLV1-40, IGKV1-16, IGKC,...
2 GO:0010628~positive regulation of gene expression
OSR2, IL33, SLC24A3, PID1, CITED2, TCF23, PDE2A, MIR27B, GATA5, FBLN1, GATA2,
KLF4, ACTG2, ACTA2, CLDN5, SFRP4, ALDH1A2, HAND2, ID1, EMILIN1

GO:0035914~skeletal muscle cell differentiation
EGR1, NR4A1, EGR2, BTG2, FOS, ATF3
3 GO:0006357~regulation of transcription from RNA polymerase II promoter
NR4A2, EGR1, CSRNP1, NR4A1, EGR2, JUN, EGR3, FOSB, FOS, JUNB, ATF3

GO:0048704~embryonic skeletal system morphogenesis
HOXA3, HOXB6, HOXA5, HOXA4, HOXB5
4 GO:0048793~pronephros development
HNF1B, PAX2
GO:0007501~mesodermal cell fate specification
SFRP2, PAX2

GO:0007275~multicellular organism development
EHF, PAEP, HOXB2, EPHA1, SPDEF
5 GO:0006357~regulation of transcription from RNA polymerase II promoter
HES6, EHF, HOXD1, IRX3, RORC, HOXB4, FOXJ1, HOXB2, IRF6, POU5F1, SPDEF

GO:0003341~cilium movement
ODAD1, RSPH4A, ODAD3, DNAI2, CFAP73, SPEF1, SPAG6, ODAD4,
CCDC103, DNAAF1, DNAH9, RSPH9, CFAP206, TTC29, CFAP100, ...
6 GO:0030317~flagellated sperm motility
DRC7, ODAD3, CFAP52, TCTE1, CFAP251, CATSPERD, TEKT1, TEKT2, ...

GO:0006508~proteolysis
ST14, MMP7, MMP1, TMPRSS3, TMPRSS4, HPN, MMP3, KLK5, PRSS22,
KLK8, MMP9, KLK11, WFDC2, KLK7, MMP10, C2, PRSS21, MMP11, ..
7 GO:0051301~cell division
UBE2C, CDCA5, CDCA8, BUB1B, ZWINT, AURKB, AURKA, CDC20, CCNB2, ...
GO:0031424~keratinization
KRT81, KRT17, KRT8, KRT7, SFN, KRT5, KRT6A

Figure 2

Figure 3

Figure 4

Figure 6

Figure 7