1 **Designing a robust whole-cell biosensor for detection of toxic metals**
2 **using intein splicing inhibition of *Mycobacterium tuberculosis* SufB protein**
3

Ashwaria Mehra[1†], Ananya Nanda[1,2†], and Sasmita Nayak[1, 3#]

[1] School of Biotechnology, Kalinga Institute of Industrial Technology Deemed to be University,
Bhubaneswar, Odisha, India 751024.
[2] Toronto Metropolitan University, 350 Victoria St, Toronto, ON M5B 2K3, Canada
[3] Department of Microbiology, Kalinga Institute of Medical Sciences, Kalinga Institute of
Industrial Technology Deemed to be University, Bhubaneswar, Odisha, India 751024

11
12 [#] Corresponding Author: sasmita.n@kiitbiotech.ac.in

13
14 [†] These authors have contributed equally to this work and share first authorship
15

**Running Title:** Intein-based whole-cell biosensor detects toxic metals

**Abbreviation:**

Ni-NTA: Nickel-nitrilotriacetic acid

TCEP.HCl: Tris 2-carboxyl ethyl phosphine. Hydrochloric acid

IPTG: Isopropyl β-d-1-thiogalactopyranoside

Trp: Tryptophan

SDS: Sodium Dodecyl Sulfate

PAGE: Polyacrylamide gel electrophoresis

ICP-OES: Inductively Coupled Plasma Optical Emission Spectroscopy

UV-Vis: UV-visible spectroscopy

PVDF: Polyvinylidene fluoride

BSA: Bovine serum albumin

*Mtb*: *Mycobacterium tuberculosis*

WT: Wild type

SI: Splicing inactive

$Cd^{2+}$: Cadmium

$Hg^{2+}$: Mercury

$Pb^{2+}$: Lead

$Cr^{3+}$: Chromium

$Cu^{2+}$: Copper

$Zn^{2+}$: Zinc

$Pt^{4+}$: Platinum

**Abstract**

Disruption of the natural geochemical cycle by human activities has led to bioaccumulation of

metals, posing a global health threat. Hence, there is a pressing need for simple, sensitive, yet eco-

friendly biosensor setups to monitor metal contamination in the environment. Existing biosensors

are limited by poor efficiency, stability issues, and complex instrumentation requiring skilled

operators. To address these caveats, current study explores how intein-mediated protein splicing,

a spontaneous post-translational process, can be adapted for metal-biosensing by coupling metal-

dependent splicing inhibition to viability loss of native microbial cells. Toxic metal ions like $Cd^{2+}$

and $Hg^{2+}$ attenuated the splicing activity of *Mtb* SufB precursor protein over a concentration range

of 25 μM to 2 mM, while $Pb^{2+}$ and $Cr^{3+}$ failed to do so. An innovative biosensor platform was

designed for colorimetric detection of metal ions via simple Alamar Blue assay, where attenuated

*Mtb* strain (H37Ra) served as the indicator cells. Metal-induced SufB splicing inhibition led to loss

of viability of H37Ra cells, while addition of metal-specific chelators reversed the effect.

Multiplexing ability was evaluated by including known splicing inhibitors like $Cu^{2+}$, $Zn^{2+}$, and $Pt^{4+}$

over various concentration range alongside $Cd^{2+}$ and $Hg^{2+}$. The simple 96-well plate format enables

multiplexed qualitative metal detection, while colorimetric absorbance measurement ensures metal

quantification. The designed biosensor offers low-cost, user-friendly, and sensitive assay for high-

throughput metal detection, utilizing whole-cell native organisms carrying metal-sensing precursor

protein. Thus, this approach can be implemented in standard biological laboratories for robust

metal screening process in environmental and industrial effluents.

**Keywords:** Whole-cell biosensor, heavy metal ion detection, *Mycobacterium tuberculosis* SufB*,

intein-based biosensor, protein splicing inhibition.

**Introduction**

Heavy metal contamination has emerged as one of the most persistent environmental and public

health challenges of the modern era. Over 35 metals have been identified in their ore forms within

the Earth's crust, many of which are essential nutrients. Among these, 23 metals are classified as

heavy metals due to their high density (>5 g/cm³) and atomic weights (>40.0), indicating their

potential for environmental toxicity (1–3). Human activities have significantly disrupted natural

geochemical cycles, causing bioaccumulation and biomagnification through contaminated food,

water, or direct exposure, thereby posing serious health risks (3–7).

Current analytical methods, such as electrothermal atomic absorption spectrometry (ETAAS),

inductively coupled plasma-optical emission spectrometry (ICP-OES), high-pressure liquid

chromatography combined with ICP-mass spectrometry (HPLC-ICP-MS), hydride generation

atomic absorption spectrometry (HGAAS), and X-ray fluorescence spectrometry (XRF), offer

excellent sensitivity for detecting metals. Nonetheless, these methods involve complex

instrumentation, require skilled operators, generate hazardous waste, contributing further to

environmental pollution (8–11). To overcome these limitations, biosensors have emerged as a

promising alternative offering affordability, and long-term operational stability (8). According to

the International Union of Pure and Applied Chemistry (IUPAC), biosensors integrate biological

recognition elements directly with transducers to provide precise quantitative data (12, 13).

Although enzyme-based electrochemical biosensors have been widely explored, their broader

implementation is limited due to poor electron transfer efficiency, and instability, limiting their

commercial reusability (14–20). In comparison, whole-cell biosensors present a more practicable

approach by using live cell sensing units, with advantages such as stable signal generation, and

scalability. For instance, Chouteau et al. synthesized BSA encapsulated microalgae as

bioreceptors to detect metals ($Cd^{2+}$, $Zn^{2+}$, and $Pb^{2+}$) (16). However, the sensitivity of detection was

affected due to variability in algal loading and distribution within BSA membrane. In another

work, transformed protozoan cells were used as whole-cell biosensor, linking the metallothionein

promoter-based metal detection to luciferase reporter system (21). Despite the innovation, signal

variability compromised the specificity and sensitivity of heavy metal detection. In another study,

Kim and colleagues engineered *E. coli* cells carrying plasmids with metal sensing promoters to

detect metals such as $Cu^{2+}$, $Zn^{2+}$, and $Cd^{2+}$ (22). However, the $Cu^{2+}$ and $Cd^{2+}$ sensing strains were

not selective and could also detect other metals. Further, detection of $Cu^{2+}$ was hindered by the

active transport of $Cu^{2+}$ ions out of the cytoplasm by the endogenous CopA protein, suppressing

reporter gene expression. Other challenges with whole-cell biosensors include declining cell

viability causing delayed response time, leaky gene expression inducing false positive signals, and

ecological threats of using genetically modified organisms (23–29). Addressing these

shortcomings, intein-mediated protein engineering techniques open up new avenues for creating

innovative cell-based sensors (30–33).

Inteins are protein splicing elements within precursor proteins that can self-excise via multi-step

nucleophilic displacement reactions, to generate active proteins (Supplemental Fig. S1) (34, 35).

This is a post-translational process that does not require energy sources or cofactors, hence inteins

can be seamlessly integrated into biosensor units minimizing interference with sensor cell

functionality (30). Furthermore, intein splicing in precursor proteins can be regulated via

conditional protein splicing (CPS), in presence of small molecules such as metal ions and metal-

based nanoparticles (Table 1) (36–42). For instance, a recent study from our group has examined

the splicing and N-terminal cleavage activity of *Mycobacterium tuberculosis* (*Mtb*) SufB precursor

protein in the presence of metal ions such as $Cu^{2+}$, $Zn^{2+}$, $Pt^{4+}$, and $Fe^{3+}/Fe^{2+}$ (43).  Prior studies

have also reported $Zn^{2+}$ mediated splicing regulation of PI-*Sce* VMA intein, *Mtb* RecA intein, and

*Ssp* PCC6803 DnaE split intein (44, 45). Effects of metal ions, such as $Cu^{+/2+}$ and CisPt, have been

evaluated on *Mtb* RecA, *Cryptococcus neoformans* Prp8, and *Mycobacterium smegmatis* DnaBi1

intein splicing (38, 39, 44, 46–51).

116    **Table 1:** Overview of intein-harboring organisms and reported modulation of splicing activity by
117    metal ions

| | Organisms | Intein containing precursor proteins | Metal ions | References |
|---|---|---|---|---|
| **Bacteria** | *Mycobacterium tuberculosis* | RecA | $Zn^{2+}$<br>$Cd^{2+}$<br>$Co^{2+}$<br>$Ni^{2+}$<br>$Cu^{+}/Cu^{2+}$<br>CisPt | (45, 47, 52, 53) |
| | *Mycobacterium tuberculosis* | SufB | $Zn^{2+}$<br>$Cu^{2+}$<br>$Pt^{4+}$<br>$Fe^{2+/3+}$ | (43) |
| | *Mycobacterium smegmatis* | DnaB-intein1 (DnaBi1) | $Zn^{2+}$ | (49) |
| | *Synechocystis* sp. PCC6803 | *Ssp* DnaE | $Zn^{2+}$<br>$Cd^{2+}$ | (45) |
| | *Oscillatoria limnetica* | split DnaE intein | $Cu^{2+}$ | (54) |
| | *Thermosynechoc occus vulcanus* | DnaE | $Cu^{2+}$ | (54) |
| | *Nostoc* sp. PCC7120 | DnaB | $Cu^{2+}$ | (54) |
| **Fungi** | *Saccharomyces cerevisiae* | PI-*Sce*I VMA | $Zn^{2+}$ | (44) |
| | *Cryptococcus neoformans* | Prp8 | $Zn^{2+}$<br>$Cu^{2+}$<br>CisPt | (39, 51) |
| | *Cryptococcus gattii* | Prp8 | CisPt | (51) |

Conditional splicing activity of intein-containing proteins has been used to (30, 55–58) design

biosensors to detect protein~protein interaction in living cells via split GFP (52) or Luciferase

reporter system (53–55). Although several fundamental studies have reported splicing regulation

of precursor protein in presence of oxidative/nitrosative stresses (56), metal ions (43),

nanoparticles (40) correlating to pathogen survival, a versatile and native whole cell intein-based

biosensor, especially for toxic metal ion detection, is yet to be elucidated.

However, concerns pertaining to the functionality and usability of the intein-based metal-sensing

biosensor remains, such as: 1) If an eco-friendly whole cell biosensor system can be designed using

native cells without any genetic modification; 2) Whether the designed biosensor can offer an user-

friendly and low-cost approach?;  3) If the working protocol can be simple, yet effective for a

specific, and sensitive metal detection?; And finally, 4) Whether this approach can be applied for

a multiplexed metal screening process, for qualitative and quantitative detection of different metal

ions?  Current research explores how toxic metal ions, like cadmium ($Cd^{2+}$), mercury ($Hg^{2+}$),

chromium ($Cr^{3+}$), and lead ($Pb^{2+}$), affect the splicing and N-terminal cleavage reactions of *Mtb*

SufB precursor protein. The fundamental knowledge gained from these results was applied to

develop a biosensor platform to detect toxic metals, using native mycobacterial cells in a 96-well

plate format.  Splicing regulation in metal-sensitive precursor protein *Mtb* SufB induced alteration

in mycobacterial cell viability, that indicated the presence of metal ions.

*Mtb* SUF protein complex constitutes an unique pathway for Fe-S cluster assembly in

mycobacteria, and its activity is dependent on the splicing of SufB precursor, producing functional

SufB protein (60). Thus, splicing inhibition which blocks the generation of active SufB, can retard

mycobacterial growth and viability by impairing Fe-S cluster biogenesis (61). In the current work,

*in vitro* studies demonstrated significant inhibition of *Mtb* SufB precursor protein splicing in

presence of toxic metal ions such as $Cd^{2+}$, and $Hg^{2+}$ over a concentration range of 25 µM to 2 mM.

Further, time-dependent assay revealed that $Cd^{2+}$ and $Hg^{2+}$ inhibit *Mtb* SufB intein splicing as early

as 10 minutes following exposure to the metal ions.  Building on this observation, a biosensor

platform was designed using  native H37Ra (an attenuated *Mtb* strain) as the metal-indicator cells.

Alamar Blue assay performed in a 96-well microtiter plate, detected loss of cell viability in SufB-

containing H37Ra cells in presence of toxic metal ions, $Cd^{2+}$ and $Hg^{2+}$. In a control study,

*Mycobacterium smegmatis* (*M. sm*), which expresses an intein-less SufB protein, persisted as

healthy, viable cells under similar experimental conditions. These results highlight the sensitivity

of H37Ra-based biosensor platform to detect toxic metal ions over a concentration range of 25 µM

to 2 mM. The specificity of the designed biosensor was assessed by adding respective metal

chelators, which protected mycobacterial cell viability, by blocking the splicing-inhibitory activity

of metal ions. Moreover, multiplexing and quantitative potential of the biosensor platform was

validated by using various concentrations of known splicing inhibitors [$Zn^{2+}$, $Cu^{2+}$, and $Pt^{4+}$ (43)]

alongside $Cd^{2+}$ and $Hg^{2+}$. The high throughput design enables simultaneous testing of many

samples for a qualitative metal detection, while absorbance measurement from the colorimetric

Alamar Blue Assay ensures metal quantification extrapolated to a standard plot. Thus, this work

opens new avenues for developing a reliable and sensitive biosensor platform using whole-cell

mycobacteria as biological indicator for toxic metal ions as well as trace metal elements (TME) in

the environmental samples. The efficacy of this system lies in its user-friendly design, specificity,

and sensitivity, enabling multiplexed metal detection by non-genetically modified whole cell

microbes. Consequently, the working principle can be easily optimised and employed in standard

biological laboratories equipped with organisms harboring other metal-sensitive intein-carrying

precursor proteins.

**Experimental Procedure:**

*Experimental Design and Statistical Rationale*

Current study was designed to evaluate intein splicing-based whole-cell mycobacterial biosensor,

where splicing inhibition in the metal-sensing SufB precursor protein, reduced mycobacterial

viability. During *in vitro* refolding assay, purified denatured *Mtb* SufB precursor and *Mtb* SufB

splicing inactive (SI) double mutant (Cys1Ala/Asn359Ala) (splicing negative control) proteins

were renatured in the absence and presence of $CdCl_2$, $HgCl_2$, $CrCl_3$, and $PbCl_2$ over the test

concentration range (2.5 µM–2 mM). All the experiments were performed in triplicates (n=3) and

the protein products from the Coomassie-stained SDS–PAGE gels were quantified by

densitometric analysis using GelQuant.NET biochemical solutions software. The splicing and

cleavage efficiencies were calculated and expressed as percentage values after baseline correction,

using 0 hour splicing and cleavage value(s).

For evaluation of the metal biosensor, a simple Alamar Blue assay was conducted, where different

mycobacterial cells [*Mtb* H37Ra cells and *Mycobacterium smegmatis* (SufB intein-less)] were

incubated with varied concentrations of $Cd^{2+}$ (2.5 µM- 2 mM) and $Hg^{2+}$ (25 µM- 2 mM) in 96-

well microtiter plate. In a parallel experimental setup, $Cd^{2+}$ and $Hg^{2+}$ specific metal chelators were

added in a ratio of 1:1 (metal: chelator) to examine the specificity of the biosensor platform. For a

quantitative evaluation, standard plots were generated by correlating the known concentrations of

samples [$Cd^{2+}$ and $Hg^{2+}$] along the X-axis, to the respective absorbance values from the Alamar

Blue assay on the Y-axis. Test metal concentrations in the unknown samples were interpolated

from the standard curve to validate the biosensor efficacy. Multiplexing ability of the whole -cell

biosensor was assessed using known splicing inhibitors $ZnCl_2$, $CuCl_2$, and $PtCl_4$, along with $CdCl_2$

and $HgCl_2$, under identical experimental conditions.

Statistical analysis for refolding assays was performed using one-way analysis of variance

(ANOVA) to compare splicing and N-terminal cleavage efficiencies of metal treated samples

against the untreated protein control, after baseline correction using 0 h values, and results were

plotted using GraphPad Prism version 5.01 for Windows, GraphPad Software, SanDiego, CA,

U.S.A. (www.graphpad.com). The data are presented as mean ± SEM. The statistical significance

(p-value) between the control and experimental groups is denoted by * ($p < 0.05$), or ** ($p < 0.01$),

or *** ($p < 0.001$), or ****($p<0.0001$).

*Protein overexpression and purification* (62–64)

Plasmids containing the genes for *Mtb* SufB precursor and *Mtb* SufB splicing inactive (SI) double

mutant (Cys1Ala/Asn359Ala) (splicing negative control), both carrying an N-terminal 6X (His)

tag, were borrowed from Belfort Lab, SUNY, Albany, USA)(62). The test plasmids were

transformed into BL21(DE3) *E. coli* cells (NEB, C2527I), followed by IPTG (500 µM, sigma 367-

93-1) induced protein over-expression. Thereafter, lysis buffer (20 mM sodium phosphate, 0.5 M

NaCl, pH 7.4) was used to resuspend the bacterial cell pellet and cells were lysed via tip sonicator

(Sonics vibra cell VCX-130). The cell lysate was centrifuged at 16500×g for 20 minutes at 4 °C,

and the supernatant was discarded. The collected inclusion body (IB) material was then washed

three times in lysis buffer via centrifugation (16500×g for 20 minutes at 4 °C). Finally, the IB

material was solubilized for protein extraction using 8 M urea (Merck, 1084870500) buffer (20

mM sodium phosphate, 0.5 M NaCl, 8M urea, 10 mM of imidazole, pH 7.4 (MP–biochemicals-

288-32-4) and subsequently centrifuged at 16,500×g for 20 minutes at 4 °C to collect the

supernatant. Then, solubilized proteins were purified by Ni-NTA affinity column (Ni-NTA His

trap, HP GE healthcare life sciences, 17524802) (38, 45, 58, 65, 66). Before applying the samples,

the columns were equilibrated with binding buffer (20 mM sodium phosphate, 0.5 M NaCl, 8 M

urea, 10 mM imidazole, pH 7.4). Following sample loading, the columns were washed multiple

times (5 CV) using wash buffer (20 mM sodium phosphate, 0.5 M NaCl, 8 M urea, 40 mM

imidazole, pH 7.4). Subsequently, the test proteins were eluted as purified fractions using elution

buffer (20 mM sodium phosphate, 0.5 M NaCl, 8 M urea, 500 mM imidazole, pH 7.4). Protein

quantification was performed using Bradford reagent.

*Refolding assay for Mtb SufB precursor protein*

Denatured protein (2.5 μM) in elution buffer was renatured in refolding buffer (20 mM sodium

phosphate, 0.5 M NaCl, 0.5 M L-Arginine) containing 2 mM TCEP-HCl (sigma-51805-45-9) for

4 hours at 25 ºC in the absence and presence of toxic metal ions (45, 65, 66). Gradient assay for

toxic metal ions [$CdCl_2$ (SRL-99643), $HgCl_2$ (SRL-25699), $CrCl_3$ (CDH-025702), $PbCl_2$ (SRL-

69789)]  was performed using varying concentrations of toxic metal ions (2.5 μM to 2 mM). 0h

sample was collected before initiation of refolding, and the reaction was abruptly blocked by the

addition of loading dye (0.1% bromophenol blue, 50% glycerol, 5% β-mercaptoethanol, 10% SDS,

tris 6.8), followed by rapid freezing at −20 ºC. After 4h of protein renaturation, the protein samples

were resolved through 4~10% gradient SDS-PAGE, followed by staining with Coomassie brilliant

blue R-250 (93473, SRL), and densitometric analysis was performed using GelQuant.NET

biochemical solutions software. The splicing and cleavage efficiencies were calculated using the

following formulae:

231
$$\% \ of \ splicing = \frac{Total \ splicing \ products \ (LE + I)}{Total \ protein \ (P + LE + I)} \ X \ 100$$

232
$$\% \ of \ N-terminal \ cleavage = \frac{Total \ N-terminal \ cleavage \ products \ (NC + NE)}{Total \ protein \ (P + NC + NE)} X \ 100$$

[P: precursor protein, LE: Ligated exteins, I: Intein, NC: N-terminal cleavage product, NE: N-

extein]

Baseline correction was done using the 0 hours splicing and cleavage value(s). The results were

analyzed using one-way ANOVA and plotted using GraphPad Prism version 8 for Windows,

GraphPad Software, San Diego, California, USA, www.graphpad.com.

*Immunoblotting to identify splicing and N-terminal cleavage protein products*(43)

Protein immunodetection was conducted to identify of splicing and N-terminal cleavage reaction

products, using anti-6X(His) tag antibodies (Invitrogen, LOT 1902132). After resolution through

4~10% gradient SDS-PAGE, the protein products were transferred onto a PVDF membrane

(Millipore, IPVH 00010) at 50V for 2 hours at 4 °C. The blot was then blocked with 5% skim milk

for 2 hours at room temperature, followed by two washes with 1X TBST. It was incubated with

primary anti-6X(His) tag (abgenex -32-6116) at a dilution of 1:5000 at 4 °C for 16 hours. After

washing with 1X TBST, the blot was treated with a secondary antibody (anti-mouse; abgenex 11-

301) at a dilution of 1:8000 for 2 hours at 37 °C. The N-extein band was detected using a lower

dilution of the primary antibody (1:2500) and the secondary antibody (1:6000). Finally, the blot

was developed using Enhanced Chemiluminescence substrate (Abcam, ab65628) to visualize the

protein bands, and images were captured using in-house ImageQuant™ LAS 500 facility (GE

Healthcare). Blots were also developed, using 3, 3'-diaminobenzidine (DAB) (SRL-94524) for

visualization of the protein bands.

*Spectroscopic and Elemental Profiling of Protein-Metal Interactions*

*UV-Visible spectroscopy* (67–69)

Denatured proteins were renatured *in vitro* in the presence and absence of toxic metal ions at their

respective concentrations for 4 hours, as specified earlier. Subsequently, the treated samples were

subjected to UV-visible spectroscopy (Cary 100 UV-Vis Agilent Technology) scanning in the

visible region spanning 200-800nm. To rule out the interference from buffer-metal interactions,

the refolding buffer (20 mM sodium phosphate, 0.5 M NaCl, 0.5 M L-Arginine) was incubated

with metal ions ($Cd^{2+}$, $Hg^{2+}$, $Cr^{3+}$, and $Pb^{2+}$) at their respective concentrations under similar

experimental conditions. This was used as a blank for baseline correction and was scanned

alongside each sample. Absorbance versus wavelength plots were generated and analyzed using

Origin (Pro) 8.5 software (Version 8.5, OriginLab Corporation, Northampton, MA, USA) to

interpret the results.

*Tryptophan Fluorescence Spectroscopy*

The purified denatured protein (2.5 µM) was allowed to renature in the presence and absence of

metals ions (25 µM $CdCl_2$, 500 µM $HgCl_2$, 2 mM $CrCl_3$, and 2 mM $PbCl_2$) under identical

experimental conditions. Following 4 hours of refolding, the test samples were analyzed via

Fluorescence spectrometry (Cary Eclipse, Agilent Technologies) over a wavelength range of 300-

600 nm and at an excitation wavelength of 280 nm (70). To rule out the buffer-metal interactions,

the refolding buffer (20 mM sodium phosphate, 0.5 M NaCl, 0.5 M Arginine) was incubated with

metal ions at their respective concentrations at 25 °C for 4 hours and then scanned alongside each

treated sample. This served as the blank for baseline correction. The tryptophan fluorescence

spectra for the test samples were recorded and analyzed by plotting intensity vs. wavelength using

Origin (Pro) 8.5, Version 8.5, from OriginLab Corporation, Northampton, MA, USA.

*Inductively coupled plasma-optical emission spectroscopy (ICP-OES) analysis*

Following *in vitro* refolding of *Mtb* SufB precursor in the presence of test metal ions, the protein

products were resolved through 4~10% SDS-PAGE. The resulting protein bands were visualized

with Coomassie brilliant blue R-250. Subsequently, the gel bands for unspliced precursor protein,

ligated extein, and intein were excised from the gel for further analysis. Each gel band was then

treated with concentrated $H_2SO_4$ and 30% $H_2O_2$. Complete dissolution of each sample was

achieved by incubating the gel pieces in a 3:1 mixture of concentrated HCl and concentrated $HNO_3$

at 80°C for 4 hours (71, 72). The final volume of the digested solution was adjusted to 50 ml with

deionized water and then analyzed by ICP-OES (Perkin-Elner, optima 8300). Prior to analyzing

the test samples, respective toxic metal standards were calibrated in the system, and the

concentration of bound metal ions was quantified from the standard curve.

*H37Ra-based whole-cell toxic metal biosensor*

Different mycobacterial cells [*Mycobacterium tuberculosis* H37Ra strain (ATCC 25177),

*Mycobacterium smegmatis* (negative control for SufB intein-less micro-organism, ATCC

700084)] were cultured in 7H9 (Middlebrook 7H9 broth medium, M198) supplemented with

OADC (Oleic acid, dextrose, and catalase, Himedia-FD018) and allowed to grow separately for 2-

3 weeks at 37 $^0$C. CFU calculations for individual cultures were made by matching the turbidity

of the inoculated mycobacterial culture with the McFarland standard (Himedia- R092) against a

black background (73). Different bacterial cells were diluted in sterile saline to a concentration of

$10^8$ CFU/ml. H37Ra cells were incubated with varied concentrations of $Cd^{2+}$ (2.5 µM- 2 mM) and

$Hg^{2+}$ (25 µM- 2 mM) in 96-well microtiter plate. Since $Cr^{3+}$ and $Pb^{2+}$ over a test concentration

range of 25 µM- 2 mM did not exert any effect on *Mtb* SufB splicing, these metal ions were not

tested via the designed biosensor platform. The plates were sealed and incubated at 37 ºC for 4

days. Then, 10% (v/v) solution of Alamar Blue reagent (SRL-42650) was added to each well,

followed by incubation at 37 ºC for 12 hrs.(74). Untreated mycobacterial cells grown in 7H9

(supplemented with OADC) media were used as a negative control, and Alamar blue reagent alone

in 7H9 (supplemented with OADC) media served as a positive control. SufB intein-less organism,

*Mycobacterium smegmatis,* was also evaluated under similar conditions and treated as an

additional experimental control. After 12 hours of incubation, varied color changes were noted in

different wells, where blue color indicated dead cells with no viable cell growth, and pink color

correlated to metabolically active live cells. In a similar experimental setup, metal chelators

specific for $Cd^{2+}$ (Edetate Calcium Disodium, ECD, Sigma-PHR1510) and $Hg^{2+}$ (meso-2,3-

Dimercaptosuccinic acid, DMSA, Sigma-D7881) were added in a ratio of 1:1 (metal: chelator) to

determine the specificity of the designed Biosensor platform.

<u>Quantitative evaluation of *H37Ra-based whole cell toxic metal biosensor*</u>

Next,  a standard plot was generated by measuring the absorbance of the respective samples  at

570 nm, with 600 nm as a reference wavelength for quantitative analysis (74). A wide range of

test metal samples ($Cd^{2+}$, $Hg^{2+}$) of known concentration were prepared and loaded onto the 96-

well plate and Alamar Blue reagent was added under similar experimental conditions. The

absorbance of test metal samples was calculated by subtracting the reference absorbance value at

600nm from the absorbance measured at 570nm [$OD_{test} = OD_{570} - OD_{600}$]. Untreated mycobacterial

cells grown in 7H9 (supplemented with OADC) media were used as a control for normalization.

Using the normalized values, XY scattered plot was obtained where the X axis represented the

known concentrations of the test metal ions and the Y axis denoted corresponding absorbance

values. Next, to assess the applicability of the standard plot, four test samples (Unknown 1,

Unknown 2, Unknown 3, and Unknown 4) containing varying concentrations of $CdCl_2$ and $HgCl_2$

were analyzed under identical experimental conditions. Absorbance values were extrapolated onto

the standard curve to calculate the unknown metal ion concentrations, thereby validating the

sensitivity and accuracy of the biosensor platform.

*Multiplexing evaluation of H37Ra-based whole cell metal biosensor*

Bacterial cell suspension containing $10^8$ CFU/ml of *Mycobacterium tuberculosis* H37Ra were

prepared in sterile saline. Cells were incubated with a varied concentration range of $ZnCl_2$ (100

μM- 2 mM), $CuCl_2$ (100 μM- 2 mM), $PtCl_4$ (20 μM- 1 mM), $CdCl_2$ (10 μM- 2 mM), and $HgCl_2$

(25 μM- 2 mM), in  96-well microtiter plates. The plates were sealed and incubated at 37 ºC for 4

days under static conditions. Then, 10% (v/v) solution of Alamar Blue reagent (SRL-42650) was

added to each well, followed by incubation at 37 ºC for 12 hrs. Untreated H37Ra cells grown in

Middlebrook 7H9 supplemented with OADC media were used as a negative control, while Alamar

blue reagent incubated in media alone served as a positive control. SufB intein-less organism,

*Mycobacterium smegmatis,* was treated with test metal ions under similar conditions and used as

negative experimental control to assess SufB intein-specific response. Post 12 hours of incubation,

color change was noted in the wells, where blue color indicated dead cells with no viability, and

pink color indicated healthy viable cells.

To further ascertain the specificity of the biosensor platform, respective metal-chelating agents

were added in separate experimental setup, using a 1:1 molar ratio of metal ions to chelator.

Cadmium-specific chelator Edetate Calcium Disodium (ECD; Sigma-PHR1510), mercury-

specific chelator meso-2,3-Dimercaptosuccinic acid (DMSA; Sigma-D7881), and a broad-

spectrum chelator Ethylenediaminetetraacetic acid (EDTA; HiMedia-MB011), were co-incubated

with the respective metal ions under identical experimental conditions. Cell viability was evaluated

using Alamar Blue reagent as described earlier.

*Statistical Analysis*

Statistical analysis for refolding assays was performed using one-way analysis of variance

(ANOVA) to compare splicing and N-terminal cleavage efficiencies of metal treated samples

against the untreated protein control, after baseline correction using 0 h values, and results were

plotted using GraphPad Prism version 5.01 for Windows, GraphPad Software, SanDiego, CA,

U.S.A. (www.graphpad.com). The data are presented as mean $\pm$ SEM. The statistical significance

(p-value) between the control and experimental groups is denoted by * ($p < 0.05$), or ** ($p < 0.01$),

or *** ($p < 0.001$), or ****($p<0.0001$).

**Results and Discussion**

***Toxic metal ions inhibited Mtb SufB precursor protein splicing***

Environmental concentrations of toxic metal ions such as cadmium ($Cd^{2+}$), mercury ($Hg^{2+}$),

chromium ($Cr^{3+}$) and lead ($Pb^{2+}$) have been reported to vary widely. $Cd^{2+}$ concentrations typically

ranges   from 1 to 10 ppm (approximately 5 $\mu$M to 55 $\mu$M), $Hg^{2+}$ from 10 to 700 ppm

(approximately 4 $\mu$M to 3 mM), $Cr^{3+}$ from 5 ppm- 140 ppm (approximately 31 $\mu$M – 885 $\mu$M) and

$Pb^{2+}$ levels from 10 to 50 ppm (approximately 36 $\mu$M to 180 $\mu$M) [75–90]. Earlier studies have

shown that metal ion concentrations between 40 $\mu$M to 2 mM effectively inhibit intein splicing

[43, 91, 92]. Built upon these observation, *in vitro* gradient assay was performed to evaluate the

sensitivity of *Mtb* SufB precursor protein to a selected concentration range of toxic metals ions

(2.5 $\mu$M to 2 mM) (Fig. 1, and Supplemental Fig. S2-S4). *Mtb* SufB precursor protein (95.98 KDa)

follows a canonical intein splicing pathway yielding 40.2 KDa  SufB intein and 55.7 KDa ligated

exteins (functional SufB protein) as illustrated in Supplemental Fig. S1(62). N-terminal cleavage

products (65.26 KDa N-terminal cleavage product, 29.9 KDa N-extein) and C-terminal cleavage

products (70.12 KDa C-terminal cleavage product, 25.76 KDa C-extein) are the byproducts of *Mtb*

SufB splicing reaction (62).

Purified *Mtb* SufB precursor protein was refolded in presence of test metal ions and then resolved

through 4~10% gradient SDS PAGE (Fig. 1A). $Cd^{2+}$ at 25 µM, and $Hg^{2+}$ at 500 µM of minimum

effective concentration (MEC) displayed significant attenuation of splicing and N-terminal

cleavage reactions. There was a noticeable accumulation of SufB precursor protein, along with

substantial reduction in the splicing products (ligated exteins and intein). Further, $Cd^{2+}$, and $Hg^{2+}$

at their respective MEC values and higher, induced a near-complete inhibition of N-terminal

cleavage reaction and the products (N-terminal cleavage product and N-extein) (Fig. 1A).

However, $Cr^{3+}$ and $Pb^{2+}$ failed to induce any effect on *Mtb* SufB splicing or cleavage reactions

across the test concentration range (Fig. 1A). Resultant protein products were further confirmed

via western blot using anti-6X(His) tag antibody (Supplemental Fig. S2). Densitometric analysis

revealed a 12.52-fold (p<0.0001) reduction in splicing efficiency and an 18.7-fold (p<0.0001)

reduction in N-terminal cleavage efficiency with 25 µM $Cd^{2+}$ [Fig. 1B (i) and C (i)]. Similarly,

treatment with 500 µM $Hg^{2+}$ resulted in an 8-fold (p < 0.0001) reduction in splicing and a 7.8-fold

(p < 0.0001) decrease in N-terminal cleavage efficiency [Fig. 1B (ii) and 1C (ii)]. No significant

alterations in SufB splicing or N-terminal cleavage products were obtained in presence of $Cr^{3+}$ or

$Pb^{2+}$ (Fig. 1B and 1C). All the test metal ions at a lower concentration range (2.5 µM- 10 µM)

failed to exert any significant effect on the SufB splicing reactions (Supplemental Fig. S3 and S4).

In order to rule out the possibility of precursor protein degradation by test metal ions, splicing

inactive (SI) *Mtb* SufB double mutant (C1A/N359A) was refolded under similar experimental

conditions (25 µM Cd²⁺, 500 µM Hg²⁺, 2 mM Cr³⁺, and 2 mM Pb²⁺). Since Cr³⁺, and Pb²⁺ showed

no regulatory effects, highest test concentration range for both were included in the above study.

SDS-PAGE and subsequent western blot analysis using anti-6X(His) tag antibodies confirmed the

absence of protein degradation products (Supplemental Fig. S2B).

**Fig. 1. Gradient assay to determine the minimum effective concentration (MEC) of toxic metal ions on splicing and N-terminal cleavage reactions of *Mtb* SufB precursor protein. (A)** Following *in vitro* refolding of *Mtb* SufB precursor protein in the presence of (i) CdCl₂, (ii) HgCl₂, (iii) CrCl₃, and (iv) PbCl₂ at varied concentrations (25 µM-2 mM), resultant products were resolved through 4~10% gradient SDS-PAGE. Lane 1: Protein ladder; Lane 2 (T₀): splicing and N-terminal cleavage reactions at 0 h; Lane 3 (UT): splicing and N-terminal cleavage products of untreated protein sample; Lanes 4-9: splicing and N-terminal cleavage products resulting in the presence of different toxic metal ions at different concentrations (25 µM-2 mM). Lane 10 (SI): splicing inactive SufB double mutant (C1A/N359A) is used as a negative control for splicing. **(B) and (C)**

Bar graphs demonstrating splicing and N-terminal cleavage efficiency of *Mtb* SufB precursor
protein in the presence and absence of different toxic metal ions. A statistically significant
difference in splicing and N-terminal cleavage reaction was observed for B (i) and C (i) $CdCl_2$,
starting with 25 µM, and B (ii) and C (ii) $HgCl_2$ at 500 µM concentrations, respectively. For $CrCl_3$
[B (iii) and C (iii)], and $PbCl_2$ [B (iv) and C (iv)], no significant difference in splicing and N-
terminal cleavage reaction were observed between untreated and metal-treated samples. Graphs
were plotted after densitometric analyses using GelQuant.Net biochemical solutions. All the
experiments were performed in triplicate, and error bars represent (±1) SEM from 3 independent
sets of experiments. The data shown in Fig. 1B, and 1C are extracted from Fig. 1A. M: Prestained
protein molecular weight ladder, P: Precursor, NC: N-terminal cleavage product, LE: Ligated
extein, I: intein, NE: N-extein.

Further, a time-dependent splicing assay was performed to examine *Mtb* SufB splicing and

cleavage activity over a period of 10 min- 4 hr., elucidating remarkable inhibitory roles of $Cd^{2+}$

(25 µM) and $Hg^{2+}$ (500 µM) as early as 10 minutes (Supplemental Fig. S5). Next, to evaluate the

possible physical interaction between *Mtb* SufB protein and the metal ions, a series of experiments

were conducted, as shown in the upcoming section. For the said interaction study, the MEC for

$Cd^{2+}$, $Hg^{2+}$ (25 µM, 500 µM, respectively), which inhibited *Mtb* SufB splicing was selected. Since,

$Cr^{3+}$, and $Pb^{2+}$ did not exhibit any such regulatory activity, highest value of the test concentration

range (2 mM) was chosen to examine possible interaction with SufB protein.

**Protein~ metal interaction study**

*UV-Visible spectroscopic analysis*

Biomolecules such as proteins exhibit a strong affinity for metal ions, which can critically

influence their structure and function (93). Prior studies have demonstrated that toxic metal ions

such as $Cd^{2+}$, $Hg^{2+}$, $Cr^{3+}$, and $Pb^{2+}$ can interact with calcium and zinc-binding sites in proteins,

thereby influencing their biological activity (94, 95). UV-visible spectroscopic analysis was

conducted to investigate the interaction between the *Mtb* SufB precursor protein and toxic metal

ions under treated and untreated conditions. Absorbance spectra for *Mtb* SufB protein were

recorded over 200-800 nm range at 25 °C (Supplemental Fig. S6). A characteristic absorbance

peak near 280 nm was detected, primarily attributable to aromatic residues such as tryptophan and

tyrosine (96, 97). Upon treatment with $Cd^{2+}$, an increase in absorbance intensity was noted,

suggesting an alteration in conformation due to metal coordination [Supplemental Fig. S6 (i)].

Previous studies have mentioned that enhanced absorbance may result from the interaction

between $Cd^{2+}$ and electron-donating groups such as amide, imidazole, and amine (68, 98, 99).

Likewise, samples treated with $Hg^{2+}$ also exhibited an increase in absorbance intensity

[Supplemental Fig. S6 (ii)]. This enhancement is likely attributed to the strong affinity of $Hg^{2+}$ for

nucleophilic functional groups, particularly the sulfhydryl (–SH) groups of cysteine residues (100,

101). Although there were sharp increments in absorbance peak following interaction of $Cr^{3+}$ and

$Pb^{2+}$ with *Mtb* SufB, they failed to exert any effects on SufB splicing reactions as shown earlier

(Fig. 1). Moreover, a strong buffer-$Pb^{2+}$ interaction was observed from the results, which may have

weakened likely interaction with the protein. Although prior works have shown interaction of $Cr^{3+}$,

and $Pb^{2+}$ with thiol, glutathione groups within protein structure including oxygen, nitrogen, and

sulfur containing amino acids (7, 102, 103), current work could not detect any effects of these

metal ions on *Mtb* SufB splicing activity [Supplemental Fig. S6 (iii and iv)]. To further substantiate

such interactions, tryptophan fluorescence spectroscopy was performed, as explained in the

forthcoming section.

***Tryptophan Fluorescence spectroscopy reveals metal-induced conformational changes in Mtb***

***SufB***

Tryptophan fluorescence spectroscopy is a widely utilized technique for monitoring alterations in

protein conformation and assessing local structural dynamics upon interaction with ligands or

macromolecules (104). Among the intrinsic fluorophores in proteins, aromatic amino acids—

tryptophan (Trp), tyrosine (Tyr), and phenylalanine (Phe)—contribute to fluorescence, with

tryptophan exhibiting the highest molar absorptivity (1614 cm$^{-1}$/M) and a characteristic emission

peak in the range of 340–350 nm (105). The indole ring of Trp is primarily responsible for this

emission and dominates the fluorescence profile over Tyr and Phe. Changes in the

microenvironment of tryptophan residues, such as those caused by ligand binding, protein folding,

or metal interaction, can lead to measurable shifts in fluorescence intensity and/or emission

maxima (104, 106).

Altogether, 13 Trp residues are present, spanning the entire sequence of *Mtb* SufB precursor

protein, with 5 Trp in N-extein, 5 Trp in Intein, and 3 Trp in the C-extein region (43). Upon

treatment with $Cd^{2+}$ (25 µM), an increase in fluorescence intensity was observed [Supplemental

Fig. S7 (i)], likely attributable to conformational rearrangements arising from interactions between

$Cd^{2+}$ and functional groups such as amide, imidazole, and amine moieties within the protein matrix

(68, 98, 99). These interactions may have led to reduced solvent exposure of Trp residues or

stabilized a more rigid protein conformation, which enhanced fluorescence emission (68, 107). In

contrast, $Hg^{2+}$ (500 µM) treated samples displayed a significant quenching of Trp fluorescence

[Supplemental Fig. S7 (ii)]. This effect is consistent with previous studies indicating that mercury

ions interact strongly with aromatic systems through cation–π interactions, particularly with the

indole ring of tryptophan, thereby reducing fluorescence emission (108). $Cr^{3+}$ (2mM) exposure

resulted in a slight increase in fluorescence intensity [Supplemental Fig. S7 (iii)], potentially due

to the preferential binding of $Cr^{3+}$ to carboxyl groups, which may induce localized structural

stabilization without extensive perturbation of the overall tertiary structure (109, 110). In case of

$Pb^{2+}$ (2 mM), a decrease in fluorescence intensity was observed [Supplemental Fig. S7 (iv)]. This

observation is supported by previous literature, where $Pb^{2+}$ interacts with hydroxy, amino, and

sulfhydryl groups, leading to partial protein unfolding and increased solvent accessibility of Trp

residues, thereby promoting fluorescence quenching (111, 112). Although $Cr^{3+}$, $Pb^{2+}$ have

displayed possible alteration in protein conformation, they failed to regulate *Mtb* SufB splicing

and cleavage reactions, likely due to lack of any effect on protein active site. Next, ICP-OES

analysis was conducted to further ascertain potential interaction between test metal ions and

protein backbone, as described in the following section.

***ICP-OES detects metal ions bound to SufB precursor and spliced out protein fractions***

ICP-OES is a sensitive analytical technique capable of detecting trace metal concentrations by

measuring the characteristic emission of photons from the excited atoms and ions as they return to

their ground state (113). During *in vitro* protein refolding experiment, specific concentrations of

$Cd^{2+}$ (25 µM = 4.583 µg/ml), $Cd^{2+}$ (500 µM = 91.66 µg/ml), $Hg^{+2}$ (500 µM = 135.76 µg/ml), $Cr^{+3}$

(2 mM = 361.72 µg/ml), and $Pb^{+2}$ (2 mM = 556.2 µg/ml) were added to the respective reaction

mixture. ICP-OES analysis of the metal-bound protein fractions revealed detectable levels of $Cd^{2+}$

(2 ppb ~ 0.002 µg/ml), $Hg^{2+}$ (5 ppb, ~0.005 µg/ml), and $Cr^{3+}$ (80 ppb, ~0.08 µg/ml) bound to the

unspliced *Mtb* SufB precursor protein (Table 2). These findings indicate a direct and quantifiable

physical interaction between the test metal ions and *Mtb* SufB precursor protein, which may

underlie the observed inhibition of splicing and N-terminal cleavage reactions. Further, measurable

interactions were observed between the test metal ions and various splicing products (ligated extein

and intein) (Table 2). $Cd^{2+}$ addition to SufB protein sample at 25 µM concentration yielded

negative results, while increasing the concentration to 500 µM led to positive data. At lower

concentrations of $Cd^{2+}$ (25 µM), the metal content in the protein fraction was possibly below the

instrument's limit of detection (LOD) and limit of quantification (LOQ), resulting in negative

readings. Although ICP-OES is capable of detecting metal concentrations at the ppb level, practical

detection limits are influenced by matrix complexity and background interference, often resulting

in quantification thresholds closer to the ppm range (114–116). Although 80 ppb (0.08 µg/ml) of

$Cr^{3+}$ was bound to the unspliced precursor (Table 2), it failed to exert any visible effects on *Mtb*

SufB splicing reactions (Fig. 1). No bound form of metal ions were detected in $Pb^{2+}$ treated

samples, even at 2 mM concentration (Table 2), consistent with no visible effect on *Mtb* SufB

splicing (Fig. 1).

**Table 2:** Quantitative elemental detection via ICP-OES analysis: Measurable concentration of
metal ions ($Cd^{2+}$, $Hg^{2+}$, $Cr^{3+,}$ and $Pb^{2+}$) were identified bound to *Mtb* SufB precursor and splicing
products. P: Precursor, LE: Ligated extein, I: Intein.

| Metal ions | Treated concentration | Concentration (µg/ml) | ICP-OES values (ppb) | Equivalent Concentration (µg/ml) |
|---|---|---|---|---|
| Cadmium ($Cd^{2+}$) | 25µM | 4.583 | P: -9<br>LE: -5<br>I: -8 | P: -0.009<br>LE: -0.005<br>I: -0.008 |
| Cadmium ($Cd^{2+}$) | 500 µM | 91.66 | P: 2<br>LE: 18<br>I: 1 | P: 0.002<br>LE: 0.018<br>I: 0.001 |
| Mercury ($Hg^{2+}$) | 500 µM | 135.76 | P: 5<br>LE: 3<br>I: 5 | P: 0.005<br>LE: 0.003<br>I: 0.005 |
| Chromium ($Cr^{3+}$) | 2mM | 361.72 | P: 80<br>LE: 83<br>I: 90 | P: 0.08<br>LE: 0.083<br>I: 0.09 |
| Lead ($Pb^{2+}$) | 2mM | 556.2 | P: -17<br>LE: -5<br>I: -73 | P: -0.017<br>LE: -0.005<br>I: -0.073 |

Collectively, the protein refolding assays, along with protein~metal interaction studies, suggested

possible interaction of  $Cd^{2+}$ and $Hg^{2+}$ with *Mtb* SufB precursor protein, thereby attenuating the

splicing and N-terminal cleavage efficacy. These mechanistic insights led to the development of a

whole-cell biosensing platform utilizing  H37Ra (*Mtb* attenuated strain) as the indicator cells to

detect  metal ions.

Fe-S clusters are essential cofactors for distinct metabolic pathways and cellular processes such as

respiration, DNA replication and repair, and persistence under oxidative and nitrosative stress

conditions (117–119). SUF complex offers the sole pathway for Fe-S cluster biogenesis in

mycobacteria, and its activity relies on the generation of functional SufB protein via protein

splicing (60, 61). Besides, there are earlier reports on loss of cellular viability as a result of SufB

splicing inhibition within mycobacteria (40, 43, 56). Since, toxic metal ions like of  $Cd^{2+}$ and $Hg^{2+}$

were shown to inhibit *Mtb* SufB splicing and N-terminal cleavage activity (Fig. 1), an attenuated

strain of *Mtb* (H37Ra) was used as the whole-cell viability indicator for these metal ions.

**H37Ra- based whole-cell biosensor detects toxic metal ions**

Cell-based biosensors are gaining interest as a screening platform for biologically active molecules

due to their ability to monitor the cellular functions and interactions with targets in their natural

environment (120, 121). Often, they are fabricated as genetically modified biosensors using native

receptors or enzymes as molecular recognition elements in living cells (28). Current work explores

the use of whole-cell mycobacteria (H37Ra)-based biosensor to detect toxic metal ions like $Cd^{2+}$

and $Hg^{2+}$. A 96-well plate assay was developed using Alamar Blue reagent, employing *Mtb*

(H37Ra) cells as the sensing component, and *M. sm* cells as the SufB intein-less cellular control

(56, 62, 122). *M. sm* carries three intein-bearing proteins: DnaB, which contains two inteins

(DnaBi1 and DnaBi2), as well as GyrA and PhoH (49, 64). Although splicing regulation has been

shown for DnaBi1 helicase, studies on the DnaBi2, PhoH, GyrA inteins splicing have not been

reported yet. DnaBi1 splicing is regulated by zinc and oxidative stressors (49, 64); however, its

helicase function can be compensated by other cellular helicases (123–125).

Alamar Blue reagent is a non-toxic, resazurin-based solution that functions as an indicator for cell

viability (74, 126). The blue-colored non-fluorescent resazurin (7-hydroxy-3H-phenoxazin-3-one

10-oxide) gets reduced to resorufin (7-hydroxy-3H-phenoxazin-3-one, pink colored compound)

by mitochondrial reductase in metabolically active cells (127). In a comparative study, non-viable

H37Ra were visualized in presence of $Cd^{2+}$ and $Hg^{2+}$, whereas persistence of viable *M. sm* cells

during the metal-interaction process highlighted the absence of metal sensing SufB precursor

protein. Untreated H37Ra cells grown in 7H9 (supplemented with OADC) media, and Alamar

Blue reagent alone in 7H9 (supplemented with OADC) media, served as the negative and positive

controls for the study respectively.

Fig. 2 (i) is a schematic representation of 96-well plate-based biosensor platform containing *Mtb*

H37Ra cells (indicator organisms) incubated with varied concentrations of $Cd^{2+}$ (2.5 μM -2 mM)

and $Hg^{2+}$ (25 μM -2 mM). The test concentration range for the metal ions were chosen based on

the positive results obtained during *in vitro* splicing analyses (Fig. 1, Supplemental Fig. S2-S4).

For a baseline correction, H37Ra cells were grown in the absence of toxic metal ions (negative

control). SufB intein-less, *M. sm* cells were used as additional experimental control. Unlike *Mtb*

H37Ra cells, the growth and viability of the *M. sm* cells are independent of SufB intein splicing,

and any loss of cell viability indicates metal toxicity. Fig. 2 (ii) shows that the indicator cells

(H37Ra) incubated in the presence of lower concentration of $Cd^{2+}$ (2.5-10 μM) did not lose cell

viability. All the cells were metabolically active and converted resazurin (blue) to resorufin (pink),

since $Cd^{2+}$ at these concentration (2.5-10 µM) didn't affect SufB splicing activity, generating

active SufB protein  [Supplemental Fig. S3 (i)]. H37Ra cells incubated in presence of 25 and 50

µM of $Cd^{2+}$, displayed purple color indicating a partial loss of cell viability, underscoring 25 µM

as the minimum inhibitory concentration (MIC). Cells incubated with $Cd^{2+}$ at a concentration range

of 100 µM- 2 mM, retained  blue color, indicating complete loss of cell viability. These results

correlated to inhibition of splicing and N-terminal activity of *Mtb* SufB in presence of $Cd^{2+}$ over a

concentration of 25 µM - 2 mM (Fig. 1).

Next, cells incubated in the presence of 25 µM and 50 µM of $Hg^{2+}$ resulted in healthy viable (pink)

cell population, whereas cells in the presence of 100 µM $Hg^{2+}$ exhibited purple color, highlighting

partial loss of cell viability at MIC value. Over rest of the concentration range for $Hg^{2+}$ (500 µM-

2 mM),  non-viable (blue) cells were noticed. These results were supported by SufB splicing

attenuation by $Hg^{2+}$ (500 µM-2 mM), while a lower concentration (2.5 µM-10 µM) $Hg^{2+}$  failed to

exert any such effects (Fig. 1, Supplemental Fig. S4). Although 100 µM of $Hg^{2+}$ didn't influence

SufB splicing, the N-terminal cleavage reaction was reduced significantly, marking this as the MIC

value with partial loss of cell viability. Summarizing, persistence of blue color in the presence of

$Cd^{2+}$ (100 µM until 2 mM), and $Hg^{2+}$ (500 µM until 2 mM) may be correlated to complete loss of

H37Ra viability due to inhibition of *Mtb* SufB splicing and N-terminal cleavage reactions (Fig. 1).

These results suggest the potential use of mycobacterial (H37Ra) cells as the biological indicator

for a qualitative toxic metal detection.

**Fig. 2. 96-well plate-based biosensor assay demonstrating whole cell mycobacteria (*Mtb* H37Ra strain) as the biological indicator for toxic metal ions (Cd²⁺ and Hg²⁺)** (i) Schematic representation of 96 well plate-based Biosensor platform. (ii) H37Ra cells (10⁸ CFU/ml) were seeded in the presence of varied concentration ranges of CdCl₂ (2.5 µM- 2 mM) and HgCl₂ (25 µM- 2 mM), respectively. Cells incubated in the presence of CdCl₂ (2.5 µM- 10 µM) and HgCl₂ (25 µM- 50 µM) were pink in color indicating healthy viable cells resisting the splicing inhibitory effect of the metal ions. In contrast, cells exposed to CdCl₂ (25 µM and 50 µM) and 100 µM of HgCl₂ displayed purple color suggesting partial loss of cell viability. Cells incubated with 100 µM- 2 mM of CdCl₂ and 100 µM- 2 mM of HgCl₂ remained blue, suggesting complete loss of viable cell population. For a baseline correction, test organisms were allowed to grow in the absence of toxic metal ions and denoted as live cell control (negative control). SufB intein-less organism *Mycobacterium smegmatis (M. sm)* was used as additional experimental control and treated under similar conditions. *M. sm* cells exhibited pink color across a varied range of toxic metal ions, suggesting their cell viability as an intein-independent phenomenon. The experiments were performed in triplicates to confirm the reproducibility of data. Cyan stars- complete loss of viability (blue) & Yellow stars- partial loss of viability (purple).

Next, a quantitative analysis for the toxic metal ions was performed by measuring the absorbance

of Alamar Blue reagent at 570 nm with 600 nm as a reference wavelength (74). Metal samples of

known concentrations were prepared and loaded on to the 96-well plate and treated with Alamar

Blue reagent under optimised experimental conditions [Fig. 3 (i)-(iv)].  A standard graph was

generated by plotting the absorbance values of Alamar reagent vs the known concentrations of test

metal ions ($Cd^{2+}$ and $Hg^{2+}$) [Fig. 3 (v) and (vi)]. Subsequently, a linear regression analysis was

performed to determine the equation of the line, expressed as $y = mx + c$. To further validate the

standard plot, unknown samples with varying concentrations of $Cd^{2+}$ ions and $Hg^{2+}$ ions were

prepared (Experimental Procedure). Absorbance values of the unknown samples were extrapolated

onto standard graph to determine the concentrations of $Cd^{2+}$ and $Hg^{2+}$ (Fig. 3 (v), (vi) and Table

S1). Unknown samples with concentration higher than the standard range may be adjusted via

dilution, or the standard calibration plot may be extended to higher concentration ranges (128).

However, sample concentration at a value lower than the standard range may not be determined

accurately possibly due to no effect on intein splicing and hence no loss of cell viability.

**Fig. 3. Standard curve for quantitative determination of $Cd^{2+}$ and $Hg^{2+}$ concentrations.** (i),
and (iii) Schematic representations of H37Ra-based Biosensor platform to detect $CdCl_2$ and $HgCl_2$.
(ii), and (iv) *Mtb* H37Ra cells were grown in presence of varied concentration of $CdCl_2$ and $HgCl_2$.
To establish a baseline, test organisms were allowed to grow in absence of toxic metal ions (live
cell or negative control). (v), and (vi) The absorbance values of the Alamar Reagent solutions were
plotted along the y-axis and the known concentrations of $Cd^{2+}$ and $Hg^{2+}$ were plotted along the x-
axis. A linear regression analysis was conducted to obtain the equation for the line, and used to
determine the concentration of unknown samples. The experiments were performed in triplicates

to confirm the reproducibility of data.

Next, to elucidate the specificity of the designed biosensor platform, $Cd^{2+}$-  and $Hg^{2+}$- specific

metal chelators, Edetate Calcium Disodium (ECD), and meso-2,3-Dimercaptosuccinic acid

(DMSA) were incubated alongside the test samples (129–134). Fig. 4 (i) is a schematic

representation of the biosensor platform, where indicator mycobacterial cells were exposed to $Cd^{2+}$

and $Hg^{2+}$, in presence of $Cd^{2+}$-specific chelator, ECD. Fig. 4 (ii) demonstrates that ECD effectively

resisted $Cd^{2+}$- induced loss of viability, resulting in healthy cells that transformed resazurin (blue)

to resorufin (pink). Conversely, ECD failed to restore viability of $Hg^{2+}$ treated cells, where

inhibition of SufB splicing, rendered non-viable (blue) cells. Next, mycobacterial cells were

subjected to varying concentrations of $Cd^{2+}$ and $Hg^{2+}$, supplemented with metal chelator DMSA

($Hg^{2+}$ specific) [Fig. 4 (iii) and (iv)].  $Hg^{2+}$ treated H37Ra cells remained viable in presence of

DMSA,  while $Cd^{2+}$ continued its inhibitory effect on SufB splicing inducing non-viable (blue)

cells. These observations affirmed the  specificity of the designed biosensor platform to detect

toxic metal ions $Cd^{2+}$ and $Hg^{2+}$.

**Fig. 4. Determination of the specificity of *Mtb* H37Ra-based biosensor platform using metal**
**chelators.** (i) Schematic representation of the biosensor platform including $Cd^{2+}$ specific metal
chelator (ECD) (ii) *Mtb* H37Ra cells ($10^8$ CFU/ml) were allowed to grow in presence of varied
concentration range of $CdCl_2$ and $HgCl_2$, along with ECD (1:1 ratio). ECD reversed the inhibitory
effect of $Cd^{2+}$ inducing viable (pink) cells, whereas the effect of $Hg^{2+}$ persisted (iii) Schematic
representation of the biosensor platform including $Hg^{2+}$ specific metal chelator (DMSA). (iv) *Mtb*
H37Ra cells ($10^8$ CFU/ml) were allowed to grow in the presence of varied concentration ranges of
$CdCl_2$ and $HgCl_2$, including DMSA in the ratio 1:1. DMSA blocked the inhibitory effect of $Hg^2$,
rendering pink viable cells, whereas the effect of $Cd^{2+}$ persisted. For a baseline correction, H37Ra
cells were grown in the absence of toxic metal ions as the live cell control. SufB intein-less
organism *Mycobacterium smegmatis (M. sm)* was used as additional experimental control and
treated under similar conditions. *M. sm* cells exhibited pink color across a varied range of toxic
metal ions, suggesting their cell viability as an intein-independent phenomenon. The experiments
were performed in triplicates to confirm the reproducibility of data. Cyan stars- complete loss of
viability (blue) & Yellow stars- partial loss of viable cells (purple). ECD: Edetate Calcium
disodium, DMSA: Meso-2,3-Dimercaptosuccinic acid.

Next for a  multiplex metal screening process, the efficacy of the platform was assessed in presence

of known splicing inhibitors  $Zn^{2+}$ (100 µM-2 mM), $Cu^{2+}$ (100 µM-2 mM), $Pt^{4+}$ (20 µM-1 mM)

along with $Cd^{2+}$ (10 µM-2 mM) and $Hg^{2+}$ (25 µM-2 mM) (Fig. 5) (43). Baseline correction was

achieved by growing H37Ra cells in the absence of toxic metal ions (negative control), while SufB

intein-less *M. sm* cells served as additional experimental control. Cells treated with 100 µM $ZnCl_2$

and $CuCl_2$, 20 µM $PtCl_4$, 10 µM $CdCl_2$, and 25 µM $HgCl_2$, turned pink in color, indicating intact

viability.  At intermediate concentrations, 600 µM of $ZnCl_2$ and $CuCl_2$, 40 µM $PtCl_4$, 25 µM $CdCl_2$,

and 100 µM $HgCl_2$, a transition from blue to purple color was observed, suggesting partial loss of

cell viability. Complete loss of viability was noted at higher concentration range, for $ZnCl_2$ and

$CuCl_2$ (800 µM- 2 mM), $PtCl_4$ (80 µM- 1 mM), $CdCl_2$ (100 µM- 2 mM), and $HgCl_2$ (500 µM- 2

mM) (Fig. 5 ii). Dead non-viable cells induced by $Zn^{2+}$, $Cu^{2+}$, and $Pt^{4+}$  may be correlated to

inhibition of *Mtb* SufB splicing and N-terminal cleavage reactions as published earlier (43). In

contrast, SufB intein-less organism (*M. sm*) remained viable (pink) under similar experimental

conditions,  suggesting cell viability as a splicing independent phenomenon. These observations

may also be used to rule out metal toxicity on cell viability over the test concentration range.

**Fig. 5. Evaluation of _Mtb_ H37Ra-based biosensor platform for high-throughput and multiplex detection of trace metal elements and toxic metal ions (Zn$^{2+}$, Cu$^{2+}$, Pt$^{4+}$, Cd$^{2+}$, and Hg$^{2+}$).** (i) Schematic representation of the multiplexing biosensor platform. (ii) H37Ra cells (10$^{8}$ CFU/ml) were allowed to grow in the presence of ZnCl$_2$ (100 µM- 2 mM), CuCl$_2$ (100 µM- 2 mM), PtCl$_4$ (20 µM- 1 mM), CdCl$_2$ (10 µM- 2 mM) and HgCl$_2$ (25 µM- 2 mM), respectively. Cells incubated with 100 µM of ZnCl$_2$ and CuCl$_2$, 20 µM of PtCl$_4$, 10 µM of CdCl$_2$ and 25 µM of HgCl$_2$ were pink in color indicating healthy viable cells. In contrast, cells exposed to 600 µM of ZnCl$_2$ and CuCl$_2$, 40 µM of PtCl$_4$, 25 µM of CdCl$_2$ and 100 µM of HgCl$_2$ turned purple in color suggesting partial loss of cell viability, whereas cells exposed to 600 µM- 2 mM of ZnCl$_2$ and CuCl$_2$, 80 µM- 1 mM of PtCl$_4$, 100 µM- 2 mM of CdCl$_2$ and 500 µM- 2 mM of HgCl$_2$ remained blue in color suggesting complete loss of viable cell population. For a baseline correction, H37Ra cells were grown in the absence of toxic metal ions as the live cell control (negative control). SufB intein-less organism _Mycobacterium smegmatis (M. sm)_ was used as additional experimental control and treated under similar conditions. _M. sm_ cells exhibited pink color across a varied range of toxic metal ions, suggesting their cell viability as an intein-independent phenomenon. The experiments were performed in triplicates to confirm the reproducibility of data. Cyan stars- complete loss of viability (blue) & Yellow stars- partial loss of viability (purple).

The multiplexing ability of the biosensor platform was further examined by employing metal-specific chelators along with the metal-treated samples. The Cd$^{2+}$-specific chelator ECD and Hg$^{2+}$-specific chelator DMSA were tested alongside the broad-spectrum metal chelator EDTA. Fig. 6 (i) is a schematic representation of the biosensor platform, where indicator mycobacterial cells were incubated with Cd$^{2+}$-specific chelator, ECD. Fig. 6 (ii) demonstrates that co-incubation with

ECD effectively reversed $Cd^{2+}$- induced loss of viability, restoring pink color. This effect was not

observed in cells exposed to other metal ions, where ECD could not block metal induced inhibition.

Similarly, DMSA reversed the toxic effects of $Hg^{2+}$ [Fig. 6 (iii) and (iv)], while other metal ions

continued to attenuate cell viability. Notably, Since EDTA is a broad-spectrum metal chelator, it

restored viability across all metal-treated samples [Fig. 6 (v) and (vi)]. Collectively, these findings

validate the high-throughput and multiplexing capability of the developed biosensor platform,

underscoring its potential for simultaneous detection of multiple toxic metal ions with high

specificity and sensitivity.

**Fig. 6. Determination of specificity of the *Mtb* H37Ra-based multiplexing biosensor platform.**
(i) Schematic representation of the multiplexing biosensor platform along with $Cd^{2+}$ specific metal
chelator (ECD) (ii) *Mtb* H37Ra cells ($10^8$ CFU/ml) were incubated with different concentration of
$ZnCl_2$, $CuCl_2$, $PtCl_4$, $CdCl_2$ and $HgCl_2$, along with ECD (1:1 ratio). ECD reversed the inhibitory
effect of $Cd^{2+}$ rendering viable pink cells, whereas the effects of other metal ions persisted (iii)
Schematic representation of the multiplexing biosensor platform along with $Hg^{2+}$ specific metal
chelator (DMSA). (iv) *Mtb* H37Ra cells ($10^8$ CFU/ml) were incubated with different concentration

of $ZnCl_2$ ,$CuCl_2$, $PtCl_4$ , $CdCl_2$ and $HgCl_2$, along with DMSA in the ratio 1:1. DMSA inhibited the
effect of $Hg^{2+}$ inducing viable pink cells, whereas the effect of other metal ions persisted. (v)
Schematic representation of plate-based biosensor platform in presence of common metal chelator
EDTA. (vi) *Mtb* H37Ra cells ($10^8$ CFU/ml) were allowed to grow in presence of $ZnCl_2$ ,$CuCl_2$,
$PtCl_4$ , $CdCl_2$ and $HgCl_2$ along with EDTA in the ratio 1:1. The inhibitory effects of test metal ions
on mycobacterial cells were reversed by EDTA. For a baseline correction, H37Ra cells were grown
in the absence of toxic metal ions, as the live cell control. SufB intein-less organism
*Mycobacterium smegmatis (M. sm)* was used as additional experimental control and treated under
similar conditions. *M. sm* cells exhibited pink color across a varied range of toxic metal ions,
suggesting their cell viability as an intein-independent phenomenon. The experiments were
performed in triplicates to confirm the reproducibility of data. Cyan stars- complete loss of
viability (blue) & Yellow stars- partial loss of viable cells (purple).  ECD: Edetate Calcium
disodium, DMSA: Meso-2,3-Dimercaptosuccinic acid, EDTA: Ethylenediaminetetraacetic Acid.

In toto, the designed biosensor has demonstrated the potential of whole-cell *Mtb* H37Ra as an

indicator to detect toxic metal ions as well as TMEs in test samples. The specificity of the designed

metal-sensor was shown by the reversal of the inhibitory effects in the presence of $Cd^{2+}$- and $Hg^{2+}$-

specific metal chelators, ECD and DMSA, respectively. Detection of metal ions at a concentration

of 25 μM ($Cd^{2+}$) highlights the sensitivity of the biosensor platform, although it requires further

optimization to cover the lower concentration range of various metal ions. We propose the use of

whole-cell microorganisms carrying metal-sensing intein-bearing proteins as an indicator for the

presence of toxic metal ions in environmental samples. This is a user-friendly method that can be

performed in a 96-well microtiter plate, which would provide a color change between blue and

pink while sensing metal ions. Since the activity of metal-sensing intein-carrying protein

sequences are unique for the indicator (whole-cell) organisms, in the unlikely event of

contamination in a biological facility, it should not raise concern about the validity of the results.

**Conclusion**

Heavy metal ions are persistent environmental contaminants that pose significant risks to both

ecological systems and human health. With the rising burden of metal pollution across diverse

environmental and biological matrices, there is a pressing need for sensitive, specific, and user-

friendly biosensing technology. Intein-mediated biosensors are emerging as promising tools in this

context, particularly due to their unique splicing mechanism that operates efficiently under

physiological conditions, eliminating the need for external cofactors or energy sources (34, 135).

In this study, an innovative biosensor platform was developed, employing native H37Ra cells

where intein splicing inhibition is a readout mechanism for the detection of toxic heavy metal ions

(e.g., $Cd^{2+}$ and $Hg^{2+}$) and other trace metal elements (Fig. 7). The resulting alteration in

mycobacterial viability is captured through a rapid and quantifiable resazurin-based colorimetric

assay that reports color change between blue-purple-pink. Importantly, the platform is adaptable

for multiplex detection of metal ions, including $Zn^{2+}$, $Pt^{4+}$, $Cu^{2+}$, $Cd^{2+}$, and $Hg^{2+}$, thereby offering

a high-throughput solution for complex sample screening (Fig. 5 and 6).  Further, this can be

extended to sense metal ions by other whole-cell microorganisms, harbouring unique metal-

sensing intein-containing proteins,  hence offering a qualitative as well as quantitative detection

method.

SufB is a critical component of SUF system, the exclusive pathway for Fe-S cluster assembly in

mycobacteria.  Previous studies have  also highlighted the pivotal role of *Mtb* SufB protein in Fe-

S cluster biogenesis, and mycobacterial survival (61, 136). Since the functionality of SUF protein

complex is dependent on generation of active SufB protein via splicing, splicing inhibition presents

a promising approach for antimicrobial intervention by targeting mycobacterial viability. Several

studies have shown that divalent cations ($Zn^{2+}$and $Cu^{2+}$), ROS, RNS, chlorine derivatives, small

molecules (cisplatin and diethanolamine) and zinc oxide nanoparticles can act as potent splicing

inhibitors. These agents have influenced mycobacterial survival, further validating the relevance

of targeting intein splicing for antimycobacterial drug development (40, 43, 51, 56, 58, 64, 66,

137–139). Earlier work from our group has demonstrated that metal ions such as $Cu^{2+}$, $Zn^{2+}$, and

$Pt^{4+}$ inhibit SufB splicing, establishing a correlation between metal-induced stress and altered

intein splicing activity in *Mtb* (43). Building on these findings, the current work explores

regulatory roles of toxic metal ions on *Mtb* SufB splicing and cleavage activity. Alamar Blue assay

was employed to examine the influence of TMEs and toxic metal ions on the viability of *Mtb* cells.

The outcome of this research will advance the real time application of H37Ra/*Mtb* based whole-

cell biosensor to detect wide range of metal ions. The platform described herein is robust, enabling

multiplex screening of metal ions while maintaining specificity and sensitivity. Importantly,

cellular viability loss in response to metal exposure is linked to an observable color shift between

blue and pink, providing a quantifiable readout for metal ion detection in test samples.

In this study, splicing inhibition was evaluated both *in vitro*, using purified SufB protein and *in*

*vivo* within H37Ra cells in a whole-cell biosensor platform, where the observed effects on

mycobacterial viability were correlated to SufB splicing inhibition. Given that H37Ra is a slow-

growing strain with a mean generation time of approximately 44 hours, the experimental window

was established at four days to encompass at least two full generations, permitting effective

assessment of metal-induced viability changes (140–142). Notably, the logarithmic growth phase

of mycobacteria extends from four to twenty days, ensuring that the platform remains adaptable to

varied experimental requirements. To further strengthen our hypothesis, SufB intein less organism,

*M. sm* control was included; viability of *M. sm* persisted throughout the experimental period,

suggesting that the tested metal ions do not elicit growth inhibitory or toxic effects in the absence

of SufB intein. *Mtb* contains three intein containing proteins; SufB, RecA and DnaB. While SufB

is essential for Fe-S cluster assembly, RecA and DnaB play pivotal roles during DNA replication;

taken together these proteins are critical for mycobacterial survival (143–145). Although current

study explains loss of mycobacterial cell viability as a result of inhibition of *Mtb* SufB splicing,

the possibility of RecA and DnaB splicing inhibition by the test metal ions cannot be ruled

completely.  Currently, there are no literatures to show inhibitory effect of metal ions other than

$Zn^{2+}$ on *Mtb* DnaB splicing (49). Moreover there are alternative helicases present in mycobacteria

to take over the function of DnaB (123–125). Several studies have reported regulatory roles of

$Zn^{2+}$, $Cd^{2+}$, $Co^{2+}$, $Ni^{2+}$, $Cu^+$/$Cu^{2+}$, CisPt  ions on *Mtb* RecA intein splicing (45, 47, 58, 137).

However, current study has not evaluated the RecA intein inhibition in mycobacteria. Hence, this

approach offers a multiplexing biosensor unit which can accommodate cell viability detection via

splicing inhibition in many organisms carrying different metal-sensitive intein containing proteins

simultaneously.

The use of non-genetically modified *Mtb* H37Ra cells ensures biosecurity compliance while

providing a biologically relevant sensing chassis. These mycobacterial cells are not only easy to

cultivate and maintain but also exhibit remarkable long-term viability, supporting extended shelf-

life and storage without specialized conditions. The platform also demonstrates good sensitivity in

detecting metal ions over a broad dynamic range, and substrate specificity, making it suitable for

metal ion screening in standard laboratory setups.

However, some limitations persist such as complex environmental or biological samples

containing mixtures of multiple metal ions may result in signal overlap or cross-reactivity with

variable binding affinities. Furthermore, while microbial whole-cell biosensors offer an excellent

alternative for initial screening and field deployment, advanced physicochemical instruments

remain crucial when trace-level quantification is required, where intein activity may not be

efficiently regulated.

In summary, our findings highlight the potential of whole cell intein-based biosensors as user-

friendly, and cost-effective tools for the detection of toxic metal and trace metal ions. Future work

should aim to refine their activity, enhancing environmental robustness, and advancing the

platform from lab-scale to deployable technologies. This can be accelerated through collaborative

efforts across academic, industrial, and governmental sectors.

**Fig. 7. Schematic representation explaining the principle of *Mtb* H37Ra-based multiplexing**

**biosensor platform.** Whole-cell mycobacteria (H37Ra) were used as biological indicator for the

detection of toxic metal ions ($Cd^{2+}$ and $Hg^{2+}$) and other trace metal elements (TME). (i) Intein

splicing under normal physiological conditions generates functional SufB protein, while viable
mycobacterial cells convert resazurin (blue) to resorufin (pink). (ii) In the presence of toxic metal
ions $Cd^{2+}$ and $Hg^{2+}$ and TME, SufB intein splicing is inhibited resulting in non-viable
mycobacterial cells, thus there is no conversion of resazurin (blue) to resorufin (pink).

## Data availability

905 The data underlying the present article are available in the article and in its online Supplementary

906 materials. Additional data will be made available on request if needed.

## Supplemental Data

908 This article contains supplemental data.

## Acknowledgment

910 The plasmids used in this study are borrowed from Prof. Marlene Belfort's Lab, SUNY, Albany,

911 NY, USA. Our sincere gratitude to the Central Instrumentation Facility (CIF) at OUAT (Odisha

912 University of Agriculture & Technology), Bhubaneswar, and Dr. S.K. Dash for his help in

913 conducting the ICP-OES analysis at OUAT, Bhubaneswar.

## Author Contributions

915 **AM:** Writing – Original Draft, Writing - Review & Editing, Methodology, Investigation, Formal

916 analysis, Data curation, Visualization, Validation, Software, Resources, Funding acquisition; **AN:**

917 Conceptualization, Methodology, Investigation, Formal analysis, Data curation; Writing - Review

918 & Editing, ; **SN:** Conceptualization, Writing - Original Draft, Writing - Review & Editing, Project

919 administration, Supervision, Methodology, Investigation, Resources,  Funding acquisition.

**Funding and Additional Information**

921 Ms. Ashwaria Mehra is supported by INSPIRE fellowship (DST/INSPIRE/03/2021/002800/IF 923 190921); INSPIRE Division, DST, Government of India. This work was also supported by the 924 UGC-DAE consortium for scientific research (UGC-DAE-CSR-KC/CRS/15/IOP/08/0562), 925 Kolkata, India.

**Conflict of Interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

943
**References**

1. Velusamy, K., Periyasamy, S., Kumar, P. S., Rangasamy, G., Nisha Pauline, J. M.,
Ramaraju, P., Mohanasundaram, S., and Nguyen Vo, D. V. (2022) Biosensor for heavy
metals detection in wastewater: A review. *Food and Chemical Toxicology* 168, 113307

2. Duffus, J. H. (2002) *"heavy metals" - A meaningless term? (IUPAC technical report)*
(Walter de Gruyter GmbH)

3. Turdean, G. L., and Ozkan, S. A. (2011) Design and Development of Biosensors for the
Detection of Heavy Metal Toxicity. *International Journal of Electrochemistry* 2011,
343125

4. Singh, R., Gautam, N., Mishra, A., and Gupta, R. (2011) Heavy metals and living
systems: An overview. *Indian J Pharmacol* 43, 246–253

5. Angon, P. B., Islam, M. S., KC, S., Das, A., Anjum, N., Poudel, A., and Suchi, S. A.
(2024) Sources, effects and present perspectives of heavy metals contamination: Soil,
plants and human food chain. *Heliyon* 10, e28357

6. Alengebawy, A., Abdelkhalek, S. T., Qureshi, S. R., and Wang, M. Q. (2021) Heavy
Metals and Pesticides Toxicity in Agricultural Soil and Plants: Ecological Risks and
Human Health Implications. *Toxics* 9, 42

7. Jaishankar, M., Tseten, T., Anbalagan, N., Mathew, B. B., and Beeregowda, K. N. (2014)
Toxicity, mechanism and health effects of some heavy metals. *Interdiscip Toxicol* 7, 60–
72

8. Hassani, S., Rezaei Akmal, M., Salek Maghsoudi, A., Rahmani, S., Vakhshiteh, F.,
Norouzi, P., Ganjali, M. R., and Abdollahi, M. (2020) High-Performance Voltammetric
Aptasensing Platform for Ultrasensitive Detection of Bisphenol A as an Environmental
Pollutant. *Front Bioeng Biotechnol* 8, 574846

9. Odobašić, A., Šestan, I., Begić, S., Odobašić, A., Šestan, I., and Begić, S. (2019)
*Biosensors for Determination of Heavy Metals in Waters* (IntechOpen)

10. Ramnani, P., Saucedo, N. M., and Mulchandani, A. (2016) Carbon nanomaterial-based
electrochemical biosensors for label-free sensing of environmental pollutants.
*Chemosphere* 143, 85–98

11. Pena-Pereira, F., Wojnowski, W., and Tobiszewski, M. (2020) AGREE - Analytical
GREEnness Metric Approach and Software. *Anal Chem* 92, 10076–10082

12.   Pearson, J. E., Gill, A., and Vadgama, P. (2000) Analytical aspects of biosensors. *Ann*
*Clin Biochem* 37, 119–145

13.   Malandain, C., Fayolle, F., and Bedouelle, H. (2005) Biosensors for the Environment. *Oil*
*& Gas Science and Technology* 60, 887–897

14.   Bontidean, I., Berggren, C., Johansson, G., Csöregi, E., Mattiasson, B., Lloyd, J. R.,
Jakeman, K. J., and Brown, N. L. (1998) Detection of Heavy Metal Ions at Femtomolar
Levels Using Protein-Based Biosensors. *Anal Chem* 70, 4162–4169

15.   Blake, D. A., Jones, R. M., Blake, R. C., Pavlov, A. R., Darwish, I. A., and Yu, H. (2001)
Antibody-based sensors for heavy metal ions. *Biosens Bioelectron* 16, 799–809

16.   Chouteau, C., Dzyadevych, S., Durrieu, C., and Chovelon, J. M. (2005) A bi-enzymatic
whole cell conductometric biosensor for heavy metal ions and pesticides detection in
water samples. *Biosens Bioelectron* 21, 273–281

17.   Mohammadi, Z. B., Zhang, F., Kharazmi, M. S., and Jafari, S. M. (2023) Nano-
biocatalysts for food applications; immobilized enzymes within different nanostructures.
*Crit Rev Food Sci Nutr* 63, 11351–11369

18.   Sun, G., Wei, X., Zhang, D., Huang, L., Liu, H., and Fang, H. (2023) Immobilization of
Enzyme Electrochemical Biosensors and Their Application to Food Bioprocess
Monitoring. *Biosensors (Basel)* 13, 886

19.   Kakkar, P., and Jaffery, F. N. (2005) Biological markers for metal toxicity. *Environ*
*Toxicol Pharmacol* 19, 335–349

20.   Castillo, J., Gáspár, S., Leth, S., Niculescu, M., Mortari, A., Bontidean, I., Soukharev, V.,
Dorneanu, S. A., Ryabov, A. D., and Csöregi, E. (2004) Biosensors for life quality:
Design, development and applications. *Sens Actuators B Chem* 102, 179–194

21.   Amaro, F., Turkewitz, A. P., Martín-González, A., and Gutiérrez, J. C. (2011) Whole-cell
biosensors for detection of heavy metal ions in environmental samples based on
metallothionein promoters from Tetrahymena thermophila. *Microb Biotechnol* 4, 513–522

22.   Kim, Y., Choi, H., Shin, W. H., Oh, J. M., Koo, S. M., Kim, Y., Lee, T., Yu, B. J., and
Park, C. (2021) Development of Colorimetric Whole-Cell Biosensor for Detection of
Heavy Metals in Environment for Public Health. *International Journal of Environmental*
*Research and Public Health 2021, Vol. 18, Page 12721* 18, 12721

23.   Bousse, L. (1996) Whole cell biosensors. *Sens Actuators B Chem* 34, 270–275

24.   Stenger, D. A., Gross, G. W., Keefer, E. W., Shaffer, K. M., Andreadis, J. D., Ma, W., and
Pancrazio, J. J. (2001) Detection of physiologically active compounds using cell-based
biosensors. *Trends Biotechnol* 19, 304–309

25.   Wang, Y.-Z., Christopher, J. K., Yong, Y.-C., and Zhai, D.-D. (2019) *Nutrient Detection*
*with Whole-Cell Biosensors* (Springer, Cham)

26.   Mihaela, G., and Mihaela, G. (2021) A short review on cell-based biosensing: challenges
and breakthroughs in biomedical analysis. *J Biomed Res* 35, 255–263

27.   Raut, N., O'Connor, G., Pasini, P., and Daunert, S. (2012) Engineered cells as biosensing
systems in biomedical analysis. *Anal Bioanal Chem* 402, 3147–3159

28.   Gui, Q., Lawson, T., Shan, S., Yan, L., and Liu, Y. (2017) The Application of Whole Cell-
Based Biosensors for Use in Environmental Analysis and in Medical Diagnostics. *Sensors*
17, 1623

29.   Kannappan, S., Bhaskar, ·, Ramisetty, C. M., Chandra, B., and Ramisetty, M. (2021)
Engineered Whole-Cell-Based Biosensors: Sensing Environmental Heavy Metal
Pollutants in Water—a Review. *Applied Biochemistry and Biotechnology 2021 194:4* 194,
1814–1840

30.   Kang, C., Shrestha, K. L., Kwon, S., Park, S., Kim, J., and Kwon, Y. (2022) Intein-
Mediated Protein Engineering for Cell-Based Biosensors. *Biosensors (Basel)* 12, 283

31.   Jeon, H., Lee, E., Kim, D., Lee, M., Ryu, J., Kang, C., Kim, S., and Kwon, Y. (2018) Cell-
Based Biosensors Based on Intein-Mediated Protein Engineering for Detection of
Biologically Active Signaling Molecules. *Anal Chem* 90, 9779–9786

32.   Lee, E., Shrestha, K. L., Kang, S., Ramakrishnan, N., and Kwon, Y. (2023) Cell-Based
Sensors for the Detection of EGF and EGF-Stimulated Ca2+ Signaling. *Biosensors*
*(Basel)* 13, 383

33.   Jeon, H., Lee, M., Jang, W., and Kwon, Y. (2016) Intein-mediated protein engineering for
biosensor fabrication. *Biochip J* 10, 277–287

34.   Mills, K. V., Johnson, M. A., and Perler, F. B. (2014) Protein Splicing: How Inteins
Escape from Precursor Proteins. *Journal of Biological Chemistry* 289, 14498–14505

35.   Perler, F. B. (2005) Protein splicing mechanisms and applications. *IUBMB Life* 57, 469–
476

36.   Panda, S., Nanda, A., Nasker, S. S., Sen, D., Mehra, A., and Nayak, S. (2021) Metal effect
on intein splicing: A review. *Biochimie* 185, 53–67

37. Nichols, N. M., Benner, J. S., Martin, D. D., and Evans, T. C. (2003) Zinc ion effects on
individual Ssp DNaE intein splicing steps: Regulating pathway progression. *Biochemistry*
42, 5301–5311

38. Chan, H., Pearson, C. S., Green, C. M., Li, Z., Zhang, J., Belfort, G., Shekhtman, A., Li,
H., and Belfort, M. (2016) Exploring Intein Inhibition by Platinum Compounds as an
Antimicrobial Strategy. *Journal of Biological Chemistry* 291, 22661–22670

39. Green, C. M., Li, Z., Smith, A. D., Novikova, O., Bacot-Davis, V. R., Gao, F., Hu, S.,
Banavali, N. K., Thiele, D. J., Li, H., and Belfort, M. (2019) Spliceosomal Prp8 intein at
the crossroads of protein and RNA splicing. *PLoS Biol* 17, e3000104

40. Ojha, D. K., Mehra, A., Rout, S. S., Giri, S., and Nayak, S. (2025) Bactericidal activity of
ZnO nanoparticles–anti-TB drug combination towards the H37Rv strain and multidrug-
resistant isolates of Mycobacterium tuberculosis via SufB splicing inhibition. *Mater Adv*
6, 5242–5259

41. Gramespacher, J. A., Stevens, A. J., Nguyen, D. P., Chin, J. W., and Muir, T. W. (2017)
Intein Zymogens: Conditional Assembly and Splicing of Split Inteins via Targeted
Proteolysis. *J Am Chem Soc* 139, 8074–8077

42. Skretas, G., and Wood, D. W. (2005) Regulation of protein activity with small-molecule-
controlled inteins. *Protein Science* 14, 523–532

43. Panda, S., Nanda, A., Nasker, S. S., Mehra, A., Ojha, D. K., Mohanty, P. S., and Nayak, S.
(2023) Metal regulation of Mycobacterium tuberculosis SufB intein splicing at the host–
pathogen crossroad. *Clin Sci* 137, 1027–1048

44. Poland, B. W., Xu, M. Q., and Quiocho, F. A. (2000) Structural Insights into the Protein
Splicing Mechanism of PI-SceI. *Journal of Biological Chemistry* 275, 16408–16413

45. Mills, K. V., and Paulus, H. (2001) Reversible Inhibition of Protein Splicing by Zinc Ion.
*Journal of Biological Chemistry* 276, 10832–10838

46. Van Roey, P., Pereira, B., Li, Z., Hiraga, K., Belfort, M., and Derbyshire, V. (2007)
Crystallographic and Mutational Studies of Mycobacterium tuberculosis recA Mini-inteins
Suggest a Pivotal Role for a Highly Conserved Aspartate Residue. *J Mol Biol* 367, 162–
173

47. Zhang, L., Zheng, Y., Xi, Z., Luo, Z., Xu, X., Wang, C., and Liu, Y. (2009) Metal ions
binding to recA inteins from Mycobacterium tuberculosis. *Mol Biosyst* 5, 644–650

48. Sun, P., Ye, S., Ferrandon, S., Evans, T. C., Xu, M. Q., and Rao, Z. (2005) Crystal
Structures of an Intein from the Split dnaE Gene of Synechocystis sp. PCC6803 Reveal

the Catalytic Model Without the Penultimate Histidine and the Mechanism of Zinc Ion
Inhibition of Protein Splicing. *J Mol Biol* 353, 1093–1105

49.    Woods, D., Vangaveti, S., Egbanum, I., Sweeney, A. M., Li, Z., Bacot-Davis, V.,
Lesassier, D. S., Stanger, M., Hardison, G. E., Li, H., Belfort, M., and Lennon, C. W.
(2020) Conditional DnaB protein splicing is reversibly inhibited by zinc in mycobacteria.
*mBio* 11, 1–14

50.    Pearson, C. S., Belfort, G., Belfort, M., and Shekhtman, A. (2015) Backbone assignments
of mini-RecA intein with short native exteins and an active N-terminal catalytic cysteine.
*Biomol NMR Assign* 9, 235–238

51.    Li, Z., Fu, B., Green, C. M., Liu, B., Zhang, J., Lang, Y., Chaturvedi, S., Belfort, M., Liao,
G., and Li, H. (2019) Cisplatin protects mice from challenge of Cryptococcus neoformans
by targeting the Prp8 intein. *Emerg Microbes Infect* 8, 895–908

52.    Ozawa, T., Nogami, S., Sato, M., Ohya, Y., and Umezawa, Y. (2000) A Fluorescent
Indicator for Detecting Protein−Protein Interactions in Vivo Based on Protein Splicing.
*Anal Chem* 72, 5151–5157

53.    Ozawa, T., Kaihara, A., Sato, M., Tachihara, K., and Umezawa, Y. (2001) Split
Luciferase as an Optical Probe for Detecting Protein−Protein Interactions in Mammalian
Cells Based on Protein Splicing. *Anal Chem* 73, 2516–2521

54.    Paulmurugan, R., Umezawa, Y., and Gambhir, S. S. (2002) Noninvasive imaging of
protein–protein interactions in living subjects by using reporter protein complementation
and reconstitution strategies. *Proceedings of the National Academy of Sciences* 99,
15608–15613

55.    Kanno, A., Ozawa, T., and Umezawa, Y. (2005) Intein-Mediated Reporter Gene Assay for
Detecting Protein−Protein Interactions in Living Mammalian Cells. *Anal Chem* 78, 556–
560

56.    Topilina, N. I., Green, C. M., Jayachandran, P., Kelley, D. S., Stanger, M. J., Piazza, C.
L., Nayak, S., and Belfort, M. (2015) SufB intein of Mycobacterium tuberculosis as a
sensor for oxidative and nitrosative stresses. *Proc Natl Acad Sci U S A* 112, 10348–10353

57.    Zhang, L., Xiao, N., Pan, Y., Zheng, Y., Pan, Z., Luo, Z., Xu, X., and Liu, Y. (2010)
Binding and inhibition of copper ions to RecA inteins from mycobacterium tuberculosis.
*Chemistry - A European Journal* 16, 4297–4306

58.    Zhang, L., Zheng, Y., Callahan, B., Belfort, M., and Liu, Y. (2011) Cisplatin Inhibits
Protein Splicing, Suggesting Inteins as Therapeutic Targets in Mycobacteria. *Journal of
Biological Chemistry* 286, 1277–1282

59.    Dassa, B., Amitai, G., Caspi, J., Schueler-Furman, O., and Pietrokovski, S. (2007) Trans
1106           protein splicing of cyanobacterial split inteins in endogenous and exogenous
1107           combinations. *Biochemistry* 46, 322–330

1108    60.    Huet, G., Daffé, M., and Saves, I. (2005) Identification of the Mycobacterium tuberculosis
1109           SUF machinery as the exclusive mycobacterial system of [Fe-S] cluster assembly:
1110           Evidence for its implication in the pathogen's survival. *J Bacteriol* 187, 6137–6146

1111    61.    Huet, G., Castaing, J. P., Fournier, D., Daffé, M., and Saves, I. (2006) Protein splicing of
1112           SufB is crucial for the functionality of the Mycobacterium tuberculosis SUF machinery. *J*
1113           *Bacteriol* 188, 3412–3414

1114    62.    Panda, S., Nanda, A., Sahu, N., Ojha, D. K., Pradhan, B., Rai, A., Suryawanshi, A. R.,
1115           Banavali, N., and Nayak, S. (2022) SufB intein splicing in Mycobacterium tuberculosis is
1116           influenced by two remote conserved N-extein histidines. *Biosci Rep* 42, 20212207

1117    63.    Friedel, K., Popp, M. A., Matern, J. C. J., Gazdag, E. M., Thiel, I. V., Volkmann, G.,
1118           Blankenfeldt, W., and Mootz, H. D. (2018) A functional interplay between intein and
1119           extein sequences in protein splicing compensates for the essential block B histidine. *Chem*
1120           *Sci* 10, 239–251

1121    64.    Kelley, D. S., Lennon, C. W., Li, Z., Miller, M. R., Banavali, N. K., Li, H., and Belfort,
1122           M. (2018) Mycobacterial DnaB helicase intein as oxidative stress sensor. *Nat Commun* 9,
1123           1–15

1124    65.    Mills, K. V., Lew, B. M., Jiang, S. Q., and Paulus, H. (1998) Protein splicing in trans by
1125           purified N- and C-terminal fragments of the Mycobacterium tuberculosis RecA intein.
1126           *Proc Natl Acad Sci U S A* 95, 3543–3548

1127    66.    Gangopadhyay, J. P., Jiang, S. qin, and Paulus, H. (2003) An in vitro screening system for
1128           protein splicing inhibitors based on green fluorescent protein as an indicator. *Anal Chem*
1129           75, 2456–2462

1130    67.    Nanda, A., Nasker, S. S., Kushwaha, A. K., Ojha, D. K., Dearden, A. K., Nayak, S. K.,
1131           and Nayak, S. (2021) Gold Nanoparticles Augment N-Terminal Cleavage and Splicing
1132           Reactions in Mycobacterium tuberculosis SufB. *Front Bioeng Biotechnol* 9, 773303

1133    68.    Chemiluminescence, B., Li, J., Xing, H., Wang, E., Zhao, J., Li, D., Ma, J., Suhartono, E.,
1134           Thalib, I., Aflanie, I., Noor, Z., and Idroes, R. (2018) in *IOP Conference Series: Materials*
1135           *Science and Engineering* (IOP Publishing), p 012008.

1136    69.    Grzyska, P. K., Hausinger, R. P., and Proshlyakov, D. A. (2010) Metal and substrate
1137           binding to an Fe(II) dioxygenase resolved by UV spectroscopy with global regression
1138           analysis. *Anal Biochem* 399, 64–71

70.  Plotnikova, O. A., Mel'nikov, A. G., Mel'nikov, G. V., and Gubina, T. I. (2016)
Quenching of tryptophan fluorescence of bovine serum albumin under the effect of ions of
heavy metals. *Opt Spectrosc* 120, 65–69

71.  Oliveira, A., Baccana, N., and Cadore, S. (2012) Evaluation of metal ions in rice samples:
extraction and direct determination by ICP OES. *J Braz Chem Soc* 23, 838–845

72.  Harrington, J. M., Young, D. J., Essader, A. S., Sumner, S. J., and Levine, K. E. (2014)
Analysis of human serum and whole blood for mineral content by ICP-MS and ICP-OES:
Development of a mineralomics method. *Biol Trace Elem Res* 160, 132–142

73.  Punjabi, K., Mehta, S., Chavan, R., Chitalia, V., Deogharkar, D., and Deshpande, S.
(2018) Efficiency of biosynthesized silver and zinc nanoparticles against multi-drug
resistant pathogens. *Front Microbiol* 9, 376058

74.  Longhin, E. M., El Yamani, N., Rundén-Pran, E., and Dusinska, M. (2022) The alamar
blue assay in the context of safety testing of nanomaterials. *Frontiers in Toxicology* 4,
981701

75.  Yang, L., Zhang, Y., Wang, F., Luo, Z., Guo, S., and Strähle, U. (2020) Toxicity of
mercury: Molecular evidence. *Chemosphere* 245, 125586

76.  Aljerf, L., Almasri, N., and Access, O. (2018) Mercury toxicity: ecological features of
organic phase of mercury in biota-Part I. *Inorganic Chemical Sciences* 3,

77.  Budnik, L. T., and Casteleyn, L. (2019) Mercury pollution in modern times and its socio-
medical consequences. *Science of The Total Environment* 654, 720–734

78.  Dash, H. R., Mangwani, N., and Das, S. (2014) Characterization and potential application
in mercury bioremediation of highly mercury-resistant marine bacterium Bacillus
thuringiensis PW-05. *Environmental Science and Pollution Research* 21, 2642–2653

79.  Tang, W., Zhang, W., Zhao, Y., Zhang, H., and Shan, B. (2017) Basin-scale
comprehensive assessment of cadmium pollution, risk, and toxicity in riverine sediments
of the Haihe Basin in north China. *Ecol Indic* 81, 295–301

80.  White, P. J., and Brown, P. H. (2010) Plant nutrition for sustainable development and
global health. *Ann Bot* 105, 1073–1080

81.  Losada Ros, M. T., Al-Enezi, E., Cesarini, E., Canonico, B., Bucci, C., Martins, M. V. A.,
Papa, S., and Frontalini, F. (2020) Assessing the Cadmium Effects on the Benthic
Foraminifer Ammonia cf. parkinsoniana: An Acute Toxicity Test. *Water (Basel)* 12, 1018

82. Dong, F., Zhu, X., Qian, W., Wang, P., and Wang, J. (2020) Combined effects of CO2-
driven ocean acidification and Cd stress in the marine environment: Enhanced tolerance of
Phaeodactylum tricornutum to Cd exposure. *Mar Pollut Bull* 150, 110594

83. Atamaleki, A., Sadani, M., Raoofi, A., Miri, A., Bajestani, S. G., Fakhri, Y., Heidarinejad,
Z., and Mousavi Khaneghah, A. (2020) The concentration of potentially toxic elements
(PTEs) in eggs: A global systematic review, meta-analysis and probabilistic health risk
assessment. *Trends Food Sci Technol* 95, 1–9

84. Stankovic, S., Kalaba, P., and Stankovic, A. R. (2014) Biota as toxic metal indicators.
*Environ Chem Lett* 12, 63–84

85. Joe, M. H., Lee, K. H., Lim, S. Y., Im, S. H., Song, H. P., Lee, I. S., and Kim, D. H.
(2012) Pigment-based whole-cell biosensor system for cadmium detection using
genetically engineered Deinococcus radiodurans. *Bioprocess Biosyst Eng* 35, 265–272

86. Guertin, J., Jacobs, J. A., and Avakian, C. P. (2004) *Chromium(VI) Handbook* (CRC
Press). 1st Ed.

87. Dawra, N., and Dabas, N. (2024) Advances in spectrophotometric determination of
Chromium(III) and Chromium(VI) in water: a review. *Int J Environ Anal Chem* 104,
2994–3015

88. Sheehan, P. J., Meyer, D. M., Sauer, M. M., and Paustenbach, D. J. (1991) Assessment of
the human health risks posed by exposure to chromium-contaminated soils. *Journal of
Toxicology and Environmental Health, Part A Current Issues* 32, 161–201

89. Mishra, S., and Bharagava, R. N. (2016) Toxic and genotoxic effects of hexavalent
chromium in environment and its bioremediation strategies. *Journal of Environmental
Science and Health, Part C* 34, 1–32

90. Feleafel, M. N., and Mirdad, Z. M. (2013) Hazard and Effects of Pollution by Lead on
Vegetable Crops. *J Agric Environ Ethics* 26, 547–567

91. Ghosh, I., Sun, L., and Xu, M. Q. (2001) Zinc Inhibition of Protein trans-Splicing and
Identification of Regions Essential for Splicing and Association of a Split Intein*. *Journal
of Biological Chemistry* 276, 24051–24058

92. Tharappel, A. M., Li, Z., and Li, H. (2022) Inteins as Drug Targets and Therapeutic Tools.
*Front Mol Biosci* 9, 821146

93. Alhazmi, H. A., Alam, M. S., Albratty, M., Najmi, A., Abdulhaq, A. A., Hassani, R.,
Ahsan, W., and Qramish, A. N. (2023) Binding Investigation of Some Important Metal
Ions Copper (I), Nickel (II), and Aluminium (III) with Bovine Serum Albumin Using
Valid Spectroscopic Techniques. *J Chem* 2023, 2581653

94.    Permyakov, E. A., and Gonçalves, A. M. M. (2021) Metal Binding Proteins. *Encyclopedia*
1205           *1*, 261–292

1206    95.    Woźnicka, O., Rybarska, J., Jagusiak, A., Konieczny, L., Stopa, B., and Roterman, I.
1207           (2017) Metal ions introduced to proteins by supramolecular ligands. *Self-Assembled*
1208           *Molecules - New Kind of Protein Ligands: Supramolecular Ligands*, 61–76

1209    96.    Antosiewicz, J. M., and Shugar, D. (2016) UV–Vis spectroscopy of tyrosine side-groups
1210           in studies of protein structure. Part 2: selected applications. *Biophys Rev* 8, 163–177

1211    97.    Biter, A. B., Pollet, J., Chen, W. H., Strych, U., Hotez, P. J., and Bottazzi, M. E. (2019) A
1212           method to probe protein structure from UV absorbance spectra. *Anal Biochem* 587,
1213           113450

1214    98.    Dieaconu, M., Ioanid, A., Iftimie, S., and Antohe, S. UV-ABSORPTION MECHANISMS
1215           OF Ni 2+-BINDING BOVINE SERUM ALBUMIN. *Dig J Nanomater Biostruct* 7, 1125–
1216           1138

1217    99.    View of Study of immunoglobulin 'G' with ultra violet spectroscopy in Duchene muscular
1218           dystrophy and Alzheimer's disease

1219    100.   Bjørklund, G., Dadar, M., Mutter, J., and Aaseth, J. (2017) The toxicology of mercury:
1220           Current research and emerging trends. *Environ Res* 159, 545–554

1221    101.   Ajsuvakova, O. P., Tinkov, A. A., Aschner, M., Rocha, J. B. T., Michalke, B., Skalnaya,
1222           M. G., Skalny, A. V., Butnariu, M., Dadar, M., Sarac, I., Aaseth, J., and Bjørklund, G.
1223           (2020) Sulfhydryl groups as targets of mercury toxicity. *Coord Chem Rev* 417, 213343

1224    102.   Sall, M. L., Diaw, A. K. D., Gningue-Sall, D., Efremova Aaron, S., and Aaron, J. J. (2020)
1225           Toxic heavy metals: impact on the environment and human health, and treatment with
1226           conducting organic polymers, a review. *Environmental Science and Pollution Research*
1227           27, 29927–29942

1228    103.   Gonick, H. C. (2011) Lead-Binding Proteins: A Review. *J Toxicol* 2011, 686050

1229    104.   Fossum, C. J., Johnson, B. O. V., Golde, S. T., Kielman, A. J., Finke, B., Smith, M. A.,
1230           Lowater, H. R., Laatsch, B. F., Bhattacharyya, S., and Hati, S. (2023) Insights into the
1231           Mechanism of Tryptophan Fluorescence Quenching due to Synthetic Crowding Agents: A
1232           Combined Experimental and Computational Study. *ACS Omega* 8, 44820–44830

1233    105.   Liu, Y., Xu, J., Han, L., Liu, Q., Yang, Y., Li, Z., Lu, Z., Zhang, H., Guo, T., and Liu, Q.
1234           (2020) Theoretical Research on Excited States: Ultraviolet and Fluorescence Spectra of
1235           Aromatic Amino Acids. *Interdiscip Sci* 12, 530–536

1236 106. Yammine, A., Gao, J., and Kwan, A. H. (2019) Tryptophan Fluorescence Quenching
1237           Assays for Measuring Protein-ligand Binding Affinities: Principles and a Practical Guide.
1238           *Bio Protoc* 9,

1239 107. Zheng, Z., Kawakami, K., Zhang, D., Negishi, L., Abomosallam, M., Asakura, T., Nagata,
1240           K., and Suzuki, M. (2021) Identification and functional analysis of cadmium-binding
1241           protein in the visceral mass of Crassostrea gigas. *Sci Rep* 11, 1–14

1242 108. Springfield, E., Willis, A., Merle, J., Mazlo, J., and Ngu-Schwemlein, M. (2021)
1243           Spectroscopic and Theoretical Studies of Hg(II) Complexation with Some Dicysteinyl
1244           Tetrapeptides. *Bioinorg Chem Appl* 2021, 9911474

1245 109. Crouse, H. F., Petrunak, E. M., Donovan, A. M., Merkle, A. C., Swartz, B. L., and Basu,
1246           S. (2011) Static and dynamic quenching of tryptophan fluorescence in various proteins by
1247           a chromium (III) complex. *Spectroscopy Letters* 44, 369–374

1248 110. Shrivastava, H. Y., and Nair, B. U. (2004) Fluorescence resonance energy transfer from
1249           tryptophan to a chromium(III) complex accompanied by non-specific cleavage of
1250           albumin: a step forward towards the development of a novel photoprotease. *J Inorg*
1251           *Biochem* 98, 991–994

1252 111. Liu, Y., Zhang, L., Liu, R., and Zhang, P. (2012) Spectroscopic identification of
1253           interactions of Pb 2+ with bovine serum albumin. *J Fluoresc* 22, 239–245

1254 112. Gen-hua Wu Study on the interaction of Pb2+ and bovine serum albumin by fluorescence.

1255 113. Luis, G., Rubio, C., Revert, C., Espinosa, A., González-Weller, D., Gutiérrez, A. J., and
1256           Hardisson, A. (2015) Dietary intake of metals from yogurts analyzed by inductively
1257           coupled plasma optical emission spectrometry (ICP-OES). *Journal of Food Composition*
1258           *and Analysis* 39, 48–54

1259 114. Wiltsche, H., and Wolfgang, M. (2020) Merits of microwave plasmas for optical emission
1260           spectrometry – characterization of an axially viewed microwave-sustained, inductively
1261           coupled, atmospheric-pressure plasma (MICAP). *J Anal At Spectrom* 35, 2369–2377

1262 115. Baraud, F., Zaiter, A., Porée, S., and Leleyter, L. (2020) New approach for determination
1263           of Cd, Cu, Cr, Ni, Pb, and Zn in sewage sludges, fired brick, and sediments using two
1264           analytical methods by microwave-induced plasma optical spectrometry and induced
1265           coupled plasma optical spectrometry. *SN Appl Sci* 2, 1–10

1266 116. Wegener Antje, Marc Wiesmann, Peter Luschas, Kerstin Vogel, and Matthias Pursch
1267           (2019) SEC-ICP-OES Hyphenation: Speciation and Quantification of
1268           Polydimethylsiloxanes at Trace levels. *Spectroscopy*, 38–46

117. Pandey, M., Talwar, S., Bose, S., and Pandey, A. K. (2018) Iron homeostasis in
Mycobacterium tuberculosis is essential for persistence. *Sci Rep* 8, 1–9

118. Tripathi, A., Anand, K., Das, M., O'Niel, R. A., Sabarinath, P. S., Thakur, C., Reddy, R.
R. L., Rajmani, R. S., Chandra, N., Laxman, S., and Singh, A. (2022) Mycobacterium
tuberculosis requires SufT for Fe-S cluster maturation, metabolism, and survival in vivo.
*PLoS Pathog* 18, e1010475

119. Saini, V., Farhana, A., Glasgow, J. N., and Steyn, A. J. C. (2012) Iron sulfur cluster
proteins and microbial regulation: implications for understanding tuberculosis. *Curr Opin*
*Chem Biol* 16, 45–53

120. Wang, P., Xu, G., Qin, L., Xu, Y., Li, Y., and Li, R. (2005) Cell-based biosensors and its
application in biomedicine. *Sens Actuators B Chem* 108, 576–584

121. Gupta, N., Renugopalakrishnan, V., Liepmann, D., Paulmurugan, R., and Malhotra, B. D.
(2019) Cell-based biosensors: Recent trends, challenges and future perspectives. *Biosens*
*Bioelectron* 141, 111435

122. Perler, F. B. (2002) InBase: the Intein Database. *Nucleic Acids Res* 30, 383–384

123. Yakovleva, L., and Shuman, S. (2012) Mycobacterium smegmatis SftH exemplifies a
distinctive clade of superfamily II DNA-dependent ATPases with 3′ to 5′ translocase and
helicase activities. *Nucleic Acids Res* 40, 7465–7475

124. Ordonez, H., and Shuman, S. (2013) Mycobacterium smegmatis Lhr Is a DNA-dependent
ATPase and a 3′-to-5′ DNA Translocase and Helicase That Prefers to Unwind 3′-Tailed
RNA:DNA Hybrids. *Journal of Biological Chemistry* 288, 14125–14134

125. Uson, M. L., Ordonez, H., and Shuman, S. (2015) Mycobacterium smegmatis HelY Is an
RNA-Activated ATPase/dATPase and 3′-to-5′ Helicase That Unwinds 3′-Tailed RNA
Duplexes and RNA:DNA Hybrids. *J Bacteriol* 197, 3057–3065

126. Rampersad, S. N. (2012) Multiple Applications of Alamar Blue as an Indicator of
Metabolic Function and Cellular Health in Cell Viability Bioassays. *Sensors* 12, 12347–
12360

127. O'Brien, J., Wilson, I., Orton, T., and Pognan, F. (2000) Investigation of the Alamar Blue
(resazurin) fluorescent dye for the assessment of mammalian cell cytotoxicity. *Eur J*
*Biochem* 267, 5421–5426

128. McMillan, J. E. (2013) *Principles of Analytical Validation* (Elsevier Inc.)

129. Howard, J. A., Kuznietsova, H., Dziubenko, N., Aigle, A., Natuzzi, M., Thomas, E.,
Lysenko, V., David, L., Brichart, T., Lux, F., and Tillement, O. (2023) Combating lead

and cadmium exposure with an orally administered chitosan-based chelating polymer. *Sci*
*Rep* 13, 1–14

130. Saleh, S. M., El-Tawil, O. S., Mahmoud, M. B., Abd El-Rahman, S. S., El-Saied, E. M.,
and Noshy, P. A. (2024) Do Nanoparticles of Calcium Disodium EDTA Minimize the
Toxic Effects of Cadmium in Female Rats? *Biol Trace Elem Res* 202, 2228–2240

131. Flora, S. J. S., and Pachauri, V. (2010) Chelation in Metal Intoxication. *Int J Environ Res*
*Public Health* 7, 2745–2788

132. Bethesda (2012) *LiverTox* (National Institute of Diabetes and Digestive and Kidney
Diseases)

133. George, G. N., Prince, R. C., Gailer, J., Buttigieg, G. A., Denton, M. B., Harris, H. H., and
Pickering, I. J. (2004) Mercury binding to the chelation therapy agents DMSA and DMPS
and the rational design of custom chelators for mercury. *Chem Res Toxicol* 17, 999–1006

134. Aaseth, J., Ajsuvakova, O. P., Skalny, A. V., Skalnaya, M. G., and Tinkov, A. A. (2018)
Chelator combination as therapeutic strategy in mercury and lead poisonings. *Coord Chem*
*Rev* 358, 1–12

135. Martí-Baena, Q., Pascuet-Fontanet, A., Berjaga-Buisan, T., Caravaca-Rodríguez, M.,
Puig-Costa-Jussà, J., Sanchez-Mejias, A., Ivančić, D., Mogas-Díez, S., Güell, M., and
Macia, J. (2024) Intein-mediated thyroid hormone biosensors: towards controlled delivery
of hormone therapy. *Frontiers in Systems Biology* 4, 1270071

136. Chahal, H. K., Dai, Y., Saini, A., Ayala-Castro, C., and Outten, F. W. (2009) The SufBCD
Fe−S Scaffold Complex Interacts with SufA for Fe−S Cluster Transfer. *Biochemistry* 48,
10644–10653

137. Zhang, L., Xiao, N., Pan, Y., Zheng, Y., Pan, Z., Luo, Z., Xu, X., and Liu, Y. (2010)
Binding and Inhibition of Copper Ions to RecA Inteins from Mycobacterium tuberculosis.
*Chemistry – A European Journal* 16, 4297–4306

138. Shaloam R, D. (2015) Preclinical Assessment of Low Doses of Cisplatin in the
Management of Acute Promyelocytic Leukemia. *Int J Cancer Res Mol Mech* 1,

139. Lennon, C. W., Wahl, D., Goetz, J. R., and Joel Weinberger, I. (2021) Reactive Chlorine
Species Reversibly Inhibit DnaB Protein Splicing in Mycobacteria. *Microbiol Spectr* 9,

140. Dillon, N. A., Lamont, E. A., Rather, M. A., and Baughn, A. D. (2025) Oxidative stress
drives potent bactericidal activity of pyrazinamide against Mycobacterium tuberculosis.
*Elife* 14,

141. Zhang, M., Gong, J., Lin, Y., and Barnes, P. F. (1998) Growth of Virulent and
AvirulentMycobacterium tuberculosis Strains in Human Macrophages. *Infect Immun* 66,
794–799

142. Revised National TB Control Programme Training Manual for Mycobacterium
tuberculosis

143. Davis, E. O., Sedgwick, S. G., and Colston, M. J. (1991) Novel structure of the recA locus
of Mycobacterium tuberculosis implies processing of the gene product. *J Bacteriol* 173,
5653–5662

144. Saves, I., Westrelin, F., Daffé, M., and Masson, J. M. (2001) Identification of the first
eubacterial endonuclease coded by an intein allele in the pps1 gene of mycobacteria.
*Nucleic Acids Res* 29, 4310–4318

145. Schneider, R. F., Hallstrom, K., DeMott, C., and McDonough, K. A. (2024) Conditional
protein splicing of the Mycobacterium tuberculosis RecA intein in its native host. *Sci Rep*
14, 1–14

1348

1349

1350

1351

1352

1353

1354

1355

1356

1357

1358

1359

1360

1361

1362

**i)**

|  | Zn²⁺ |  |  | Cu²⁺ |  |  | Pt⁴⁺ |  |  | Cd²⁺ | Hg²⁺ |  |

H37Ra — *M. sm* — H37Ra — *M. sm* — H37Ra — *M. sm* H37Ra H37Ra

| Zn 100 μM | Zn 2 mM | *M. Sm*+ Zn 100 μM | Cu 100 μM | Cu 2 mM | *M. Sm*+ Cu 100 μM | Pt 20 μM | Pt 1 mM | *M. Sm*+ Pt 20 μM | Cd 10 μM | Hg 25 μM | +ve Control |
| Zn 100 μM | Zn 2 mM | *M. Sm*+ Zn 100 μM | Cu 100 μM | Cu 2 mM | *M. Sm*+ Cu 100 μM | Pt 20 μM | Pt 1 mM | *M. Sm*+ Pt 20 μM | Cd 10 μM | Hg 25 μM | +ve Control |
| Zn 600 μM |  | *M. Sm*+ Zn 600 μM | Cu 600 μM |  | *M. Sm*+ Cu 600 μM | Pt 40 μM |  | *M. Sm*+ Pt 40 μM | Cd 25 μM | Hg 100 μM | +ve Control |
| Zn 600 μM |  | *M. Sm*+ Zn 600 μM | Cu 600 μM |  | *M. Sm*+ Cu 600 μM | Pt 40 μM |  | *M. Sm*+ Pt 40 μM | Cd 25 μM | Hg 100 μM | -ve Control |
| Zn 800 μM |  | *M. Sm*+ Zn 2 mM | Cu 800 μM |  | *M. Sm*+ Cu 2 mM | Pt 80 μM |  | *M. Sm*+ Pt 1 mM | Cd 100 μM | Hg 500 μM | -ve Control |
| Zn 800 μM |  | *M. Sm*+ Zn 2 mM | Cu 800 μM |  | *M. Sm*+ Cu 2 mM | Pt 80 μM |  | *M. Sm*+ Pt 1 mM | Cd 100 μM | Hg 500 μM | -ve Control |
| Zn 1 mM |  |  | Cu 1 mM |  |  | Pt 100 μM |  |  | Cd 2 mM | Hg 2 mM |  |
| Zn 1 mM |  |  | Cu 1 mM |  |  | Pt 100 μM |  |  | Cd 2 mM | Hg 2 mM |  |

+ve control

-ve control

**ii)**

|  | Zn²⁺ |  |  | Cu²⁺ |  |  | Pt⁴⁺ |  |  | Cd²⁺ | Hg²⁺ |

H37Ra — *M. sm* — H37Ra — *M. sm* — H37Ra — *M. sm* H37Ra H37Ra

+ve control

-ve control

**SufB Intein**

**(i)**

**Live cells**

**Functional SufB protein**

Splicing of *Mtb* SufB precursor protein within viable cells

**Alamar Reagent**

Mitochondrial reductase

- Cadmium (Cd²⁺)
- Mercury (Hg²⁺)
- TME

**Dead cells**

Splicing inhibition within non-viable cells

**(ii)**

**Alamar Reagent**

Mitochondrial reductase