1 **Designing a robust whole-cell biosensor platform for detection of toxic metals**
2 **using intein splicing inhibition of *Mycobacterium tuberculosis* SufB protein**
3

Ashwaria Mehra[1†], Ananya Nanda[1,2†], and Sasmita Nayak[1, 3*]

[1] School of Biotechnology, Kalinga Institute of Industrial Technology Deemed to be University,
Bhubaneswar, Odisha, India 751024.
[2]Toronto Metropolitan University, 350 Victoria St, Toronto, ON M5B 2K3, Canada
[3] Department of Microbiology, Kalinga Institute of Medical Sciences, Kalinga Institute of
Industrial Technology Deemed to be University, Bhubaneswar, Odisha, India 751024
Corresponding Author: sasmita.n@kiitbiotech.ac.in

12
13 [†]These authors have contributed equally to this work and share first authorship

14
Abstract:

Heavy metal contamination has emerged as a persistent environmental and public health challenge.

Although existing analytical methods provide high sensitivity, they often require complex

instrumentation with limited on-site applications, and generate hazardous waste, raising

environmental concerns. At present, whole-cell biosensors provide an empirical alternative,

utilizing cellular machinery for sensitive detection of metals. In a relevant context, intein-mediated

protein splicing, a spontaneous post-translational process, can be adapted for biosensing by

coupling metal-dependent splicing inhibition with whole-cell sensing units. Current research

elucidates how toxic metal ions, including cadmium (Cd^{2+}), mercury (Hg^{2+}), chromium (Cr^{3+}), and

lead (Pb^{2+}), influence the splicing and N-terminal cleavage reactions of *Mycobacterium*

tuberculosis (*Mtb*) SufB precursor protein. While Cd^{2+} and Hg^{2+} significantly attenuated the

splicing activity of *Mtb* SufB precursor protein over a concentration range of 25 μM to 2 mM, Pb^{2+}

and Cr^{3+} failed to do so. Next, an innovative biosensor platform was designed to detect these metal

ions by utilizing splicing inhibition in metal-sensitive *Mtb* SufB precursor protein within

attenuated *Mtb* (H37Ra) cells. Metal-induced splicing inhibition led to loss of viability in H37Ra

indicator cells, while specificity of this detection system was confirmed by adding respective metal

chelators. Multiplexing and quantitative efficacy of the designed biosensor was evaluated by

including known splicing inhibitors like Cu^{2+}, Zn^{2+}, and Pt^{4+} over various concentration range

alongside Cd^{2+} and Hg^{2+}. The developed biosensor offers a cost-effective, user-friendly, specific,

and sensitive assay that does not require advanced instrumentation facility. It can be employed in

standard biological laboratories utilizing whole-cell microorganisms harbouring precursor proteins

with metal-sensing intein sequences. This approach holds potential for multiplexing metal

screening in environmental and industrial effluents, significantly enhancing toxic metal

monitoring capabilities.

Keywords: Whole-cell biosensor, heavy metal ion detection, *Mycobacterium tuberculosis* SufB*,

intein-based biosensor, protein splicing inhibition.

Heavy metal contamination has emerged as one of the most persistent environmental and public

health challenges of the modern era. Over 35 metals have been identified in their ore forms within

the Earth's crust, many of which are essential nutrients. Among these, 23 metals are classified as

heavy metals due to their high density (>5 g/cm³) and atomic weights (>40.0), indicating their

potential for environmental toxicity.[1–3] Human activities have significantly disrupted natural

geochemical cycles, causing bioaccumulation and biomagnification through contaminated food,

water, or direct exposure, thereby posing serious health risks.[3–7]

Current analytical methods, such as electrothermal atomic absorption spectrometry (ETAAS),

inductively coupled plasma-optical emission spectrometry (ICP-OES), high-pressure liquid

chromatography combined with ICP-mass spectrometry (HPLC-ICP-MS), hydride generation

atomic absorption spectrometry (HGAAS), and X-ray fluorescence spectrometry (XRF), offer

excellent sensitivity for detecting metals. Nonetheless, these methods involve complex

instrumentation, require skilled operators, generate hazardous waste, contributing further to

environmental pollution.[8–11] To overcome these limitations, biosensors have emerged as a

promising alternative offering affordability, and long-term operational stability.[8] According to the

International Union of Pure and Applied Chemistry (IUPAC), biosensors integrate biological

recognition elements directly with transducers to provide precise quantitative data.[12,13] Although

enzyme-based electrochemical biosensors have been widely explored, their broader

implementation is limited due to poor electron transfer efficiency, and instability, limiting their

commercial reusability.[14–20] In comparison, whole-cell biosensors present a more practicable

approach by using live cell sensing units, with advantages such as stable signal generation, and

scalability. Despite these strengths, declining cell viability causing delayed response time remain

challenging.[21–26] Addressing these shortcomings, intein-mediated protein engineering techniques

open up new avenues for creating innovative cell-based sensors.[27–30]

Inteins are protein splicing elements within precursor proteins that can self-excise via multi-step

nucleophilic displacement reactions, to generate active proteins.[31,32] This is a post-translational

process that does not require energy sources or cofactors, hence inteins can be seamlessly

integrated into biosensor units minimizing interference with sensor cell functionality.[27]

Furthermore, intein splicing can be regulated via conditional protein splicing (CPS), in presence

of small molecules such as metal ions and metal-based nanoparticles (Table 1).[33–37] For instance,

a recent study from our group has examined the splicing and N-terminal cleavage activity of

Mycobacterium tuberculosis (*Mtb*) SufB precursor protein in the presence of metal ions such as

Cu^{2+}, Zn^{2+}, Pt^{4+}, and Fe^{3+}/Fe^{2+}.[38] Prior studies have also reported Zn^{2+} mediated splicing regulation

of PI-*Sce* VMA intein, *Mtb* RecA intein, and *Ssp* PCC6803 DnaE split intein.[39,40] Effects of metal

ions, such as $Cu^{+/2+}$ and CisPt, have been evaluated on *Mtb* RecA, *Cryptococcus neoformans* Prp8,

and *Mycobacterium smegmatis* DnaBi1 intein splicing.[35,36,39,41–46]

Table 1: Overview of intein-harboring organisms and reported modulation of splicing activity by
metal ions

Name of Organisms		Intein containing precursor proteins	Metal ions	References
Bacteria	*Mycobacterium tuberculosis*	RecA	Zn^{2+} Cd^{2+} Co^{2+} Ni^{2+} Cu^+/Cu^{2+} CisPt	40,42,47,48
	Mycobacterium tuberculosis	SufB	Zn^{2+} Cu^{2+} Pt^{4+} $Fe^{2+/3+}$	38
	Mycobacterium smegmatis	DnaB-intein1 (DnaBi1)	Zn^{2+}	44
	Synechocystis sp. PCC6803	*Ssp* DnaE	Zn^{2+} Cd^{2+}	40
	Oscillatoria limnetica	split DnaE intein	Cu^{2+}	49
	Thermosynechococcus vulcanus	DnaE	Cu^{2+}	49
	Nostoc sp. PCC7120	DnaB	Cu^{2+}	49
Fungi	*Saccharomyces cerevisiae*	PI-*Sce*I VMA	Zn^{2+}	39
	Cryptococcus neoformans	Prp8	Zn^{2+} Cu^{2+} CisPt	36,46

Cryptococcus gattii	Prp8	CisPt	46

Conditional splicing activity of intein-containing proteins has been widely used to develop

biosensors with unique functionalities.[27,50–52] However*,* a versatile intein-based metal-sensing

biosensor, especially for toxic metals ions, is yet to be elucidated. Current research explores how

toxic metal ions, like cadmium (Cd^{2+}), mercury (Hg^{2+}), chromium (Cr^{3+}), and lead (Pb^{2+}), affect

the splicing and N-terminal cleavage reactions of *Mtb* SufB precursor protein. Furthermore, a

whole-cell biosensor system using mycobacterial cells has been developed and evaluated to detect

the presence of toxic metal ions via splicing regulation in metal-sensitive precursor protein, *Mtb*

SufB.

SUF protein complex constitutes a unique pathway for Fe-S cluster assembly in mycobacteria, and

its activity is dependent on the splicing of *Mtb* SufB precursor, producing functional SufB

protein.[53] Thus, splicing inhibition which blocks the generation of active SufB, can impair

mycobacterial growth and viability.[54] In the current work, *in vitro* studies demonstrated significant

inhibition of *Mtb* SufB precursor protein splicing in presence of toxic metal ions such as Cd^{2+}, and

Hg^{2+} over a concentration range of 25 μM to 2 mM. Further, time-dependent assay revealed that

Cd^{2+} and Hg^{2+} could inhibit *Mtb* SufB intein splicing as early as 10 minutes following exposure to

the metal ions. Building on this observation, an *in vitro* biosensor platform was designed using

H37Ra (an attenuated *Mtb* strain) as metal-indicator cells. Alamar Blue assay performed in a 96-

well microtiter plate, detected loss of cell viability in SufB-containing H37Ra cells in presence of

toxic metal ions, Cd^{2+} and Hg^{2+}. In contrast, *Mycobacterium smegmatis* (*M. sm*), which expresses

an intein-less SufB protein, persisted as healthy, viable cells under similar experimental conditions.

Hence, current study highlights the sensitivity of H37Ra-based biosensor platform to detect toxic

metal ions via inhibiting the generation of essential protein SufB in mycobacteria. Moreover, the

specificity of the biosensor was demonstrated by addition of respective metal chelators which

protected mycobacterial cell viability via blocking the splicing-inhibitory roles of the metal ions.

Since, Zn^{2+}, Cu^{2+}, and Pt^{4+} are also known inhibitors of *Mtb* SufB splicing, multiplexing and

quantitative potential of this platform was validated using various concentrations of these metal

ions alongside Cd^{2+} and Hg^{2+}. Thus, this work opens new avenues for developing a reliable and

sensitive *in vitro* biosensor platform using whole-cell mycobacteria as biological indicator for

toxic metal ions as well as trace metal elements (TME) in the environmental samples. The efficacy

of this system lies in its cost-effectiveness, user-friendly design, high specificity, and sensitivity,

enabling multiplexed metal detection. It is particularly useful as it can detect toxic metal ions at

low concentration (25 µM) within just 10 minutes. Consequently, it can be easily employed in

standard biological laboratories equipped with organisms harboring metal-sensitive intein-

carrying precursor proteins.

Materials and Methods:

Protein overexpression and purification [55–57]

Plasmids containing the genes for *Mtb* SufB precursor and *Mtb* SufB splicing inactive (SI) double

mutant (Cys1Ala/Asn359Ala) (splicing negative control),[55] both carrying an N-terminal 6X (His)

tag, were borrowed from Belfort Lab, SUNY, Albany, USA. The test plasmids were transformed

into BL21(DE3) *E. coli* cells (NEB, C2527I), followed by IPTG (500 µM, sigma 367-93-1)

induced protein over-expression. Thereafter, lysis buffer (20 mM sodium phosphate, 0.5 M NaCl,

pH 7.4) was used to resuspend the bacterial cell pellet and cells were lysed via tip sonicator (Sonics

vibra cell VCX-130). The cell lysate was centrifuged at 16500×g for 20 minutes at 4 °C, and the

supernatant was discarded. The collected inclusion body (IB) material was then washed three times

in lysis buffer via centrifugation (16500×g for 20 minutes at 4 °C). Finally, the IB material was

solubilized for protein extraction using 8 M urea (Merck, 1084870500) buffer (20 mM sodium

phosphate, 0.5 M NaCl, 8M urea, 10 mM of imidazole, pH 7.4 (MP–biochemicals-288-32-4) and

subsequently centrifuged at 16,500×g for 20 minutes at 4 °C to collect the supernatant. Then,

solubilized proteins were purified by Ni-NTA affinity column (Ni-NTA His trap, HP GE

healthcare life sciences, 17524802).[35,40,48,58,59] Before applying the samples, the columns were

equilibrated with binding buffer (20 mM sodium phosphate, 0.5 M NaCl, 8 M urea, 10 mM

imidazole, pH 7.4). Following sample loading, the columns were washed multiple times (5 CV)

using wash buffer (20 mM sodium phosphate, 0.5 M NaCl, 8 M urea, 40 mM imidazole, pH 7.4).

Subsequently, the test proteins were eluted as purified fractions using elution buffer (20 mM

sodium phosphate, 0.5 M NaCl, 8 M urea, 500 mM imidazole, pH 7.4). Protein quantification was

performed using Bradford reagent.

Refolding assay for Mtb SufB precursor protein

Denatured protein (2.5 µM) in elution buffer was renatured in refolding buffer (20 mM sodium

phosphate, 0.5 M NaCl, 0.5 M L-Arginine) containing 2 mM TCEP-HCl (sigma-51805-45-9) for

4 hours at 25 ºC in the absence and presence of toxic metal ions.[40,58,59] Gradient assay for toxic

metal ions [$CdCl_2$ (SRL-99643), $HgCl_2$ (SRL-25699), $CrCl_3$ (CDH-025702), $PbCl_2$ (SRL-69789)]

was performed using varying concentrations of toxic metal ions (2.5 µM to 2 mM). 0h sample was

collected before initiation of refolding, and the reaction was abruptly blocked by the addition of

loading dye (0.1% bromophenol blue, 50% glycerol, 5% β-mercaptoethanol, 10% SDS, tris 6.8),

followed by rapid freezing at −20 ºC. After 4h of protein renaturation, the protein samples were

resolved through 4~10% gradient SDS-PAGE, followed by staining with Coomassie brilliant blue

R-250 (93473, SRL), and densitometric analysis was performed using GelQuant.NET biochemical

solutions software. The splicing and cleavage efficiencies were calculated using the following

formulae:

$$\% \: of \: splicing = \frac{Total \: splicing \: products \: (LE + I)}{Total \: protein \: (P + LE + I)} \: X \: 100$$

$$\% \: of \: N - terminal \: cleavage = \frac{Total \: N - terminal \: cleavage \: products \: (NC + NE)}{Total \: protein \: (P + NC + NE)} X \: 100$$

[P: precursor protein, LE: Ligated exteins, I: Intein, NC: N-terminal cleavage product, NE: N-

extein]

Baseline correction was done using the 0 hours splicing and cleavage value(s). The results were

analyzed using one-way ANOVA and plotted using GraphPad Prism version 8 for Windows,

GraphPad Software, San Diego, California, USA, www.graphpad.com.

Immunoblotting to identify splicing and N-terminal cleavage protein products[38]

Protein immunodetection was conducted to identify of splicing and N-terminal cleavage reaction

products, using anti-6X(His) tag antibodies (Invitrogen, LOT 1902132). After resolution through

4~10% gradient SDS-PAGE, the protein products were transferred onto a PVDF membrane

(Millipore, IPVH 00010) at 50V for 2 hours at 4 °C. The blot was then blocked with 5% skim milk

for 2 hours at room temperature, followed by two washes with 1X TBST. It was incubated with

primary anti-6X(His) tag (abgenex -32-6116) at a dilution of 1:5000 at 4 °C for 16 hours. After

washing with 1X TBST, the blot was treated with a secondary antibody (anti-mouse; abgenex 11-

301) at a dilution of 1:8000 for 2 hours at 37 °C. The N-extein band was detected using a lower

dilution of the primary antibody (1:2500) and the secondary antibody (1:6000). Finally, the blot

was developed using Enhanced Chemiluminescence substrate (Abcam, ab65628) to visualize the

protein bands, and images were captured using in-house ImageQuant™ LAS 500 facility (GE

Healthcare). Blots were also developed, using 3, 3'-diaminobenzidine (DAB) (SRL-94524) for

visualization of the protein bands.

Spectroscopic and Elemental Profiling of Protein-Metal Interactions

UV-Visible spectroscopy [60–62]

Denatured proteins were renatured _in vitro_ in the presence and absence of toxic metal ions at their

respective concentrations for 4 hours, as specified earlier. Subsequently, the treated samples were

subjected to UV-visible spectroscopy (Cary 100 UV-Vis Agilent Technology) scanning in the

visible region spanning 200-800nm. To rule out the interference from buffer-metal interactions,

the refolding buffer (20 mM sodium phosphate, 0.5 M NaCl, 0.5 M L-Arginine) was incubated

with metal ions (Cd^{2+}, Hg^{2+}, Cr^{3+}, and Pb^{2+}) at their respective concentrations under similar

experimental conditions. This was used as a blank for baseline correction and was scanned

alongside each sample. Absorbance versus wavelength plots were generated and analyzed using

Origin (Pro) 8.5 software (Version 8.5, OriginLab Corporation, Northampton, MA, USA) to

interpret the results.

Tryptophan Fluorescence Spectroscopy

The purified denatured protein (2.5 μM) was allowed to renature in the presence and absence of

metals ions (25 μM $CdCl_2$, 500 μM $HgCl_2$, 2 mM $CrCl_3$, and 2 mM $PbCl_2$) under identical

experimental conditions. Following 4 hours of refolding, the test samples were analyzed via

Fluorescence spectrometry (Cary Eclipse, Agilent Technologies) over a wavelength range of 300-

600 nm and at an excitation wavelength of 280 nm.[63] To rule out the buffer-metal interactions, the

refolding buffer (20 mM sodium phosphate, 0.5 M NaCl, 0.5 M Arginine) was incubated with

metal ions at their respective concentrations at 25 °C for 4 hours and then scanned alongside each

treated sample. This served as the blank for baseline correction. The tryptophan fluorescence

spectra for the test samples were recorded and analyzed by plotting intensity vs. wavelength using

Origin (Pro) 8.5, Version 8.5, from OriginLab Corporation, Northampton, MA, USA.

Inductively coupled plasma-optical emission spectroscopy (ICP-OES) analysis

Following *in vitro* refolding of *Mtb* SufB precursor in the presence of test metal ions, the protein

products were resolved through 4~10% SDS-PAGE. The resulting protein bands were visualized

with Coomassie brilliant blue R-250. Subsequently, the gel bands for unspliced precursor protein,

ligated extein, and intein were excised from the gel for further analysis. Each gel band was then

treated with concentrated H_2SO_4 and 30% H_2O_2. Complete dissolution of each sample was

achieved by incubating the gel pieces in a 3:1 mixture of concentrated HCl and concentrated HNO_3

at 80°C for 4 hours.[64,65] The final volume of the digested solution was adjusted to 50 ml with

deionized water and then analyzed by ICP-OES (Perkin-Elner, optima 8300). Prior to analyzing

the test samples, respective toxic metal standards were calibrated in the system, and the

concentration of bound metal ions was quantified from the standard curve.

In vitro whole-cell (H37Ra) toxic metal biosensor

Different mycobacterial cells [*Mycobacterium tuberculosis* H37Ra strain (ATCC 25177),

Mycobacterium smegmatis (negative control for SufB intein-less micro-organism, ATCC

700084)] were cultured in 7H9 (Middlebrook 7H9 broth medium, M198) supplemented with

OADC (Oleic acid, dextrose, and catalase, Himedia-FD018) and allowed to grow separately for 2-

3 weeks at 37 ^{0}C. CFU calculations for individual cultures were made by matching the turbidity

of the inoculated mycobacterial culture with the McFarland standard (Himedia- R092) against a

black background.[66] Different bacterial cells were diluted in sterile saline to a concentration of

10^8 CFU/ml. H37Ra cells were incubated with varied concentrations of Cd^{2+} (2.5 µM- 2 mM) and

Hg^{2+} (25 µM- 2 mM) in 96-well microtiter plate. Since Cr^{3+} and Pb^{2+} over a test concentration

range of 25 µM- 2 mM did not exert any effect on *Mtb* SufB splicing, these metal ions were not

tested via the *in vitro* biosensor platform. The plates were sealed and incubated at 37 ºC for 4

days. Then, 10% (v/v) solution of Alamar Blue reagent (SRL-42650) was added to each well,

followed by incubation at 37 ºC for 12 hrs.[67] Untreated mycobacterial cells grown in 7H9

(supplemented with OADC) media were used as a negative control, and Alamar blue reagent alone

in 7H9 (supplemented with OADC) media served as a positive control. SufB intein-less organism,

Mycobacterium smegmatis, was also evaluated under similar conditions and treated as an

additional experimental control. After 12 hours of incubation, varied colour changes were noted

in different wells, where blue colour indicated dead cells with no viable cell growth, and pink

colour correlated to metabolically active live cells. In a similar experimental setup, metal chelators

specific for Cd^{2+} (Edetate Calcium Disodium, ECD, Sigma-PHR1510) and Hg^{2+} (meso-2,3-

Dimercaptosuccinic acid, DMSA, Sigma-D7881) were added in a ratio of 1:1 (metal: chelator) to

determine the specificity of the designed *in vitro* Biosensor platform.

Quantitative evaluation of *(H37Ra)-based whole cell toxic metal biosensor*

Next, a standard plot was generated by measuring the absorbance of the respective samples at

570 nm, with 600 nm as a reference wavelength for quantitative analysis.[67] A wide range of test

metal samples (Cd^{2+}, Hg^{2+}) of known concentration were prepared and loaded onto the 96-well

plate and Alamar Blue reagent was added under similar experimental conditions. The absorbance

of test metal samples was calculated by subtracting the reference absorbance value at 600nm from

the absorbance measured at 570nm [$OD_{test} = OD_{570} - OD_{600}$]. Untreated mycobacterial cells grown

in 7H9 (supplemented with OADC) media were used as a control for normalization. Using the

normalized values, XY scattered plot was obtained where the X axis represented the known

concentrations of the test metal ions and the Y axis denoted corresponding absorbance values.

Next, to assess the applicability of the standard plot, four test samples (Unknown 1, Unknown 2,

Unknown 3, and Unknown 4) containing varying concentrations of $CdCl_2$ and $HgCl_2$ were

analyzed under identical experimental conditions. Absorbance values were extrapolated onto the

standard curve to calculate the unknown metal ion concentrations, thereby validating the

sensitivity and accuracy of the biosensor platform.

Multiplexing evaluation of (H37Ra)-based whole cell metal biosensor

Bacterial cell suspension containing 10^8 CFU/ml of *Mycobacterium tuberculosis* H37Ra were

prepared in sterile saline. Cells were incubated with a varied concentration range of $ZnCl_2$ (100

μM- 2 mM), $CuCl_2$ (100 μM- 2 mM), $PtCl_4$ (20 μM- 1 mM), $CdCl_2$ (10 μM- 2 mM), and $HgCl_2$

(25 μM- 2 mM), in 96-well microtiter plates. The plates were sealed and incubated at 37 ºC for 4

days under static conditions. Then, 10% (v/v) solution of Alamar Blue reagent (SRL-42650) was

added to each well, followed by incubation at 37 ºC for 12 hrs. Untreated H37Ra cells grown in

Middlebrook 7H9 supplemented with OADC media were used as a negative control, while Alamar

blue reagent incubated in media alone served as a positive control. SufB intein-less organism,

Mycobacterium smegmatis, was treated with test metal ions under similar conditions and used as

negative experimental control to assess SufB intein-specific response. Post 12 hours of incubation,

colour change was noted in the wells, where blue colour indicated dead cells with no viability, and

pink colour indicated healthy viable cells.

To further ascertain the specificity of the biosensor platform, respective metal-chelating agents

were added in separate experimental setup, using a 1:1 molar ratio of metal ions to chelator.

Cadmium-specific chelator Edetate Calcium Disodium (ECD; Sigma-PHR1510), mercury-

specific chelator meso-2,3-Dimercaptosuccinic acid (DMSA; Sigma-D7881), and a broad-

spectrum chelator Ethylenediaminetetraacetic acid (EDTA; HiMedia-MB011), were co-incubated

with the respective metal ions under identical experimental conditions. Cell viability was evaluated

using Alamar Blue reagent as described earlier.

Results and Discussion

Toxic metal ions inhibited Mtb SufB precursor protein splicing

Environmental concentrations of toxic metal ions such as cadmium (Cd^{2+}), mercury (Hg^{2+}),

chromium (Cr^{3+}) and lead (Pb^{2+}) have been reported to vary widely. Cd^{2+} concentrations typically

ranges   from 1 to 10 ppm (approximately 5 µM to 55 µM), Hg^{2+} from 10 to 700 ppm

(approximately 4 µM to 3 mM), Cr^{3+} from 5 ppm- 140 ppm (approximately 31 µM – 885 µM) and

Pb^{2+} levels from 10 to 50 ppm (approximately 36 µM to 180 µM).[68–83] Earlier studies have shown

that metal ion concentrations between 40 µM to 2 mM effectively inhibit intein splicing.[38,84,85]

Hence, an *in vitro* gradient assay was performed to evaluate the sensitivity of *Mtb* SufB precursor

protein over a selected concentration range of toxic metals ions (2.5 µM to 2 mM) (Figure 1, and

Figures S2-S4, Supporting Information). *Mtb* SufB precursor protein (95.98 KDa) follows a

canonical intein splicing pathway yielding 40.2 KDa SufB intein and 55.7 KDa ligated exteins

(functional SufB protein) as illustrated in Figure S1, Supporting Information.[55] N-terminal

cleavage products (65.26 KDa N-terminal cleavage product, 29.9 KDa N-extein) and C-terminal

cleavage products (70.12 KDa C-terminal cleavage product, 25.76 KDa C-extein) are the

byproducts of *Mtb* SufB splicing reaction. [55]

Purified *Mtb* SufB precursor protein was refolded in presence of test metal ions and then resolved

through 4~10% gradient SDS PAGE (Figure 1A). Cd^{2+} at 25 µM, and Hg^{2+} at 500 µM of minimum

effective concentration (MEC) displayed significant attenuation of splicing and N-terminal

cleavage reactions. There was a noticeable accumulation of SufB precursor protein, along with

substantial reduction in the splicing products (ligated exteins and intein). Further, Cd^{2+}, and Hg^{2+}

at their respective MEC values and higher, induced a near-complete inhibition of N-terminal

cleavage reaction and the products (N-terminal cleavage product and N-extein) (Figure 1A).

However, Cr^{3+} and Pb^{2+} failed to induce any effect on *Mtb* SufB splicing or cleavage reactions

across the test concentration range (Figure 1A). Resultant protein products were further confirmed

via western blot using anti-6X(His) tag antibody (Figure S2, Supporting Information).

Densitometric analysis revealed a 12.52-fold (p<0.0001) reduction in splicing efficiency and an

18.7-fold (p<0.0001) reduction in N-terminal cleavage efficiency with 25 µM Cd^{2+} [Figure 1B (i)

and C (i)]. Similarly, treatment with 500 µM Hg^{2+} resulted in an 8-fold (p < 0.0001) reduction in

splicing and a 7.8-fold (p < 0.0001) decrease in N-terminal cleavage efficiency [Figure 1B (ii) and

1C (ii)]. No significant alterations in SufB splicing or N-terminal cleavage products were obtained

in presence of Cr^{3+} or Pb^{2+} (Figure 1B and 1C). All the test metal ions at a lower concentration

range (2.5 µM- 10 µM) failed to exert any significant effect on the SufB splicing reactions (Figures

S3 and S4, Supporting Information). In order to rule out the possibility of precursor protein

degradation by test metal ions, *in vitro* refolding assays were also conducted using splicing inactive

(SI) *Mtb* SufB double mutant (C1A/N359A) under similar experimental conditions (25 µM Cd^{2+},

500 µM Hg^{2+}, 2 mM Cr^{3+}, and 2 mM Pb^{2+}). Since Cr^{3+}, and Pb^{2+} showed no regulatory effects,

highest test concentration range for both were included in the above study. SDS-PAGE and

subsequent western blot analysis using anti-6X(His) tag antibodies confirmed the absence of

protein degradation products (Figures S2B, Supporting Information).

Figure 1. Gradient assay to determine the minimum effective concentration (MEC) of toxic metal ions on splicing and N-terminal cleavage reactions of *Mtb* SufB precursor protein. (A) Following *in vitro* refolding of *Mtb* SufB precursor protein in the presence of (i) $CdCl_2$, (ii) $HgCl_2$, (iii) $CrCl_3$, and (iv) $PbCl_2$ at varied concentrations (25 μM-2 mM), resultant products were resolved through 4~10% gradient SDS-PAGE. Lane 1: Protein ladder; Lane 2 (T_0): splicing and N-terminal cleavage reactions at 0 h; Lane 3 (UT): splicing and N-terminal cleavage products of untreated protein sample; Lanes 4-9: splicing and N-terminal cleavage products resulting in the presence of different toxic metal ions at different concentrations (25 μM-2 mM). Lane 10 (SI): splicing inactive SufB double mutant (C1A/N359A) is used as a negative control for splicing. **(B) and (C)**

Bar graphs demonstrating splicing and N-terminal cleavage efficiency of *Mtb* SufB precursor protein in the presence and absence of different toxic metal ions. A statistically significant difference in splicing and N-terminal cleavage reaction was observed for B (i) and C (i) CdCl$_2$, starting with 25 µM, and B (ii) and C (ii) HgCl$_2$ at 500 µM concentrations, respectively. For CrCl$_3$ [B (iii) and C (iii)], and PbCl$_2$ [B (iv) and C (iv)], no significant difference in splicing and N-terminal cleavage reaction were observed between untreated and metal-treated samples. Graphs were plotted after densitometric analyses using GelQuant.Net biochemical solutions. All the experiments were performed in triplicate, and error bars represent (±1) SEM from 3 independent sets of experiments. The data shown in Figures 1B, and 1C are extracted from Figure 1A. M: Prestained protein molecular weight ladder, P: Precursor, NC: N-terminal cleavage product, LE: Ligated extein, I: intein, NE: N-extein.

Further, a time-dependent splicing assay was performed to examine *Mtb* SufB splicing and cleavage activity over a period of 10 min- 4 hr., elucidating remarkable inhibitory roles of Cd^{2+} (25 µM) and Hg^{2+} (500 µM) as early as 10 minutes (Figure S5, Supporting information). Next, to evaluate the possible physical interaction between *Mtb* SufB protein and the metal ions, a series of experiments were conducted, as shown in the upcoming section. For the said interaction study, the MEC for Cd^{2+}, Hg^{2+} (25 µM, 500 µM, respectively), which inhibited *Mtb* SufB splicing was selected. Since, Cr^{3+}, and Pb^{2+} did not exhibit any such regulatory activity, highest value of the test concentration range (2 mM) was chosen to examine possible interaction with SufB protein.

Protein~ metal interaction study

UV-Visible spectroscopic analysis

Biomolecules such as proteins exhibit a strong affinity for metal ions, which can critically influence their structure and function.[86] Prior studies have demonstrated that toxic metal ions such as Cd^{2+}, Hg^{2+}, Cr^{3+}, and Pb^{2+} can interact with calcium and zinc-binding sites in proteins, thereby influencing their biological activity.[87,88] UV-visible spectroscopic analysis was conducted to investigate the interaction between the *Mtb* SufB precursor protein and toxic metal ions under treated and untreated conditions. Absorbance spectra for *Mtb* SufB protein were recorded over

200-800 nm range at 25 °C (Figure S6, Supporting Information). A characteristic absorbance peak

near 280 nm was detected, primarily attributable to aromatic residues such as tryptophan and

tyrosine.[89,90] Upon treatment with Cd^{2+}, an increase in absorbance intensity was noted, suggesting

an alteration in conformation due to metal coordination (Figure S6 (i), Supporting Information).

Previous studies have mentioned that enhanced absorbance may result from the interaction

between Cd^{2+} and electron-donating groups such as amide, imidazole, and amine.[61,91,92] Likewise,

samples treated with Hg^{2+} also exhibited an increase in absorbance intensity (Figure S6 (ii),

Supporting Information). This enhancement is likely attributed to the strong affinity of Hg^{2+} for

nucleophilic functional groups, particularly the sulfhydryl (–SH) groups of cysteine residues.[93,94]

Although there were sharp increments in absorbance peak following interaction of Cr^{3+} and Pb^{2+}

with *Mtb* SufB, they failed to exert any effects on SufB splicing reactions as shown earlier (Figure

1). Moreover, a strong buffer-Pb^{2+} interaction was observed from the results, which may have

weakened likely interaction with the protein. Although prior works have shown interaction of Cr^{3+},

and Pb^{2+} with thiol, glutathione groups within protein structure including oxygen, nitrogen, and

sulfur containing amino acids, [7,95,96] current work could not detect any effects of these metal ions

on *Mtb* SufB splicing activity (Figure S6 (iii and iv), Supporting Information). To further

substantiate such interactions, tryptophan fluorescence spectroscopy was performed, as explained

in the forthcoming section.

Tryptophan Fluorescence spectroscopy reveals metal-induced conformational changes in Mtb

SufB

Tryptophan fluorescence spectroscopy is a widely utilized technique for monitoring protein

conformational alterations and assessing their local structural dynamics upon interaction with

ligands or other macromolecules.[97] Among the intrinsic fluorophores in proteins, aromatic amino

acids—tryptophan (Trp), tyrosine (Tyr), and phenylalanine (Phe)—contribute to fluorescence,

with tryptophan exhibiting the highest molar absorptivity (1614 cm^{-1}/M) and a characteristic

emission peak typically in the range of 340–350 nm.[98] The indole ring of Trp is primarily

responsible for this emission and dominates the fluorescence profile over Tyr and Phe. Changes in

the microenvironment of tryptophan residues, such as those caused by ligand binding, protein

folding, or metal interaction, can lead to measurable shifts in fluorescence intensity and/or

emission maxima.[97,99]

Altogether, 13 Trp residues are present, spanning the entire sequence of *Mtb* SufB precursor

protein, with 5 Trp in N-extein, 5 Trp in Intein, and 3 Trp in the C-extein region.[38] Upon treatment

with Cd^{2+} (25 µM), an increase in fluorescence intensity was observed (Figure S7 (i), Supporting

Information), likely attributable to conformational rearrangements arising from interactions

between Cd^{2+} and functional groups such as amide, imidazole, and amine moieties within the

protein matrix.[61,91,92] These interactions may have led to reduced solvent exposure of Trp residues

or stabilized a more rigid protein conformation, which can enhance fluorescence emission. [61,100]

In contrast, Hg^{2+} (500 µM) treated samples displayed a significant quenching of Trp fluorescence

(Figure S7 (ii), Supporting Information). This effect is consistent with previous studies indicating

that mercury ions interact strongly with aromatic systems through cation–π interactions,

particularly with the indole ring of tryptophan, thereby reducing fluorescence emission.[101] Cr^{3+}

(2mM) exposure resulted in a slight increase in fluorescence intensity (Figure S7 (iii), Supporting

Information), potentially due to the preferential binding of Cr^{3+} to carboxyl groups, which may

induce localized structural stabilization without extensive perturbation of the overall tertiary

structure.[102,103] In case of Pb^{2+} (2 mM), a decrease in fluorescence intensity was observed (Figure

S7 (iv), Supporting Information). This observation is supported by previous literature, where Pb^{2+}

interacts with hydroxy, amino, and sulfhydryl groups, leading to partial protein unfolding and

increased solvent accessibility of Trp residues, thereby promoting fluorescence quenching.[104,105]

Although Cr^{3+}, Pb^{2+} have displayed possible alteration in protein conformation, they failed to

regulate *Mtb* SufB splicing and cleavage reactions, likely due to lack of any effect on protein active

site. Next, ICP-OES analysis was conducted to further ascertain potential interaction between test

metal ions and protein backbone, as described in the following section.

ICP-OES detects metal ions bound to SufB precursor and spliced out protein fractions

ICP-OES is a sensitive analytical technique capable of detecting trace metal concentrations by

measuring the characteristic emission of photons from the excited atoms and ions as they return to

their ground state.[106] During *in vitro* protein refolding experiment, specific concentrations of Cd^{2+}

(25 μM = 4.583 μg/ml), Cd^{2+} (500 μM = 91.66 μg/ml), Hg^{+2} (500 μM = 135.76 μg/ml), Cr^{+3} (2

mM = 361.72 μg/ml), and Pb^{+2} (2 mM = 556.2 μg/ml) were added to the respective reaction

mixture. ICP-OES analysis of the metal-bound protein fractions revealed detectable levels of Cd^{2+}

(2 ppb ~ 0.002 μg/ml), Hg^{2+} (5 ppb, ~0.005 μg/ml), and Cr^{3+} (80 ppb, ~0.08 μg/ml) bound to the

unspliced *Mtb* SufB precursor protein (Table 2). These findings indicate a direct and quantifiable

physical interaction between the toxic metal ions and the *Mtb* SufB precursor protein, which may

underlie the observed inhibition of splicing and N-terminal cleavage reactions. Further, measurable

interactions were observed between the test metal ions and various splicing products (ligated extein

and intein) (Table 2). Cd^{2+} addition to SufB protein sample at 25 μM concentration yielded

negative results, while increasing the concentration to 500 μM led to positive data. At lower

concentrations of Cd^{2+} (25 μM), the metal content in the protein fraction was possibly below the

instrument's limit of detection (LOD) and limit of quantification (LOQ), resulting in negative

readings. Although ICP-OES is capable of detecting metal concentrations at the ppb level, practical

detection limits are influenced by matrix complexity and background interference, often resulting

in quantification thresholds closer to the ppm range.[107–109] Although 80 ppb (0.08 µg/ml) of Cr^{3+}

was bound to the unspliced precursor (Table 2), it failed to exert any visible effects on *Mtb* SufB

splicing reactions. No bound form of metal ions was detected in Pb^{2+} treated samples, even at 2

mM concentration, consistent with no visible effect on *Mtb* SufB splicing.

Table 2: Quantitative elemental detection via ICP-OES analysis: Measurable concentration of
metal ions (Cd^{2+}, Hg^{2+}, $Cr^{3+,}$ and Pb^{2+}) were identified bound to *Mtb* SufB precursor and splicing
products. P: Precursor, LE: Ligated extein, I: Intein.

Metal ions	Treated concentration	Concentration (µg/ml)	ICP-OES values (ppb)	Equivalent Concentration (µg/ml)
Cadmium (Cd^{2+})	25µM	4.583	P: -9 LE: -5 I: -8	P: -0.009 LE: -0.005 I: -0.008
Cadmium (Cd^{2+})	500 µM	91.66	P: 2 LE: 18 I: 1	P: 0.002 LE: 0.018 I: 0.001
Mercury (Hg^{2+})	500 µM	135.76	P: 5 LE: 3 I: 5	P: 0.005 LE: 0.003 I: 0.005
Chromium (Cr^{3+})	2mM	361.72	P: 80 LE: 83 I: 90	P: 0.08 LE: 0.083 I: 0.09
Lead (Pb^{2+})	2mM	556.2	P: -17 LE: -5 I: -73	P: -0.017 LE: -0.005 I: -0.073

Collectively, the gradient and time-dependent splicing assays, along with protein~metal

interaction studies, suggested possible interaction of Cd^{2+} and Hg^{2+} with *Mtb* SufB precursor

protein, thereby attenuating the splicing and N-terminal cleavage efficacy. These mechanistic

insights led to the development of an *in vitro* whole-cell biosensing platform utilizing H37Ra (*Mtb*

attenuated strain) cells, as the metal indicator to detect metal ions.

Fe-S clusters are essential cofactors for distinct metabolic pathways and cellular processes such as

respiration, metabolism, DNA replication and repair, antibiotic resistance and persistence under

oxidative and nitrosative stress conditions.[110–112] SUF complex offers the sole pathway for Fe-S

cluster biogenesis in mycobacteria, and its activity relies on generation of functional SufB via

protein splicing.[53,54] Besides, there are earlier reports on loss of cellular viability as a result of SufB

splicing inhibition within mycobacteria.[37,38,46,48,57,59,113–116] Since, toxic metal ions like of Cd^{2+}

and Hg^{2+} were shown to inhibit *Mtb* SufB splicing and N-terminal cleavage activity, an attenuated

strain of *Mtb* (H37Ra) was used as the whole-cell viability indicator for these metal ions.

H37Ra- based whole-cell biosensor detects toxic metal ions

Cell-based biosensors are gaining interest as a screening platform for biologically active molecules

due to their ability to monitor the cellular functions and interactions with targets in their natural

environment.[117,118] Often, they are fabricated as genetically encoded biosensors using native

receptors or enzymes as molecular recognition elements in living cells.[26] Current work explores

the use of whole-cell mycobacteria (H37Ra)-based biosensor to detect toxic metal ions like Cd^{2+}

and Hg^{2+}. An *in vitro* 96-well plate assay was developed using Alamar Blue reagent, employing

Mtb (H37Ra) cells as the sensing component, and *M. sm* cells as the SufB intein-less cellular

control.[55,115,119] *M. sm* carries three intein-bearing proteins: DnaB, which contains two inteins

(DnaBi1 and DnaBi2), as well as GyrA and PhoH.[44,57] Although splicing regulation has been

reported for DnaBi1 helicase, studies on the DnaBi2, PhoH, GyrA inteins splicing have not been

explored yet. DnaBi1 splicing is regulated by zinc and oxidative stressors;[44,57] however, its helicase

function can be compensated by other cellular helicases.[120–122]

Alamar Blue reagent is a non-toxic, resazurin-based solution that functions as an indicator for cell

viability.[67,123] The blue-coloured non-fluorescent resazurin (7-hydroxy-3H-phenoxazin-3-one 10-

oxide) gets reduced to resorufin (7-hydroxy-3H-phenoxazin-3-one, pink coloured compound) by

mitochondrial reductase in metabolically active cells.[124] In a comparative study, non-viable H37Ra

were visualized in presence of Cd^{2+} and Hg^{2+}, whereas persistence of viable *M. sm* cells during the

metal-interaction process highlighted the absence of metal sensing SufB precursor protein.

Untreated H37Ra cells grown in 7H9 (supplemented with OADC) media, and Alamar Blue reagent

alone in 7H9 (supplemented with OADC) media, served as the negative and positive controls for

the study respectively.

Figure 2 (i) is a schematic representation of *in vitro* 96-well plate-based biosensor platform

containing *Mtb* H37Ra cells (indicator organisms) incubated with varied concentrations of Cd^{2+}

(2.5 µM -2 mM) and Hg^{2+} (25 µM -2 mM). The test concentration range for the metal ions were

chosen based on the positive results obtained during *in vitro* splicing analyses (Figure 1, Figures

S2-S4, Supporting Information). For a baseline correction, H37Ra cells were grown in the absence

of toxic metal ions (negative control). SufB intein-less, *M. sm* cells were used as additional

experimental control. Unlike *Mtb* H37Ra cells, the growth and viability of the *M. sm* cells are

independent of SufB intein splicing, and any loss of cell viability indicates metal toxicity. Figure

2 (ii) shows that the indicator cells (H37Ra) incubated in the presence of lower concentration of

Cd^{2+} (2.5-10 µM) did not lose cell viability. All the cells were metabolically active and converted

resazurin (blue) to resorufin (pink), since Cd^{2+} at these concentration (2.5-10 µM) didn't affect

SufB splicing activity, generating active SufB protein (Figure S3 (i), Supporting Information).

H37Ra cells incubated in presence of 25 and 50 µM of Cd^{2+}, displayed purple colour indicating a

partial loss of cell viability, underscoring 25 µM as the minimum inhibitory concentration (MIC).

Cells incubated with Cd^{2+} at a concentration range of 100 µM- 2 mM, retained blue colour,

indicating complete loss of cell viability. These results correlated to inhibition of splicing and N-

terminal activity of *Mtb* SufB in presence of Cd^{2+} over a concentration of 25 µM - 2 mM (Figure

1).

Next, cells incubated in the presence of 25 µM and 50 µM of Hg^{2+} resulted in healthy viable (pink)

cell population, whereas cells in the presence of 100 µM Hg^{2+} exhibited purple colour, highlighting

partial loss of cell viability at MIC value. Over rest of the concentration range for Hg^{2+} (500 µM-

2 mM), non-viable (blue) cells were noticed. These results were supported by SufB splicing

attenuation by Hg^{2+} (500 µM-2 mM), while a lower concentration (2.5 µM-10 µM) Hg^{2+} failed to

exert any such effects (Figure 1, Figure S4, Supporting Information). Although 100 µM of Hg^{2+}

didn't influence SufB splicing, the N-terminal cleavage reaction was reduced significantly,

marking this as the MIC value with partial loss of cell viability. Summarizing, persistence of blue

colour in the presence of Cd^{2+} (100 µM until 2 mM), and Hg^{2+} (500 µM until 2 mM) may be

correlated to complete loss of H37Ra viability due to inhibition of *Mtb* SufB splicing and N-

terminal cleavage reactions (Figure 1). These results suggest the potential of mycobacterial

(H37Ra) cells as the biological indicator for toxic metal detection.

Figure 2. 96-well plate-based biosensor assay demonstrating whole cell mycobacteria (*Mtb* H37Ra strain) as the biological indicator for toxic metal ions (Cd^{2+} and Hg^{2+}) (i) Schematic representation of 96 well plate-based Biosensor platform. (ii) H37Ra cells (10^8 CFU/ml) were seeded in the presence of varied concentration ranges of CdCl$_2$ (2.5 µM- 2 mM) and HgCl$_2$ (25 µM- 2 mM), respectively. Cells incubated in the presence of CdCl$_2$ (2.5 µM- 10 µM) and HgCl$_2$ (25 µM- 50 µM) were pink in colour indicating healthy viable cells resisting the splicing inhibitory effect of the metal ions. In contrast, cells exposed to CdCl$_2$ (25 µM and 50 µM) and 100 µM of HgCl$_2$ displayed purple colour suggesting partial loss of cell viability. Cells incubated with 100 µM-2 mM of CdCl$_2$ and 100 µM- 2 mM of HgCl$_2$ remained blue, suggesting complete loss of viable cell population. For a baseline correction, test organisms were allowed to grow in the absence of toxic metal ions and denoted as live cell control (negative control). SufB intein-less organism *Mycobacterium smegmatis (M. sm)* was used as additional experimental control and treated under similar conditions. *M. sm* cells exhibited pink colour across a varied range of toxic metal ions, suggesting their cell viability as an intein-independent phenomenon. The experiments were performed in triplicates to confirm the reproducibility of data. Cyan stars- complete loss of viability (blue) & Yellow stars- partial loss of viability (purple).

Next, a quantitative analysis for the toxic metal ions was performed by measuring the absorbance

of Alamar Blue reagent at 570 nm with 600 nm as a reference wavelength.[67] Metal samples of

known concentrations were prepared and loaded on to the 96-well plate and treated with Alamar

Blue reagent under optimised experimental conditions [Figures 3 (i)-(iv)]. A standard graph was

generated by plotting the absorbance values of Alamar reagent vs the known concentrations of test

metal ions (Cd^{2+} and Hg^{2+}) [Figures 3 (v) and (vi)]. Subsequently, a linear regression analysis was

performed to determine the equation of the line, expressed as y = mx + c. To further validate the

standard plot, unknown samples with varying concentrations of Cd^{2+} ions and Hg^{2+} ions were

prepared (Materials and Methods). Absorbance values of the unknown samples were extrapolated

onto standard graph to determine the concentrations of Cd^{2+} and Hg^{2+} (Figures 3 (v), (vi) and Table

S1, Supporting Information). Unknown samples with concentration higher than the standard range

may be adjusted via dilution, or the standard calibration plot may be extended to higher

concentration ranges.[125] However sample concentration at a value lower than the standard range

cannot be determined accurately possibly due to no effect on intein splicing and hence no loss of

cell viability.

Figure 3. Standard curve for quantitative determination of Cd^{2+} and Hg^{2+} concentrations.
(i), and (iii) Schematic representations of H37Ra-based Biosensor platform to detect $CdCl_2$ and
$HgCl_2$. (ii), and (iv) *Mtb* H37Ra cells were grown in presence of varied concentration of $CdCl_2$
and $HgCl_2$. To establish a baseline, test organisms were allowed to grow in absence of toxic metal
ions (live cell or negative control). (v), and (vi) The absorbance values of the Alamar Reagent

solutions were plotted along the y-axis and the known concentrations of Cd^{2+} and Hg^{2+} were
plotted along the x-axis. A linear regression analysis was conducted to obtain the equation for the
line, and used to determine the concentration of unknown samples. The experiments were
performed in triplicates to confirm the reproducibility of data.

Next, to elucidate the specificity of the designed biosensor platform, Cd^{2+}- and Hg^{2+}- specific

metal chelators, Edetate Calcium Disodium (ECD), and meso-2,3-Dimercaptosuccinic acid

(DMSA) were incubated alongside the test samples.[126–131] Figure 4 (i) is a schematic representation

of the *in vitro* biosensor platform, where indicator mycobacterial cells were exposed to Cd^{2+} and

Hg^{2+}, in presence of Cd^{2+}-specific chelator, ECD. Figure 4 (ii) demonstrates that ECD effectively

resisted Cd^{2+}- induced loss of viability, resulting in healthy cells that transformed resazurin (blue)

to resorufin (pink). Conversely, ECD failed to restore viability of Hg^{2+} treated cells, where

inhibition of SufB splicing, rendered non-viable (blue) cells. Next, mycobacterial cells were

subjected to varying concentrations of Cd^{2+} and Hg^{2+}, supplemented with metal chelator DMSA

(Hg^{2+} specific) [Figure 4 (iii) and (iv)]. Hg^{2+} treated H37Ra cells remained viable in presence of

DMSA, while Cd^{2+} continued its inhibitory effect on SufB splicing inducing non-viable (blue)

cells. These observations affirmed the specificity of the designed biosensor platform to detect

toxic metal ions Cd^{2+} and Hg^{2+}.

Figure 4. Determination of the specificity of *Mtb* H37Ra-based biosensor platform using metal chelators. (i) Schematic representation of the biosensor platform including Cd^{2+} specific metal chelator (ECD) (ii) *Mtb* H37Ra cells (10^8 CFU/ml) were allowed to grow in presence of varied concentration range of $CdCl_2$ and $HgCl_2$, along with ECD (1:1 ratio). ECD reversed the inhibitory effect of Cd^{2+} inducing viable (pink) cells, whereas the effect of Hg^{2+} persisted (iii) Schematic representation of the biosensor platform including Hg^{2+} specific metal chelator (DMSA). (iv) *Mtb* H37Ra cells (10^8 CFU/ml) were allowed to grow in the presence of varied concentration ranges of $CdCl_2$ and $HgCl_2$, including DMSA in the ratio 1:1. DMSA blocked the inhibitory effect of Hg^2, rendering pink viable cells, whereas the effect of Cd^{2+} persisted. For a baseline correction, H37Ra cells were grown in the absence of toxic metal ions as the live cell control. SufB intein-less organism *Mycobacterium smegmatis (M. sm)* was used as additional experimental control and treated under similar conditions. *M. sm* cells exhibited pink colour across a varied range of toxic metal ions, suggesting their cell viability as an intein-independent phenomenon. The experiments were performed in triplicates to confirm the reproducibility of data. Cyan stars- complete loss of viability (blue) & Yellow stars- partial loss of viable cells (purple). ECD: Edetate Calcium disodium, DMSA: Meso-2,3-Dimercaptosuccinic acid.

Next for a multiplex metal screening process, the efficacy of the platform was assessed in presence

of known splicing inhibitors Zn^{2+} (100 μM-2 mM), Cu^{2+} (100 μM-2 mM), Pt^{4+} (20 μM-1 mM)

along with Cd^{2+} (10 μM-2 mM) and Hg^{2+} (25 μM-2 mM) (Figure 5).[38] Baseline correction was

achieved by growing H37Ra cells in the absence of toxic metal ions (negative control), while SufB

intein-less *M. sm* cells served as additional experimental control. Cells treated with 100 μM $ZnCl_2$

and $CuCl_2$, 20 μM $PtCl_4$, 10 μM $CdCl_2$, and 25 μM $HgCl_2$, turned pink in colour, indicating intact

viability. At intermediate concentrations, 600 μM of $ZnCl_2$ and $CuCl_2$, 40 μM $PtCl_4$, 25 μM $CdCl_2$,

and 100 μM $HgCl_2$, a transition from blue to purple colour was observed, suggesting partial loss

of cell viability. Complete loss of viability was noted at higher concentration range, for $ZnCl_2$ and

$CuCl_2$ (800 μM- 2 mM), $PtCl_4$ (80 μM- 1 mM), $CdCl_2$ (100 μM- 2 mM), and $HgCl_2$ (500 μM- 2

mM) (Figure 5 ii). Dead non-viable cells induced by Zn^{2+}, Cu^{2+}, and Pt^{4+} may be correlated to

inhibition of *Mtb* SufB splicing and N-terminal cleavage reactions as published earlier.[38] In

contrast, SufB intein-less organism (*M. sm*) remained viable (pink) under similar experimental

conditions, suggesting cell viability as a splicing independent phenomenon. These observations

may also be used to rule out metal toxicity on cell viability over the test concentration range.

Figure 5. Evaluation of *Mtb* H37Ra-based biosensor platform for high-throughput and multiplex detection of trace metal elements and toxic metal ions (Zn^{2+}, Cu^{2+}, Pt^{4+}, Cd^{2+}, and Hg^{2+}). (i) Schematic representation of the multiplexing biosensor platform. (ii) H37Ra cells (10^8 CFU/ml) were allowed to grow in the presence of $ZnCl_2$ (100 µM- 2 mM), $CuCl_2$ (100 µM- 2 mM), $PtCl_4$ (20 µM- 1 mM), $CdCl_2$ (10 µM- 2 mM) and $HgCl_2$ (25 µM- 2 mM), respectively. Cells incubated with 100 µM of $ZnCl_2$ and $CuCl_2$, 20 µM of $PtCl_4$, 10 µM of $CdCl_2$ and 25 µM of $HgCl_2$ were pink in colour indicating healthy viable cells. In contrast, cells exposed to 600 µM of $ZnCl_2$ and $CuCl_2$, 40 µM of $PtCl_4$, 25 µM of $CdCl_2$ and 100 µM of $HgCl_2$ turned purple in colour suggesting partial loss of cell viability, whereas cells exposed to 600 µM- 2 mM of $ZnCl_2$ and $CuCl_2$, 80 µM- 1 mM of $PtCl_4$, 100 µM- 2 mM of $CdCl_2$ and 500 µM- 2 mM of $HgCl_2$ remained blue in colour suggesting complete loss of viable cell population. For a baseline correction, H37Ra cells were grown in the absence of toxic metal ions as the live cell control (negative control). SufB intein-less organism *Mycobacterium smegmatis (M. sm)* was used as additional experimental control and treated under similar conditions. *M. sm* cells exhibited pink colour across a varied range of toxic metal ions, suggesting their cell viability as an intein-independent phenomenon. The experiments were performed in triplicates to confirm the reproducibility of data. Cyan stars- complete loss of viability (blue) & Yellow stars- partial loss of viability (purple).

Further assessment of the multiplexing biosensor platform was done by employing metal-specific chelators along with the metal-treated samples. The Cd^{2+}-specific chelator ECD and Hg^{2+}-specific chelator DMSA were tested alongside the broad-spectrum metal chelator EDTA. Figure 6 (i) is a

schematic representation of the *in vitro* biosensor platform, where indicator mycobacterial cells

were incubated with Cd^{2+}-specific chelator, ECD. Figure 6 (ii) demonstrates that co-incubation

with ECD effectively reversed Cd^{2+}- induced loss of viability, restoring pink colour. This effect

was not observed in cells exposed to other metal ions, where ECD could not block metal induced

inhibition. Similarly, DMSA reversed the toxic effects of Hg^{2+} [Figure 6 (iii) and (iv)], while other

metal ions continued to attenuate cell viability. Notably, Since EDTA is a broad-spectrum metal

chelator, it restored viability across all metal-treated samples [Figure 6 (v) and (vi)]. Collectively,

these findings validate the high-throughput and multiplexing capability of the developed biosensor

platform, underscoring its potential for simultaneous detection of multiple toxic metal ions with

high specificity and sensitivity.

Figure 6. Determination of specificity of the *Mtb* H37Ra-based multiplexing biosensor platform. (i) Schematic representation of the multiplexing biosensor platform along with Cd^{2+} specific metal chelator (ECD) (ii) *Mtb* H37Ra cells (10^8 CFU/ml) were incubated with different concentration of $ZnCl_2$, $CuCl_2$, $PtCl_4$, $CdCl_2$ and $HgCl_2$, along with ECD (1:1 ratio). ECD reversed the inhibitory effect of Cd^{2+} rendering viable pink cells, whereas the effects of other metal ions persisted (iii) Schematic representation of the multiplexing biosensor platform along with Hg^{2+} specific metal chelator (DMSA). (iv) *Mtb* H37Ra cells (10^8 CFU/ml) were incubated with different concentration of $ZnCl_2$, $CuCl_2$, $PtCl_4$, $CdCl_2$ and $HgCl_2$, along with DMSA in the ratio 1:1. DMSA inhibited the effect of Hg^{2+} inducing viable pink cells, whereas the effect of other metal ions persisted. (v) Schematic representation of plate-based biosensor platform in presence of common metal chelator EDTA. (vi) *Mtb* H37Ra cells (10^8 CFU/ml) were allowed to grow in presence of $ZnCl_2$, $CuCl_2$, $PtCl_4$, $CdCl_2$ and $HgCl_2$ along with EDTA in the ratio 1:1. The inhibitory effects of test metal ions on mycobacterial cells were reversed by EDTA. For a baseline correction, H37Ra cells were grown in the absence of toxic metal ions, as the live cell control. SufB intein-less organism *Mycobacterium smegmatis (M. sm)* was used as additional experimental control and treated under similar conditions. *M. sm* cells exhibited pink colour across a varied range of toxic metal ions, suggesting their cell viability as an intein-independent phenomenon. The experiments were performed in triplicates to confirm the reproducibility of data. Cyan stars- complete loss of viability (blue) & Yellow stars- partial loss of viable cells (purple). ECD: Edetate Calcium disodium, DMSA: Meso-2,3-Dimercaptosuccinic acid, EDTA: Ethylenediaminetetraacetic Acid.

In toto, the designed biosensor has demonstrated the potential of whole-cell *Mtb* H37Ra as an indicator to detect toxic metal ions as well as TMEs in test samples. The specificity of the designed metal-sensor was shown by the reversal of the inhibitory effects in the presence of Cd^{2+}- and Hg^{2+}-specific metal chelators, ECD and DMSA, respectively. Detection of metal ions at a concentration of 25 µM (Cd^{2+}) highlights the sensitivity of the biosensor platform, although it requires further optimization to cover the lower concentration range of various metal ions. We propose the use of whole-cell microorganisms carrying metal-sensing intein-bearing proteins as an indicator for the presence of toxic metal ions in environmental samples. This is a user-friendly method that can be performed in a 96-well microtiter plate, which would provide a colour change between blue and pink while sensing metal ions. Since the activity of metal-sensing intein-carrying protein

sequences are unique for the indicator (whole-cell) organisms, in the unlikely event of

contamination in a biological facility, it should not raise concern about the validity of the results.

Conclusion

Heavy metal ions are persistent environmental contaminants that pose significant risks to both

ecological systems and human health. With the rising burden of metal pollution across diverse

environmental and biological matrices, there is a pressing need for sensitive, specific, and user-

friendly biosensing technology. Intein-mediated biosensors are emerging as promising tools in this

context, particularly due to their unique splicing mechanism that operates efficiently under

physiological conditions, eliminating the need for external cofactors or energy sources.[31,132] In this

study, *Mtb* H37Ra -based *in vitro* biosensor platform was developed, employing intein splicing

inhibition as a readout mechanism for the detection of toxic heavy metal ions (e.g., Cd^{2+} and Hg^{2+})

and other trace metal elements (Figure 7). The resulting alteration in mycobacterial viability is

captured through a rapid and quantifiable resazurin-based colorimetric assay that reports colour

change between blue-purple-pink. Importantly, the platform is adaptable for multiplex detection

of metal ions, including Zn^{2+}, Pt^{4+}, Cu^{2+}, Cd^{2+}, and Hg^{2+}, thereby offering a high-throughput

solution for complex sample screening (Figures 5 and 6). Further, this can be extended to sense

metal ions by other whole-cell microorganisms, harbouring unique metal-sensing intein-

containing proteins, hence offering a qualitative as well as quantitative detection method.

SufB is a critical component of SUF system, the exclusive pathway for Fe-S cluster assembly in

mycobacteria. Previous studies have also highlighted the pivotal role of *Mtb* SufB protein in Fe-

S cluster biogenesis, and mycobacterial survival.[54,133] Since the functionality of SUF protein

complex is dependent on generation of active SufB protein via splicing, splicing inhibition

presents a promising approach for antimicrobial intervention by targeting mycobacterial viability.

Several studies have shown that divalent cations (Zn^{2+} and Cu^{2+}), ROS, RNS, chlorine derivatives,

small molecules (cisplatin and diethanolamine) and zinc oxide nanoparticles can act as potent

splicing inhibitors. These agents have influenced mycobacterial survival, further validating the

relevance of targeting intein splicing for antimycobacterial drug development.[37,38,46,48,57,59,113–116]

Earlier work from our group has demonstrated that metal ions such as Cu^{2+}, Zn^{2+}, and Pt^{4+} inhibit

SufB splicing, establishing a correlation between metal-induced stress and altered intein splicing

activity in *Mtb*.[38] Building on these findings, the current work explores regulatory roles of toxic

metal ions on *Mtb* SufB splicing and cleavage activity. Alamar blue assay was employed to

examine the influence of TMEs and toxic metal ions on the viability of *Mtb* cells. The outcome of

this research will advance the real time application of H37Ra based whole-cell biosensor to detect

wide range of metal ions. The platform described herein is robust, enabling multiplex screening of

metal ions while maintaining specificity and sensitivity. Importantly, cellular viability loss in

response to metal exposure is linked to an observable colour shift between blue and pink, providing

a quantifiable readout for metal ion detection in test samples.

In this study, splicing inhibition was evaluated both *in vitro*, using purified SufB protein and *in*

vivo within H37Ra cells in a whole-cell biosensor platform, where the observed effects on

mycobacterial viability were correlated to SufB splicing inhibition. Given that H37Ra is a slow-

growing strain with a mean generation time of approximately 44 hours, the experimental window

was established at four days to encompass at least two full generations, permitting effective

assessment of metal-induced viability changes.[134–136] Notably, the logarithmic growth phase of

mycobacteria extends from four to twenty days, ensuring that the platform remains adaptable to

varied experimental requirements. To further strengthen our hypothesis, SufB intein less organism,

M. sm control was included; viability of *M. sm* persisted throughout the experimental period,

suggesting that the tested metal ions do not elicit growth inhibitory or toxic effects in the absence

of SufB intein. *Mtb* contains three intein containing proteins; SufB, RecA and DnaB. While SufB

is essential for Fe-S cluster assembly, RecA and DnaB play pivotal roles during DNA replication;

taken together these proteins are critical for mycobacterial survival.[137–139] Although current study

explains loss of mycobacterial cell viability as a result of inhibition of *Mtb* SufB splicing, the

possibility of RecA and DnaB splicing inhibition by the test metal ions cannot be ruled completely.

Currently, there are no literatures to show inhibitory effect of metal ions other than Zn^{2+} on *Mtb*

DnaB splicing.[44] Moreover there are alternative helicases present in mycobacteria to take over the

function of DnaB.[120–122] Several studies have reported regulatory roles of Zn^{2+}, Cd^{2+}, Co^{2+}, Ni^{2+},

Cu^+/Cu^{2+}, CisPt ions on *Mtb* RecA intein splicing.[40,42,48,113] However, current study has not

evaluated the RecA intein inhibition in mycobacteria. Hence, this approach offers a multiplexing

biosensor unit which can accommodate cell viability detection via splicing inhibition in many

organisms carrying different metal-sensitive intein containing proteins simultaneously.

The use of non-genetically modified *Mtb* H37Ra cells ensures biosecurity compliance while

providing a biologically relevant sensing chassis. These mycobacterial cells are not only easy to

cultivate and maintain but also exhibit remarkable long-term viability, supporting extended shelf-

life and storage without specialized conditions. The platform also demonstrates good sensitivity in

detecting metal ions over a broad dynamic range, and substrate specificity, making it suitable for

metal ion screening in standard laboratory setups.

However, some limitations persist such as complex environmental or biological samples

containing mixtures of multiple metal ions may result in signal overlap or cross-reactivity with

variable binding affinities. Furthermore, while microbial whole-cell biosensors offer an excellent

alternative for initial screening and field deployment, advanced physicochemical instruments

remain crucial when trace-level quantification is required, where intein activity may not be

efficiently regulated.

In summary, our findings highlight the potential of whole cell intein-based biosensors as user-

friendly, and cost-effective tools for the detection of toxic metal and trace metal ions. Future work

should aim to refine their activity, enhancing environmental robustness, and advancing the

platform from lab-scale to deployable technologies. This can be accelerated through collaborative

efforts across academic, industrial, and governmental sectors.

Figure 7. Schematic representation explaining the principle of *Mtb* H37Ra-based
multiplexing biosensor platform. Whole-cell mycobacteria (H37Ra) were used as biological

indicator for the detection of toxic metal ions (Cd^{2+} and Hg^{2+}) and other trace metal elements
(TME). (i) Intein splicing under normal physiological conditions generates functional SufB
protein, while viable mycobacterial cells convert resazurin (blue) to resorufin (pink). (ii) In the
presence of toxic metal ions Cd^{2+} and Hg^{2+} and TME, SufB intein splicing is inhibited resulting in
non-viable mycobacterial cells, thus there is no conversion of resazurin (blue) to resorufin (pink).

Acknowledgments

The plasmids used in this study are borrowed from Prof. Marlene Belfort's Lab, SUNY, Albany,

NY, USA. Our sincere gratitude to the Central Instrumentation Facility (CIF) at OUAT (Odisha

University of Agriculture & Technology), Bhubaneswar, and Dr. S.K. Dash for his help in

conducting the ICP-OES analysis at OUAT, Bhubaneswar.

Funding

Ms. Ashwaria Mehra is supported by INSPIRE fellowship (DST/INSPIRE/03/2021/002800/IF

190921); INSPIRE Division, DST, Government of India. This work was also supported by the

UGC-DAE consortium for scientific research (UGC-DAE-CSR-KC/CRS/15/IOP/08/0562),

Kolkata, India.

Author Contributions

AM: Writing – Original Draft, Writing - Review & Editing, Methodology, Investigation, Formal

analysis, Data curation, Visualization, Validation, Software, Resources, Funding acquisition; **AN:**

Conceptualization, Methodology, Investigation, Formal analysis, Data curation; Writing - Review

& Editing, ; **SN:** Conceptualization, Writing - Original Draft, Writing - Review & Editing, Project

administration, Supervision, Methodology, Investigation, Resources, Funding acquisition.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Abbreviations:

Ni-NTA: Nickel-nitrilotriacetic acid
TCEP.HCl: Tris 2-carboxyl ethyl phosphine. Hydrochloric acid

IPTG: Isopropyl β-d-1-thiogalactopyranoside

Trp: Tryptophan

SDS: Sodium Dodecyl Sulfate

PAGE: Polyacrylamide gel electrophoresis

ICP-OES: Inductively Coupled Plasma Optical Emission Spectroscopy

UV-Vis: UV-visible spectroscopy

PVDF: Polyvinylidene fluoride

BSA: Bovine serum albumin

Mtb: *Mycobacterium tuberculosis*

WT: Wild type

SI: Splicing inactive

Cd^{2+}: Cadmium

Hg^{2+}: Mercury

Pb^{2+}: Lead

Cr^{3+}: Chromium

References

(1)   Velusamy, K.; Periyasamy, S.; Kumar, P. S.; Rangasamy, G.; Nisha Pauline, J. M.;
860   Ramaraju, P.; Mohanasundaram, S.; Nguyen Vo, D. V. Biosensor for Heavy Metals
861   Detection in Wastewater: A Review. *Food and Chemical Toxicology* **2022**, *168*, 113307.
862   https://doi.org/10.1016/J.FCT.2022.113307.

863   (2)   Duffus, J. H. *"heavy Metals" - A Meaningless Term? (IUPAC Technical Report)*; Walter
864   de Gruyter GmbH, 2002; Vol. 74.
865   https://www.degruyterbrill.com/document/doi/10.1351/pac200274050793/html (accessed
866   2025-06-16).

867   (3)   Turdean, G. L.; Ozkan, S. A. Design and Development of Biosensors for the Detection of
868   Heavy Metal Toxicity. *International Journal of Electrochemistry* **2011**, *2011* (1), 343125.
869   https://doi.org/10.4061/2011/343125.

870   (4)   Singh, R.; Gautam, N.; Mishra, A.; Gupta, R. Heavy Metals and Living Systems: An
871   Overview. *Indian J Pharmacol* **2011**, *43* (3), 246–253. https://doi.org/10.4103/0253-
872   7613.81505.

873   (5)   Angon, P. B.; Islam, M. S.; KC, S.; Das, A.; Anjum, N.; Poudel, A.; Suchi, S. A. Sources,
874   Effects and Present Perspectives of Heavy Metals Contamination: Soil, Plants and Human
875   Food Chain. *Heliyon* **2024**, *10* (7), e28357.
876   https://doi.org/10.1016/J.HELIYON.2024.E28357.

877   (6)   Alengebawy, A.; Abdelkhalek, S. T.; Qureshi, S. R.; Wang, M. Q. Heavy Metals and
878   Pesticides Toxicity in Agricultural Soil and Plants: Ecological Risks and Human Health
879   Implications. *Toxics* **2021**, *9* (3), 42. https://doi.org/10.3390/TOXICS9030042.

880   (7)   Jaishankar, M.; Tseten, T.; Anbalagan, N.; Mathew, B. B.; Beeregowda, K. N. Toxicity,
881   Mechanism and Health Effects of Some Heavy Metals. *Interdiscip Toxicol* **2014**, *7* (2),
882   60–72. https://doi.org/10.2478/INTOX-2014-0009.

883   (8)   Hassani, S.; Rezaei Akmal, M.; Salek Maghsoudi, A.; Rahmani, S.; Vakhshiteh, F.;
884   Norouzi, P.; Ganjali, M. R.; Abdollahi, M. High-Performance Voltammetric Aptasensing
885   Platform for Ultrasensitive Detection of Bisphenol A as an Environmental Pollutant. *Front
886   Bioeng Biotechnol* **2020**, *8*, 574846. https://doi.org/10.3389/FBIOE.2020.574846.

887   (9)   Odobašić, A.; Šestan, I.; Begić, S.; Odobašić, A.; Šestan, I.; Begić, S. *Biosensors for
888   Determination of Heavy Metals in Waters*; IntechOpen, 2019.
889   https://doi.org/10.5772/INTECHOPEN.84139.

890   (10)   Ramnani, P.; Saucedo, N. M.; Mulchandani, A. Carbon Nanomaterial-Based
891   Electrochemical Biosensors for Label-Free Sensing of Environmental Pollutants.
892   *Chemosphere* **2016**, *143*, 85–98. https://doi.org/10.1016/J.CHEMOSPHERE.2015.04.063.

893   (11)   Pena-Pereira, F.; Wojnowski, W.; Tobiszewski, M. AGREE - Analytical GREEnness
894   Metric Approach and Software. *Anal Chem* **2020**, *92* (14), 10076–10082.
895   https://doi.org/10.1021/acs.analchem.0c01887.

896   (12)   Pearson, J. E.; Gill, A.; Vadgama, P. Analytical Aspects of Biosensors. *Ann Clin Biochem*
897   **2000**, *37* (2), 119–145. https://doi.org/10.1258/0004563001899131.

898   (13)   Malandain, C.; Fayolle, F.; Bedouelle, H. Biosensors for the Environment. *Oil & Gas*
899   *Science and Technology* **2005**, *60* (6), 887–897. https://doi.org/10.2516/OGST:2005062.

900   (14)   Bontidean, I.; Berggren, C.; Johansson, G.; Csöregi, E.; Mattiasson, B.; Lloyd, J. R.;
901   Jakeman, K. J.; Brown, N. L. Detection of Heavy Metal Ions at Femtomolar Levels Using
902   Protein-Based Biosensors. *Anal Chem* **1998**, *70* (19), 4162–4169.
903   https://doi.org/10.1021/ac9803636.

904   (15)   Blake, D. A.; Jones, R. M.; Blake, R. C.; Pavlov, A. R.; Darwish, I. A.; Yu, H. Antibody-
905   Based Sensors for Heavy Metal Ions. *Biosens Bioelectron* **2001**, *16* (9–12), 799–809.
906   https://doi.org/10.1016/S0956-5663(01)00223-8.

907   (16)   Chouteau, C.; Dzyadevych, S.; Durrieu, C.; Chovelon, J. M. A Bi-Enzymatic Whole Cell
908   Conductometric Biosensor for Heavy Metal Ions and Pesticides Detection in Water
909   Samples. *Biosens Bioelectron* **2005**, *21* (2), 273–281.
910   https://doi.org/10.1016/J.BIOS.2004.09.032.

911   (17)   Mohammadi, Z. B.; Zhang, F.; Kharazmi, M. S.; Jafari, S. M. Nano-Biocatalysts for Food
912   Applications; Immobilized Enzymes within Different Nanostructures. *Crit Rev Food Sci*
913   *Nutr* **2023**, *63* (32), 11351–11369. https://doi.org/10.1080/10408398.2022.2092719.

914   (18)   Sun, G.; Wei, X.; Zhang, D.; Huang, L.; Liu, H.; Fang, H. Immobilization of Enzyme
915   Electrochemical Biosensors and Their Application to Food Bioprocess Monitoring.
916   *Biosensors (Basel)* **2023**, *13* (9), 886. https://doi.org/10.3390/BIOS13090886.

917   (19)   Kakkar, P.; Jaffery, F. N. Biological Markers for Metal Toxicity. *Environ Toxicol*
918   *Pharmacol* **2005**, *19* (2), 335–349. https://doi.org/10.1016/J.ETAP.2004.09.003.

919   (20)   Castillo, J.; Gáspár, S.; Leth, S.; Niculescu, M.; Mortari, A.; Bontidean, I.; Soukharev, V.;
920   Dorneanu, S. A.; Ryabov, A. D.; Csöregi, E. Biosensors for Life Quality: Design,
921   Development and Applications. *Sens Actuators B Chem* **2004**, *102* (2), 179–194.
922   https://doi.org/10.1016/J.SNB.2004.04.084.

923 (21) Bousse, L. Whole Cell Biosensors. *Sens Actuators B Chem* **1996**, *34* (1–3), 270–275.
924    https://doi.org/10.1016/S0925-4005(96)01906-5.

925 (22) Stenger, D. A.; Gross, G. W.; Keefer, E. W.; Shaffer, K. M.; Andreadis, J. D.; Ma, W.;
926    Pancrazio, J. J. Detection of Physiologically Active Compounds Using Cell-Based
927    Biosensors. *Trends Biotechnol* **2001**, *19* (8), 304–309. https://doi.org/10.1016/S0167-
928    7799(01)01690-0.

929 (23) Wang, Y.-Z.; Christopher, J. K.; Yong, Y.-C.; Zhai, D.-D. *Nutrient Detection with Whole-*
930    *Cell Biosensors*; Springer, Cham, 2019. https://doi.org/10.1007/978-3-319-47405-2_110-
931    1.

932 (24) Mihaela, G.; Mihaela, G. A Short Review on Cell-Based Biosensing: Challenges and
933    Breakthroughs in Biomedical Analysis. *J Biomed Res* **2021**, *35* (4), 255–263.
934    https://doi.org/10.7555/JBR.34.20200128.

935 (25) Raut, N.; O'Connor, G.; Pasini, P.; Daunert, S. Engineered Cells as Biosensing Systems in
936    Biomedical Analysis. *Anal Bioanal Chem* **2012**, *402* (10), 3147–3159.
937    https://doi.org/10.1007/s00216-012-5756-6.

938 (26) Gui, Q.; Lawson, T.; Shan, S.; Yan, L.; Liu, Y. The Application of Whole Cell-Based
939    Biosensors for Use in Environmental Analysis and in Medical Diagnostics. *Sensors* **2017**,
940    *17* (7), 1623. https://doi.org/10.3390/S17071623.

941 (27) Kang, C.; Shrestha, K. L.; Kwon, S.; Park, S.; Kim, J.; Kwon, Y. Intein-Mediated Protein
942    Engineering for Cell-Based Biosensors. *Biosensors (Basel)* **2022**, *12* (5), 283.
943    https://doi.org/10.3390/BIOS12050283.

944 (28) Jeon, H.; Lee, E.; Kim, D.; Lee, M.; Ryu, J.; Kang, C.; Kim, S.; Kwon, Y. Cell-Based
945    Biosensors Based on Intein-Mediated Protein Engineering for Detection of Biologically
946    Active Signaling Molecules. *Anal Chem* **2018**, *90* (16), 9779–9786.
947    https://doi.org/10.1021/acs.analchem.8b01481.

948 (29) Lee, E.; Shrestha, K. L.; Kang, S.; Ramakrishnan, N.; Kwon, Y. Cell-Based Sensors for
949    the Detection of EGF and EGF-Stimulated Ca2+ Signaling. *Biosensors (Basel)* **2023**, *13*
950    (3), 383. https://doi.org/10.3390/BIOS13030383/S1.

951 (30) Jeon, H.; Lee, M.; Jang, W.; Kwon, Y. Intein-Mediated Protein Engineering for Biosensor
952    Fabrication. *Biochip J* **2016**, *10* (4), 277–287. https://doi.org/10.1007/S13206-016-0404-4.

953 (31) Mills, K. V.; Johnson, M. A.; Perler, F. B. Protein Splicing: How Inteins Escape from
954    Precursor Proteins. *Journal of Biological Chemistry* **2014**, *289* (21), 14498–14505.
955    https://doi.org/10.1074/JBC.R113.540310.

(32)   Perler, F. B. Protein Splicing Mechanisms and Applications. *IUBMB Life* **2005**, *57* (7), 469–476. https://doi.org/10.1080/15216540500163343.

(33)   Panda, S.; Nanda, A.; Nasker, S. S.; Sen, D.; Mehra, A.; Nayak, S. Metal Effect on Intein Splicing: A Review. *Biochimie* **2021**, *185*, 53–67. https://doi.org/10.1016/J.BIOCHI.2021.03.006.

(34)   Nichols, N. M.; Benner, J. S.; Martin, D. D.; Evans, T. C. Zinc Ion Effects on Individual Ssp DNaE Intein Splicing Steps: Regulating Pathway Progression. *Biochemistry* **2003**, *42* (18), 5301–5311. https://doi.org/https://doi.org/10.1021/bi020679e.

(35)   Chan, H.; Pearson, C. S.; Green, C. M.; Li, Z.; Zhang, J.; Belfort, G.; Shekhtman, A.; Li, H.; Belfort, M. Exploring Intein Inhibition by Platinum Compounds as an Antimicrobial Strategy. *Journal of Biological Chemistry* **2016**, *291* (43), 22661–22670. https://doi.org/10.1074/JBC.M116.747824.

(36)   Green, C. M.; Li, Z.; Smith, A. D.; Novikova, O.; Bacot-Davis, V. R.; Gao, F.; Hu, S.; Banavali, N. K.; Thiele, D. J.; Li, H.; Belfort, M. Spliceosomal Prp8 Intein at the Crossroads of Protein and RNA Splicing. *PLoS Biol* **2019**, *17* (10), e3000104. https://doi.org/10.1371/JOURNAL.PBIO.3000104.

(37)   Ojha, D. K.; Mehra, A.; Rout, S. S.; Giri, S.; Nayak, S. Bactericidal Activity of ZnO Nanoparticles–Anti-TB Drug Combination towards the H37Rv Strain and Multidrug-Resistant Isolates of Mycobacterium Tuberculosis via SufB Splicing Inhibition. *Mater Adv* **2025**, *6* (15), 5242–5259. https://doi.org/10.1039/D4MA01224K.

(38)   Panda, S.; Nanda, A.; Nasker, S. S.; Mehra, A.; Ojha, D. K.; Mohanty, P. S.; Nayak, S. Metal Regulation of Mycobacterium Tuberculosis SufB Intein Splicing at the Host–Pathogen Crossroad. *Clin Sci* **2023**, *137* (14), 1027–1048. https://doi.org/10.1042/CS20230307.

(39)   Poland, B. W.; Xu, M. Q.; Quiocho, F. A. Structural Insights into the Protein Splicing Mechanism of PI-SceI. *Journal of Biological Chemistry* **2000**, *275* (22), 16408–16413. https://doi.org/10.1074/JBC.275.22.16408.

(40)   Mills, K. V.; Paulus, H. Reversible Inhibition of Protein Splicing by Zinc Ion. *Journal of Biological Chemistry* **2001**, *276* (14), 10832–10838. https://doi.org/10.1074/JBC.M011149200.

(41)   Van Roey, P.; Pereira, B.; Li, Z.; Hiraga, K.; Belfort, M.; Derbyshire, V. Crystallographic and Mutational Studies of Mycobacterium Tuberculosis RecA Mini-Inteins Suggest a Pivotal Role for a Highly Conserved Aspartate Residue. *J Mol Biol* **2007**, *367* (1), 162–173. https://doi.org/10.1016/J.JMB.2006.12.050.

990   (42)   Zhang, L.; Zheng, Y.; Xi, Z.; Luo, Z.; Xu, X.; Wang, C.; Liu, Y. Metal Ions Binding to
991   RecA Inteins from Mycobacterium Tuberculosis. *Mol Biosyst* **2009**, *5* (6), 644–650.
992   https://doi.org/10.1039/B903144H.

993   (43)   Sun, P.; Ye, S.; Ferrandon, S.; Evans, T. C.; Xu, M. Q.; Rao, Z. Crystal Structures of an
994   Intein from the Split DnaE Gene of Synechocystis Sp. PCC6803 Reveal the Catalytic
995   Model Without the Penultimate Histidine and the Mechanism of Zinc Ion Inhibition of
996   Protein Splicing. *J Mol Biol* **2005**, *353* (5), 1093–1105.
997   https://doi.org/10.1016/J.JMB.2005.09.039.

998   (44)   Woods, D.; Vangaveti, S.; Egbanum, I.; Sweeney, A. M.; Li, Z.; Bacot-Davis, V.;
999   Lesassier, D. S.; Stanger, M.; Hardison, G. E.; Li, H.; Belfort, M.; Lennon, C. W.
1000   Conditional DnaB Protein Splicing Is Reversibly Inhibited by Zinc in Mycobacteria. *mBio*
1001   **2020**, *11* (4), 1–14. https://doi.org/10.1128/mBio.01403-20.

1002   (45)   Pearson, C. S.; Belfort, G.; Belfort, M.; Shekhtman, A. Backbone Assignments of Mini-
1003   RecA Intein with Short Native Exteins and an Active N-Terminal Catalytic Cysteine.
1004   *Biomol NMR Assign* **2015**, *9* (2), 235–238. https://doi.org/10.1007/s12104-014-9581-z.

1005   (46)   Li, Z.; Fu, B.; Green, C. M.; Liu, B.; Zhang, J.; Lang, Y.; Chaturvedi, S.; Belfort, M.;
1006   Liao, G.; Li, H. Cisplatin Protects Mice from Challenge of Cryptococcus Neoformans by
1007   Targeting the Prp8 Intein. *Emerg Microbes Infect* **2019**, *8* (1), 895–908.
1008   https://doi.org/10.1080/22221751.2019.1625727.

1009   (47)   Zhang, L.; Xiao, N.; Pan, Y.; Zheng, Y.; Pan, Z.; Luo, Z.; Xu, X.; Liu, Y. Binding and
1010   Inhibition of Copper Ions to RecA Inteins from Mycobacterium Tuberculosis. *Chemistry -*
1011   *A European Journal* **2010**, *16* (14), 4297–4306.
1012   https://doi.org/https://doi.org/10.1002/chem.200903584.

1013   (48)   Zhang, L.; Zheng, Y.; Callahan, B.; Belfort, M.; Liu, Y. Cisplatin Inhibits Protein
1014   Splicing, Suggesting Inteins as Therapeutic Targets in Mycobacteria. *Journal of*
1015   *Biological Chemistry* **2011**, *286* (2), 1277–1282.
1016   https://doi.org/10.1074/JBC.M110.171124.

1017   (49)   Dassa, B.; Amitai, G.; Caspi, J.; Schueler-Furman, O.; Pietrokovski, S. Trans Protein
1018   Splicing of Cyanobacterial Split Inteins in Endogenous and Exogenous Combinations.
1019   *Biochemistry* **2007**, *46* (1), 322–330. https://doi.org/10.1021/bi0611762.

1020   (50)   Gramespacher, J. A.; Stevens, A. J.; Nguyen, D. P.; Chin, J. W.; Muir, T. W. Intein
1021   Zymogens: Conditional Assembly and Splicing of Split Inteins via Targeted Proteolysis. *J*
1022   *Am Chem Soc* **2017**, *139* (24), 8074–8077. https://doi.org/10.1021/JACS.7B02618.

1023   (51)   Skretas, G.; Wood, D. W. Regulation of Protein Activity with Small-Molecule-Controlled
1024   Inteins. *Protein Science* **2005**, *14* (2), 523–532. https://doi.org/10.1110/PS.04996905.

(52) Ozawa, T.; Nogami, S.; Sato, M.; Ohya, Y.; Umezawa, Y. A Fluorescent Indicator for
Detecting Protein−Protein Interactions in Vivo Based on Protein Splicing. *Anal Chem*
2000, *72* (21), 5151–5157. https://doi.org/10.1021/AC000617Z.

(53) Huet, G.; Daffé, M.; Saves, I. Identification of the Mycobacterium Tuberculosis SUF
Machinery as the Exclusive Mycobacterial System of [Fe-S] Cluster Assembly: Evidence
for Its Implication in the Pathogen's Survival. *J Bacteriol* **2005**, *187* (17), 6137–6146.
https://doi.org/10.1128/jb.187.17.6137-6146.2005.

(54) Huet, G.; Castaing, J. P.; Fournier, D.; Daffé, M.; Saves, I. Protein Splicing of SufB Is
Crucial for the Functionality of the Mycobacterium Tuberculosis SUF Machinery. *J*
Bacteriol **2006**, *188* (9), 3412–3414. https://doi.org/10.1128/JB.188.9.3412-3414.2006.

(55) Panda, S.; Nanda, A.; Sahu, N.; Ojha, D. K.; Pradhan, B.; Rai, A.; Suryawanshi, A. R.;
Banavali, N.; Nayak, S. SufB Intein Splicing in Mycobacterium Tuberculosis Is
Influenced by Two Remote Conserved N-Extein Histidines. *Biosci Rep* **2022**, *42* (3),
20212207. https://doi.org/10.1042/BSR20212207/230724.

(56) Friedel, K.; Popp, M. A.; Matern, J. C. J.; Gazdag, E. M.; Thiel, I. V.; Volkmann, G.;
Blankenfeldt, W.; Mootz, H. D. A Functional Interplay between Intein and Extein
Sequences in Protein Splicing Compensates for the Essential Block B Histidine. *Chem Sci*
2018, *10* (1), 239–251. https://doi.org/10.1039/C8SC01074A.

(57) Kelley, D. S.; Lennon, C. W.; Li, Z.; Miller, M. R.; Banavali, N. K.; Li, H.; Belfort, M.
Mycobacterial DnaB Helicase Intein as Oxidative Stress Sensor. *Nat Commun* **2018**, *9* (1),
1–15. https://doi.org/10.1038/S41467-018-06554-X.

(58) Mills, K. V.; Lew, B. M.; Jiang, S. Q.; Paulus, H. Protein Splicing in Trans by Purified N-
and C-Terminal Fragments of the Mycobacterium Tuberculosis RecA Intein. *Proc Natl*
Acad Sci U S A **1998**, *95* (7), 3543–3548. https://doi.org/10.1073/pnas.95.7.3543.

(59) Gangopadhyay, J. P.; Jiang, S. qin; Paulus, H. An in Vitro Screening System for Protein
Splicing Inhibitors Based on Green Fluorescent Protein as an Indicator. *Anal Chem* **2003**,
75 (10), 2456–2462. https://doi.org/10.1021/ac020756b.

(60) Nanda, A.; Nasker, S. S.; Kushwaha, A. K.; Ojha, D. K.; Dearden, A. K.; Nayak, S. K.;
Nayak, S. Gold Nanoparticles Augment N-Terminal Cleavage and Splicing Reactions in
Mycobacterium Tuberculosis SufB. *Front Bioeng Biotechnol* **2021**, *9*, 773303.
https://doi.org/10.3389/FBIOE.2021.773303.

(61) Chemiluminescence, B.; Li, J.; Xing, H.; Wang, E.; Zhao, J.; Li, D.; Ma, J.; Suhartono, E.;
Thalib, I.; Aflanie, I.; Noor, Z.; Idroes, R. Study of Interaction between Cadmium and
Bovine Serum Albumin with UV-Vis Spectrocopy Approach. In *IOP Conference Series:*

Materials Science and Engineering; IOP Publishing, 2018; Vol. 350, p 012008.
https://doi.org/10.1088/1757-899X/350/1/012008.

(62)　Grzyska, P. K.; Hausinger, R. P.; Proshlyakov, D. A. Metal and Substrate Binding to an
Fe(II) Dioxygenase Resolved by UV Spectroscopy with Global Regression Analysis. *Anal*
Biochem **2010**, *399* (1), 64–71. https://doi.org/10.1016/J.AB.2009.11.022.

(63)　Plotnikova, O. A.; Mel'nikov, A. G.; Mel'nikov, G. V.; Gubina, T. I. Quenching of
Tryptophan Fluorescence of Bovine Serum Albumin under the Effect of Ions of Heavy
Metals. *Opt Spectrosc* **2016**, *120* (1), 65–69.
https://doi.org/https://doi.org/10.1134/S0030400X16010148.

(64)　Oliveira, A.; Baccana, N.; Cadore, S. Evaluation of Metal Ions in Rice Samples:
Extraction and Direct Determination by ICP OES. *J Braz Chem Soc* **2012**, *23* (5), 838–
845. https://doi.org/10.1590/S0103-50532012000500008.

(65)　Harrington, J. M.; Young, D. J.; Essader, A. S.; Sumner, S. J.; Levine, K. E. Analysis of
Human Serum and Whole Blood for Mineral Content by ICP-MS and ICP-OES:
Development of a Mineralomics Method. *Biol Trace Elem Res* **2014**, *160* (1), 132–142.
https://doi.org/10.1007/s12011-014-0033-5.

(66)　Punjabi, K.; Mehta, S.; Chavan, R.; Chitalia, V.; Deogharkar, D.; Deshpande, S.
Efficiency of Biosynthesized Silver and Zinc Nanoparticles against Multi-Drug Resistant
Pathogens. *Front Microbiol* **2018**, *9* (SEP), 376058.
https://doi.org/https://doi.org/10.3389/fmicb.2018.02207.

(67)　Longhin, E. M.; El Yamani, N.; Rundén-Pran, E.; Dusinska, M. The Alamar Blue Assay
in the Context of Safety Testing of Nanomaterials. *Frontiers in Toxicology* **2022**, *4*,
981701. https://doi.org/10.3389/FTOX.2022.981701.

(68)　Yang, L.; Zhang, Y.; Wang, F.; Luo, Z.; Guo, S.; Strähle, U. Toxicity of Mercury:
Molecular Evidence. *Chemosphere* **2020**, *245*, 125586.
https://doi.org/10.1016/J.CHEMOSPHERE.2019.125586.

(69)　Aljerf, L.; Almasri, N.; Access, O. Mercury Toxicity: Ecological Features of Organic
Phase of Mercury in Biota-Part I. *Inorganic Chemical Sciences* **2018**, *3* (3).
https://doi.org/10.32474/AOICS.2018.03.000157.

(70)　Budnik, L. T.; Casteleyn, L. Mercury Pollution in Modern Times and Its Socio-Medical
Consequences. *Science of The Total Environment* **2019**, *654*, 720–734.
https://doi.org/10.1016/J.SCITOTENV.2018.10.408.

(71)　Dash, H. R.; Mangwani, N.; Das, S. Characterization and Potential Application in
Mercury Bioremediation of Highly Mercury-Resistant Marine Bacterium Bacillus

1093   Thuringiensis PW-05. *Environmental Science and Pollution Research* **2014**, *21* (4), 2642–
1094   2653. https://doi.org/10.1007/s11356-013-2206-8.

1095 (72) Tang, W.; Zhang, W.; Zhao, Y.; Zhang, H.; Shan, B. Basin-Scale Comprehensive
1096   Assessment of Cadmium Pollution, Risk, and Toxicity in Riverine Sediments of the Haihe
1097   Basin in North China. *Ecol Indic* **2017**, *81*, 295–301.
1098   https://doi.org/10.1016/J.ECOLIND.2017.06.011.

1099 (73) White, P. J.; Brown, P. H. Plant Nutrition for Sustainable Development and Global
1100   Health. *Ann Bot* **2010**, *105* (7), 1073–1080. https://doi.org/10.1093/AOB/MCQ085.

1101 (74) Losada Ros, M. T.; Al-Enezi, E.; Cesarini, E.; Canonico, B.; Bucci, C.; Martins, M. V. A.;
1102   Papa, S.; Frontalini, F. Assessing the Cadmium Effects on the Benthic Foraminifer
1103   Ammonia Cf. Parkinsoniana: An Acute Toxicity Test. *Water (Basel)* **2020**, *12* (4), 1018.
1104   https://doi.org/10.3390/W12041018.

1105 (75) Dong, F.; Zhu, X.; Qian, W.; Wang, P.; Wang, J. Combined Effects of CO2-Driven Ocean
1106   Acidification and Cd Stress in the Marine Environment: Enhanced Tolerance of
1107   Phaeodactylum Tricornutum to Cd Exposure. *Mar Pollut Bull* **2020**, *150*, 110594.
1108   https://doi.org/10.1016/J.MARPOLBUL.2019.110594.

1109 (76) Atamaleki, A.; Sadani, M.; Raoofi, A.; Miri, A.; Bajestani, S. G.; Fakhri, Y.;
1110   Heidarinejad, Z.; Mousavi Khaneghah, A. The Concentration of Potentially Toxic
1111   Elements (PTEs) in Eggs: A Global Systematic Review, Meta-Analysis and Probabilistic
1112   Health Risk Assessment. *Trends Food Sci Technol* **2020**, *95*, 1–9.
1113   https://doi.org/10.1016/J.TIFS.2019.11.003.

1114 (77) Stankovic, S.; Kalaba, P.; Stankovic, A. R. Biota as Toxic Metal Indicators. *Environ*
1115   *Chem Lett* **2014**, *12* (1), 63–84. https://doi.org/10.1007/s10311-013-0430-6.

1116 (78) Joe, M. H.; Lee, K. H.; Lim, S. Y.; Im, S. H.; Song, H. P.; Lee, I. S.; Kim, D. H. Pigment-
1117   Based Whole-Cell Biosensor System for Cadmium Detection Using Genetically
1118   Engineered Deinococcus Radiodurans. *Bioprocess Biosyst Eng* **2012**, *35* (1–2), 265–272.
1119   https://doi.org/10.1007/s00449-011-0610-3.

1120 (79) Guertin, J.; Jacobs, J. A.; Avakian, C. P. *Chromium(VI) Handbook*, 1st ed.; CRC Press,
1121   2004. https://doi.org/10.1201/9780203487969.

1122 (80) Dawra, N.; Dabas, N. Advances in Spectrophotometric Determination of Chromium(III)
1123   and Chromium(VI) in Water: A Review. *Int J Environ Anal Chem* **2024**, *104* (13), 2994–
1124   3015. https://doi.org/10.1080/03067319.2022.2076224.

1125 (81) Sheehan, P. J.; Meyer, D. M.; Sauer, M. M.; Paustenbach, D. J. Assessment of the Human
1126   Health Risks Posed by Exposure to Chromium-contaminated Soils. *Journal of Toxicology*

and *Environmental Health, Part A Current Issues* **1991**, *32* (2), 161–201.
https://doi.org/10.1080/15287399109531476.

(82) Mishra, S.; Bharagava, R. N. Toxic and Genotoxic Effects of Hexavalent Chromium in Environment and Its Bioremediation Strategies. *Journal of Environmental Science and Health, Part C* **2016**, *34* (1), 1–32. https://doi.org/10.1080/10590501.2015.1096883.

(83) Feleafel, M. N.; Mirdad, Z. M. Hazard and Effects of Pollution by Lead on Vegetable Crops. *J Agric Environ Ethics* **2013**, *26* (3), 547–567. https://doi.org/10.1007/S10806-012-9403-1.

(84) Ghosh, I.; Sun, L.; Xu, M. Q. Zinc Inhibition of Protein Trans-Splicing and Identification of Regions Essential for Splicing and Association of a Split Intein*. *Journal of Biological Chemistry* **2001**, *276* (26), 24051–24058. https://doi.org/10.1074/JBC.M011049200.

(85) Tharappel, A. M.; Li, Z.; Li, H. Inteins as Drug Targets and Therapeutic Tools. *Front Mol Biosci* **2022**, *9*, 821146. https://doi.org/10.3389/FMOLB.2022.821146.

(86) Alhazmi, H. A.; Alam, M. S.; Albratty, M.; Najmi, A.; Abdulhaq, A. A.; Hassani, R.; Ahsan, W.; Qramish, A. N. Binding Investigation of Some Important Metal Ions Copper (I), Nickel (II), and Aluminium (III) with Bovine Serum Albumin Using Valid Spectroscopic Techniques. *J Chem* **2023**, *2023* (1), 2581653. https://doi.org/10.1155/2023/2581653.

(87) Permyakov, E. A.; Gonçalves, A. M. M. Metal Binding Proteins. *Encyclopedia* **2021**, *1* (1), 261–292. https://doi.org/10.3390/ENCYCLOPEDIA1010024.

(88) Woźnicka, O.; Rybarska, J.; Jagusiak, A.; Konieczny, L.; Stopa, B.; Roterman, I. Metal Ions Introduced to Proteins by Supramolecular Ligands. *Self-Assembled Molecules - New Kind of Protein Ligands: Supramolecular Ligands* **2017**, 61–76. https://doi.org/10.1007/978-3-319-65639-7_4.

(89) Antosiewicz, J. M.; Shugar, D. UV–Vis Spectroscopy of Tyrosine Side-Groups in Studies of Protein Structure. Part 2: Selected Applications. *Biophys Rev* **2016**, *8* (2), 163–177. https://doi.org/10.1007/s12551-016-0197-7.

(90) Biter, A. B.; Pollet, J.; Chen, W. H.; Strych, U.; Hotez, P. J.; Bottazzi, M. E. A Method to Probe Protein Structure from UV Absorbance Spectra. *Anal Biochem* **2019**, *587*, 113450. https://doi.org/10.1016/J.AB.2019.113450.

(91) Dieaconu, M.; Ioanid, A.; Iftimie, S.; Antohe, S. UV-ABSORPTION MECHANISMS OF Ni 2+-BINDING BOVINE SERUM ALBUMIN. *Dig J Nanomater Biostruct 7* (3), 1125–1138.

(92)   *View of Study of immunoglobulin 'G' with ultra violet spectroscopy in Duchene muscular dystrophy and Alzheimer's disease.* https://journals.scholarpublishing.org/index.php/BJHR/article/view/100/98 (accessed 2025-06-16).

(93)   Bjørklund, G.; Dadar, M.; Mutter, J.; Aaseth, J. The Toxicology of Mercury: Current Research and Emerging Trends. *Environ Res* **2017**, *159*, 545–554. https://doi.org/10.1016/J.ENVRES.2017.08.051.

(94)   Ajsuvakova, O. P.; Tinkov, A. A.; Aschner, M.; Rocha, J. B. T.; Michalke, B.; Skalnaya, M. G.; Skalny, A. V.; Butnariu, M.; Dadar, M.; Sarac, I.; Aaseth, J.; Bjørklund, G. Sulfhydryl Groups as Targets of Mercury Toxicity. *Coord Chem Rev* **2020**, *417*, 213343. https://doi.org/10.1016/J.CCR.2020.213343.

(95)   Sall, M. L.; Diaw, A. K. D.; Gningue-Sall, D.; Efremova Aaron, S.; Aaron, J. J. Toxic Heavy Metals: Impact on the Environment and Human Health, and Treatment with Conducting Organic Polymers, a Review. *Environmental Science and Pollution Research* **2020**, *27* (24), 29927–29942. https://doi.org/10.1007/s11356-020-09354-3.

(96)   Gonick, H. C. Lead-Binding Proteins: A Review. *J Toxicol* **2011**, *2011* (1), 686050. https://doi.org/10.1155/2011/686050.

(97)   Fossum, C. J.; Johnson, B. O. V.; Golde, S. T.; Kielman, A. J.; Finke, B.; Smith, M. A.; Lowater, H. R.; Laatsch, B. F.; Bhattacharyya, S.; Hati, S. Insights into the Mechanism of Tryptophan Fluorescence Quenching Due to Synthetic Crowding Agents: A Combined Experimental and Computational Study. *ACS Omega* **2023**, *8* (47), 44820–44830. https://doi.org/10.1021/acsomega.3c06006.

(98)   Liu, Y.; Xu, J.; Han, L.; Liu, Q.; Yang, Y.; Li, Z.; Lu, Z.; Zhang, H.; Guo, T.; Liu, Q. Theoretical Research on Excited States: Ultraviolet and Fluorescence Spectra of Aromatic Amino Acids. *Interdiscip Sci* **2020**, *12* (4), 530–536. https://doi.org/10.1007/s12539-020-00395-3.

(99)   Yammine, A.; Gao, J.; Kwan, A. H. Tryptophan Fluorescence Quenching Assays for Measuring Protein-Ligand Binding Affinities: Principles and a Practical Guide. *Bio Protoc* **2019**, *9* (11). https://doi.org/10.21769/BIOPROTOC.3253.

(100) Zheng, Z.; Kawakami, K.; Zhang, D.; Negishi, L.; Abomosallam, M.; Asakura, T.; Nagata, K.; Suzuki, M. Identification and Functional Analysis of Cadmium-Binding Protein in the Visceral Mass of Crassostrea Gigas. *Sci Rep* **2021**, *11* (1), 1–14. https://doi.org/10.1038/s41598-021-90882-4.

1193   (101) Springfield, E.; Willis, A.; Merle, J.; Mazlo, J.; Ngu-Schwemlein, M. Spectroscopic and
1194   Theoretical Studies of Hg(II) Complexation with Some Dicysteinyl Tetrapeptides.
1195   *Bioinorg Chem Appl* **2021**, *2021* (1), 9911474. https://doi.org/10.1155/2021/9911474.

1196   (102) Crouse, H. F.; Petrunak, E. M.; Donovan, A. M.; Merkle, A. C.; Swartz, B. L.; Basu, S.
1197   Static and Dynamic Quenching of Tryptophan Fluorescence in Various Proteins by a
1198   Chromium (III) Complex. *Spectroscopy Letters* **2011**, *44* (5), 369–374.
1199   https://doi.org/10.1080/00387010.2010.546470.

1200   (103) Shrivastava, H. Y.; Nair, B. U. Fluorescence Resonance Energy Transfer from Tryptophan
1201   to a Chromium(III) Complex Accompanied by Non-Specific Cleavage of Albumin: A
1202   Step Forward towards the Development of a Novel Photoprotease. *J Inorg Biochem* **2004**,
1203   *98* (6), 991–994. https://doi.org/10.1016/J.JINORGBIO.2004.02.014.

1204   (104) Liu, Y.; Zhang, L.; Liu, R.; Zhang, P. Spectroscopic Identification of Interactions of Pb 2+
1205   with Bovine Serum Albumin. *J Fluoresc* **2012**, *22* (1), 239–245.
1206   https://doi.org/10.1007/s10895-011-0950-z.

1207   (105) Gen-hua Wu. *Study on the interaction of Pb2+ and bovine serum albumin by*
1208   *fluorescence*. https://europepmc.org/article/med/15852867 (accessed 2025-06-16).

1209   (106) Luis, G.; Rubio, C.; Revert, C.; Espinosa, A.; González-Weller, D.; Gutiérrez, A. J.;
1210   Hardisson, A. Dietary Intake of Metals from Yogurts Analyzed by Inductively Coupled
1211   Plasma Optical Emission Spectrometry (ICP-OES). *Journal of Food Composition and*
1212   *Analysis* **2015**, *39*, 48–54. https://doi.org/10.1016/J.JFCA.2014.11.013.

1213   (107) Wiltsche, H.; Wolfgang, M. Merits of Microwave Plasmas for Optical Emission
1214   Spectrometry – Characterization of an Axially Viewed Microwave-Sustained, Inductively
1215   Coupled, Atmospheric-Pressure Plasma (MICAP). *J Anal At Spectrom* **2020**, *35* (10),
1216   2369–2377. https://doi.org/10.1039/D0JA00293C.

1217   (108) Baraud, F.; Zaiter, A.; Porée, S.; Leleyter, L. New Approach for Determination of Cd, Cu,
1218   Cr, Ni, Pb, and Zn in Sewage Sludges, Fired Brick, and Sediments Using Two Analytical
1219   Methods by Microwave-Induced Plasma Optical Spectrometry and Induced Coupled
1220   Plasma Optical Spectrometry. *SN Appl Sci* **2020**, *2* (9), 1–10.
1221   https://doi.org/10.1007/s42452-020-03220-0.

1222   (109) Wegener Antje; Marc Wiesmann; Peter Luschas; Kerstin Vogel; Matthias Pursch. *SEC-*
1223   *ICP-OES Hyphenation: Speciation and Quantification of Polydimethylsiloxanes at Trace*
1224   *levels*. Spectroscopy. https://www.spectroscopyonline.com/view/sec-icp-oes-hyphenation-
1225   speciation-and-quantification-polydimethylsiloxanes-trace-levels (accessed 2025-08-01).

(110) Pandey, M.; Talwar, S.; Bose, S.; Pandey, A. K. Iron Homeostasis in Mycobacterium
Tuberculosis Is Essential for Persistence. *Sci Rep* **2018**, *8* (1), 1–9.
https://doi.org/10.1038/S41598-018-35012-3.

(111) Tripathi, A.; Anand, K.; Das, M.; O'Niel, R. A.; Sabarinath, P. S.; Thakur, C.; Reddy, R.
R. L.; Rajmani, R. S.; Chandra, N.; Laxman, S.; Singh, A. Mycobacterium Tuberculosis
Requires SufT for Fe-S Cluster Maturation, Metabolism, and Survival in Vivo. *PLoS*
Pathog **2022**, *18* (4), e1010475. https://doi.org/10.1371/JOURNAL.PPAT.1010475.

(112) Saini, V.; Farhana, A.; Glasgow, J. N.; Steyn, A. J. C. Iron Sulfur Cluster Proteins and
Microbial Regulation: Implications for Understanding Tuberculosis. *Curr Opin Chem Biol*
2012, *16* (1–2), 45–53. https://doi.org/10.1016/J.CBPA.2012.03.004.

(113) Zhang, L.; Xiao, N.; Pan, Y.; Zheng, Y.; Pan, Z.; Luo, Z.; Xu, X.; Liu, Y. Binding and
Inhibition of Copper Ions to RecA Inteins from Mycobacterium Tuberculosis. *Chemistry –*
A European Journal **2010**, *16* (14), 4297–4306.
https://doi.org/10.1002/CHEM.200903584.

(114) Shaloam R, D. Preclinical Assessment of Low Doses of Cisplatin in the Management of
Acute Promyelocytic Leukemia. *Int J Cancer Res Mol Mech* **2015**, *1* (3).
https://doi.org/10.16966/2381-3318.113.

(115) Topilina, N. I.; Green, C. M.; Jayachandran, P.; Kelley, D. S.; Stanger, M. J.; Piazza, C.
L.; Nayak, S.; Belfort, M. SufB Intein of Mycobacterium Tuberculosis as a Sensor for
Oxidative and Nitrosative Stresses. *Proc Natl Acad Sci U S A* **2015**, *112* (33), 10348–
10353. https://doi.org/10.1073/pnas.1512777112.

(116) Lennon, C. W.; Wahl, D.; Goetz, J. R.; Joel Weinberger, I. Reactive Chlorine Species
Reversibly Inhibit DnaB Protein Splicing in Mycobacteria. *Microbiol Spectr* **2021**, *9* (2).
https://doi.org/10.1128/SPECTRUM.00301-21.

(117) Wang, P.; Xu, G.; Qin, L.; Xu, Y.; Li, Y.; Li, R. Cell-Based Biosensors and Its
Application in Biomedicine. *Sens Actuators B Chem* **2005**, *108* (1–2), 576–584.
https://doi.org/10.1016/J.SNB.2004.11.056.

(118) Gupta, N.; Renugopalakrishnan, V.; Liepmann, D.; Paulmurugan, R.; Malhotra, B. D.
Cell-Based Biosensors: Recent Trends, Challenges and Future Perspectives. *Biosens*
Bioelectron **2019**, *141*, 111435. https://doi.org/10.1016/J.BIOS.2019.111435.

(119) Perler, F. B. InBase: The Intein Database. *Nucleic Acids Res* **2002**, *30* (1), 383–384.
https://doi.org/10.1093/NAR/30.1.383.

(120) Yakovleva, L.; Shuman, S. Mycobacterium Smegmatis SftH Exemplifies a Distinctive
Clade of Superfamily II DNA-Dependent ATPases with 3′ to 5′ Translocase and Helicase

Activities. *Nucleic Acids Res* **2012**, *40* (15), 7465–7475.
https://doi.org/10.1093/NAR/GKS417.

(121) Ordonez, H.; Shuman, S. Mycobacterium Smegmatis Lhr Is a DNA-Dependent ATPase
and a 3′-to-5′ DNA Translocase and Helicase That Prefers to Unwind 3′-Tailed
RNA:DNA Hybrids. *Journal of Biological Chemistry* **2013**, *288* (20), 14125–14134.
https://doi.org/10.1074/JBC.M113.466854.

(122) Uson, M. L.; Ordonez, H.; Shuman, S. Mycobacterium Smegmatis HelY Is an RNA-
Activated ATPase/DATPase and 3′-to-5′ Helicase That Unwinds 3′-Tailed RNA Duplexes
and RNA:DNA Hybrids. *J Bacteriol* **2015**, *197* (19), 3057–3065.
https://doi.org/10.1128/JB.00418-15.

(123) Rampersad, S. N. Multiple Applications of Alamar Blue as an Indicator of Metabolic
Function and Cellular Health in Cell Viability Bioassays. *Sensors* **2012**, *12* (9), 12347–
12360. https://doi.org/10.3390/S120912347.

(124) O'Brien, J.; Wilson, I.; Orton, T.; Pognan, F. Investigation of the Alamar Blue (Resazurin)
Fluorescent Dye for the Assessment of Mammalian Cell Cytotoxicity. *Eur J Biochem*
2000, *267* (17), 5421–5426. https://doi.org/10.1046/j.1432-1327.2000.01606.x.

(125) McMillan, J. E. *Principles of Analytical Validation*; Elsevier Inc., 2013.
https://doi.org/10.1016/B978-0-444-59378-8.00011-6.

(126) Howard, J. A.; Kuznietsova, H.; Dziubenko, N.; Aigle, A.; Natuzzi, M.; Thomas, E.;
Lysenko, V.; David, L.; Brichart, T.; Lux, F.; Tillement, O. Combating Lead and
Cadmium Exposure with an Orally Administered Chitosan-Based Chelating Polymer. *Sci
Rep* **2023**, *13* (1), 1–14. https://doi.org/10.1038/s41598-023-28968-4.

(127) Saleh, S. M.; El-Tawil, O. S.; Mahmoud, M. B.; Abd El-Rahman, S. S.; El-Saied, E. M.;
Noshy, P. A. Do Nanoparticles of Calcium Disodium EDTA Minimize the Toxic Effects
of Cadmium in Female Rats? *Biol Trace Elem Res* **2024**, *202* (5), 2228–2240.
https://doi.org/10.1007/s12011-023-03842-6.

(128) Flora, S. J. S.; Pachauri, V. Chelation in Metal Intoxication. *Int J Environ Res Public
Health* **2010**, *7* (7), 2745–2788. https://doi.org/10.3390/IJERPH7072745.

(129) Bethesda. *LiverTox*; National Institute of Diabetes and Digestive and Kidney Diseases,
2012.

(130) George, G. N.; Prince, R. C.; Gailer, J.; Buttigieg, G. A.; Denton, M. B.; Harris, H. H.;
Pickering, I. J. Mercury Binding to the Chelation Therapy Agents DMSA and DMPS and
the Rational Design of Custom Chelators for Mercury. *Chem Res Toxicol* **2004**, *17* (8),
999–1006. https://doi.org/10.1021/tx049904e.

1294   (131) Aaseth, J.; Ajsuvakova, O. P.; Skalny, A. V.; Skalnaya, M. G.; Tinkov, A. A. Chelator
1295   Combination as Therapeutic Strategy in Mercury and Lead Poisonings. *Coord Chem Rev*
1296   **2018**, *358*, 1–12. https://doi.org/10.1016/J.CCR.2017.12.011.

1297   (132) Martí-Baena, Q.; Pascuet-Fontanet, A.; Berjaga-Buisan, T.; Caravaca-Rodríguez, M.;
1298   Puig-Costa-Jussà, J.; Sanchez-Mejias, A.; Ivančić, D.; Mogas-Díez, S.; Güell, M.; Macia,
1299   J. Intein-Mediated Thyroid Hormone Biosensors: Towards Controlled Delivery of
1300   Hormone Therapy. *Frontiers in Systems Biology* **2024**, *4*, 1270071.
1301   https://doi.org/10.3389/FSYSB.2024.1270071.

1302   (133) Chahal, H. K.; Dai, Y.; Saini, A.; Ayala-Castro, C.; Outten, F. W. The SufBCD Fe−S
1303   Scaffold Complex Interacts with SufA for Fe−S Cluster Transfer. *Biochemistry* **2009**, *48*
1304   (44), 10644–10653. https://doi.org/10.1021/BI901518Y.

1305   (134) Dillon, N. A.; Lamont, E. A.; Rather, M. A.; Baughn, A. D. Oxidative Stress Drives
1306   Potent Bactericidal Activity of Pyrazinamide against Mycobacterium Tuberculosis. *Elife*
1307   **2025**, *14*. https://doi.org/10.7554/ELIFE.105614.1.

1308   (135) Zhang, M.; Gong, J.; Lin, Y.; Barnes, P. F. Growth of Virulent and
1309   AvirulentMycobacterium Tuberculosis Strains in Human Macrophages. *Infect Immun*
1310   **1998**, *66* (2), 794–799. https://doi.org/10.1128/IAI.66.2.794-799.1998.

1311   (136) *Revised National TB Control Programme Training Manual for Mycobacterium*
1312   *tuberculosis*.

1313   (137) Davis, E. O.; Sedgwick, S. G.; Colston, M. J. Novel Structure of the RecA Locus of
1314   Mycobacterium Tuberculosis Implies Processing of the Gene Product. *J Bacteriol* **1991**,
1315   *173* (18), 5653–5662. https://doi.org/10.1128/JB.173.18.5653-5662.1991.

1316   (138) Saves, I.; Westrelin, F.; Daffé, M.; Masson, J. M. Identification of the First Eubacterial
1317   Endonuclease Coded by an Intein Allele in the Pps1 Gene of Mycobacteria. *Nucleic Acids*
1318   *Res* **2001**, *29* (21), 4310–4318. https://doi.org/10.1093/NAR/29.21.4310.

1319   (139) Schneider, R. F.; Hallstrom, K.; DeMott, C.; McDonough, K. A. Conditional Protein
1320   Splicing of the Mycobacterium Tuberculosis RecA Intein in Its Native Host. *Sci Rep*
1321   **2024**, *14* (1), 1–14. https://doi.org/10.1038/S41598-024-71248-Y.

1322

1323

i)

	Zn²⁺			Cu²⁺			Pt⁴⁺			Cd²⁺	Hg²⁺	
	H37Ra		*M. sm*	H37Ra		*M. sm*	H37Ra		*M. sm*	H37Ra	H37Ra	

SufB Intein

(i)

Live cells

Functional SufB protein

Splicing of *Mtb* SufB precursor protein within viable cells

Alamar Reagent

Mitochondrial reductase

- Cadmium (Cd²⁺)

- Mercury (Hg²⁺)

- TME

(ii)

Dead cells

Splicing inhibition within non-viable cells

Alamar Reagent

Mitochondrial reductase