Sustained dynamics of saccadic inhibition and adaptive oculomotor responses

during continuous exploration

Abbreviated title: Saccadic Inhibition and Oculomotor Adaptation

Celeste Cafaro[1#], Giovanni Cirillo[2#], Giovanna Vermiglio[3], Carlo Cavaliere[4], Alessio

Fracasso[5,6*], Antimo Buonocore[1*]

[1]Department of Educational, Psychological and Communication Sciences, Suor Orsola

Benincasa University, 80135, Naples, Italy;

[2]Division of Human Anatomy - Neuronal Networks Morphology and Systems Biology

Lab, Department of Mental, Physical Health and Preventive Medicine, University of

Campania "Luigi Vanvitelli", 80138, Naples, Italy;

[3]Department of Biomedical, Dental, Morphological and Functional Imaging Sciences,

University of Messina, 98122, Messina, Italy;

[4]IRCCS Synlab SDN, 80143, Naples, Italy;

[5]Department of Psychology, University of Padua, 35131, Padua, Italy;

[6]School of Psychology and Neuroscience, University of Glasgow, Glasgow, G12 8QE5,

Scotland, UK;

[#] Co-first authorship.

* Co-last authorship.

Correspondence: antimo.buonocore@unisob.na.it

Number of pages: 42

Number of figures: 6

Number of words for abstract (250), introduction (726), and discussion (1494)

Conflict of interest statement: The authors declare no competing financial interests.

Acknowledgments:

A.B., C.C., G.C., and G.V are supported by a grant from the Italian Minister of Research and University (PRIN 2022_PNRR-P2022ST78T). C.C. and G.C. were supported by #NEXTGENERATIONEU (NGEU) and funded by the Ministry of University and Research (MUR), National Recovery and Resilience Plan (NRRP), project MNESYS (PE0000006) – A Multiscale integrated approach to the study of the nervous system in health and disease (DN. 1553 11.10.2022). A.F. was supported by a grant from the Biotechnology and Biology Research Council (BBSRC, grant number: BB/S006605/1) and the Bial Foundation (Bial Foundation Grants Program; Grant id: A-29315, number: 203/2020, grant edition: G-15516). The authors used AI-assisted technology for grammar checking of the manuscript text.

Abstract

In natural environments, stimuli often recur across time and space, requiring the visual system to remain sensitive to novelty while managing predictability. A central question in systems neuroscience is how motor systems adapt to repeated sensory events without compromising responsiveness. We investigated this adaptive capacity using saccadic inhibition (SI)—a reflexive suppression of eye movements triggered by sudden visual transients—as a probe of oculomotor dynamics during naturalistic viewing. Human participants (N = 21) freely explored visual arrays while brief gaze-contingent flashes appeared five times at random intervals, either foveally or parafoveally. SI reliably occurred ~120 ms post-flash across repetitions and locations, indicating robust sensory-driven inhibition. However, the rebound phase—reflecting saccade reprogramming—showed a progressive decline. In a second experiment (N = 19), only the first or the fifth flash was visible on each trial. In this case, neither inhibition nor rebound was altered, suggesting that the rebound decline is driven by repeated sensory stimulation rather than exploration time. This dissociation reveals selective habituation of motor re-engagement mechanisms, while reflexive inhibitory gating remains stable. We propose that inhibition is mediated by circuitry that transiently suppresses saccade initiation and resists habituation. By contrast, the weakening rebound reflects a separate, habituation-prone route that reduces saccade generation to irrelevant events. Functionally, this imbalance implies a recalibration within the saccade generator, preserving inhibitory capacity while constraining motor output. Our findings uncover a distinct form of oculomotor habituation and demonstrate how SI reveals dynamic decoupling of sensory input and motor output under repeated stimulation.

Keywords: Saccades, Saccadic inhibition, Adaptation, Habituation, Free viewing task,

Brainstem

Significance Statement

The ability to interrupt and resume eye movements in response to environmental

changes is fundamental to visual exploration. We investigated how this process unfolds

across repeated visual transients in naturalistic conditions. Our findings show that

saccadic inhibition remains stable across time, whereas the subsequent rebound phase

habituates. This dissociation suggests that distinct processes mediate reflexive

interruption and habituation of motor recovery. Our data point to the dynamic

modulation of motor gating circuits as a mechanism for optimizing oculomotor behavior.

These results deepen our understanding of visual-motor coordination, provide insight

about the constrains governing the underlying system circuitry, and may inform clinical

tools for assessing sensorimotor adaptability in health and disease.

Introduction

Exploring the environment is essential for survival, requiring the nervous system to continuously integrate dynamic sensory input with motor plans (Hayhoe, 2017). Vision plays a dominant role in this process, and eye movements serve to actively sample the visual world, redirecting gaze toward relevant stimuli. While these movements are often goal-directed, they are also shaped by sudden external events. The visual system must therefore remain responsive to novelty while adapting to environmental regularities (Pascucci et al., 2019). Within this perception–action loop (Fuster, 2004), repeated stimulation is common. In response, the brain might engage sensory adaptation—a largely automatic reduction in neural responsiveness to persistent or fast recurring inputs—that enhances coding efficiency and aligns perception with context (Webster, 2011; Solomon and Kohn, 2014).

Adaptation occurs across various visual features, including contrast, orientation, motion, and complex forms like faces (Sekuler and Ganz, 1963; Blakemore and Campbell, 1969; Georgeson and Harris, 1984; Greenlee and Heitger, 1988; Anstis et al., 1998; Webster and MacLeod, 2011). From a neural perspective, adaptation reduces firing in repeatedly stimulated circuits, from the retina to higher-order areas (Smirnakis et al., 1997; Kohn, 2007; Benucci et al., 2013; Torrado Pacheco et al., 2019). Oculomotor areas also adapt: the superior colliculus (SC) shows attenuated responses to repeated transients (Boehnke et al., 2011; Basso and May, 2017), and the frontal eye fields (FEF) exhibit strong adaptation to sequential flashes (Bruce and Goldberg, 1985; Mayo and Sommer, 2008).

External events can also recur over longer time scales. Whereas fast adaptation reflects transient sensory adjustments, habituation denotes a slower, learned reduction in behavioral responses to repeated, irrelevant stimuli (Boehnke et al., 2011). Unlike adaptation, habituation engages higher-order modulation shaped by attention, expectation, and context, thereby supporting goal-directed behavior in environments rich with distractors (Burleson et al., 2025). A central question in systems neuroscience is how motor circuits respond to repeated sensory stimulation, balancing responses subject to adaptation or habituation with the need to react rapidly and flexibly to external stimuli.

Saccadic inhibition (SI) provides a paradigm to probe visuomotor control: an abrupt visual transient suppresses eye movements ~90 ms later, followed by a rebound at 120–150 ms reflecting motor reprogramming (Reingold and Stampe, 1999, 2002, 2004; Buonocore and McIntosh, 2008; Edelman and Xu, 2009; Bompas and Sumner, 2011). This inhibitory–rebound cycle is shaped by stimulus and task factors (e.g., location, salience, attention, content) (Bompas and Sumner, 2011; Buonocore and McIntosh, 2012, 2013; Buonocore et al., 2017a; Taylor et al., 2024) and extends beyond saccades to microsaccades, smooth pursuit, and even fixation drift (Rolfs et al., 2008; Kerzel et al., 2010; Buonocore et al., 2017b; Buonocore et al., 2019; Malevich et al., 2020). Together, these findings identify reflex-like inhibition as a unifying mechanism at the core of oculomotor control.

Neurally, SI is thought to act downstream of the SC, potentially involving omnipause neurons (OPNs) that gate saccade initiation (Buonocore and Hafed, 2023). Subcortical structure such as the SC and cortical intervention of FEF, LIP and other

motor areas might support the reinstatement of the eye movements. While SI has been

well studied in isolated trials, its evolution under repeated stimulation—as it might occur

during natural viewing—remains poorly understood. Evidence indicates that the visual

component driving inhibition adapts to rapidly presented visual stimuli, resulting in

reduced effects (Shan and Edelman, 2023). However, it is not yet established how the

oculomotor system responds to repeated flashes over longer time scales that more

closely approximate natural viewing conditions.

We hypothesize that SI reflects two distinct mechanisms: a fast, visually driven

inhibition phase subject to short-timescale adaptation (Shan and Edelman, 2023), and a

motor-driven rebound phase vulnerable to slower habituation. Rapid successive

transients may reduce inhibition through sensory adaptation, whereas at longer intervals

(~1 s) inhibition remains stable while the motor system disengages from post-inhibition

recovery. In this study, participants freely explored visual scenes while gaze-contingent

transients appeared foveally or parafoveally. If inhibition and rebound were tightly

coupled, both would decline together, implying a single governing process. Instead,

evidence for their dissociation would support independent mechanisms: sensory

processes regulating inhibitory strength and motor processes regulating recovery

dynamics, enabling selective tuning of responsiveness without uniformly suppressing

downstream motor output. By dissociating these components of oculomotor behavior,

our findings advance understanding of how the brain balances responsiveness with

efficiency under repeated interference.

Materials and methods

Participants

Experiments 1 and 2, were eye tracking experiments on human participants. Experiment 1 included a sample of twenty-one participants (17 females and 4 males) while Experiment 2, included a sample of nineteen participants (14 females and 5 males). All participants had normal or corrected-to-normal vision, reported no neurological or visual impairments, and were naïve to the purpose of the experiment. The sample comprised university students from the Department of Educational, Psychological and Communication Sciences from the Psychology Degree Course of Suor Orsola Benincasa University. Written informed consent was obtained from all participants in compliance with the 2024 Declaration of Helsinki. The study received ethical approval from the local ethics committee of Suor Orsola Benincasa University, Naples.

Apparatus and stimuli

In both Experiment 1 and 2, participants were seated with their eyes aligned to the center of the display at a viewing distance of 65 cm, with their heads stabilized using a chin and forehead rest, to avoid head movements and ensure that the eyes were positioned approximately in front of the center of the screen. Experimental sessions were conducted in a dark room, illuminated only by the experimental display monitor and the operator's monitor, which was positioned behind the participant and facing

away from them. Eye movements were recorded using a LiveTrack Lightning eye tracker (Cambridge Research System) sampling both eyes at 500 Hz and determining the direction of gaze based on the pupil and Purkinje centers. Before the start of the experiment, a binocular calibration was performed using nine locations arranged in a three-by-three grid. The calibration procedure was performed with the LiveTrack Viewer software and calibrations were accepted if the error for each eye was below 0.5 degrees of visual angle. Both eyes were then tracked throughout the entire experiment. The experimental protocol and stimuli were generated using MATLAB R2023a (MathWorks, Inc.) in combination with Psychophysics Toolbox 3.0.14 (Brainard, 1997; Pelli, 1997). All stimuli were presented in black on a mid-gray background on a 27-inch LCD monitor with a resolution of 1920 by 1080 pixels and a refresh rate of 144 Hz (6.9 ms). A small blue fixation dot, with a radius of about 0.30 degrees (10 pixels) served as a fixation point. Stimuli consisted of screens displaying randomly generated geometric figures. Depending on the task (see Procedure), each screen contained either horizontal/vertical lines or circles/squares presented in completely random positions. All the geometric shapes were about two degrees in size. The number of elements was randomly determined on each trial between 19 and 29. In particular, one display might contain more horizontal than vertical lines (or vice versa), with a difference in number from 1 to 3 elements. The same logic applied to circles and squares. During the stimulus presentation, a black circle two degrees in size appeared at various retinal locations, either within the fovea (Experiment 1 and 2) or 5.6 degrees in the periphery (only Experiment 1), always on the left side of the fovea. Manual responses to the shape

number estimation were collected on a five-button USB response pad from The Black

Box Toolkit.

Procedure

Both experiments consisted of four blocks, each comprising two tasks: lines (Task one)

or shape (Task two) number estimation. At the beginning of each block, a brief on-

screen sentence informed participants whether the upcoming comparison on the

number of elements would involve lines or shapes. In each task, participants were

required to explore the visual display and make a judgment about the numerosity of

geometric items. Specifically, participants were instructed to scan the entire screen to

estimate whether the display contained more vertical or horizontal lines (Task one), or

more circles than squares (Task two). The sequence of tasks was randomized across

blocks for each participant. Participants initiated each task by pressing the central

button on a five-button response box. On each trial, they were first asked to fixate on a

blue dot presented at the center of the screen. After a variable delay between 1 to 1.5

seconds from fixation onset, a stimulus array was displayed for approximately 12

seconds. During the visual exploration phase, brief transient circles were presented for

80 ms, contingent on gaze position. In Experiment 1, each trial included five such

transients, with an inter-stimulus interval (ISI) ranging from 0.7 to 1.2 seconds.

Depending on the condition, the transient circle was presented either at the fovea or at

a fixed eccentricity along the horizontal meridian, about 5.6 degrees away from the

fovea (see Apparatus and stimuli). In Experiment 2, the transient circle appeared either

as the first or fifth stimulus, according to the experimental condition. The remaining four circles were flagged as "invisible", meaning that a stimulus trigger was only sent for time alignment (See Experimental design and statistical analysis).

In both experiments, following the stimulus presentation, a response screen appeared. Participants were asked to indicate whether they had seen more circles or more squares, or more horizontal than vertical lines, using two designated buttons on the response box. The left button was used to report more circles or more vertical lines, while the right button was used for more squares or more horizontal lines. No time limit was imposed for responses. After a response was given, the next trial began automatically. The experiments comprised 120 trials, organized into four blocks of 30 trials each, with 60 trials dedicated to each geometric task. In Experiment 1, across the entire experiment, 600 flashes were presented—300 at foveal and 300 at parafoveal locations—with 60 flashes delivered at each of five flash orders. In Experiment 2, a total of 120 visible flashes were presented, 60 flashes presented as first and 60 flashes presented as fifth. The remaining 480 flashes were classified as "invisible". A break was provided after the first two blocks to allow participants to rest their eyes. The full session lasted approximately 30 minutes, including the break. Participants resumed the experiment by pressing the central button on the response box. At the end of the session, all participants were debriefed about the purpose of the study.

Figure 1. Procedure and stimuli used in the experiment.

Participants fixated on a central dot for a variable duration (1 to1.5 s). A display containing geometric shapes, either lines – Task 1, or circle/square – Task 2, was then presented for about 12 s. During visual exploration, a gaze-contingent black circle flashed for 80 ms either five times per trial (Experiment 1) or it was visible only on the first or fifth presentations (Experiment 2). At the end of each trial, participants reported the number of figures observed using a button box. In Experiment 1, the transient stimulus could appear either at the fovea (blue) or 5.6 degrees of visual angle along the horizontal meridian in the parafovea (red), to the left of fixation. In Experiment 2, the transient was always presented at the fovea.

Data preprocessing

For both experiments, continuous raw eye position data (recorded in Fick coordinates: longitude and azimuth) were segmented into trials and stored as matrices. Eye velocity was computed using a custom band-limited differentiator filter (Hafed and Chen, 2016), which combines smoothing and numerical differentiation via a symmetric kernel. Acceleration was derived as the temporal derivative of the velocity signal. Saccades were detected automatically based on velocity and acceleration thresholds of 30 degrees per second and 900 degrees per second square. Each trial was then manually inspected to correct false positives and missed detections. Blinks, identified by abrupt

discontinuities in eye position, were manually marked and excluded. Saccade amplitude

was calculated as the Euclidean distance of displacement across horizontal and vertical

components. Peak velocities were calculated as the maximum of the velocity trace.

Experimental design and statistical analysis

To examine how saccadic responses evolved across repeated visual transients, we first

computed saccadic inhibition profiles separately for each condition (foveal vs.

parafoveal flash) and flash order (flash 1 to 5). For saccadic inhibition profile, we refer to

the change in saccade rate, measured in saccades per seconds (Hz), relative to the

onset of the visual transient (Reingold and Stampe, 2002; Buonocore and McIntosh,

2008). First, we extracted all saccades within each trial and aligned them relative to

flash onset, by subtracting the time of each flash from the time of each saccade.

Saccades with a peak velocity exceeding 600 degrees per second (Experiment 1:

0.37%; Experiment 2: 0.38%) and those with an amplitude greater than 20 degrees of

visual angle (Experiment 1: 2.45%; Experiment 2: 2.42%) were excluded from the rest

of the analysis. Next, for each trial, we constructed histograms of saccade frequency

within a time window spanning from 200 milliseconds before to 600 milliseconds after

each flash onset, using a bin width of 10 milliseconds. These histograms were then

converted into saccade rates per second by dividing the counts by the bin width and

multiplying by 1000. The resulting histograms were grouped by participant and

condition, distinguishing between foveal and parafoveal flashes (Experiment 1), and

between flashes one through five (Experiment 1 and 2). In Experiment 2, we also

constructed histograms for "invisible" flashes by aligning the saccades generated during

visual exploration with the timestamp of the so called "invisible" flash, that was not drawn on screen. This procedure allowed to visualize the flat distribution of saccades aligned the trigger event. For each participant and condition, the histograms were averaged and smoothed using the MATLAB function "smoothdata" with the parameter "movemean" and a window of seven samples. This smoothing procedure corresponds to a moving average over 70 milliseconds, centered on the fourth sample, and advanced in 10-millisecond steps. Edge effects were handled automatically by the function, which initiates full smoothing from the fourth sample onward. At the end of preprocessing, each participant contributed ten smoothed histograms—one for each flash-location and flash order condition in Experiment 1, and for flash visibility and flash order condition in Experiment 2. The averaged histograms were used to extract key temporal and dynamic features of the saccadic response.

To quantitatively assess the dynamics of saccadic inhibition, we extracted a set of descriptive parameters, means and standard deviations, from the saccade histograms of each participant and condition during critical time periods of 50 ms durations. These parameters captured both the overall magnitude of the response and the timing of key features in the inhibition profile. In Experiment 1, we first measured the baseline saccade frequency (i.e. *Baseline frequency*), defined as the average rate of saccades in the pre-flash interval from −100 to −50 ms, which served as a reference for ongoing oculomotor activity before the onset of the transient. Following the flash, we identified the minimum saccade frequency (i.e. *Inhibition frequency*) within a predefined window (90 to 140 ms), reflecting the depth of the inhibitory response, as well as the latency to this minimum (i.e. *Inhibition latency*), indexing the time point at which

inhibition reached its peak. We then turned to the rebound phase, capturing the transient increase in saccade frequency that typically follows inhibition (i.e. *Rebound frequency*). Here, we measured the maximum rebound frequency within a post-inhibition window (200 to 250 ms). In Experiment 2, all the parameters were the same, but we observed a small shift of about 20 ms in the saccadic inhibition profile. For this reason, to better capture inhibition and rebound frequencies, we computed the former between 110 and 160 ms and the latter between 240 and 290 ms after flash onset. All these temporal windows are visually indicated by shaded rectangles in Figures 4 and 5, providing a structured summary of the key stages of the saccadic inhibition response alongside a table with all the measures extracted in Experiment 4, 5 and 6.

Each of these dependent variables (*Inhibition frequency, Inhibition latency, Rebound frequency*) was analyzed using linear mixed-effects models (LME). In Experiment 1, fixed effects included *Flash location* (foveal versus parafoveal), *Flash order* (from the first to the fifth flash), their interaction, and the pre-flash *Baseline frequency* as continuous covariate. A random intercept for each Participant was included to account for inter-individual variability. For the dependent variables *Inhibition frequency*, *Inhibition latency* and *Rebound frequency* the general model structure was as follows, in Wilkinson notation (Wilkinson and Rogers, 1973):

Eq. 1:

$$Y = \beta 0 + \beta 1\, Flash\ location\ * \beta 2\, Flash\ order\ + \beta 3\, Baseline\ frequency\ + \ (1\,|\,Participant)$$

321   To examine whether rebound dynamics were explained by prior inhibition, we tested

322   *Rebound frequency* as dependent variable with a second model, extending the model in

323   Eq.1 by adding *Inhibition frequency* as a continuous predictor (Eq. 2):

324   Eq. 2:

$$Y = \beta 0 + \beta 1\,Flash\,location\,*\beta 2\,Flash\,order\,+\,\beta 3\,Baseline\,frequency\,+\,\beta 4\,Inhibition\,frequency$$
$$+\,(1\,|\,Participant)$$

In Experiment 2, we compared *Rebound frequency* between the first and fifth flash

using a paired-samples t-test. Additionally, we compared *Rebound frequency* for the

fifth flash between Experiment 1 and Experiment 2 using an independent-samples t-

test.

Results

Saccadic inhibition remains consistent across saccades of varying amplitudes,

directions, and velocities.

We first characterized qualitatively eye movement behavior by analyzing standard

saccade parameters across all trials (Figure 2). In Experiment 1, a total of 31,559

saccades were retained after manual screening to exclude detection errors (see Data

preprocessing). As expected in free-viewing tasks (Foulsham et al., 2008; Foulsham

and Kingstone, 2010), eye movements were predominantly horizontal (Figure 2A).

Saccade amplitudes had a mode at approximately 2.5 degrees of visual angle (Figure

2B), consistent with serial object inspection. The main sequence (Figure 2C) exhibited

the typical nonlinear relationship between saccade amplitude and peak velocity (Bahill

et al., 1975; Buonocore et al., 2017b).

Figure 2. Global oculomotor behavior during free viewing (Experiment 1).

(A) The distribution of saccade directions shows a predominant horizontal bias in visual exploration, as

commonly observed in free-viewing tasks. (B) The distribution of saccade amplitudes is highly positively

skewed, with a mode at ~2.5 degrees of visual angle, consistent with serial object inspection strategies

that favor smaller saccades. (C) From the main sequence, it is possible to appreciate how peak velocity

scales nonlinearly with amplitude, reflecting the canonical relationship. Note that the main sequence is

plot in log-log coordinates.

Temporal characteristics, including saccade and fixation durations (Figure 3A), were

also in line with established profiles for unconstrained visual exploration. Critically,

despite substantial variability in saccade properties highlighted in Figure 2, a consistent

pattern of saccadic inhibition emerged when saccades were time-locked to flash onset.

A robust suppression of saccade frequency was observed approximately 120 ms post-

flash across all participants (Figure 3B), confirming the reliability of SI under naturalistic

viewing conditions. These results support the notion that saccadic inhibition is a stable

and generalizable feature of oculomotor control, largely invariant to fluctuations in

saccade amplitude, direction, or velocity (Reingold and Stampe, 2000, 2004).

Figure 3. Distributions of saccade and fixation durations.

A) Saccade and fixation durations fell within the expected range for unconstrained exploration. (B) When

time-locked to flash onset (magenta vertical bar), saccades showed a robust suppression at ~120 ms,

confirming reliable saccadic inhibition during free viewing. Each dot represents a single saccade. The

dataset includes 31,559 saccades after manual screening, with colors indicating individual subjects.

Foveal flashes elicit stronger inhibition than parafoveal ones

We next investigated how the spatial location of visual transients modulates the

temporal dynamics of saccadic inhibition and rebound. Specifically, we compared the

effects of foveal and parafoveal flashes by analyzing their associated inhibition profiles

(see *Experimental Design and Statistical Analysis*). Both flash types induced a

pronounced inhibition peaking around 120 ms after onset (Figure 4A and B), with gray

lines indicating consistent effects across individual participants. At visual inspection,

foveal flashes (blue line, Figure 4A) produced stronger and more temporally precise

inhibition compared to parafoveal flashes (red line, Figure 4B, see statistics below). This

was followed by a sharper and more pronounced rebound in saccade frequency,

mirroring the inhibition phase and again favoring foveal stimulation.

Figure 4. Saccadic inhibition profiles in foveal and parafoveal flashes (Experiment 1)

Saccadic inhibition profiles for foveal (panel A, blue) and parafoveal (panel B, red) flashes showed robust

reduction in saccade generation ~120 ms after onset. Foveal flashes produced stronger and more

temporally precise suppression, followed by a sharper rebound in saccade rate ~100 ms later. Gray lines

represent individual participants, demonstrating consistent effects across subjects. Shaded blue and red

rectangles indicate the time windows used to compute group averages for baseline, inhibition, rebound,

and post-rebound measures (see *Experimental Design and Statistical Analysis* and Figure 2).

In Figure 5A and B, we further partition inhibition profiles across five successive flashes

(dark to brown lines indicating the first to fifth flash). Notably, the magnitude and latency

of saccadic inhibition remained remarkably stable across repetitions, regardless of flash

location. This suggests that while spatial position modulates the strength of the

inhibitory response, the temporal dynamics of the mechanism are preserved across

repeated stimulation. In contrast, a clear pattern emerged in the rebound: later flashes

increasingly hindered the generation of new saccades, weakening the rebound phase of

the saccadic inhibition profile. In addition, baseline levels appeared to be affected by

flash repetition, with later saccades starting from a reduced frequency. In the next

sections, we provide statistical modeling of the saccadic inhibition parameters described

in the *Experimental design and statistical analysis* section.

Figure 5. Saccadic inhibition profiles and parameters across flash order (Experiment 1).

Saccadic inhibition profiles across flash order (1–5) showed a robust reduction in saccade generation ~120 ms after onset. (A) Foveal flashes elicited stronger and more temporally precise suppression. The rebound in saccade rate occurred ~100 ms later, being clearly marked for the first flash but progressively less pronounced for subsequent flashes. (B) Parafoveal flashes followed the same pattern, though with an overall weaker rebound. Shaded rectangles indicate the time windows used to compute group averages for baseline, inhibition, rebound, and post-rebound measures (see *Experimental Design and Statistical Analysis* and Figure 2). (C) Mean saccade frequency (Hz) during the inhibition window and (D) during the rebound window, plotted across five successive flashes (flash order 1–5). Data are shown

separately for foveal (blue) and parafoveal (orange) flash locations. While inhibition frequency remains

stable across flashes, rebound frequency tend to diminish.

Statistical modeling of inhibition frequency. Inhibition frequency is not modulated by

flash repetition

We statistically tested these observations using linear mixed-effects models (LME) on

parameters extracted within predefined time windows (see shaded rectangles in Figures

3, 4, 5, and in the *Experimental Design and Statistical Analysis* section for a full

description of dependent measures). To quantify how the strength of saccadic inhibition

varied across conditions, we modeled *Inhibition frequency*—defined as the mean

saccade rate during the inhibitory dip—as a function of *Baseline frequency*, *Flash*

location, *Flash order*, and their interaction, with random intercepts for each participant to

account for repeated measures (Eq. 1). The model intercept was 1.36 Hz (p < 0.001),

representing the average saccade frequency during inhibition under the reference

condition (i.e., first flash, central location, and average baseline rate). This value reflects

a typical reduction of ~1.5 Hz compared to the pre-flash baseline (~3 Hz, Figure 5),

indicating robust oculomotor inhibition. A significant main effect of *Flash location* (β =

0.63, p < 0.001) revealed that inhibition was weaker when the flash occurred in the

parafoveal visual field: saccade frequency during the inhibition period was higher at

parafoveal compared to central locations (Figure 5A and B, blue vs. red lines and Figure

5C and D, all red dots above blue dots for all Flash orders). This supports the presence

of spatial specificity in the strength of inhibition, consistent with known differences in

foveal magnification across cortical and subcortical circuits (Cynader and Berman,

1972; Robinson, 1972; Benson et al., 2012; Hafed and Chen, 2016; Kupers et al., 2022;

Fracasso et al., 2023).

Although baseline saccade frequency showed a declining trend across flash

repetitions (Figure 5)—possibly reflecting fatigue or habituation — it did not significantly

modulate inhibition frequency (β = 0.03, p = .545). Similarly, neither the main effect of

Flash order (all p > 0.15) nor the Flash order by location interaction (all p > 0.07)

reached significance. These results suggest that neither repetition nor individual

baseline oculomotor state reliably influenced the depth of inhibition under these

conditions. Finally, random effects revealed modest between-subject variability (SD =

0.51 Hz), indicating that average inhibition strength was relatively stable across

participants. Overall, these findings support the interpretation that saccadic inhibition is

a robust, reflex-like phenomenon—largely invariant to both recent stimulation history

and individual differences in baseline saccade rates.

Statistical modeling of inhibition latency. Inhibition onset scales with the baseline level of

oculomotor activity

We next fit a linear mixed-effects model to predict Inhibition latency. The

intercept was 121.93 ms (p < 0.001), consistent with classical estimates of saccadic

inhibition timing under baseline conditions (Reingold and Stampe, 2002; Buonocore and

McIntosh, 2008; Edelman and Xu, 2009; Bompas and Sumner, 2011), but with a slight

delay, probably due to the free viewing nature of the task and the persistency of the

visual transient which lasted 80 ms, probably widening the inhibitory dip. A significant

negative effect of *Baseline frequency* (β = –5.23, p = 0.027) indicated that participants

with higher pre-flash saccade rates tended to exhibit earlier inhibition. This suggests

that the onset of inhibition scales with the initial level of oculomotor activity: participants

with more active oculomotor states may generate stronger or more efficient inhibitory

signals, reaching the suppression threshold faster. Conversely, lower baseline rates

may reflect a less excitable oculomotor system requiring more time to initiate inhibition.

A fixed-threshold mechanism, modulated by ongoing activity, could account for this

variability in latency (Dorris and Munoz, 1998). Neither *Flash order* (all p > 0.20) nor

Flash location (p = 0.80) yielded significant main effects, suggesting that these factors

alone did not influence inhibition timing. However, the *Flash order by location* interaction

for the fifth flash was significant (β = –23.89 ms, p = 0.048), pointing to a potential trend

whereby later flashes at parafoveal locations were associated with slightly earlier

inhibition.

Statistical modeling of inhibition frequency. Rebound responses decline with repetition.

Unlike inhibition, rebound responses demonstrated clear habituation over time.

Rebound frequency peaked approximately 250–300 ms after flash onset, followed by a

gradual decline in amplitude across successive flashes (Figure 5A, B, C and D). The

intercept was 3.49 Hz (p < 0.001), reflecting the average saccade frequency during the

rebound period under reference conditions. This value marks the typical post-inhibitory

overshoot in eye movement activity (Reingold and Stampe, 2002; Buonocore and McIntosh, 2008; Edelman and Xu, 2009; Bompas and Sumner, 2011). A significant main effect of Flash order was observed in particular for the third and fifth repetition (Flash order3: $\beta = -0.42$, p = 0.04; Flash order$_5$: $\beta = -0.68$, p = 0.0016), indicating a progressive reduction in rebound frequency over time. This decline suggests that the oculomotor system became less responsive across repeated flashes—possibly reflecting habituation. A significant main effect of *Flash location* ($\beta = -0.56$, p = 0.008) confirmed that rebound strength was location-dependent, with reduced reactivation at parafoveal sites. Additionally, higher baseline saccade frequency predicted stronger rebound responses ($\beta = 0.20$, p = 0.003), suggesting that participants with more active oculomotor baselines not only inhibited earlier (see section above) but also re-engaged motor activity more robustly after suppression.

Inhibition and rebound are dissociable processes.

To determine whether rebound responses were simply a delayed consequence of stronger inhibition, we included *Inhibition frequency* as a predictor in the rebound model (see Eq. 2). *Inhibition frequency* did not have any effect ($\beta = -0.15$, p = 0.098), suggesting no predictive value. Furthermore, partial correlations between inhibition and rebound—controlling for baseline saccade frequency—were near zero across all conditions. These results support the notion that inhibition and rebound are functionally dissociable, reflecting distinct neural mechanisms: inhibition appears reflexive and sensory-driven, while rebound may reflect adaptive motor reactivation, potentially modulated by top-down influences.

Crucially, *Inhibition frequency* did not significantly predict rebound strength,

indicating that these two components of the saccadic response—reflexive suppression

and subsequent reactivation—can vary independently. The absence of a systematic

relationship between inhibition depth and rebound amplitude argues against a simple

compensatory mechanism, and instead supports the view that these reflect distinct

processes with separable sensitivities. Overall, these findings indicate that while

rebound dynamics are sensitive to stimulus location, baseline activation, and stimulus

repetition, they are not determined by the preceding degree of inhibition, further

supporting their dissociation.

Rebound suppression shows experiment-specific modulation but no clear temporal

decay.

One possible explanation for the pattern of results observed in Experiment 1 is that the

oculomotor system becomes less reactive over time during the free-viewing task,

regardless of flash presentation. In this scenario, when probed, the sensory-driven

inhibition remains intact, but the motor reprogramming component would gradually

decrease over time. Crucially, this change would occur regardless on the number of

flashes presented and simply related to the time dedicated to the task (*time-on-task*).

This would alter the shape of the saccadic inhibition profile, yielding a stable inhibition

level, a progressively reduced rebound across flash order, and a decrease in baseline

frequency. To test this hypothesis, we conducted a second experiment, identical to the

Experiment 1, but where only the first or the fifth flash was presented. If the effect

observed in Experiment 1 reflects habituation and time-on-task, motor rebound levels

would be comparable between the first and fifth flash in Experiment 2. In contrast, if the

effect were entirely due to time-on-task, differences between the first and fifth flash

should still be observed under these conditions.

In Experiment 2, 31,559 saccades were retained after excluding detection errors.

As in Experiment 1, saccade metrics conformed to known patterns of free viewing,

including predominantly horizontal directions, a peak amplitude around 2.5 degrees of

visual angle, a typical main sequence, and canonical temporal dynamics. Crucially, a

reliable suppression of saccade frequency was observed around 130 ms after flash

onset across participants, confirming the robustness of saccadic inhibition under

naturalistic conditions.

To assess whether the rebound response varies with flash position, we

compared *Rebound frequency* for the fifth flash between Experiments 1 and 2 (Figure

6). *Rebound frequency* differed significantly between experiments for the fifth flash ($t = -$

2.35, $df = 38$, $p = 0.02$). Moreover, while a clear reduction from the first to the fifth flash

was observed in Experiment 1, no such difference was found in Experiment 2 ($t = 1.5$, df

$= 18$, $p = 0.15$). This pattern is inconsistent with an explanation based on time-on-task

effects, which would predict within-experiment differences even when only one flash

was shown. Instead, the findings support a habituation account, specific to repeated

flash stimulation, whereby repeated flash exposure progressively weakens the rebound

phase of saccadic inhibition.

Figure 6. Saccadic inhibition profiles for first and fifth flashes in Experiment 1 and Experiment 2.

(A) Experiment 1: SI profiles for the first (dark line) and fifth (light line) flash within a trial sequence.

Inhibition depth remained stable, but rebound frequency was significantly reduced for the fifth flash. (B)

Experiment 2: SI profiles when only the first or the fifth flash was presented per trial. No significant

difference emerged between conditions, indicating that rebound suppression in Experiment 1 reflects

repeated exposure rather than the temporal position of the flash. Shaded rectangles indicate analysis

windows for baseline, inhibition, rebound, and post-rebound measures. In panel B, the thin lines refer to

the "invisible" flashes that did not elicit any inhibition. As expected, when no stimuli were presented the

baseline level remained constant across the entire time window.

Discussion

The present study investigated how saccadic inhibition evolves across repeated visual

stimulation during naturalistic free viewing. Across two experiments, we found that the

initial inhibition phase following a visual transient remained stable across repeated

flashes, whereas the subsequent rebound—characterized by a saccade rate increase above baseline—showed progressive attenuation (Experiment 1). Simple time-on-task effects (Shan and Edelman, 2023) could not explain the observed pattern: both inhibition and rebound magnitudes were comparable between the first and last transient when they were presented in isolation (Experiment 2). On the other hand, the rebound generated by the fifth flash in isolation was stronger than that generated by the fifth repeated flash (Experiment 1 vs. Experiment 2). Thus, we suggest that the decline in rebound in Experiment 1 reflects habituation induced by repeated sensory transients.

Sensory Inhibition and Motor rebound reflect distinct processes

The decoupling of inhibition and rebound is evidence that distinct mechanisms underlie the two phases of the oculomotor response to sudden visual changes. Specifically, the initial inhibition reflects a resilient, reflex-like suppression process (Reingold and Stampe, 2002; Buonocore and McIntosh, 2008; Edelman and Xu, 2009; Bompas and Sumner, 2011), whereas the rebound reflects an adaptive, experience-dependent modulation. Altogether, our findings demonstrate a functional dissociation between inhibition and rebound in oculomotor behavior during continuous free viewing.

Our results extend previous work on SI adaptation (Shan and Edelman, 2023) by showing that the inhibitory phase is subjected to adaptation only within a certain time window. Unlike the rapid attenuation of the inhibition reported at short ISIs (~0.2 s), the absence of SI decline at longer flash intervals (~0.7–1.2 s) suggests that low-level sensory adaptation recovers rapidly over time, in the order or a few hundreds of milliseconds (Wark et al., 2007). This mechanism may be particularly relevant under

free-viewing conditions, so that inhibition remains stable to support the reinstatement of

saccades, which are often redirected toward alternative new visual targets (Buonocore

et al., 2017a).

Habituation in the motor system.

This pattern of stable inhibition contrasted with weakening motor output resembles

habituation in other motor systems, where early sensory gating remains preserved while

subsequent motor responses decline. For example, in the vestibulo-ocular reflex

(VOR)—which stabilizes vision by producing eye movements opposite to head

movements—the basic sensory pathway that detects head motion remains intact.

However, when the system is exposed to repeated or prolonged stimulation, the

strength of the motor response gradually decreases. This reduction in "motor gain"

means that the eyes no longer fully counteract the head movement, resulting in

progressively smaller compensatory eye movements (Lisberger et al., 1994). In another

sensory domain, the acoustic startle reflex—the rapid whole-body flinch triggered by

sudden loud sounds—illustrates a similar pattern (Yeomans and Frankland, 1995). The

initial inhibitory process that suppresses background activity remains consistently

engaged, ensuring that the sound is detected. However, when the stimulus is repeated,

the strength of the motor response (the actual body movement) gradually diminishes. In

other words, the system continues to register the sound but produces a smaller and

smaller behavioral reaction over time (Davis, 1984; Koch, 1999; Zheng and Schmid,

2023).

At the neuronal level, these behavioral effects correspond to distinct sensory and

motor dynamics. In the superior colliculus (SC), responses to repeated visual transients

decrease despite stable sensory detection (Boehnke et al., 2011). In brainstem circuits

underlying the startle reflex, neurons exhibit early inhibition of auditory input alongside

diminished motor output with repetition (Davis, 1984; Geyer and Braff, 1987). Across

these systems, inhibitory gating mechanisms remain intact to preserve responsiveness,

whereas motor re-engagement is selectively attenuated. By analogy, the dissociation

we observed between stable inhibition and weakening rebound in SI likely reflects the

same principle: robust sensory suppression coupled with an adaptive, experience-

dependent reduction in motor reprogramming under repeated stimulation.

Neural mechanisms.

While the current manuscript centers around behavioral (oculomotor) data, we can

speculate as to what might be the most likely neural mechanism mediating the observed

results. The persistence of SI across repetitions is consistent with existing models

proposing subcortical circuits gating saccade execution. Within the oculomotor system,

omnipause neurons (OPNs) in the pontine reticular formation exert tonic inhibition over

saccade-generating burst neurons (Fuchs and Luschei, 1972; Luschei and Fuchs, 1972;

Keller, 1974; Büttner-Ennever and Büttner, 1978; Evinger et al., 1982; Büttner-Ennever

et al., 1988; Büttner-Ennever et al., 1999; Takahashi et al., 2022), maintain tonic firing

during fixation and pausing just prior to saccade initiation. Recently, we have suggested

that OPNs may receive direct sensory input, allowing rapid responses to exogenous

stimuli and raising the threshold for saccade initiation (Buonocore and Hafed, 2023).

This would explain the rapid onset of SI following a new stimulus. Increases in OPN

activity following visual flashes might reflect a hard-wired gating mechanism that

operates independently of behavioral relevance. Supporting this, the timing and

magnitude of SI in our data remained invariant even under repeated stimulation,

consistent with evidence that either direct OPNs reactivation or visual bursts in the SC

via projections to OPNs can rapidly trigger suppression of saccade generation. In

contrast, the rebound may rely on mechanisms disengaging OPN activity and re-

exciting burst neurons. Repeated stimulation may reduce the efficiency of this

disengagement, leading to weaker reactivation of saccade commands. Such selective

habituation echoes findings in the SC (Boehnke et al., 2011; Basso and May, 2017) and

frontal eye fields (FEF) (Bruce and Goldberg, 1985; Mayo and Sommer, 2008), where

responses to repeated transients diminish despite preserved detection.

These findings imply a recalibration of the balance between inhibitory stability

and reorienting flexibility within brainstem circuits. Our work provides a systems-level

perspective on how sensorimotor loops evolve during active vision. SI thus emerges

from dynamic interactions between SC, FEF, and OPNs. While SC and FEF provide

excitatory signals for orienting, OPNs contribute an inhibitory signal that prevents

premature eye movements. By ensuring that saccades are executed only after sufficient

sensory processing, this interaction adapts visual behavior to environmental demands.

Variations in attentional state, relevance, or motivational salience may gradually

disengage the system from reflexive orienting responses .

Interestingly, the temporal structure of the rebound included dynamics consistent

with rhythmicity in oculomotor control (see Figure 3A, B and Figure 4A, B). Recurrent

fluctuations in saccade likelihood following inhibition may be shaped by beta- or alpha-band oscillations in frontoparietal areas, which have been linked to saccadic timing and attentional cycles (VanRullen, 2016; Gaillard et al., 2020). This rhythmic post-inhibitory signature may support a framework in which the brain evaluates salience in periodic windows, reconfiguring sensorimotor coupling accordingly (Helfrich et al., 2018; Fiebelkorn and Kastner, 2019).

Conclusion

Together, these findings reveal a selective habituation of motor re-engagement in SI, whereas sensory-driven inhibition remains robust. This dissociation shows that the oculomotor system recalibrates to repeated stimulation by preserving inhibitory fidelity while attenuating rebound, thereby filtering out irrelevant events. More broadly, SI offers a window into how sensory and motor components of eye-movement control adapt independently, with implications for both basic mechanisms and clinical applications. Functionally, a system that maintains transient suppression yet reduces redundant reprogramming may balance sensory prioritization with motor efficiency. In cluttered or predictable environments, minimizing rebound to repeated flashes could optimize exploration by limiting distraction and conserving resources.

This separation also holds clinical relevance. Disorders such as Parkinson's disease and ADHD are characterized by deficits in oculomotor inhibition and variability in response timing (Antoniades and Spering, 2024). In Parkinson's disease (PD), other than the emerging of fatigue in movement control (Di Vico et al., 2021) also oculomotor abnormalities emerge early in the disease course. Patients typically exhibit saccadic

hypometria, prolonged latencies, and reduced peak velocities (Pretegiani and Optican, 2017). Performance on tasks probing reflexive oculomotor control—such as visually guided prosaccades—is often relatively preserved, whereas antisaccade tasks consistently reveal impaired voluntary inhibition and strategy reconfiguration, with increased error rates and longer latencies, including at early drug-naïve stages (Antoniades et al., 2015). Against this backdrop, our results point to a clinically relevant dissociation that is translatable to clinical implementation for PD assessment: the SI dip (transient, stimulus-driven suppression of saccades) appears preserved, whereas the motor rebound is probably delayed and/or reduced in magnitude and time relative to controls—consistent with a framework in which sensory-driven inhibitory gating remains intact while motor re-engagement is selectively compromised (Fooken et al., 2022; Buonocore and Hafed, 2023). This pattern aligns with evidence that some oculomotor behaviors relying on superior colliculus–brainstem pathways are spared in PD, whereas behaviors requiring greater fronto–basal ganglia engagement are disproportionately affected. The ability to quantify the stability of SI and the adaptability of rebound may yield sensitive markers for early detection of dysfunctions affecting automatic and voluntary control systems.

Future work should investigate neural signatures of this dissociation using EEG or fMRI and assess whether rebound adaptation generalizes across modalities and tasks. By exploring the interaction between SI and sensory adaptation, we aim to uncover the principles that regulate visual exploration. This knowledge could inform both theoretical models of oculomotor control and translational applications in perception and clinical neuroscience.

References

Anstis S, Verstraten FA, Mather G (1998) The motion aftereffect. Trends Cogn Sci 2:111-117.

Antoniades CA, Spering M (2024) Eye movements in Parkinson's disease: from neurophysiological mechanisms to diagnostic tools. Trends Neurosci 47:71-83.

Antoniades CA, Demeyere N, Kennard C, Humphreys GW, Hu MT (2015) Antisaccades and executive dysfunction in early drug-naive Parkinson's disease: The discovery study. Mov Disord 30:843-847.

Bahill AT, Clark MR, Stark L (1975) The main sequence, a tool for studying human eye movements. Mathematical Biosciences 24:191-204.

Basso MA, May PJ (2017) Circuits for Action and Cognition: A View from the Superior Colliculus. Annual Review of Vision Science 3:197-226.

Benson V, Ietswaart M, Milner D (2012) Eye Movements and Verbal Report in a Single Case of Visual Neglect. PLoS ONE 7:e43743.

Benucci A, Saleem AB, Carandini M (2013) Adaptation maintains population homeostasis in primary visual cortex. Nat Neurosci 16:724-729.

Blakemore C, Campbell FW (1969) Adaptation to spatial stimuli. J Physiol 200:11p-13p.

Boehnke SE, Berg DJ, Marino RA, Baldi PF, Itti L, Munoz DP (2011) Visual adaptation and novelty responses in the superior colliculus. Eur J Neurosci 34:766-779.

Bompas A, Sumner P (2011) Saccadic inhibition reveals the timing of automatic and voluntary signals in the human brain. J Neurosci 31:12501-12512.

Brainard DH (1997) The Psychophysics Toolbox. Spatial Vision 10:433-436.

Bruce CJ, Goldberg ME (1985) Primate frontal eye fields. I. Single neurons discharging

before saccades. J Neurophysiol 53:603-635.

Buonocore A, McIntosh RD (2008) Saccadic inhibition underlies the remote distractor

effect. Experimental brain research Experimentelle Hirnforschung

Expérimentation cérébrale 191:117-122.

Buonocore A, McIntosh RD (2012) Modulation of saccadic inhibition by distractor size

and location. Vision Research 69:32-41.

Buonocore A, McIntosh RD (2013) Attention modulates saccadic inhibition magnitude.

Quarterly journal of experimental psychology (2006) 66:1051-1059.

Buonocore A, Hafed ZM (2023) The inevitability of visual interruption. J Neurophysiol

130:225-237.

Buonocore A, Purokayastha S, McIntosh RD (2017a) Saccade Reorienting Is Facilitated

by Pausing the Oculomotor Program. Journal of cognitive neuroscience 29:2068-

2080.

Buonocore A, Skinner J, Hafed ZM (2019) Eye Position Error Influence over "Open-

Loop" Smooth Pursuit Initiation. The Journal of Neuroscience 39:2709-2721.

Buonocore A, Chen C-Y, Tian X, Idrees S, Münch TA, Hafed ZM (2017b) Alteration of

the microsaccadic velocity-amplitude main sequence relationship after visual

transients: implications for models of saccade control. Journal of

Neurophysiology 117:1894-1910.

Burleson B, Turatto M, Plomp G, Pascucci D (2025) Anticipatory and reactive

mechanisms of habituation to visual distractors. Sci Rep 15:22953.

Büttner-Ennever JA, Büttner U (1978) A cell group associated with vertical eye

751    movements in the rostral mesencephalic reticular formation of the monkey. Brain

752    Research 151:31-47.

753 Büttner-Ennever JA, Cohen B, Pause M, Fries W (1988) Raphe nucleus of the pons

754    containing omnipause neurons of the oculomotor system in the monkey, and its

755    homologue in man. Journal of Comparative Neurology 267:307-321.

756 Büttner-Ennever JA, Horn AKE, Henn V, Cohen B (1999) Projections from the superior

757    colliculus motor map to omnipause neurons in monkey. Journal of Comparative

758    Neurology 413:55-67.

759 Cynader M, Berman N (1972) Receptive-field organization of monkey superior

760    colliculus. J Neurophysiol 35:187-201.

761 Davis M (1984) The Mammalian Startle Response. In: Neural Mechanisms of Startle

762    Behavior (Eaton RC, ed), pp 287-351. Boston, MA: Springer US.

763 Di Vico IA, Cirillo G, Tessitore A, Siciliano M, Venturelli M, Falup-Pecurariu C, Tedeschi

764    G, Morgante F, Tinazzi M (2021) Fatigue in hypokinetic, hyperkinetic, and

765    functional movement disorders. Parkinsonism Relat Disord 86:114-123.

766 Dorris MC, Munoz DP (1998) Saccadic probability influences motor preparation signals

767    and time to saccadic initiation. J Neurosci 18:7015-7026.

768 Edelman JA, Xu KZ (2009) Inhibition of Voluntary Saccadic Eye Movement Commands

769    by Abrupt Visual Onsets. Journal of Neurophysiology 101:1222-1234.

770 Evinger C, Kaneko CR, Fuchs AF (1982) Activity of omnipause neurons in alert cats

771    during saccadic eye movements and visual stimuli. J Neurophysiol 47:827-844.

Fiebelkorn IC, Kastner S (2019) A Rhythmic Theory of Attention. Trends Cogn Sci

23:87-101.

Fooken J, Patel P, Jones CB, McKeown MJ, Spering M (2022) Preservation of Eye

Movements in Parkinson's Disease Is Stimulus- and Task-Specific. J Neurosci

42:487-499.

Foulsham T, Kingstone A (2010) Asymmetries in the direction of saccades during

perception of scenes and fractals: effects of image type and image features.

Vision Res 50:779-795.

Foulsham T, Kingstone A, Underwood G (2008) Turning the world around: patterns in

saccade direction vary with picture orientation. Vision Res 48:1777-1790.

Fracasso A, Buonocore A, Hafed ZM (2023) Peri-Saccadic Orientation Identification

Performance and Visual Neural Sensitivity Are Higher in the Upper Visual Field. J

Neurosci 43:6884-6897.

Fuchs AF, Luschei ES (1972) Unit activity in the brainstem related to eye movement.

Possible inputs to the motor nuclei. Bibl Ophthalmol 82:17-27.

Fuster JM (2004) Upper processing stages of the perception-action cycle. Trends Cogn

Sci 8:143-145.

Gaillard C, Ben Hadj Hassen S, Di Bello F, Bihan-Poudec Y, VanRullen R, Ben Hamed

S (2020) Prefrontal attentional saccades explore space rhythmically. Nat

Commun 11:925.

Georgeson MA, Harris MG (1984) Spatial selectivity of contrast adaptation: models and

data. Vision Res 24:729-741.

Geyer MA, Braff DL (1987) Startle habituation and sensorimotor gating in schizophrenia

and related animal models. Schizophr Bull 13:643-668.

Greenlee MW, Heitger F (1988) The functional role of contrast adaptation. Vision Res

28:791-797.

Hafed ZM, Chen CY (2016) Sharper, Stronger, Faster Upper Visual Field

Representation in Primate Superior Colliculus. Curr Biol 26:1647-1658.

Hayhoe MM (2017) Vision and Action. Annu Rev Vis Sci 3:389-413.

Helfrich RF, Fiebelkorn IC, Szczepanski SM, Lin JJ, Parvizi J, Knight RT, Kastner S

(2018) Neural Mechanisms of Sustained Attention Are Rhythmic. Neuron 99:854-

865.e855.

Keller EL (1974) Participation of Medial Pontine Reticular-Formation in Eye-Movement

Generation in Monkey. Journal of Neurophysiology 37:316-332.

Kerzel D, Born S, Souto D (2010) Inhibition of Steady-State Smooth Pursuit and Catch-

Up Saccades by Abrupt Visual and Auditory Onsets. Journal of Neurophysiology

104:2573-2585.

Koch M (1999) The neurobiology of startle. Prog Neurobiol 59:107-128.

Kohn A (2007) Visual adaptation: physiology, mechanisms, and functional benefits. J

Neurophysiol 97:3155-3164.

Kupers ER, Benson NC, Carrasco M, Winawer J (2022) Asymmetries around the visual

field: From retina to cortex to behavior. PLoS Comput Biol 18:e1009771.

Lisberger SG, Pavelko TA, Broussard DM (1994) Neural basis for motor learning in the

vestibuloocular reflex of primates. I. Changes in the responses of brain stem

neurons. J Neurophysiol 72:928-953.

Luschei ES, Fuchs AF (1972) Activity of brain stem neurons during eye movements of

818   alert monkeys. J Neurophysiol 35:445-461.

819   Malevich T, Buonocore A, Hafed ZM (2020) Rapid stimulus-driven modulation of slow

820   ocular position drifts. eLife 9:e57595.

821   Mayo JP, Sommer MA (2008) Neuronal Adaptation Caused by Sequential Visual

822   Stimulation in the Frontal Eye Field. Journal of Neurophysiology 100:1923-1935.

823   Pascucci D, Mancuso G, Santandrea E, Della Libera C, Plomp G, Chelazzi L (2019)

824   Laws of concatenated perception: Vision goes for novelty, decisions for

825   perseverance. PLoS Biol 17:e3000144.

826   Pelli DG (1997) The VideoToolbox software for visual psychophysics: transforming

827   numbers into movies. Spatial Vision 10:437-442.

828   Pretegiani E, Optican LM (2017) Eye Movements in Parkinson's Disease and Inherited

829   Parkinsonian Syndromes. Front Neurol 8:592.

830   Reingold EM, Stampe DM (1999) Saccadic Inhibition in Complex Visual Tasks. In, pp

831   249-255. Boston, MA: Springer US.

832   Reingold EM, Stampe DM (2000) Saccadic Inhibition and Gaze Contingent Research

833   Paradigms. In, pp 119-145: Elsevier.

834   Reingold EM, Stampe DM (2002) Saccadic Inhibition in Voluntary and Reflexive

835   Saccades. Journal of cognitive neuroscience 14:371-388.

836   Reingold EM, Stampe DM (2004) Saccadic Inhibition in Reading. Journal of

837   Experimental Psychology: Human Perception and Performance 30:194-211.

838   Robinson DA (1972) Eye movements evoked by collicular stimulation in the alert

839   monkey. Vision Res 12:1795-1808.

Rolfs M, Kliegl R, Engbert R (2008) Toward a model of microsaccade generation: The

case of microsaccadic inhibition. Journal of Vision 8:5-5.

Sekuler RW, Ganz L (1963) Aftereffect of seen motion with a stabilized retinal image.

Science 139:419-420.

Shan Y, Edelman JA (2023) The reduction of saccadic inhibition by distractor repetition.

J Neurophysiol 130:619-627.

Smirnakis SM, Berry MJ, Warland DK, Bialek W, Meister M (1997) Adaptation of retinal

processing to image contrast and spatial scale. Nature 386:69-73.

Solomon SG, Kohn A (2014) Moving sensory adaptation beyond suppressive effects in

single neurons. Curr Biol 24:R1012-1022.

Takahashi M, Sugiuchi Y, Na J, Shinoda Y (2022) Brainstem Circuits Triggering

Saccades and Fixation. J Neurosci 42:789-803.

Taylor R, Buonocore A, Fracasso A (2024) Saccadic "inhibition" unveils the late

influence of image content on oculomotor programming. Exp Brain Res

242:2281-2294.

Torrado Pacheco A, Tilden EI, Grutzner SM, Lane BJ, Wu Y, Hengen KB, Gjorgjieva J,

Turrigiano GG (2019) Rapid and active stabilization of visual cortical firing rates

across light-dark transitions. Proc Natl Acad Sci U S A 116:18068-18077.

VanRullen R (2016) Perceptual Cycles. Trends Cogn Sci 20:723-735.

Wark B, Lundstrom BN, Fairhall A (2007) Sensory adaptation. Curr Opin Neurobiol

17:423-429.

Webster MA (2011) Adaptation and visual coding. J Vis 11.

Webster MA, MacLeod DI (2011) Visual adaptation and face perception. Philos Trans R

Soc Lond B Biol Sci 366:1702-1725.

Wilkinson GN, Rogers CE (1973) Symbolic Description of Factorial Models for Analysis

of Variance. Journal of the Royal Statistical Society Series C (Applied Statistics)

22:392-399.

Yeomans JS, Frankland PW (1995) The acoustic startle reflex: neurons and

connections. Brain Res Brain Res Rev 21:301-314.

Zheng A, Schmid S (2023) A review of the neural basis underlying the acoustic startle

response with a focus on recent developments in mammals. Neurosci Biobehav

Rev 148:105129.