TITLE: Predator induced phenotypes are inherited over four generations in genetically identical

Daphnia lumholtzi

RUNNING TITLE: Transgenerational Inheritance in *Daphnia lumholtzi*

Authors:

Shannon N. Snyder[1,*],

Walker C. Meyer[1],

Shanie L. Jorgenson[1]

Ethan B. Contreras[1]

Tea Bland[1]

William A. Cresko[1,2,*]

Affiliations:

[1] Institute of Ecology & Evolution, University of Oregon, Eugene, OR, 97403, USA

[2] Department of Bioengineering, Knight Campus for Accelerating Scientific Impact, Eugene,

OR, 97403, USA

*Corresponding Authors (ssnyder3@uoregon.edu & wcresko@uoregon.edu)

ORCID:

SNS: 0000-0001-7049-2950

WCM: 0009-0003-7225-4296

SLJ: 0009-0007-0144-2504

EBC: 0009-0001-1019-8263

TB: 0009-0007-1456-0569

WAC: 0000-0002-3496-8074

Keywords: Transgenerational inheritance, Phenotypic plasticity, Non-genetic inheritance,

Epigenetic Inheritance, Armor Morphology, Bayesian hierarchical modeling, Inducible defenses

Data availability statement: The raw data and full analysis scripts that support the findings of

this study are openly available in Markdown on GitHub

(https://github.com/shannonsnyder/daphnia-transgenerational-plasticity).

Author contributions: **Conceptualization and Methodology:** SNS and WAC conceived and

designed the study. **Investigation:** SNS, WCM, and SLJ reared the animals and collected the

imaging data. **Data Collection and Curation:** SNS, EBC and TB curated the morphological

measurements. **Formal Analysis:** SNS and WAC analyzed the data. **Visualization:** SNS created

the data visualizations. **Writing – Original Draft:** SNS wrote the manuscript. **Writing –**

Review & Editing: WAC provided revisions. All authors approved the final manuscript.

Funding: This work was funded by the National Science Foundation Grant OPP-2015301 (to

WAC), University of Oregon Research Excellence funds (WAC), University of Oregon Office of

the Vice President for Research and Innovation (OVPRI) seed funding (SNS and WAC). WCM

received funding and mentorship from the Knight Campus Undergraduate Scholars program at

the University of Oregon. SLJ and EC received funding from the University of Oregon Graduate

Students in Ecology and Evolution Student Organization. SLJ received funding from the McNair

Scholar Program. EC and WCM received funding from University of Oregon OVRPI.

Conflict of Interest: The authors declare no conflict of interest.

Acknowledgements: We would like to thank M. Lynch for his contribution of the *Daphnia*

lumholtzi (SAG) clone utilized for this study. We would also like to thank Mark Currey for his

assistance with stickleback husbandry and the Cresko lab for valuable feedback. Finally, we

thank Emily Willams for collection and maintenance of *Daphnia* clones prior to their arrival in

the Cresko lab.

57
58

Teaser Text

Can the ghosts of environments past shape organismal phenotypes of future generations? In an

experiment with clonal *Daphnia lumholtzi*, we reveal that predator-induced changes in body

shape can persist across several generations, even when the predator signal is long gone.

Exposing only the first generation to predator cues, we tracked morphological shifts through five

generations of genetically identical individuals. We found that these induced plastic phenotypes

persisted to the F4 generation before fading in F5. This discovery supports the importance of

non-genetic inheritance and underscores the powerful interplay between ancestral and current

environments in shaping organismal form and function.

Abstract

69   Environmental variation can induce phenotypic changes through adaptation of

populations and acclimation of individuals. Whereas adaptation can lead to persistent changes

within a population via allele frequency shifts, induced plastic phenotypes have been thought to

be ephemeral and lasting only through the affected generation. A growing number of studies

have shown that induced phenotypes may be inherited, but it is unclear how persistent the

inheritance of induced plastic phenotypes is across organisms. To directly address these

questions, we measured the phenotypic effects of exposure to a predator cue in the water flea,

Daphnia lumholtzi, which reproduces clonally allowing the isolation of the effects of direct

environmental exposure from allelic variation. We measured the phenotypic effects of exposure

to fish conditioned media in the F0, F3, F4, and F5 generations of genetically identical

generations following cue exposure only in the F0 generation. We found evidence of predation

induced morphologies into the F4 generation, with phenotypes receding in the F5 generation,

four generations after cue removal. These results highlight the importance of the interaction

between current and ancestral environments in determining phenotypic variation and highlights

the importance in future studies of characterizing the molecular mechanisms of within and

transgenerational phenotypic plasticity.

Introduction

Organisms respond to environmental variation through changes in morphology, behavior, and life history traits (Donelson et al., 2018; Szűcs et al., 2019; West-Eberhard, 2005) . These phenotypic responses may arise from shifts in allele frequencies across generations or from non-genetic mechanisms acting directly on individuals, such as developmental noise (Levis & Pfennig, 2019; Uller et al., 2018; West-Eberhard, 2020). Increased phenotypic variation can enhance the probability that some offspring will express trait values suited to novel environmental conditions (Botero et al., 2015; Simons, 2009; Tufto, 2015). However, this variation also increases the likelihood of mismatched phenotypes, reducing individual fitness for those with the maladaptive traits. While genetic adaptation can improve population-level fitness over time, subsequent environmental shifts may render previously adaptive traits maladaptive (Lande, 2009; Radchuk et al., 2019).

Regularly experienced environmental variation presents a particular challenge for evolving populations of organisms because it can lead to persistent mismatch of organisms and the environment, and thus ongoing genetic load (Ghalambor et al., 2007; Via et al., 1995). An alternative biological strategy for dealing with such variable conditions is the evolution of plastic traits (Hendry, 2016) in which phenotypic expression covaries with environmental variation (Bradshaw, 1965; Schlichting & Levin, 1986). Phenotypic plasticity allows a single genotype to consistently produce different phenotypes depending on the environment, without requiring genetic change (Pfennig et al., 2010; Pigliucci et al., 2006). Plasticity includes developmental plasticity (Nettle & Bateson, 2015; Uller et al., 2018), where early-life cues shape long-term traits, and acclimatory plasticity (Angilletta, 2009; Beaman et al., 2016), involving reversible

physiological changes in response to short-term fluctuations, illustrating a range of temporal

scales over which environmentally induced phenotypic changes can occur.

Plasticity within a generation has been well documented over numerous organisms and

environmental conditions (Agrawal et al., 1999a; Corl et al., 2018; Forsman, 2015). Although

plasticity itself is often an adaptive phenotype that has been crafted by natural selection, the trait

distributions produced by plasticity within one generation have historically been thought to be

ephemeral with little consequence for the phenotypes of future generations (West-Eberhard,

2020). However, modern molecular genetic and genomic analyses have documented that

variation in plastic traits is mediated by changes in gene expression (Hales et al., 2017) and are

hypothesized to involve epigenetic regulation (St-Cyr & McGowan, 2015). These molecular

mechanisms lead to the possibility that induced plastic phenotypes may be inherited across

generations (Harris et al., 2012a; Romero-Mujalli et al., 2024).

Transgenerational inheritance of induced plastic trait values is expected to evolve when

the parental environment has some covariance with the offspring environment (Agrawal et al.,

1999a). Although the inherited phenotypes should return to the parental state in the absence of

the inducing cues over generations, the rate at which this occurs is hypothesized to be determined

by the number of generations of which the environment shifts (Furrow & Feldman, 2014; Proulx

& Teotónio, 2017). In this way the evolution of inherited phenotypic plasticity provides a

mechanism for organisms and their offspring to respond adaptively to changing environments

without significant genetic load (Ghalambor et al., 2007). A small number of studies have

documented that induced phenotypes may be inherited (Hellmann et al., 2020; Tariel et al.,

2020), but it is unclear how common this is across organisms. Even less is known about how

long environmentally induced phenotypes persist across generations once the environmental cue

is removed (Norouzitallab et al., 2014).

The freshwater microcrustacean *Daphnia* is well suited for studying transgenerational

plasticity and addressing these questions (Harris et al., 2012b; Seda & Petrusek, 2011;

Yampolsky et al., 2014). The concept of plasticity was introduced in the early 1900s by Richard

Woltereck**,** who demonstrated that head size in *Daphnia* varied with nutrient availability, and

proposed that environmentally induced traits could be inherited (Woltereck, 1913). *Daphnia's*

parthenogenetic reproduction produces genetically identical offspring, eliminating variation due

to recombination, genetic background, and parental sex (Harris et al., 2012b). These

ecoresponsive crustaceans exhibit plastic responses to a range of environmental stressors,

including cyanobacteria (Dao et al., 2018; Gustafsson & Hansson, 2004; Schwarzenberger &

Martin-Creuzburg, 2021), temperature (Geerts et al., 2015; Müller et al., 2018), and predation,

affecting morphology (Horstmann et al., 2021), behavior (Beklioglu et al., 2006), and life history

traits (Adamczuk, 2020; Agrawal et al., 1999b). Classic examples include inducible helmets,

spines, and neckteeth in response to predator cues (Horstmann et al., 2021; Laforsch & Tollrian,

2004). Several of these traits persist across clonal generations, such as helmet length in *D.*

cucullata (Agrawal et al., 1999a) and clutch size in *D. ambigua* (Walsh, Cooley, et al., 2014),

suggesting a non-genetic mechanism of inheritance.

The number of generations over which induced traits persist in *Daphnia* in the absence of

the original environmental cue remains poorly resolved. Most studies track only one or two

generations post-exposure, such as helmet length in *D. cucullata* which disappears by the second

generation (F2) (Agrawal et al., 1999a; Laforsch & Tollrian, 2004), and clutch size in *D.*

ambigua which persists to the third generation (F3) (Walsh, Cooley, et al., 2014). Molecular

studies show differential gene expression persisting into F2 after predator cue removal (Hales et

al., 2017), and methylation differences between exposed F0 and naïve F1 individuals (Feiner et

al., 2022; Hearn et al., 2019; Jeremias et al., 2018; Trijau et al., 2018). These findings suggest

that induced traits can persist beyond the directly exposed generation and decline gradually, but

the duration of this persistence and its evolutionary significance remain unclear (Feinberg &

Irizarry, 2010; Ho & Burggren, 2010).

A challenge with such studies is that embryos exist within the mother's brood chamber,

making it possible that they too are directly exposed to the environmental cue (Agrelius &

Dudycha, 2025). Furthermore, those embryos may have developing germ cells that can similarly

be exposed, making it necessary to examine beyond the third (F2) generation (Shahmohamadloo

et al., 2025). Here we directly address the question of how long induced phenotypes can be

inherited in *Daphnia lumholtzi* with a specific focus on periods after which direct exposure must

not have occurred. We measured predator-induced morphological traits in *D. lumholtzi* across the

fourth (F3), fifth (F4), and sixth (F5) generations following initial exposure in F0. *D. lumholtzi*

develops elongated head and tailspines in response to fish predation (Dzialowski et al., 2003),

traits thought to contribute to its invasive success in North America (Engel & Tollrian, 2009;

Sousa et al., 2017; Work & Gophen, 1999) as other *Daphnia* species typically produce helmets

(Horstmann et al., 2021). Previous work in *D. lumholtzi*, shows that maternal exposure can

accelerate defensive trait development in F1 (Graeve et al., 2021) and that cyanobacteria-induced

life history effects persist for at least one generation (Dao et al., 2018), but no study has

examined the long-term persistence of predator-induced morphology in *D. lumholtzi*. We found

that induced defensive traits persisted through four unexposed generations, with measurable

effects disappearing in F5. These results demonstrate that transgenerational plasticity has a

limited temporal duration, with trait-specific patterns of decline that reflect the gradual loss of

ancestral environmental influence (Geoghegan & Spencer, 2012; Heard & Martienssen, 2014).

Methods

Experimental Overview (Fig. 1). To explore the dynamics of predation induced phenotypes

within and across generations, we reared five lineages of predation exposed, and four lineages of

unexposed, *Daphnia lumholtzi* for five generations following initial predation exposure and cue

removal. *Daphnia* were individually grown in 50 mL beakers containing 30 mL of fish

conditioned medium or standard artificial lake medium (COMBO)(Kilham et al., 1998). Every

other day during the F0, F3, F4 and F5 generations, individual *D. lumholtzi* were imaged for

morphological measurements using ImageJ (Schindelin et al., 2012). On this same schedule the

media was exchanged, and algae was replenished with 85,000 cells/mL of a 1:1 mixture of

Scenedesmus obliquus and *Chlorella vulgaris*, hereby referred to as the standard algae mixture.

See Fig. 1 for experimental schematic and sample sizes.

***Daphnia Lumholtzi* Clone.** *Daphnia lumholtzi* clones were collected in the summer of 2019

from Saguaro Lake, AZ, USA. This clone was specifically chosen for its ability to develop

headspines and tailspines in response to vertebrate predation. These predator response

phenotypes are only found in *D. lumholtzi*, whereas other species develop different armor

morphologies.

Common Garden Rearing to Standardize Generational Exposures. *D. lumholtzi* clones were

reared for 6 months in artificial lake medium (COMBO) and fed *Scenedesmus obliquus* and

Chlorella vulgaris ad libitum to dull residual effects of lake predation. Prior to experimental

perturbations, ancestral *D. lumholtzi* were reared for three generations in COMBO medium and

fed a standard algae mixture of *S. obliquus* and *C. vulgaris* to minimize the influence of maternal

effects on experimental generations.

Fish and Control Media Preparation. To prepare fish (induction) medium, 10L of artificial

lake medium was conditioned with five *Gasterosteus aculeatus* for three days prior to filtration

(0.45 μm). Fish were fed 10 *D. lumholtzi* and 10 *Daphnia magna* each day to maximize the

inducing effect. To create control medium, artificial lake medium without fish conditioning was

created in congruence with fish medium. Media was frozen and thawed overnight prior to use.

Initial F0 Predation Exposure and Experimental Conditions. From the third generation of

common garden reared *Daphnia*, one mother was selected due to observed clutch survivability

and size. Nine neonates born within 24 hours from the same mother were isolated and randomly

assigned to treatment groups, either fish exposure or control. Animals were reared individually in

a 50 mL plastic beaker with 30mL of the appropriate medium and fed 5 mL standard algae

mixture every other day. Individual *Daphnia* were staged, imaged, and returned to fresh media

with algae every other day.

Propagation across Generations. Upon reproduction, up to six neonates were randomly

collected and reared individually in 50 mL beakers containing 30 mL of control media only. If

the first clutch did not produce a viable lineage, the next viable clutch was utilized. See Fig. 1 for

samples sizes.

Imaging and Data Collection. Every other day, animals in the F0, F3, F4, and F4 generation

were staged on a glass slide and imaged using `cellSenseEntry` (Olympus Life Science). Scale

bars were added to images during acquisition for calibration during measurements.

Morphological measurements of *headspine* (tip of spine to top of compound eye), *body length*

(top of eye to base of tail), *tailspine* (base of tail to tip of tailspine), *total body length* (tip of

headspine to tip of tailspine) were collected (see Fig.2) using `ImageJ` (Schindelin et al., 2012).

All images were calibrated to a consistent scale using the 'set global scale' function in `ImageJ`,

which converts pixel distances to micrometers based on the embedded scale bar. See Figure 2 for

measurement details.

Statistical Analysis.

Phenotype Transmission Across Generations. Morphological variation was analyzed using

both frequentist and Bayesian mixed-effects models in `R` (R Core Team, 2021), using the `lme4`

(Bates et al., 2015a) and `brms` (Bürkner, 2017) packages respectively. For each generation (*F0*,

F3, *F4*, *F5*) and each trait (*headspine*, *tailspine*), we tested for the effects of predation *treatment*,

with *body length* as a covariate, and a random intercept for sample identity to account for nested

sampling. The model formula used was the following:

232   `spine ~ treatment + bodylength + (1|sample)`

We selected this model structure based on established principles in *Daphnia* morphological

analysis and our experimental design. *Body length* was included as a covariate because spine

length scales allometrically with body size in *Daphnia* and failing to account for this relationship

can confound treatment effects with size-related variation (Laforsch & Tollrian, 2004; Petrusek

et al., 2009). *Treatment* (predator exposure vs. control) was included as the main fixed effect to

test our primary hypothesis of transgenerational inheritance of induced defenses. The random

intercept for sample identity accounts for the hierarchical structure of our data, where multiple

individuals were measured from each experimental replicate, addressing potential pseudo-

replication issues common in experimental ecology (Freeberg & Lucas, 2009; Hurlbert, 1984).

We did not include *age* as a covariate due to its high correlation with *body length* (Pearson's R =

0.85; Supp. Fig. 1). Linear mixed models were evaluated using Kenward-Roger F-tests and *post*

hoc Tukey tests to assess treatment effects. Conditional and marginal R² values, AIC, and BIC

were calculated to assess model fit (See Supplemental Table 1 for model diagnostic summary).

We implemented Bayesian hierarchical models using the `brms` (Bürkner, 2017) package to

estimate treatment effects on spine morphology while accounting for individual-level variation.

These models included fixed effects for *treatment*, *body length*, and random intercepts for

individual identity to capture repeated measures. The hierarchical structure allows partial pooling

across individuals, improving estimation in the presence of sparse data. Posterior distributions

were used to quantify uncertainty, and model diagnostics (e.g., Rhat, ESS, posterior predictive

checks) confirmed convergence and reliability. Marginal and conditional R² values were

computed using the `performance` (Lüdecke et al., 2021) package to assess model fit. See

Supplemental Figures 3 and 4 for full diagnostic plots for headspine and tailspine models across

generations.

Models were fit using a Student's t-distribution to account for potential outliers and

heavy-tailed data which are common in ecological trait data. This choice was supported by

posterior predictive checks and traceplot diagnostics indicating stable convergence and reliable

inference. We used weakly informative priors and ran three chains with 5,000 iterations each

(2,000 warm-up), ensuring convergence ($\hat{R} < 1.01$ for all parameters). Posterior distributions

were summarized to extract treatment effect estimates and 95% credible intervals. Bayesian R²

values (marginal and conditional) were calculated to assess model fit (See Supplemental Table 2

for model diagnostic summary). Posterior predictions were generated using `posterior_epred()`

(Gabry et al., 2025) to estimate treatment effects at the mean body length for each generation,

controlling for sample-level variation. Models were evaluated using leave-one-out- cross

validation and WAIC (Vehtari et al., 2017). Bayesian and frequentist estimates were highly

consistent across generations and traits, with overlapping intervals and similar effect sizes,

supporting the robustness of our findings across inferential frameworks. Diagnostic plots for

Bayesian models are shown in Supplemental Figures 3 and 4

Age-Dependent Morphological Plasticity (Delta). To assess how morphological traits

change across *age* and *treatment*, we calculated within-individual morphological change (Δ) for

body length, *total body size*, *headspine*, and *tailspine*. Δ values were computed as the difference

between successive measurements for each individual (Δ`trait = trait_current -`

`trait_previous`), capturing short-term plastic responses over time. Age trajectories (*Δ trait* vs.

age) were visualized using scatterplots with LOESS-smoothed trend lines and 95% confidence

intervals. To statistically evaluate the effect of induction on morphological plasticity, we

modeled Δ values using linear mixed-effects models `lme4` (Bates et al., 2015b) and `lmerTest`

package (Kuznetsova et al., 2017), with *treatment* and *age* as fixed effects and individual identity

as a random intercept to account for repeated measures `(formula: delta value ~ treatment`

`+ age + (1|sample))`. We initially explored more complex temporal modeling approaches,

including natural splines with varying degrees of freedom (df = 2 to 4) to capture potential non-

linear age-related changes in growth patterns (`formula: delta value ~ treatment +`

`ns(age, df = 2) + (1|sample))`. However, these models suffered from overparameterization

given our sample size, and model significance was sensitive to the choice of degrees of freedom,

indicating unstable parameter estimates. Consequently, we adopted the more parsimonious linear

approach with *age* as a continuous covariate, which provided robust and interpretable results

while adequately capturing the age-related trends in our data. Model significance was assessed

using ANOVA to compare treatment effects. Results were visualized in a 2×2 panel plot

showing estimated trait changes and associated standard errors across treatments.

Size Selective Survivorship Analysis. To test whether size-selective mortality, rather than

treatment, could explain differences in spine lengths, we constructed mixed-effects linear models

using binary survival status. Survival was defined as the presence or absence of a subsequent

timepoint for each individual. Separate models were fit for *headspine* and *tailspine* lengths as a

function of *body length*, with a random intercept for sample identity (`spine ~ bodylength +`

`(1|sample)`). Residuals from these models represent spine length variation independent of body

size. To visualize trait dynamics across age and survival status, we plotted body length and spine

residuals using LOESS-smoothed trend lines with 95% confidence intervals. Survivors and non-

survivors were distinguished by color, and traits were faceted for comparison.

To further evaluate survival probability, we constructed a binomial generalized additive

model (GAM) with *survival status* as the response variable and *age*, *body length*, *spine* lengths,

treatment, and the interaction between *age* and *treatment* as predictors (`status ~ bodylength`

`+ headspine + tailspine + age * treatment`). Trait values representing the 10th, 50th,

and 90th percentiles were used to generate survival predictions across age. Predicted survival

probabilities and 95% confidence intervals were plotted to assess how trait variation influences

survivorship over time.

Statistical Software and Reproducibility. All statistical analyses and visualizations were

conducted in `R` (R Core Team, 2021). Data manipulation and cleaning were performed using the

`dplyr` package (Wickham et al., 2022) , and visualizations were generated using `ggplot2`

(Wickham, 2016), with additional formatting and layout support from `patchwork` (Pedersen,

2025), `gghighcontrast` (Grosenbach, 2021) , and `Rmisc` (Hope, 2022) for summary statistics.

Estimated marginal means and post hoc comparisons were calculated using `emmeans` (Searle et

al., 2025).

All analyses were conducted using `Rstudio` (Posit Team, 2025) in `RMarkdown`

(Allaire et al., 2019) and `Quarto` (Allaire et al., 2025), ensuring full reproducibility of code,

figures, and statistical outputs. Annotated Quarto files and datasets are available on GitHub at

https://github.com/shannonsnyder/daphnia-transgenerational-plasticity.

Results

Ontogenetic Trajectories Demonstrate Environmental Effects Targeted to Defensive Traits

Daphnia lumholtzi exhibited distinct developmental trajectories across their full lifespan

for body length, headspine, and tailspine structures (Fig. 3). Body length increased rapidly as the

animals aged in both control and induced lineages, reaching an early asymptote (~2,500 μm)

around 25 days. This growth pattern was strongly correlated with age (R = 0.85; Supp. Fig. S1),

and no significant differences were observed between treatments, indicating that predator cue

exposure did not affect overall somatic growth.

In contrast, defensive spine traits displayed age-dependent plasticity, with early-life

elongation followed by a gradual reduction in spine length. Headspine length peaked (~850 μm)

within the first 10 days and declined steadily to ~400 μm in older individuals (Fig. 3B). Tailspine

length reached a delayed maximum (~1,800 μm) around 20 days before declining to values

comparable to those observed in early developmental stages (~700 μm) (Fig. 3C). These patterns

suggest that predator-induced plasticity is most pronounced during early developmental stages

and diminishes progressively across the lifespan of *Daphnia*, consistent with a life-stage–specific

defense strategy.

Together, these ontogenetic analyses demonstrate that transgenerational inheritance of

predator-induced traits is specific to defensive structures and occurs independently of changes in

body size or growth rate, supporting the hypothesis that environmental induction targets adaptive

traits without altering general somatic development.

Predator cue exposure results in an increase in headspine and tailspine length in the

exposed generation.

Postembryonic predator cue exposure resulted in a substantial increase in both headspine

and tailspine length relative to unexposed controls. In the F0 generation, with predation cues

provided from the first post-embryonic day, both linear and Bayesian models estimate significant

increases in induced defense trait lengths (Fig. 4; Tables 1,2). The linear mixed-effects model

provided an estimated increase in headspine length of 68.98 ± 27.19 μm (F-stat = 6.44,

$p = 0.035$), closely aligning with the Bayesian estimate of 68.94 ± 32.26 μm. These differences

correspond to a roughly 10% increase in headspine length over control *Daphnia* never exposed

to fish conditioned medium. Tailspine length also increased dramatically in the exposed F0

lineage relative to controls, by 114.73 ± 31.93 μm (F-stat = 12.91 , $p = 0.008$; linear model) and

100.65 ± 39.75 μm (Bayesian model), representing a roughly 10% increase. Because we noted

that most of the morphological changes occur within the first 20 days of life, before spine growth

plateaus (Fig. 4), we modeled these effects on *Daphnia* less than 20 days of age.

Gen	Spine	Est. Difference (μm)	Std_Error	R^2 marginal	R^2 conditional	F statistic	P value
F0	Headspine	68.98	27.19	0.58	0.60	6.44	0.035
F0	Tailspine	114.73	31.93	0.93	0.94	12.91	0.008
F3	Headspine	37.02	17.61	0.51	0.65	4.42	0.038
F3	Tailspine	8.19	18.28	0.95	0.98	0.20	0.655
F4	Headspine	43.81	20.19	0.48	0.66	4.71	0.034
F4	Tailspine	85.45	34.14	0.90	0.98	6.26	0.015
F5	Headspine	-5.79	21.41	0.51	0.53	0.07	0.789
F5	Tailspine	-24.21	39.07	0.93	0.97	0.38	0.539

354
Table 1. Linear mixed-effects model estimates of treatment-induced changes in spine length across
generations in Daphnia. Models were fit separately by generation for individuals ≤ 20 days old. Each row
represents a unique combination of generation and spine type. "Gen" indicates the generation of the experimental
cohort (e.g., F0, F3, etc.), and "Spine" refers to the type of spine measured, either headspine or tailspine. "Est.
Difference (μm)" is the estimated mean difference in spine length (in micrometers) between predator-induced and
control treatments, where positive values indicate longer spines in induced individuals. "Std_Error" is the standard
error of the estimated difference. "R² marginal" and "R² conditional" represent the proportion of variance explained
by fixed effects alone and by both fixed and random effects, respectively. The "F statistic" and associated "P value"
test the significance of the treatment effect; p-values < 0.05 indicate statistically significant differences. See
Methods for full model specification and random effect structure.

Gen	Spine	Est. Difference (μm)	Std_Error	R2_marginal	R2_conditional
F0	Headspine	68.94	32.26	0.60	0.047
F0	Tailspine	100.65	39.75	0.95	0.003
F3	Headspine	37.95	18.19	0.65	0.025
F3	Tailspine	6.21	18.24	0.98	0.001
F4	Headspine	44.58	20.79	0.66	0.025
F4	Tailspine	85.82	34.29	0.98	0.001
F5	Headspine	-6.78	22.28	0.56	0.034
F5	Tailspine	-19.34	29.32	0.95	0.004

Table 2. Posterior mean estimates of treatment-induced changes in spine length across generations in *Daphnia*, based on Bayesian hierarchical models. Models were fit separately by generation for individuals ≤ 20 days old. Each row represents a unique combination of generation and spine type. "Gen" indicates the generation of the experimental cohort, and "Spine" refers to the type of spine measured - either headspine or tailspine. "Est. Difference (μm)" is the posterior mean estimate of the difference in spine length (in micrometers) between predator-induced and control treatments, with positive values indicating elongation in induced individuals. "Std_Error" is the standard error of the posterior estimate. "R^2 marginal" and "R^2 conditional" represent the proportion of variance explained by fixed effects alone and by both fixed and random effects, respectively. See Methods for full model specification and prior structure. Note: Lower R^2 values in some models likely reflect limited variation among clonal individuals, consistent with the use of genetically identical lineages in this study.

Defensive Traits Are Inherited Through F4 but Disappear in F5.

Consistent with well-established within-generation plastic responses, F0 *Daphnia* reared in predator cue environments exhibited robust and statistically significant increases in both headspine and tailspine length relative to controls. Frequentist estimates indicated increases of 68.98 μm ($p = 0.035$) for headspine and 114.73 μm ($p = 0.008$) for tailspine, with Bayesian estimates closely corroborating these results (headspine: 68.94 ± 32.26; tailspine: 100.65 ±39.75; Tables 1 and 2).

Strikingly, phenotypic differences persisted for multiple generations after removal from the inducing environment. F3 represents the first generation that was not exposed to predator cues either directly or indirectly. When F0 mothers are exposed, their developing embryos (F1) and the germline cells that will become the F2 generation are also exposed. Only by the F3

generation are the effects of induction potentially transmitted through purely epigenetic

mechanisms.

Despite two intervening generations without exposure, F3 individuals from the induced

lineage exhibited a significant increase in headspine length relative to controls. Frequentist

analysis estimated a 37.02 μm increase (SE = 17.61, F = 4.42, p = 0.038), with Bayesian

inference yielding a nearly identical effect size (37.95 μm, SE = 18.19). This represents

approximately 50% retention of the original F0 headspine response. In contrast, tailspine length

in F3 showed no significant difference between lineages (frequentist: 8.19 μm, p = 0.655;

Bayesian: 6.21 μm), indicating trait-specific attenuation of the induced phenotype.

In the F4 generation, also completely unexposed to predator cues, both headspine and

tailspine lengths were again significantly elevated in the induced *Daphnia* as compared to the

control *Daphnia*. Frequentist estimates showed a 43.81 μm increase in headspine length (p =

0.034) and an 85.45 μm increase in tailspine length (p = 0.015), with Bayesian estimates closely

matching (headspine: 44.58 μm; tailspine: 85.82 μm). These values reflect retention of about

50% of the original F0 effect sizes, similar to the F3 estimates (Tables 1 and 2).

By the F5 generation all measurable differences between induced and control lineages

had disappeared. Frequentist analysis estimated a negligible difference in headspine length (–

5.79 μm, p = 0.789) and tailspine length (–24.21 μm, p = 0.539), with Bayesian estimates

similarly non-significant. These findings indicate that the induced phenotypes ultimately reverted

to baseline in the absence of continued environmental cues over 5 unexposed generations. See

Supplemental Figure 2 for raw trait values.

Taken together, these results demonstrate that predator-induced morphological plasticity

can be inherited for at least four generations, but this inheritance is finite and trait-dependent.

The disappearance of measurable trait differences in F5 underscores the transient nature of this

plasticity and suggests that the mechanism of transgenerational epigenetic inheritance decays

over time without environmental cue reinforcement within several generations (see Fig. 4).

Differences in Growth Rate and Trait Distributions Do Not Explain Induced Trait

Expression.

To evaluate whether the observed morphological differences between induced and

control lineages could be attributed to variation in growth dynamics or trait distributions, rather

than transgenerational induction, we assessed whether treatment effects could be explained by

differences in developmental timing, size-selective mortality, or trait-based survival.

The most pronounced morphological changes occurred early in life, prior to 20 days of

age (Figure 4). This period includes the onset of reproduction (typically 7–12 days), during

which *D. lumholtzi* mothers are not yet fully grown. Morphological change peaked around this

time for all traits, particularly for body length and total body size.

Importantly no significant differences in growth rate were detected between treatment

groups for any of the four traits examined (body length, headspine, tailspine, and total body

length), indicating that predator exposure did not alter the rate of somatic or defensive trait

development (Figure 5). Specifically, treatment effects were non-significant for body length (Est.

$= 7.73 \pm 16.72$ μm, $F = 0.22$, $p = 0.642$), total body size (Est. $= 0.78 \pm 33.30$ μm, $F = 0.00$, $p =$

0.981), headspine (Est. $= -2.00 \pm 6.33$ μm, $F = 0.10$, $p = 0.751$), and tailspine (Est. $= -4.96 \pm$

12.11 μm, F = 0.17, p = 0.681). These results suggest that the observed morphological

differences are not due to accelerated or delayed growth in induced individuals but instead reflect

differences in trait expression independent of growth rate.

To assess whether trait distributions might reflect selective mortality, such as the

preferential survival of individuals with longer spines, we compared growth trajectories of

survivors and non-survivors during the first 20 days of life. Plotting developmental trajectories

for body length, headspine residuals, and tailspine residuals (controlling for body size) by

survival status (Fig. 6) showed that while survivors tended to be slightly larger in body length,

residual spine lengths showed overlapping distributions and trajectories between survivors and

non-survivors. These patterns indicate that trait-based selection is unlikely to account for the

observed morphological differences.

The absence of an effect of spine length on survivorship was further supported by a

generalized linear model (GLM) predicting survival status from body length, headspine,

tailspine, age, and treatment. The model revealed that body length was positively associated with

survival ($\beta = 0.0030$, $p < 0.001$), while headspine and tailspine lengths were not significant

predictors (headspine: $\beta = -0.0010$, $p = 0.058$; tailspine: $\beta = -0.0007$, $p = 0.110$). Treatment had

a significant positive effect on survival ($\beta = 0.678$, $p < 0.001$), but the interaction between age

and treatment was not significant ($\beta = -0.0145$, $p = 0.096$). The model explained a substantial

portion of deviance (ΔDeviance = 328.8; AIC = 2554.1), but no evidence emerged for size- or

trait-selective mortality based on spine length (Figure 7).

Discussion

Induced Defenses Are Robust, Age-Specific, and Not Driven by Selection

Exposure to predator cues in the F0 generation triggered strong morphological defenses in *Daphnia lumholtzi*, with significant elongation of both headspine and tailspine structures. Notably, post-embryonic exposure alone was sufficient to induce these traits in our clone collected from Arizona, USA, contrasting with previous findings in other *D. lumholtzi* populations suggesting embryonic exposure was necessary for trait development (Graeve et al., 2021; Laforsch & Tollrian, 2004). This finding underscores the differences in inducibility between clones of the same species and highlights the need to assess transgenerational inheritance in a variety of systems (Auld et al., 2010; Scoville & Pfrender, 2010; Shahmohamadloo et al., 2025). This is further illustrated by the characterization of "uninducible" *D. lumholtzi* clones which constitutively express defense phenotypes in the absence of predation cues (Engel & Tollrian, 2009), collected just miles away from our Saguaro Lake, AZ clone at Canyon Lake, AZ, USA.

Importantly, these trait differences were not attributable to size-selective or trait-selective mortality. Survivors and non-survivors showed overlapping developmental trajectories, and survival probability was primarily associated with body length, not spine morphology. This confirms that the observed phenotypic differences are induced responses rather than artifacts of differential survivorship among individuals based upon armor morphology (Dieckmann et al., 2006; Kingsolver et al., 2001).

We also found that induced spine lengths were most prominent in early life and declined with age, a pattern distinct from somatic growth. This suggests stage-specific optimization of defenses, likely targeting periods of highest vulnerability to predation (Nagano & Yoshida, 2020;

E. E. Werner & Gilliam, 1984). The age-dependent regression of defensive traits, including

previously unreported headspine shortening with age, further highlights the dynamic nature of

plastic responses and the influence of genetic background (DeWitt et al., 1998; Scheiner, 1993;

Via et al., 1995).

Induced Phenotypes Persists Across Generations but Fades Without Cues

Our results demonstrate that predator-induced defenses are not confined to the directly

exposed generation, nor the immediate offspring of the exposed generation. Headspine and

tailspine elongation persisted through four unexposed generations, with measurable effects in F3

and F4 despite the absence of continued induction and lack of indirect exposure through

germline conditioning. This multigenerational persistence supports the hypothesis that

environmental information can be retained and expressed across generations via non-genetic

mechanisms (Gapp et al., 2014; Heard & Martienssen, 2014; Rechavi et al., 2014).

By the F5 generation, however, trait differences between induced and control lineages

had disappeared. This regression to baseline aligns with evolutionary expectations that plastic

traits fade when environmental cues vanish, minimizing fitness costs associated with

unnecessary defenses (DeWitt et al., 1998; Van Buskirk & Steiner, 2009). The trait-specific

dynamics, headspine persistence in F3, tailspine resurgence in F4, suggest that transgenerational

plasticity is not uniform and may fluctuate across lineages (Bonduriansky & Day, 2009; Herman

& Sultan, 2011; Uller et al., 2013).

Mechanisms of Non-genetic Inheritance and Evolutionary Implications

The persistence and gradual decline of induced traits across generations lead to the strong

testable hypothesis of epigenetic inheritance as a likely mechanism. DNA methylation, histone

modification, and other chromatin-level changes may facilitate the transmission of plasticity

without altering the genetic code (Banerjee & Roychoudhury, 2017; Casadesús, 2016;

Davidowitz, 2009; Ho & Burggren, 2010; Li & Zhang, 2014; Niederhuth et al., 2016; Robichaud

et al., 2012; M. S. Werner et al., 2023; Yaniv, 2014). This pattern is consistent with incomplete

erasure of epigenetic marks and has been observed in other invertebrate and vertebrate systems

(Dias & Ressler, 2014; Gapp et al., 2014; Grishkevich & Yanai, 2013). In *Caenorhabditis*

elegans, the histone mark H3K27me3 was shown to persist across generations, influencing gene

expression and development in offspring (Kaneshiro et al., 2019). In the freshwater snail, *Physa*

acuta, predator-induced gene expression changes were maintained across two generations,

supporting epigenetic regulation (Dejeux et al., 2025). In mice, paternal diet altered offspring

DNA methylation and metabolic gene expression (Carone et al., 2010). In humans, trauma-

related DNA methylation patterns were found to persist across three generations (Mulligan et al.,

2025). Similarly, in *Daphnia magna*, differential DNA methylation patterns induced by

environmental stressors have been documented to persist for multiple generations before

returning to baseline (Asselman et al., 2016; Vandegehuchte et al., 2010)

Our findings underscore the evolutionary relevance of environmental history even in the

absence of genetic variation (Jablonka & Lamb, 20062; Laland et al., 2015; Pigliucci & Müller,

2010). Transgenerational plasticity allows organisms to respond to ancestral conditions,

potentially buffering populations against short-term environmental fluctuations (Donelson et al.,

2018; Uller et al., 2018). However, because our study used a single clone, future work should

explore responses across diverse genotypes (Auld et al., 2010). The existence of "uninducible"

clones with constitutive defenses raises intriguing questions about genetic assimilation and the

long-term fate of plastic traits (Lande, 2009; West-Eberhard, 2003).

Conclusions

In evolutionary genetics, it is still unclear how general the non-genetic inheritance of

induced phenotypes is across organisms, and when it occurs, how long the induced phenotypes

will persist in the absence of the signal. Our work contributes to answering both questions by

demonstrating that predator-induced morphological defenses in *Daphnia lumholtzi* can persist for

up to four generations through non-genetic mechanisms but ultimately regress in the absence of

continued environmental cues. This finding aligns with previous work in *Daphnia*, such as clutch

size persistence to F3 in *D. ambigua* (Walsh, Whittington, et al., 2014), helmet length inheritance

to F2 in *D. cucullata* (Agrawal et al., 1999a) and multi-generational erosion of plastic traits in *D.*

pulicaria (Landy et al., 2024), but is novel in demonstrating morphological trait persistence

through four unexposed generations. Were this phenomenon to occur in humans, it would be

equivalent to the environmental conditions experienced by our great-grandparents in the 19th

century shaping our phenotypes today, a striking reminder of how ancestral environments may

echo across generations. This transgenerational plasticity may buffer populations against short-

term ecological fluctuations, while its eventual attenuation minimizes the costs of unnecessary

trait expression (Herman & Sultan, 2011; Uller et al., 2013). Future research should aim to

identify the molecular basis of this inheritance (Klosin et al., 2017; Trijau et al., 2018) and

explore its evolutionary consequences across diverse genetic backgrounds and dynamic

environments.

Acknowledgements

We would like to thank Michael Lynch for his contribution of SAG *Daphnia lumholtzi* clones

utilized in this study. We would also like to thank Mark Currey for this assistance in stickleback

fish husbandry and handling and Emily Williams for the collection and maintenance of *D.*

lumholtzi lineages prior to 2020.

FIGURES

Figure 1. Schematic illustrating the experimental setup allowing for direct testing of
transgenerational epigenetic inheritance by induced defenses by comparing lineages with
and without grandparental exposure to predator cues. Offspring (F0) from a single
unexposed mother were split into two lineages: control (top), where individuals where
continuously reared in artificial lake media, and induced (bottom), where F0 individuals were
exposed to fish-conditioned media indicated in green. Only the F0 induced lineage is exposed to
fish conditioned media. Sample size for each generation and treatment is indicated. Subsequent
generation (F3-F5) are from both lineages but are maintained in artificial lake media. Lineages
were propagated for six generations and measured every other day during the F0, F3, F4, and F5
generations. Schematic distinguishes plasticity from persistent heritable effects.

565
566
Figure 2. Inducible morphological defense in Daphnia in response to predator cues.
Daphnia reared in fish-conditioned media develop significantly elongated headspine and
tailspines compared to control. Representative lateral images of uninduced animal reared in
artificial lake media (left) and induced animal, reared in fish conditioned media (right).
Individuals exposed to fish cues (right) exhibit pronounced elongation of the headspine and
tailspine, consistent with a predator-induced defense phenotype. Green lines indicate locations
and dimensions for standard measurements of headspine length, body length, and tailspine
length, detailed as follows: Headspine was measured from tip of spine to top of compound eye,
body length was measured from top of compound eye to base of tailspine, tailspine was
measured from base to tip of spine. Total body size was measured from top of headspine to tip of
tailspine. The scale bar corresponds to 500 μm.

578
Figure 3. Trajectories of somatic and defensive traits reveal divergence in late-life
dynamics. Scatterplots display individual measurements of (A) body length, (B) total body size,
(C) headspine length, and (D) tailspine length (in microns) as a function of age (in days). Green
line represents locally averaged (GAM) trends for induced lineage (F0 exposed to fish-
conditioned media); purple line represents locally averages (GAM) trends for control lineage (no
F0 exposure). Somatic traits - body length (A) and total body size (B) - increase rapidly during
early juvenile stages, then reach a plateau in mid-life, with little elongation even at advanced
ages. In contrast, defensive traits - headspine (C) and tailspine (D) – regress later in life. These
results illustrate that the timing and persistence of morphological development vary substantially
between defensive and somatic phenotypes. Morphological trajectories demonstrate distinct
patterns in defense trait expression, but not body growth, between lineages, consistent with the
inheritance of plastic response initiated by ancestral exposure.
591

Figure 4. Predicted spine lengths by generation and treatment from linear and
Bayesian models. Predicted mean headspine length (top row) and tailspine length
(bottom row) for *Daphnia* in control and induced lineages across generations
(F0,F3,F4,F5). Points represent model-predicted means ± standard error, with black
circles for linear mixed-effect model estimates and open grey circles for Bayesian
hierarchical model posterior means. Both model frameworks show increased spine length
in induced (fish-conditioned) lineages and yield qualitatively similar inferences,
supporting robustness of transgenerational treatment effects. Estimates are based on data
from individuals ≤ 20 days old. See tables 1 and 2 for underlying estimates and model
parameters.

610
611
Figure 5. Growth of defensive and somatic traits are most pronounced early in life,
followed by a stabilization with age and were unaffected by treatment. Changes in
morphological measurements (body length, total body size, headspine, and tailspine) across the
lifespan (days). Each panel shows the mean growth (Δ; in μm) from the previous timepoint for
the indicated trait. Morphological changes are largest early in life and stabilize as individuals
age, as shown by the bold smoothed trend line with 95% confidence intervals (shaded area).

629
Figure 6. Developmental trajectories during periods of concentrated growth do not
indicate treatment or size selective survivorship. Body length, head spine (residual), and tail
spine (residual) measurements over first 20 days of age for individuals grouped by survival status
(purple for those who survived to the next timepoint - survivors, teal for those that perished).
Residual values for head and tail spines are calculated from linear mixed-effects models that
account for body length, isolating variation in spine length independent of body size.
Both survivors and non-survivors display overlapping developmental trajectories and trait
distributions, providing no evidence that size-selective or trait-selective mortality underlies
observed group differences.
639
640
641
642

643
Figure 7. Survivorship and morphological trait distributions do not support size-selective mortality as a driver of morphological differences. Modeled probability of survival to the next timepoint as a function of age, body length, spine length, and treatment, based on a generalized linear model. High body length values are associated with a greater likelihood of survival at most ages, whereas spine lengths show minimal effect. This suggests that while body length does relate to survival probability in the model, size-selective mortality is not strong enough to cause large differences in defensive morpholgies.

651
652

Model	Generation	Trait	Marginal R^2	Conditional R^2	AIC	BIC
Linear	F0	headspine	0.584	0.597	881.039	892.492
Linear	F0	tailspine	0.934	0.942	870.549	882.001
Linear	F3	headspine	0.514	0.650	3,651.041	3,669.772
Linear	F3	tailspine	0.950	0.981	3,485.125	3,503.856
Linear	F4	headspine	0.477	0.659	3,315.954	3,334.286
Linear	F4	tailspine	0.903	0.984	3,225.705	3,244.037
Linear	F5	headspine	0.507	0.526	2,454.164	2,470.852
Linear	F5	tailspine	0.931	0.966	2,427.085	2,443.773

Supplemental Table 1: Goodness-of-fit and diagnostic statistics for linear mixed-effects
models. Models were fit independently for each combination of experimental generation (F0,
F3, F4, F5) and morphological trait (headspine, tailspine). The proportion of variance explained
by the models is presented as Marginal R^2 (variance explained by fixed effects alone) and
Conditional R^2 (variance explained by both fixed and random effects). Model fit was assessed
using the Akaike Information Criterion (AIC) and the Bayesian Information Criterion (BIC),
where lower values indicate a better trade-off between model fit and complexity. The
consistently high Conditional R^2 values, particularly for the tailspine trait, indicate that the
random effects structure captured a significant portion of the unexplained variance.
662

Generation	Trait	Max Rhat	Min Bulk ESS	Min Tail ESS	LOOIC	WAIC	p(Direction)	Marginal R²
F0	headspine	1.00	10025	14059	889.10	888.85	0.98	0.60
F0	tailspine	1.00	9045	15377	854.56	854.44	0.99	0.95
F3	headspine	1.00	10641	21785	3,634.53	3,632.48	0.98	0.65
F3	tailspine	1.00	11265	18072	3,408.11	3,401.26	0.62	0.98
F4	headspine	1.00	11813	21833	3,296.61	3,294.12	0.98	0.66
F4	tailspine	1.00	5359	10355	3,111.48	3,101.43	0.99	0.98
F5	headspine	1.00	7366	13477	2,451.79	2,451.35	0.63	0.56
F5	tailspine	1.00	7322	9952	2,274.57	2,270.67	0.75	0.95

Supplemental Table 2: Bayesian Model Diagnostics and Fit Summary. Diagnostics are
presented for the Bayesian mixed-effects models fit separately for each combination of
experimental generation (F0–F5) and trait (headspine, tailspine). Model convergence was
assessed using the maximum potential scale reduction factor (Max Rhat), which should be
approximately 1.0, and the minimum effective sample sizes for the bulk (Min Bulk ESS) and
tails (Min Tail ESS) of the posterior distributions, with higher values indicating more reliable
estimates. Model fit and predictive accuracy were compared using the Leave-One-Out
Information Criterion (LOOIC) and the Widely Applicable Information Criterion (WAIC); lower
values suggest better out-of-sample fit. The probability of direction, p(Direction), represents the
posterior probability that the treatment effect is positive. The Marginal R² is the proportion of
variance explained by the model's fixed effects. All models demonstrated convergence.

674
675
Supplemental Figure 1. Ontogenetic growth trajectories for Daphnia body length across
treatments show high correlation between age and body length. Scatterplot of individual
body length measurements (µm) as a function of age (days) for control (open circles) and
induced (filled circles) treatment. The solid line shows the overall fitted growth curve (R=0.85)
across all data. Growth patterns are similar between treatment, indicating that environmental
induction principally affects defensive traits rather than somatic traits in this clone.
682

683
Supplemental Figure 2. Transgenerational patterns of headspine and tailspine length in Daphnia
following ancestral predator cue exposure. Mean (± standard error) headspine (top panels) and
tailspine (bottom panels) values are shown for control and induced lineages across four non-
exposed generations (F3–F5) and the originally exposed generation (F0). Each panel displays
pairwise comparisons between control (open circle) and induced (closed circle) groups for the
indicated generation. Significant induction of both headspine and tailspine length is evident in
predator-cue–induced F0 animals, with differential persistence of these defensive traits in
subsequent generations. While headspine elongation remains significant through F3 and F4
before declining in F5, tailspine induction fluctuates, with significant retention re-emerging in F4
but lost by F5. These data illustrate the finite and dynamic pattern of transgenerational plasticity
in response to environmental cues. See Methods for statistical details and Tables 1,2 for full
effect sizes.
696

697
Supplemental Figure 3: **Bayesian model diagnostic plots for the headspine trait across**
generations. Each row represents an independent model. The panels display posterior predictive
checks (model fit), residuals vs. fitted values (homoscedasticity), MCMC traceplots
(convergence), and Quantile-Quantile (QQ) plots (normality of residuals). The successful
convergence of the MCMC chains and the adequate fit shown by the predictive checks and
residual plots support the validity of the models for inference.
704
705
706

707
Supplemental Figure 4: Bayesian model diagnostic plots for the tailspine trait across
generations. Each row represents an independent model. Plots include posterior predictive
checks (model fit), residuals vs. fitted values (homoscedasticity), MCMC traceplots
(convergence), and Quantile-Quantile (QQ) plots (normality of residuals). The posterior
predictive checks show good overlap between observed and simulated data, and the traceplots
indicate stable, well-mixed chains. The residuals show no major patterns and are approximately
normally distributed, supporting the validity of the models.
715
716
717

<div align="center">Literature Cited</div>

Adamczuk, M. (2020). Population dynamics and life history traits of Daphnia magna across thermal regimes of environments. *Science of The Total Environment*, *723*, 137963. https://doi.org/10.1016/J.SCITOTENV.2020.137963

Agrawal, A. A., Laforsch, C., & Tollrian, R. (1999a). Transgenerational induction of defences in animals and plants. *Nature 1999 401:6748*, *401*(6748), 60–63. https://doi.org/10.1038/43425

Agrawal, A. A., Laforsch, C., & Tollrian, R. (1999b). Transgenerational induction of defences in animals and plants. *Nature 1999 401:6748*, *401*(6748), 60–63. https://doi.org/10.1038/43425

Agrelius, T. C., & Dudycha, J. L. (2025). Maternal effects in the model system Daphnia: the ecological past meets the epigenetic future. *Heredity*, *134*(2), 142–154. https://doi.org/10.1038/S41437-024-00742-W

Allaire, J., Teague, C., Scheidegger, C., Yihui, X., Dervieux, C., & Woodhull, G. (2025). *Quarto*. https://github.com/quarto-dev/quarto-cli

Allaire, J., Xie, Y., Dervieux, C., Grolemund, Garrett., McPherson, J., Luraschi, J., Ushey, K., Atkins, A., Wickham, H., Cheng, J., Chang, W., & Iannnone, R. (2019). *rmarkdown: Dynamic Documents for R* (p. 303). CRC Press, Taylor & Francis Group. https://github.com/rstudio/rmarkdown

Angilletta, M. J. (2009). Thermal Adaptation: A Theoretical and Empirical Synthesis. *Thermal Adaptation: A Theoretical and Empirical Synthesis*, 1–302. https://doi.org/10.1093/ACPROF:OSO/9780198570875.001.1

Asselman, J., De Coninck, D. I. M., Pfrender, M. E., & De Schamphelaere, K. A. C. (2016). Gene body methylation patterns in daphnia are associated with gene family size. *Genome Biology and Evolution*, *8*(4), 1185–1196. https://doi.org/10.1093/gbe/evw069

Auld, J. R., Agrawal, A. A., & Relyea, R. A. (2010). Re-evaluating the costs and limits of adaptive phenotypic plasticity. *Proceedings. Biological Sciences*, *277*(1681), 503–511. https://doi.org/10.1098/RSPB.2009.1355

Banerjee, A., & Roychoudhury, A. (2017). Epigenetic regulation during salinity and drought stress in plants: Histone modifications and DNA methylation. *Plant Gene*, *11*, 199–204. https://doi.org/10.1016/J.PLGENE.2017.05.011

Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2015a). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*(1). https://doi.org/10.18637/JSS.V067.I01

Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2015b). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*(1). https://doi.org/10.18637/JSS.V067.I01

Beaman, J. E., White, C. R., & Seebacher, F. (2016). Evolution of Plasticity: Mechanistic Link between Development and Reversible Acclimation. *Trends in Ecology & Evolution*, *31*(3), 237–249. https://doi.org/10.1016/J.TREE.2016.01.004

Beklioglu, M., Telli, M., & Gozen, A. G. (2006). Fish and mucus-dwelling bacteria interact to produce a kairomone that induces diel vertical migration in Daphnia. *Freshwater Biology*, *51*(12), 2200–2206. https://doi.org/10.1111/j.1365-2427.2006.01642.x

Bonduriansky, R., & Day, T. (2009). Nongenetic inheritance and its evolutionary implications. *Annual Review of Ecology, Evolution, and Systematics*, *40*(Volume 40, 2009), 103–125. https://doi.org/10.1146/ANNUREV.ECOLSYS.39.110707.173441/CITE/REFWORKS

Botero, C. A., Weissing, F. J., Wright, J., & Rubenstein, D. R. (2015). Evolutionary tipping points in the capacity to adapt to environmental change. *Proceedings of the National Academy of Sciences of the United States of America*, *112*(1), 184–189. https://doi.org/10.1073/PNAS.1408589111

Bradshaw, A. D. (1965). Evolutionary Significance of Phenotypic Plasticity in Plants. *Advances in Genetics*, *13*(C), 115–155. https://doi.org/10.1016/S0065-2660(08)60048-6

Bürkner, P. C. (2017). brms: An R package for Bayesian multilevel models using Stan. *Journal of Statistical Software*, *80*. https://doi.org/10.18637/JSS.V080.I01

Carone, B. R., Fauquier, L., Habib, N., Shea, J. M., Hart, C. E., Li, R., Bock, C., Li, C., Gu, H., Zamore, P. D., Meissner, A., Weng, Z., Hofmann, H. A., Friedman, N., & Rando, O. J. (2010). Paternally induced transgenerational environmental reprogramming of metabolic gene expression in mammals. *Cell*, *143*(7), 1084–1096. https://doi.org/10.1016/j.cell.2010.12.008

Casadesús, J. (2016). Bacterial DNA methylation and methylomes. *Advances in Experimental Medicine and Biology*, *945*, 35–61. https://doi.org/10.1007/978-3-319-43624-1_3/FIGURES/3

Corl, A., Bi, K., Luke, C., Challa, A. S., Stern, A. J., Sinervo, B., & Nielsen, R. (2018). The Genetic Basis of Adaptation following Plastic Changes in Coloration in a Novel Environment. *Current Biology*, *28*(18), 2970-2977.e7. https://doi.org/10.1016/j.cub.2018.06.075

Dao, T. S., Vo, T. M. C., Wiegand, C., Bui, B. T., & Dinh, K. V. (2018). Transgenerational effects of cyanobacterial toxins on a tropical micro-crustacean Daphnia lumholtzi across three generations. *Environmental Pollution (Barking, Essex : 1987)*, *243*(Pt B), 791–799. https://doi.org/10.1016/J.ENVPOL.2018.09.055

Davidowitz, G. (2009). Ecological Developmental Biology: Integrating Epigenetics, Medicine and Evolution. Scott F. Gilbert and David Epel. *Integrative and Comparative Biology*, *49*(6), 727–729. https://doi.org/10.1093/ICB/ICP106

Dejeux, L., Saclier, N., Tariel-Adam, J., Hoareau, M., Lefébure, T., Konecny, L., Plénet, S., & Luquet, E. (2025). Transcriptomic basis of within- and trans-generational predator-induced plasticity in the freshwater snail Physa acuta. *Heredity*, *134*(7), 439–449. https://doi.org/10.1038/S41437-025-00775-9;TECHMETA

DeWitt, T. J., Sih, A., & Wilson, D. S. (1998). Costs and limits of phenotypic plasticity. *Trends in Ecology and Evolution*, *13*(2), 77–81. https://doi.org/10.1016/S0169-5347(97)01274-3

Dias, B. G., & Ressler, K. J. (2014). Parental olfactory experience influences behavior and neural structure in subsequent generations. *Nature Neuroscience*, *17*(1), 89–96. https://doi.org/10.1038/NN.3594

Dieckmann, U., Heino, M., & Parvinen, K. (2006). The adaptive dynamics of function-valued traits. *Journal of Theoretical Biology*, *241*(2), 370–389. https://doi.org/10.1016/j.jtbi.2005.12.002

Donelson, J. M., Salinas, S., Munday, P. L., & Shama, L. N. S. (2018). Transgenerational plasticity and climate change experiments: Where do we go from here? *Global Change Biology*, *24*(1), 13–34. https://doi.org/10.1111/GCB.13903

Dzialowski, A. R., Lennon, J. T., O'Brien, W. J., & Smith, V. H. (2003). Predator-induced phenotypic plasticity in the exotic cladoceran Daphnia lumholtzi. *Freshwater Biology*, *48*(9), 1593–1602. https://doi.org/10.1046/J.1365-2427.2003.01111.X

Engel, K., & Tollrian, R. (2009). Inducible defences as key adaptations for the successful
invasion of Daphnia lumholtzi in North America? *Proceedings of the Royal Society B:*
Biological Sciences, *276*(1663), 1865. https://doi.org/10.1098/RSPB.2008.1861
Feinberg, A. P., & Irizarry, R. A. (2010). Stochastic epigenetic variation as a driving force of
development, evolutionary adaptation, and disease. *Proceedings of the National Academy of*
Sciences, *107*(suppl_1), 1757–1764. https://doi.org/10.1073/PNAS.0906183107
Feiner, N., Radersma, R., Vasquez, L., Ringnér, M., Nystedt, B., Raine, A., Tobi, E. W.,
Heijmans, B. T., & Uller, T. (2022). Environmentally induced DNA methylation is
inherited across generations in an aquatic keystone species. *IScience*, *25*(5), 104303.
https://doi.org/10.1016/J.ISCI.2022.104303
Forsman, A. (2015). Rethinking phenotypic plasticity and its consequences for individuals,
populations and species. *Heredity*, *115*(4), 276–284. https://doi.org/10.1038/HDY.2014.92
Freeberg, T. M., & Lucas, J. R. (2009). Pseudoreplication is (still) a problem. *Journal of*
Comparative Psychology (Washington, D.C. : 1983), *123*(4), 450–451.
https://doi.org/10.1037/A0017031
Furrow, R. E., & Feldman, M. W. (2014). Genetic variation and the evolution of epigenetic
regulation. *Evolution; International Journal of Organic Evolution*, *68*(3), 673–683.
https://doi.org/10.1111/EVO.12225
Gabry, J., Goodrich, B., Lysy, M., & JOhnson, A. (2025). *rstantools: Tools for Developing R*
Packages Interfacing with 'Stan (2.5.0). https://mc-stan.org/rstantools/
Gapp, K., Jawaid, A., Sarkies, P., Bohacek, J., Pelczar, P., Prados, J., Farinelli, L., Miska, E., &
Mansuy, I. M. (2014). Implication of sperm RNAs in transgenerational inheritance of the
effects of early trauma in mice. *Nature Neuroscience*, *17*(5), 667–669.
https://doi.org/10.1038/NN.3695
Geerts, A. N., Vanoverbeke, J., Vanschoenwinkel, B., Van Doorslaer, W., Feuchtmayr, H.,
Atkinson, D., Moss, B., Davidson, T. A., Sayer, C. D., & De Meester, L. (2015). Rapid
evolution of thermal tolerance in the water flea Daphnia. *Nature Climate Change*, *5*(7),
665–668. https://doi.org/10.1038/NCLIMATE2628;SUBJMETA
Geoghegan, J. L., & Spencer, H. G. (2012). Population-epigenetic models of selection.
Theoretical Population Biology, *81*(3), 232–242. https://doi.org/10.1016/j.tpb.2011.08.001
Ghalambor, C. K., McKay, J. K., Carroll, S. P., & Reznick, D. N. (2007). Adaptive versus non-
adaptive phenotypic plasticity and the potential for contemporary adaptation in new
environments. *Functional Ecology*, *21*(3), 394–407. https://doi.org/10.1111/J.1365-
2435.2007.01283.X
Graeve, A., Janßen, M., Villalba de la Pena, M., Tollrian, R., & Weiss, L. C. (2021). Higher,
Faster, Better: Maternal Effects Shorten Time Lags and Increase Morphological Defenses in
Daphnia lumholtzi Offspring Generations. *Frontiers in Ecology and Evolution*, *9*, 324.
https://doi.org/10.3389/FEVO.2021.637421/BIBTEX
Grishkevich, V., & Yanai, I. (2013). The genomic determinants of genotype × environment
interactions in gene expression. *Trends in Genetics*, *29*(8), 479–487.
https://doi.org/10.1016/j.tig.2013.05.006
Grosenbach, G. (2021). *A High Contrast Theme for ggplot - Topfunky Blog*.
https://topfunky.com/2021/ggplot-high-contrast-theme-gghighcontrast/
Gustafsson, S., & Hansson, L. A. (2004). Development of tolerance against toxic cyanobacteria
in Daphnia. *Aquatic Ecology*, *38*(1), 37–44.
https://doi.org/10.1023/B:AECO.0000020985.47348.5E/METRICS

Hales, N. R., Schield, D. R., Andrew, A. L., Card, D. C., Walsh, M. R., & Castoe, T. A. (2017).
Contrasting gene expression programs correspond with predator-induced phenotypic
plasticity within and across generations in Daphnia. *Molecular Ecology*, *26*(19), 5003–
5015. https://doi.org/10.1111/mec.14213

Harris, K. D. M., Bartlett, N. J., & Lloyd, V. K. (2012a). Daphnia as an Emerging Epigenetic
Model Organism. *Genetics Research International*, *2012*, 1–8.
https://doi.org/10.1155/2012/147892

Harris, K. D. M., Bartlett, N. J., & Lloyd, V. K. (2012b). Daphnia as an Emerging Epigenetic
Model Organism. *Genetics Research International*, *2012*, 1–8.
https://doi.org/10.1155/2012/147892

Heard, E., & Martienssen, R. A. (2014). Transgenerational epigenetic inheritance: Myths and
mechanisms. *Cell*, *157*(1), 95–109. https://doi.org/10.1016/j.cell.2014.02.045

Hearn, J., Pearson, M., Blaxter, M., Wilson, P. J., & Little, T. J. (2019). Genome-wide
methylation is modified by caloric restriction in Daphnia magna. *BMC Genomics*, *20*(1).
https://doi.org/10.1186/s12864-019-5578-4

Hellmann, J. K., Bukhari, S. A., Deno, J., & Bell, A. M. (2020). Sex-specific transgenerational
plasticity I: Maternal and paternal effects on sons and daughters. *BioRxiv*, 763862.
https://doi.org/10.1101/763862

Hendry, A. P. (2016). Key Questions on the Role of Phenotypic Plasticity in Eco-Evolutionary
Dynamics. *The Journal of Heredity*, *107*(1), 25–41.
https://doi.org/10.1093/JHERED/ESV060

Herman, J. J., & Sultan, S. E. (2011). Adaptive transgenerational plasticity in plants: Case
studies, mechanisms, and implications for natural populations. *Frontiers in Plant Science*,
2(DEC), 18242. https://doi.org/10.3389/FPLS.2011.00102/BIBTEX

Ho, D. H., & Burggren, W. W. (2010). Epigenetics and transgenerational transfer: a
physiological perspective. *The Journal of Experimental Biology*, *213*(1), 3–16.
https://doi.org/10.1242/JEB.019752

Hope, R. M. (2022). *Rmisc: Ryan Miscellaneous*. https://cran.r-
project.org/web/packages/Rmisc/index.html

Horstmann, M., Tollrian, R., & Weiss, L. C. (2021). Thwarting predators? A three-dimensional
perspective of morphological alterations in the freshwater crustacean Daphnia. *PLOS ONE*,
16(7), e0254263. https://doi.org/10.1371/JOURNAL.PONE.0254263

Hurlbert, S. H. (1984). Pseudoreplication and the Design of Ecological Field Experiments.
Ecological Monographs, *54*(2), 187–211. https://doi.org/10.2307/1942661

Jablonka, Eva., & Lamb, M. J. . (20062). *Evolution in four dimensions : genetic, epigenetic,*
behavioral, and symbolic variation in the history of life. MIT Press.

Jeremias, G., Barbosa, J., Marques, S. M., De Schamphelaere, K. A. C., Van Nieuwerburgh, F.,
Deforce, D., Gonçalves, F. J. M., Pereira, J. L., & Asselman, J. (2018). Transgenerational
Inheritance of DNA Hypomethylation in Daphnia magna in Response to Salinity Stress.
Environmental Science & Technology, *52*(17), 10114–10123.
https://doi.org/10.1021/ACS.EST.8B03225

Kaneshiro, K. R., Rechtsteiner, A., & Strome, S. (2019). Sperm-inherited H3K27me3 impacts
offspring transcription and development in C. elegans. *Nature Communications*, *10*(1), 1–9.
https://doi.org/10.1038/S41467-019-09141-W;TECHMETA

Kilham, S. S., Kreeger, D. A., Lynn, S. G., Goulden, C. E., & Herrera, L. (1998). COMBO: a
defined freshwater culture medium for algae and zooplankton. *Hydrobiologia 1998 377:1*,
377(1), 147–159. https://doi.org/10.1023/A:1003231628456
Kingsolver, J. G., Hoekstra, H. E., Hoekstra, J. M., Berrigan, D., Vignieri, S. N., Hill, C. E.,
Hoang, A., Gibert, P., & Beerli, P. (2001). The strength of phenotypic selection in natural
populations. *The American Naturalist*, *157*(3), 245–261. https://doi.org/10.1086/319193
Klosin, A., Casas, E., Hidalgo-Carcedo, C., Vavouri, T., & Lehner, B. (2017). Transgenerational
transmission of environmental information in C. elegans. *Science (New York, N.Y.)*,
356(6335), 320–323. https://doi.org/10.1126/SCIENCE.AAH6412
Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in
Linear Mixed Effects Models. *Journal of Statistical Software*, *82*(13), 1–26.
https://doi.org/10.18637/JSS.V082.I13
Laforsch, C., & Tollrian, R. (2004). Embryological aspects of inducible morphological defenses
in Daphnia. *Journal of Morphology*, *262*(3), 701–707. https://doi.org/10.1002/JMOR.10270
Laland, K. N., Uller, T., Feldman, M. W., Sterelny, K., Müller, G. B., Moczek, A., Jablonka, E.,
& Odling-Smee, J. (2015). The extended evolutionary synthesis: its structure, assumptions
and predictions. *Proceedings of the Royal Society B: Biological Sciences*, *282*(1813).
https://doi.org/10.1098/RSPB.2015.1019
Lande, R. (2009). Adaptation to an extraordinary environment by evolution of phenotypic
plasticity and genetic assimilation. *Journal of Evolutionary Biology*, *22*(7), 1435–1446.
https://doi.org/10.1111/J.1420-9101.2009.01754.X
Landy, J. A., Muh, S., Parajuli, B., Munch, S. B., & Walsh, M. R. (2024). Multi-generation
exposure to a novel environmental signal shapes the expression of plasticity in offspring in
resurrected Daphnia. *Biological Journal of the Linnean Society*, *143*(4).
https://doi.org/10.1093/BIOLINNEAN/BLAE091
Levis, N. A., & Pfennig, D. W. (2019). Plasticity-led evolution: evaluating the key prediction of
frequency-dependent adaptation. *Proceedings. Biological Sciences*, *286*(1897).
https://doi.org/10.1098/RSPB.2018.2754
Li, E., & Zhang, Y. (2014). DNA methylation in mammals. *Cold Spring Harbor Perspectives in
Biology*, *6*(5). https://doi.org/10.1101/CSHPERSPECT.A019133
Lüdecke, D., Ben-Shachar, M., Patil, I., Waggoner, P., & Makowski, D. (2021). performance:
An R Package for Assessment, Comparison and Testing of Statistical Models. *Journal of
Open Source Software*, *6*(60), 3139. https://doi.org/10.21105/JOSS.03139
Müller, M. F., Colomer, J., & Serra, T. (2018). Temperature-driven response reversibility and
short-term quasi-acclimation of Daphnia magna. *PLoS ONE*, *13*(12).
https://doi.org/10.1371/JOURNAL.PONE.0209705
Mulligan, C. J., Quinn, E. B., Hamadmad, D., Dutton, C. L., Nevell, L., Binder, A. M., Panter-
Brick, C., & Dajani, R. (2025). Epigenetic signatures of intergenerational exposure to
violence in three generations of Syrian refugees. *Scientific Reports*, *15*(1), 1–13.
https://doi.org/10.1038/S41598-025-89818-Z;SUBJMETA
Nagano, M., & Yoshida, T. (2020). Size-selective predation accounts for intra- and inter-specific
variation of inducible morphological defense of Daphnia. *Ecosphere*, *11*(7), e03192.
https://doi.org/10.1002/ECS2.3192
Nettle, D., & Bateson, M. (2015). Adaptive developmental plasticity: what is it, how can we
recognize it and when can it evolve? *Proceedings of the Royal Society B: Biological
Sciences*, *282*(1812). https://doi.org/10.1098/RSPB.2015.1005

Niederhuth, C. E., Bewick, A. J., Ji, L., Alabady, M. S., Kim, K. Do, Li, Q., Rohr, N. A.,
    Rambani, A., Burke, J. M., Udall, J. A., Egesi, C., Schmutz, J., Grimwood, J., Jackson, S.
    A., Springer, N. M., & Schmitz, R. J. (2016). Widespread natural variation of DNA
    methylation within angiosperms. *Genome Biology 2016 17:1*, *17*(1), 1–19.
    https://doi.org/10.1186/S13059-016-1059-0

Norouzitallab, P., Baruah, K., Vandegehuchte, M., Van Stappen, G., Catania, F., Vanden
    Bussche, J., Vanhaecke, L., Sorgeloos, P., & Bossier, P. (2014). Environmental heat stress
    induces epigenetic transgenerational inheritance of robustness in parthenogenetic Artemia
    model. *FASEB Journal : Official Publication of the Federation of American Societies for
    Experimental Biology*, *28*(8), 3552–3563. https://doi.org/10.1096/FJ.14-252049

Pedersen, T. L. (2025). *The Composer of Plots [R package patchwork version 1.3.2]*.
    https://CRAN.R-project.org/package=patchwork

Petrusek, A., Tollrian, R., Schwenk, K., Haas, A., & Laforsch, C. (2009). A "crown of thorns" is
    an inducible defense that protects Daphnia against an ancient predator. *Proceedings of the
    National Academy of Sciences*, *106*(7), 2248–2252.
    https://doi.org/10.1073/PNAS.0808075106

Pfennig, D. W., Wund, M. A., Snell-Rood, E. C., Cruickshank, T., Schlichting, C. D., &
    Moczek, A. P. (2010). Phenotypic plasticity's impacts on diversification and speciation.
    Trends in Ecology and Evolution, *25*(8), 459–467.
    https://doi.org/10.1016/j.tree.2010.05.006

Pigliucci, M., Murren, C. J., & Schlichting, C. D. (2006). Phenotypic plasticity and evolution by
    genetic assimilation. *Journal of Experimental Biology*, *209*(12), 2362–2367.
    https://doi.org/10.1242/JEB.02070

Pigliucci, & Müller. (2010). *Evolution, the Extended Synthesis* .

Postit Team. (2025). *RStudio: Integrated Development Environment for R*. Postit Software.
    http://www.posit.co/.

Proulx, S. R., & Teotónio, H. (2017). What kind of maternal effects can be selected for in
    fluctuating environments? *American Naturalist*, *189*(6), E118–E137.
    https://doi.org/10.5061/DRYAD.58416

R Core Team. (2021). R: A language and environment for statisticalcomputing. *R Foundation for
    Statistical Computing*.

Radchuk, V., Reed, T., Teplitsky, C., van de Pol, M., Charmantier, A., Hassall, C., Adamík, P.,
    Adriaensen, F., Ahola, M. P., Arcese, P., Miguel Avilés, J., Balbontin, J., Berg, K. S.,
    Borras, A., Burthe, S., Clobert, J., Dehnhard, N., de Lope, F., Dhondt, A. A., … Kramer-
    Schadt, S. (2019). Adaptive responses of animals to climate change are most likely
    insufficient. *Nature Communications*, *10*(1), 1–14. https://doi.org/10.1038/S41467-019-
    10924-4;TECHMETA

Rechavi, O., Houri-Ze'Evi, L., Anava, S., Goh, W. S. S., Kerk, S. Y., Hannon, G. J., & Hobert,
    O. (2014). Starvation-induced transgenerational inheritance of small RNAs in C. elegans.
    Cell, *158*(2), 277–287. https://doi.org/10.1016/j.cell.2014.06.020

Robichaud, N. F., Sassine, J., Beaton, M. J., & Lloyd, V. K. (2012). The Epigenetic Repertoire
    of Daphnia magna Includes Modified Histones . *Genetics Research International*, *2012*, 1–
    7. https://doi.org/10.1155/2012/174860

Romero-Mujalli, D., Fuchs, L. I. R., Haase, M., Hildebrandt, J. P., Weissing, F. J., & Revilla, T.
    A. (2024). Emergence of phenotypic plasticity through epigenetic mechanisms. *Evolution
    Letters*, *8*(4), 561–574. https://doi.org/10.1093/EVLETT/QRAE012

Scheiner, S. M. (1993). Genetics and evolution of phenotypic plasticity. *Annual Review of Ecology and Systematics*, *24*(1), 35–68. https://doi.org/10.1146/ANNUREV.ES.24.110193.000343/CITE/REFWORKS

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. *Nature Methods 2012 9:7*, *9*(7), 676–682. https://doi.org/10.1038/nmeth.2019

Schlichting, C. D., & Levin, D. A. (1986). Phenotypic plasticity: an evolving plant character. *Biological Journal of the Linnean Society*, *29*(1), 37–47. https://doi.org/10.1111/J.1095-8312.1986.TB01769.X

Schwarzenberger, A., & Martin-Creuzburg, D. (2021). Daphnia's Adaptive Molecular Responses to the Cyanobacterial Neurotoxin Anatoxin-α Are Maternally Transferred. *Toxins*, *13*(5). https://doi.org/10.3390/TOXINS13050326

Scoville, A. G., & Pfrender, M. E. (2010). Phenotypic plasticity facilitates recurrent rapid adaptation to introduced predators. *Proceedings of the National Academy of Sciences of the United States of America*, *107*(9), 4260–4263. https://doi.org/10.1073/PNAS.0912748107

Searle, S. R., Speed, F. M., & Milliken, G. A. (2025). Estimated Marginal Means, aka Least-Squares Means [R package emmeans version 1.11.2-8]. *American Statistician*, *34*(4), 216–221. https://doi.org/10.1080/00031305.1980.10483031

Seda, J., & Petrusek, A. (2011). Daphnia as a model organism in limnology and aquatic biology: some aspects of its reproduction and development. *Journal of Limnology*, *70*(2), 335–386. https://doi.org/10.4081/jlimnol.2011.335

Shahmohamadloo, R. S., Fryxell, J. M., & Rudman, S. M. (2025). Transgenerational epigenetic inheritance increases trait variation but is not adaptive. *Evolution*, *79*(6), 1033–1043. https://doi.org/10.1093/EVOLUT/QPAF050

Simons, A. M. (2009). Fluctuating natural selection accounts for the evolution of diversification bet hedging. *Proceedings of the Royal Society B: Biological Sciences*, *276*(1664), 1987–1992. https://doi.org/10.1098/RSPB.2008.1920

Sousa, F. D. R., Palaoro, A. V., Elmoor-Loureiro, L. M. A., & Kotov, A. A. (2017). Predicting the invasive potential of the cladoceran Daphnia lumholtzi Sars, 1885 (Crustacea: Cladocera: Daphniidae) in the Neotropics: Are generalists threatened and relicts protected by their life-history traits? *Journal of Limnology*, *76*(2), 272–280. https://doi.org/10.4081/JLIMNOL.2016.1571

St-Cyr, S., & McGowan, P. O. (2015). Programming of stress-related behavior and epigenetic neural gene regulation in mice offspring through maternal exposure to predator odor. *Frontiers in Behavioral Neuroscience*, *9*(JUNE), 145810. https://doi.org/10.3389/FNBEH.2015.00145/BIBTEX

Szűcs, G., Sója, A., Péter, M., Sárközy, M., Bruszel, B., Siska, A., Földesi, I., Szabó, Z., Janáky, T., Vígh, L., Balogh, G., & Csont, T. (2019). Prediabetes Induced by Fructose-Enriched Diet Influences Cardiac Lipidome and Proteome and Leads to Deterioration of Cardiac Function prior to the Development of Excessive Oxidative Stress and Cell Damage. *Oxidative Medicine and Cellular Longevity*, *2019*. https://doi.org/10.1155/2019/3218275

Tariel, J., Plénet, S., & Luquet, É. (2020). Transgenerational plasticity of inducible defences: Combined effects of grand-parental, parental and current environments. *Ecology and Evolution*, *10*(5), 2367–2376. https://doi.org/10.1002/ECE3.6046

Trijau, M., Asselman, J., Armant, O., Adam-Guillermin, C., De Schamphelaere, K. A. C., & Alonzo, F. (2018). Transgenerational DNA Methylation Changes in Daphnia magna Exposed to Chronic γ Irradiation. *Environmental Science and Technology*, *52*(7), 4331–4339. https://doi.org/10.1021/acs.est.7b05695

Tufto, J. (2015). Genetic evolution, plasticity, and bet-hedging as adaptive responses to temporally autocorrelated fluctuating selection: A quantitative genetic model. *Evolution*, *69*(8), 2034–2049. https://doi.org/10.1111/EVO.12716

Uller, T., Moczek, A. P., Watson, R. A., Brakefield, P. M., & Laland, K. N. (2018). Developmental Bias and Evolution: A Regulatory Network Perspective. *Genetics*, *209*(4), 949–966. https://doi.org/10.1534/GENETICS.118.300995

Uller, T., Nakagawa, S., & English, S. (2013). Weak evidence for anticipatory parental effects in plants and animals. *Journal of Evolutionary Biology*, *26*(10), 2161–2170. https://doi.org/10.1111/JEB.12212

Van Buskirk, J., & Steiner, U. K. (2009). The fitness costs of developmental canalization and plasticity. *Journal of Evolutionary Biology*, *22*(4), 852–860. https://doi.org/10.1111/J.1420-9101.2009.01685.X

Vandegehuchte, M. B., De Coninck, D., Vandenbrouck, T., De Coen, W. M., & Janssen, C. R. (2010). Gene transcription profiles, global DNA methylation and potential transgenerational epigenetic effects related to Zn exposure history in Daphnia magna. *Environmental Pollution*, *158*(10), 3323–3329. https://doi.org/10.1016/j.envpol.2010.07.023

Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing*, *27*(5), 1413–1432. https://doi.org/10.1007/S11222-016-9696-4

Via, S., Gomulkiewicz, R., De Jong, G., Scheiner, S. M., Schlichting, C. D., & Van Tienderen, P. H. (1995). Adaptive phenotypic plasticity: consensus and controversy. *Trends in Ecology & Evolution*, *10*(5), 212–217. https://doi.org/10.1016/S0169-5347(00)89061-8

Walsh, M. R., Cooley, F., Biles, K., & Munch, S. B. (2014). Predator-induced phenotypic plasticity within- and across-generations: A challenge for theory? *Proceedings of the Royal Society B: Biological Sciences*, *282*(1798). https://doi.org/10.1098/rspb.2014.2205

Walsh, M. R., Whittington, D., & Funkhouser, C. (2014). Thermal Transgenerational Plasticity in Natural Populations of Daphnia. *Integrative and Comparative Biology*, *54*(5), 822–829. https://doi.org/10.1093/ICB/ICU078

Werner, E. E., & Gilliam, J. F. (1984). The ontogenetic niche and species interactions in size-structured populations. *Annual Review of Ecology and Systematics. Vol. 15*, *15*(Volume 15,), 393–425. https://doi.org/10.1146/ANNUREV.ES.15.110184.002141/CITE/REFWORKS

Werner, M. S., Loschko, T., King, T., Reich, S., Theska, T., Franz-Wachtel, M., Macek, B., & Sommer, R. J. (2023). Histone 4 lysine 5/12 acetylation enables developmental plasticity of Pristionchus mouth form. *Nature Communications 2023 14:1*, *14*(1), 1–14. https://doi.org/10.1038/s41467-023-37734-z

West-Eberhard, M. J. (2003). Developmental Plasticity and Evolution. *Developmental Plasticity and Evolution*. https://doi.org/10.1093/OSO/9780195122343.001.0001

West-Eberhard, M. J. (2005). Developmental plasticity and the origin of species differences. *Proceedings of the National Academy of Sciences of the United States of America*, *102 Suppl 1*(Suppl 1), 6543–6549. https://doi.org/10.1073/PNAS.0501844102

West-Eberhard, M. Jane. (2020). *Developmental plasticity and evolution*.

Wickham, H. (2016). ggplot2. *Ggplot2*. https://doi.org/10.1007/978-0-387-98141-3

Wickham, H., François, R., Henry, L., & Müller, K. (2022). *dplyr: A Grammar of Data*
Manipulation. https://github.com/tidyverse/dplyr

Woltereck, R. (1913). Weitere experimentelle untersuchungen über Artänderung, speziell über
das Wesen quantitativer Artunterschiede bei Daphniden. *Zeitschrift Für Induktive*
Abstammungs- Und Vererbungslehre, *9*(1), 146–146. https://doi.org/10.1007/BF01876686

Work, K. A., & Gophen, M. (1999). Factors which affect the abundance of an invasive
cladoceran, Daphnia lumholtzi, in U.S. reservoirs. *Freshwater Biology*, *42*(1), 1–10.
https://doi.org/10.1046/J.1365-2427.1999.00449.X

Yampolsky, L. Y., Zeng, E., Lopez, J., Williams, P. J., Dick, K. B., Colbourne, J. K., & Pfrender,
M. E. (2014). Functional genomics of acclimation and adaptation in response to thermal
stress in Daphnia. *BMC Genomics*, *15*(1), 1–12. https://doi.org/10.1186/1471-2164-15-
859/TABLES/2

Yaniv, M. (2014). Chromatin remodeling: From transcription to cancer. In *Cancer Genetics*
(Vol. 207, Issue 9, pp. 352–357). Elsevier Inc.
https://doi.org/10.1016/j.cancergen.2014.03.006

1099