TITLE: Non-genetic inheritance of induced defense morphologies across multiple unexposed

generations of *Daphnia lumholtzi*

RUNNING TITLE: Transgenerational Inheritance in *Daphnia lumholtzi*

Authors:

Shannon N. Snyder[1,*],

Walker C. Meyer[1],

Shanie L. Jorgenson[1]

Ethan B. Contreras[1]

Tea Bland[1]

William A. Cresko[1,2,*]

Affiliations:

[1] Institute of Ecology & Evolution, University of Oregon, Eugene, OR, 97403, USA

[2] Department of Bioengineering, Knight Campus for Accelerating Scientific Impact, Eugene,

OR, 97403, USA

*Corresponding Authors (ssnyder3@uoregon.edu)

<u>ORCID</u>:

SNS: 0000-0001-7049-2950

WCM: 0009-0003-7225-4296

SLJ: 0009-0007-0144-2504

EBC: 0009-0001-1019-8263

TB: 0009-0007-1456-0569

WAC: 0000-0002-3496-8074

<u>**Keywords**</u>: Transgenerational inheritance, Phenotypic plasticity, Transgenerational Plasticity,

Non-genetic inheritance, Epigenetic Inheritance, Armor Morphology, Bayesian hierarchical

modeling, Inducible defenses

<u>**Data availability statement**</u>: The raw data and full analysis scripts that support the findings of

this study are openly available in Markdown on GitHub

(https://github.com/shannonsnyder/daphnia-transgenerational-plasticity).

<u>**Author contributions**</u>: **Conceptualization and Methodology:** SNS and WAC conceived and

designed the study. **Investigation:** SNS, WCM, and SLJ reared the animals and collected the

imaging data. **Data Collection and Curation:** SNS, EBC and TB curated the morphological

measurements. **Formal Analysis:** SNS and WAC analyzed the data. **Visualization:** SNS created

the data visualizations. **Writing – Original Draft:** SNS wrote the manuscript. **Writing –**

Review & Editing: WAC provided revisions. All authors approved the final manuscript.

<u>**Funding**</u>: This work was funded by the National Science Foundation Grant OPP-2015301 (to

WAC), University of Oregon Research Excellence funds (WAC), University of Oregon Office of

the Vice President for Research and Innovation (OVPRI) seed funding (SNS and WAC). WCM

received funding and mentorship from the Knight Campus Undergraduate Scholars program at

the University of Oregon. SLJ and EBC received funding from the University of Oregon

Graduate Students in Ecology and Evolution Student Organization. SLJ received funding from

the McNair Scholar Program. EBC and WCM received funding from University of Oregon

OVRPI.

Conflict of Interest: The authors declare no conflict of interest.

Acknowledgements: We would like to thank M. Lynch for his contribution of the *Daphnia*

lumholtzi (SAG) utilized for this study. We would also like to thank Mark Currey for his

assistance with stickleback husbandry and the Cresko lab for valuable feedback. Finally, we

thank Emily Willams for collection and maintenance of *Daphnia* prior to their arrival in the

Cresko lab.

59
60

Teaser Text

Can the ghosts of environments past shape organismal phenotypes of future generations? In an experiment with clonal *Daphnia lumholtzi*, we reveal that predator-induced changes in body shape in one generation can persist across several unexposed generations, even when the predator signal is long gone. Exposing only the first generation to predator cues, we tracked morphological shifts through five generations of genetically identical individuals. We found that these induced plastic phenotypes persisted to the F4 generation before fading in F5. This discovery supports the importance of non-genetic inheritance and underscores the powerful interplay between ancestral and current environments in shaping organismal form and function.

Abstract

Environmental variation can induce phenotypic changes through adaptation of

populations and acclimation of individuals. While genetic adaptation creates persistent change

within a population via allele frequency shifts, reversible plastic phenotypes can be inherited

through non-genetic mechanisms. However, most transgenerational plasticity studies examine

generations where direct embryonic or germline exposure to environmental cues cannot be

excluded. Empirical evidence for persistence definitively beyond the critical threshold for

distinguishing true transgenerational plasticity from in utero exposure remains scarce,

particularly for vertebrate predator-induced defenses.

We measured phenotypic effects of vertebrate predator exposure in the clonally

reproducing water flea, *Daphnia lumholtzi*, isolating environmental effects from genetic

variation. We exposed the F0 generation to fish conditioned media, then measured

morphological defenses in definitively unexposed F3, F4, and F5 generations characterizing the

temporal dynamics of non-genetic inheritance. Predator-induced morphologies persisted through

F4 before receding in F5, demonstrating that non-genetic effects extend well beyond the

embryonically exposed F1 and germ cell exposed F2 generations. This provides rare empirical

evidence for transgenerational plasticity lasting through F4, without confounding genetic

variation. We also examined ontogenetic patterns of somatic and defensive trait development,

revealing trait-specific temporal dynamics in transgenerational effect expression and decay.

These results highlight how current and ancestral environments interact to determine

phenotypic variation across generations and underscores the ecological significance of non-

genetic inheritance in natural populations, particularly for understanding population responses to

environmental change and predator reintroduction. Characterizing molecular mechanisms

underlying transgenerational phenotypic plasticity remains critical for predicting persistence and

ecological consequences of non-genetic inheritance.

Introduction

Organisms respond to environmental variation via changes in morphology, behavior, and

life history traits (Donelson et al., 2018; Szűcs et al., 2019; West-Eberhard, 2005). These

phenotypic responses may arise from shifts in allele frequencies across generations or from non-

genetic mechanisms acting within individuals (Levis & Pfennig, 2019; Uller et al., 2018; West-

Eberhard, 2020). Increased phenotypic variation can enhance the probability that some offspring

will express trait values suited to novel environmental conditions (Botero et al., 2015; Simons,

2009; Tufto, 2015). However, this variation also increases the likelihood of mismatched

phenotypes, reducing individual fitness. While genetic adaptation can improve population-level

fitness over time, subsequent environmental shifts may render previously adaptive traits

maladaptive (Lande, 2009; Radchuk et al., 2019).

Regularly experienced environmental variation presents a particular challenge for

evolving populations because it can lead to persistent mismatch of organisms and the

environment, and thus ongoing genetic load (Ghalambor et al., 2007; Via et al., 1995). An

alternative biological strategy for dealing with such variable conditions is the evolution of plastic

traits (Hendry, 2016) in which phenotypic expression covaries with environmental variation

(Bradshaw, 1965; Schlichting & Levin, 1986). Phenotypic plasticity allows a single genotype to

consistently produce different phenotypes depending on the environment, without requiring

genetic change (Pfennig et al., 2010; Pigliucci et al., 2006). Plasticity includes developmental

plasticity (Nettle & Bateson, 2015; Uller et al., 2018), where early-life cues shape long-term

traits, and acclimatory plasticity (Angilletta, 2009; Beaman et al., 2016), involving reversible

physiological changes in response to short-term fluctuations, illustrating a range of temporal

scales over which environmentally induced phenotypic changes can occur.

Plasticity within a generation has been well documented across numerous taxa and

environmental conditions (Agrawal et al., 1999; Corl et al., 2018; Forsman, 2015). Although

plasticity itself is often an adaptive phenotype that has been crafted by natural selection, the trait

distributions produced by plasticity within one generation have historically been thought to be

ephemeral with little consequence for the phenotypes of future generations (West-Eberhard,

2020). However, modern analyses have documented that variation in plastic traits is mediated by

changes in gene expression (Hales et al., 2017) and is hypothesized to involve epigenetic

regulation (St-Cyr & McGowan, 2015), raising the possibility that induced plastic phenotypes

may be inherited across generations (Harris et al., 2012a; Romero-Mujalli et al., 2024). This

potential for non-genetic inheritance exemplifies the need to integrate genetic and non-genetic

mechanisms of information transmission across generations (inclusive heritability, Danchin &

Wagner, 2010) in evolutionary biology.

Transgenerational inheritance of induced plastic traits from parent to offspring is

expected to evolve when the parental environment covaries with the offspring environment

(Agrawal et al., 1999). Although induced plastic phenotypes should return to the baseline state in

the absence of the inducing cues over generations, the rate at which this occurs is hypothesized

to be determined by the frequency of which the environment shifts (Furrow & Feldman, 2014;

Proulx & Teotónio, 2017). The evolution of inherited phenotypic plasticity therefore provides a

mechanism for organisms and their offspring to respond adaptively to changing environments

without significant genetic load (Ghalambor et al., 2007).

Studies have documented that induced phenotypes can be inherited (Bell & Hellmann,

2019; MacLeod et al., 2022; Moore et al., 2019; Neylan et al., 2025; Tariel et al., 2020; Yin et

al., 2019), but the prevalence and duration of this phenomenon remain unclear, especially under

142 fluctuating environmental conditions (Norouzitallab et al., 2014). A particular challenge is

143 distinguishing true transgenerational inheritance from extended maternal or embryonic effects.

144 Because embryos often exist within the maternal body or egg, they may be directly exposed to

145 the environmental cue (Agrelius & Dudycha, 2025; Skinner, 2008). Furthermore, these embryos

146 may have developing germ cells that can be similarly exposed, necessitating the examination of

147 generations sufficiently removed from the initial exposure to rule out direct effects. The number

148 of generations required to differentiate maternal effects from true transgenerational inheritance

149 varies by organism depending on developmental timing and reproductive mode, but empirical

150 data demonstrating persistence beyond the last generation with possible germline exposure

151 remains limited, partially because embryonic and germ line exposure is often conflated with true

152 transgenerational plasticity (Agrelius & Dudycha, 2025; Skinner, 2008). In fact, a 2020 meta-

153 analysis of transgenerational phenotypic plasticity (TGP) in predator-prey interactions found that

154 only 13% of studies (seven total studies) extended beyond the F0 and F1 generations (Tariel et

155 al., 2020), with studies examining F3+ generations remaining notably rare. Despite these

156 advances, critical questions remain: How long do induced traits persist without ongoing cue

157 exposure? At what generation do transgenerational effects decay? And can we definitively

158 distinguish transgenerational plasticity from extended maternal effects?

159   Convincing examples of non-genetic inheritance that extend beyond temporal limits of

160 potential direct exposure include a handful of examinations in *Caenorhabditis elegans* (Greer et

161 al., 2011; S. W. Kim et al., 2013; Kishimoto et al., 2017; Klosin et al., 2017; Remy, 2010; M.

162 Wang et al., 2019), *Mus musculus* (Anway et al., 2005; Skinner et al., 2013; Van Steenwyk et al.,

163 2018), *Drosophila melanogaster* (Panacek et al., 2011), *Danio rerio* (Alfonso et al., 2019), *Ovis*

164   *aries* (Kizilaslan et al., 2025), and the keystone aquatic species, *Daphnia* (Shahmohamadloo et

165   al., 2025).

166   Often overlooked in such studies is the influence of genetic variation. Asexual

reproduction provides a particularly powerful system for studying non-genetic transgenerational

effects because it eliminates confounding variation due to recombination, genetic background,

and parental sex. Examples of such studies include a striking example in aphids, which examined

27 generations of the inheritance of predation induced phenotypes (Sentis et al., 2018), clonal

and hermaphroditic freshwater snail systems (Dejeux et al., 2025; Smithson et al., 2020; Tariel et

al., 2020), in which alterations in shells shape and thickness can be inherited across generations,

and hermaphroditic *C.elegans* (Baugh & Day, 2020). These examples remain relatively few,

leaving it unclear whether true transgenerational inheritance is rare or understudied.

175   The freshwater microcrustacean *Daphnia* is especially well suited for studying

transgenerational plasticity and addressing these questions (Ebert, 2022; Harris et al., 2012; Seda

& Petrusek, 2011; Yampolsky et al., 2014). The concept of plasticity was introduced in the early

1900s by Richard Woltereck**,** who demonstrated that head size in *Daphnia* varied with nutrient

availability, and proposed that environmentally induced traits could be inherited (Woltereck,

1913). *Daphnia's* parthenogenetic reproduction produces genetically identical offspring,

eliminating confounding variation due to recombination, genetic background, and parental sex

(Harris et al., 2012). *These ecoresponsive crustaceans* exhibit plastic responses to a range of

environmental stressors, including cyanobacteria (Dao et al., 2018; Gustafsson & Hansson, 2004;

Schwarzenberger & Martin-Creuzburg, 2021), temperature (Geerts et al., 2015; Landy et al.,

2024; Müller et al., 2018), and predation (Hebert & Grewe, 1985; Tollrian & Harvell, 1999),

affecting morphology (Horstmann et al., 2021), behavior (Beklioglu et al., 2006), and life history

traits (Adamczuk, 2020; Agrawal et al., 1999). Examples include inducible helmets, spines, and

neckteeth in response to predator cues (Horstmann et al., 2021; Laforsch & Tollrian, 2004).

Many of these traits persist across clonal generations, such as helmet length in *D. cucullata*

(Agrawal et al., 1999) and clutch size in *D. ambigua* (Walsh et al., 2014), suggesting a non-

genetic mechanism of inheritance.

The number of generations over which induced traits persist in *Daphnia* in the absence of

the original environmental cue is still being resolved. While phenotypic effects extending

through the F1-F2 generations (maternal effects) in *Daphnia* are well-documented, evidence for

persistence beyond F2 (which eliminates direct embryonic exposure; (Agrelius & Dudycha,

2025) remains limited. Most documented cases track only one or two generations post-exposure,

such as helmet length in *D. cucullata* (Agrawal et al., 1999; Laforsch & Tollrian, 2004), and age

of maturation in *D. ambigua* (Walsh et al., 2016). Many of these studies have predominantly

focused on invertebrate predation, such as *Chaoborus* larvae (Horstmann et al., 2022),

Bythotrephes longimanus (spiny water flea) (Landy et al., 2024), *Leptodora* (Baludo et al.,

2024), *Notonecta* (Horstmann et al., 2023; Weiss et al., 2015), and *Triops* (Pietrzak et al., 2020),

or focus on various toxicological stressors (Funke et al., 2024; J. Kim & Choi, 2023; Liao et al.,

2023; Liu et al., 2020; Maggio & Jenkins, 2022a, 2022b; Martins & Guilhermino, 2018a, 2018b;

Shahmohamadloo et al., 2025; Song et al., 2022; Vandegehuchte, Lemière, et al., 2010; J. Wang

et al., 2024). However, studies examining the transgenerational responses to vertebrate predation

beyond generations with possible germline exposure remain notably scarce (Tariel et al., 2020).

This gap is particularly important because vertebrate predators impose strong, and often

more temporally consistent selective pressure on prey populations (Batabyal, 2023; Salo et al.,

2010; Wooster, 1994), making transgenerational responses to fish predation critical for

understanding adaptive potential in natural systems, and relevant to predator reintroduction

efforts (Bracis & Wirsing, 2021). Molecular studies suggest long term effects are possible, with

differential gene expression persisting after predator cue removal (Hales et al., 2017), and

methylation differences between naive and unexposed individuals across various stressors

(Feiner et al., 2022; Hearn et al., 2019; Jeremias et al., 2018; Trijau et al., 2018). However, the

duration of phenotypic persistence and its evolutionary significance remain unclear (Bird, 2024;

Feinberg, 2007; Feinberg & Irizarry, 2010; Fitz-James & Cavalli, 2022; Ho & Burggren, 2010;

Klibaner-Schiff et al., 2024), requiring empirical evidence beyond generations with possible

germline exposure.

Daphnia lumholtzi develops elongated head and tailspine in response to fish predation

(Dzialowski et al., 2003), traits that effectively reduce predation risk and represent a clear

example of adaptive phenotypic plasticity (Engel et al., 2014). These distinctive spines are

thought to contribute to the species' invasive success in North America (Engel & Tollrian, 2009;

Sousa et al., 2017; Work & Gophen, 1999) where *D. lumholtzi* often displaces native North

American *Daphnia* populations (Johnson & Havel, 2001). Unlike *D. lumholtzi,* many North

American *Daphnia* species typically produce broad, helmet-like adornments rather than

elongated spines (Horstmann et al., 2021) exemplifying *D. lumholtzi's* unique defensive strategy.

These distinctive morphological defenses, combined with *Daphnia's* parthenogenetic

reproduction, make *D. lumholtzi* an ideal system for tracking transgenerational effects across

multiple generations without genetic confounds.

We measured predator induced morphological traits in *D. lumholtzi* across multiple

genetically identical generations following initial exposure, focusing on F3-F5 cohorts where

neither the individuals themselves nor their germ cells (as developing gametes in F1-F2) could

have experienced direct exposure to predator cues. Previous work in *D. lumholtzi* shows that

continued multigenerational maternal exposure can accelerate defensive trait development

(Graeve et al., 2021) and that cyanobacteria-induced life history effects persist for at least one

generation (Dao et al., 2018), but no study has examined the long-term persistence of vertebrate

predator induced morphology in *Daphnia lumholtzi*, particularly beyond the generations with

possible direct exposure. By focusing on F3-F5 generations, our observations document

transgenerational inheritance via necessarily non-genetic mechanisms rather than extended

effects of direct exposure. Additionally, deriving all experimental lineages from a single

founding mother of the Saguaro Lake *D. lumholtzi* clone eliminates confounding effects of

standing genetic variation, enabling us to fully isolate non-genetic inheritance.

We found that induced traits persisted through multiple unexposed generations, with

measurable effects gradually declining to undetectable levels in F5. We identify distinct

ontogenetic trajectories in defensive versus somatic traits across the Daphnia's lifespan. A

critical question is whether transgenerational morphological difference reflect inherited plasticity

versus size-selective mortality of non-induced phenotypes. To address this alternative

hypothesis, we explicitly test for evidence of trait-selective or size-selective mortality and report

overlapping survival trajectories between different sized defensive structures, indicating that

elongated spines are indeed inherited, rather than a selective artifact. These results demonstrate

that true transgenerational plasticity can be inherited across clonal generations, but exhibits

limited temporal persistence, with trait-specific patterns of decline that reflect the gradual loss of

ancestral environmental influence (Geoghegan & Spencer, 2012; Heard & Martienssen, 2014).

Methods

Experimental Overview (Figure 1). To explore the dynamics of predation induced phenotypes

within and across generations, we reared individual Daphnia with ancestral predation exposure in

the F0 generation, alongside unexposed *Daphnia lumholtzi* individuals for five generations

following initial predation exposure only in the predator F0 generation (total generations: F0-F5;

only F0 exposed). These individuals were the progeny of a single founding mother, whose

genetically identical daughters (F0) were randomly assigned to treatment or control to establish

replicates, and propagated for an additional four generations without subsequent exposure. This

design isolated environmental effects from confounding genetic variation while maintaining

independent biological replicates.

We focused morphological measurements on F0, F3, F4, and F5 generations. F0

established baseline induction effects, while F3-F5 represented generations with no direct

embryonic or germline exposure to predator cues - essential for distinguishing true

transgenerational inheritance from extended maternal effects (Agrelius & Dudycha, 2025;

Skinner, 2008). F1 and F2 generations were propagated but not measured, as phenotypic effects

in these cohorts are well-documented and can result from direct embryonic exposure. While this

design limits F0 sample size, our primary objective was characterizing persistence of effects in

F3-F5 individuals rather than recapitulating characterized F0-F2 responses.

Daphnia were individually grown in 50 mL beakers containing 30 mL of fish conditioned

medium or control medium (COMBO) (Kilham et al., 1998), enabling replication at the

individual level. Every other day, until death, during the F0, F3, F4 and F5 generations,

individual *D. lumholtzi* were imaged for morphological measurements using `ImageJ` (Schindelin

et al., 2012). This experimental framework also allowed us to examine survivorship (as animals

were imaged until they died) and life-long ontogenetic trajectories. On this same schedule the

media was exchanged, and algae was replenished with 85,000 cells/mL of a 1:1 mixture of

Scenedesmus obliquus and *Chlorella vulgaris*, hereafter referred to as the standard algae mixture.

Variation in sample size between treated and control individuals and across generations

reflects differential survival and reproductive success. Predator-exposed individuals showed

higher survival and fecundity in some generations, resulting in asymmetric sample sizes.

Bayesian hierarchical models can maintain robustness despite imbalance and small sample sizes.

See Figure 1 for experimental schematic and sample sizes; See Table 1 for expanded sample

sizes, including number of repeated measures observations on individuals.

This study focuses on a single genetic background to isolate non-genetic effects. Future

work examining multiple clones and populations will be necessary to assess the generalizability

of these transgenerational effects.

Clonal Genetic Background: *Daphnia Lumholtzi*. *Daphnia lumholtzi* was collected in the

summer of 2019 from Saguaro Lake, AZ, USA. We utilized one Saguaro Lake Daphnia, as our

primary objective is to minimize confounding genetic variation, rather than generalize a

population's ability to respond to predator cues. This species and genetic background were

specifically chosen for its ability to develop unique headspines and tailspines in response to

vertebrate predation and for their ecological significance as an invasive North American species,

capable of displacing native *Daphnia* species (Engel et al., 2014b; Engel & Tollrian, 2009; Sousa

et al., 2017; Work & Gophen, 1999).

Common Garden Rearing to Standardize Generational Exposures. After lake collection, *D.*

lumholtzi were reared for 6 months in artificial lake medium (COMBO) and fed *Scenedesmus*

obliquus and *Chlorella vulgaris* ad libitum to dull residual effects of lake predation. Prior to

experimental perturbations, *D. lumholtzi* were common garden reared for three generations in

COMBO medium (Kilham et al., 1998) and fed a standard algae mixture of *S. obliquus* and *C.*

vulgaris to minimize the influence of maternal effects on experimental generations.

Fish and Control Media Preparation. To prepare fish (induction) medium, 10L of artificial

lake medium (COMBO) was conditioned with five *Gasterosteus aculeatus* for three days prior to

filtration (0.45 μm). Fish were fed 10 live *D. lumholtzi* and 10 *D. magna* daily to provide both

conspecific and heterospecific prey cues, maximizing predation kairomone concentration. To

create control medium, artificial lake medium without fish conditioning was created in parallel

with fish medium. Media was frozen and thawed overnight prior to use.

Initial F0 Predation Exposure and Experimental Conditions. From the third generation of

common garden reared *Daphnia*, one mother was selected due to observed clutch survivability

and clutch size. Nine neonates born within a 12 hour period from the same mother were isolated

and randomly assigned to treatment groups, either fish exposure or control. Animals were reared

individually in 50 mL plastic beakers with 30mL of the appropriate medium and fed 5 mL

standard algae mixture every other day, ensuring replication at the individual level. Individual

Daphnia were staged, imaged, and returned to fresh media with algae every other day for the

duration of their lives.

Propagation across Generations. Upon reproduction, up to six neonates from each mother's

first clutch were randomly collected and reared individually in 50 mL beakers containing 30 mL

control medium. If individuals from the mother's first clutch did not survive to reproduce,

neonates from the mother's subsequent clutch were used instead. This ensured propagation of all

lineages while maintaining consistency in clutch selection across treatments. See Figure 1 and

Table 1 for samples sizes.

Imaging and Data Collection. Every other day, animals in the F0, F3, F4, and F5 generation

were staged on a glass slide and imaged using `cellSenseEntry` (Olympus Life Science). Scale

bars were added to images during acquisition for calibration during measurements.

Morphological measurements of *headspine* (tip of spine to top of compound eye), *body length*

(top of eye to base of tail), *tailspine* (base of tail to tip of tailspine), *total body length* (tip of

headspine to tip of tailspine) were collected (see Figure 2) using `ImageJ` (Schindelin et al.,

2012). All images were calibrated to a consistent scale using the 'set global scale' function in

`ImageJ`, which converts pixel distances to micrometers based on the embedded scale bar. See

Figure 2 for measurement details.

Ontogenetic Trajectories. To illustrate general patterns of ontogenetic development in somatic

and defensive traits, Figure 3 presents scatterplots pooled across generations, using `ggplot2`

(Wickham, 2016). Generation- and treatment-specific effects are subsequently analyzed in detail

in our primary statistical models (see Primary Hypothesis section below).

Statistical Analysis.

Primary Hypothesis: Phenotype Transmission Across Generations. Because individuals were

measured repeatedly within generations, we used hierarchical mixed models to account for non-

independence. We employed complementary Bayesian hierarchical and frequentist mixed-effects

approaches to ensure robust inference. Bayesian models provide full posterior distributions

enabling probabilistic statements about treatment effects and improved estimation with sparse

data through partial pooling (Gelman & Hill, 2006). Frequentist models provide p-value-based

hypothesis testing for readers familiar with traditional frameworks. Importantly, both approaches

yielded consistent effect size estimates and overlapping confidence/credible intervals across all

generations and traits (Table 2, Supplemental Table 2), supporting the reliability of our

conclusions regardless of statistical framework.

Morphological variation was analyzed using both frequentist and Bayesian mixed-effects

models in R (R Core Team, 2021), using the `lme4` (Bates et al., 2015) and `brms` (P. C. Bürkner,

2017) packages respectively. For each generation (*F0, F3, F4, F5*) and each trait (*headspine,*

tailspine), we tested for the effects of predation treatment, with *body length* as a covariate, and a

random intercept for individual identity to account for repeated measures on individuals. The

model formula used was:

362   `spine ~ treatment + bodylength + (1|individual)`

where *individual* represents the individual Daphnia identity, accounting for repeated measures on

the same organism over time. The *bodylength* term is a covariate accounting for differences in

body length that contribute to spine length, and treatment represents whether the F0 progenitor

was exposed to predation (for F3, F4 and F5); F0 treated animals were directly exposed. Spine

refers to either the measured head or tailspine length.

We selected this model structure based on established principles in *Daphnia*

morphological analysis and our experimental design. *Body length* was included as a covariate

because spine length scales allometrically with body size in *Daphnia* and failing to account for

this relationship can confound treatment effects with size-related variation (Laforsch & Tollrian,

2004; Petrusek et al., 2009). *Treatment* (predator exposure vs. control) was included as the main

fixed effect to test our primary hypothesis of transgenerational inheritance of induced defenses.

The random intercept for individual identity *(1|individual)* accounts for repeated measurements

on the same individuals, addressing potential pseudo-replication issues common in experimental

ecology (Freeberg & Lucas, 2009; Hurlbert, 1984). We included *body length* rather than *age* as a

covariate because they are highly correlated (Pearson's r = 0.85; Supplemental Figure 1), and

body length more directly captures the allometric relationship between body size and spine

development (Laforsch & Tollrian, 2004).

Bayesian Hierarchical Models. We implemented Bayesian hierarchical models with the model

structure described above. The hierarchical structure allows partial pooling across individuals,

improving estimation in the presence of sparse data. This method also improves upon *p*-value

based frequentist methods by providing a full posterior distribution for the effect size, allowing

probabilistic statements about parameter values rather than relying on a binary significant/non-

significant framework. Models were fit using a Student's t-distribution to account for potential

outliers and heavy-tailed data, which are common in ecological trait data. We used weakly

informative priors and ran three chains with 5,000 iterations each (2,000 warm-up), ensuring

convergence ($\hat{R} < 1.01$ for all parameters). Posterior distributions were summarized to extract

treatment effect estimates and 95% credible intervals. We generated posterior predictions at the

mean body length using `posterior_epred()` from the `posterior` package (Bürkner et al.,

2025), isolating treatment effects while controlling for individual-level variation. These

predictions are visualized in Figure 4. We determined adequate model fit and convergence using

$\hat{R}$ values, effective sample sizes (ESS), leave-one-out cross-validation (LOOIC), WAIC

(Supplemental Table 1; (Vehtari et al., 2021) , trace plots, and posterior predictive checks

(Supplemental Figure 2 and 3; Gabry et al., 2019). Marginal and conditional R² values were

computed using the `performance` package (Lüdecke et al., 2021) to assess model fit. Treatment

effect estimates, including standard errors, confidence intervals, probability of direction (PD),

and R² values are presented in Table 2. The predicted effect of treatment on defensive traits

across generations is visualized in Figure 4.

Frequentist Mixed-Effects Models. To provide familiar p-value-based hypothesis testing, we

also conducted frequentist analysis using the `lme4` (Bates et al., 2015) R package with the same

model formula. Linear mixed models were evaluated using Kenward-Roger F-tests and post hoc

Tukey tests to assess treatment effects. Conditional and marginal R² values, AIC, and BIC were

calculated to assess model fit (Supplemental Table 3). Bayesian and frequentist estimates were

highly consistent across generations and traits, with overlapping intervals and similar effect sizes

(Table 2; Supplemental Table 2). For transparency, raw means comparisons are presented in

Supplemental Figure 5.

Primary Analysis: Testing for Size-Dependent Treatment Effects. To specifically assess the

effect of treatment on *body length* across generations, we again utilized Bayesian hierarchical

modeling using the following formula:

```
bodylength ~ treatment + headspine + tailspine + (1|individual)
```

413   This model determines whether *treatment* affects body size independent of spine elongation. We

414   used *spine* lengths (and not *age*) as covariates because *age* and *body length* are highly correlated

415   (Pearson's r =0.85, Supplemental Figure 1) and because our spine models used *body length* (not

416   *age*) as a covariate, making this specification more consistent. As with the spine models, we used

417   weakly informative priors and a Student's t distribution and ran 2000 warmup and 4000

418   iterations using 4 chains and 4 cores utilizing the `brms` R package (Bürkner, 2017). Trace plots

419 and posterior predictive checks confirmed model convergence and adequate model fit and are

420 included in the supplemental materials for reference (Supplemental Figure 6).

421 ***Secondary Analysis: Age-Dependent Morphological Plasticity (Delta).*** To assess how

422 morphological traits change across *age* and *treatment*, we calculated within-individual

423 morphological change (Δ) for *body length*, *total body size*, *headspine*, and *tailspine*. Δ values

424 were computed as the difference between successive measurements for each individual (`Δtrait`

425 `= trait_current – trait_previous`), capturing short-term plastic responses over time. Age

426 trajectories (*Δ trait* vs. *age*) were visualized using scatterplots with LOESS-smoothed trend lines

427 and 95% confidence intervals (Figure 5). To statistically evaluate the effect of induction on

428 morphological plasticity, we modeled Δ values using linear mixed-effects models, `lme4` (Bates

429 et al., 2015) and `lmerTest` package (Kuznetsova et al., 2017), with *treatment* and *age* as fixed

430 effects and individual identity as a random intercept to account for repeated measures:

431   `delta value ~ treatment + age + (1|individual)`

432 Because this analysis addressed a secondary question about short-term morphological plasticity

433 rather than our primary hypothesis on spine inheritance, we opted for a parsimonious linear

434 mixed-effects approach. This strategy ensures computational efficiency for secondary analyses

435 while providing the level of inferential detail appropriate for each research question. We initially

436 explored more complex temporal modeling approaches, including natural splines with varying

437 degrees of freedom (df = 2 to 4) to capture potential non-linear age-related changes in growth

438 patterns:

439   `delta value ~ treatment + ns(age, df = 2) + (1|individual)`

440 However, these models suffered from overparameterization, and model significance was

441 sensitive to the choice of degrees of freedom, indicating unstable parameter estimates.

442 Consequently, we adopted the more parsimonious linear approach with *age* as a continuous

443   covariate, which provided robust and interpretable results while adequately capturing the age-

444   related trends in our data. Results were visualized in Table 3, including the estimated effect of

445   treatment, standard error, F-statistic, and *p*-value. For a more granular approach, we also

446   modeled these effects by generation and present the same parameters (Supplemental Table 4).

447   ***Secondary Analysis: Size Selective Survivorship Analysis.*** To test whether size-selective

448   mortality, rather than treatment, could explain differences in spine lengths, we constructed

449   mixed-effects linear models using binary survival status. Because this analysis addressed a

450   secondary question about size-selective mortality rather than our primary hypothesis on spine

451   inheritance, we opted for a parsimonious frequentist mixed-effects approach. Bayesian modeling

452   was unnecessary here because the goal was to test a simple association between body size and

453   spine length, and linear mixed-effects models provided interpretable estimates while accounting

454   for repeated measures. This approach balances computational efficiency with the depth of

455   inference appropriate for each research question. *Survival status* was defined as the presence or

456   absence of a subsequent timepoint for each individual. Separate models were fit for *headspine*

457   and *tailspine* lengths as a function of *body length*, with a random intercept for individual identity.

458
```
spine ~ bodylength + (1|individual)
```

459   Residuals from these models represent spine length variation independent of body size. To

460   visualize trait dynamics across age and survival status, we plotted body length and spine

461   residuals using LOESS-smoothed trend lines with 95% confidence intervals. Survivors and non-

462   survivors were distinguished by color, and traits were faceted for comparison (Figure 6).

463   To further evaluate survival probability, we constructed a binomial generalized additive

464   model (GAM) with *survival status* as the response variable and *age*, *body length*, *spine* lengths,

465   treatment, and the interaction between *age* and *treatment* as predictors.

466
```
status ~ bodylength + headspine + tailspine + age * treatment
```

467   Trait values representing the 10th, 50th, and 90th percentiles were used to generate survival

468   predictions across *age*. Predicted survival probabilities and 95% confidence intervals were

469   plotted to assess how trait variation influences survivorship over time (Figure 7).

470   ***Sample Size Considerations.*** Due to attrition and differences in fecundity and survivorship

471   across 6 generations, the F5 generation suffered from lower sample sizes comparatively,

472   particularly in the control animals. Although we employ Bayesian hierarchical models, which are

473   better equipped to handle unbalanced designs and sparse sample sizes, we conducted a

474   retrospective power analysis using the mean effect size from the F3 and F4 generations

475   (Supplemental Table 6) using the `pwr` R package and the `pwr.t.test` function (Champely et

476   al., 2020).

478   **Statistical Software and Reproducibility.** All statistical analyses and visualizations were

479   conducted in **`R`** (R Core Team, 2021). Data manipulation and cleaning were performed using the

480   **`dplyr`** package (Wickham et al., 2022) , and visualizations were generated using `ggplot2`

481   (Wickham, 2016), with additional formatting and layout support from `patchwork` (Pedersen,

482   2019), `gghighcontrast` (Grosenbach, 2021) **,** and `Rmisc` (Hope, 2022) for summary

483   statistics. Estimated marginal means and post hoc comparisons were calculated using `emmeans`

484   (Lenth & Piaskowski, 2017).

485   All analyses were conducted using `Rstudio` (Posit Team, 2025) in `RMarkdown` (Allaire

486   et al., 2019) and `Quarto` (Allaire et al., 2025), ensuring full reproducibility of code, figures, and

487   statistical outputs. Annotated Quarto files and datasets are available on GitHub at

488   https://github.com/shannonsnyder/daphnia-transgenerational-plasticity.

490   Table 1: Sample sizes across experimental generations and treatment showing number of

491   individuals and repeated observation details.

Generation	Treatment	Individuals	Observations	Mean Obs/Ind
F0	induced	5	66	13.2
	control	4	34	8.5
F3	induced	74	480	6.5
	control	58	124	2.1
F4	induced	63	556	8.8
	control	17	131	7.7
F5	induced	41	500	12.2
	control	8	45	5.6

492
493   F0 generation began with 9 founder individuals (4 control and 5 predator induced individuals) produced in the same
494   clutch by the same mother. All subsequent generations (F1-F5) were reared in control medium only. 'Individuals' =
495   unique Daphnia measured within that generation/treatment; 'Observations' = total measurements across repeated
496   sampling (every other day) within generation/treatment; 'Mean Obs/Ind' = average number of repeated measurements
497   per individual. F1 and F2 generations were propagated in control medium but not measured, as their phenotypes can
498   result from direct embryonic exposure (see Methods for detailed rationale). Variation in sample size across treatments
499   and generations reflects biological differences in survival and reproductive success between predator-exposed and
500   control lineages. Note the reduced sample sizes in F5, particularly among control individuals, which is addressed with
501   a retrospective power analysis (Supplemental Table 5).

502   **Results**

503   **Defensive and Somatic Traits Display Distinct Ontogenetic Patterns.** *Daphnia*

504   *lumholtzi* exhibited distinct developmental trajectories across their lifespan for body length,

505   headspine, and tailspine structures, regardless of treatment (Figure 3). Defensive spine traits

506   displayed age-dependent plasticity, with early-life elongation followed by gradual reduction in

507   spine length across all individuals, independent of treatment (Fig. 3B-C). Headspine length

508   peaked (~850 μm) within the first 10 days and declined steadily to ~400 μm in older individuals

509   (Figure 3B). Tailspine length reached a delayed maximum (~1,800 μm) around 20 days before

510   declining to values comparable to those observed in early developmental stages (~700 μm)

511   (Figure 3C). These general developmental patterns provide the ontogenetic context within which

512   treatment-specific differences (Fig. 4) should be interpreted.

513
514   **Predator cue exposure results in an increase in headspine and tailspine length in the**

515   **exposed generation.** Because most morphological changes occur within the first 20 days of life,

516   before spine growth plateaus (Figure 3), we modeled treatment effects on individuals less than

517   20 days of age. Consistent with previously documented within-generation plastic responses in

518   other species, F0 *Daphnia lumholtzi* reared in predator cue environments exhibited robust and

519   statistically significant increases in both headspine and tailspine length relative to controls,

520   indicating postembryonic predator cue exposure was sufficient to produce a substantial increase

521   in both headspine and tailspine length. In the F0 generation, with predation cues provided from

522   the first post-embryonic day, Bayesian models estimate significant increases in induced defense

523   trait lengths (Figure 4; Table 2). Specifically, we report an estimated increase in headspine length

524   of nearly 70 μm (95% CI: [1.67,132.90]). These differences correspond to a roughly 10%

525   increase in headspine length in exposed Daphnia over control Daphnia. Tailspine length also

526   increased dramatically in the exposed F0 individuals, by nearly 100 µm in directly exposed

527   Daphnia (95% CI: [15.35,181.54]), also representing a roughly 10% increase.

528
529   **Table 2. Posterior mean estimates of treatment-induced changes in spine length across**
530   **generations**
531

Spine	Generation	Estimate	SE	95% CrI	PD	R² (marg.)
Head Spine	F0	68.54	32.90	[1.67, 132.9]	**97.7%**	60.4%
	F3	38.19	18.32	[3.1, 74.71]	**98.5%**	64.9%
	F4	44.36	20.45	[3.99, 84.58]	**98.4%**	65.8%
	F5	-7.07	22.32	[-51.23, 36.78]	62.6%	56.5%
Tail Spine	F0	99.71	41.28	[15.35, 181.54]	**98.6%**	94.5%
	F3	5.81	18.35	[-30.24, 41.81]	62.5%	98%
	F4	85.57	34.76	[16.95, 154.28]	**99.3%**	98.3%
	F5	-19.21	29.07	[-77.17, 37.5]	74.7%	94.8%

532   Models were fit separately by generation for individuals <20 days old. Each row represents a unique combination of
533   generation and spine type. 'Estimate' is the posterior mean difference in spine length (µm) between predator-induced
534   and control treatments, with positive values indicating elongation in induced individuals. 'SE' is the posterior standard
535   deviation. '95% CrI' denotes the 95% credible interval for the posterior distribution. 'PD' (Probability of Direction)
536   represents the posterior probability that the effect is in the estimated direction (positive indicating higher trait values in
537   treated, and negative indicating lower trait values in treated, compared to control). 'R² (marg.)' is the marginal
538   Bayesian R², indicating variance explained by fixed effects only. See Methods for full model specification and prior
539   structure.
540

541   **Defensive Traits Are Inherited Through F4.** Phenotypic differences persisted for multiple

542   generations after removal from the inducing environment, in generations sufficiently removed

543   from embryonic and germline exposure. When F0 mothers are exposed, their developing F1

544   embryos are directly exposed in the brood chamber, and the germline cells within F1 embryos

545   that will produce F2 are also exposed. Thus, F2 represents the last generation with potential

546   germline exposure. Only by the F3 generation are any observed effects transmitted through

547 purely non-genetic mechanisms, as F3 individuals and their germ cells had no direct contact with

548 predator cues.

549   Despite two intervening generations without exposure, F3 individuals with ancestral

550 predation exposure in F0 exhibited a significant increase in headspine length. Bayesian inference

551 yielded an effect size of almost 40 μm (95% CI: [3.10,74.71]). This represents ~56% retention of

552 the F0 headspine response magnitude (38.19 μm vs. 68.54 μm). In contrast, tailspine length in F3

553 showed no significant difference between treated and control individuals ($\beta = 5.81$ μm, 95% CI:

554 [-30.24, 41.81]), indicating trait-specific attenuation of the tailspine below a statistically

555 detectable threshold.

556   In the F4 generation, also completely unexposed to predator cues, both headspine and

557 tailspine lengths were again significantly elevated in the induced animals as compared to the

558 control individuals. Ancestral predator exposure increased the headspine length by 44.36 μm

559 (95% CI: [3.99,84.58]) and the tailspine length by 85.57 μm (95% CI: [16.95,154.28]).

560 Therefore, F4 headspine retained ~65% of the F0 response magnitude (44.36 vs. 68.54 μm),

561 while F4 tailspine retained ~86% (85.57 vs. 99.71 μm), demonstrating trait-specific patterns of

562 transgenerational persistence (Table 2). Tailspine showed a distinct pattern from headspine: non-

563 significant in F3 but strongly significant in F4, suggesting potential generational fluctuations in

564 trait expression or differential transgenerational mechanisms between the two defensive

565 structures.

566   By the F5 generation all measurable differences between induced and control individuals

567 had disappeared. We report an estimate of -7.07 (indicating treated animals have a shorter

568 headspine; 95% CI: [-51.23,36.78]) for the headspine and -19.21 μm (95% CI: [-77.17, 37.50]

569   for tailspine length (Table 2, Figure 4). Estimated differences were small with 95% credible

570   intervals widely overlapping zero for both headspine (95% CrI: [-51.23, 36.78]) and tailspine

571   (95% CrI: [-77.17, 37.50]), indicating no detectable treatment effect. These findings indicate that

572   the induced phenotypes ultimately reverted to a statistically undetectable baseline in the absence

573   of continued environmental cues over 5 unexposed generations. Attrition across six generations

574   resulted in a small sample size for this generation, making it difficult to discern if we lack the

575   statistical power to detect a subtle effect.

576   Model fit was strong across all analyses, with marginal R^2 values ranging from 56-66%

577   for headspine models and 94-98% for tailspine models, indicating that treatment and body length

578   together explain substantial variance in spine morphology. Frequentist linear mixed-effects

579   models yielded highly consistent results (Supplemental Table 2), with overlapping confidence

580   intervals and similar effect size estimates, confirming the robustness of our inferences across

581   statistical frameworks.

582   These results provide rare empirical evidence of vertebrate predator-induced

583   transgenerational plasticity persisting through the F4 generation. The temporal trajectory of

584   treatment effects across generations (Figure 4) reveals both the persistence and ultimate decay of

585   transgenerational plasticity. Headspine effects declined from F0 through F5, while tailspine

586   showed an unexpected resurgence in F4 after non-significant F3 effects, highlighting trait-

587   specific dynamics. The persistence of these effects through four generations, particularly in a

588   species with rapid generation times (~2 weeks), demonstrates substantial temporal stability of

589   non-genetic inheritance.

590   **Treatment Effects on Body Length are Minimal.** In contrast to defensive traits, body

591   length increased rapidly as the animals aged in both control and induced individuals, reaching an

592   early asymptote (~2,500 μm) around 25 days. This growth pattern was strongly correlated with

593   age (Pearson's r = 0.85; Supplemental Figure S1). Visual inspection of ontogenetic patterns

594   (Figure 3) showed overlap between body length trajectories between control and induced

595   animals, likely demonstrating trait-specific inheritance of defensive morphologies independent of

596   general body size effects. Indeed, confidence intervals for the Bayesian models illustrating the

597   effect of treatment on body length across generations largely overlap zero, with probability of

598   direction (PD) estimates hovering around 50% (PD > 97.5% typically indicates strong directional

599   evidence), and estimated body size differences between -2.63 and 0.46 μm (treated - control;

600   Table 3), except for the F4 generation. Notably, F4 confidence intervals narrowly overlapped

601   zero (-122.54, 2.22) with an estimated body length difference of -61.45 μm, and a PD of 97.1%,

602   just shy of the conventional 97.5%, providing some support for a smaller body size due to

603   predation exposure in this generation only. Effect sizes were negligible across most generations

604   (Cohen's d = -0.05 to 0.03), with only F4 showing a small to medium effect (d = -1.16), further

605   supporting that ancestral predation exposure primarily affects defensive structures rather than

606   general body size.

607   Together, these ontogenetic analyses demonstrate that transgenerational inheritance of

608   predator-induced traits is largely specific to defensive structures and occurs independently of

609   changes in body size. There is, however, some evidence of smaller body sizes due to ancestral

610   predation exposure in the F4 generation alone.

612   **Table 3. Treatment effects on body length controlling for spine morphology**.

Generation	Est. Difference (μm)	95% CrI	Cohen's d	PD
F0	-0.53	[-99.45, 95.21]	< 0.01	50.3%
F3	-2.53	[-39.42, 34.84]	-0.05	55.1%
F4	-61.45	[-122.54, 2.22]	-1.16	97.1%
F5	0.46	[-93.78, 89.51]	0.03	51.7%

613 Treatment effects on body length across generations. Bayesian hierarchical models tested whether ancestral
614 predator exposure affected body size independent of spine elongation, controlling for headspine and tailspine length,
615 and with a random intercept for individual identity. Negative estimates indicate smaller body size in predator-exposed
616 individuals. See Methods for full model specification. 'Generation' refers to the generation examined, 'Est. Difference
617 (μm)' is the estimated effect of treated – control, therefore negative values represent a smaller estimated body size in
618 treated animals. '95% CrI' depicts the 95% credible intervals. For interpretability, we include the estimated effect sizes
619 in the 'Cohen's d' column. 'PD' (Probability of Direction) represents the posterior probability that the treatment effect is
620 in the estimated direction. Values >97.5% indicate strong evidence for the effect direction; values near 50% indicate
621 high uncertainty.
622

623 These results demonstrate that predator-induced morphological plasticity can be inherited

624 through at least four generations and is largely concentrated to defensive spines rather than body

625 length. This inheritance is finite and trait-dependent, with detectable headspine elongation

626 through the F4 generations. The decrease of effect sizes and the disappearance of measurable

627 trait differences in F5 underscores the transient nature of this plasticity and suggests that the

628 mechanism of transgenerational epigenetic inheritance decays over time without environmental

629 cue reinforcement (Figure 4).

630 **Differences in Growth Rate and Trait Distributions Do Not Explain Induced Trait**

631 **Expression**.

632 To evaluate whether the observed morphological differences between induced and

633 control individuals could be attributed to variation in growth dynamics (rather than final trait

634 size) or trait distributions, we assessed whether treatment effects could be explained by

635 differences in developmental timing, size-selective mortality, or trait-based survival.

636   The most pronounced morphological changes occurred early in life, prior to 20 days of

637   age (Figure 3). This period includes the onset of reproduction (typically 7–12 days), during

638   which *D. lumholtzi* mothers are not yet fully grown. Morphological change peaked around this

639   time for all traits, particularly for body length and total body size (Figure 5).

640   Importantly, no significant differences in growth rate were detected between treatment

641   groups for any of the four traits examined (body length, headspine, tailspine, and total body

642   size), indicating that predator exposure did not alter the rate of somatic or defensive trait

643   development (Table 4). Specifically, treatment effects were non-significant for body length (Est.

644     $= 5.70 \pm 7.94$ µm, $F = 0.52$, $p = 0.47$), total body size (Est. $= 9.87 \pm 19.20$ µm, $F = 0.27$, $p =$

645   0.61), headspine (Est. $= 6.51 \pm 4.57$ µm, $F = 2.06$, $p = 0.15$), and tailspine (Est. $= 1.88 \pm 7.57$

646   µm, $F = 0.06$, $p = 0.80$). Generation-specific analyses similarly revealed no treatment effects on

647   growth rates in any individual generation (all $p > 0.05$; Supplemental Table 4), confirming that

648   differences in defensive trait expression persist consistently across F0-F5 independent of

649   developmental timing. These results suggest that the observed morphological differences are not

650   due to accelerated or delayed growth in induced individuals but instead reflect differences in

651   final trait expression independent of growth rate.

652   **Table 4: Growth rate is not influenced by treatment**

Trait	Est. Difference (µm)	Std. Error	F statistic	P value
Body Length	5.70	7.94	0.52	0.469
Total Body Size	9.87	19.20	0.27	0.605
Headspine	6.51	4.57	2.06	0.151
Tailspine	1.88	7.57	0.06	0.802

653 Results of a linear model estimating the effect of treatment on trait change between timepoints. Analysis includes all
654 individuals measured during the first 20 days of life across all generations (n = 183 induced, n = 87 control) 'Trait'
655 represents the morphology examined, the 'Est Difference (μm)' represents the model estimated difference (treated –
656 control; positive values indicate larger estimates in treated animals compared to control). 'Std. Error' describes the
657 standard error of the estimate, and the F-statistic and *p*-value for the corresponding morphology are represented in the
658 'F statistic' and 'P value' columns.

659   To assess whether trait distributions might reflect selective mortality, such as the

660 preferential survival of individuals with longer spines, we compared growth trajectories of

661 survivors and non-survivors during the first 20 days of life. Plotting developmental trajectories

662 for body length, headspine residuals, and tailspine residuals (residuals control for body size) by

663 survival status (Figure 6) showed that while survivors tended to have a larger body length,

664 **residual spine lengths showed overlapping distributions and trajectories** between survivors

665 and non-survivors. Overlapping distributions of spine residuals between survivors and non-

666 survivors (Figure 6) indicate that trait-based selection is unlikely to account for the observed

667 morphological differences.

668   The absence of an effect of spine length on survivorship was further supported by a

669 generalized linear model (GLM) predicting survival status from body length, headspine,

670 tailspine, age, and treatment. The model revealed that **body length was positively associated**

671 **with survival** ($\beta = 0.0030$, $p < 0.001$), while **headspine and tailspine lengths were not**

672 **significant predictors** (headspine: $\beta = -0.0010$, $p = 0.058$; tailspine: $\beta = -0.0007$, $p = 0.110$).

673 Individuals from predator-exposed lineages showed significantly higher survival than controls (β

674 $= 0.678$, $p < 0.001$), suggesting that transgenerational defensive trait expression may confer

675 fitness benefits even in the absence of direct predator exposure. Importantly, this survival

676 advantage was independent of spine length, as neither headspine nor tailspine significantly

677 predicted survival status. The model explained a substantial portion of deviance (ΔDeviance =

678   328.8; AIC = 2554.1), but **no evidence emerged for size- or trait-selective mortality based on**

679   **spine length** (Figure 7).

680   **Power Analysis.** F5 sample sizes were reduced due to natural attrition across six

681   generations (n=8 control individuals, 41 induced; Table 1). Retrospective power analysis using

682   mean effect sizes from F3-F4 generations yielded estimated power of 0.30 for headspine and

683   0.66 for tailspine (Supplemental Table 5). These estimates are conservative because they do not

684   account for repeated measures (average 12.2 observations per individual for induced, 5.6 for

685   control; Table 1), which increase effective sample size. While we cannot definitively distinguish

686   between trait decay and insufficient power in F5, the pattern across F0-F5 suggests gradual

687   attenuation consistent with temporal decay of transgenerational effects.

688   These findings demonstrate that transgenerational plasticity in *D. lumholtzi* operates

689   primarily on defensive structures, persists for at least four generations following ancestral

690   exposure, and is not due to size-selective or trait-selective mortality.

691   **Discussion**

692   Our findings demonstrate that predator-induced morphological defenses in *Daphnia*

693   *lumholtzi* persist through at least four unexposed generations via non-genetic mechanisms.

694   Critically, measurable effects in F3, F4, and (to a lesser extent) F5 occurred in cohorts with no

695   direct, embryonic, or germline exposure to predator cues, a necessary criterion for distinguishing

696   transgenerational plasticity from extended maternal effects (Agrelius & Dudycha, 2025; Skinner,

697   2008). This temporal persistence is notable given that F3 represents the first generation where

698   developing germ cells had no contact with the inducing signal, and F4-F5 are even further

699   removed. These data contribute to a limited but growing body of evidence demonstrating that

700   environmentally induced phenotypes can be inherited across multiple generations through

701   epigenetic or other non-genetic mechanisms.

702   Our experimental design intentionally employed a single clonal lineage to isolate non-

703   genetic effects from standing genetic variation, enabling us to definitively demonstrate

704   transgenerational inheritance independent of allelic differences. However, this design limits

705   inference to one genetic background. The existence of 'uninducible' *D. lumholtzi* clones with

706   constitutive defenses (Engel & Tollrian, 2009), even from nearby populations in Arizona,

707   demonstrates substantial genetic variation in plasticity and defense strategies within this species.

708   Future work must examine whether the 4-generation persistence we observed is representative

709   across clones or population-specific, and whether genotype-by-environment interactions

710   influence the temporal dynamics of transgenerational effects (Auld et al., 2010; Scoville &

711   Pfrender, 2010).

713   **Induced Phenotypes Persist Across Generations but Fade Without Cues**

714   Headspine and tailspine elongation persisted through four unexposed genetically identical

715   generations, with measurable effects in F3 and F4 despite the absence of continued induction.

716   This transgenerational persistence supports the hypothesis that environmental information can be

717   retained and expressed across generations via non-genetic mechanisms (Gapp et al., 2014; Heard

718   & Martienssen, 2014; Rechavi et al., 2014).

719   By the F5 generation, trait differences between induced and control individuals were not

720   detectable. This regression to baseline aligns with evolutionary expectations that plastic traits

721   fade when environmental cues vanish, minimizing fitness costs associated with unnecessary

722   defenses (DeWitt et al., 1998; Van Buskirk & Steiner, 2009).

723   Headspine and tailspine displayed divergent transgenerational trajectories, suggesting

724   trait-specific non-genetic regulation. Headspine effects declined rather monotonically across

725   generations (68 μm → 38 μm → 44 μm → -7 μm), consistent with gradual erosion of epigenetic

726   marks. In contrast, tailspine showed non-significant effects in F3 (6 μm) but pronounced effects

727   in F4 (86 μm), approaching F0 magnitude (100 μm). This "resurgence" pattern could reflect: (1)

728   differential stability of epigenetic marks regulating the two traits, (2) compensatory

729   developmental mechanisms favoring tailspine expression when headspine inheritance weakens,

730   or (3) stochastic fluctuations in small sample sizes. These trait-specific dynamics highlight that

731   transgenerational plasticity is not uniformly expressed across correlated traits and may involve

732   independent regulatory mechanisms (Bonduriansky & Day, 2009; Herman & Sultan, 2011; Uller

733   et al., 2013). Future work should examine whether these two defensive structures are regulated

734   by distinct epigenetic pathways.

735   Retrospective power analysis for F5 yielded estimated power of 0.30 for headspine and

736   0.66 for tailspine, based on mean effect sizes from F3-F4 generations. While we cannot

737   definitively distinguish between true decay and insufficient power in F5, several lines of

738   evidence support the former: (1) the decline in effect sizes across F0-F4, (2) credible intervals for

739   F5 estimates centered near zero rather than showing directional trends, and (3) the biological

740   expectation that transgenerational effects should eventually attenuate without environmental

741   reinforcement. Achieving high statistical power in ecological studies, especially with

742   multigenerational designs is inherently challenging (Kimmel et al., 2023) but our use of repeated

743   measures and hierarchical modeling maximized inferential strength given practical constraints.

744   The 5-generation persistence of predator-induced defenses has implications for

745   population dynamics in variable predation regimes. In systems with fluctuating predator densities

746   (seasonal fish migrations, predator reintroductions, reproductive timing), prey populations may

747   remain partially defended for 4-5 generations (several months in this *Daphnia* species) after

748   predator removal. This "ecological memory" could buffer populations during temporary predator

749   absences but may incur costs if defenses persist longer than threats. Fitness consequences,

750   including fecundity and survival across generations will be the focus of future work in our

751   predator-prey system.

753   ***Daphnia lumholtzi* Induced Defenses Are Robust, Age-Specific, and Not Driven by Selection**

754   Exposure to predator cues in the F0 generation triggered strong morphological defenses

755   in *Daphnia lumholtzi*, with significant elongation of both headspine and tailspine structures.

756   Post-embryonic exposure alone was sufficient to induce these traits in our Arizona clone,

757   demonstrating that embryonic priming is not universally required for defense induction in this

758   genus (Laforsch & Tollrian, 2004). This finding underscores the differences in inducibility

759   between clones and species and highlights the need to assess transgenerational inheritance in a

760   variety of systems (Auld et al., 2010; Scoville & Pfrender, 2010; Shahmohamadloo et al., 2025).

761   This is further illustrated by the characterization of "uninducible" *D. lumholtzi* clones which

762   constitutively express defense phenotypes even in the absence of predation cues (Engel &

763   Tollrian, 2009).

764   Importantly, these trait differences were not attributable to size-selective or trait-selective

765   mortality. Survivors and non-survivors showed overlapping developmental trajectories, and

766   survival probability was primarily associated with body length, not spine morphology. This

767   confirms that the observed phenotypic differences are induced responses rather than artifacts of

768   differential survivorship among individuals based upon armor morphology (Dieckmann et al.,

769   2006; Kingsolver et al., 2001).

770   We also found that spine lengths were most prominent in early life and declined with age,

771   a pattern distinct from somatic growth. This suggests stage-specific optimization of defenses,

772   likely targeting periods of highest vulnerability to predation (Nagano & Yoshida, 2020; E. E.

773   Werner & Gilliam, 1984). Smaller, younger Daphnia likely experience higher predation rates

774   from gape-limited fish predators (Nagano & Yoshida, 2020), making early-life defenses

775   particularly adaptive. The age-dependent decline in spine length we observe may represent

776   adaptive reallocation from defense to reproduction as individuals reach body sizes less

777   vulnerable to predation. This ontogenetic trajectory, previously unreported for *D. lumholtzi*

778   headspines, parallels patterns in other inducible defense systems where defenses are most

779   pronounced during vulnerable life stages (Nagano & Yoshida, 2020) and documented tailspine

780   shortening with age in *Daphnia* (Gu et al., 2021).

782   **Mechanisms of Non-genetic Inheritance and Evolutionary Implications**

783   The persistence and gradual decline of induced traits across generations strongly suggests

784   epigenetic inheritance as a likely mechanism. Having established the phenotypic baseline and

785   generational persistence, future mechanistic work can now uncover the molecular mechanisms

786   underlying these patterns. Candidate mechanisms include, DNA methylation, histone

787   modification, and chromatin-level changes that facilitate the transmission of plasticity without

788   altering the genetic code (Banerjee & Roychoudhury, 2017; Casadesús, 2016; Davidowitz, 2009;

789   Ho & Burggren, 2010; Li & Zhang, 2014; Niederhuth et al., 2016; Robichaud et al., 2012; M. S.

790   Werner et al., 2023; Yaniv, 2014). The temporal decay pattern we observed is consistent with

791   incomplete erasure of epigenetic marks (Dias & Ressler, 2014; Gapp et al., 2014; Grishkevich &

792   Yanai, 2013), as documented in other invertebrate and vertebrate systems including

793   *Caenorhabditis elegans* (Kaneshiro et al., 2019), the asexual freshwater snail, *Physa acuta*

794   (Dejeux et al., 2025), *Daphnia magna* (Asselman et al., 2016; Vandegehuchte, De Coninck, et

795   al., 2010), as well as vertebrate systems including mice (Carone et al., 2010) and humans

796   (Mulligan et al., 2025). By identifying these generational phenotypic dynamics, we establish a

797   foundation for uncovering the molecular mechanisms underlying generation-specific adaptive

798   phenotypes, a key priority for future research.

799   Our focus on vertebrate predation addresses an important gap in transgenerational

800   plasticity research. While numerous studies examine *Daphnia* responses to invertebrate predators

801   such as *Chaoborus* larvae (Agrawal et al., 1999; Horstmann et al., 2022), studies of vertebrate

802   predator-induced transgenerational effects remain scarce (Tariel et al., 2020). This gap is

803   consequential because vertebrate predators typically exert more consistent selective pressure than

804   invertebrate predators (Batabyal, 2023; Wooster, 1994) and are central to current conservation

805   efforts. Our findings have direct relevance for predator reintroduction and rewilding efforts,

806   where prey populations may exhibit defenses for 4-5 generations (~2-3 months in *D. lumholtzi*)

807   following ancestral exposure. Understanding the temporal dynamics of these transgenerational

808   responses, especially in a keystone aquatic species, could improve predictions of predator-prey

809   dynamics and ecosystem restoration outcomes (Bracis & Wirsing, 2021; Delibes-Mateos et al.,

810   2019).

811   Our findings align with theoretical predictions that transgenerational plasticity should

812   evolve when parental and offspring environments are positively correlated (Agrawal et al., 1999;

813   Herman & Sultan, 2011). The 4-generation persistence we observe (~2-3 months in this system)

814   matches timescales over which predator densities fluctuate in natural lakes due to seasonal

815   migrations, overwinter mortality, or breeding events. However, the eventual decay by F5 is also

816   adaptive as prolonged expression of costly defenses in predator-free environments would reduce

817   fitness through allocation trade-offs (DeWitt et al., 1998; Van Buskirk & Steiner, 2009). The

818   temporal dynamics we document therefore suggest fine-tuning of inheritance duration to match

819   ecological predictability, though testing this hypothesis would require comparative studies across

820   populations experiencing different predation regimes.

821   Our findings underscore the evolutionary relevance of environmental history in the

822   absence of genetic variation (Jablonka & Lamb, 2006; Laland et al., 2015; Pigliucci & Müller,

823   2010). Transgenerational plasticity allows organisms to respond to ancestral conditions,

824   potentially buffering populations against short-term environmental fluctuations (Donelson et al.,

825   2018; Uller et al., 2018). Because our study intentionally used a single *Daphnia* population,

826   future work should explore responses across diverse genotypes (Auld et al., 2010). The existence

827   of "uninducible" clones with constitutive defenses raises intriguing questions about genetic

828   assimilation and the long-term fate of plastic traits (Lande, 2009; West-Eberhard, 2003).

830   **Conclusions**

831   This work advances understanding of transgenerational plasticity in a keystone aquatic

832   genus by documenting vertebrate predator-induced morphological effects in key generations

833   sufficiently removed from embryonic and germline exposure, therefore revealing trait-specific

834   temporal persistence in defensive structures. Our findings align with previous work documenting

835   clutch size persistence in *D. ambigua* (Walsh et al., 2014), helmet length inheritance in *D.*

836   *cucullata* (Agrawal et al., 1999), and multi-generational erosion of plastic traits in *D. pulicaria*

837   (Landy et al., 2024), while crucially extending the temporal window of observation for predator-

838   induced effects to key, previously undocumented generations.

839   This transgenerational plasticity may buffer populations against short-term ecological

840   fluctuations, while its eventual attenuation minimizes the costs of unnecessary trait expression

841   (Herman & Sultan, 2011; Uller et al., 2013). Future research should integrate molecular profiling

842   (DNA methylation, chromatin modifications, differential transcription) across F0-F5 generations

843   with multi-clone comparative studies to determine both the mechanistic basis and generality of

844   these transgenerational effects. Such integrative approaches will advance understanding of how

845   organisms respond to environmental variation across generational timescales and the role of non-

846   genetic inheritance in evolutionary adaptation to changing environments.

FIGURES

Figure 1. Schematic illustrating the experimental setup allowing for direct testing of transgenerational epigenetic inheritance by induced defenses by comparing individuals with and without grandparental exposure to predator cues. Offspring (F0) from a single unexposed mother were split into two genetically identical groups: control (top), where individuals where continuously reared in artificial lake medium, and induced (bottom), where F0 individuals were exposed to fish-conditioned medium indicated in green. Only the F0 induced generation is exposed to fish conditioned media. Sample size for each generation and treatment is indicated (N ind.), and the number of observations (repeated measures, N.obs) is indicated. Subsequent generations (F3-F5) from both treatment group and control are maintained in artificial lake media. Generations were propagated for six generations using the first viable (successful reproduction) and measured every other day during the F0, F3, F4, and F5 generations.

865
866
867 **Figure 2. Measurement of inducible morphological defenses in Daphnia in response to**
868 **predator cues.** Daphnia reared in fish-conditioned media develop significantly elongated
869 headspines and tailspines compared to control animals. Representative lateral images of
870 morphological measurements are indicated, detailed as follows: Headspine was measured from tip
871 of spine to top of compound eye, body length was measured from top of compound eye to base of
872 tailspine, tailspine was measured from base to tip of spine. Total body size was measured from top
873 of headspine to tip of tailspine.

874
875 **Figure 3. Trajectories of somatic and defensive traits reveal divergence in late-life dynamics.**
876 Scatterplots display individual measurements of (A) body length, (B) total body size, (C)
877 headspine length, and (D) tailspine length (in µm) as a function of age (in days). Green line
878 represents locally averaged (LOESS) trends for induced individuals (F0 exposed to fish-
879 conditioned media); purple line represents locally averages (LOESS) trends for control individuals
880 (no F0 exposure). Somatic traits - body length (A) and total body size (B) - increase rapidly during
881 early juvenile stages, then reach a plateau in mid-life, with little elongation even at advanced ages.
882 In contrast, defensive traits - headspine (C) and tailspine (D) – regress later in life. Morphological
883 trajectories demonstrate distinct patterns in defense trait expression, but not body growth, between
884 control and treated individuals, consistent with the inheritance of plastic response initiated by
885 ancestral exposure.
886

887
888 **Figure 4. Predicted spine lengths by generation and treatment.** Predicted mean
889 headspine length (top row) and tailspine length (bottom row) for *Daphnia* in control and
890 induced individuals across generations (F0, F3 ,F4, F5). Points represent model-predicted
891 means ± standard error. Estimates are based on data from individuals ≤ 20 days old when
892 defensive traits are most pronounced (Figure 3). See Table 2 for underlying estimates and
893 parameters, and Supplemental Table 1 for model diagnostics.
894
895
896
897

898
899
900

901
902 **Figure 5. Growth of defensive and somatic traits are most pronounced early in life, followed**
903 **by a stabilization with age, and were unaffected by treatment.** Changes in morphological
904 measurements (body length, total body size, headspine, and tailspine) across the lifespan (days).
905 Each panel shows the mean growth (Δ; in μm) from the previous timepoint for the indicated trait.
906 Morphological changes are largest early in life and stabilize as individuals age, as shown by the
907 bold smoothed trend line with 95% confidence intervals (shaded area). Induced/treated animals
908 are represented by the solid line, and control animals are indicated by the dashed line.

920
921 **Figure 6. Developmental trajectories during periods of concentrated growth do not indicate**
922 **treatment or size selective survivorship.** Body length, head spine (residual), and tail spine
923 (residual) measurements over first 20 days of age for individuals grouped by survival status (dark
924 grey for those who survived to the next timepoint - survivors, light grey for those that perished).
925 Residual values for head and tail spines are calculated from linear mixed-effects models that
926 account for body length, isolating variation in spine length independent of body size.
927 Both survivors and non-survivors display overlapping developmental trajectories and trait
928 distributions, providing no evidence that size-selective or trait-selective mortality underlies
929 observed group differences.
930
931
932
933

934
935 **Figure 7. Survivorship and morphological trait distributions do not support size-selective**
936 **mortality as a driver of morphological differences.** Modeled probability of survival to the next
937 timepoint as a function of age, body length, spine length, and treatment, based on a generalized
938 linear model. Survival probability was predicted at 10th (low - purple), 50th (mean – blue), and
939 90th (high- yellow) percentiles of trait values, with dashed lines representing induced/treated
940 animals and solid lines representing control. Body length values are associated with a greater
941 likelihood of survival at most ages, whereas different spine lengths show minimal effect on
942 predicted survival, regardless of treatment. This suggests that while body length does relate to
943 survival probability in the model, size-selective mortality is not strong enough to cause large
944 differences in defensive morpholgies.
945
946

949   **Supplemental Table 1: Bayesian Model Diagnostics and Fit Summary.**

Generation	Trait	Max R	Min Bulk ESS	Min Tail ESS	LOOIC	WAIC	p(Direction)	Marginal R²
F0	Head spine	1.00	10025	14059	889.10	888.85	0.98	0.60
	Tail spine	1.00	9045	15377	854.56	854.44	0.99	0.95
F3	Head spine	1.00	10641	21785	3,634.53	3,632.48	0.98	0.65
	Tail spine	1.00	11265	18072	3,408.11	3,401.26	0.62	0.98
F4	Head spine	1.00	11813	21833	3,296.61	3,294.12	0.98	0.66
	Tail spine	1.00	5359	10355	3,111.48	3,101.43	0.99	0.98
F5	Head spine	1.00	7366	13477	2,451.79	2,451.35	0.63	0.56
	Tails pine	1.00	7322	9952	2,274.57	2,270.67	0.75	0.95

950   Diagnostics are presented for the Bayesian mixed-effects models fit separately for each combination of experimental
951   generation (F0–F5) and trait (headspine, tailspine). Model convergence was assessed using the maximum potential
952   scale reduction factor (Max Rhat), which should be approximately 1.0, and the minimum effective sample sizes for the
953   bulk (Min Bulk ESS) and tails (Min Tail ESS) of the posterior distributions, with higher values indicating more reliable
954   estimates. Model fit and predictive accuracy were compared using the Leave-One-Out Information Criterion (LOOIC)
955   and the Widely Applicable Information Criterion (WAIC); lower values suggest better out-of-sample fit. The probability
956   of direction, p(Direction), represents the posterior probability that the treatment effect is positive. The Marginal R² is the
957   proportion of variance explained by the model's fixed effects. All models demonstrated convergence.

970
971
972
973   **Supplemental Table 2: Linear estimates for headspine and tailspine length**

Spine	Generation	Est. Difference (µm)	Std. Error	R² marginal	R² conditional	P value
Head Spine	F0	68.98	27.19	58.4%	59.7%	**< 0.05**
	F3	37.02	17.61	51.4%	65%	**< 0.05**
	F4	43.81	20.19	47.7%	65.9%	**< 0.05**
	F5	-5.79	21.41	50.7%	52.6%	0.789
Tail Spine	F0	114.73	31.93	93.4%	94.2%	**< 0.01**
	F3	8.19	18.28	95%	98.1%	0.655
	F4	85.45	34.14	90.3%	98.4%	**< 0.05**
	F5	-24.21	39.07	93.1%	96.6%	0.539

974   Linear mixed-effects model results for headspine and tailspine length across generations. Models were fit identically
975   to Bayesian models for comparison. Treatment effects reach significance ($p < 0.05$) in F0, F3, and F4 but not F5 for
976   headspine, and in F0 and F4 for tailspine, consistent with Bayesian results. See main text for Bayesian results.
977
978

Supplemental Table 3: Goodness-of-fit and diagnostic statistics for linear mixed-effects models.

Spine	Generation	Est. (µm)	Std. Error	t-value	Marginal R²	Conditional R²	AIC	BIC	Log-Likelihood
Headspine	F0	68.98	27.19	2.54	58.4%	59.7%	881.0	892.5	-435.5
	F3	37.02	17.61	2.10	51.4%	65%	3651.0	3669.8	-1820.5
	F4	43.81	20.19	2.17	47.7%	65.9%	3316.0	3334.3	-1653.0
	F5	-5.79	21.41	-0.27	50.7%	52.6%	2454.2	2470.9	-1222.1
Tailspine	F0	114.73	31.93	3.59	93.4%	94.2%	870.5	882.0	-430.3
	F3	8.19	18.28	0.45	95%	98.1%	3485.1	3503.9	-1737.6
	F4	85.45	34.14	2.50	90.3%	98.4%	3225.7	3244.0	-1607.9
	F5	-24.21	39.07	-0.62	93.1%	96.6%	2427.1	2443.8	-1208.5

Models were fit independently for each combination of experimental generation (F0, F3, F4, F5) and morphological trait (headspine, tailspine). The estimate represents the effect of treatment in µm. Std. error indicates the standard error, and the t-values measures the difference between the sample and population means. The proportion of variance explained by the models is presented as Marginal R² (variance explained by fixed effects alone) and Conditional R² (variance explained by both fixed and random effects). Model fit was assessed using the Akaike Information Criterion (AIC) and the Bayesian Information Criterion (BIC), where lower values indicate a better trade-off between model fit and complexity.

991 **Supplemental Table 4: Generation-specific growth rate analysis.**

Generation	Trait	Estimate	SE	t value	P value
F0	Body Length	-25.592741	31.10285	-0.82284239	0.413
	Total Body Size	-7.091392	79.88130	-0.08877412	0.929
	Headspine	22.142802	22.23214	0.99598168	0.322
	Tailspine	3.423961	32.21077	0.10629862	0.916
F3	Body Length	16.060122	20.28191	0.79184464	0.429
	Total Body Size	56.924441	61.25339	0.92932721	0.361
	Headspine	-1.379195	16.10992	-0.08561156	0.932
	Tailspine	31.772759	18.21731	1.74409681	0.082
F4	Body Length	-16.610240	14.82382	-1.12051036	0.263
	Total Body Size	-39.774020	33.47196	-1.18827879	0.235
	Headspine	-9.445478	7.87350	-1.19965430	0.231
	Tailspine	-18.663554	14.63455	-1.27530766	0.203
F5	Body Length	-14.346394	27.29111	-0.52568011	0.599
	Total Body Size	-13.165803	66.40782	-0.19825680	0.843
	Headspine	-14.225783	13.31968	-1.06802748	0.286
	Tailspine	2.735027	25.45705	0.10743693	0.914

992 Mixed-effects models tested treatment effects on morphological growth rates (Δ from previous timepoint) separately
993 for each generation. Model: delta ~ treatment + age + (1|individual). Each row represents a unique combination of
994 generation and trait. Significant treatment effects (p < 0.05) in specific generations would indicate generation-
995 dependent differences in developmental strategy, however no significant differences were detected.
996
997

998   **Supplemental Table 5: Retrospective power analysis for F5 generation.**

Spine	Effect Sizes (µm)			F5 Sample Sizes		Power Analysis			
	F3 Effect	F4 Effect	Avg Effect (µm)	F5 n Control	F5 n Induced	Cohen's d	Power	Needed n (80%)	Est. Power
Head Spine	37.0	43.8	40.4	8	41	0.57	0.30	49	**29.8%**
Tail Spine	8.2	85.4	46.8	8	41	0.96	0.66	19	**66.4%**

999   Effect sizes for head and tail spine traits were averaged across F3 and F4 generations and used as benchmarks to
1000   estimate statistical power for detecting similar effects in F5 given actual sample sizes (n = 8 control, n = 41 induced).
1001   Cohen's *d* represents standardized effect size (treatment effect divided by pooled SD). Columns report observed
1002   effect sizes (µm), sample sizes, calculated power, and the estimated sample size required to achieve 80% power.

1003
1004
1005 **Supplemental Figure 1. Ontogenetic growth trajectories for Daphnia body length across**
1006 **treatments show high correlation between age and body length.** Scatterplot of individual
1007 body length measurements (μm) as a function of age (days) for control (open circles) and
1008 induced (filled circles) treatment across all generations. The solid line shows the overall fitted
1009 growth curve (Pearson's r =0.85) across all data. Growth patterns are similar between treatment,
1010 indicating that environmental induction principally affects defensive traits rather than somatic
1011 traits in this clone.
1012

1013
1014 **Supplemental Figure 2**: **Bayesian model diagnostic plots for the headspine trait across**
1015 **generations.** Each row represents an independent model. The panels display posterior predictive
1016 checks (model fit), residuals vs. fitted values (homoscedasticity), MCMC traceplots
1017 (convergence), and Quantile-Quantile (QQ) plots (normality of residuals). The successful
1018 convergence of the MCMC chains and the adequate fit shown by the predictive checks and
1019 residual plots support the validity of the models for inference.
1020
1021

1022
1023 **Supplemental Figure 3: Bayesian model diagnostic plots for the tailspine trait across**
1024 **generations.** Each row represents an independent model. Plots include posterior predictive
1025 checks (model fit), residuals vs. fitted values (homoscedasticity), MCMC traceplots
1026 (convergence), and Quantile-Quantile (QQ) plots (normality of residuals). The posterior
1027 predictive checks show good overlap between observed and simulated data, and the traceplots
1028 indicate stable, well-mixed chains. The residuals show no major patterns and are approximately
1029 normally distributed, supporting the validity of the models.
1030
1031

1032
1033 **Supplemental Figure 4: Predicted spine lengths by generation and treatment from linear**
1034 **mixed-effects models.** Predicted mean headspine length (top row) and tailspine length (bottom
1035 row) for Daphnia in control and induced individuals across generations (F0, F3, F4, F5). Points
1036 represent model-predicted means ± standard error from linear mixed-effects models.

1037
1038

1039
1040 **Supplemental Figure 5. Transgenerational patterns of headspine and tailspine length in**
1041 **Daphnia following ancestral predator cue exposure.** Mean (± standard error) headspine (top
1042 panels) and tailspine (bottom panels) values are shown for control and induced individuals across
1043 four non-exposed generations (F3–F5) and the originally exposed generation (F0). Each panel
1044 displays pairwise comparisons between control and induced groups for the indicated generation.
1045 Significant induction of both headspine and tailspine length is evident in predator-cue–induced
1046 F0 animals, with differential persistence of these defensive traits in subsequent generations.
1047 These data illustrate the finite and dynamic pattern of transgenerational plasticity in response to
1048 environmental cues. Note: this data is included for transparency, but does not include correction
1049 for body sizes, and violates independence assumptions as there are repeated measures.
1050

1051
1052 **Supplemental Figure 6: Model diagnostics for body length Bayesian models across**
1053 **generations.** For each generation: posterior predictive check (top left), MCMC traceplots (top
1054 right), residuals vs fitted (bottom left), and QQ plot (bottom right).
1055

References

Adamczuk, M. (2020). Population dynamics and life history traits of Daphnia magna across thermal regimes of environments. *Science of The Total Environment*, *723*, 137963. https://doi.org/10.1016/J.SCITOTENV.2020.137963

Agrawal, A. A., Laforsch, C., & Tollrian, R. (1999a). Transgenerational induction of defences in animals and plants. *Nature 1999 401:6748*, *401*(6748), 60–63. https://doi.org/10.1038/43425

Agrawal, A. A., Laforsch, C., & Tollrian, R. (1999b). Transgenerational induction of defences in animals and plants. *Nature 1999 401:6748*, *401*(6748), 60–63. https://doi.org/10.1038/43425

Agrelius, T. C., & Dudycha, J. L. (2025). Maternal effects in the model system Daphnia: the ecological past meets the epigenetic future. *Heredity*, *134*(2), 142–154. https://doi.org/10.1038/S41437-024-00742-W

Alfonso, S., Blanc, M., Joassard, L., Keiter, S. H., Munschy, C., Loizeau, V., Bégout, M. L., & Cousin, X. (2019). Examining multi- and transgenerational behavioral and molecular alterations resulting from parental exposure to an environmental PCB and PBDE mixture. *Aquatic Toxicology (Amsterdam, Netherlands)*, *208*, 29–38. https://doi.org/10.1016/J.AQUATOX.2018.12.021

Allaire, J., Teague, C., Scheidegger, C., Yihui, X., Dervieux, C., & Woodhull, G. (2025). *Quarto*. https://github.com/quarto-dev/quarto-cli

Allaire, J., Xie, Y., Dervieux, C., Grolemund, Garrett., McPherson, J., Luraschi, J., Ushey, K., Atkins, A., Wickham, H., Cheng, J., Chang, W., & Iannnone, R. (2019). *rmarkdown: Dynamic Documents for R* (p. 303). CRC Press, Taylor & Francis Group. https://github.com/rstudio/rmarkdown

Angilletta, M. J. (2009). Thermal Adaptation: A Theoretical and Empirical Synthesis. *Thermal Adaptation: A Theoretical and Empirical Synthesis*, 1–302. https://doi.org/10.1093/ACPROF:OSO/9780198570875.001.1

Anway, M. D., Cupp, A. S., Uzumcu, N., & Skinner, M. K. (2005). Epigenetic transgenerational actions of endocrine disruptors and male fertility. *Science (New York, N.Y.)*, *308*(5727), 1466–1469. https://doi.org/10.1126/SCIENCE.1108190

Asselman, J., De Coninck, D. I. M., Pfrender, M. E., & De Schamphelaere, K. A. C. (2016). Gene body methylation patterns in daphnia are associated with gene family size. *Genome Biology and Evolution*, *8*(4), 1185–1196. https://doi.org/10.1093/gbe/evw069

Auld, J. R., Agrawal, A. A., & Relyea, R. A. (2010). Re-evaluating the costs and limits of adaptive phenotypic plasticity. *Proceedings. Biological Sciences*, *277*(1681), 503–511. https://doi.org/10.1098/RSPB.2009.1355

Baludo, M. Y., Octorina, P., Beckerman, A., & Straile, D. (2024). Antipredator responses of three Daphnia species within the D. longispina species complex to two invertebrate predators. *Ecology and Evolution*, *14*(1), e10841. https://doi.org/10.1002/ECE3.10841;REQUESTEDJOURNAL:JOURNAL:20457758

Banerjee, A., & Roychoudhury, A. (2017). Epigenetic regulation during salinity and drought stress in plants: Histone modifications and DNA methylation. *Plant Gene*, *11*, 199–204. https://doi.org/10.1016/J.PLGENE.2017.05.011

Batabyal, A. (2023). Predator-prey systems as models for integrative research in biology: the value of a non-consumptive effects framework. *The Journal of Experimental Biology*, *226*(19). https://doi.org/10.1242/JEB.245851

Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2015a). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*(1). https://doi.org/10.18637/JSS.V067.I01

Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2015b). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *67*(1), 1–48. https://doi.org/10.18637/JSS.V067.I01

Baugh, L. R., & Day, T. (2020). Nongenetic inheritance and multigenerational plasticity in the nematode C. elegans. *ELife*, *9*, e58498. https://doi.org/10.7554/ELIFE.58498

Beaman, J. E., White, C. R., & Seebacher, F. (2016). Evolution of Plasticity: Mechanistic Link between Development and Reversible Acclimation. *Trends in Ecology & Evolution*, *31*(3), 237–249. https://doi.org/10.1016/J.TREE.2016.01.004

Beklioglu, M., Telli, M., & Gozen, A. G. (2006). Fish and mucus-dwelling bacteria interact to produce a kairomone that induces diel vertical migration in Daphnia. *Freshwater Biology*, *51*(12), 2200–2206. https://doi.org/10.1111/j.1365-2427.2006.01642.x

Bell, A. M., & Hellmann, J. K. (2019). An Integrative Framework for Understanding the Mechanisms and Multigenerational Consequences of Transgenerational Plasticity. *Annual Review of Ecology, Evolution, and Systematics*, *50*(Volume 50, 2019), 97–118. https://doi.org/10.1146/ANNUREV-ECOLSYS-110218-024613/CITE/REFWORKS

Bird, A. (2024). Transgenerational epigenetic inheritance: a critical perspective. *Frontiers in Epigenetics and Epigenomics*, *2*, 1434253. https://doi.org/10.3389/FREAE.2024.1434253

Bonduriansky, R., & Day, T. (2009). Nongenetic inheritance and its evolutionary implications. *Annual Review of Ecology, Evolution, and Systematics*, *40*(Volume 40, 2009), 103–125. https://doi.org/10.1146/ANNUREV.ECOLSYS.39.110707.173441/CITE/REFWORKS

Botero, C. A., Weissing, F. J., Wright, J., & Rubenstein, D. R. (2015). Evolutionary tipping points in the capacity to adapt to environmental change. *Proceedings of the National Academy of Sciences of the United States of America*, *112*(1), 184–189. https://doi.org/10.1073/PNAS.1408589111

Bracis, C., & Wirsing, A. J. (2021). Prey Foraging Behavior After Predator Introduction Is Driven by Resource Knowledge and Exploratory Tendency. *Frontiers in Ecology and Evolution*, *9*, 698370. https://doi.org/10.3389/FEVO.2021.698370/BIBTEX

Bradshaw, A. D. (1965). Evolutionary Significance of Phenotypic Plasticity in Plants. *Advances in Genetics*, *13*(C), 115–155. https://doi.org/10.1016/S0065-2660(08)60048-6

Bürkner, P. C. (2017). brms: An R package for Bayesian multilevel models using Stan. *Journal of Statistical Software*, *80*. https://doi.org/10.18637/JSS.V080.I01

Bürkner, P., Gabry, J., & Vehtari, A. (2025). *posterior: Tools for Working with Posterior Distributions* (R package version 1.6.1). https://mc-stan.org/posterior/

Carone, B. R., Fauquier, L., Habib, N., Shea, J. M., Hart, C. E., Li, R., Bock, C., Li, C., Gu, H., Zamore, P. D., Meissner, A., Weng, Z., Hofmann, H. A., Friedman, N., & Rando, O. J. (2010). Paternally induced transgenerational environmental reprogramming of metabolic gene expression in mammals. *Cell*, *143*(7), 1084–1096. https://doi.org/10.1016/j.cell.2010.12.008

Casadesús, J. (2016). Bacterial DNA methylation and methylomes. *Advances in Experimental Medicine and Biology*, *945*, 35–61. https://doi.org/10.1007/978-3-319-43624-1_3/FIGURES/3

Champely, S., Ekstrom, C., Dalgaard, P., Gill, J., Anandkumar, A., Ford, C., Volcic, R., & De Rosario, H. (2020). *Package "pwr" Title Basic Functions for Power Analysis*. https://github.com/heliosdrm/pwr

Corl, A., Bi, K., Luke, C., Challa, A. S., Stern, A. J., Sinervo, B., & Nielsen, R. (2018). The Genetic Basis of Adaptation following Plastic Changes in Coloration in a Novel Environment. *Current Biology*, *28*(18), 2970-2977.e7. https://doi.org/10.1016/j.cub.2018.06.075

Danchin, É., & Wagner, R. H. (2010). Inclusive heritability: combining genetic and non-genetic information to study animal behavior and culture. *Oikos*, *119*(2), 210–218. https://doi.org/10.1111/J.1600-0706.2009.17640.X

Dao, T. S., Vo, T. M. C., Wiegand, C., Bui, B. T., & Dinh, K. V. (2018). Transgenerational effects of cyanobacterial toxins on a tropical micro-crustacean Daphnia lumholtzi across three generations. *Environmental Pollution (Barking, Essex : 1987)*, *243*(Pt B), 791–799. https://doi.org/10.1016/J.ENVPOL.2018.09.055

Davidowitz, G. (2009). Ecological Developmental Biology: Integrating Epigenetics, Medicine and Evolution. Scott F. Gilbert and David Epel. *Integrative and Comparative Biology*, *49*(6), 727–729. https://doi.org/10.1093/ICB/ICP106

Dejeux, L., Saclier, N., Tariel-Adam, J., Hoareau, M., Lefébure, T., Konecny, L., Plénet, S., & Luquet, E. (2025a). Transcriptomic basis of within- and trans-generational predator-induced plasticity in the freshwater snail Physa acuta. *Heredity 2025 134:7*, *134*(7), 439–449. https://doi.org/10.1038/s41437-025-00775-9

Dejeux, L., Saclier, N., Tariel-Adam, J., Hoareau, M., Lefébure, T., Konecny, L., Plénet, S., & Luquet, E. (2025b). Transcriptomic basis of within- and trans-generational predator-induced plasticity in the freshwater snail Physa acuta. *Heredity*, *134*(7), 439–449. https://doi.org/10.1038/S41437-025-00775-9;TECHMETA

DeWitt, T. J., Sih, A., & Wilson, D. S. (1998). Costs and limits of phenotypic plasticity. *Trends in Ecology and Evolution*, *13*(2), 77–81. https://doi.org/10.1016/S0169-5347(97)01274-3

Dias, B. G., & Ressler, K. J. (2014). Parental olfactory experience influences behavior and neural structure in subsequent generations. *Nature Neuroscience*, *17*(1), 89–96. https://doi.org/10.1038/NN.3594

Dieckmann, U., Heino, M., & Parvinen, K. (2006). The adaptive dynamics of function-valued traits. *Journal of Theoretical Biology*, *241*(2), 370–389. https://doi.org/10.1016/j.jtbi.2005.12.002

Donelson, J. M., Salinas, S., Munday, P. L., & Shama, L. N. S. (2018). Transgenerational plasticity and climate change experiments: Where do we go from here? *Global Change Biology*, *24*(1), 13–34. https://doi.org/10.1111/GCB.13903

Dzialowski, A. R., Lennon, J. T., O'Brien, W. J., & Smith, V. H. (2003). Predator-induced phenotypic plasticity in the exotic cladoceran Daphnia lumholtzi. *Freshwater Biology*, *48*(9), 1593–1602. https://doi.org/10.1046/J.1365-2427.2003.01111.X

Ebert, D. (2022). Daphnia as a versatile model system in ecology and evolution. *EvoDevo 2022 13:1*, *13*(1), 1–13. https://doi.org/10.1186/S13227-022-00199-0

Engel, K., Schreder, T., & Tollrian, R. (2014a). Morphological defences of invasive Daphnia lumholtzi protect against vertebrate and invertebrate predators. *Journal of Plankton Research*, *36*(4), 1140–1145. https://doi.org/10.1093/PLANKT/FBU023

Engel, K., Schreder, T., & Tollrian, R. (2014b). Morphological defences of invasive Daphnia lumholtzi protect against vertebrate and invertebrate predators. *Journal of Plankton Research*, *36*(4), 1140–1145. https://doi.org/10.1093/PLANKT/FBU023

Engel, K., & Tollrian, R. (2009). Inducible defences as key adaptations for the successful invasion of Daphnia lumholtzi in North America? *Proceedings of the Royal Society B: Biological Sciences*, *276*(1663), 1865. https://doi.org/10.1098/RSPB.2008.1861

Feinberg, A. P. (2007). Phenotypic plasticity and the epigenetics of human disease. *Nature*, *447*(7143), 433–440. https://doi.org/10.1038/NATURE05919

Feinberg, A. P., & Irizarry, R. A. (2010). Stochastic epigenetic variation as a driving force of development, evolutionary adaptation, and disease. *Proceedings of the National Academy of Sciences*, *107*(suppl_1), 1757–1764. https://doi.org/10.1073/PNAS.0906183107

Feiner, N., Radersma, R., Vasquez, L., Ringnér, M., Nystedt, B., Raine, A., Tobi, E. W., Heijmans, B. T., & Uller, T. (2022). Environmentally induced DNA methylation is inherited across generations in an aquatic keystone species. *IScience*, *25*(5), 104303. https://doi.org/10.1016/J.ISCI.2022.104303

Fitz-James, M. H., & Cavalli, G. (2022). Molecular mechanisms of transgenerational epigenetic inheritance. *Nature Reviews Genetics 2022 23:6*, *23*(6), 325–341. https://doi.org/10.1038/s41576-021-00438-5

Forsman, A. (2015). Rethinking phenotypic plasticity and its consequences for individuals, populations and species. *Heredity*, *115*(4), 276–284. https://doi.org/10.1038/HDY.2014.92

Freeberg, T. M., & Lucas, J. R. (2009). Pseudoreplication is (still) a problem. *Journal of Comparative Psychology (Washington, D.C. : 1983)*, *123*(4), 450–451. https://doi.org/10.1037/A0017031

Funke, E., Webb, L., & Wolinska, J. (2024). The effect of microplastics on Daphnia fitness – Systematic review and meta-analysis. *Freshwater Biology*, *69*(4), 461–476. https://doi.org/10.1111/FWB.14228;WEBSITE:WEBSITE:PERICLES;JOURNAL:JOURNAL:13652427;WGROUP:STRING:PUBLICATION

Furrow, R. E., & Feldman, M. W. (2014). Genetic variation and the evolution of epigenetic regulation. *Evolution; International Journal of Organic Evolution*, *68*(3), 673–683. https://doi.org/10.1111/EVO.12225

Gabry, J., Simpson, D., Vehtari, A., Betancourt, M., & Gelman, A. (2019). Visualization in Bayesian Workflow. *Journal of the Royal Statistical Society Series A: Statistics in Society*, *182*(2), 389–402. https://doi.org/10.1111/rssa.12378

Gapp, K., Jawaid, A., Sarkies, P., Bohacek, J., Pelczar, P., Prados, J., Farinelli, L., Miska, E., & Mansuy, I. M. (2014). Implication of sperm RNAs in transgenerational inheritance of the effects of early trauma in mice. *Nature Neuroscience*, *17*(5), 667–669. https://doi.org/10.1038/NN.3695

Geerts, A. N., Vanoverbeke, J., Vanschoenwinkel, B., Van Doorslaer, W., Feuchtmayr, H., Atkinson, D., Moss, B., Davidson, T. A., Sayer, C. D., & De Meester, L. (2015). Rapid evolution of thermal tolerance in the water flea Daphnia. *Nature Climate Change*, *5*(7), 665–668. https://doi.org/10.1038/NCLIMATE2628;SUBJMETA

Gelman, A., & Hill, J. (2006). Data Analysis Using Regression and Multilevel/Hierarchical Models. *Data Analysis Using Regression and Multilevel/Hierarchical Models*. https://doi.org/10.1017/CBO9780511790942

Geoghegan, J. L., & Spencer, H. G. (2012). Population-epigenetic models of selection. *Theoretical Population Biology*, *81*(3), 232–242. https://doi.org/10.1016/j.tpb.2011.08.001

Ghalambor, C. K., McKay, J. K., Carroll, S. P., & Reznick, D. N. (2007). Adaptive versus non-adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. *Functional Ecology*, *21*(3), 394–407. https://doi.org/10.1111/J.1365-2435.2007.01283.X

Graeve, A., Janßen, M., Villalba de la Pena, M., Tollrian, R., & Weiss, L. C. (2021). Higher, Faster, Better: Maternal Effects Shorten Time Lags and Increase Morphological Defenses in Daphnia lumholtzi Offspring Generations. *Frontiers in Ecology and Evolution*, *9*, 324. https://doi.org/10.3389/FEVO.2021.637421/BIBTEX

Greer, E. L., Maures, T. J., Ucar, D., Hauswirth, A. G., Mancini, E., Lim, J. P., Benayoun, B. A., Shi, Y., & Brunet, A. (2011). Transgenerational epigenetic inheritance of longevity in Caenorhabditis elegans. *Nature 2011 479:7373*, *479*(7373), 365–371. https://doi.org/10.1038/nature10572

Grishkevich, V., & Yanai, I. (2013). The genomic determinants of genotype × environment interactions in gene expression. *Trends in Genetics*, *29*(8), 479–487. https://doi.org/10.1016/j.tig.2013.05.006

Grosenbach, G. (2021). *A High Contrast Theme for ggplot - Topfunky Blog*. https://topfunky.com/2021/ggplot-high-contrast-theme-gghighcontrast/

Gu, L., Xu, Y., Yang, T., Qin, S., Zhang, L., Sun, Y., Huang, Y., & Yang, Z. (2021). Predator-induced allometric changes in the tail spine length of Daphnia: a distinct resource allocation strategy. *Journal of Plankton Research*, *43*(6), 884–893. https://doi.org/10.1093/PLANKT/FBAB063

Gustafsson, S., & Hansson, L. A. (2004). Development of tolerance against toxic cyanobacteria in Daphnia. *Aquatic Ecology*, *38*(1), 37–44. https://doi.org/10.1023/B:AECO.0000020985.47348.5E/METRICS

Hales, N. R., Schield, D. R., Andrew, A. L., Card, D. C., Walsh, M. R., & Castoe, T. A. (2017). Contrasting gene expression programs correspond with predator-induced phenotypic plasticity within and across generations in Daphnia. *Molecular Ecology*, *26*(19), 5003–5015. https://doi.org/10.1111/mec.14213

Harris, K. D. M., Bartlett, N. J., & Lloyd, V. K. (2012a). Daphnia as an Emerging Epigenetic Model Organism. *Genetics Research International*, *2012*, 1–8. https://doi.org/10.1155/2012/147892

Harris, K. D. M., Bartlett, N. J., & Lloyd, V. K. (2012b). Daphnia as an Emerging Epigenetic Model Organism. *Genetics Research International*, *2012*, 1–8. https://doi.org/10.1155/2012/147892

Heard, E., & Martienssen, R. A. (2014). Transgenerational epigenetic inheritance: Myths and mechanisms. *Cell*, *157*(1), 95–109. https://doi.org/10.1016/j.cell.2014.02.045

Hearn, J., Pearson, M., Blaxter, M., Wilson, P. J., & Little, T. J. (2019). Genome-wide methylation is modified by caloric restriction in Daphnia magna. *BMC Genomics*, *20*(1). https://doi.org/10.1186/s12864-019-5578-4

Hebert, P. D. N., & Grewe, P. M. (1985). Chaoborus-induced shifts in the morphology of Daphnia ambigua1. *Limnology and Oceanography*, *30*(6), 1291–1297. https://doi.org/10.4319/LO.1985.30.6.1291

Hendry, A. P. (2016). Key Questions on the Role of Phenotypic Plasticity in Eco-Evolutionary Dynamics. *The Journal of Heredity*, *107*(1), 25–41. https://doi.org/10.1093/JHERED/ESV060

Herman, J. J., & Sultan, S. E. (2011). Adaptive transgenerational plasticity in plants: Case studies, mechanisms, and implications for natural populations. *Frontiers in Plant Science*, *2*(DEC), 18242. https://doi.org/10.3389/FPLS.2011.00102/BIBTEX

Ho, D. H., & Burggren, W. W. (2010). Epigenetics and transgenerational transfer: a physiological perspective. *The Journal of Experimental Biology*, *213*(1), 3–16. https://doi.org/10.1242/JEB.019752

Hope, R. M. (2022). *Rmisc: Ryan Miscellaneous*. https://cran.r-project.org/web/packages/Rmisc/index.html

Horstmann, M., Luncke, J., Weiss, L. C., & Tollrian, R. (2023). Long-term analysis and effectiveness of inducible behavioural and morphological traits of Daphnia longicephala in response to Notonecta. *Fundamental and Applied Limnology*, *196*(3–4), 287–299. https://doi.org/10.1127/FAL/2022/1472

Horstmann, M., Tollrian, R., & Weiss, L. C. (2021). Thwarting predators? A three-dimensional perspective of morphological alterations in the freshwater crustacean Daphnia. *PLOS ONE*, *16*(7), e0254263. https://doi.org/10.1371/JOURNAL.PONE.0254263

Horstmann, M., Weiss, L. C., & Tollrian, R. (2022). Specific Turbulence- and Chaoborus-Induced Morphotypes Affect the Streamlining Properties of Daphnia cucullata. *Frontiers in Ecology and Evolution*, *9*, 825765. https://doi.org/10.3389/FEVO.2021.825765/BIBTEX

Hurlbert, S. H. (1984). Pseudoreplication and the Design of Ecological Field Experiments. *Ecological Monographs*, *54*(2), 187–211. https://doi.org/10.2307/1942661

Jablonka, Eva., & Lamb, M. J. . (20062). *Evolution in four dimensions : genetic, epigenetic, behavioral, and symbolic variation in the history of life*. MIT Press.

Jeremias, G., Barbosa, J., Marques, S. M., De Schamphelaere, K. A. C., Van Nieuwerburgh, F., Deforce, D., Gonçalves, F. J. M., Pereira, J. L., & Asselman, J. (2018). Transgenerational Inheritance of DNA Hypomethylation in Daphnia magna in Response to Salinity Stress. *Environmental Science & Technology*, *52*(17), 10114–10123. https://doi.org/10.1021/ACS.EST.8B03225

Johnson, J. L., & Havel, J. E. (2001). Competition Between Native and Exotic Daphnia: In situ Experiments. *Journal of Plankton Research*, *23*(4), 373–387. https://doi.org/10.1093/PLANKT/23.4.373

Kaneshiro, K. R., Rechtsteiner, A., & Strome, S. (2019). Sperm-inherited H3K27me3 impacts offspring transcription and development in C. elegans. *Nature Communications*, *10*(1), 1–9. https://doi.org/10.1038/S41467-019-09141-W;TECHMETA

Kilham, S. S., Kreeger, D. A., Lynn, S. G., Goulden, C. E., & Herrera, L. (1998). COMBO: a defined freshwater culture medium for algae and zooplankton. *Hydrobiologia 1998 377:1*, *377*(1), 147–159. https://doi.org/10.1023/A:1003231628456

Kim, J., & Choi, J. (2023). Trans- and Multigenerational Effects of Isothiazolinone Biocide CMIT/MIT on Genotoxicity and Epigenotoxicity in Daphnia magna. *Toxics*, *11*(4), 388. https://doi.org/10.3390/TOXICS11040388/S1

Kim, S. W., Kwak, J. Il, & An, Y. J. (2013). Multigenerational Study of Gold Nanoparticles in Caenorhabditis elegans: Transgenerational Effect of Maternal Exposure. *Environmental Science and Technology*, *47*(10), 5393–5399. https://doi.org/10.1021/ES304511Z

Kimmel, K., Avolio, M. L., & Ferraro, P. J. (2023). Empirical evidence of widespread exaggeration bias and selective reporting in ecology. *Nature Ecology & Evolution 2023 7:9*, *7*(9), 1525–1536. https://doi.org/10.1038/s41559-023-02144-3

Kingsolver, J. G., Hoekstra, H. E., Hoekstra, J. M., Berrigan, D., Vignieri, S. N., Hill, C. E., Hoang, A., Gibert, P., & Beerli, P. (2001). The strength of phenotypic selection in natural populations. *The American Naturalist*, *157*(3), 245–261. https://doi.org/10.1086/319193

Kishimoto, S., Uno, M., Okabe, E., Nono, M., & Nishida, E. (2017). Environmental stresses induce transgenerationally inheritable survival advantages via germline-to-soma communication in Caenorhabditis elegans. *Nature Communications 2017 8:1*, *8*(1), 1–12. https://doi.org/10.1038/ncomms14031

Kizilaslan, M., Braz, C. U., Townsend, J., Taylor, T., Crenshaw, T. D., & Khatib, H. (2025). Transgenerational Epigenetic and Phenotypic Inheritance Across Five Generations in Sheep. *International Journal of Molecular Sciences*, *26*(13), 6412. https://doi.org/10.3390/IJMS26136412/S1

Klibaner-Schiff, E., Simonin, E. M., Akdis, C. A., Cheong, A., Johnson, M. M., Karagas, M. R., Kirsh, S., Kline, O., Mazumdar, M., Oken, E., Sampath, V., Vogler, N., Wang, X., & Nadeau, K. C. (2024). Environmental exposures influence multigenerational epigenetic transmission. *Clinical Epigenetics 2024 16:1*, *16*(1), 145-. https://doi.org/10.1186/S13148-024-01762-3

Klosin, A., Casas, E., Hidalgo-Carcedo, C., Vavouri, T., & Lehner, B. (2017). Transgenerational transmission of environmental information in C. elegans. *Science (New York, N.Y.)*, *356*(6335), 320–323. https://doi.org/10.1126/SCIENCE.AAH6412

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in Linear Mixed Effects Models. *Journal of Statistical Software*, *82*(13), 1–26. https://doi.org/10.18637/JSS.V082.I13

Laforsch, C., & Tollrian, R. (2004). Embryological aspects of inducible morphological defenses in Daphnia. *Journal of Morphology*, *262*(3), 701–707. https://doi.org/10.1002/JMOR.10270

Laland, K. N., Uller, T., Feldman, M. W., Sterelny, K., Müller, G. B., Moczek, A., Jablonka, E., & Odling-Smee, J. (2015). The extended evolutionary synthesis: its structure, assumptions and predictions. *Proceedings of the Royal Society B: Biological Sciences*, *282*(1813). https://doi.org/10.1098/RSPB.2015.1019

Lande, R. (2009). Adaptation to an extraordinary environment by evolution of phenotypic plasticity and genetic assimilation. *Journal of Evolutionary Biology*, *22*(7), 1435–1446. https://doi.org/10.1111/J.1420-9101.2009.01754.X

Landy, J. A., Muh, S., Parajuli, B., Munch, S. B., & Walsh, M. R. (2024). Multi-generation exposure to a novel environmental signal shapes the expression of plasticity in offspring in resurrected Daphnia. *Biological Journal of the Linnean Society*, *143*(4). https://doi.org/10.1093/BIOLINNEAN/BLAE091

Lenth, R. V., & Piaskowski, J. (2017). *emmeans: Estimated Marginal Means, aka Least-Squares Means*. https://doi.org/10.32614/CRAN.package.emmeans

Levis, N. A., & Pfennig, D. W. (2019). Plasticity-led evolution: evaluating the key prediction of frequency-dependent adaptation. *Proceedings. Biological Sciences*, *286*(1897). https://doi.org/10.1098/RSPB.2018.2754

Li, E., & Zhang, Y. (2014). DNA methylation in mammals. *Cold Spring Harbor Perspectives in Biology*, *6*(5). https://doi.org/10.1101/CSHPERSPECT.A019133

Liao, H., Gao, D., Junaid, M., Liu, S., Kong, C., Chen, X., Pan, T., Zheng, Q., Ai, W., Chen, G., & Wang, J. (2023). Parental exposure to polystyrene nanoplastics and di(2-ethylhexyl) phthalate induces transgenerational growth and reproductive impairments through

1371   bioaccumulation in Daphnia magna. *Science of The Total Environment*, *882*, 163657.
1372   https://doi.org/10.1016/J.SCITOTENV.2023.163657

1373 Liu, Z., Cai, M., Wu, D., Yu, P., Jiao, Y., Jiang, Q., & Zhao, Y. (2020). Effects of nanoplastics at
1374   predicted environmental concentration on Daphnia pulex after exposure through multiple
1375   generations. *Environmental Pollution*, *256*, 113506.
1376   https://doi.org/10.1016/J.ENVPOL.2019.113506

1377 Lüdecke, D., Ben-Shachar, M., Patil, I., Waggoner, P., & Makowski, D. (2021). performance:
1378   An R Package for Assessment, Comparison and Testing of Statistical Models. *Journal of
1379   Open Source Software*, *6*(60), 3139. https://doi.org/10.21105/JOSS.03139

1380 MacLeod, K. J., Monestier, C., Ferrari, M. C. O., McGhee, K. E., Sheriff, M. J., & Bell, A. M.
1381   (2022). Predator-induced transgenerational plasticity in animals: a meta-analysis. *Oecologia
1382   2022 200:3*, *200*(3), 371–383. https://doi.org/10.1007/S00442-022-05274-W

1383 Maggio, S. A., & Jenkins, J. J. (2022a). Multi- and Trans-Generational Effects on Daphnia
1384   Magna of Chlorpyrifos Exposures. *Environmental Toxicology and Chemistry*, *41*(4), 1054–
1385   1065. https://doi.org/10.1002/ETC.5283

1386 Maggio, S. A., & Jenkins, J. J. (2022b). Multi- and Trans-Generational Effects on Daphnia
1387   Magna of Chlorpyrifos Exposures. *Environmental Toxicology and Chemistry*, *41*(4), 1054–
1388   1065.
1389   https://doi.org/10.1002/ETC.5283;SUBPAGE:STRING:ABSTRACT;WEBSITE:WEBSIT
1390   E:PERICLES;ISSUE:ISSUE:DOI

1391 Martins, A., & Guilhermino, L. (2018a). Transgenerational effects and recovery of microplastics
1392   exposure in model populations of the freshwater cladoceran Daphnia magna Straus. *Science
1393   of The Total Environment*, *631–632*, 421–428.
1394   https://doi.org/10.1016/J.SCITOTENV.2018.03.054

1395 Martins, A., & Guilhermino, L. (2018b). Transgenerational effects and recovery of microplastics
1396   exposure in model populations of the freshwater cladoceran Daphnia magna Straus. *Science
1397   of The Total Environment*, *631–632*, 421–428.
1398   https://doi.org/10.1016/J.SCITOTENV.2018.03.054

1399 Moore, M. P., Whiteman, H. H., & Martin, R. A. (2019). A mother's legacy: the strength of
1400   maternal effects in animal populations. *Ecology Letters*, *22*(10), 1620–1628.
1401   https://doi.org/10.1111/ELE.13351;WGROUP:STRING:PUBLICATION

1402 Müller, M. F., Colomer, J., & Serra, T. (2018). Temperature-driven response reversibility and
1403   short-term quasi-acclimation of Daphnia magna. *PLoS ONE*, *13*(12).
1404   https://doi.org/10.1371/JOURNAL.PONE.0209705

1405 Mulligan, C. J., Quinn, E. B., Hamadmad, D., Dutton, C. L., Nevell, L., Binder, A. M., Panter-
1406   Brick, C., & Dajani, R. (2025). Epigenetic signatures of intergenerational exposure to
1407   violence in three generations of Syrian refugees. *Scientific Reports*, *15*(1), 1–13.
1408   https://doi.org/10.1038/S41598-025-89818-Z;SUBJMETA

1409 Nagano, M., & Yoshida, T. (2020a). Size-selective predation accounts for intra- and inter-
1410   specific variation of inducible morphological defense of Daphnia. *Ecosphere*, *11*(7),
1411   e03192. https://doi.org/10.1002/ECS2.3192

1412 Nagano, M., & Yoshida, T. (2020b). Size-selective predation accounts for intra- and inter-
1413   specific variation of inducible morphological defense of Daphnia. *Ecosphere*, *11*(7),
1414   e03192. https://doi.org/10.1002/ECS2.3192

Nettle, D., & Bateson, M. (2015). Adaptive developmental plasticity: what is it, how can we recognize it and when can it evolve? *Proceedings of the Royal Society B: Biological Sciences*, *282*(1812). https://doi.org/10.1098/RSPB.2015.1005

Neylan, I. P., Salgado, A. L., Vaidya, R. V., Guillory, C., & Kelly, M. W. (2025). Context matters: A meta-analysis of the variable impact of transgenerational and developmental plasticity on responses to stress. *Functional Ecology*, *39*(11), 3105–3123. https://doi.org/10.1111/1365-2435.70168;WEBSITE:WEBSITE:BESJOURNALS;REQUESTEDJOURNAL:JOURNAL:13652435;WGROUP:STRING:PUBLICATION

Niederhuth, C. E., Bewick, A. J., Ji, L., Alabady, M. S., Kim, K. Do, Li, Q., Rohr, N. A., Rambani, A., Burke, J. M., Udall, J. A., Egesi, C., Schmutz, J., Grimwood, J., Jackson, S. A., Springer, N. M., & Schmitz, R. J. (2016). Widespread natural variation of DNA methylation within angiosperms. *Genome Biology 2016 17:1*, *17*(1), 1–19. https://doi.org/10.1186/S13059-016-1059-0

Norouzitallab, P., Baruah, K., Vandegehuchte, M., Van Stappen, G., Catania, F., Vanden Bussche, J., Vanhaecke, L., Sorgeloos, P., & Bossier, P. (2014). Environmental heat stress induces epigenetic transgenerational inheritance of robustness in parthenogenetic Artemia model. *FASEB Journal : Official Publication of the Federation of American Societies for Experimental Biology*, *28*(8), 3552–3563. https://doi.org/10.1096/FJ.14-252049

Panacek, A., Prucek, R., Safarova, D., Dittrich, M., Richtrova, J., Benickova, K., Zboril, R., & Kvitek, L. (2011). Acute and chronic toxicity effects of silver nanoparticles (NPs) on Drosophila melanogaster. *Environmental Science & Technology*, *45*(11), 4974–4979. https://doi.org/10.1021/ES104216B

Pedersen, T. L. (2019). *patchwork: The Composer of Plots*. https://doi.org/10.32614/CRAN.package.patchwork

Petrusek, A., Tollrian, R., Schwenk, K., Haas, A., & Laforsch, C. (2009). A "crown of thorns" is an inducible defense that protects Daphnia against an ancient predator. *Proceedings of the National Academy of Sciences*, *106*(7), 2248–2252. https://doi.org/10.1073/PNAS.0808075106

Pfennig, D. W., Wund, M. A., Snell-Rood, E. C., Cruickshank, T., Schlichting, C. D., & Moczek, A. P. (2010). Phenotypic plasticity's impacts on diversification and speciation. *Trends in Ecology and Evolution*, *25*(8), 459–467. https://doi.org/10.1016/j.tree.2010.05.006

Pietrzak, B., Rabus, M., Religa, M., Laforsch, C., & Dańko, M. J. (2020). *Phenotypic plasticity of senescence in Daphnia under predation impact: no ageing acceleration when the perceived risk decreases with age*. https://doi.org/10.1098/rsos.191382

Pigliucci, M., Murren, C. J., & Schlichting, C. D. (2006). Phenotypic plasticity and evolution by genetic assimilation. *Journal of Experimental Biology*, *209*(12), 2362–2367. https://doi.org/10.1242/JEB.02070

Pigliucci, & Müller. (2010). *Evolution, the Extended Synthesis* .

Postit Team. (2025). *RStudio: Integrated Development Environment for R*. Postit Software. http://www.posit.co/.

Proulx, S. R., & Teotónio, H. (2017). What kind of maternal effects can be selected for in fluctuating environments? *American Naturalist*, *189*(6), E118–E137. https://doi.org/10.5061/DRYAD.58416

R Core Team. (2021). R: A language and environment for statisticalcomputing. *R Foundation for Statistical Computing*.

Radchuk, V., Reed, T., Teplitsky, C., van de Pol, M., Charmantier, A., Hassall, C., Adamík, P., Adriaensen, F., Ahola, M. P., Arcese, P., Miguel Avilés, J., Balbontin, J., Berg, K. S., Borras, A., Burthe, S., Clobert, J., Dehnhard, N., de Lope, F., Dhondt, A. A., … Kramer-Schadt, S. (2019). Adaptive responses of animals to climate change are most likely insufficient. *Nature Communications*, *10*(1), 1–14. https://doi.org/10.1038/S41467-019-10924-4;TECHMETA

Rechavi, O., Houri-Ze'Evi, L., Anava, S., Goh, W. S. S., Kerk, S. Y., Hannon, G. J., & Hobert, O. (2014). Starvation-induced transgenerational inheritance of small RNAs in C. elegans. *Cell*, *158*(2), 277–287. https://doi.org/10.1016/j.cell.2014.06.020

Remy, J. J. (2010). Stable inheritance of an acquired behavior in caenorhabditis elegans. *Current Biology*, *20*(20), R877–R878. https://doi.org/10.1016/j.cub.2010.08.013

Robichaud, N. F., Sassine, J., Beaton, M. J., & Lloyd, V. K. (2012). The Epigenetic Repertoire of Daphnia magna Includes Modified Histones . *Genetics Research International*, *2012*, 1–7. https://doi.org/10.1155/2012/174860

Romero-Mujalli, D., Fuchs, L. I. R., Haase, M., Hildebrandt, J. P., Weissing, F. J., & Revilla, T. A. (2024). Emergence of phenotypic plasticity through epigenetic mechanisms. *Evolution Letters*, *8*(4), 561–574. https://doi.org/10.1093/EVLETT/QRAE012

Salo, P., Banks, P. B., Dickman, C. R., & Korpimäki, E. (2010). Predator manipulation experiments: Impacts on populations of terrestrial vertebrate prey. *Ecological Monographs*, *80*(4), 531–546. https://doi.org/10.1890/09-1260.1;WEBSITE:WEBSITE:ESAJOURNALS;JOURNAL:JOURNAL:15577015;WGROUP:STRING:PUBLICATION

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. *Nature Methods 2012 9:7*, *9*(7), 676–682. https://doi.org/10.1038/nmeth.2019

Schlichting, C. D., & Levin, D. A. (1986). Phenotypic plasticity: an evolving plant character. *Biological Journal of the Linnean Society*, *29*(1), 37–47. https://doi.org/10.1111/J.1095-8312.1986.TB01769.X

Schwarzenberger, A., & Martin-Creuzburg, D. (2021). Daphnia's Adaptive Molecular Responses to the Cyanobacterial Neurotoxin Anatoxin-α Are Maternally Transferred. *Toxins*, *13*(5). https://doi.org/10.3390/TOXINS13050326

Scoville, A. G., & Pfrender, M. E. (2010). Phenotypic plasticity facilitates recurrent rapid adaptation to introduced predators. *Proceedings of the National Academy of Sciences of the United States of America*, *107*(9), 4260–4263. https://doi.org/10.1073/PNAS.0912748107

Seda, J., & Petrusek, A. (2011). Daphnia as a model organism in limnology and aquatic biology: some aspects of its reproduction and development. *Journal of Limnology*, *70*(2), 335–386. https://doi.org/10.4081/jlimnol.2011.335

Sentis, A., Bertram, R., Dardenne, N., Ramon-Portugal, F., Espinasse, G., Louit, I., Negri, L., Haeler, E., Ashkar, T., Pannetier, T., Cunningham, J. L., Grunau, C., Le Trionnaire, G., Simon, J. C., Magro, A., Pujol, B., Hemptinne, J. L., & Danchin, E. (2018). Evolution without standing genetic variation: change in transgenerational plastic response under

1505   persistent predation pressure. *Heredity, 121*(3), 266–281. https://doi.org/10.1038/s41437-
1506   018-0108-8
1507 Shahmohamadloo, R. S., Fryxell, J. M., & Rudman, S. M. (2025). Transgenerational epigenetic
1508   inheritance increases trait variation but is not adaptive. *Evolution, 79*(6), 1033–1043.
1509   https://doi.org/10.1093/EVOLUT/QPAF050
1510 Simons, A. M. (2009). Fluctuating natural selection accounts for the evolution of diversification
1511   bet hedging. *Proceedings of the Royal Society B: Biological Sciences, 276*(1664), 1987–
1512   1992. https://doi.org/10.1098/RSPB.2008.1920
1513 Skinner, M. K. (2008). What is an epigenetic transgenerational phenotype?: F3 or F2.
1514   *Reproductive Toxicology, 25*(1), 2–6. https://doi.org/10.1016/J.REPROTOX.2007.09.001
1515 Skinner, M. K., Haque, C. G. B. M., Nilsson, E., Bhandari, R., & McCarrey, J. R. (2013).
1516   Environmentally Induced Transgenerational Epigenetic Reprogramming of Primordial
1517   Germ Cells and the Subsequent Germ Line. *PLOS ONE, 8*(7), e66318.
1518   https://doi.org/10.1371/JOURNAL.PONE.0066318
1519 Smithson, M., Thorson, J. L. M., Sadler-Riggleman, I., Beck, D., Skinner, M. K., & Dybdahl, M.
1520   (2020). Between-Generation Phenotypic and Epigenetic Stability in a Clonal Snail. *Genome
1521   Biology and Evolution, 12*(9), 1604. https://doi.org/10.1093/GBE/EVAA181
1522 Song, J., Kim, C., Na, J., Sivri, N., Samanta, P., & Jung, J. (2022a). Transgenerational effects of
1523   polyethylene microplastic fragments containing benzophenone-3 additive in Daphnia
1524   magna. *Journal of Hazardous Materials, 436*, 129225.
1525   https://doi.org/10.1016/J.JHAZMAT.2022.129225
1526 Song, J., Kim, C., Na, J., Sivri, N., Samanta, P., & Jung, J. (2022b). Transgenerational effects of
1527   polyethylene microplastic fragments containing benzophenone-3 additive in Daphnia
1528   magna. *Journal of Hazardous Materials, 436*, 129225.
1529   https://doi.org/10.1016/J.JHAZMAT.2022.129225
1530 Sousa, F. D. R., Palaoro, A. V., Elmoor-Loureiro, L. M. A., & Kotov, A. A. (2017). Predicting
1531   the invasive potential of the cladoceran Daphnia lumholtzi Sars, 1885 (Crustacea:
1532   Cladocera: Daphniidae) in the Neotropics: Are generalists threatened and relicts protected
1533   by their life-history traits? *Journal of Limnology, 76*(2), 272–280.
1534   https://doi.org/10.4081/JLIMNOL.2016.1571
1535 St-Cyr, S., & McGowan, P. O. (2015). Programming of stress-related behavior and epigenetic
1536   neural gene regulation in mice offspring through maternal exposure to predator odor.
1537   *Frontiers in Behavioral Neuroscience, 9*(JUNE), 145810.
1538   https://doi.org/10.3389/FNBEH.2015.00145/BIBTEX
1539 Szűcs, G., Sója, A., Péter, M., Sárközy, M., Bruszel, B., Siska, A., Földesi, I., Szabó, Z., Janáky,
1540   T., Vígh, L., Balogh, G., & Csont, T. (2019). Prediabetes Induced by Fructose-Enriched
1541   Diet Influences Cardiac Lipidome and Proteome and Leads to Deterioration of Cardiac
1542   Function prior to the Development of Excessive Oxidative Stress and Cell Damage.
1543   *Oxidative Medicine and Cellular Longevity, 2019*. https://doi.org/10.1155/2019/3218275
1544 Tariel, J., Plénet, S., & Luquet, É. (2020a). Transgenerational Plasticity in the Context of
1545   Predator-Prey Interactions. *Frontiers in Ecology and Evolution, 8*, 319.
1546   https://doi.org/10.3389/FEVO.2020.548660/BIBTEX
1547 Tariel, J., Plénet, S., & Luquet, É. (2020b). Transgenerational plasticity of inducible defences:
1548   Combined effects of grand-parental, parental and current environments. *Ecology and
1549   Evolution, 10*(5), 2367–2376. https://doi.org/10.1002/ECE3.6046

Tariel, J., Plénet, S., & Luquet, É. (2020c). Transgenerational plasticity of inducible defences: Combined effects of grand-parental, parental and current environments. *Ecology and Evolution*, *10*(5), 2367. https://doi.org/10.1002/ECE3.6046

Tollrian, R., & Harvell, C. D. (1999). *The ecology and evolution of inducible defenses*. Princeton University Press, NJ.

Trijau, M., Asselman, J., Armant, O., Adam-Guillermin, C., De Schamphelaere, K. A. C., & Alonzo, F. (2018). Transgenerational DNA Methylation Changes in Daphnia magna Exposed to Chronic γ Irradiation. *Environmental Science and Technology*, *52*(7), 4331–4339. https://doi.org/10.1021/acs.est.7b05695

Tufto, J. (2015). Genetic evolution, plasticity, and bet-hedging as adaptive responses to temporally autocorrelated fluctuating selection: A quantitative genetic model. *Evolution*, *69*(8), 2034–2049. https://doi.org/10.1111/EVO.12716

Uller, T., Moczek, A. P., Watson, R. A., Brakefield, P. M., & Laland, K. N. (2018). Developmental Bias and Evolution: A Regulatory Network Perspective. *Genetics*, *209*(4), 949–966. https://doi.org/10.1534/GENETICS.118.300995

Uller, T., Nakagawa, S., & English, S. (2013). Weak evidence for anticipatory parental effects in plants and animals. *Journal of Evolutionary Biology*, *26*(10), 2161–2170. https://doi.org/10.1111/JEB.12212

Van Buskirk, J., & Steiner, U. K. (2009). The fitness costs of developmental canalization and plasticity. *Journal of Evolutionary Biology*, *22*(4), 852–860. https://doi.org/10.1111/J.1420-9101.2009.01685.X

Van Steenwyk, G., Roszkowski, M., Manuella, F., Franklin, T. B., & Mansuy, I. M. (2018). Transgenerational inheritance of behavioral and metabolic effects of paternal exposure to traumatic stress in early postnatal life: evidence in the 4th generation. *Environmental Epigenetics*, *4*(2), dvy023. https://doi.org/10.1093/EEP/DVY023

Vandegehuchte, M. B., De Coninck, D., Vandenbrouck, T., De Coen, W. M., & Janssen, C. R. (2010). Gene transcription profiles, global DNA methylation and potential transgenerational epigenetic effects related to Zn exposure history in Daphnia magna. *Environmental Pollution*, *158*(10), 3323–3329. https://doi.org/10.1016/j.envpol.2010.07.023

Vandegehuchte, M. B., Lemière, F., Vanhaecke, L., Vanden Berghe, W., & Janssen, C. R. (2010). Direct and transgenerational impact on Daphnia magna of chemicals with a known effect on DNA methylation. *Comparative Biochemistry and Physiology - C Toxicology and Pharmacology*, *151*(3), 278–285. https://doi.org/10.1016/J.CBPC.2009.11.007

Vehtari, A., Gelman, A., Simpson, D., Carpenter, B., & Bürkner, P.-C. (2021). Rank-Normalization, Folding, and Localization: An Improved Rˆ for Assessing Convergence of MCMC (with Discussion). *Bayesian Analysis*, *16*(2). https://doi.org/10.1214/20-BA1221

Via, S., Gomulkiewicz, R., De Jong, G., Scheiner, S. M., Schlichting, C. D., & Van Tienderen, P. H. (1995). Adaptive phenotypic plasticity: consensus and controversy. *Trends in Ecology & Evolution*, *10*(5), 212–217. https://doi.org/10.1016/S0169-5347(00)89061-8

Walsh, M. R., Castoe, T., Holmes, J., Packer, M., Biles, K., Walsh, M., Munch, S. B., & Post, D. M. (2016). Local adaptation in transgenerational responses to predators. *Proceedings of the Royal Society B: Biological Sciences*, *283*(1823). https://doi.org/10.1098/RSPB.2015.2271;PAGE:STRING:ARTICLE/CHAPTER

Walsh, M. R., Cooley, F., Biles, K., & Munch, S. B. (2014). Predator-induced phenotypic plasticity within- and across-generations: A challenge for theory? *Proceedings of the Royal Society B: Biological Sciences*, *282*(1798). https://doi.org/10.1098/rspb.2014.2205

1596 Wang, J., Li, M., Gu, L., Sun, Y., Zhu, X., & Yang, Z. (2024). Transgenerational Effects of
1597 Atrazine on Daphnia magna Based on Life-History Traits and Population Dynamics.
1598 *Bulletin of Environmental Contamination and Toxicology 2024 114:1*, *114*(1), 15-.
1599 https://doi.org/10.1007/S00128-024-03983-X

1600 Wang, M., Nie, Y., Liu, Y., Dai, H., Wang, J., Si, B., Yang, Z., Cheng, L., Liu, Y., Chen, S., &
1601 Xu, A. (2019). Transgenerational effects of diesel particulate matter on Caenorhabditis
1602 elegans through maternal and multigenerational exposure. *Ecotoxicology and*
1603 *Environmental Safety*, *170*, 635–643. https://doi.org/10.1016/J.ECOENV.2018.12.027

1604 Weiss, L. C., Leimann, J., & Tollrian, R. (2015). Predator-induced defences in Daphnia
1605 longicephala: location of kairomone receptors and timeline of sensitive phases to trait
1606 formation. *Journal of Experimental Biology*, *218*(18), 2918–2926.
1607 https://doi.org/10.1242/JEB.124552

1608 Werner, E. E., & Gilliam, J. F. (1984). The ontogenetic niche and species interactions in size-
1609 structured populations. *Annual Review of Ecology and Systematics. Vol. 15*, *15*(Volume
1610 15,), 393–425.
1611 https://doi.org/10.1146/ANNUREV.ES.15.110184.002141/CITE/REFWORKS

1612 Werner, M. S., Loschko, T., King, T., Reich, S., Theska, T., Franz-Wachtel, M., Macek, B., &
1613 Sommer, R. J. (2023). Histone 4 lysine 5/12 acetylation enables developmental plasticity of
1614 Pristionchus mouth form. *Nature Communications 2023 14:1*, *14*(1), 1–14.
1615 https://doi.org/10.1038/s41467-023-37734-z

1616 West-Eberhard, M. J. (2003). Developmental Plasticity and Evolution. *Developmental Plasticity*
1617 *and Evolution*. https://doi.org/10.1093/OSO/9780195122343.001.0001

1618 West-Eberhard, M. J. (2005). Developmental plasticity and the origin of species differences.
1619 *Proceedings of the National Academy of Sciences of the United States of America*, *102*
1620 *Suppl 1*(Suppl 1), 6543–6549. https://doi.org/10.1073/PNAS.0501844102

1621 West-Eberhard, M. Jane. (2020). *Developmental plasticity and evolution*.

1622 Wickham, H. (2016). ggplot2. *Ggplot2*. https://doi.org/10.1007/978-0-387-98141-3

1623 Wickham, H., François, R., Henry, L., & Müller, K. (2022). *dplyr: A Grammar of Data*
1624 *Manipulation*. https://github.com/tidyverse/dplyr

1625 Woltereck, R. (1913). Weitere experimentelle untersuchungen über Artänderung, speziell über
1626 das Wesen quantitativer Artunterschiede bei Daphniden. *Zeitschrift Für Induktive*
1627 *Abstammungs- Und Vererbungslehre*, *9*(1), 146–146. https://doi.org/10.1007/BF01876686

1628 Wooster, D. (1994). Predator impacts on stream benthic prey. *Oecologia*, *99*(1–2), 7–15.
1629 https://doi.org/10.1007/BF00317077

1630 Work, K. A., & Gophen, M. (1999). Factors which affect the abundance of an invasive
1631 cladoceran, Daphnia lumholtzi, in U.S. reservoirs. *Freshwater Biology*, *42*(1), 1–10.
1632 https://doi.org/10.1046/J.1365-2427.1999.00449.X

1633 Yampolsky, L. Y., Zeng, E., Lopez, J., Williams, P. J., Dick, K. B., Colbourne, J. K., & Pfrender,
1634 M. E. (2014). Functional genomics of acclimation and adaptation in response to thermal
1635 stress in Daphnia. *BMC Genomics*, *15*(1), 1–12. https://doi.org/10.1186/1471-2164-15-
1636 859/TABLES/2

1637 Yaniv, M. (2014). Chromatin remodeling: From transcription to cancer. In *Cancer Genetics*
1638 (Vol. 207, Issue 9, pp. 352–357). Elsevier Inc.
1639 https://doi.org/10.1016/j.cancergen.2014.03.006

1640 Yin, J., Zhou, M., Lin, Z., Li, Q. Q., & Zhang, Y. Y. (2019). Transgenerational effects benefit
1641 offspring across diverse environments: a meta-analysis in plants and animals. *Ecology*

1642   *Letters, 22*(11), 1976–1986.

1643   https://doi.org/10.1111/ELE.13373;WGROUP:STRING:PUBLICATION

1644