Oxidation and allocation of nectar amino acids during butterfly flight

ABSTRACT

Flying animals face extreme energetic demands, relying mainly on carbohydrates and lipids, with
occasional contributions from proteins and amino acids. In nectar-feeding species like butterflies
and hummingbirds, sugars are the primary fuel, yet the extent to which other nectar-derived
nutrients, like amino acids, are used for flight or retained for other functions remains unclear.
Using [13]C-labeled nectar, we tracked the metabolic fate of sugars and amino acids during flight
in *Pieris rapae* butterflies. We found that proline and glycine, two of the most abundant nectar
amino acids, were oxidized alongside sugars. Importantly, flight intensity modulated nutrient
allocation from nectar: high-flight females incorporated less glycine into tissues, implying
diversion toward flight, while threonine deposition in abdomens increased, reflecting
prioritization for reproduction and storage. These findings reveal the complex role of nectar-
derived nutrients in supporting locomotion and reproduction, while showing how nectar use can
modulate trade-offs between flight and fecundity.

16 INTRODUCTION

Powered flight is a key adaptation across animals, allowing insects, birds, and bats to forage,
escape predators, locate mates, and disperse over long distances (Anderson & Ruxton, 2020;
Dudley, 2002; Ruaux et al., 2020). Yet flight is also the most energetically costly mode of
locomotion, far exceeding the metabolic demands of running or swimming (Bale et al., 2014;
Schmidt-Nielsen, 1972). In insects, for example, flight can demand up to 100 times the resting
metabolic rate (Anderson & Ruxton, 2020; Dudley, 2002; Ruaux et al., 2020). Thus, meeting
these energetic demands requires efficient fuel use, with flying animals relying primarily on
carbohydrates and lipids, and, in some cases, proteins and amino acids.The relative contribution
of each fuel depends on the species, flight intensity, ecological context, and resource availability
(Weber, 2011).

For nectar-feeding animals such as butterflies, bees, and hummingbirds, nectar sugars are the

dominant energy source during flight (O'Brien, 1999; Suarez et al., 2011). Yet, nectar is a

complex resource that also contains small but physiologically relevant amounts of amino acids

(Baker & Baker, 1986; Nicolson & Thornburg, 2007). While studies, especially in insects,

indicate that some amino acids are oxidized from internal stores during flight (Teulier et al.,

2016; Stec et al., 2021; Weeda et al., 1979), it is still unclear whether amino acids obtained

directly from nectar are similarly used in flight.

Beyond serving as a fuel for flight, nectar can also contribute to reproduction and long-term

somatic functions. In many butterflies and moths, for example, nectar sugars as well as amino

acids are incorporated into proteins and directly support oogenesis, supplementing or in some

cases even replacing nutrient reserves acquired during larval development (Jervis & Boggs,

2005; Levin et al., 2017; Mevi□Schütz & Erhardt, 2005). Similarly, in bird pollinators,

especially those that rely heavily on nectar, amino acids in nectar have been suggested to help

supplement their nitrogen requirements (Nicolson, 2007; Roguz et al., 2019). This dual role of

nectar-derived resources may accentuate tradeoffs between flight and reproduction, as amino

acids oxidized to sustain flight are no longer available for egg production. Thus, understanding

whether nectar amino acids are metabolized during flight, or instead retained for tissue

incorporation, is critical to link flight energetics with fitness and help clarify the physiological

mechanisms underlying frequently observed trade-offs between flight and reproduction

(Schmidt-Wellenburg et al., 2008; Tigreros & Davidowitz, 2019).

Here, we investigate the metabolic fate of nectar-derived nutrients in *Pieris rapae*, a common

butterfly that relies on floral nectar throughout its adult life. Using [13]C-labeled sugars and amino

acids (proline, glycine, and threonine), we examined nutrient oxidation during flight as well as

nutrient deposition in thoracic and abdominal tissues under varying flight conditions. By

distinguishing which nectar components serve as immediate fuels and which are retained for

other functions, we provide new insight into the role of nectar composition in supporting the

energetics of insect flight.

MATERIALS AND METHODS

Study organism and Rearing

The cabbage white butterfly (*Pieris rapae*) is a widespread species whose adults feed on floral nectar, serving as pollinators for a variety of plants while obtaining sugars and amino acids as nutritional rewards (Alm et al., 1990; Corbera et al., 2018; Hongfang et al., 2010; Rader et al., 2009). Like other pollinators, female *P. rapae* can detect and preferentially consume nectars enriched in amino acids (Alm et al., 1990). This species faces high energetic demands due to its investment in both flight and reproduction: females fly an average of 0.7 km per day (Scott, 1987) and lay an average of 278 ± 172 eggs over a short adult lifespan of about 12.2 ± 4.0 days (Kimura & Tsubaki, 1986).

For this study, *P. rapae* were obtained from a laboratory colony established from a wild population in Utah, USA. Larvae were reared in a greenhouse on *Brassica oleracea*. Pupae and adults were maintained in a walk-in environmental chamber at 22 °C, 50% relative humidity, and a 16:8 h light–dark cycle.

Nectar Treatments

In this study, we used artificial nectars that replicated the sugar and amino acid composition of *Lantana camara* (Verbenaceae), a high-quality nectar source frequently visited by diverse butterfly species. The composition of *L. camara* nectar has been well characterized and is widely used in studies of nectar preference in *P. rapae* (Alm et al., 1990) and other Lepidoptera (Erhardt & Rusterholz, 1998; Jervis & Boggs, 2005; Mevi-Schütz & Erhardt, 2003). Natural L. camara nectar typically contains ~1 M sugars and ~10 mM amino acids, yielding an approximate 1:100 amino acid-to-sugar ratio. Of the amino acid pool, roughly 6% comprises three essential amino acids, with threonine being the most abundant, while the remaining 94% consists of eight nonessential amino acids, dominated by proline and glycine.

To track use of specific nectar's nutrients by female *P. rapae*, we created five lantana-mimicking nectar treatments. Four contained a single ^{13}C-labeled compound—^{13}C☐-glycine, ^{13}C☐-proline, ^{13}C☐-threonine, or ^{13}C-sucrose (enriched with cane sugar)—while the fifth served as a control,

82 containing beet sugar and only unlabeled amino acids ($\delta^{13}C \approx -26.5‰$). The labeled amino acids

83 were selected because they are abundant in *L. camara* nectar and in other butterfly-pollinated

84 flowers (Mevi-Schütz & Erhardt, 2003; Roguz et al., 2019). All isotope tracers were obtained

85 from Cambridge Isotope Laboratories, Inc. (Tewksbury, MA, USA).

Experiment 1. Oxidation of nectar nutrients during flight

To assess oxidation of nectar nutrients during flight, females were fed 48 h post-emergence with

10 µl of either a Control or one of four $^{13}C\square$-labelled nectars: $^{13}C\square$-glycine, $^{13}C\square$-proline, $^{13}C\square$-

threonine, or ^{13}C-sucrose. Four hours after feeding, each female was placed in a respirometry

chamber consisting of a 500 ml syringe fitted with an injection port and adjusted to provide a

200 ml chamber volume. This volume was sufficient to allow flight while enabling $CO\square$

accumulation to ≥ 300 ppm. The syringe air was replaced with dried, $CO\square$-free air, sealed, and

females were flown for 4 min under natural light, with gentle shaking to prevent resting. A 20 ml

breath sample was then analyzed for $\delta^{13}CO\square$ using a cavity ring-down spectrometer (G2201-i,

Picarro Inc., Sunnyvale, CA) coupled to a SSIM2 Small Sample Isotope Module (Picarro Inc.,

Sunnyvale, CA).

Experiment 2. The effect of flight on nectar nutrients depositions

To determine where nectar-derived amino acids are deposited, we used a different group of adult

females (48 h post-emergence) which were randomly assigned to one of two flight treatments—

Low or High flight intensity—and fed either a control nectar or one of four $^{13}C\square$-labelled

nectars. For the Low flight intensity group, females were kept in a $30 \times 30 \times 30$ cm mesh cage

lined with paper towel inside a climate-controlled walk-in chamber (LD 16:8 h ,22°C at 50%

RH). For the High flight intensity group, females were kept under the same conditions but were

additionally stimulated to fly once per day, for five continuous minutes, by gently touching them

with a fine paintbrush (Niitepõld & Boggs, 2015). Preliminary trials showed that longer flight

durations caused extreme exhaustion, with butterflies becoming unresponsive. All females were

fed once daily with 10 µl of their assigned nectar treatment. After three consecutive days of the

flight and nectar treatments, individuals were euthanized by freezing at −20 °C. Thoraces and

abdomens were dissected, dried at 50 °C for 48 h, and homogenized. $\delta^{13}C$ values were then

measured using cavity ring-down spectroscopy (CRDS; Picarro stable isotope analyzer, G2201-i)

coupled with an A0201 combustion module and A0301 gas interface.

Statistical Analysis

To assess nectar-derived nutrient oxidation (Experiment 1) and tissue deposition (Experiment 2),

we used Dunnett's test to compare the $\delta^{13}C$ values of breath or tissue samples from females fed

each ^{13}C-labeled nectars against those of control females fed unlabeled nectar.

Because $\delta^{13}C$ values can naturally change with tissue type and treatment, we used $\delta^{13}C$ of control

females to estimate atom % excess (APE), which is a unit of enrichment (McCue et al., 2011;

Slater et al., 2001). We used ^{13}C concentrations of both control and labelled tissues expressed in

$\delta^{13}C_{VPDB}$ to calculate the atom percent (AP) and then Atom Percent Excess (APE), defined as

APE = AP(label) – AP(control) (Slater et al., 2001).

The effects of flight on nutrient allocation to thoracic and abdominal tissues were analyzed using

two-way ANOVAs with flight treatment, nectar nutrient treatment (^{13}C-labeled vs. control), and

their interaction as fixed factors. When significant interactions were detected, we estimated

predicted values and performed pairwise comparisons of simple effects using the *emmeans*

package in R, applying a Tukey adjustment for multiple testing. All statistical analyses were

conducted in R 3.6.0. and assumptions of normality and homogeneity of variance were verified

prior to analysis.

RESULTS AND DISCUSSION

Nectar is a critical resource for flying pollinators, providing the energy needed to support their

high metabolic demands during flight (Brien, 1999; Suarez, 2005; Welch et al., 2006). While it is

well established that nectar sugars serve as the primary fuel for flight, far less is known about the

roles of non-sugar components, like amino acids. Here we investigated the use of nectar-derived

nutrients, including essential and non-essential amino acids, as metabolic fuels during flight and

how flight activity may influences their allocation to different tissues in female *Pieris rapae*.

Oxidation of nectar nutrients during flight

Females that fed on nectar containing ^{13}C-labeled sugars,

proline, or glycine showed significant ^{13}CO☐ enrichment

(more positive δ^{13}C values) in breath compared to controls

fed unlabeled nectar (**Fig. 1**), indicating active oxidation of

both sugars and non-essential amino acids. Although

previous studies have suggested that nectar amino acids

help meet the high metabolic demands of flying

nectarivores (Carter et al., 2006; Levin et al., 2017; Stec et

al., 2021), our results provide the first direct evidence that

nectar amino acids are actually oxidized during flight. At

the same time, no significant oxidation of threonine by

flying females was observed (**Fig. 1**), suggesting that

essential amino acids may be less likely to contribute to

locomotion, at least under our experimental conditions.

Fig. 1 Boxplots show the δ^{13}C values from exhaled breath during flight of *P. rapae* females. Significant differences between the δ^{13}C from control (blue boxplot) and females that consumed ^{13}C-labelled nectars indicate that a nectar-derived nutrient (sugars, glycine, proline, or threonine) was oxidized during flight.

These results add to the growing body of evidence

highlighting the importance of nectar amino acids for flying animals, and in particular for insect

pollinators. Proline and glycine are common constituents of floral nectar (Baker & Baker, 1973;

Rusterholz & Erhardt, 1998) and can act as a phagostimulant in pollinator insects species (Lim et

al., 2019; Ruedenauer et al., 2019). Several studies have shown that proline, in particular, is

selectively oxidized in insect flight muscle, especially during the initial 30 seconds of flight (Stec

et al., 2021; Teulier et al., 2016; Weeda et al., 1979). Though glycine's role in flight is less

defined, its consumption has been linked to neuromodulation and memory in insects (Parkinson

et al., 2025), suggesting a dual function: fuelling flight and supporting cognitive processes
essential for effective foraging.

The effect of flight on nectar nutrients depositions

Gravimetric measurements of the impact of flight on thorax and abdomen resources did not
indicate any differences (both thorax and abdomen $p >$ 0.05). In contrast, isotopic analysis of
control, unlabeled butterflies, revealed significant changes in tissue carbon composition due to
flight intensity. Here, females subjected to high-intensity flight showed enriched $\delta^{13}C$ values in
both thorax and abdomen compared to low-flight individuals (ANOVA: $F_{\square,\square\square} = 6.26$, $P =$
0.02), consistent with increased oxidation and depletion of lipid stores. This is in line with
previous studies showing that butterflies rely on both carbohydrates and lipids to meet the high
energy costs of flight, and that sustained flight can rapidly deplete internal energy, often leading
to flight fecundity tradeoffs (Tigreros &
Davidowitz, 2019).

Previous studies have shown that nectar in
general and its amino acids in particular can be
rapidly incorporated, within minutes, into
Lepidoptera flight muscle and developing
oocytes (DeFino & Davidowitz, 2024; Levin et
al., 2017). Given that the thorax serves as the
primary site of fuel combustion during flight
(Dudley, 2002; Treidel et al., 2024), it can be
expected that incoming amino acids and sugars
are preferentially oxidized rather than retained.
Here, we demonstrate that flight activity
significantly alters the allocation of nectar
nutrients across female tissues ($F_{\square,\square\square\square} = 5.33$,
$P = 0.002$). In the thorax, the presence of nectar-

Fig. 2 Effects of flight, high vs. low intensity, on the use of nectar-derived nutrients in abdomen (left panel) and thorax (right panel) tissues of *P. rapae* females. The y-axis represents the estimated marginal means ± 95CI of the atom percent excess (APE ^{13}C (%)), with higher APE values indicating greater incorporation of nectar derived sucrose, non-essential amino acids (glycine and proline) and an essential amino acid (threonine).

derived nutrients declined with increasing flight intensity ($F_{\square,\square\square} = 4.99$, $P = 0.03$; **Fig. 2**),

consistent with their increased metabolic oxidation or mobilization. Interestingly, while most

labelled amino acids were detected in both thorax and abdomen tissues (**Fig. S1**), proline was

notably absent in the thorax under high flight conditions, and it only approached significance in

low-flight males ($P = 0.09$; **Fig. S1**). This supports the hypothesis that proline is rapidly oxidized

during flight, leaving little available for tissue incorporation.

Patterns of nutrient incorporation and use were distinct in the abdomen, reflecting its key roles in

nutrient storage and reproduction (**Fig. 2**). We found that the incorporation of nectar nutrients

varied by type ($F_{3,51} = 4.9$, $P = 0.01$): glycine decreased (p = 0.002), while threonine increased (p

= 0.03) with higher flight activity (**Fig. 2; Table S1**). Interestingly, proline and sucrose

incorporation remained unchanged (**Fig. 2; Table S1**). As the primary site of nutrient storage and

reproduction, the abdomen likely channels nectar-derived nutrients toward oogenesis and long-

term metabolic demands (Boggs, 2009; O'Brien et al., 2002). The decline in abdominal glycine,

an amino acid significantly oxidized during flight (**Fig. 1**), suggests that this non-essential amino

acid may be diverted to meet immediate energy demands, possibly contributing to a trade-off

with female fecundity. Conversely, the increase in threonine, an essential amino acid, may reflect

its selective prioritization for egg provisioning. These findings support the idea that adult

Lepidoptera use nectar to buffer against nutrient limitations, whether from poor larval diets

(Mevi-Schütz & Erhardt, 2005; O'Brien et al., 2000) or increased metabolic demands from

flight.

Collectively, our findings demonstrate that *P. rapae* females utilize both sugars and specific

amino acids—particularly proline and glycine—as metabolic fuels during flight. Moreover,

intense flight alters the fate and allocation of nectar-derived nutrients in adult tissues, likely

reflecting a trade-off between immediate energy needs and resource storage. These insights

contribute to a growing understanding of how nectar composition affects insect energetics and

fitness, highlighting the ecological importance of amino acids in adult diets, particularly under

conditions of high energy demand.

REFERENCES

Alm, J., Ohnmeiss, T. E., Lanza, J., & Vriesenga, L. (1990). Preference of cabbage white butterflies and honey bees for nectar that contains amino acids. *Oecologia*, *84*, 53–57. https://doi.org/10.1007/BF00665594

Anderson, S. C., & Ruxton, G. D. (2020). The evolution of flight in bats: a novel hypothesis. In *Mammal Review* (Vol. 50, Issue 4, pp. 426–439). John Wiley & Sons, Ltd. https://doi.org/10.1111/mam.12211

Baker, H. G., & Baker, I. (1973). Amino-acids in nectar and their evolutionary significance. In *Nature* (Vol. 241, Issue 5391, pp. 543–545). Nature Publishing Group. https://doi.org/10.1038/241543b0

Baker, H. G., & Baker, I. (1986). The occurrence and significance of amino acids in floral nectar. *Plant Systematics and Evolution*. https://doi.org/10.1007/BF02430273

Bale, R., Hao, M., Bhalla, A. P. S., & Patankar, N. A. (2014). Energy efficiency and allometry of movement of swimming and flying animals. *Proceedings of the National Academy of Sciences of the United States of America*, *111*(21), 7517–7521. https://doi.org/10.1073/pnas.1310544111

Boggs, C. L. (2009). Understanding insect life histories and senescence through a resource allocation lens. In *Functional Ecology*. https://doi.org/10.1111/j.1365-2435.2009.01527.x

Brien. (1999). *Fuel use flexibility in a nectarivorous hawkmoth*. https://jeb.biologists.org/content/jexbio/202/4/441.full.pdf

Carter, C., Shafir, S., Yehonatan, L., Palmer, R. G., & Thornburg, R. (2006). A novel role for proline in plant floral nectars. *Naturwissenschaften*, *93*(2), 72–79. https://doi.org/10.1007/s00114-005-0062-1

Corbera, J., Alvarez-Cros, C., & Stefanescu, C. (2018). Evidence of butterfly wing pollination in the martagon lily Lilium martagon L. *Butlletí ICHN*, *82*, 117–120. https://www.researchgate.net/publication/327968095_Evidence_of_butterfly_wing_pollination_in_the_martagon_lily_Lilium_martagon_L

DeFino, N., & Davidowitz, G. (2024). Crop-emptying rate and nectar resource allocation in a nectivorous pollinator. *Journal of Insect Physiology*, *154*, 104617. https://doi.org/10.1016/j.jinsphys.2024.104617

Dudley, R. (2002). The Biomechanics of Insect Flight: Form, Function, Evolution. In *Princeton University Press*.

Erhardt, A., & Rusterholz, H. P. (1998). Do peacock butterflies (Inachis io L.) detect and prefer nectar
amino acids and other nitrogenous compounds? *Oecologia*, *117*(4), 536–542.
https://doi.org/10.1007/s004420050690

Hongfang, Z., Liqiang, L., Zhongjian, L., & Yibo, L. (2010). The butterfly Pieris rapae resulting in the
reproductive success of two transplanted orchids in a botanical garden. *Biodiversity Science*, *18*(1),
11. https://doi.org/10.3724/sp.j.1003.2010.011

Jervis, M. A., & Boggs, C. L. (2005). Linking nectar amino acids to fitness in female butterflies. In
Trends in Ecology and Evolution (Vol. 20, Issue 11, pp. 585–587).
https://doi.org/10.1016/j.tree.2005.08.015

Kimura, K., & Tsubaki, Y. (1986). Female size and age-specific fecundity in the small white butterfly,
Pieris rapae crucivora Boisduval (Lepidoptera: Pieridae). *Population Ecology*, *28*(2), 295–304.
https://doi.org/10.1007/BF02515456

Levin, E., McCue, M. D., & Davidowitz, G. (2017). More than just sugar: Allocation of nectar amino
acids and fatty acids in a lepidopteran. *Proceedings of the Royal Society B: Biological Sciences*.
https://doi.org/10.1098/rspb.2016.2126

Lim, S., Jung, J., Yunusbaev, U., Ilyasov, R., & Kwon, H. W. (2019). Characterization and its implication
of a novel taste receptor detecting nutrients in the honey bee, Apis mellifera. *Scientific Reports 2019*
9:1, *9*(1), 1–13. https://doi.org/10.1038/s41598-019-46738-z

McCue, M. D., Smith, A., McKinney, R., Rewald, B., Pinshow, B., & McWilliams, S. R. (2011). A mass
balance approach to identify and compare differential routing of 13C-labeled carbohydrates, lipids,
and proteins in vivo. *Physiological and Biochemical Zoology*, *84*(5), 506–513.
https://doi.org/10.1086/661638

Mevi-Schütz, & Erhardt. (2005). Amino Acids in Nectar Enhance Butterfly Fecundity: A Long-Awaited
Link. *The American Naturalist*. https://doi.org/10.2307/3473471

Mevi-Schütz, J., & Erhardt, A. (2003). Effects of nectar amino acids on fecundity of the wall brown
butterfly (Lasiommata megera L.). *Basic and Applied Ecology*, *4*(5), 413–421.
https://doi.org/10.1078/1439-1791-00183

Mevi□Schütz, J., & Erhardt, A. (2005). Amino Acids in Nectar Enhance Butterfly Fecundity: A
Long□Awaited Link. *The American Naturalist*, *165*(4), 411–419. https://doi.org/10.1086/429150

Nicolson, S. W. (2007). Amino acid concentrations in the nectars of southern African bird-pollinated

flowers, especially Aloe and Erythrina. *Journal of Chemical Ecology*, *33*(9), 1707–1720. https://doi.org/10.1007/s10886-007-9342-x

Nicolson, S. W., & Thornburg, R. W. (2007). Nectar chemistry. In S. W. Nicolson, M. Nepi, & E. Pacini (Eds.), *Nectaries and Nectar* (pp. 215–264). Springer US. https://doi.org/10.1007/978-1-4020-5937-7_5

Niitepõld, K., & Boggs, C. L. (2015). Effects of increased flight on the energetics and life history of the butterfly speyeria mormonia. *PLoS ONE*, *10*(10), e0140104. https://doi.org/10.1371/journal.pone.0140104

O'Brien, D. M. (1999). Fuel use in flight and its dependence on nectar feeding in the hawkmoth Amphion floridensis. *The Journal of Experimental Biology*, *202*(Pt 4), 441–451. https://doi.org/10.1111/j.1754-9485.2011.02309.x

O'Brien, D. M., Fogel, M. L., & Boggs, C. L. (2002). Renewable and nonrenewable resources: amino acid turnover and allocation to reproduction in Lepidoptera. *Proceedings of the National Academy of Sciences of the United States of America*, *99*(7), 4413–4418. https://doi.org/10.1073/pnas.072346699

O'Brien, D. M., Schrag, D. P., & Martínez Del Rio, C. (2000). Allocation to reproduction in a hawkmoth: A quantitative analysis using stable carbon isotopes. *Ecology*. https://doi.org/10.1890/0012-9658(2000)081[2822:ATRIAH]2.0.CO;2

Rader, R., Howlett, B. G., Cunningham, S. A., Westcott, D. A., Newstrom-Lloyd, L. E., Walker, M. K., Teulon, D. A. J., & Edwards, W. (2009). Alternative pollinator taxa are equally efficient but not as effective as the honeybee in a mass flowering crop. *Journal of Applied Ecology*, *46*(5), 1080–1087. https://doi.org/10.1111/j.1365-2664.2009.01700.x

Roguz, K., Bajguz, A., Chmur, M., Gołębiewska, A., Roguz, A., & Zych, M. (2019). Diversity of nectar amino acids in the Fritillaria (Liliaceae) genus: ecological and evolutionary implications. *Scientific Reports*, *9*(1), 1–12. https://doi.org/10.1038/s41598-019-51170-4

Ruaux, G., Lumineau, S., & De Margerie, E. (2020). The development of flight behaviours in birds. *Proceedings of the Royal Society B: Biological Sciences*, *287*(1929). https://doi.org/10.1098/rspb.2020.0668

Ruedenauer, F. A., Leonhardt, S. D., Lunau, K., & Spaethe, J. (2019). Bumblebees are able to perceive amino acids via chemotactile antennal stimulation. *Journal of Comparative Physiology A:*

Neuroethology, Sensory, Neural, and Behavioral Physiology, *205*(3), 321–331. https://doi.org/10.1007/s00359-019-01321-9

Rusterholz, H. P., & Erhardt, A. (1998). Effects of elevated CO2 on flowering phenology and nectar production of nectar plants important for butterflies of calcareous grasslands. *Oecologia*, *113*(3), 341–349. https://doi.org/10.1007/s004420050385

Schmidt-Nielsen, K. (1972). Locomotion: Energy cost of swimming, flying, and running. In *Science* (Vol. 177, Issue 4045, pp. 222–228). American Association for the Advancement of Science. https://doi.org/10.1126/science.177.4045.222

Schmidt-Wellenburg, C. A., Visser, G. H., Biebach, B., Delhey, K., Oltrogge, M., Wittenzellner, A., Biebach, H., & Kempenaers, B. (2008). Trade-off between migration and reproduction: Does a high workload affect body condition and reproductive state? *Behavioral Ecology*, *19*(6), 1351–1360. https://doi.org/10.1093/beheco/arn066

Scott, A. J. (1987). The Butterflies of North America: A Natural History and Field Guide. In P. J. DeVries & M. C. Singer (Eds.), *The Quarterly Review of Biology* (Vol. 62, Issue 3). Stanford University Press. https://doi.org/10.1086/415568

Slater, C., Preston, T., & Weaver, L. T. (2001). Stable isotopes and the international system of units. *Rapid Communications in Mass Spectrometry*, *15*(15), 1270–1273. https://doi.org/10.1002/rcm.328

Stec, N., Saleem, A., & Darveau, C. A. (2021). Proline as a sparker metabolite of oxidative metabolism during the flight of the bumblebee, bombus impatiens. *Metabolites*, *11*(8), 511. https://doi.org/10.3390/metabo11080511

Suarez, R. K. (2005). Energy metabolism in orchid bee flight muscles: carbohydrate fuels all. *Journal of Experimental Biology*, *208*(18), 3573–3579. https://doi.org/10.1242/jeb.01775

Suarez, R. K., Herrera M., L. G., & Welch, K. C. (2011). The sugar oxidation cascade: Aerial refueling in hummingbirds and nectar bats. *Journal of Experimental Biology*, *214*(2), 172–178. https://doi.org/10.1242/jeb.047936

Teulier, L., Weber, J. M., Crevier, J., & Darveau, C. A. (2016). Proline as a fuel for insect flight: Enhancing carbohydrate oxidation in hymenopterans. *Proceedings of the Royal Society B: Biological Sciences*, *283*(1834). https://doi.org/10.1098/rspb.2016.0333

Tigreros, N., & Davidowitz, G. (2019). Flight-fecundity tradeoffs in wing-monomorphic insects. In *Advances in Insect Physiology*. https://doi.org/10.1016/bs.aiip.2019.02.001

Treidel, L. A., Deem, K. D., Salcedo, M. K., Dickinson, M. H., Bruce, H. S., Darveau, C. A., Dickerson,
B. H., Ellers, O., Glass, J. R., Gordon, C. M., Harrison, J. F., Hedrick, T. L., Johnson, M. G.,
Lebenzon, J. E., Marden, J. H., Niitepõld, K., Sane, S. P., Sponberg, S., Talal, S., … Wold, E. S.
(2024). Insect Flight: State of the Field and Future Directions. *Integrative and Comparative Biology*,
64(2), 533–555. https://doi.org/10.1093/icb/icae106

Weber, J.-M. (2011). Metabolic fuels: regulating fluxes to select mix. *Journal of Experimental Biology*,
214(2), 286–294. https://doi.org/10.1242/jeb.047050

Weeda, E., Kort, C. A. D., & Beenakkers, A. M. T. (1979). Fuels for energy metabolism in the Colorado
potato beetle, _Leptinotarsa decemlineata_ Say. *Journal of Insect Physiology*, *25*(12), 951–955.
file:///y:/7305.pdf

Welch, K. C., Bakken, B. H., Martínez Del Rio, C., & Suarez, R. K. (2006). Hummingbirds fuel hovering
flight with newly ingested sugar. *Physiological and Biochemical Zoology*, *79*(6), 1082–1087.
https://doi.org/10.1086/507665

351