Robust height-diameter allometries for 41 European tree species: stand characteristics and structure matter

Natacha Savine[*], Thomas Cordonnier, Gauthier Ligot, Patrick Vallet

Natacha Savine
Univ. Grenoble Alpes, INRAE, LESSEM, 2 rue de la Papeterie-BP 76, F-38402 St-Martin-d'Hères, France
orcid: 0009-0002-5978-3487

Thomas cordonnier
Office National des Forêts, Recherche Développement et Innovation, 21 rue du Muguet, 39100 Dole, France
orcid: 0000-0003-3684-4662

Gauthier Ligot
TERRA Teaching and Research Centre, Gembloux Agro-Bio Tech, University of Liège, Gembloux, Belgium
orcid: 0000-0002-5508-4358

Patrick Vallet
Univ. Grenoble Alpes, INRAE, LESSEM, F-38402 St-Martin-d'Hères, France
orcid: 0000-0003-2649-9447

*Corresponding author: natacha.savine@inrae.fr

Abstract

Tree height is a key variable for assessing forest functioning and resources (e.g volume, carbon stocks) at both tree and stand levels. However, direct field measurements are costly and time-consuming. Developing accurate and unbiased height-diameter allometries applicable over large spatial scales is therefore crucial for forest research and management.

Based on the French National Forest Inventory, with 269,460 tree-height observations on 49,120 plots measured, we developed generalized, species-specific height-diameter allometries that integrate stand dendrometrical characteristics and stand structure. The models cover 41 European tree species and the four main stand structures: even-aged, uneven-aged, coppice-with-standards and coppice. To enhance local accuracy, we developed an optional recalibration method at the plot level, assessing how many additional trees should be measured and which ones to select to maximize the improvement of the model. By integrating basal area as stand density and quadratic mean diameter as an indicator of stand development stage, our models enable us to assess the influence stand dendrometrical characteristics on these allometries and evaluate how different species responded to competition.

Results indicate that with increasing competition, tree height tends to be higher for a given diameter, and that stand structure significantly influences 28 out of the 41 species. Local recalibration showed that measuring just one to six trees (among the largest and thinnest diameter) per plot reduced prediction error by 10 to 70%, depending on species.

This study provides a useful, robust and scalable tool for forest research and management, for the most widespread species in Europe, while offering precision for local applications.

Keywords: Height-diameter allometry, European tree species, stand structure, National Forest Inventory, nonlinear mixed effects model, local recalibration.

Declarations

Funding

65 The French National Forest Office funded this work under the grant "Régénération, croissance et production des
66 forêts de plaine, 2022-2024"

Conflicts of interest

69 The authors declare that they have no known competing financial interests or personal relationships that could
70 have appeared to influence the work reported in this paper.

Availability of data and material

73 Data from the French National Forest Inventory are available online.

Author's contributions

76 Natacha Savine: Writing – original draft, Validation, Methodology, Investigation, Data curation,
77 Conceptualization.
78
79 Thomas Cordonnier: Writing – review & editing, Supervision, Methodology, Investigation, Conceptualization.
80
81 Gauthier Ligot: Writing – review & editing, Supervision, Methodology, Investigation.
82
83 Patrick Vallet: Writing – review & editing, Supervision, Methodology, Investigation, Funding acquisition,
84 Conceptualization.
85

Acknowledgments:

87 The authors wish to thank the French National Forest Office, who funded this work under the grant

88 "Régénération, croissance et production des forêts de plaine, 2022-2024". The authors also thank the National

89 Forest Inventory for making the data available free of charge.

1. Introduction

Forest management involves not only assessing existing resources but also evaluating how they have changed, or will change, over time. In this respect, along with diameter, tree height is a key variable to assess forest ecosystem functioning and forest ecosystem services such as carbon storage or wood production (Hunter et al., 2013; Teobaldelli et al., 2009). Studying height variability across individuals can be key to understanding important ecological processes, such as competitive and facilitation interactions between individuals.

However, accurately measuring tree height is often time-consuming and technically challenging especially on slopes or stands with low visibility. As a consequence, tree height measurements are often performed on a limited sample of trees.

In biological organisms, allometries defines how the size of an organ changes in relation to the size of another one, or to the size of the whole organism. These allometries are used to understand how variables are interconnected, and can serve as an alternative for variables that are difficult to measure directly. In forestry, allometric models can be used to estimate tree characteristics such as biomass (Pilli et al., 2006) or volume (Vallet et al., 2006), being common examples. For these two allometric models, the most accurate models include height measurements, or height estimates. Tree height can also be estimated indirectly through allometric relationships, most commonly by using diameter at breast height (dbh), a variable that is much easier to obtain in the field (Ciceu et al., 2023).

Height-diameter allometries are known to differ across species, due to biological constraints, limitations and growth strategies inherent to each species (Vázquez-Veloso et al., 2025). Allometries can therefore be species-specific (Adame et al., 2008; Ciceu et al., 2020; Huang et al., 1992) or can take species characteristics directly into account in the model (Sharma et al., 2019b).

It is also essential to consider the specific environmental conditions influencing individual growth. When trees grow in proximity to other individuals, they compete for resources in different ways. In dense areas, where space is limited, trees may allocate more resources to vertical growth, which increases height, rather than to radial growth. This phenomenon can be integrated into allometric models by considering the social status of the tree (Aussenac et al., 2023; Trouvé et al., 2015). Social status can be used per se in the models, or can be included as a competitive index as in Mugasha et al. (2019).

The dendrometrical characteristics of the stand also play a role in shaping height-diameter allometries. Stand density variables such as basal area, for example, are commonly used to account for competition. As a result, trees

growing in highly competitive environments present greater height for a given diameter compared to those in less

competitive ones.

Ciceu et al (2023) highlighted that dominant height is used to reflect the effect of stand developmental stage in

most studies working on height-diameter allometry. However, to obtain this variable, the height of a subset of trees

must be measured. The widespread inclusion of dominant height in empirical models is likely due to the strong

correlation between dominant height and individual tree heights. Alternative variables, such as the quadratic mean

diameter, also reflect stand development stage (Aussenac et al., 2023; Ciceu et al., 2020; Feldpausch et al., 2011;

Gonçalves, 2022; Mugasha et al., 2019; Temesgen and v. Gadow, 2004).

The structure of the stand (even-aged, uneven-aged and other configurations) also influences height-diameter

allometry(Gonçalves, 2022). Different stand structures reflect different stand dynamics and result in diverse tree

diameter distributions. For instance, even-aged stands are characterized by individuals of similar ages and heights,

whereas uneven-aged stands exhibit more diverse ages and height structures, potentially resulting in different

allometric patterns.

Throughout the diversity of environmental conditions and forest structures, the French metropolitan forest area is

covered by 58% of even-aged stands. This leaves 42% of the stands that are managed under other regimes such as

uneven-aged, coppice or coppice-with-standards systems (described in figure S1, Supplementary Materials). The

even-aged structure has been the focus of most studies on forest structure, as it is one of the most represented and

constitutes a simple silvicultural system (Bettinger et al., 2017). However, since structure plays an important role

in growth, especially with different competitive relationships, it is necessary to study how height-diameter

allometries are affected by each structure specifically. This could prevent overestimations or underestimations

when assessing forest resources (Gonçalves, 2022).

Eventually, allometric models can be developed at different scales, from local to continental. While large-scale

models can provide higher genericity, they may introduce estimation bias when applied to specific local conditions.

Studies by Sullivan et al. (2018), Mensah et al. (2018) and Kearsley et al. (2017) have shown that locally calibrated

models perform better than large scale models. To improve the genericity and performance of these models, it can

be beneficial to incorporate local factors such as soil fertility, water availability or climate, thus allowing the model

to adapt to specific conditions.

In this study, we developed height-diameter allometries for 41 species. We hypothesized that this relationship

would be influenced by individual-tree and stand characteristics such as stand basal area, mean diameter and stand

structure (even-aged, uneven-aged, etc.). We attempted to provide the most practical and generic models possible,

for the most species, and for different stand structures. We also developed a method to locally adapt large-scale

models, with as few field measurements as possible.

2. Material

2.1 Measurement protocol

We used data collected by the French National Forest Inventory (NFI) between 2005 and 2022 (IGN, 2023). The

plots were located throughout the French metropolitan area, on a systematic sampling grid of 1x1 km. Each year,

one twentieth of the grid was sampled, representing approximately 6000 plots per year. The plots are composed

of three concentric subplots with 6-, 9- and 15-meter radii. Very small trees (dbh [7.5-12.5cm [ ) and small trees

(dbh [12.5-22.5cm [ ) are measured in the 6-meter radius subplots, medium trees (dbh [22.5 – 37.5cm [ ) are

measured in the 9-meter radius subplots, and large (dbh [37.5 – 52.5 cm [ ) and very large trees (dbh over 52.5

cm) are measured in the 15-meter radius subplots. To account for the concentric circle sampling design, the NFI

assigns a weight to each tree based on its diameter class (very small, small, medium, large and very large). These

weights are then used to calculate stand-level variables, such as quadratic mean diameter and basal area.

Until 2013, height was measured for six trees per diameter class and per species. From 2014 to 2015, only one tree

was measured per category and per species. In 2016, the very-small-tree category was added and additional height

measurements were carried out on very small trees. During the inventories, height and diameter are measured,

along with some other tree variables such as the vegetative state (living or dead) or stem singularities (broken,

abnormal, pollarded). The complete measurement protocol is described in Dalmasso et al. (2014).

From 2005 to 2013, the NFI classified stand structures in five categories based on the characteristics of the stand,

combining the regeneration method (coppice, coppice-with-standards, even-aged forest, uneven-aged forest, no

structure) and vertical distribution. From 2014 to 2022, the NFI used different stand structure classifications, which

we aggregated to distinguish four stand structures: even-aged (EA), uneven-aged (UA), coppice-with-standards

(CWS) and coppice (C) (aggregation criteria are described in Supplementary Materials, table S1).

2.2 Dataset sub-samples

We selected plots dominated by one species and for which stand structure was available. The dominant species

had to represent at least 75% of the plot basal area, considering only living trees. We discarded plots for which

stand structure was not available (structure not registered, undefined, or registered as no structure). Within the

selected plots, we excluded trees with special stem characteristics, such as broken or pollarded trees. This

eliminated 1.38% of the dataset. Additionally, we selected species for which there were at least one hundred tree

records. Only the combinations of species and stand structure for which we had at least ten tree records were

analyzed.

Our sample contained tree records for 51 species in all: 30 species in all four stand structures, 16 species in three

structures (14 not present in coppice stands, 2 not present in uneven-aged stands), and five species in two stand

structures (even-aged and coppice-with-standards standards). We also discarded the few tree records of

coniferous species in coppice stands.

Our final models were fitted for 41 out of the 51 species. For ten species, the fitting procedure did not reach

convergence, most likely because we had too few observations or because individual height was not evenly

distributed across diameter ranges.

The final calibration dataset contained 269,460 measured trees on 49,120 temporary plots. All 41 species are

presented in the Supplementary Materials, table S2, and the ten most sampled species are described in table 1.

Species	Number of trees	Number of plots	Number of trees per structure				DBH (cm)			Height (m)		
			EA	UA	CWS	C	min	mean	max	min	mean	max
Abies alba	19,055	2,361	12,502	3,772	2,781	0	7.48	36.91	129.23	2.4	21.90	46.2
Castanea sativa	14,334	2,670	2,993	417	7,050	3,874	7.48	27.46	262.92	2.2	15.40	34.8
Fagus sylvatica	28,135	4,402	16,437	1,302	9,201	1,195	7.48	33.52	174.12	1.3	20.21	47.1
Picea abies	19,176	2,664	14,933	2,127	2,116	0	7.51	32.14	127.01	1.6	21.68	48.4
Pinus pinaster	20,695	4,075	16,204	551	3,940	0	7.48	30.61	111.09	2.6	16.93	39.1
Pinus sylvestris	21,865	3,572	15,409	1,988	4,468	0	7.48	27.22	98.96	1.7	13.97	41.2
Pseudotsuga menziesii	13,691	2,151	11,809	207	1,675	0	7.51	33.20	140.69	2.0	23.62	49.4
Quercus petraea	31,927	5,772	16,779	575	13,370	1,203	7.48	35.64	137.35	1.9	20.95	44.0
Quercus pubescens	20,689	4,655	5,527	879	9,188	5,095	7.48	20.80	192.26	1.8	11.39	30.8
Quercus robur	28,049	5,691	13,362	935	12,762	990	7.48	35.68	164.12	2.2	18.83	43.2

204   *Table 1:Description of the sampling characteristics of the ten most sampled species in the calibration dataset.*

205   *EA: Even-aged, UA: Uneven-aged, CWS: Coppice-With-Standards, C: Coppice. The full list is available in*

206   *Supplementary Materials, Table S1.*

208 2.3 Validation datasets

In addition to our calibration dataset, we used two other independent datasets for validation purposes. These

datasets enabled us to test both the performance and the robustness of our models, and were not used for the

calibration. To assess prediction bias, we conducted a t-test on the residuals (observed minus predicted heights)

to determine whether the mean of the prediction error differed significantly from zero.

The internal validation dataset contains NFI data collected in 2023 (IGN, 2023). This data was collected

following the same protocol as the calibration dataset. It includes a large number of height measurements from a

large number of species and for the various stand structures. By applying the same criteria as for the calibration

dataset, we selected 8,771 height measurements from 2,548 plots, covering 40 species.

This second dataset will be referred to as the NFI_2023 dataset, and is described in Supplementary Materials,

table S3.

We also used an external dataset for validation, which included six species. This data was collected following a

different protocol than that of the NFI. The trees were measured between 1920 and 1955. The plots were sampled

among pure highly-productivity even-aged permanent plots established by the former French Forest

Administration. The measurement and selection protocol are described in Vallet et al. (2006).

This third dataset will be referred to as the FFA dataset, and is described in table 2.

Species	Number of trees	Number of plots	dbh (cm)			Height (m)		
			min	mean	max	min	mean	max
Abies alba	762	28	7.64	31.99	102.81	9.9	23.45	39.4
Fagus sylvatica	1,293	50	2.86	24.54	79.58	5.4	22.16	42.5
Pinus sylvestris	389	17	6.68	22.40	53.16	9.0	19.18	28.5
Picea abies	309	8	15.28	32.27	70.98	10.9	26.45	41.3
Pseudotsuga menziesii	224	7	4.77	14.95	34.06	5.5	15.12	25.0
Quercus petraea	1,222	58	4.46	23.15	89.45	6.0	20.55	39.5

227 *Table 2: Description of the French Forest Administration validation dataset*

229 3. Method

230 3.1 General principles of the study

Following the general principles for fitting allometric relationships (Mehtätalo et al., 2015; Zhou et al., 2023), we

developed generic models: for each species, the same equation was calibrated, incorporating both tree social status

(dbh/D_g) and stand variables. Stand basal area (BA) was used as a proxy for stand density, and quadratic mean

diameter (D_g) for development stage. We also included the effect of stand structure (even-aged, uneven-aged,

coppice-with-standards and coppice) on the height-diameter allometry. Our models were tested for accuracy and

robustness on both validation datasets.

Finally, to improve the height estimates for a given forest plot, we developed a local recalibration method based

on a limited number of height measurements.

3.2 Functions

3.2.1 General equation

Allometries can take many forms and curve shapes (Mehtätalo et al., 2015). To develop our models, we considered

commonly used functions (Ciceu et al., 2023). We selected three three-parameter sigmoidal functions: the

Chapman – Richards (Huang et al., 1992; Richards, 1959), Hossfeld II (Elfving and Kiviste, 1997; Vallet and

Perot, 2016) and Weibull (Weibull, 1951) functions.

As we aimed to develop generic models, we fitted the same equation to all the species. By comparing the Akaike

Information Criterion (AIC)(Akaike, 1974), we selected the Hossfeld II function (Eq.1), which gave the best

adjustments.

$$y = 1.3 + a_s * \frac{1}{1+\frac{b_s}{x^{c_s}}} \quad (1)$$

Parameters a_s, b_s and c_s (Eq.1) were species dependent. In this function, the height equals 1.3m when the diameter

tends to zero, and the sum of $1.3 + a_s$ is the asymptotic height value for large diameters. c_s is a shape coefficient,

and b_s is linked by a power function to the diameter at half asymptotic height (dha) as dha $= b_s^{1/c_s}$. The social

status of the tree was used as the explicative variable in order to take intra-plot competition into account.

3.2.2 Inclusion of stand variables in relation to tree growth and biological

constraints

When developing our models, we considered practical constraints related to fieldwork. Diameter measurements

for all trees can be rapidly and efficiently measured, and are necessary when evaluating stand characteristics such

as basal area and quadratic mean diameter. The use of a stand development stage variable, usually characterized

by its dominant height, can significantly improve the fit of height-diameter allometries (Adame et al., 2008; Ciceu

et al., 2023; Patrício et al., 2022). However, the quadratic mean diameter can be used as a substitute (Vallet and

Perot, 2016). The quadratic diameter has the advantage of not requiring any field measurements other than tree

diameter. Therefore, we chose to use basal area and quadratic mean diameter as the sole calculated stand variables,

thereby simplifying the necessary field measurements.

We developed our models to quantify the influence of stand structure on height-diameter allometry. The even-

aged stand structure was used as the reference condition. Other structures (uneven-aged, coppice-with-standards

and coppice) were added as multiplicative adjustments representing their relative effect compared to the even-aged

reference.

• a_s parameter

Stand variables were introduced into the asymptotic height parameter (a_s) as formulated in equation 2:

$$a_s = a_{s1} * \left(1 + a_{s,UA} + a_{s,CWS} + a_{s,C}\right) * \left(1 + a_{s2} * BA\right) * \left(1 - e^{-a_{s3}*Dg^{a_{s4}}}\right) \text{ (2)}$$

Where a_{s1} is the reference value corresponding to the even-aged structure, and $a_{s,UA}$, $a_{s,CWS}$ and $a_{s,C}$ are the

parameters expressing the relative difference in asymptotic height for uneven-aged, coppice-with-standards and

coppice stand structures, respectively, included as dummy variables. BA is the basal area (in m²/ha) of the stand,

D_g is the quadratic mean diameter (in cm) of the dominant species of the stand, and a_{s2}, a_{s3}, and a_{s4} their

corresponding parameters.

The quadratic mean diameter (D_g) was integrated into the model to reflect the biological assumption that dominant

height, and therefore asymptotic height, increases as the stand ages. This effect was modeled with a negative

exponential form. This form represents the relationship between quadratic diameter and dominant height, and also

reflects the saturating nature of height growth over time.

The addition of basal area reflects its demonstrated influence on maximum tree height, as supported by del Río et

al (2019) and Gonçalves (2022). Following similar modeling approaches (Adame et al., 2008; Temesgen and v.

Gadow, 2004), we used a linear form to incorporate basal area into the asymptotic parameter. This allowed us to

evaluate both the statistical significance and biological impact of stand basal area on individual tree height.

• b_s parameter

The b_s parameter, as the diameter at half asymptotic height, was developed as follows (equation 3):

$$b_s = b_{s1} * \left(1 + b_{s,UA} + b_{s,CWS} + b_{s,C}\right) * \left(e^{-b_{s2}*BA}\right) \text{ (3)}$$

Where b_{s1} is the reference value corresponding to the even-aged structure, $b_{s,UA}$, $b_{s,CWS}$ and $b_{s,C}$ are the

structure-specific adjustment parameters, added as dummy variables.

Structure-specific effects were added similarly to those for a_s. b_{s2} is the parameter associated with basal area. The

exponential form for basal area was chosen to satisfy two criteria: i) the b parameter must remain positive to ensure

realistic and biologically acceptable height estimates; ii) as basal area increases, diameter growth is limited due to

competition, but height growth may continue at a similar rate (Eichhorn law, (Decourt, 1973)). Therefore, for a

given diameter, small-diameter trees will be taller in a denser stand.

3.2.3 Final calibrated equation

When all stand variables are taken into account, equation 4 results as follows:

$$h_{tot} = 1,3 + (1 + \lambda_i) * a_{s1} * \left(1 + a_{s,UA} + a_{s,CWS} + a_{s,C}\right) * (1 + a_{s2} * BA) * \left(1 - \right.$$

$$\left. exp\left(-a_{s3} * D_g{}^{a_{s4}}\right)\right) * \frac{1}{1 + \frac{b_{s1}*(1 + b_{s,UA} + b_{s,CWS} + b_{s,C})*(exp(-b_{s2}*BA))}{\left(\frac{dbh}{D_g}\right)^{c_s}}} \quad (4)$$

Where h_{tot} is height in m, BA is basal area in m²/ha, and D_g is quadratic mean diameter in cm. λ_i refers to the

random effect calibrated (described in 3.3.1). When applying the model, its value can either be 0 (using only the

fixed effects of the model) or estimated using local recalibration (described in 3.4).

3.3   Statistical choices

To take spatial dependency, due to trees sampled per plots, into account, we used non-linear mixed-effect models

(NLME) (Pinheiro and Bates, 2006). The model included a plot random effect λ_i, where i identified each plot

(equation 4). The λ_i parameter will henceforth be referred to as the site parameter.

The residual model variance was not constant as tree height variance generally increases with tree size. Moreover,

as the trees were inventoried in subplots of different radii, tree height variance was typically smaller for the smallest

trees in each category, and higher for the largest. To take these two effects into account, we used a varIdent variance

function from the "nlme" package (Pinheiro and Bates, 2006). We defined ten variance groups by splitting each

of the five NFI diameter categories (very small trees, small trees, medium trees, large trees, and very large trees)

at their median value.

3.4   Local recalibration process

3.4.1 Principle

The local recalibration approach consisted in estimating the random site parameter (λ_i) with data that could easily

be collected in the field: the height of a small subsample of trees. This allowed the height-diameter allometric

relationship to be adjusted to each plot with only a few height measurements, as illustrated in figure 1, and

demonstrated in Supplementary Materials, section C.

Figure 1: Example of local recalibration for a given Fagus sylvatica plot. The three trees with the largest diameter

and the one tree with the smallest diameter were used to estimate the site parameter associated to the plot - FFA

dataset.

We estimated $\lambda_{m,j}$ with the mean-based approach shown in equation 5, relying only on a subset of N_j trees from

the plot j:

$$\lambda_{m,j} = \frac{1}{N_j}\sum\left(\frac{h_{tot,ij}-1,3}{\hat{h_{0,ij}}-1,3}\right) - 1 \quad (5)$$

We then assessed the optimal tree subsample that would allow the site parameter to be accurately estimated and

reduce the root-mean-square error (RMSE) at best, while minimizing field effort.

3.4.2 Sub-samples and performance evaluation

We estimated $\lambda_{m,j}$ with different samples of tree height. These samples varied in terms of size (number of trees)

and in terms of relative tree diameter. We selected one to six trees according to three sampling strategies:

selecting only the thinnest trees, only the largest trees, or a combination of both diameter classes. For this last

strategy, we evaluated two scenarios. In the first scenario, large-diameter trees were predominant, following the

distribution: 1 large; 1 large with 1 small; 2 large with 1 small; 3 large with 1 small; 3 large with 2 small and 4

large with 2 small. In the second scenario, small-diameter trees were predominant and distributed as: 1 small; 1

small with 1 large; 2 small with 1 large; 3 small with 1 large; 3 small with 2 large and 4 small with 2 large.

We applied this method to all FFA dataset plots, that had at least 13 tree records. The six largest diameter trees

and the six thinnest, which were used for site parameter estimation (following the different combination

described above), were then removed from the dataset. We then calculated the RMSE associated to the height

prediction for each type of sub-samples, for all remaining individuals. We then identified the best diameter-class

combinations based on the root-mean-square error RMSE difference relative to (i) the RMSE obtained when

height was predicted using fixed effects only – and (ii) the RMSE obtained using the optimal site-parameter

value, where height was predicted using $\lambda_{m,j}$ as the site parameter and all available trees.

The best performing combination was then re-applied to the FFA dataset. In this second step, only the trees

involved in the selected combination (ie., 6 trees of the largest, smallest or a mix of both), were removed from

each plot, instead of systematically removing the six smallest and six largest diameter trees. RMSE was then

calculated using all remaining individuals. To evaluate the efficiency of the local recalibration method, and

identify the optimal number of trees needed, we calculated the relative RMSE improvement (RRI) as the

difference between the fixed-effect model RMSE and the recalibrated model RMSE, divided by the fixed-effect

RMSE.

4. Results

4.1 Calibration

Our models highlight a positive influence of quadratic mean diameter and basal area on the asymptotic height, and

a negative impact of basal area on the diameter at half asymptotic height (i.e., half asymptotic height is reached

for smaller diameters) (Table 3 and table S5).

The effect of stand structure on the allometric curve was species-dependent. Parameter estimates (figure 2)

revealed a predominantly negative effect of coppice structure on asymptotic height compared to even-aged stands,

and a strong similarity between even-aged and coppice-with-standards stands. In contrast, the effect of an uneven-

aged structure on asymptotic height differed by species group, being positive for conifers and negative for

broadleaved species.

For the diameter at half asymptotic height, structural effects were substantially more mixed, with positive, negative

and non-significant responses observed across species. All effects for the ten most represented species are

summarized in Figure 3 and in the Supplementary Materials (figures S2 - S11).

Species	Estimation and Statistics parameter	a_{s1}	$a_{s,UA}$	$a_{s,CWS}$	$a_{s,C}$	a_{s2}	a_{s3}	a_{s4}	b_{s1}	$b_{s,UA}$	$b_{s,CWS}$	$b_{s,C}$	b_{s2}	c_s	Residual deviation	Deviation of random effects
Abies alba	estimate	59.27	0.1295	0.0106	-	0.0003	0.0200	1.0637	0.7621	0.5553	0.0832	-	0.0094	1.5730	2.04	0.13
	significativity	***	***	.	-	.	***	***	***	***	***	-	***	***		
	stddev	2.64	0.01113	0.01003	-	0.00027	0.00108	0.03178	0.02310	0.02907	0.02282	-	0.00051	0.01615		
Castanea sativa	estimate	21.36	-0.0798	-0.0260	0.0096	0.0031	0.0762	1.0608	0.3808	0.0305	-0.0671	0.1576	0.0097	1.5726	1.67	0.21
	significativity	***	**	*	.	***	***	***	***	.	*	***	***	***		
	stddev	0.52	0.02736	0.01280	0.01654	0.00045	0.00926	0.05363	0.02140	0.08879	0.03017	0.04726	0.00085	0.03545		
Fagus sylvatica	estimate	49.75	0.0412	-0.0338	-0.1867	0.0005	0.0318	0.9863	0.4410	0.5103	0.0530	0.1154	0.0029	1.5511	2.38	0.19
	significativity	***	*	***	***	.	***	***	***	***	**	.	***	***		
	stddev	1.48	0.01789	0.00757	0.01948	0.00028	0.00134	0.02421	0.01293	0.05093	0.01709	0.06320	0.00053	0.01841		
Picea abies	estimate	39.91	0.07737	0.01371	-	0.00246	0.01509	1.26384	0.43086	0.65289	0.06797	-	0.00797	1.93804	1.92	0.15
	significativity	***	***	.	-	***	***	.	***	***	*	-	***	***		
	stddev	1.06	0.01273	0.01118	-	0.00028	0.00107	0.03250	0.01399	0.03865	0.02768	-	0.00049	0.02503		
Pinus pinaster	estimate	35.06	0.23787	-0.01374	-	0.00092	0.00967	1.40117	0.47802	1.17399	0.04229	-	0.00921	1.42384	1.18	0.15
	significativity	***	***	.	-	**	***	***	***	***	.	-	***	***		
	stddev	0.71	0.02923	0.00899	-	0.00030	0.00036	0.01761	0.01922	0.08918	0.02406	-	0.00065	0.02941		
Pinus sylvestris	estimate	103.55	-0.08569	0.01118	-	0.01033	0.00743	0.91605	0.32923	0.26842	0.04402	-	0.00518	1.81832	2.32	0.25
	significativity	.	***	.	-	***	.	***	***	***	.	-	***	***		
	stddev	86.87	0.01670	0.01188	-	0.00063	0.00534	0.06061	0.01348	0.05030	0.02640	-	0.00072	0.03242		
Pseudotsuga menziesii	estimate	45.07	0.20845	0.01852	-	0.00160	0.00718	1.45561	0.39143	0.97330	0.05445	-	0.00652	1.74230	1.85	0.13
	significativity	***	***	.	-	***	***	***	***	***	.	-	***	***		
	stddev	1.01	0.0377	0.0111	-	0.0003	0.0004	0.0234	0.0153	0.1216	0.0299	-	0.0005	0.0338		
Quercus petraea	estimate	32.63	-0.0534	-0.0294	-0.0721	0.0060	0.0330	1.0137	0.1874	0.5170	-0.0003	0.6569	0.0061	2.1660	2.02	0.15
	significativity	***	**	***	***	***	***	***	***	***	.	***	***	***		
	stddev	0.54	0.01642	0.00470	0.01659	0.00032	0.00121	0.01770	0.00609	0.08860	0.01602	0.09265	0.00080	0.02605		
Quercus pubescens	estimate	22.58	-0.06542	0.00810	-0.11512	0.01122	0.04416	0.99286	0.42095	0.06926	0.02891	0.16874	0.00583	1.70921	1.27	0.24
	significativity	***	**	.	***	***	***	***	***	.	.	***	***	***		
	stddev	1.2	0.02237	0.01155	0.01390	0.00071	0.00293	0.04387	0.01796	0.06035	0.02370	0.04067	0.00090	0.03229		
Quercus robur	estimate	30.27	-0.08605	0.00685	-0.00597	0.00396	0.03890	0.99302	0.27664	0.16480	0.02373	0.28099	0.00844	1.91118	2.11	0.17
	significativity	***	***	.	.	***	***	***	***	**	.	***	***	***		
	stddev	0.55	0.01471	0.00620	0.02071	0.00033	0.00176	0.02094	0.00985	0.06260	0.01900	0.08086	0.00080	0.02957		

*Table 3: Model parameters and statistics for fitting of equation 4 for the ten most represented species. Significance: ***: p-value<0.001, **: p-value <0.01, *: p-value <0.05*

Figure 2: Stand structure effects on asymptotic height and diameter at half asymptotic height, compared to the

even-aged reference. Estimates and associated standard deviations are presented by structure and species. Due

to high non-significant values, estimates and standard deviations for Quercus cerris are not presented for

parameters $a_{s,CWS}$ and $b_{s,CWS}$. Standard deviation was not represented for Pinus pinea for $b_{s,CWS}$, nor for Corylus

avellana for $b_{s,C}$.

Figure 3: Presentation of the Quercus petraea model, for varying basal area, quadratic diameter and stand

characteristics. Predictions are limited to the range of diameter values observed for each structure in the

calibration dataset.

4.2 Validation (fixed effects only)

We calculated the errors associated to $\hat{h_0}$. Without the site parameter, height estimates were contained between

three and five meters for the calibration dataset. Table 4 highlights similar results for model errors between the

calibration and NFI_2023 datasets. As expected, the results for the FFA dataset were less similar to the calibration

dataset. A bias in estimation can be observed when predicted height is confronted to observed height (figure 4).

Models' residual graphs are presented in supplementary materials, section G. Results of the one-sample t-test

showed a significant bias of -0.22m on the NFI_2023 dataset. For the FFA dataset, the t-test showed a significant

difference of 2.75m.

Species	Calibration dataset		NFI_2023		FFA	
	$RMSE_0$(m)	Nb obs.	$RMSE_0$(m)	Nb obs.	$RMSE_0$(m)	Nb obs.
Abies alba	3.5551	19 055	3.5644	606	2.7906	618
Fagus sylvatica	4.535	28 135	5.1654	853	5.0553	816
Picea abies	3.7235	19 176	3.8088	390	4.4948	287
Pinus sylvestris	3.8592	21 865	4.2631	496	5.3985	246
Pseudotsuga menziesii	3.3942	13 691	3.3494	475	2.611	212
Quercus petraea	3.5309	31 927	3.6339	1 187	3.5484	830

392 *Table 4: Validation results for 6 species across all datasets. For the calibration dataset and NFI_2023, results*

393 *aggregate all stand structure types. Only even-aged structures are present in the FFA dataset.*

394 *Figure 4: Observed height compared to predicted height for the whole NFI_2023 dataset (A – 40 species; 8771*

395 *observations) and the FFA dataset (B – 6 species; 3009 observations). The black line represents perfect*

396 *prediction (1:1). The red line shows the overall trend in the data (lowess regression).*

4.3   Local recalibration

398 The selection of a combination of trees with a predominance in trees with the largest diameter reduced RMSE the

399 most (figure 5). From one to three trees, adding an additional tree provided a strong reduction in RMSE; thereafter,

400 adding more trees only slightly improved the reduction in RMSE.

Figure 5: RMSE values for tree height prediction with recalibration, with different tree sub-samples based on

number and diameter class of the trees on the calibration dataset, per species (Quercus petraea (A) and Pinus

sylvestris (B))

The root-mean-square error of tree height predictions following local recalibration applied to the FFA dataset, with

sub-samples of one to six trees with a predominance of large diameter trees, are presented in table 5.

Estimating the site parameter with all trees present on the plot (optimal value) consistently improved prediction

accuracy across species, with relative RMSE improvement (RRI) ranging from 20 to 70 %, compared to fixed

effects.

When the site parameter was estimated from a sub-sample of three to four trees (2 large with 1small and 3large

with 1 small), substantial improvements in RMSE generally resulted, compared to using fixed effects only. The

RRI ranges from 10 to 70%, except for *Abies alba*, for which the use of sub-samples led to higher RMSEs than

the fixed-effects baseline. This is likely due to *Abies alba* plots displaying a height-diameter relationship closer to

those of the calibration dataset than those of the other species.

Site parameter estimates		Abies alba		Fagus sylvatica		Picea abies		Pinus sylvestris		Pseudotsuga menziesii		Quercus petraea	
		RMSE (m)	RRI (%)	RMSE (m)	RRI (%)	RMSE (m)	RRI (%)	RMSE (m)	RRI (%)	RMSE (m)	RRI (%)	RMSE (m)	RRI (%)
	Fixed effects	2.66	-	5.06	-	4.40	-	5.46	-	2.65	-	3.51	-
	Optimal value	1.98	25.76	1.84	63.60	2.42	44.86	1.21	77.77	1.26	52.39	1.57	55.45
Number of trees selected among the largest	1	3.00	-12.64	3.44	31.92	2.87	34.82	1.90	65.21	2.57	2.82	2.15	38.94
	2	3.35	-25.56	2.29	54.79	3.70	15.95	1.53	72.02	2.81	-6.04	1.89	46.33
	3	2.82	-5.66	2.06	59.38	3.08	29.95	1.33	75.68	2.41	8.81	1.76	49.95
	4	2.69	-0.85	2.15	57.57	2.80	36.27	1.31	75.97	2.22	16.08	1.77	49.57
	5	2.50	6.29	2.02	60.04	2.95	32.92	1.30	76.24	2.17	17.85	1.76	49.99
	6	2.42	9.06	2.05	59.52	2.74	37.75	1.28	76.51	1.99	24.96	1.69	52.03
Number of trees		480		652		245		176		176		648	

Table 5: Local recalibration results for the FFA dataset, compared to fixed-effects-only height prediction. RMSE

in meters and relative RMSE improvement (RRI) in percentage, for height predictions with fixed effects, optimal

RMSE value, and sub-samples of one to six trees selected among the largest and smallest diameters with

predominant large trees, for the FFA dataset. RRI is calculated relative to the fixed-effects baseline.

5. Discussion

5.1 Effects of stand characteristics on height-diameter allometry

5.1.1 Influence of stand structure

Stand structure had a significant effect on height-diameter allometry for 30 species. However, a few species,

such as *Populus tremula*, showed no significant parameter estimates for any of the stand structures. Overall, the

stand structure parameter estimates (figure 2) highlight the influence of forest management on allometric

relationships.

In coppice stands, results were consistent with the regeneration method (vegetative sprouting from existing root

system), which generally leads to reduced tree height growth compared to even-aged stands for a given diameter.

However, the diameter at half asymptotic height (dha) was often greater than in even-aged conditions. This shift

suggests a reallocation of resources toward radial rather than vertical growth, a pattern observed in previous

studies (Corcuera et al., 2006; Deforce and Haneca, 2015).

In uneven-aged stands, competition between individuals is generally reduced compared to that in even-aged

stands, because density is lower. As highlighted by del Rìo et al. 2019, higher competition leads to greater tree

heights for a given diameter, due to the increased drive for vertical growth under denser conditions. This pattern

is consistent with the fact that reduced competition, particularly for light in uneven-aged stands, allows

individuals to allocate more resources to radial growth than to height. As a result, trees in uneven-aged stands

tend to be of smaller height for a given diameter, leading to a negative effect on asymptotic height compared to

trees in even-aged stands. Our results for broadleaved species were consistent with this observation. However,

coniferous species showed a positive effect on asymptotic height, which can partly be explained by the

correlation between asymptotic height and dha parameters: the diameter range for these species corresponds to a

section of the curves where height is greater for even-aged stands even if the asymptotic height is lower (the

same effect as in Supplementary Materials, section F).

Additionally, our results indicate an increase in dha in uneven-aged stands, which is consistent with lesser light

competition compared to even-aged stands. Under high competition, such as in even-aged stands, trees prioritize

height growth over radial growth, shifting the allometry curve to the left. Therefore, for a given diameter (among

small trees), trees in uneven-aged stands tend to be shorter than those in even-aged stands (trends for parameters

a and b in the HossfledII equation (Vallet and Perot, 2016)).

451   5.1.2 Influence of basal area

Tree height for a given diameter can increase with stand density. Basal area (BA), often used as a proxy for stand

density, also serves as an index for competition among trees. As basal area increases, competition for resources

intensifies, and trees tend to allocate more energy to vertical growth in order to maintain access to light.

Consequently, for a given diameter, tree height is expected to be greater in denser stands, as highlighted by del

Río et al. (2019) and Gonçalves (2022).

In our models, this effect is reflected through an increase in asymptotic height and a corresponding decrease in the

diameter at half asymptotic height (parameters a_{s2} and b_{s2}), consistent with our expectations. As a result, the

height-diameter allometric curve shifts upward: for a given diameter, predicted height increases with basal area

(Figure 2). However, this trend did not apply uniformly across all species. For *Corylus avellana*, *Pinus halepensis*

and *Tilia cordata*, basal area had a significant negative effect on asymptotic tree height. As illustrated in figure

S12 (Supplementary Materials, section F), in the case of *Tilia cordata*, the high effect of basal area over diameter

at half asymptotic height, associated with a lack of observation for large diameters, tended to bend the allometric

curve to the left, inducing a reduction in asymptotic height by mathematical construction. Hence, as for some

effects in uneven-aged stands, for the observed range of diameters of these species, the height is still higher with

increasing BA even if the asymptotic parameter is negative.

In addition to its effect on asymptotic height, increased competition directly influences diameter growth. As emphasized by Aleinikovas et al. (2014), increased competition has a negative effect on tree diameter growth. As a consequence, the height-diameter curve is shifted to the left, indicating greater heights for a given diameter.

5.1.3 Influence of quadratic mean diameter

Quadratic mean diameter provides insights into the developmental stage of the stand (Aussenac et al., 2023; Sharma et al., 2019a). The silvicultural system applied to each structure will induce different growth dynamics. For example, even-aged stands are typically allowed to grow for several decades, enabling trees to reach greater heights and diameters. In contrast, coppice systems imply periodic cutting, and the trees regenerate from stump sprouts, resulting in shorter growth periods, and more limited development in size. In both uneven-aged and coppice structures, Dg values tended to remain relatively stable over time, due to the continuous or cyclical nature of their management regimes.

Across all species studied, Dg had a positive effect on the asymptotic height in the height diameter allometry. As Dg increased, the height asymptote also increased. This pattern aligns with findings in the literature, which report a strong relationship between Dg, dominant height, and height-diameter allometry (Schmidt et al., 2018; Sharma et al., 2019a). The increase in asymptotic height with Dg reflects a typical stand development trajectory, where older and more developed stands support taller dominant trees.

5.2 Optimization of model accuracy and predictive performance

5.2.1 Quality of the models

Using National Forest Inventory (NFI) data enabled the inclusion of a wide range of species and tree records, covering diverse climatic, structural and edaphic conditions across the French mainland territory. The robustness of the dataset increases the confidence in applying our models across Europe as well, as was proven possible for studies on growth (Mahnken et al., 2022). Additionally, the large size of the dataset reduced the influence of outliers during calibration, enhancing parameter stability and reliability. The RMSE obtained for each species are in line with other studies, where the models selected had a RMSE between 1 and 3.5 meters (Adame et al., 2008; Gómez-García et al., 2015; Mehtätalo et al., 2015; Vázquez-Veloso et al., 2025).

To assess the efficiency of including a stand structure effect in our models, we evaluated the RMSE associated with applying the even-aged calibrated model to plots with different stand structures and compared it to structure-specific model errors. Estimations were done using only fixed effects. The relative error difference

calculated for the calibration dataset (Table 6) indicates a strong improvement in model accuracy with structure-

specific models for uneven-aged and coppice stand structures. In contrast, applying the even-aged model to

coppice-with-standards plots revealed a high degree of similarity between the two structures, as reflected by the

relatively small RMSE differences. As mentioned in Gonçalves (2022), using even-aged models to predict tree

height in uneven-aged stands can lead to estimation errors of up to 20%. Our models show similar degrees of

difference between estimation errors (table 6). Our results were obtained with only the fixed effects of our

models, which explains the negative result for *Pseudotsuga menziesii*. Thus, adding stand structure to our models

improved the accuracy of estimated tree height across diverse management systems, and provided insights into

less studied structures and how they differ from the even-aged structure.

Finally, model bias observed on validation datasets can largely be attributed to the lack of site-specific information.

This limitation arises from applying a generalized model to a specific region, a known source of error when site

fertility or climate diverges from the average conditions of the calibration dataset (Mensah et al., 2018).

Nevertheless, the low bias observed on the NFI_2023 dataset suggests high consistency with the calibration data.

In contrast, the underestimation of tree height in the FFA dataset was expected, as these plots concern more fertile

areas than do the calibration plots.

Species	Even-aged model applied to stand structure		
	Uneven-aged	Coppice-with-standards	Coppice
Abies alba	+6.8	+1.2	
Castanea sativa	+15.5	+1.9	+2.0
Fagus sylvatica	+11.7	+5.3	+50.8
Picea abies	+8.4	+0.2	
Pinus pinaster	+9.8	+3.6	
Pinus sylvestris	+26.4	-0.3	
Pseudotsuga menziesii	-1.2	-0.3	
Quercus petraea	+16.5	+3.9	+24.8
Quercus pubescens	+10.9	-0.2	+21.8
Quercus robur	+19.5	-0.2	+6.4

513   *Table 6: Relative difference in RMSE (%) when applying models calibrated for even-aged plots to stands with*

514   *other structures. Positive values indicate higher prediction error for the even-aged model than for structure-*

515   *specific models, while negative values indicate better accuracy for the even-aged model. These results are based*

516   *on the calibration dataset.*

517   5.2.2 Benefits of local recalibration

This approach significantly improved performance on the FFA dataset, especially where site fertility deviated from

the average. Our recalibration strategy focused on a predominant number of trees with the largest diameters, in

line with recommendations from Sullivan et al. (2018), who showed that using large-diameter trees improves

recalibration outcomes. Furthermore, as shown by Sharma and Breidenbach (2015), including trees from different

species when estimating site parameters induces error. We therefore performed species-specific recalibrations.

5.3   Limits

We chose to develop a single model structure that could be adapted to different stand types by adding structure

parameters. While this approach simplifies generalization and cross-species comparison, it may have limitations.

For example, coppice systems, which are managed through frequent cutting cycles, pose particular modeling

challenges and can complicate convergence when the same stand variables as those applied to other structures

are used. In uneven-aged stands, the management process tends to maintain basal area and quadratic mean

diameter within a narrow range, unlike in even-aged stands. Consequently, using the same variables to

characterize structurally distinct stands may reduce model relevance or performance in some cases.

We also used a single functional form across species, selected based on the overall AIC comparison. For a few

less-represented species, alternative functional forms would have produced lower AIC values. Developing

species-specific functions could improve model accuracy, but maintaining a consistent equation enables

parameter comparisons across species and the development of generic models for practical use.

We also used species-specific models to test each species' response to competition and stand dynamics.

5.4   Implications for forest modelling and management

The models developed in this study account for both stand structure and dendrometrical characteristics,

providing highly accurate height estimates across different management systems (even-aged, uneven-aged,

coppice and coppice-with-standards). This represents a notable advancement in modeling, and can improve

estimates of other critical forest variables such as timber volume and carbon stocks.

Our validation results show strong predictive performance in plots with characteristics similar to the calibration

dataset. In plots with high fertility or atypical conditions, applying local recalibration reduced bias (see practical

method in Supplementary Materials, section C). The number of trees required for recalibration can be adjusted

depending on the desired accuracy of the height estimates, but as few as one to four trees could already strongly

improve the results (Table 5).

6. Conclusion

Based on data from more than 40,000 stands surveyed by the French National Forest Inventory, we developed height-diameter allometries for 41 European species that take into account stand characteristics. We found that stand structure (even-aged, uneven-aged, coppice-with-standards and coppice) did significantly impact height-diameter allometry, as did the quadratic mean diameter and the basal area of the stand. Given the large diversity of climates, stand structures and soil fertility present in the French metropolitan area, the models developed here could be applied all over Europe. Moreover, to address the loss of local accuracy when applying a large-scale model at stand level, we developed a recalibration method requiring only a few height measurements on the largest trees in the stand. Measuring one to four trees can considerably reduce estimation errors.

References

Adame, P., del Río, M., Cañellas, I., 2008. A mixed nonlinear height–diameter model for pyrenean oak (*Quercus pyrenaica* Willd.). For. Ecol. Manag. 256, 88–98. https://doi.org/10.1016/j.foreco.2008.04.006

Akaike, H., 1974. A new look at the statistical model identification. IEEE Trans. Autom. Control 19, 716–723. https://doi.org/10.1109/TAC.1974.1100705

Aleinikovas, M., Linkevičius, E., Kuliešis, A., Rolhle, H., Schroeder, J., 2014. The Impact of Competition for Growing Space on Diameter, Basal Area and Height Growth in Pine Trees. Balt. For. 20.

Aussenac, R., Monnet, J.-M., Klopčič, M., Hawryło, P., Socha, J., Mahnken, M., Gutsch, M., Cordonnier, T., Vallet, P., 2023. Diameter, height and species of 42 million trees in three European landscapes generated from field data and airborne laser scanning data. Open Res. Eur. 3, 32. https://doi.org/10.12688/openreseurope.15373.2

Bettinger, P., Boston, K., Siry, J.P., Grebner, D.L., 2017. Estimation and Projection of Stand and Forest Conditions, in: Forest Management and Planning. Elsevier, pp. 87–111. https://doi.org/10.1016/B978-0-12-809476-1.00004-7

Ciceu, A., Chakraborty, D., Ledermann, T., 2023. Examining the transferability of height–diameter model calibration strategies across studies. For. Int. J. For. Res. cpad063. https://doi.org/10.1093/forestry/cpad063

Ciceu, A., Garcia-Duro, J., Seceleanu, I., Badea, O., 2020. A generalized nonlinear mixed-effects height–diameter model for Norway spruce in mixed-uneven aged stands. For. Ecol. Manag. 477, 118507. https://doi.org/10.1016/j.foreco.2020.118507

Corcuera, L., Camarero, J.J., Sisó, S., Gil-Pelegrín, E., 2006. Radial-growth and wood-anatomical changes in overaged Quercus pyrenaica coppice stands: functional responses in a new Mediterranean landscape. Trees 20, 91–98. https://doi.org/10.1007/s00468-005-0016-4

Dalmasso, M., Derrière, N., Wurpillot, S., 2014. Les données de l'inventaire forestier : Etat des lieux et évolution [WWW Document]. URL https://inventaire-forestier.ign.fr/IMG/pdf/LP_1.pdf (accessed 7.18.24).

Decourt, N., 1973. Production primaire, production utile : méthodes d'évaluation, indices de productivité. Ann. Sci. For. 30, 219–237. https://doi.org/10.1051/forest/19730302

Deforce, K., Haneca, K., 2015. Tree-ring analysis of archaeological charcoal as a tool to identify past woodland management: The case from a 14th century site from Oudenaarde (Belgium). Quat. Int., Charcoal: resource and ubiquitous proxy 366, 70–80. https://doi.org/10.1016/j.quaint.2014.05.056

del Río, M., Bravo-Oviedo, A., Ruiz-Peinado, R., Condés, S., 2019. Tree allometry variation in response to intra- and inter-specific competitions. Trees 33, 121–138. https://doi.org/10.1007/s00468-018-1763-3

Elfving, B., Kiviste, A., 1997. Construction of site index equations for *Pinus sylvestris* L. using permanent plot data in Sweden. For. Ecol. Manag. 98, 125–134. https://doi.org/10.1016/S0378-1127(97)00077-7

Feldpausch, T.R., Banin, L., Phillips, O.L., Baker, T.R., Lewis, S.L., Quesada, C.A., Affum-Baffoe, K., Arets, E.J.M.M., Berry, N.J., Bird, M., Brondizio, E.S., de Camargo, P., Chave, J., Djagbletey, G., Domingues, T.F., Drescher, M., Fearnside, P.M., França, M.B., Fyllas, N.M., Lopez-Gonzalez, G., Hladik, A., Higuchi, N., Hunter, M.O., Iida, Y., Salim, K.A., Kassim, A.R., Keller, M., Kemp, J., King, D.A., Lovett, J.C., Marimon, B.S., Marimon-Junior, B.H., Lenza, E., Marshall, A.R., Metcalfe, D.J., Mitchard, E.T.A., Moran, E.F., Nelson, B.W., Nilus, R., Nogueira, E.M., Palace, M., Patiño, S., Peh, K.S.-H., Raventos, M.T., Reitsma, J.M., Saiz, G., Schrodt, F., Sonké, B., Taedoumg, H.E., Tan, S., White, L., Wöll, H., Lloyd, J., 2011. Height-diameter allometry of tropical forest trees. Biogeosciences 8, 1081–1106. https://doi.org/10.5194/bg-8-1081-2011

Gómez-García, E., Fonseca, T.F., Crecente-Campo, F., Almeida, L.R., Diéguez-Aranda, U., Huang, S., Marques, C.P., 2015. Height-diameter models for maritime pine in Portugal: a comparison

of basic, generalized and mixed-effects models. IForest - Biogeosciences For. 9, 72.
https://doi.org/10.3832/ifor1520-008
Gonçalves, A.C., 2022. Stand Structure Impacts on Forest Modelling. Appl. Sci. 12, 6963.
https://doi.org/10.3390/app12146963
Huang, S., Titus, S.J., Wiens, D.P., 1992. Comparison of nonlinear height–diameter functions for
major Alberta tree species. Can. J. For. Res. 22, 1297–1304. https://doi.org/10.1139/x92-172
Hunter, M.O., Keller, M., Vitoria, D., Morton, D.C., 2013. Tree height and tropical forest biomass
estimation. Biogeosciences 108385-8399. https://doi.org/10.5194/bg-10-8385-2013
IGN, 2023. IGN – Inventaire forestier national français, Données brutes, Campagnes annuelles 2005 et
suivantes [WWW Document]. URL https://inventaire-forestier.ign.fr/dataIFN/ (accessed
5.27.25).
Kearsley, E., Moonen, P.C., Hufkens, K., Doetterl, S., Lisingo, J., Boyemba Bosela, F., Boeckx, P.,
Beeckman, H., Verbeeck, H., 2017. Model performance of tree height-diameter relationships
in the central Congo Basin. Ann. For. Sci. 74, 1–13. https://doi.org/10.1007/s13595-016-0611-
0
Mehtätalo, L., de-Miguel, S., Gregoire, T.G., 2015. Modeling height-diameter curves for prediction.
Can. J. For. Res. 45, 826–837. https://doi.org/10.1139/cjfr-2015-0054
Mensah, S., Pienaar, O.L., Kunneke, A., du Toit, B., Seydack, A., Uhl, E., Pretzsch, H., Seifert, T.,
2018. Height – Diameter allometry in South Africa's indigenous high forests: Assessing
generic models performance and function forms. For. Ecol. Manag. 410, 1–11.
https://doi.org/10.1016/j.foreco.2017.12.030
Mugasha, W.A., Mauya, E.W., Njana, A.M., Karlsson, K., Malimbwi, R.E., Ernest, S., 2019. Height-
Diameter Allometry for Tree Species in Tanzania Mainland. Int. J. For. Res. 2019, 4832849.
https://doi.org/10.1155/2019/4832849
Patrício, M.S., Dias, C.R.G., Nunes, L., 2022. Mixed-effects generalized height-diameter model: A
tool for forestry management of young sweet chestnut stands. For. Ecol. Manag. 514, 120209.
https://doi.org/10.1016/j.foreco.2022.120209
Pilli, R., Anfodillo, T., Carrer, M., 2006. Towards a functional and simplified allometry for estimating
forest biomass. For. Ecol. Manag. 237, 583–593. https://doi.org/10.1016/j.foreco.2006.10.004
Pinheiro, J.C., Bates, D., 2006. Mixed-Effects Models in S and S-PLUS | SpringerLink [WWW
Document]. URL https://link.springer.com/book/10.1007/b98882 (accessed 8.15.24).
Richards, F.J., 1959. A Flexible Growth Function for Empirical Use. J. Exp. Bot. 10, 290–301.
https://doi.org/10.1093/jxb/10.2.290
Schmidt, M., Breidenbach, J., Astrup, R., 2018. Longitudinal height-diameter curves for Norway
spruce, Scots pine and silver birch in Norway based on shape constraint additive regression
models. For. Ecosyst. 5, 9. https://doi.org/10.1186/s40663-017-0125-8
Sharma, R.P., Breidenbach, J., 2015. Modeling height-diameter relationships for Norway spruce, Scots
pine, and downy birch using Norwegian national forest inventory data. For. Sci. Technol. 11,
44–53. https://doi.org/10.1080/21580103.2014.957354
Sharma, R.P., Štefančík, I., Vacek, Z., Vacek, S., 2019a. Generalized Nonlinear Mixed-Effects
Individual Tree Diameter Increment Models for Beech Forests in Slovakia. Forests 10, 451.
https://doi.org/10.3390/f10050451
Sharma, R.P., Vacek, Z., Vacek, S., Kučera, M., 2019b. Modelling individual tree height–diameter
relationships for multi-layered and multi-species forests in central Europe. Trees 33, 103–119.
https://doi.org/10.1007/s00468-018-1762-4
Sullivan, M.J.P., Lewis, S.L., Hubau, W., Qie, L., Baker, T.R., Banin, L.F., Chave, J., Cuni-Sanchez,
A., Feldpausch, T.R., Lopez-Gonzalez, G., Arets, E., Ashton, P., Bastin, J., Berry, N.J.,
Bogaert, J., Boot, R., Brearley, F.Q., Brienen, R., Burslem, D.F.R.P., De Canniere, C.,
Chudomelová, M., Dančák, M., Ewango, C., Hédl, R., Lloyd, J., Makana, J., Malhi, Y.,
Marimon, B.S., Junior, B.H.M., Metali, F., Moore, S., Nagy, L., Vargas, P.N., Pendry, C.A.,
Ramírez-Angulo, H., Reitsma, J., Rutishauser, E., Salim, K.A., Sonké, B., Sukri, R.S.,
Sunderland, T., Svátek, M., Umunay, P.M., Martinez, R.V., Vernimmen, R.R.E., Torre, E.V.,
Vleminckx, J., Vos, V., Phillips, O.L., 2018. Field methods for sampling tree height for
tropical forest biomass estimation. Methods Ecol. Evol. 9, 1179–1189.
https://doi.org/10.1111/2041-210X.12962

Temesgen, H., v. Gadow, K., 2004. Generalized height–diameter models—an application for major
tree species in complex stands of interior British Columbia. Eur. J. For. Res. 123, 45–51.
https://doi.org/10.1007/s10342-004-0020-z

Teobaldelli, M., Somogyi, Z., Migliavacca, M., Usoltsev, V.A., 2009. Generalized functions of
biomass expansion factors for conifers and broadleaved by stand age, growing stock and site
index. For. Ecol. Manag. 257, 1004–1013. https://doi.org/10.1016/j.foreco.2008.11.002

Trouvé, R., Bontemps, J.-D., Seynave, I., Collet, C., Lebourgeois, F., 2015. Stand density, tree social
status and water stress influence allocation in height and diameter growth of Quercus petraea
(Liebl.) | Tree Physiology | Oxford Academic [WWW Document]. URL
https://academic.oup.com/treephys/article/35/10/1035/2364497 (accessed 8.8.24).

Vallet, P., Dhôte, J.-F., Moguédec, G.L., Ravart, M., Pignard, G., 2006. Development of total
aboveground volume equations for seven important forest tree species in France. For. Ecol.
Manag. 229, 98–110. https://doi.org/10.1016/j.foreco.2006.03.013

Vallet, P., Perot, T., 2016. Tree diversity effect on dominant height in temperate forest. For. Ecol.
Manag. 381, 106–114. https://doi.org/10.1016/j.foreco.2016.09.024

Vázquez-Veloso, A., Yang, S.-I., Bullock, B.P., Bravo, F., 2025. One model to rule them all: A
nationwide height–diameter model for 91 Spanish forest species. For. Ecol. Manag. 595,
122981. https://doi.org/10.1016/j.foreco.2025.122981

Weibull, W., 1951. A Statistical Distribution Function of Wide Applicability. J. Appl. Mech.

Zhou, X., Kutchartt, E., Hernández, J., Corvalán, P., Promis, Á., Zwanzig, M., 2023. Determination of
optimal tree height models and calibration designs for Araucaria araucana and Nothofagus
pumilio in mixed stands affected to different levels by anthropogenic disturbance in South-
Central Chile. Ann. For. Sci. 80, 18. https://doi.org/10.1186/s13595-023-01185-9

704