The social, decoupled self: interpersonal synchronization of breathing alters intrapersonal cardiorespiratory coupling

Ivana Konvalinka[1*], Natalie Sebanz[2], Guenther Knoblich[2]

[1] Section for Cognitive Systems, DTU Compute, Technical University of Denmark, Kongens Lyngby, Denmark

[2] Department of Cognitive Science, Central European University, Vienna, Austria

*** Corresponding author:**

Ivana Konvalinka

Section for Cognitive Systems, DTU Compute, Technical University of Denmark

Richard Petersens Plads, B321, room 107

2800 Kongens Lyngby, Denmark

ivako@dtu.dk

Abstract

People synchronize their periodic behavioural and physiological rhythms with each other during social interaction. While this interpersonal synchronization has largely been associated with positive effects such as social bonding, some evidence suggests that it may also impair self-regulation and disrupt intrapersonal coordination. Because respiration and heart rhythms are weakly coupled within individuals, we investigated whether synchronizing breathing with another person alters intra-personal cardiorespiratory coupling. Across two experiments, participants synchronized their breathing reciprocally (bidirectional interaction), unidirectionally with a confederate, or with pre-recorded breathing signals, while respiration and electrocardiography (ECG) were continuously measured. Relative phase analyses revealed that bidirectional breathing synchronization induced in-phase synchronization of heart rhythms between individuals. Critically, interpersonal synchronization coincided with cardiorespiratory decoupling: respiration and heart rhythms became more out-of-phase during interaction compared to resting baselines and the unidirectional condition. Moreover, stronger interpersonal respiratory synchronization predicted greater intra-personal cardiorespiratory decoupling. These findings provide the first evidence that interpersonal physiological synchronization entails a trade-off with intra-personal physiological coupling, perturbing the phase relationship within one's own physiological system. We propose that aligning one's physiological rhythms with others strengthens self–other coupling but weakens intrapersonal coupling, pointing to a physiological mechanism of self-decoupling during social interaction.

Keywords: social interaction, interpersonal physiological synchronization, biofeedback, intrapersonal cardiorespiratory coupling, self-decoupling

1   Introduction:

When people engage in social interaction or simply receive visual or auditory feedback of one another, they tend to synchronize their bodily rhythms with each other both spontaneously and intentionally[1–3]. For example, people's applause becomes synchronized following a theatre or opera performance[4]; they synchronize their walking strides when walking side-by-side[5]; and even their postures become coordinated when engaged in conversation with one another[6]. Such interpersonal synchronization has been proposed to promote self-other integration and merging[7,8]. Beyond overt motor behaviour, more recent research shows that interacting individuals also synchronize their periodic physiological signals, such as heart rhythms and respiration[9–12].

Interpersonal physiological synchronization has been observed across a wide range of contexts[11]: mother-infant dyads show synchronized fluctuations in heart rate and respiration during face-to-face interaction[13,14]; maternal and fetal rhythms show coupling during periods of paced maternal breathing[15]; romantic couples[16] and attracted blind dates[17] exhibit heart rhythm synchronization; and performers and spectators synchronize heart rhythms during collective rituals such as fire-walking, and this interpersonal synchronization is modulated by the relationship between them[9]. Behavioural coupling also modulates interpersonal physiological synchronization. For example, choir singers who sing in unison synchronize their respiration rhythms, and as a result also heart rhythms[18]; and behavioural movement synchronization has also been linked to heart rhythm synchronization[19]. Breathing rhythms also synchronize during conversation, as a result of interpersonal coupling, even though the speakers take turns to speak and listen[20,21]. A recent study also found a causal relationship between respiratory synchronization and behavioural synchronization, which was further shaped by interpersonal differences in resting rhythms, such that partners with more similar rhythms achieved better synchronization[22].

Interpersonal behavioural and physiological synchronization have been proposed to follow the same dynamical principles as synchronization of other biological (e.g., fire-flies) or physical (e.g., pendulums) systems, arising from adjustments of rhythms between interacting self-sustained periodic behaviours through phase and frequency locking[23]. As a result, the underlying dynamics of human interpersonal synchronization have been widely studied using nonlinear models of coupled oscillators[16,24–26].

However, the effects of human interpersonal synchronization extend beyond the mechanics of their coupled dynamics. Much research suggests that interpersonal behavioural and physiological synchronization lead to positive social outcomes, promoting social connection, cohesion, and prosociality[27–31]. Yet a smaller body of work indicates that such effects may not always be beneficial to the self[12]. Empirical studies have linked behavioural interpersonal synchronization to poorer self-regulation of affect[32] and self-decoupling[26,33], but no research has directly examined how interpersonal physiological synchronization affects one's own intrapersonal physiological rhythms.

This question may be particularly interesting in the context of respiration, which is both under voluntary control and strongly linked to cardiac dynamics, given that respiration and heart rhythms are weakly coupled to one another[34,35]. This suggests that synchronization of breathing may influence not only self-other integration but also the coupling of one's own internal physiological rhythms. In other words, adjusting one's physiological rhythms to another person's through *inter*personal physiological synchronization may affect the phase relationship in one's own *intra*personal cardiorespiratory system, modulating the coupling between one's respiration and heart rhythms. Understanding the consequences of interpersonal synchronization on intrapersonal "self" coupling could provide insights into dynamic blurring of self and other that underlies interpersonal coupling.

The underlying mechanisms that modulate interpersonal synchronization are still debated, but have been linked to self and other processes. In the context of movement synchronization, as people move in and out of synchronization, they balance self-other integration and segregation, shifting attentional focus between their own and others' actions[26,36,37]. Maintaining a clear distinction between self and other is considered crucial for joint action[38], as it allows individuals to flexibly switch between self-other integration with segregation. Indeed, the degree to which people adapt to one another varies, with distinct interpersonal strategies emerging during coordination. For example, partners may mutually adapt[39], one may lead while the other follows[40], or both may maintain their own rhythm[41], effectively decoupling from the other by focusing on their own actions. In a study of musicians performing a finger-tapping task, most achieved interpersonal synchronization via mutual adaptation, whereas drummers tended to ignore their partner's timing and maintain their own tempo – yet still synchronized like two independent metronomes[41]. A mathematical model capturing these strategies revealed that mutual adaption and "follower" behaviour were characterized by strong interpersonal coupling of self-other action-perception loops and weakened within-personal coupling, reflecting dynamic self-other integration. Conversely, the non-adaptive drummer or "leader" behaviour was characterized by weak interpersonal coupling but strong intra-personal action-perception coupling, reflecting self-other segregation. This model thus suggests that strong coupling to others, leading to interpersonal synchronization, coincides with decoupling within the self.

Consistent with this prediction, Zimmermann and colleagues[33] found neural evidence of self-decoupling during interactive finger movement coordination. When participants mutually synchronized their movements, EEG functional connectivity was reduced compared to when they improvised alone, suggesting decreased intra-brain synchronization when integrating one's actions with those of another, potentially due to reduced self-monitoring.

However, we add caution to the interpretation of this finding as the neural mechanism underlying task-based reduced intra-brain synchronization remains unclear. Conversely, being observed by the other led to a widespread increase in functional connectivity, particularly over frontal and centro-parietal regions, suggesting neural effects of heightened self-focus.

Yet the functional effects of interpersonal synchronization on the self remain understudied. Theoretical work proposes that social interaction should also allow for moments of independence and disengagement from the other[42], and that excessive social attunement, as during prolonged interpersonal synchronization, may impair self-regulation[32]. Supporting this view, one study found that interpersonal movement synchrony during dyadic improvisation was associated with reduced self-regulation of affect, whereas intra-personal synchrony – measured as the coordination between one's own limbs – opposed these effects[32].

Taken together, these studies suggest that excessively monitoring others' behaviour and bodily rhythms in social settings may result in decoupled intra-personal rhythms, and consequently, poorer self-regulation. Extending these insights from motor behaviour to physiological rhythms raises new questions: does synchronizing one's breathing with another person's pull one away from their natural respiratory rhythms, affecting their own intra-personal cardiorespiratory synchronization? Strong interpersonal respiratory coupling may require adjustments that weaken the stable phase relationship between heartbeats and breathing cycles observed in resting states. Investigating this trade-off between interpersonal and intra-personal coupling offers a novel way to link theories of self-other integration and segregation to physiological mechanisms, and to understand how the social self is dynamically balanced with the physiological self.

In this study, we investigate how synchronized breathing, achieved through real-time visual biofeedback of other people's breathing rhythms, influences one's own cardiorespiratory coupling. Previous analysis of the respiration data from this study showed that synchronization

of breathing rhythms is facilitated not only by mutual interaction, but through emergence of more predictable breathing rhythms during mutual interaction[10]. Here, we analyze the effect of interpersonal breathing synchronization on interpersonal heart rhythm synchronization and intra-personal cardiorespiratory coupling. We contrast conditions of bidirectional versus unidirectional coupling, and compare cardiorespiratory coupling during interaction versus pre-interaction baselines, across two experiments. Given that respiration and heart rhythms are weakly coupled to each other, we expected to find interpersonal heart rhythm synchronization, particularly during bidirectional interaction. Building on previous theoretical work showing that mutual adaptation during interpersonal behavioural synchronization results in the increase in self-other integration but weakening of one's own action perception loops, we expected to see decreased cardiorespiratory coupling during bidirectional interaction compared to both unidirectional interaction and baseline (i.e., breathing at rest) conditions. Furthermore, we expected cardiorespiratory coupling to decrease with increasing interpersonal coupling of breathing rhythms, with the assumption that more self-other integration would yield more self-decoupling.

2   Materials & methods:

2.1   Participants

Twenty-six participants (17 females) took part in Experiment 1 in pairs. One pair was excluded because the participants knew each other. The remaining pairs were strangers to each other. Twenty-four participants, gender-matched to those in Experiment 1, participated in Experiment 2. The sample size for synchronization measures was determined according to studies of interpersonal coordination at the time the study was designed[19,40]. We also ran a post-hoc power analysis based on the effect sizes from the analyses of interpersonal synchronization of breathing rhythms, reported in Konvalinka et al.[10], which estimated sample sizes of 5-26 participants needed in order to achieve 80% power. However, given that the effect of interpersonal synchronization on cardiorespiratory coupling has not been studied before, our sample size was more exploratory for this study.

All participants gave written, informed consent prior to participation, and ethical approval was obtained from the United Ethical Review Committee for Research in Psychology, in Hungary (EPKEB).

2.2   Task and procedure

In experiment 1, the participants were seated in separate rooms, and did not meet each other prior to the experiment. They were each fitted with a BIOPAC wireless MP150 Bionomadix respiration belt and 3-lead ECG electrodes, and all four signals were synchronized through the BIOPAC Acknowledge software. The participants were told they would breathe together with two different people, and their instructions were to synchronize their breathing to the breathing of the other person(s). They interacted in two conditions: bidirectional condition, in which both participants received real-time visual biofeedback of the other person's breathing signal, but not their own; and unidirectional condition, in which the participants both received real-time

biofeedback of the confederate's breathing signal, while a confederate was asked to breathe at rest and did not receive any feedback. The visual signals were presented via the Acknowledge interface. Each breathing interaction lasted 5 minutes, and a 5 minute resting baseline was recorded prior to each interaction. We thus used the baseline conditions to compare cardiorespiratory coupling – the phase relationship between one's breathing and heart rhythm signals – during interaction (bidirectional vs. unidirectional) versus baseline (i.e., while breathing at rest). The order of the interaction conditions was counter-balanced across pairs.

Moreover, we used the data collected as part of experiment 2 in order to see if we could replicate cardiorespiratory coupling effects from experiment 1 - contrasting the interaction condition to the baseline. In experiment 2, the breathing data that emerged in the bidirectional interaction condition in experiment 1 was used as visual feedback, which was originally designed to test the effect of reciprocity versus emergence of predictable breathing patterns on interpersonal synchronization, reported in Konvalinka et al.[10]. Participants were thus recruited individually, and asked to breathe in synchrony to the signals from experiment 1 (i.e., each person interacted with one person's breathing signal), but were told there was another participant in a different room who was producing these signals. The pre-recorded signals were resampled to the frame rate of the monitor and presented using MATLAB's Psychtoolbox. This offline control condition lasted 5 minutes, and a 5 minute resting baseline was recorded prior to the interaction.

2.3   Phase analysis of heart rhythm and cardiorespiratory coupling

The respiration and ECG data were all recorded at a sampling rate of 1000 Hz. The R-peaks were extracted from each person's signal using a peak detection algorithm, and further investigated for artefacts such as missing beats or falsely detected beats using manual detection. Any falsely detected beats were removed, and missing beats were interpolated using linear

interpolation across preceding beats. In order to align the heart rate data to respiration data, the RR-intervals were converted to heart rate, resampled at 1000 Hz and interpolated using piecewise cubic spline interpolation (using interp1 function in MATLAB), and finally detrended. The respiration data were also detrended, and all the signals were normalized by computing z-scores.

The phase was estimated using the Hilbert transform, as in Konvalinka et al. [10]. To quantify heart rhythm synchronization, relative phase was computed between pairs of signals. In the bidirectional interaction and baseline conditions, this was done by subtracting the phase time series from one participant from that of the other interacting participant. For the unidirectional condition, each participant's phase signal was referenced to the confederate's phase signal. In experiment 2, the relative phase was obtained by subtracting each participant's phase signal from the corresponding participant's signal in the bidirectional condition of experiment 1 (i.e., the participant whose signal they synchronized their breathing to).

For measures of cardiorespiratory synchronization, the resampled and interpolated heart rate signals were subtracted from the time-aligned respiration signals for each participant.

Across both heart rhythm and cardiorespiratory measures, relative phase values were converted to a scale spanning -180 to +180 degrees. The distribution of these values was then quantified by computing the percentage of occurrences within $2°$ intervals. In the absence of synchronization, the distribution is expected to be approximately uniform across all bins. By contrast, in-phase synchronization is reflected by a concentration of values around $0°$, anti-phase coordination by peaks near $±180°$, and consistent phase lags by peaks at intermediate angles between $-180°$ and $0°$ (or $0°$ and $180°$).

Following procedures from Konvalinka et al.[10], we statistically compared the bidirectional (12 pairs) condition to the baseline/unidirectional (24 pairs) conditions by dividing the bidirectional recordings into two equal segments for measures of respiratory and

heart rhythm interpersonal synchronization. The first segment was assigned to participant 1 and the second to participant 2, effectively yielding synchronization data across 24 pairs. Relative phase was then calculated separately for each segment, with the division corresponding to the first and second halves of the interaction period (2.5 minutes each). For the analysis of the offline control condition in experiment 2 and cardiorespiratory coupling, such a division was not necessary, given that we had data for 24 pairs and 24 participants, respectively.

From the relative phase distributions, we computed in-phase synchronization to quantify heart rhythm synchronization, defined as % occurrence of relative phase in the -10:10 degree window. In order to quantify cardiorespiratory coupling, we computed the mean relative phase of the relative phase distributions between respiration and cardiac rhythms.

2.4   Statistical analyses

Statistical analyses were carried out using linear mixed-effects models (LMMs). For each LMM, we compared the full model with reduced versions, using a stepwise approach based on Akaike's information criterion (AIC) model selection strategy. The best performing model was then evaluated using the Type III Analysis of Variance with Satterthwaite's method. Pairwise comparisons were conducted using estimated marginal means, with p-values adjusted for multiple comparisons using Tukey's method.

We also measured the relationship between interpersonal respiratory synchronization in the bidirectional (experiment 1) and offline control conditions (experiment 2), and cardiorespiratory coupling. We fit linear regression models with in-phase occurrence between people's respiration rhythms as a predictor variable, and cardiorespiratory mean relative phase as a response variable.

3   Results

3.1   Bidirectional breathing synchronization led to heart rhythm synchronization

The distribution of relative phase was calculated between the two participants' heart rhythms in the bidirectional and unidirectional interaction conditions, i.e., when participants synchronized their breathing to each other or to a confederate's breathing, respectively, as well as in the first resting baseline condition prior to the interactions. The participants' heart rhythms synchronized in-phase only in the bidirectional interaction condition (example data shown in Fig. 1B), while the baseline and the unidirectional condition resembled unform relative phase distributions, as seen in Fig. 1C. This was confirmed using the Rayleigh test for uniformity, which was significant for the bidirectional interaction condition ($z = 6.33$, $p = 0.0017$), but not for the baseline ($z = 0.88$, $p = 0.12$) or the unidirectional condition ($z = 0.52$, $p = 0.66$). A linear mixed-effects model with the condition type as a fixed effect and participants as a random intercept revealed a significant main effect of condition on heart rhythm in-phase synchronization (% occurrence from -10 degrees to +10 degrees), $F(2,46) = 8.14$, $p = .0009$. The bidirectional breathing condition showed a significant increase in in-phase synchronization compared to the baseline ($\beta = 3.89$, SE = 1.09, *t-ratio* = 3.55, $p = .0025$) and the unidirectional breathing condition ($\beta = 3.76$, SE = 1.09, *t-ratio* = 3.44, $p = .0036$). There was no statistical difference between the baseline and the unidirectional condition ($\beta = -0.13$, SE = 1.09, *t-ratio* = -0.12, $p = .99$).

3.2   People's cardiorespiratory rhythms decoupled out-of-phase during interpersonal synchronization of respiration rhythms

The distribution of relative phase was calculated between each participant's respiration and heart rhythms during the bidirectional and unidirectional interaction conditions and the resting baseline conditions that preceded them, as shown in Fig. 2A, capturing cardiorespiratory

coupling. All the conditions exhibited cardiorespiratory synchronization, which were centered at a negative relative phase difference, indicating the heart rate increases and decreases (or RR interval decreases and increases) slightly preceded inhalation and exhalation, respectively. We extracted the mean relative phase between respiratory and cardiac rhythms, as seen in Fig. 2B, in order to capture the degree of decoupling. A linear mixed-effects model with the condition (interaction vs. baseline), directionality (bidirectional vs. unidirectional), and condition x directionality interaction as fixed effects (the best performing LMM), and participants as a random intercept, revealed a significant main effect of condition on the cardiorespiratory mean relative phase $F(1,69) = 25.11$, $p < .0001$, and a significant condition x directionality interaction $F(1,69) = 8.11$, $p = .0058$, but no effect of directionality $F(1,69) = 2.38$, $p = .13$. Pairwise comparisons revealed a significant decrease in the cardiorespiratory mean relative phase during the bidirectional interaction condition compared to the preceding baseline ($\beta = -20.05$, SE = 3.61, $t\text{-}ratio = -5.56$, $p < .0001$), as well as compared to the unidirectional interaction ($\beta = -11.20$, SE = 3.61, $t\text{-}ratio = -3.11$, $p < .014$). There was no difference in the cardiorespiratory mean relative phase between the unidirectional interaction and the preceding baseline ($\beta = -5.52$, SE = 3.61, $t\text{-}ratio = 1.53$, $p < .43$).

To test whether the decoupling could be explained by the variations in the breathing frequency, which showed high variability as reported in Konvalinka et al.[10], we added the breathing frequency as a covariate to the model, both as a fixed effect on its own as well as in interaction with the condition and directionality. However, there was no statistical difference in model comparisons between the original simpler model (*LMM1*: meanrelphase ~ Condition * Directionality + (1|Subject)) and the more complex models with frequency as a fixed effect (*LMM2*: meanrelphase ~ Condition * Directionality + Frequency + (1|Subject), $p = 0.59$; *LMM3*: meanrelphase ~ Condition * Directionality * Frequency + (1|Subject), $p = 0.34$).

Hence, the effect of bidirectional interaction on cardiorespiratory coupling could not be explained by the breathing frequency.

To test whether the interaction had an effect on the stability of the cardiorespiratory phase relationship, we tested the effect of condition, directionality, and condition x directionality interaction on the PLV between the respiration and cardiac rhythms. Model comparisons revealed the simplest model, with only the condition as a fixed effect and participants as a random intercept, to be the best performing model. The condition (interaction vs. baseline) did not have a significant effect on the PLVs, $F(1,71) = 0.05$, $p = .83$.

To investigate whether the cardiorespiratory decoupling effect would be replicated with another group of participants, who synchronized their respiration to the respiration signals that emerged in the bidirectional condition in Experiment 1, we tested the effect of the condition (interaction vs. baseline) on the mean relative phase in Experiment 2. Fig. 3 suggests that people's cardiorespiratory rhythms also decoupled during interaction in this scenario, showing a peak in the distribution of relative phase in the negative direction once again. The relative phase distribution for the baseline condition was centered closer to 0 degrees, suggesting in-phase cardiorespiratory synchronization during the baseline. A linear mixed-effects model with the condition as the fixed effect and participants as random intercepts, showed that synchronizing one's own breathing to another person's (from the bidirectional condition) decreased the mean relative phase in comparison to the resting baseline (Fig. 3B), $F(1,23) = 5.75$, $p = .025$, showing evidence of cardiorespiratory decoupling. This effect could not be explained by the breathing frequency alone, as shown through model comparisons when adding frequency as a covariate or fixed effect in the linear mixed-effects model.

3.3   Higher *inter*personal synchronization of respiration rhythms was associated with higher *intra*personal cardiorespiratory decoupling

In order to test the relationship between interpersonal synchronization of respiration rhythms and intrapersonal cardiorespiratory decoupling, we fit linear regression models between the in-phase synchronization (% occurrence from -10 to +10 degrees relative phase) of the paired participants' respiration rhythms and their cardiorespiratory mean relative phase, in both the bidirectional interaction condition (experiment 1) and offline control condition (experiment 2). As can be seen in Fig. 4A, the more in-phase synchronized the participants were in the bidirectional interaction condition, the more out-of-phase their cardiorespiratory rhythms were ($r^2 = 0.22$, $p = 0.019$). We found the same relationship in the offline control condition as seen in Fig. 4B ($r^2 = 0.43$, $p = 0.0005$), indicating that higher interpersonal synchronization is associated with higher intrapersonal decoupling.

4   Discussion

We investigated how synchronizing breathing with another person modulates interpersonal heart rhythm synchronization and intra-personal cardiorespiratory coupling. Given that breathing is weakly coupled to heart rate[34,35], such that heart rate increases during breathing inhalation and decreases during exhalation, we expected to find interpersonal heart rhythm synchronization when two people synchronized their breathing together. As expected, during bidirectional interaction, when people reciprocally synchronized their breathing together, interpersonal heart rhythm synchronization emerged. Heart rhythm synchronization did not emerge in the unidirectional interaction, between participants and a confederate. Interestingly, interpersonal synchronization of physiological rhythms led to a decoupling of intrapersonal cardiorespiratory rhythms, with a larger out-of-phase relationship between respiration and heart rate during bidirectional interaction compared to breathing at rest (or compared to synchronizing one's breathing to a confederate's breathing in the unidirectional interaction). Moreover, the more people synchronized their breathing with another person, the more decoupled their intrapersonal cardiorespiratory rhythms became. This was shown both in experiment 1 where participants synchronized reciprocally, as well as in experiment 2, when participants synchronized their breathing rhythms unidirectionally to the breathing rhythms that emerged from reciprocal interaction in experiment 1. This indicates that interpersonal physiological synchronization leads to a decoupling of one's own physiological rhythms.

This study provides empirical evidence that synchronizing one's own bodily rhythms with another person, which requires self-other integration, results in a self-decoupling within one's own physiological system. It also indicates that synchronized breathing pulls people away from their own natural physiological rhythms, which may weaken the stable phase relationship within their own cardiorespiratory system, i.e., between their breathing and heart

rhythm cycles. The more one synchronizes with another, the more one is presumably pulled away from their own natural rhythms, which may explain the linear association between the amount of interpersonal synchronization and intrapersonal decoupling. This empirical evidence builds on previous neural evidence of self-decoupling[33], as well as theoretical findings within computational modeling of interpersonal behavioural synchronization. Such models have shown that interpersonal action coordination where one person adapts their behaviour to another's leads to a strengthening in between-personal action-perception loops but a weakening in one's own within-personal action-perception coupling[26,43].

It could be argued that the frequency of breathing plays a role in the cardiorespiratory phase relationship, as shown in previous work[44,45], such that e.g., slower breathing may result in a larger out-of-phase relationship between respiration and heart rate. However, the breathing frequency did not affect the cardiorespiratory phase relationship, or moderate the effect of interpersonal synchronization on cardiorespiratory coupling. Ideally, we could have better disassociated the effect of interpersonal synchronization and the breathing frequency by having a wider range of produced breathing frequencies that people had to synchronize to. However, due to our small sample size and uncontrolled breathing frequencies, this is a limitation of this work.

Interestingly, even the baseline conditions in experiment 1 showed a slight out-of-phase cardiorespiratory relationship, while the baseline condition in experiment 2 did not. This may be explained by carryover effects from the interaction conditions, as the order of bidirectional and unidirectional conditions was counterbalanced across pairs. Half of the baselines preceding each condition were recorded after one of the interaction conditions. For example, if one person started with the unidirectional interaction condition, the baseline preceding the bidirectional condition was recorded after the unidirectional interaction and before the bidirectional one.

Notably, in experiment 2, all the baselines were recorded first, as participants only engaged in one interaction. However, the extent to which self-decoupling carries over after interpersonal synchronization remains to be investigated in future research.

This research also builds on previous work by Galbusera and colleagues[32] showing that interpersonal synchronization impedes self-regulation of affect. Similarly, previous research suggests that weakened self-regulation may modulate interpersonal synchrony. For example, anxious parents have been reported to have higher physiological synchrony with their infants than less anxious parents, due to greater self-reactivity to small-scale fluctuations in their infants' arousal[46]. In other words, when their infants experience increased arousal, instead of downregulating, the anxious parents experience increased arousal as well, which may reflect poorer self-regulation.

The question remains whether the physiological self-decoupling we observe here is indicative of weakened self-regulation; and in addition, what the consequences of such self-decoupling within the physiological system might be. Are we less able to regulate our arousal to different situations when we have synchronized our physiological rhythms with other people's rhythms? Is this mechanism related to self-monitoring, such that decreased self-monitoring, as when integrating oneself with others in coordination tasks, leads to physiological self-decoupling and less self-regulation? Unfortunately, as we did not include any measures of self-regulation in this study, we cannot link the physiological basis of self-decoupling to such measures.

Conclusion

The present study is the first to show intrapersonal physiological effects of interpersonal synchronization, suggesting that when we synchronize our physiological rhythms with others, our own within-personal physiological rhythms become decoupled through an altered phase relationship. This is in line with previous research showing decoupling of one's own action-perception loops during action synchronization, and in line with previous proposals that the mechanism underlying interpersonal synchronization may involve self and other blurring and merging. Future work could investigate how physiological decoupling during interpersonal synchronization impacts self-regulation and self-other blurring, and more directly, what consequences such decoupling has for the self.

Acknowledgements:

This work is supported by the Carlsberg Foundation's Semper Ardens: Accelerate grant, "*Self, other, and we: mechanisms and dynamics of self-other integration in social observation an interaction*", CF22-1251.

Author contributions:

IK: conceptualization, funding acquisition, project administration, formal analysis, data curation, software, visualization, writing – original draft, writing – review & editing. NS: conceptualization, visualization, resources, supervision, writing – review & editing. GK: conceptualization, visualization, resources, supervision, writing – review & editing.

Competing interest statement:

We declare that we have no competing interests.

References

1. Koban L, Ramamoorthy A, Konvalinka I. Why Do We Fall into Sync with Others? Interpersonal Synchronization and the Brain's Optimization Principle. *Soc Neurosci*. 2019;14(1):17470919.2017.1400463. doi:10.1007/s10854-017-7511-4
2. Schmidt RC, Richardson MJ. Dynamics of interpersonal coordination. In: *Coordination: Neural, Behavioral and Social Dynamics*. Berlin, Heidelberg: Springer Berlin Heidelberg; 2008:281-308.
3. Hoehl S, Fairhurst M, Schirmer A. Interactional synchrony: signals, mechanisms and benefits. *Soc Cogn Affect Neurosci*. 2021;16(1-2):5-18. doi:10.1093/scan/nsaa024
4. Néda Z, Ravasz E, Brechet Y, Vicsek T, Barabási AL. The sound of many hands clapping. *Nature*. 2000;403(6772):849-850. doi:10.1038/35002660
5. van Ulzen NR, Lamoth CJC, Daffertshofer A, Semin GR, Beek PJ. Characteristics of instructed and uninstructed interpersonal coordination while walking side-by-side. *Neurosci Lett*. 2008;432(2):88-93. doi:10.1016/j.neulet.2007.11.070
6. Shockley K, Santana MV, Fowler CA. Mutual interpersonal postural constraints are involved in cooperative conversation. *J Exp Psychol Hum Percept Perform*. 2003;29(2):326-332. doi:10.1037/0096-1523.29.2.326
7. Hove MJ, Risen JL. It's All in the Timing: Interpersonal Synchrony Increases Affiliation. *Soc Cogn*. 2009;27(6):949-960. doi:10.1521/soco.2009.27.6.949
8. Sadaphal DP, Blum CR, Keller PE, Fitch WT. Beyond perfect synchrony: shared interpersonal rhythmic timing enhances self-other merging judgements. *R Soc Open Sci*. 2025;12(3). doi:10.1098/rsos.241501
9. Konvalinka I, Xygalatas D, Bulbulia J, et al. Synchronized arousal between performers and related spectators in a fire-walking ritual. *Proc Natl Acad Sci U S A*. 2011;108(20):8514-8519.
10. Konvalinka I, Sebanz N, Knoblich G. The role of reciprocity in dynamic interpersonal coordination of physiological rhythms. *Cognition*. 2023;230:105307. doi:10.1016/j.cognition.2022.105307
11. Palumbo R V., Marraccini ME, Weyandt LL, et al. Interpersonal Autonomic Physiology: A Systematic Review of the Literature. *Personality and Social Psychology Review*. Published online 2016:1-43. doi:10.1177/1088868316628405
12. Mayo O, Gordon I. In and out of synchrony—Behavioral and physiological dynamics of dyadic interpersonal coordination. *Psychophysiology*. 2020;57(6). doi:10.1111/psyp.13574
13. Feldman R, Magori-Cohen R, Galili G, Singer M, Louzoun Y. Mother and infant coordinate heart rhythms through episodes of interaction synchrony. *Infant Behav Dev*. 2011;34(4):569-577. doi:10.1016/j.infbeh.2011.06.008
14. McFarland DH, Fortin AJ, Polka L. Physiological measures of mother–infant interactional synchrony. *Dev Psychobiol*. 2020;62(1):50-61. doi:10.1002/dev.21913
15. Van Leeuwen P, Geue D, Thiel M, et al. Influence of paced maternal breathing on fetal–maternal heart rate coordination. *Proceedings of the National Academy of Sciences*. 2009;106(33):13661-13666. doi:10.1073/pnas.0901049106

16. Helm JL, Sbarra D, Ferrer E. Assessing cross-partner associations in physiological responses via coupled oscillator models. *Emotion*. 2012;12(4):748-762. doi:10.1037/a0025036

17. Prochazkova E, Sjak-Shie E, Behrens F, Lindh D, Kret ME. Physiological synchrony is associated with attraction in a blind date setting. *Nat Hum Behav*. Published online November 1, 2021. doi:10.1038/s41562-021-01197-3

18. Müller V, Lindenberger U. Cardiac and respiratory patterns synchronize between persons during choir singing. *PLoS One*. 2011;6(9). doi:10.1371/journal.pone.0024893

19. Noy L, Levit-Binun N, Golland Y. Being in the zone: physiological markers of togetherness in joint improvisation. *Front Hum Neurosci*. 2015;9:187. doi:10.3389/fnhum.2015.00187

20. McFarland DH. Respiratory Markers of Conversational Interaction. *Journal of Speech Language and Hearing Research*. 2001;44(1):128. doi:10.1044/1092-4388(2001/012)

21. Rochet-Capellan A, Fuchs S. Take a breath and take the turn: how breathing meets turns in spontaneous dialogue. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 2014;369(1658):20130399. doi:10.1098/rstb.2013.0399

22. Yi W, Palmer C. Respiratory synchrony and individual differences causally influence dyadic interpersonal coordination. *bioRxiv*. Published online May 8, 2025. doi:10.1101/2025.05.02.651958

23. Pikovsky A, Rosenblum M, Kurths J. Synchronization: A Universal Concept in Nonlinear Sciences. *Cambridge Nonlinear Science Series 12*. Published online 2003:432. doi:10.1063/1.1554136

24. Haken H, Kelso J a. S, Bunz H. A theoretical model of phase transitions in human hand movements. *Biol Cybern*. 1985;51:347-356. doi:10.1007/BF00336922

25. Schmidt RC, O'Brien B. Evaluating the Dynamics of Unintended Interpersonal Coordination. *Ecological Psychology*. 1997;9(3):189-206. doi:10.1207/s15326969eco0903_2

26. Heggli OA, Cabral J, Konvalinka I, Vuust P, Kringelbach ML. A Kuramoto model of self-other integration across interpersonal synchronization strategies. *PLoS Comput Biol*. 2019;15(10). doi:10.1371/journal.pcbi.1007422

27. Marsh KL, Richardson MJ, Schmidt RC. Social Connection Through Joint Action and Interpersonal Coordination. *Top Cogn Sci*. 2009;1(2):320-339. doi:10.1111/j.1756-8765.2009.01022.x

28. Cirelli LK, Einarson KM, Trainor LJ. Interpersonal synchrony increases prosocial behavior in infants. *Dev Sci*. 2014;17(6):1003-1011. doi:10.1111/desc.12193

29. Wiltermuth SS, Heath C. Synchrony and Cooperation. *Psychol Sci*. 2009;20(1):1-5. doi:10.1111/j.1467-9280.2008.02253.x

30. Launay J, Tarr B, Dunbar RIM. Synchrony as an Adaptive Mechanism for Large-Scale Human Social Bonding. *Ethology*. 2016;122(10):779-789. doi:10.1111/eth.12528

31. Tomashin A, Gordon I, Wallot S. Interpersonal Physiological Synchrony Predicts Group Cohesion. *Front Hum Neurosci*. 2022;16. doi:10.3389/fnhum.2022.903407

32. Galbusera L, Finn MTM, Tschacher W, Kyselo M. Interpersonal synchrony feels good but impedes self-regulation of affect. *Sci Rep*. 2019;9(1):14691. doi:10.1038/s41598-019-50960-0

33. Zimmermann M, Lomoriello AS, Konvalinka I. Intra-individual behavioural and neural signatures of audience effects and interactions in a mirror-game paradigm. . *R Soc Open Sci*. Published online 2022.

34. Schäfer C, Rosenblum MG, Kurths J, Abel HH. Heartbeat synchronized with ventilation. *Nature*. 1998;392(6673):239-240. doi:10.1038/32567

35. Yasuma F, Hayano J ichiro. Respiratory Sinus Arrhythmia. *Chest*. 2004;125(2):683-690. doi:10.1378/chest.125.2.683

36. Keller PE, Novembre G, Loehr J. Musical Ensemble Performance. In: *Shared Representations*. Cambridge University Press; 2016:280-310. doi:10.1017/CBO9781107279353.015

37. Liebermann-Jordanidis H, Novembre G, Koch I, Keller PE. Simultaneous self-other integration and segregation support real-time interpersonal coordination in a musical joint action task. *Acta Psychol (Amst)*. 2021;218:103348. doi:10.1016/j.actpsy.2021.103348

38. Novembre G, Ticini LF, Schutz-Bosbach S, Keller PE. Distinguishing Self and Other in Joint Action. Evidence from a Musical Paradigm. *Cerebral Cortex*. 2012;22(12):2894-2903. doi:10.1093/cercor/bhr364

39. Konvalinka I, Vuust P, Roepstorff A, Frith CD. Follow you, follow me: continuous mutual prediction and adaptation in joint tapping. *Quarterly journal of experimental psychology*. 2010;63:2220-2230. doi:10.1080/17470218.2010.497843

40. Konvalinka I, Bauer M, Stahlhut C, Hansen LK, Roepstorff A, Frith CD. Frontal alpha oscillations distinguish leaders from followers: Multivariate decoding of mutually interacting brains. *Neuroimage*. 2014;94:79-88. doi:10.1016/j.neuroimage.2014.03.003

41. Heggli OA, Konvalinka I, Kringelbach ML, Vuust P. Musical interaction is influenced by underlying predictive models and musical expertise. *Sci Rep*. 2019;9(1). doi:10.1038/s41598-019-47471-3

42. Kyselo M. The body social: an enactive approach to the self. *Front Psychol*. 2014;5. doi:10.3389/fpsyg.2014.00986

43. Heggli OA, Konvalinka I, Kringelbach ML, Vuust P. A metastable attractor model of self–other integration (MEAMSO) in rhythmic synchronization. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 2021;376(1835). doi:10.1098/rstb.2020.0332

44. Lancaster G, Debevec T, Millet GP, et al. Relationship between cardiorespiratory phase coherence during hypoxia and genetic polymorphism in humans. *J Physiol*. 2020;598(10):2001-2019. doi:10.1113/JP278829

45. Lehrer PM, Gevirtz R. Heart rate variability biofeedback: how and why does it work? *Front Psychol*. 2014;5. doi:10.3389/fpsyg.2014.00756

46. Smith CG, Jones EJH, Charman T, Clackson K, Mirza FU, Wass S V. Anxious parents show higher physiological synchrony with their infants. *Psychol Med*. 2022;52(14):3040-3050. doi:10.1017/S0033291720005085

Figure legends

Fig. 1. A. Experimental manipulation. In the bidirectional breathing condition, each participant receives real-time biofeedback of the other participant's respiration signal in real-time. Both their respiration and ECG signals are measured, which are weakly coupled to each other. B. Example respiration signals (top), resampled RR intervals (middle), and the phases of the resampled RR-interval signals (bottom) from the two participants in the bidirectional breathing condition. C. Heart rhythm synchronization: relative phase distribution between the two participants' heart rhythm signals (i.e., resampled RR intervals) during the baseline (black), bidirectional (blue), and unidirectional (red) conditions.

Fig. 2. Cardiorespiratory coupling in Experiment 1, showing effects of interpersonal synchronization of respiration rhythms on intra-personal synchronization of cardiorespiratory rhythms. **A.** Distribution of relative phase (from -180 to 180 degrees) across the conditions types (baseline vs. interaction) and directionality (bidirectional vs. unidirectional). The baselines correspond to the resting baseline recorded prior to each of the interaction conditions.

B. Box plots of the mean relative phase across the four conditions (baseline prior to bidirectional, bidirectional interaction, baseline prior to unidirectional, unidirectional interaction).

Fig. 3. Cardiorespiratory coupling in Experiment 2. **A.** Distribution of relative phase (from -180 to 180 degrees) across the resting baseline and interaction (offline control) conditions. **B.** Box plots of the mean relative phase during baseline and interaction conditions across all participants.

Fig. 4. Relationship between interpersonal synchronization of respiration rhythms (in-phase % occurrence, from -10 to +10 degrees) and intrapersonal cardiorespiratory decoupling (cardiorespiratory mean relative phase) during the **A.** bidirectional interaction condition (experiment 1) and **B.** offline control condition (experiment 2).