Functional heterogeneity of beta bursts in childhood reveals a dimensional neural signature of motor skill

Holly Rayson[1,2,3*], Sarah A. Gerson[4*], Jennifer Keating[5], Catherine R.G. Jones[4], Catherine Purcell[6], James J Bonaiuto[1,2,3**], Ross E Vanderwert[4,7**]

1) Marc Jeannerod Institute of Cognitive Sciences, ISC, CNRS UMR5229, Lyon, France
2) Université Claude Bernard Lyon 1, Université de Lyon, France
3) Hospices Civils de Lyon, Lyon, France
4) School of Psychology, Cardiff University Centre for Human Developmental Science (CUCHDS), Cardiff, United Kingdom
5) School of Social Sciences, Cardiff University, Cardiff, United Kingdom
6) School of Healthcare Sciences, Cardiff University, Cardiff, UK
7) School of Psychology, Cardiff University Brain Research Imaging Centre (CUBRIC), Cardiff, United Kingdom
* authors contributed equally
** authors contributed equally

Correspondence: Holly Rayson <holly.rayson1@gmail.com>

Funding: This work was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (ERC consolidator grant 864550) and British Council Springboard programme (grant 525715).

Abstract
Activity in the beta frequency band appears to play a critical role in sensorimotor processing, with transient 'bursts' exhibiting distinct waveform motifs that may reflect different neural computations. However, how these dynamics relate to motor development and neurodevelopmental conditions remains unclear. Here, we used EEG to investigate sensorimotor beta bursts in children with and without developmental coordination disorder (DCD). We characterized beta bursts during action execution and observation of gross and fine motor actions. Conventional spectral features and overall burst rates did not differ between groups. In contrast, children with DCD showed altered rate modulation of specific burst waveform motifs, including atypical hemispheric lateralization and disrupted interhemispheric connectivity. Critically, motif-specific burst rates were systematically related to motor performance across individuals, with similar relations observed across groups. These findings indicate that waveform-specific beta burst dynamics track graded variation in motor performance, rather than mapping selectively onto diagnostic status. Beta burst variability may therefore reflect the functional expression of motor performance, rather than a disorder-specific neural mechanism underlying DCD. This work highlights the importance of beta burst waveform diversity for understanding variability in sensorimotor function across development.

1. Introduction

Sensorimotor beta activity (13-30 Hz in adults) has long been associated with motor control and is traditionally conceptualized as a sustained oscillatory rhythm that modulates in power prior to, during, and after movement (Pfurtscheller, 1981; Pfurtscheller et al., 1996). However, more recent work indicates that beta activity occurs in transient, burst-like events, rather than as a continuous oscillation (Feingold et al., 2015; Sherman et al., 2016; Little et al., 2019). These *beta bursts* are temporally sparse and behaviorally relevant, predicting response time, motor errors, and stopping performance in adults (Little et al., 2019; Wessel, 2020; West et al., 2023), and are modulated not only during action execution, but also during observation of others' movements (Rayson et al., 2023), implicating them in motor simulation and internal model engagement (Jeannerod, 2001; Press et al., 2011). Developmental work further suggests that beta activity occurs as bursts from infancy, with age-related shifts in peak frequency, timing, duration, and hemispheric lateralization (Gaetz et al., 2010; Cheyne et al., 2014; Rayson et al., 2022, 2023; Rier et al., 2024; Wilkinson et al., 2024). These findings indicate that maturation of sensorimotor function may involve progressive refinement of transient beta events rather than simple changes in sustained oscillatory power. Although the precise computational role of beta bursts remains unresolved, their developmental tuning and behavioral relevance suggest that burst dynamics provide a useful framework for understanding how transient beta activity contributes to sensorimotor processing (Rayson et al., 2025).

Most studies implicitly treat beta bursts as homogenous events, yet recent work demonstrates that individual bursts vary systematically in their time-domain waveform shape. Rather than defining burst types using spectral features (e.g., peak frequency, power, or duration), waveform-based approaches represent each burst as a vector of voltage (or magnetic field strength in the case of magnetoencephalography) across time and apply dimensionality reduction (e.g., PCA) to identify axes of shape variability (Rayson et al., 2023; Szul et al., 2023; Papadopoulos et al., 2024; Agouram et al., 2025). A "motif" in this context refers to the characteristic waveform shape at the extreme ends of a given principal component, that is, bursts with high positive or negative loadings that share reproducible configurations of peaks, troughs, asymmetries, or temporal skew relative to the mean burst waveform. Motifs are therefore not discrete physiological generators, but statistical patterns of shape variation within the burst population. These motifs differ in their rate dynamics and lateralization, and accumulating evidence suggests that they index distinct functional processes (Rayson et al., 2023; Szul et al., 2023; Agouram et al., 2025) such as behavioural error (Moreau et al., 2026).

Developmental findings support the functional relevance of this variability. Infant bursts share a similar canonical waveform with adults but are temporally longer; after

accounting for duration differences, their waveforms align closely in shape (Rayson et al.,
2023). Moreover, variability around the mean waveform is structured and conserved
across age, while motif-specific rate modulation becomes increasingly precise from
infancy to adulthood (Rayson et al., 2023), consistent with progressive specialization of
sensorimotor computations. Examining waveform-defined motifs across children
spanning a range of motor abilities, including those with developmental coordination
disorder (Wilson et al., 2017) therefore provides a principled framework for testing
whether specific dimensions of burst shape variability track individual differences in motor
skill and interhemispheric organization.

Recent evidence indicates that beta bursts with distinct waveform shapes exhibit
dissociable patterns of rate modulation and hemispheric lateralization during movement,
with some motifs showing increasingly precise lateralization across early development
(Rayson et al., 2023). These findings suggest that waveform-defined burst types may
reflect distinct computational roles within distributed sensorimotor networks, potentially
including interhemispheric coordination. Motor ability varies substantially across
development, and individual differences in coordination are associated with variability in
hemispheric specialization and functional connectivity within sensorimotor systems
(Querne et al., 2008; Tallet et al., 2013; Brown-Lum and Zwicker, 2015; McLeod et al.,
2016). Studying children with a wide range of motor proficiency, including those meeting
criteria for developmental coordination disorder (Wilson et al., 2017), therefore provides
a principled way to test whether specific waveform-defined beta burst motifs scale with
motor performance and interhemispheric dynamics. While DCD is defined categorically
at the clinical level, variation in motor performance and its neural substrates may
nonetheless be continuous across individuals. Under this framework, DCD therefore
serves as a model of pronounced motor difficulty within a broader continuum, enabling us
to examine how variation in burst motif dynamics relates to motor coordination.

Here, we characterized sensorimotor beta burst dynamics in children spanning a
broad range of motor abilities during action execution and observation, including children
with and without DCD. We explicitly tested for differences between groups in the
modulation, lateralization, and interhemispheric connectivity of waveform-defined burst
motifs, and additionally examined whether motif-specific burst rates were related to
individual differences in motor performance. We show that specific burst motifs display
altered rate modulation and hemispheric asymmetry during movement in children with
DCD, and that motif-specific burst rates are systematically related to motor performance
across individuals. Notably, although these effects differentiate groups, the relationship
between burst dynamics and motor performance is continuous across the full sample.
These findings suggest that variability in waveform-specific beta burst dynamics reflects
graded differences in the organization of interhemispheric sensorimotor communication,
along a continuum of motor performance.

2. Methods

2.1 Participants

Twenty-three 8- to 12-year-old children with DCD (M = 10.88, SD = 1.56; 18 male) were recruited via social media and the Dyspraxia Foundation, and 33 typically developing children (M = 9.85, SD = 1.30; 17 male) were recruited via social media as a comparison group. Participants in the DCD group were required to have received a clinical diagnosis of DCD, confirmed for this study via parent report and converted total score on the Movement Assessment Battery for Children 2nd Edition (MABC-2; Henderson et al., 2007) at or below the 16th percentile at the time of the testing session. In line with the Leeds Consensus Statement (2006), participants with a diagnosis of DCD who had co-occurring neurodevelopmental conditions were not excluded from participating, but additional diagnoses were noted. In the final sample, co-occurring conditions were autism (n = 3), ADHD (n = 1), dyslexia (n = 1), and autism, dyslexia, developmental language disorder and learning difficulty (n = 1). Children were not recruited to the comparison group if they had a diagnosis of a neurodevelopmental condition or developmental delay.

Nine children from the comparison group were excluded from the analyses of group differences as they scored below the 25th percentile on the MABC-2. This was to ensure we were not including children in the comparison group who did not have normative motor development (French et al., 2018). A further six participants were excluded during EEG processing due to insufficient data (DCD n = 4; comparison n = 2). The final sample therefore included 19 children with DCD (M_{age} = 11.17 years, SD = 1.28; 14 male; 14 right-handed) and 22 children without DCD (M_{age} = 9.78 years, SD = 1.40; 12 male; 21 right-handed). Full testing and demographic details are available in Keating et al., (2023); however we note here that the comparison group had significantly higher IQ scores (p = .011) (assessed by the Weschler Abbreviated Scales of Intelligence, 2nd edition [WASI-II]; Weschler, 2011) and were significantly younger than the DCD group (p = .002). Written informed consent was obtained from the participant's caregiver before the start of the experiment. Each child also provided written assent to take part. Ethical approval was granted by the Ethics Committee at BLINDED FOR REVIEW. Participants received travel expenses and a small prize.

2.2 Movement Assessment Battery for Children

The Movement Assessment Battery for Children 2nd Edition (MABC-2; Henderson et al., 2007) test component was used to assess motor skills. The MABC-2 has good internal consistency and test re-test reliability (Henderson et al., 2007; Wuang et al., 2012). The assessment consists of eight tasks organized into three subsets: manual dexterity, aiming and catching, and static and dynamic balance. Age-adjusted standard scores were calculated for each subtest, and a total difficulty score was created from which percentiles can be derived. A total score below the 5th percentile indicates

significant motor difficulties, a score between the 5th and 16th percentile indicates a risk
of motor difficulties, and a score above the 25th percentile indicates age-expected motor
performance.

2.3 Task and Stimuli

Video stimuli for the EEG motor tasks were filmed using a handheld camera fixed
to a tripod. The camera was positioned overlooking the model's shoulder while she
performed the actions to best match the participant's perspective of the actions as if they
were performing them, while also providing a clear view of the entire action. We chose an
ego-centric perspective because previous studies have suggested this elicits greater
sensorimotor activity (Jackson et al., 2006; Fu and Franz, 2014; Drew et al., 2015).

All videos were trimmed to fit within a 10-second period, with three types of videos
in total (gross motor, fine motor, and non-biological movement). The gross motor video
displayed a model sitting with the hand starting at rest on the edge of a table, extending
the arm to reach out and tap a red square, and then returning the arm to rest. This action
took approximately 2 seconds and was repeated 5 times. The fine motor video displayed
a hand holding a red pen and tracing a flower figure (based on the Movement Assessment
Battery for Children; Henderson et al., 1992). The tracing lasted approximately 20
seconds and was split into two separate videos; therefore, half of the fine motor action
observation trials began with an empty flower and showed the hand tracing halfway
around and the other half began with the hand already halfway around the flower and
continued tracing to the end of the flower. The non-biological movement videos were of
kaleidoscope patterns taken from Pixabay (www.pixabay.com). All videos of actions were
performed by a right-handed person and were mirrored to make the actions appear to be
performed by a left-handed model for left-handed participants (n = 6). All videos were
presented on the center of the screen, subtending a visual angle of ~28° (horizontal) and
~14° (vertical). Example stimuli are displayed in Figure 1 and are available at
https://osf.io/vnup9/.

The EEG task consisted of five conditions adapted from Hobson and Bishop (2016;
Figure 1). Two conditions were execution trials: Execution Gross motor (ExG) and
Execution Fine motor (ExF). These began with an instruction screen describing the action
with an image of the action stimulus (i.e. a red square for ExG/the flower for ExF). Once
the participant was ready, the experimenter initiated the trial with a button press. During
the trial, the command 'GO' was displayed beneath the action stimulus for 10 seconds
while participants performed the action. A 'STOP' command was then displayed for 2
seconds. Three conditions consisted of observation trials using the videos described
above. These began with a fixation cross. Once the participant was ready, the
experimenter initiated the trial with a button press. During the trial, either the Observation
Gross motor (ObG), Observation Fine motor (ObF), or the non-biological movement
(kaleidoscope; Kal) video played for 10 seconds. Participants completed 40 trials of each

condition split across two blocks. Trials were presented in a random order and timing was
controlled by E-Prime 3.0 software (Psychology Software Tools).

158
Figure 1. Example Trials in the EEG Task. Observe trials began with a fixation cross to cue participants
to remain still and observe the videos for 10 seconds. Example screenshots of the videos are shown for
Kaleidoscope (a); the hand reaching for the red square for Gross Motor (b); and a hand tracing a flower for
Fine Motor (c) trials. Execution trials began with action instructions to prepare participants; specifically to
"Tap the square!" for Gross Motor (d) or "Trace the flower!" for Fine Motor (e) trials. The "Go" screen was
then displayed for 10 seconds to initiate participants execution and then the "Stop" screen was displayed
for 2 seconds to end the trial.

2.4 Procedure

Each child was prepared for the EEG acquisition, including placement of the EEG
cap and preparation of electrodes with electrogel, and then brought into the testing room.
Their caregiver was invited to observe the testing session from an adjoining room while
they completed questionnaires on a tablet computer (see Keating et al., 2023 for details).
During placement of the EEG cap, the child completed the Edinburgh Handedness
Inventory - Short Form (Veale, 2014) to determine whether they were dominantly left- or
right-handed. Once the EEG was fully set-up, the task was explained to the participant.

Participants sat at a table with a red square affixed to the surface across from a
17" computer screen approximately 80 cm away. The session began by measuring
resting-state EEG as the child sat with their eyes first open for 30 seconds, and then their
eyes closed for 30 seconds. This procedure was repeated three times.

Following resting measures, children practiced the tasks via completion of two
observation trials (one ObG and one ObF) and two execution trials (one ExG and one
ExF). Participants then completed the motor task, with the entire session lasting around

25 minutes. Once the EEG session was complete, the EEG cap was removed, and the
participants completed the MABC-2 and WASI-II (in this fixed order). Children could take
breaks during testing as needed, with the whole session lasting approximately 100
minutes.

2.5 EEG Acquisition

Continuous EEG was recorded with the BrainVision actiCHamp Plus system (Brain
Products GmbH, Gilching, Germany) from 32 channels placed according to the
International 10-20 System, and referenced online to the vertex electrode (Cz). ECI
Electro-Gel (GVB geliMED GmbH, Germany) was used to reduce impedances, which
were kept below 20 kΩ. Data were sampled at 500 Hz. Video recordings of all EEG
sessions were coded to identify trials in which participants were not watching a video or
completing the required action. Any trials where participants were visibly not attending or
were making off-task movements were removed prior to analysis (Observation: *Mdn* = 4,
range 0-13 trials; Execution: *Mdn* = 2, range 0-8 trials).

2.6 Data preprocessing

EEG data were preprocessed in MATLAB (R2018a) using a custom version of the
MADE pipeline (Debnath et al., 2020), augmented with artifact-detection routines from
the NEAR pipeline (Kumaravel et al., 2022). Data were high-pass filtered at 1 Hz and low-
pass filtered at 100 Hz using EEGLAB (v2021.1; Delorme and Makeig, 2004) Hamming-
window FIR filters (high-pass cutoff at 0.5 Hz; low-pass cutoff at 105 Hz, reflecting the
transition-band specification in the filter design). Prior to cleaning, discontinuous
segments outside the task were removed by trimming the continuous recording to 1.5 s
before the first event and 1.5 s after the last event.

Bad channels were identified using NEAR's local outlier factor (LOF) procedure,
with an initial LOF cutoff of 2.5 and adaptive thresholding enabled (the cutoff was
incremented when >10% of channels were flagged); flat-line detection was also enabled
(5 s tolerance), while the periodogram-based detector was disabled (comparison: 0 - 3
channels, *M* = 0.24, *SD* = 0.66; DCD: 0 - 12 channels, *M* = 0.70, *SD* = 2.51). Artifact
Subspace Reconstruction (ASR) was then applied to correct non-stereotyped artifacts
(correction mode, burst criterion = 13). Stereotyped artifacts were subsequently removed
using ICA, computed on a copy of the dataset segmented into 1 s epochs; epochs
exceeding ±1000 µV were rejected prior to ICA, and channels with >20% rejected 1 s
epochs were removed from the ICA copy and correspondingly excluded from the original
data before transferring ICA weights. Artifactual independent components were identified
using the Adjusted-ADJUST plugin with default parameters (Mognon et al., 2011; Leach
et al., 2020) and removed from the original continuous dataset (comparison: 1 - 19
components, *M* = 8.03, *SD* = 4.66; DCD: 1 - 14 components, *M* = 7.74, *SD* = 2.83).

Following ICA-based correction, data were epoched into task segments time-
locked to condition markers (Kal, ObF, ObG, ExF, ExG) using a -8 to 15 s window. Artifact
rejection was performed at the epoch level using a ±150 µV threshold; epochs were
rejected if ≥10% of channels exceeded threshold (comparison: 1 - 99 epochs, M = 34.24,
SD = 29.12; DCD: 9 - 99 epochs, M = 31.96, SD = 27.79), otherwise the affected channels
were interpolated within that epoch. After artifact rejection, remaining missing channels
were interpolated (spherical interpolation) and data were average re-referenced. Finally,
50 Hz line noise was removed using an iterative Zapline procedure (de Cheveigné, 2020)
implemented in the MEEGKit package (https://nbara.github.io/python-meegkit/), using a
window size of 10 Hz for polynomial fitting and 2.5 Hz for noise peak removal and
interpolation. Participants with fewer than 5 trials for a condition were excluded from
analysis of that condition. All source code for preprocessing is available at
https://github.com/danclab/dcd_bursts/tree/main/preprocessing.

2.7 Data-Driven Definition of the Beta Band

All analyses focused on the C3 and C4 electrodes, overlying left and right
sensorimotor cortex, which are the canonical loci of movement-related beta modulation
(Pfurtscheller, 1981; Pfurtscheller et al., 1996). To define the frequency limits of the beta
band in a data-driven manner, power spectral densities (PSDs) were computed for each
participant from 0.5 to 40 Hz using Welch's method (1 s Hann windows, 50% overlap, 10×
zero-padding). PSDs were parameterized using specparam (Donoghue et al., 2020) to
separate periodic (oscillatory) and aperiodic components.

Periodic components were averaged across C3 and C4 electrodes and then
across participants from both groups. Spectral peaks were identified using an iterative
peak-fitting procedure in which the frequency-domain periodic spectrum was fit with a
Gaussian at its global maximum, subtracted, and re-evaluated until no additional peaks
remained. Peak frequency bands were defined using the full width at half maximum
(FWHM) of each Gaussian, retaining only peaks with a minimum FWHM of 1 Hz below
10 Hz or 3 Hz above 10 Hz. This procedure yielded a beta range of 14-31.25 Hz, which
was used in all subsequent analyses.

2.8 Conventional Beta Power Analysis

To characterize canonical beta modulation, time-frequency (TF) representations
of each trial were computed using an adaptive superlet transform (Moca et al., 2021), with
varying central frequencies (1-50 Hz) and a fixed number of cycles (4 cycles) within a
Gaussian envelope. The wavelet order (i.e., the multiplier for the number of cycles in the
wavelet) varied linearly from 1 to 40 over the frequency range. Time-frequency
representations were averaged across the defined beta band and baseline-corrected
relative to -7 to -1 s using percent signal change. Group-level time courses were
compared using cluster-based permutation testing (Maris & Oostenveld, 2007), with

between-group comparisons performed via independent-samples cluster tests and
within-group modulation assessed using one-sample cluster tests against zero.

To complement time-resolved analyses, beta power was averaged within
predefined temporal windows (baseline: -7 to -1 s; task: 1-9 s; post-task: 11-14 s), and
analyzed using linear mixed-effects models to assess differences across groups and task
windows within each hemisphere and execution condition. For each hemisphere
(contralateral, ipsilateral) and execution condition (ExF, ExG), window-averaged beta
power was modeled as a function of group (DCD, comparison), window, and their
interaction, with participant included as a random intercept. Models were fit separately for
each hemisphere and condition. Statistical significance of fixed effects was evaluated
using Type III Wald χ^2 tests (car v3.1.0; Fox and Weisberg, 2018). Where appropriate,
pairwise post hoc comparisons were performed using Tukey-corrected estimated
marginal means (lsmeans v2.30; Lenth et al., 2020). All models were fit in R (v4.2.2; R
Core Team, 2022) using the lme4 package (v1.1.30; Bates et al., 2014).

2.9 Lagged Hilbert Autocoherence

To assess whether beta activity reflected sustained rhythmic oscillations or
transient burst-like dynamics, we computed lagged Hilbert autocoherence (LHaC) over
the 1-40 Hz range during the 0-10 s execution interval. LHaC quantifies the degree to
which oscillatory amplitude fluctuations remain temporally coherent across lags; lower
values in the beta band relative to neighboring frequencies indicate reduced sustained
rhythmicity and greater burst-like structure (Zhang et al., 2025). For each participant and
hemisphere, LHaC was computed across temporal lags (0.1-4.5 cycles). Group-level
LHaC spectra were obtained by averaging across participants within hemisphere.

2.10 Beta Burst Detection Algorithm

The lagged Hilbert autocoherence analysis indicated that beta activity over
sensorimotor cortex was weakly rhythmic and therefore better characterized as transient
burst events rather than sustained oscillations. We therefore analyzed beta activity at the
level of individual burst events using a recently developed adaptive burst detection
algorithm (Rayson et al., 2023; Szul et al., 2023) designed to capture transient oscillatory
events over a wide amplitude range without relying on a fixed global threshold (Figure 2).
For each trial, time-frequency (TF) representations were computed using the superlet
transform (Moca et al., 2021), which optimally balances time and frequency resolution,
and is thus particularly suited for detecting transient bursts within narrow frequency
bands. As for conventional power analyses, an adaptive superlet transform was used,
with varying central frequencies (1-50 Hz) and a fixed number of cycles (4 cycles) within
a Gaussian envelope. The wavelet order (i.e., the multiplier for the number of cycles in
the wavelet) varied linearly from 1 to 40 over the frequency range.

Figure 2. Iterative beta burst detection and waveform extraction pipeline. Raw EEG time series are transformed into the time–frequency domain using the superlet transform. Power spectral densities are computed and parameterized to separate periodic and aperiodic components, allowing subtraction of the aperiodic fit from the TF representation. Burst detection proceeds iteratively by identifying the global TF maximum, fitting a two-dimensional Gaussian, subtracting it from the residual TF matrix, and repeating until no peaks exceed a noise threshold (2 SD). For each detected burst, a 260 ms time-domain waveform is extracted, phase-aligned using the Hilbert transform, checked for temporal consistency with the TF peak, DC-corrected, and retained for further analysis.

To isolate oscillatory burst activity, PSDs were derived from each trial's TF decomposition and parameterized using specparam to estimate the aperiodic (1/f-like) component of the spectrum. This step is critical because aperiodic activity contributes broadband power that can vary across frequency and time, and if left unaccounted for, can bias burst detection by inflating apparent oscillatory peaks or distorting their frequency and duration estimates. The aperiodic fit was therefore subtracted from the TF representation, yielding a residual TF matrix that more selectivity reflects band-limited power (Brady and Bardouille, 2022). Burst detection was then performed directly in this residual TF space. At each iteration, the algorithm identified the global maximum in the residual TF matrix, defined jointly over the time and frequency dimensions. A two-dimensional Gaussian was then fit to this peak, with one dimension capturing the burst's temporal extent (duration) and the other capturing its spectral extent (frequency span).

The fitted Gaussian provides estimates of peak time, duration, peak frequency, frequency span, and peak amplitude, and was subsequently subtracted from the residual TF matrix. The algorithm then proceeded to the next iteration on the updated residual. Iteration continued until no remaining TF peak exceeded a noise floor defined as 2 standard deviations above the mean residual power across all time and frequency bins, recomputed in every iteration. To minimize edge effects, burst detection was applied to TF data spanning 11-34.25 Hz, and only bursts with peak frequencies within the predefined beta band (14-31.25 Hz) were retained for further analysis.

For each detected burst, the corresponding time-domain waveform was extracted
from the minimally filtered EEG signal (1 Hz high-pass, 100 Hz low-pass) using a 260 ms
window centered on the TF peak time (Szul et al., 2023). To ensure consistent alignment
of waveform features across bursts, waveforms were phase-aligned by band-pass
filtering each burst within its detected frequency span, computing the instantaneous
phase using the Hilbert transform, and re-centering the unfiltered waveform on the phase
minimum closest to the TF peak (Boto et al., 2022). Bursts were excluded if the phase-
aligned peak deviated by more than 30 ms from the TF-detected peak time. Finally, the
DC offset was removed to yield the accepted burst waveform used in subsequent
analyses. All code for the burst detection algorithm is available at
https://github.com/danclab/burst_detection.

2.8 Burst analysis

We first compared beta burst features (duration, peak amplitude, peak frequency,
and frequency span) between the comparison and DCD groups to ensure that any
subsequent group differences were not driven by conventional time-frequency
characteristics. Statistical significance was assessed using nonparametric permutation
tests (N = 100,000), in which group labels were randomly shuffled and the observed
difference in mean feature values was evaluated against the resulting null distribution.

Overall burst rate was then computed separately for contralateral and ipsilateral
sensorimotor electrodes during fine (ExF) and gross (ExG) motor execution. For each
participant and condition, burst occurrences were binned in 0.2 s intervals from -8 to 15
s, converted to rates (Hz), and normalized by the number of trials. The resulting time
series were smoothed with a Gaussian kernel (σ = 2 bins) and baseline-corrected by
subtracting the mean rate during the pre-task interval (-7 to -1 s). Group differences in
burst-rate dynamics were evaluated using cluster-based permutation testing (1,000
permutations) on the baseline-corrected time series, controlling for multiple comparisons
across time.

We then used Principal Component Analysis (PCA) to characterize variability in
burst waveform shape and to determine whether this variability differed between the
comparison and DCD groups. Unlike conventional time-frequency features (e.g., peak
frequency or amplitude), which are derived from spectral representations, waveform
analysis operates directly on the raw time-domain signal of each detected burst. PCA
identifies orthogonal axes that capture the dominant dimensions of waveform variability
across bursts. These components represent statistical directions of variance and are not
assumed to correspond to discrete physiological generators. Throughout the manuscript,
we use the term "motif" as a descriptive shorthand for the waveform shapes associated
with extreme loadings on a given component.

Burst waveforms were extracted during the detection procedure and pooled across
conditions for PCA. We specified 20 components to ensure that the decomposition
captured the full range of observable waveform variability. This dimensionality was
chosen conservatively: across multiple datasets analyzed with this framework,
permutation-based significance testing has consistently identified no more than
approximately 10-12 components as exceeding chance levels of explained variance
(Rayson et al., 2023; Szul et al., 2023; Papadopoulos et al., 2024; Agouram et al., 2025).
Specifying 20 components therefore provides a stable upper bound that exceeds the
number of meaningful components without constraining the solution a priori.

We first conducted separate PCAs for each group (after robust scaling), treating
each time point as a feature and defining the mean burst waveform as the origin of a 20-
dimensional space. Each burst was represented by its scores along each principal
component, indicating its deviation from the mean waveform along that dimension. To
assess whether the structure of waveform variability differed between groups, we
compared eigenvectors using Spearman correlation and observed high correspondence
across groups. A single global PCA was therefore fit to all bursts pooled across
participants, enabling direct comparison within a shared representational space.

To determine which components captured meaningful waveform variance, we
compared the variance explained by each component in the real data with null
distributions obtained from 100 iterations of temporally shuffled data, in which time points
within each waveform were permuted to destroy temporal structure while preserving
marginal distributions (Vieira, 2012). P-values were defined as the proportion of shuffled
solutions explaining greater variance than the empirical component. After Bonferroni
correction ($\alpha = 0.0025$), PCs 1-11 were retained for further analysis.

To identify waveform motifs that were modulated during movement, we compared
principal component (PC) scores between baseline (-7 to -1 s) and task (1 to 9 s) windows
within each execution condition (ExF, ExG). For each subject, bursts were assigned to
100 ms time bins, and mean PC scores were computed per bin, normalized by trial count,
and smoothed (Gaussian kernel, $\sigma = 2.5$ bins). This yielded a time-resolved estimate of
the average contribution of each PC to the burst population over the course of the trial.
For each PC, we then computed the difference between baseline and task window means
separately for each subject and group. Statistical significance was assessed using paired
permutation tests (sign-flipping across subjects; 1,000 permutations). PCs showing a
significant baseline-task difference ($p < 0.05$) were considered task-modulated. This
analysis identified PCs 3, 6, and 8 as significantly modulated during execution. Because
changes in mean PC score may reflect altered prevalence of bursts at different positions
along that dimension rather than purely within-burst shape changes, we next quantified

waveform-specific burst rates by stratifying bursts according to their PC score and
examining condition-dependent changes in their occurrence.

Importantly, because principal components define continuous dimensions of
waveform variation, group differences are not expected to manifest uniformly across all
portions of a component's score distribution; instead, they may appear selectively at one
extreme of that dimension, reflecting a redistribution in the prevalence of bursts with
particularly high or low loadings rather than a global shift in mean waveform shape.
Therefore, for each PC that exhibited task-related modulation, we quantified waveform-
specific burst rate by stratifying bursts into tertiles of PC score (based on the global
percentiles of that PC across all bursts). Burst rates were then estimated separately for
contralateral and ipsilateral sensorimotor electrodes and for each execution condition
(ExF, ExG). Within each subject, bursts were binned in 200 ms intervals over the full trial
epoch (-8 to 15 s), converted to rates (Hz) by dividing by bin width, and smoothed with a
Gaussian kernel (σ = 2 bins) at the single-trial level before averaging across trials. For
each tertile, the resulting burst-rate time course was baseline-corrected by subtracting
the subject-specific mean rate in the baseline window (-7 to -1 s), yielding a baseline-
referenced rate time series (ΔHz) for each subject, hemisphere, and condition. Between-
group differences in these baseline-corrected burst-rate time series were assessed using
cluster-based permutation testing (two-sample; 1,000 permutations; two-tailed t-threshold
based on the pooled degrees of freedom). Cluster significance was evaluated over the
analysis interval (-7 to 14 s), and clusters shorter than 20 samples were discarded to
reduce spurious detections.

Waveform-specific burst rates were further analyzed using linear mixed-effects
models applied to window-averaged, baseline-referenced burst rates as a targeted follow-
up to the time-resolved permutation tests, allowing us to quantify whether group
differences reflected a consistent shift in burst rate within specific waveform-defined burst
motifs. For each burst tertile, hemisphere, and execution condition (ExF, ExG), we
computed a subject-level summary measure defined as the difference between mean
burst rate during the task window (1-9 s) and the baseline window (-7 to -1 s). These
values were then entered into mixed-effects models to assess between-group
differences. Models were fit separately for each PC (PC6, PC8), tertile, hemisphere
(contralateral, ipsilateral), and execution condition. In each model, burst rate was
modeled as a function of group (DCD, comparison), with participant included as a random
intercept to account for repeated measures. Statistical significance of fixed effects was
assessed using Type III Wald χ^2 tests (car v3.1.0; Fox and Weisberg, 2018). Model fitting
was performed in R (v4.2.2; R Core Team, 2022) using the lme4 package (v1.1.30; Bates
et al., 2014).

2.9 Relationship between waveform-specific burst rates and motor skill (MABC-2)

To test whether execution-related, waveform-specific beta burst dynamics were
associated with motor skill, we related baseline-referenced burst rates to MABC-2 total
scores. Analyses were performed on ipsilateral sensorimotor electrode burst-rate time
courses from the execution condition and PC identified in the group-difference analyses
(PC6 during ExG; PC8 during ExF). For each PC, bursts were stratified into tertiles based
on the global percentiles of PC score across all bursts. Within each participant and tertile,
burst rates were computed by binning burst peak times in 200 ms intervals, converting
counts to rates (Hz), smoothing at the single-trial level (Gaussian kernel; $\sigma = 2$ bins),
averaging across trials, and baseline-referencing by subtracting the mean rate in the
baseline window (-7 to -1 s). A single summary value per participant was then obtained
as the mean baseline-referenced burst rate during the early execution period (1 to 6 s).

Associations between burst-rate summaries and MABC-2 scores were assessed
using Spearman rank correlations pooling participants across groups, and including
participants from the comparison group with MABC-2 scores below the 25th percentile.
To visualize and complement the rank-based effects, we fit separate group-wise linear
models and tested the slope using OLS with HC3 heteroskedasticity-consistent standard
errors. For slopes with $p < 0.05$, we plotted regression lines with 95% bootstrap
confidence intervals (10,000 iterations; resampling participants with replacement).

2.10 Granger Causality Analysis
To test whether group differences in waveform-specific burst dynamics were
accompanied by altered directed inter-hemispheric interactions, we estimated Granger
causality (GC) from motif-specific burst-count time series rather than from the continuous
beta envelope. This choice was motivated by the aim of quantifying directed coupling
specifically between discrete burst events stratified by waveform motif (PC-score bins),
whereas beta-envelope signals pool across motifs and can conflate changes in burst
probability with changes in within-burst amplitude.

GC was computed only for the PC-condition pairs that showed robust motif-specific
group effects in the burst-rate time series (PC6 during ExG; PC8 during ExF). For each
analysis (PC6-ExG; PC8-ExF), bursts were first stratified into PC tertiles (T1-T3) using
global PC-score percentiles. Within each subject, bursts were then converted into trial-
wise point-process count series by binning burst peak times into 10 ms bins over the
movement period (0-10 s post-onset), separately for ipsilateral and contralateral
sensorimotor electrodes. This yielded six units per subject (ipsi T1-T3; contra T1-T3),
each represented as an array of shape *units × trials × time bins*.

Directed connectivity was estimated using a point-process GC framework
implemented as a negative-binomial GLM with constant dispersion ($\alpha = 1.0$), following the
logic of point-process Granger methods (Kim et al., 2011; Agouram et al., 2025). For each

source→target unit pair, we fit (i) a full model predicting the target's binned counts from
an intercept, the lagged activity of the source, and the lagged activity of all other units
(fully conditional GC), and (ii) a reduced model identical to the full model but omitting the
source term. GC strength was defined as the likelihood-ratio statistic, GC = 2(LL_{full} -
$LL_{reduced}$). To aid interpretation, GC values were signed by the sign of the source
coefficient in the full model (positive = excitatory influence; negative = inhibitory influence,
in the point-process sense). Because the burst detector prevents temporally overlapping
bursts within a channel, within-hemisphere (and within-unit) dependencies are trivially
constrained; therefore, within-hemisphere connections were excluded a priori and
inference focused on inter-hemispheric connections.

History lags were selected per source-target pair by cross-validation over trials,
choosing the lag that maximized held-out predictive performance. Candidate lags were
1-9 bins, corresponding to 10-90 ms given the 10 ms bin size, and model selection used
3-fold cross-validation. Statistical significance of within-group GC was assessed non-
parametrically using 1,000 permutations. For each permutation and subject, trial labels
were shuffled to disrupt across-unit temporal alignment for off-diagonal (inter-unit)
interactions, while within-trial time points were shuffled to generate null values for
diagonal terms; null GC matrices were then averaged across subjects to form a group-
level null distribution for each connection. Two-sided p-values were computed as the
proportion of permuted GC values with absolute magnitude exceeding the observed
absolute GC value, and multiple comparisons across connections were controlled using
Benjamini–Hochberg FDR (α = 0.05) within each PC-condition analysis. Between-group
differences were tested by subject-label permutation (10,000 permutations), recomputing
the group-mean signed-GC difference (DCD - comparison) for each connection and
deriving two-sided p-values from the resulting null distribution; significant group-
difference connections were retained for visualization and interpretation after the same
within-analysis multiple-comparisons control.

Source code for all of these analyses is available at
https://github.com/danclab/dcd_bursts.

3. Results
3.1 Time-frequency-based burst features do not differ between DCD and
comparison groups
We first examined conventional beta power dynamics over sensorimotor cortex
(C3/C4), baseline-corrected to the pre-task interval. Cluster-based permutation testing of
the continuous time series revealed task-related beta modulation only during motor
execution (Figure 3, see Figure S1 for observation conditions). During fine motor
execution (ExF), significant contralateral beta modulation was observed in the
comparison group, whereas no reliable clusters were detected in the DCD group. During

gross motor execution (ExG), significant beta modulation was present ipsilaterally in both groups. No significant between-group differences emerged in any condition from the time-resolved analyses. Importantly, task epochs lasted 10 s and were not time-locked to individual movement events, limiting the temporal specificity of conventional power estimates and motivating a complementary window-based analysis.

We therefore averaged beta power within three predefined windows (baseline, task, and post-task) and analysed trial-wise values using linear mixed-effects models. During contralateral ExF, there was a significant main effect of window ($\chi^2(2) = 20.77$, $p < 0.001$) and a group × window interaction ($\chi^2(2) = 7.26$, $p = 0.026$), but no main effect of group ($\chi^2(1) = 1.87$, $p = 0.171$). Post hoc contrasts indicated reduced beta power during the task relative to baseline in both groups (both $p \leq 0.002$), consistent with movement-related beta desynchronization, with a larger reduction in the comparison group; however, no group differences were observed within any individual window (all $p > 0.10$). Ipsilateral ExF showed a similar interaction ($\chi^2(2) = 6.29$, $p = 0.043$) without a main effect of group ($\chi^2(1) = 1.87$, $p = 0.171$).

For ExG, both hemispheres exhibited a robust main effect of window (contralateral: $\chi^2(2) = 12.46$, $p = 0.002$; ipsilateral: $\chi^2(2) = 15.23$, $p < 0.001$), again reflecting reduced beta power during task relative to baseline (all $p \leq 0.0004$), but neither hemisphere showed a main effect of group (both $p \geq 0.94$) nor a group × window interaction (both $p \geq 0.23$). Importantly, there was no consistent increase in post-task beta power relative to baseline in any condition, indicating the absence of a reliable beta rebound at the group level. These findings indicate that conventional beta power captures expected movement-related desynchronization but does not reveal stable group differences.

Figure 3. Task-related modulation of sensorimotor beta power during fine and gross motor execution. Time courses of baseline-corrected beta power (% change from baseline) over sensorimotor cortex during execution of fine (ExF) and gross (ExG) actions, shown separately for the hemisphere contralateral (left column) and ipsilateral (right column) to the hand used. Solid lines depict group means for the comparison (green) and DCD (purple) groups; shaded regions indicate ±SEM across participants. Vertical dashed lines mark task onset (0 s) and offset (10 s); the horizontal dashed line indicates baseline (0%). Colored asterisks denote significant within-group deviations from baseline (one-sample permutation tests, $p < 0.05$).

We next assessed whether sensorimotor beta activity exhibited sustained rhythmic structure using lagged Hilbert autocoherence (LHaC) (Zhang et al., 2025). Across groups, LHaC values were lower within the identified beta range (Figure 4a-b) relative to lower frequencies, indicating limited long-range periodicity and supporting the characterization of beta activity as transient rather than sustained rhythmic oscillations (Figure 4c-d). No qualitative differences between groups were apparent in the spectral or autocoherence profiles.

We then applied an iterative burst detection algorithm (Rayson et al., 2023; Szul et al., 2023) to identify candidate beta burst events and compared time-frequency-based burst features between groups, pooling bursts detected over left and right sensorimotor cortex (C3/C4). There were no group differences in burst duration (FWHM time; $M_{comparison}$ = 0.420 s, SD = 0.049; M_{DCD} = 0.418 s, SD = 0.053; p = 0.868), peak amplitude ($M_{comparison}$ = 0.943 µV, SD = 0.321; M_{DCD} = 0.940 µV, SD = 0.500; p = 0.979), peak frequency ($M_{comparison}$ = 19.54 Hz, SD = 1.17; M_{DCD} = 19.66 Hz, SD = 1.39; p = 0.762), or frequency span (FWHM frequency; $M_{comparison}$ = 1.46 Hz, SD = 0.09; M_{DCD} = 1.45 Hz, SD = 0.13; p

557 = 0.964; Figure 4e-h). Finally, we examined overall burst rate across execution (ExF,
558 ExG) conditions, expressed relative to the pre-task baseline. Burst rate did not differ
559 between groups in any condition (all p < 0.152). Conventional time-frequency burst
560 metrics and burst incidence therefore do not distinguish DCD from comparison
561 participants, motivating a more detailed analysis of burst waveform structure.

562
Figure 4. Spectral characteristics and time-frequency burst features do not differ between groups.
(a) Log power spectral density over central sensorimotor electrodes (C3/C4), averaged across participants,
for comparison (green) and DCD (purple) groups; shaded areas indicate ±SEM. (b) Periodic (aperiodic-
corrected) power spectra derived using spectral parameterization. Vertical dashed lines in (a-b) mark the
beta band used for burst detection (14-31.25 Hz). (c-d) Mean lagged Hilbert autocoherence (LHaC) as a
function of frequency (≤35 Hz) and lag (cycles), averaged across hemispheres, shown for the comparison
group (c) and DCD group (d); white dashed lines indicate the beta band. Color scale is matched across
groups. (e-h) Distributions of burst features pooled across task conditions: duration (FWHM in time; ms)
(e), peak amplitude (µV) (f), peak frequency (Hz) (g), and frequency span (FWHM in frequency; Hz) (h).
Across all spectral and burst-level metrics, groups did not differ significantly.

573
3.2 Sensorimotor cortical bursts share similar waveform motifs across groups
Because time-frequency-based beta burst features are derived from time-
frequency decomposition of the burst event time series, and the correspondence between
temporal and time-frequency-based burst features is not one-to-one (Jones, 2016; Szul
et al., 2023), we then focused our analysis on burst waveform shapes. In line with
previous work, we found that sensorimotor cortical bursts in both groups had a wavelet-
like median waveform shape with a strong central negative deflection and surrounding
positive deflections (Figure 5a; Sherman et al., 2016; Bonaiuto et al., 2021; Brady and
Bardouille, 2022; Rayson et al., 2023; Szul et al., 2023). To assess group differences in
burst waveform shape, we next applied PCA to the burst waveforms, an approach
previously shown to capture functionally relevant waveform motifs (Rayson et al., 2023;
Szul et al., 2023). We ran separate PCAs on burst waveforms from each group and
compared the resulting eigenvectors. All PCs were highly similar between groups (ρ =
0.85 - 1.0; Figure 5b). For nearly all PCs, the most similar component from the comparison
group PCA was the corresponding one of the DCD group PCA, indicating that the

waveform motif and relative percentage of variance explained was very similar across groups.

Figure 5. Burst waveform motifs and task-related modulation revealed by PCA. (a) Median burst waveforms over sensorimotor cortex for comparison (green) and DCD (purple) groups, showing highly similar canonical beta burst morphology. (b) Absolute Spearman correlation coefficients between eigenvectors obtained from separate PCAs performed on comparison and DCD bursts, demonstrating strong correspondence of waveform components across groups. (c) Proportion of variance explained by each principal component (bars). Grey shading indicates components exceeding the permutation-based significance threshold derived from waveform shuffling. Overlaid lines show the mean absolute change in PC score from baseline during execution of fine and gross actions (averaged across groups). Blue asterisks mark components exhibiting significant execution-related modulation. (d) For each significantly modulated component, mean waveform shapes of bursts falling within each score tertile (T1-T3; colored lines), plotted against the overall mean waveform (black), illustrating distinct waveform motifs captured along each principal axis.

3.3 Specific burst waveform motifs are modulated by action execution tasks

Given the similarity in waveform shape and variability between comparison and DCD bursts, we next performed a single global PCA on all sensorimotor burst waveforms to represent bursts within a common low-dimensional space. Components retained for further analysis were determined using a temporal-shuffling permutation procedure, which identified those PCs whose variance exceeded that expected under waveform phase randomization (PCs 1-11, all $p < 0.001$). For each retained component, we then quantified condition-related modulation by computing, separately for each group, the difference in mean PC score between execution (ExF, ExG) and pre-task baseline, averaging within subjects and testing significance using paired permutation tests. This analysis revealed selective modulation during gross motor execution (ExG) in the comparison group only. Specifically, PC3 ($p = 0.019$), PC6 ($p = 0.042$), and PC8 ($p = 0.045$) showed significant baseline-to-task changes in the comparison group. No PCs showed significant modulation during fine motor execution (ExF), and no components reached significance in the DCD group for either condition. Each of these components

captures distinct deviations from the mean burst waveform, reflecting variation in the
depth of the central negative deflection and the relative prominence of surrounding peaks.
Because changes in mean PC score are ambiguous with respect to burst incidence (i.e.,
an increase in mean score could reflect increased occurrence of high-scoring bursts or
decreased occurrence of low-scoring bursts), we next examined these components in
terms of waveform-specific burst rate.

626 *3.4 Waveform-specific burst rates differ between DCD and comparison groups*

For each PC that showed task-related modulation, we quantified waveform-
specific burst rate by binning bursts into tertiles of PC score and computing the mean
burst rate (Hz) within three predefined epoch windows (baseline, task, and post-task).
Burst-rate time courses were estimated separately for contralateral and ipsilateral
sensorimotor electrodes during ExF and ExG. Cluster-based permutation testing of the
burst-rate time series revealed ipsilateral group differences that were selective to bursts
with specific waveform shapes, with significant between-group effects for the first and
third tertiles of PC6 during ExG (Figure 6) and the third tertile of PC8 during ExF (Figure
7); no other PCs or hemispheres showed reliable task-related group differences.

Importantly, these effects reflected opposing task-related modulations across the
extremes of the PC dimensions rather than uniform changes in overall bursting. Cluster-
based permutation tests of the time-resolved burst rate revealed cross-over patterns in
the ipsilateral hemisphere during movement execution. For PC6 (ExG, ipsilateral), the
lowest-score tertile (T1) showed a decrease in burst rate during movement in the
comparison group but an increase in the DCD group, whereas the highest-score tertile
(T3) showed the inverse pattern (Figure 6). A similar divergence was observed for PC8
(ExF, ipsilateral), where the highest-score tertile (T3) exhibited a task-related decrease in
burst rate in the comparison group and a concurrent increase in the DCD group (Figure
7). These patterns indicate a redistribution of burst occurrence toward opposite ends of
the waveform dimensions between groups.

Figure 6. Waveform-specific burst-rate differences between groups along PC6 during gross motor execution (ExG). Left column: Mean waveform shapes of bursts falling into each tertile (T1-T3; color-coded from cyan to magenta) of PC6 scores, shown relative to the overall mean waveform (black), illustrating the distinct waveform motifs captured along this principal axis. Middle and right panels: Time-resolved, baseline-corrected burst rates (ΔHz) for comparison (solid lines) and DCD (dashed lines) groups during gross motor execution (ExG), plotted separately for the hemisphere contralateral and ipsilateral to the dominant hand. Shaded regions indicate ±SEM across participants. Vertical dashed lines mark task onset (0 s) and offset (10 s); the horizontal dashed line indicates baseline (0 Hz). Red asterisks denote significant between-group clusters identified using cluster-based permutation testing ($p < 0.05$).

Figure 7. Waveform-specific burst-rate differences between groups along PC8 during fine motor execution (ExF). Left column: Mean waveform shapes of bursts falling into each tertile (T1-T3; color-coded from cyan to magenta) of PC8 scores, shown relative to the overall mean waveform (black), illustrating the distinct waveform motifs captured along this principal axis. Middle and right panels: Time-resolved, baseline-corrected burst rates (ΔHz) for comparison (solid lines) and DCD (dashed lines) groups during fine motor execution (ExF), plotted separately for the hemisphere contralateral and ipsilateral to the dominant hand. Shaded regions indicate ±SEM across participants. Vertical dashed lines mark task onset (0 s) and offset (10 s); the horizontal dashed line indicates baseline (0 Hz). Red asterisks denote significant between-group clusters identified using cluster-based permutation testing ($p < 0.05$).

To quantify these effects, we performed follow-up linear mixed-effects analyses on window-averaged burst rates (task minus baseline), focusing on PCs and conditions identified in the time-resolved analysis. For PC6 during gross motor execution (ExG), significant between-group differences were confined to the ipsilateral hemisphere and were selective to the extreme tertiles. In the lowest-score tertile (T1), burst rate differed significantly between groups ($\chi^2(1) = 8.59$, $p = 0.003$), with the DCD group showing a higher task-related increase relative to the comparison group. In contrast, the highest-score tertile (T3) showed the opposite pattern ($\chi^2(1) = 5.55$, $p = 0.018$), reflecting a reduction in burst rate in the DCD group relative to controls. No group effects were observed for the middle tertile (T2) or in the contralateral hemisphere (all $p > 0.12$). For

PC8 during fine motor execution (ExF), a significant group effect was observed only for the ipsilateral highest-score tertile (T3) ($\chi^2(1) = 8.96$, $p = 0.0028$). Here, the comparison group exhibited a task-related decrease in burst rate, whereas the DCD group showed a relative increase. No significant group effects were present for lower tertiles, the contralateral hemisphere, or during ExG (all $p > 0.13$). Together, these analyses converge to show that execution-related group differences are waveform-specific, ipsilaterally expressed, and restricted to the extremes of particular waveform dimensions (PC6 and PC8). Rather than reflecting a uniform shift in burst rate, these effects indicate a redistribution of burst occurrence across distinct waveform motifs in DCD, consistent with altered engagement of specific beta burst subtypes during motor execution.

3.5 Waveform-specific burst rates along PC6 and PC8 correlate with motor skill

Given that execution-related group differences were selective to ipsilateral burst subtypes defined by PC6 (ExG) and PC8 (ExF), we next asked whether these task-related, baseline-corrected burst-rate changes scaled with motor performance across individuals (Figure 9). To capture variability across the full sample, we included all participants, including comparison-group children with MABC-2 scores below the 25th percentile. We quantified, for each participant, the mean task-baseline change in burst rate for bursts in each tertile of PC score (ipsilateral hemisphere; ExG for PC6 and ExF for PC8), and related these values to MABC-2 total score using Spearman rank correlations.

For PC6 during ExG, motor performance was strongly related to the two extreme tertiles in opposite directions: T1 burst-rate change was negatively associated with MABC-2 ($\rho = -0.55$, $p < 0.001$), whereas T3 burst-rate change was positively associated with MABC-2 ($\rho = 0.54$, $p < 0.001$). Robust (HC3) group-wise linear fits were consistent with this pattern: for PC6-T1, slopes were negative in both groups (comparison: slope = -28.85, $p = 0.020$; DCD: slope = -47.03, $p = 0.039$), and for PC6-T3, slopes were positive in both groups (comparison: slope = 30.63, $p < 0.001$; DCD: slope = 38.07, $p = 0.023$). This mirrors the redistribution observed in the group contrasts: better motor performance corresponded to fewer low-PC6 (T1) bursts and more high-PC6 (T3) bursts during ExG in both groups. For PC8 during ExF, the association with motor performance was specific to the highest tertile (T3), where task-baseline burst-rate change was negatively correlated with MABC-2 ($\rho = -0.42$, $p = 0.004$). However, this relationship was not reliable within either group alone using robust linear fits (comparison: slope = -23.52, $p = 0.099$; DCD: slope = 9.22, $p = 0.760$), suggesting that the effect is expressed primarily across the pooled sample rather than as a consistent within-group slope.

Figure 9. **Waveform-specific burst rates predict motor performance in a motif- and condition-dependent manner.** Scatter plots show the relationship between baseline-corrected burst rate (ipsilateral hemisphere) and MABC-2 total score for bursts falling into each tertile (T1-T3) of the indicated principal component (PC6, PC8) during execution conditions. Points represent individual participants (comparison: green; DCD: purple). Solid lines indicate group-wise linear fits, with shaded regions denoting bootstrapped 95% confidence intervals (10,000 iterations); fits are shown where $p < 0.05$.

3.6 Waveform-specific interhemispheric connectivity differences in DCD during motor execution

Figure 10. Waveform-specific, hemisphere-resolved Granger causality during execution for PC6 (ExG) and PC8 (ExF). Signed Granger causality (GC) matrices computed from burst-rate time series (0-10 s post-movement onset) for bursts stratified by waveform tertile within a given principal component. Units correspond to ipsilateral and contralateral sensorimotor electrodes, separated by black lines; within each hemisphere, rows and columns represent PC tertiles (T1-T3; waveform motifs illustrated). Top row (PC6, ExG): (a) DCD group; (b) comparison group; (c) between-group difference (DCD - comparison). Bottom row (PC8, ExF): (d) DCD group; (e) comparison group; (f) between-group difference (DCD - comparison). Colored cells indicate significant GC values after permutation testing with FDR correction (within-group panels) or significant group differences from subject-label permutation (right panels); non-significant connections are set to zero. Warmer colors denote stronger positive signed GC (directional influence), and cooler colors denote negative signed GC.

To test whether motif- and hemisphere-specific differences in burst dynamics were accompanied by altered directed interactions between sensorimotor cortices, we examined signed Granger causality (GC) between ipsilateral and contralateral burst-rate time series. GC was computed separately for each waveform tertile within a given PC and execution condition. Within-hemisphere interactions were excluded because the burst detection procedure precludes temporal overlap and therefore trivially constrains intra-hemispheric coupling. Statistical significance was assessed using permutation testing with FDR correction for within-group effects and subject-label permutation for between-group contrasts.

For PC6 during gross motor execution (ExG), clear group differences in interhemispheric connectivity were observed (Figure 10a-c). Two directed connections differed significantly between groups. Connectivity from ipsilateral T1 to contralateral T2 was inhibitory in the comparison group but excitatory in the DCD group, indicating a reversal in the sign of directional influence. In addition, connectivity from contralateral T3

to ipsilateral T1 was excitatory in the comparison group but inhibitory in DCD, again
reflecting a qualitative change in coupling rather than a simple attenuation. These effects
show that, for PC6, DCD is associated with sign-inverted interhemispheric interactions
between bursts at opposing ends of the waveform dimension. For PC8 during fine motor
execution (ExF), group differences were more circumscribed (Figure 10d-f). A significant
effect was observed for the connection from ipsilateral T1 to contralateral T3, which was
inhibitory in DCD but not significant in the comparison group. No additional
interhemispheric connections showed reliable group differences for PC8. Overall, the
connectivity analyses reveal that group differences in beta burst dynamics are
accompanied by selective disruptions of interhemispheric communication, expressed in
a waveform-dependent manner and, in some cases, involving reversals in the sign of
directed influence between sensorimotor hemispheres during motor execution.

4. Discussion

This study applied waveform-based beta burst analyses to investigate how
sensorimotor burst motifs relate to motor performance in childhood. Conventional time-
frequency measures, including beta power and burst spectral features, did not differ
between groups, and the overall structure of burst waveform variability was highly similar
in children with and without developmental coordination disorder. Instead, differences
emerged only when examining burst dynamics along specific waveform dimensions.
Movement execution was associated with systematic redistribution of bursts along these
dimensions, and the prevalence and lateralization of particular motifs co-varied with motor
coordination. Children with DCD were disproportionately represented at one end of this
motif-defined axis, but the relation between waveform-specific burst rate and MABC-2
scores followed a similar slope across groups. These findings indicate that beta bursts
comprise functionally heterogeneous events (Rayson et al., 2023, 2025; Szul et al., 2023;
Zich et al., 2023; Moreau et al., 2026) whose motif-specific dynamics track continuous
variation in motor competence, with DCD providing a clinically informative model of
extreme motor difficulty rather than a categorically distinct beta phenotype.

Our results further suggest that different beta burst waveform motifs contribute
differently to sensorimotor processing during action execution. Rather than reflecting
discrete burst classes, the principal components derived from waveform analysis capture
continuous dimensions of burst shape variability along which bursts can be redistributed
depending on task demands and motor performance. Group differences emerged
selectively at the extremes of these dimensions: during gross motor execution, bursts
with low versus high PC6 scores showed opposing changes in rate across groups, while
during fine motor execution a similar divergence was observed for bursts occupying the
highest tertile of PC8. Importantly, these effects did not involve global changes in burst
incidence or spectral burst features. Instead, motor-related differences reflected shifts in
the relative prevalence of bursts occupying specific regions of the waveform variability

space. This pattern suggests that sensorimotor beta bursts form a heterogeneous
population of transient events whose functional contribution depends on their waveform
motif.

A notable feature of these effects was their localization to the ipsilateral
hemisphere. Although sensorimotor beta activity is often interpreted primarily in terms of
contralateral motor control, ipsilateral sensorimotor cortex plays an important role in
coordinating activity across hemispheres during movement (Serrien et al., 2006). The
motif-specific redistribution observed here may therefore reflect differences in how
transient beta events participate in interhemispheric sensorimotor communication
(Seedat et al., 2020). Consistent with this interpretation, directed connectivity analyses
revealed motif-specific differences in interhemispheric interactions, including changes in
the sign of influence for particular burst tertiles. These findings suggest that the
coordination of activity between hemispheres depends not only on the timing of beta
bursts but also on the waveform motifs through which these events are expressed.

The relation between burst dynamics and motor performance further supports a
dimensional interpretation at the level of brain–behavior mapping. Although children with
DCD exhibited poorer motor performance overall, the association between waveform-
specific burst rate and MABC-2 score was comparable across groups, indicating that
similar neural-behavioral relationships are expressed across the full sample. This pattern
suggests that variation in sensorimotor beta burst dynamics may index sensorimotor
performance, rather than an underlying process specific to DCD.

The present findings therefore complement existing accounts of motor impairment
in DCD that emphasize atypical sensorimotor connectivity and hemispheric specialization
(Tallet et al., 2013; McLeod et al., 2014, 2016). Previous work has linked these differences
to alterations in corpus callosum-mediated motor coordination and interhemispheric
inhibition. The motif-specific redistribution observed here provides a potential
physiological substrate for these network-level differences. If distinct beta burst motifs
reflect different modes of sensorimotor processing, then altered recruitment of these
motifs could influence how information is exchanged between hemispheres during action
planning and execution. Such differences may ultimately contribute to the motor planning
and adaptation difficulties often observed in DCD (Subara-Zukic et al., 2022). More
broadly, these findings suggest that the functional role of beta bursts in motor control may
depend on the distribution of waveform motifs within the burst population rather than on
overall burst incidence alone.

Task demands may also shape how these motif dynamics are expressed. The
motif-specific differences observed here varied between fine and gross motor conditions,
suggesting that the recruitment of particular burst motifs depends on the structure of the

motor task. Fine motor actions require precise and sustained control of a specific effector,
whereas gross motor actions often involve broader muscle synergies and more
distributed sensorimotor engagement. Differences in effector specificity and bilateral
coordination may therefore influence how burst motifs are distributed across hemispheres
and how strongly ipsilateral sensorimotor cortex contributes to task execution. In children
with DCD, altered motif recruitment during these tasks may reflect differences in the ability
to coordinate sensorimotor activity across hemispheres when task demands require
flexible integration of bilateral motor signals, for example when stabilizing one hand while
executing a precise action with the other, or when suppressing unintended mirror
movements.

The present study has several strengths. The inclusion of execution, observation,
and non-biological motion conditions allowed us to isolate sensorimotor-specific effects
and demonstrate that motif-related differences were not simply driven by visual or
attentional factors. Moreover, this study represents the first application of waveform-
based beta burst analysis to a neurodevelopmental population, demonstrating the utility
of this approach for characterizing subtle differences in sensorimotor network dynamics
that are not apparent using conventional spectral measures. Nevertheless, several
limitations should be noted. The sample size, while comparable to previous EEG studies
in DCD, was relatively modest, and replication in larger cohorts will be important. In
addition, the present analyses focused on central electrodes overlying sensorimotor
cortex; beta bursts arising from associative or prefrontal regions may also contribute to
motor planning and control (Wessel, 2020; Diesburg et al., 2021; Liljefors et al., 2024).
Finally, because the experimental design used relatively long task epochs, the temporal
relation between individual movements and burst events could not be examined with high
precision. Future studies using event-locked designs will be important for understanding
how motif-specific bursts relate to individual movement components and error signals.

In conclusion, our findings demonstrate that sensorimotor beta burst dynamics in
childhood vary systematically along waveform-defined dimensions that relate to motor
performance. Rather than reflecting global differences in beta activity, motor coordination
differences were associated with motif-specific redistribution of bursts within the broader
population of beta events. These results support emerging models in which the functional
heterogeneity of beta bursts plays a central role in sensorimotor computation. By linking
waveform-defined burst motifs to motor competence across development, this work
provides new insight into the neural mechanisms underlying motor coordination
differences and highlights the potential of burst-based approaches for studying atypical
motor development.

References

Agouram H, Szul MJ, Eusebio A, Daucé E, Sorrentino P, Bonaiuto JJ (2025) Beta Burst Waveform Motifs Link Cortico-Subcortical Connectivity and Clinical Improvement in Parkinson's Disease. :2025.07.09.25331208 Available at: https://www.medrxiv.org/content/10.1101/2025.07.09.25331208v1.

Bates D, Mächler M, Bolker B, Walker S (2014) Fitting linear mixed-effects models using lme4. ArXiv Prepr ArXiv14065823.

Bonaiuto J, Little S, Neymotin SA, Jones SR, Barnes GR, Bestmann S (2021) Laminar dynamics of high amplitude beta bursts in human motor cortex. NeuroImage 242:118479.

Boto E, Shah V, Hill RM, Rhodes N, Osborne J, Doyle C, Holmes N, Rea M, Leggett J, Bowtell R, Brookes MJ (2022) Triaxial detection of the neuromagnetic field using optically-pumped magnetometry: feasibility and application in children. NeuroImage 252:119027.

Brady B, Bardouille T (2022) Periodic/Aperiodic parameterization of transient oscillations (PAPTO)–Implications for healthy ageing. NeuroImage 251:118974.

Brown-Lum M, Zwicker JG (2015) Brain Imaging Increases Our Understanding of Developmental Coordination Disorder: a Review of Literature and Future Directions. Curr Dev Disord Rep 2:131–140.

Cheyne D, Jobst C, Tesan G, Crain S, Johnson B (2014) Movement-related neuromagnetic fields in preschool age children. Hum Brain Mapp 35:4858–4875.

de Cheveigné A (2020) ZapLine: A simple and effective method to remove power line artifacts. NeuroImage 207:116356.

Debnath R, Buzzell GA, Morales S, Bowers ME, Leach SC, Fox NA (2020) The Maryland analysis of developmental EEG (MADE) pipeline. Psychophysiology 57:e13580.

Delorme A, Makeig S (2004) EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. J Neurosci Methods 134:9–21.

Diesburg DA, Greenlee JD, Wessel JR (2021) Cortico-subcortical β burst dynamics underlying movement cancellation in humans. elife 10:e70270.

Donoghue T, Haller M, Peterson EJ, Varma P, Sebastian P, Gao R, Noto T, Lara AH, Wallis JD, Knight RT, Shestyuk A, Voytek B (2020) Parameterizing neural power spectra into periodic and aperiodic components. Nat Neurosci 23:1655–1665.

Drew AR, Quandt LC, Marshall PJ (2015) Visual influences on sensorimotor EEG responses during observation of hand actions. Brain Res 1597:119–128.

Feingold J, Gibson DJ, DePasquale B, Graybiel AM (2015) Bursts of beta oscillation differentiate postperformance activity in the striatum and motor cortex of monkeys performing movement tasks. Proc Natl Acad Sci U S A 112:13687–13692.

Fox J, Weisberg S (2018) An R companion to applied regression. Sage publications.

French B, Sycamore NJ, McGlashan HL, Blanchard CC, Holmes NP (2018) Ceiling effects in the Movement Assessment Battery for Children-2 (MABC-2) suggest that non-parametric scoring methods are required. PLoS One 13:e0198426.

Fu Y, Franz EA (2014) Viewer perspective in the mirroring of actions. Exp Brain Res 232:3665–3674.

Gaetz W, MacDonald M, Cheyne D, Snead OC (2010) Neuromagnetic imaging of movement-related cortical oscillations in children and adults: Age predicts post-movement beta rebound. NeuroImage 51:792–807.

Henderson SE, Sugden D, Barnett AL (1992) Movement assessment battery for children-2. Res Dev Disabil.

Henderson SE, Sugden DA, Barnett AL (2007) Movement assessment battery for children, 2nd ed. London, UK: The Psychological Corporation.

Hobson HM, Bishop DV (2016) Mu suppression–a good measure of the human mirror neuron system? Cortex J Devoted Study Nerv Syst Behav 82:290.

Jackson PL, Meltzoff AN, Decety J (2006) Neural circuits involved in imitation and perspective-taking. Neuroimage 31:429–439.

Jeannerod M (2001) Neural Simulation of Action: A Unifying Mechanism for Motor Cognition. NeuroImage 14:S103–S109.

Jones SR (2016) When brain rhythms aren't 'rhythmic': implication for their mechanisms and meaning. Curr Opin Neurobiol 40:72–80.

Keating J, Gerson SA, Jones CRG, Vanderwert RE, Purcell C (2023) Possible disrupted biological movement processing in Developmental Coordination Disorder. Cortex 168:1–13.

Kim S, Putrino D, Ghosh S, Brown EN (2011) A Granger Causality Measure for Point Process Models of Ensemble Neural Spiking Activity. PLOS Comput Biol 7:e1001110.

Kumaravel VP, Farella E, Parise E, Buiatti M (2022) NEAR: An artifact removal
pipeline for human newborn EEG data. Dev Cogn Neurosci 54:101068.

Leach SC, Morales S, Bowers ME, Buzzell GA, Debnath R, Beall D, Fox NA (2020)
Adjusting ADJUST: Optimizing the ADJUST algorithm for pediatric data using
geodesic nets. Psychophysiology 57:e13566.

Lenth R, Buerkner P, Herve M, Love J, Riebl H, Singmann H (2020) emmeans:
estimated marginal means, aka least-squares means, v1. 5.1. Vienna R Core
Team.

Liljefors J, Almeida R, Rane G, Lundström JN, Herman P, Lundqvist M (2024)
Distinct functions for beta and alpha bursts in gating of human working memory.
Nat Commun 15:8950.

Little S, Bonaiuto J, Barnes G, Bestmann S (2019) Human motor cortical beta
bursts relate to movement planning and response errors. PLOS Biol 17:e3000479.

McLeod KR, Langevin LM, Dewey D, Goodyear BG (2016) Atypical within-and
between-hemisphere motor network functional connections in children with
developmental coordination disorder and attention-deficit/hyperactivity disorder.
NeuroImage Clin 12:157–164.

McLeod KR, Langevin LM, Goodyear BG, Dewey D (2014) Functional connectivity
of neural motor networks is disrupted in children with developmental coordination
disorder and attention-deficit/hyperactivity disorder. NeuroImage Clin 4:566–575.

Moca VV, Bârzan H, Nagy-Dăbâcan A, Mureșan RC (2021) Time-frequency super-
resolution with superlets. Nat Commun 12:337.

Mognon A, Jovicich J, Bruzzone L, Buiatti M (2011) ADJUST: An automatic EEG
artifact detector based on the joint use of spatial and temporal features.
Psychophysiology 48:229–240.

Moreau Q, Szul MJ, Daligaut S, Schwartz D, Bonaiuto JJ (2026) Distinct beta burst
motifs exhibit opposing error relationships during motor adaptation.
:2026.03.06.710026 Available at:
https://www.biorxiv.org/content/10.64898/2026.03.06.710026v2.

Papadopoulos S, Darmet L, Szul MJ, Congedo M, Bonaiuto JJ, Mattout J (2024)
Surfing beta burst waveforms to improve motor imagery-based BCI. Imaging
Neurosci 2:imag–2–00391.

Pfurtscheller G (1981) Central beta rhythm during sensorimotor activities in man.
Electroencephalogr Clin Neurophysiol 51:253–264.

Pfurtscheller G, Stancák A, Neuper C (1996) Post-movement beta
synchronization. A correlate of an idling motor area? Electroencephalogr Clin
Neurophysiol 98:281–293.

Press C, Cook J, Blakemore S-J, Kilner J (2011) Dynamic Modulation of Human
Motor Activity When Observing Actions. J Neurosci 31:2792–2800.

Querne L, Berquin P, Vernier-Hauvette M-P, Fall S, Deltour L, Meyer M-E, de
Marco G (2008) Dysfunction of the attentional brain network in children with
Developmental Coordination Disorder: A fMRI study. Brain Res 1244:89–102.

R Core Team (2022) R: A language and environment for statistical computing.
Vienna, Austria: R Foundation for Statistical Computing.

Rayson H, Debnath R, Alavizadeh S, Fox N, Ferrari PF, Bonaiuto JJ (2022)
Detection and analysis of cortical beta bursts in developmental EEG data. Dev
Cogn Neurosci 54:101069.

Rayson H, Moreau Q, Gailhard S, Szul MJ, Bonaiuto JJ (2025) Beta Burst
Waveform Diversity: A Window onto Cortical Computation. The
Neuroscientist:10738584251390779.

Rayson H, Szul MJ, El-Khoueiry P, Debnath R, Gautier-Martins M, Ferrari PF, Fox
N, Bonaiuto JJ (2023) Bursting with Potential: How Sensorimotor Beta Bursts
Develop from Infancy to Adulthood. J Neurosci 43:8487–8503.

Rier L, Rhodes N, Pakenham DO, Boto E, Holmes N, Hill RM, Reina Rivero G,
Shah V, Doyle C, Osborne J, Bowtell RW, Taylor M, Brookes MJ (2024) Tracking
the neurodevelopmental trajectory of beta band oscillations with optically pumped
magnetometer-based magnetoencephalography Luo H, Colgin LL, eds. eLife
13:RP94561.

Seedat ZA, Quinn AJ, Vidaurre D, Liuzzi L, Gascoyne LE, Hunt BAE, O'Neill GC,
Pakenham DO, Mullinger KJ, Morris PG, Woolrich MW, Brookes MJ (2020) The
role of transient spectral 'bursts' in functional connectivity: A
magnetoencephalography study. NeuroImage 209:116537.

Serrien DJ, Ivry RB, Swinnen SP (2006) Dynamics of hemispheric specialization
and integration in the context of motor control. Nat Rev Neurosci 7:160–166.

Sherman MA, Lee S, Law R, Haegens S, Thorn CA, Hämäläinen MS, Moore CI,
Jones SR (2016) Neural mechanisms of transient neocortical beta rhythms:
Converging evidence from humans, computational modeling, monkeys, and mice.
Proc Natl Acad Sci 113:E4885–E4894.

Subara-Zukic E, Cole MH, McGuckian TB, Steenbergen B, Green D, Smits-
Engelsman BC, Lust JM, Abdollahipour R, Domellöf E, Deconinck FJ (2022)
Behavioral and neuroimaging research on developmental coordination disorder

(DCD): A combined systematic review and meta-analysis of recent findings. Front Psychol 13:809455.

Sugden DA (2006) Development Coordination Disorder as a specific learning difficulty. Leeds Consens Statement ESRC:2004–2005.

Szul MJ, Papadopoulos S, Alavizadeh S, Daligaut S, Schwartz D, Mattout J, Bonaiuto JJ (2023) Diverse beta burst waveform motifs characterize movement-related cortical dynamics. Prog Neurobiol 228:102490.

Tallet J, Albaret J-M, Barral J (2013) Developmental changes in lateralized inhibition of symmetric movements in children with and without Developmental Coordination Disorder. Res Dev Disabil 34:2523–2532.

Veale JF (2014) Edinburgh Handedness Inventory – Short Form: A revised version based on confirmatory factor analysis. Laterality Asymmetries Body Brain Cogn 19:164–177.

Vieira VM (2012) Permutation tests to estimate significances on Principal Components Analysis. Comput Ecol Softw 2:103.

Weschler D (2011) WASI-II: Wechsler abbreviated scale of intelligence–second edition.

Wessel JR (2020) β-bursts reveal the trial-to-trial dynamics of movement initiation and cancellation. J Neurosci 40:411–423.

West TO, Duchet B, Farmer SF, Friston KJ, Cagnan H (2023) When do bursts matter in the primary motor cortex? Investigating changes in the intermittencies of beta rhythms associated with movement states. Prog Neurobiol 221:102397.

Wilkinson CL, Yankowitz LD, Chao JY, Gutiérrez R, Rhoades JL, Shinnar S, Purdon PL, Nelson CA (2024) Developmental trajectories of EEG aperiodic and periodic components in children 2–44 months of age. Nat Commun 15:5788.

Wilson PH, Smits-Engelsman B, Caeyenberghs K, Steenbergen B, Sugden D, Clark J, Mumford N, Blank R (2017) Cognitive and neuroimaging findings in developmental coordination disorder: new insights from a systematic review of recent research. Dev Med Child Neurol 59:1117–1129.

Wuang Y, Su C, Huang M (2012) Psychometric comparisons of three measures for assessing motor functions in preschoolers with intellectual disabilities. J Intellect Disabil Res 56:567–578.

Zhang S, Szul MJ, Papadopoulos S, Massera A, Rayson H, Bonaiuto JJ (2025) Multi-scale parameterization of neural rhythmicity with lagged Hilbert autocoherence. Imaging Neurosci 3:IMAG.a.993.

1035   Zich C, Quinn AJ, Bonaiuto JJ, O'Neill G, Mardell LC, Ward NS, Bestmann S
1036   (2023) Spatiotemporal organisation of human sensorimotor beta burst activity.
1037   Elife 12:e80160.

1038 **Supplementary Material**

1039

1040

Figure S1. Task-related modulation of sensorimotor beta power during observation conditions. Time
courses of baseline-corrected beta power (% change from baseline) over sensorimotor cortex during
observation of a kaleidoscope (Kal), as well as fine (ObF) and gross (ObG) actions, shown separately for
the hemisphere contralateral (left column) and ipsilateral (right column) to the hand used. Solid lines depict
group means for the comparison (green) and DCD (purple) groups; shaded regions indicate ±SEM across
participants. Vertical dashed lines mark task onset (0 s) and offset (10 s); the horizontal dashed line
indicates baseline (0%). Colored asterisks denote significant within-group deviations from baseline (one-
sample permutation tests, $p < 0.05$).