Helicase-mediated mechanism of SSU processome maturation and disassembly

Olga Buzovetsky[1] and Sebastian Klinge[1,*]

1 Laboratory of Protein and Nucleic Acid Chemistry, The Rockefeller University, New York, New York 10065, USA

*Correspondence: klinge@rockefeller.edu (S.K.)

ABSTRACT

Eukaryotic ribosomal small subunit (SSU) assembly requires the SSU processome, a

nucleolar precursor containing the RNA chaperone U3 snoRNA. The underlying

molecular mechanisms of SSU processome maturation, remodeling, disassembly, RNA

quality control, and the transitions between states remain elusive due to a paucity of

intermediates. Here we report 16 native SSU processome structures alongside genetic

data, revealing how two helicases, the Mtr4-exosome and Dhr1, are controlled for

accurate and unidirectional ribosome biogenesis. Our data show how irreversible pre-

ribosomal RNA degradation by the redundantly tethered RNA exosome couples the

transformation of the SSU processome into a pre-40S particle during which Utp14 can

probe evolving surfaces, ultimately positioning and activating Dhr1 to unwind the U3

snoRNA and initiate nucleolar pre-40S release. This study highlights a paradigm for large

dynamic RNA-protein complexes where irreversible RNA degradation drives

compositional changes and communicates these changes to govern enzyme activity

while maintaining overall quality control.

MAIN

The nucleolus, a membrane-less organelle, is generated via eukaryotic ribosome biogenesis in a transcription-dependent manner. The maturation of ribosomal small subunit (SSU) and large subunit (LSU) precursors generates distinct phases of this multi-layered organelle, which is maintained via multi-valent RNA-protein and protein-protein interactions. Early stages of eukaryotic ribosomal small subunit (SSU/40S) assembly involve the SSU processome, a giant nucleolar particle containing approximately a quarter of all eukaryotic ribosome assembly factors and three RNAs (SSU pre-rRNA/18S, 5′ ETS and the RNA chaperone U3 snoRNA)[1–3]. The 18S is composed of four domains (5′, central, 3′ major, and 3′ minor) which are housed in separate modules. By serving as the architectural blueprint for the SSU processome, the chaperone U3 snoRNA base-pairs with parts of the SSU pre-rRNA as well as the 5′ ETS. High-resolution structural studies of SSU processome assembly and maturation in yeast and human cells have started to reveal isolated native states before and after cleavage of the mature rRNA 5′ end (site A1)[4–13]. After a poorly characterized departure of most assembly factors (SSU processome disassembly) a pre-40S particle emerges, which in addition to compacted rRNA domains still contains U3 snoRNA and the RNA helicase Dhr1[ref9]. The current absence of intermediate states highlights our limited understanding of the mechanisms that guide transitions between visualized states. The transition of the SSU processome from maturation to disassembly was suggested to involve the RNA exosome (with associated exonuclease Rrp6 and RNA helicase Mtr4)[10,12–15]. As the arbiter of RNA quality control, the RNA exosome ensures correct RNA processing. This system requires

synergistic interactions between the RNA exosome and its substrates for RNA

surveillance. Substrates that remain tethered to the RNA exosome are subject to RNA

turnover whereas correctly processed RNA substrates are released[16]. The limited number

of visualized SSU processome maturation and disassembly states has prevented a

mechanistic understanding of their interaction with the RNA exosome and how RNA

exosome function is coupled to SSU processome maturation and disassembly.

Specifically, the mechanisms of RNA exosome recruitment, substrate access, and

disengagement remain unclear[12,13].

At the end of SSU processome disassembly, the DEAH-box helicase Dhr1 (DHX37

in humans) plays a key role in the removal of U3 snoRNA from a pre-40*S* particle to

facilitate the formation of the central pseudoknot. However, despite ample studies[17–21],

the mechanism with which Dhr1 is initially recruited to the SSU processome, kept in an

inactive conformation, subsequently repositioned to engage its substrate U3 snoRNA,

and ultimately activated, remains unknown. Presently there is a lack of understanding of

how the SSU processome, the RNA exosome, and Dhr1 work in concert to bring about

unidirectional maturation and disassembly while maintaining RNA quality control

throughout the entire process, which enables completed particles to leave from the

nucleolus.

To address this, we determined native high-resolution structures of 16

Saccharomyces cerevisiae SSU processome states undergoing maturation and

disassembly with resolutions up to 2.7 Ångstroms. Together with complementary genetics

and an *in silico* protein-protein interaction screen, we have elucidated an RNA exosome-

mediated mechanism that explains both SSU processome maturation and disassembly

as a function of the 3′ to 5′ helicase activity of the RNA exosome-bound helicase Mtr4.

Within this framework SSU processome disassembly, exosome-mediated RNA

surveillance, and Dhr1 activation are tightly coordinated to maintain RNA quality control

during the entire pathway.

To determine how Dhr1-bound SSU processomes undergo maturation and

disassembly, we obtained native cryo-EM structures of particles containing Dhr1 during

early, intermediate, and late stages (**Fig. 1 and Extended Data Fig. 1**). Early assembly

states were selected using Dhr1 and the early assembly factor Kre33 as affinity baits

(Dataset 1, **Supplementary Fig. 1**). For unbiased views of all Dhr1-bound states, Dhr1

was used as the sole affinity bait (Dataset 2, **Supplementary Fig. 2**). The late

disassembly state containing activated Dhr1 (here State O) was obtained by selecting for

particles containing Dhr1 and its activator Utp14 and lacking Utp7, which is present in

earlier intermediates (Dataset 3, **Supplementary Fig. 3**).

From a total of 213,766 cryo-EM images, initial data curation and comprehensive

3D classifications yielded 16 SSU processome reconstructions (states A-O) with overall

resolutions of 2.7-6 Ångstroms (**Fig. 1 and Extended Fig. 1**). Focused refinements with

105 maps were used to obtain 16 composite maps for all structures (**Supplementary**

Figs. 4-25), which enabled model building and assignment of 50 unique assembly factors,

21 ribosomal proteins, and three RNAs (5′ ETS, 18S rRNA, and U3 snoRNA) across all

states (**Extended Data Table 1**). Here, states are defined based on distinct rRNA folding

transitions and the presence or absence of distinct assembly factors resulting in 15 unique

states (States A-O), and a substate of State A that undergoes conformational changes

only (State A*). Together these states provide unprecedented high-resolution coverage

of native SSU processome maturation and disassembly that goes far beyond prior

studies[9,10,12] (**Extended Data Fig. 1-6, Supplementary Fig. 26 and 27**).

Chronology of SSU biogenesis

The large number of observed states allows us to visualize nucleolar SSU biogenesis as

a function of RNA exosome-mediated 5′ ETS degradation, which first triggers cleavage

of the 18S rRNA at site A1 (States A-G), and subsequently orchestrates SSU processome

disassembly, rRNA compaction, and activation of Dhr1 (States H-O) (**Fig. 1, Extended**

Data Fig. 1 and 2). This is contrary to prior studies which lacked the temporal resolution

to precisely define both the chronology and causality of events leading to cleavage at site

A1 and hence the underlying mechanism (**Supplementary Fig. 26**)[9]. Initial degradation

of the 3′ end of the 5′ ETS on the exterior of the particle is followed by accommodation of

the C-terminus of Rrp6 (State C), suggesting that the RNA exosome is responsible for

this remodeling step. In State D the restructuring of the 5′ ETS in the interior of the SSU

processome allows for partial release of Krr1 and Faf1 from their binding sites. This leads

to the 3′ end of the 18S rRNA precursor (helix 45) to be chaperoned by the assembly

factor Pno1 towards the interior of the particle, thus allowing an N-terminal element of the

DEAH-box helicase Dhr1 to bind. As the assembly factor Utp14 connects Pno1 with Dhr1

(**Supplementary Fig. 27 e and i**), the integration of helix 45 near the central domain

results in the docking of the auto-inhibited Dhr1 in State E. Surprisingly we find that A1

cleavage in State G is solely triggered by the incorporation of Dim1 as this protein couples

the interrogation of helix 45 installation with the stabilization of U3 snoRNA-bound regions

within the 18S rRNA in an upward conformation. This facilitates Utp24-mediated A1

cleavage in particles that still contain the Kre33 module, which was previously thought to

precede A1 cleavage[9] (**Fig. 1, Extended Data Fig. 3**).

141   SSU processome disassembly further progresses as a function of 3′ to 5′

142   elimination of the 5′ ETS, resulting in compositional and conformational changes of the

143   particle (**Fig.1**). During these compositional changes, Utp14, a largely unstructured

144   protein containing several regulatory segments visualized in our structures (here referred

145   to as a-f) can adopt different binding conformations as a response to SSU processome

146   maturation and disassembly (**Extended Data Fig.6 a,d,e**).

147   With the processing of the 5′ hinge, distinct stages of disassembly are observed

that involve the stepwise departure of assembly factors, such as Rrp5, Sof1, and Utp7

(States H-J). In States J and K, Utp14 licenses a new binding site for Dhr1, which was

previously occupied by Utp7. The subsequent processing of the 3′ hinge (States K-L)

releases much of the UtpB complex (except for Utp18) and the helicase Dhr1. In a major

compositional remodeling step, the transition from State L to M marks the complete

elimination of the 5′ ETS, which triggers the departure of the UtpA and UtpB complexes

as well as individual assembly factors and Rrp6. This remodeling step subsequently

allows Dhr1 to be repositioned and activated by Utp14 in a state-specific manner. States

M-O therefore capture the activation of Dhr1 as further compaction of the particle triggers

the undocking of U3 snoRNP proteins and Utp20. These results suggest a direct

mechanism by which the RNA exosome could degrade the 5′ ETS to trigger large

compositional and conformational changes that transform the SSU processome into a

pre-40S particle.

Integrated nucleolar RNA surveillance

Genetic studies suggested that the yeast RNA exosome binds the SSU processome via an interaction between the RNA helicase Mtr4 arch domain and the assembly factor Utp18 via an arch-interacting motif (AIM)[14] . While low-resolution density for the RNA exosome in the proximity of the SSU processome has been observed[10,12], exactly how exosome recruitment and positioning is achieved across multiple states remains unclear. Moreover, the RNA exosome must efficiently detect and eliminate trapped SSU precursors within the nucleolus which have failed to complete biogenesis, yet how nucleolar RNA surveillance is set up to do so remains unresolved[10,12,14] . The 71 proteins present in the SSU processome together with the 14 proteins present within the RNA exosome provide a unique opportunity to screen pathway-wide protein-protein interactions *in silico* using Alphafold-Multimer[22–24]. We employed this approach to obtain an SSU processome predictome to test 3,570 unique binary interactions, which together with our high-resolution cryo-EM maps enabled the identification of many previously unknown interactions (**Fig. 2a, Extended Data Fig. 4, and Extended Data Table 2**). These data highlight key features of the SSU biogenesis pathway including the architectural roles of Sas10-domain containing complexes, tethering functions of Mpp10 for ribosomal protein S5 and Dim1, and predictions for the recruitment and activation of Dhr1/DHX37 (**Extended Data Fig. 4d,f**). In addition, mutually exclusive interactions between Kre33 and Faf1 or Bms1 are revealed, which now explain previously unassigned density for the human homologs of Kre33/Faf1 (NAT10/C1ORF13) in the human SSU processome[11] (**Extended Data Fig. 4e**).

A striking series of predicted interactions emerged from the SSU processome

predictome that suggest extensive contacts between the RNA degradation machinery and

the maturing and disassembling SSU processome (**Fig. 2a,b**). In addition to previously

known interactions between the RNA helicase Mtr4 and Utp18[ref14] and newly identified

multivalent interactions between the yeast Rrp6 C-terminus (lasso) and the SSU

processome in this study, the SSU processome predictome suggested that Utp14, the

essential activator of Dhr1, additionally serves as a bridge between Mtr4 (via Exosome

Recruitment Motif 1, ERM1), the core exosome component Rrp40 (via ERM2), and the

SSU processome (**Fig. 2a,b**). This mode of binding is similar to the binding of the RNA

exosome cofactor Mpp6[ref25], with Mpp6 motifs having a high degree of sequence similarity

to Utp14 ERM1 and ERM2 (**Extended Data Fig. 5a**).

Based on these predictions, a model emerges in which the RNA exosome contains

four contact points for the SSU processome, which in addition to Utp18 AIM and the C-

terminus of Rrp6 (lasso) include two peptides of Utp14 (ERM1 and ERM2) (**Fig. 2b**).

Synergistically, the 2.7 Ångstrom structure of State H highlights that elements of Rrp6,

Utp18 (AIM), and predicted Utp14 (ERM1 and ERM2) are all positioned in close proximity

to the 3′ end of the 5′ ETS where the Mtr4 helicase needs to unwind the 5′ETS for both

maturation and disassembly to occur (**Fig. 2c**). In State H, Utp14 occupies a strategic

position with key regulatory elements that respond to conformational changes of the entire

particle. Elements of Utp14 that will in later states provide a part of a Dhr1 binding site

(segment b) or activate Dhr1 (segments d and e) are sequestered in State H near

Sof1/Utp20 and Utp7/Utp1 respectively (**Fig. 2c,d).** Separately, the newly identified Utp14

hook (segment c) recruits the auto-inhibited Dhr1 and this inactive state is further

stabilized by N-terminal elements of Pno1 (**Supplementary Fig. 27i,l**). This arrangement

places ERM1 and ERM2 of Utp14 directly between segments b and c so that after

cleavage at site A1 all elements that bridge the SSU processome and the RNA exosome

are present near the 3′ end of the 5′ ETS (**Fig. 2c,d**).

The network consisting of Utp18, Rrp6 and Utp14 suggests that beyond exosome

catalytic function, connective redundancy may play a major role in ensuring correct

placement of the RNA exosome near the SSU processome. While prior data have

indicated that Utp18 AIM mutants are synthetically lethal in the absence of Rrp6[ref12,14],

the relative contributions of Rrp6 exonuclease activity and its ability to bind to the SSU

processome have remained unclear. Using yeast genetics we show that in the absence

of the Utp18 AIM, an inactive Rrp6 exonuclease (Y361A)[26] results in synthetic lethality,

suggesting that catalytic activity of Rrp6 becomes essential if Mtr4 helicase cannot be

correctly positioned (**Fig. 2e**). These data suggest that both positioning and the

redundancy in connections of the RNA exosome to the SSU processome is essential for

its function. Without the Utp18 AIM, three connections to the SSU processome remain

via the Rrp6 lasso and the predicted Utp14 ERM1 & ERM2 (**Fig. 2a**). Our genetic data

indicate that as long as any one of these connections is present, cell viability is

maintained. By contrast, synthetic lethality is observed if all four connections are disrupted

(**Fig. 2e**). Since the Rrp6 lasso is a multivalent binder of the SSU processome with four

binding sites, we could separately show that iterative Rrp6 C-terminal truncations in a

genetic background lacking the Utp18 and Utp14 connections gradually reduce growth

(**Extended Data Fig. 5**). The SSU processome predictome together with structural data

and genetics reveal a highly redundant nucleolar tethering network that couples SSU

processome maturation, disassembly and RNA quality control. These findings are further

consistent with the visualization of the human homolog of Rrp6 (EXOSC10) in the

immediate vicinity of nucleolar SSU precursors[27] .

Synchronized exosome-mediated disassembly

Elucidating the precise molecular mechanisms governing the controlled disassembly of

the SSU processome remains a significant challenge, as current models lack the

necessary sampling and resolution to mechanistically explain this process[9]. Here, States

H-O capture a series of intermediates that follow RNA exosome-mediated disassembly

(**Figs. 1 and 3**), allowing us to mechanistically explain how disassembly and RNA quality

control are coupled. The irreversible 3′ to 5′ elimination of the 5′ ETS by the RNA exosome

is directly coupled to the release of assembly factors in the exosome's path. These

include, among others, the Sof1-Utp7 complex at the 5′ hinge, the UtpB complex at the

3′ hinge, and finally Utp18 and the UtpA complex positioned at the start of the 5′ ETS

(**Fig. 3a**). RNA exosome-mediated compositional changes to the SSU processome result

in three key events that occur concurrently. First, the reduction of multi-valent RNA-

protein and protein-protein interactions within the SSU processome leads to release of

bound and tethered complexes. Second, the RNA exosome loses two of its interaction

sites with the SSU processome and its function transitions from substrate engagement to

substrate monitoring. Third, the changes of the SSU processome landscape provide new

binding surfaces for Utp14, which organizes both Dhr1 repositioning and activation (**Fig.**

3).

Beginning in State H, RNA exosome mediated processing of the 3′ end of the 5′

ETS alters the Sof1 binding site, weakening its interaction with Utp7 until only the C-

terminus of Utp7 remains bound in State I. During this transition, key regulatory elements

of Utp14 involved in subsequent Dhr1 recruitment (segment b) and activation (segments

d and e) are released while remaining tethered via peptides bound to Dhr1 and Pno1

(segments c and f respectively). During subsequent 5′ ETS degradation, represented in

states J and K, the Utp7 C-terminus is released, and its binding site is gradually occupied

by elements of Utp14 (segments a and b) that prepare a binding site for Dhr1 as the UtpC

complex is released. During the transition from State K to L, continued 5′ ETS processing

results in UtpB losing its primary binding site at the 3′ hinge while remaining tethered to

the SSU processome via Utp18, and Dhr1 remains connected to the particle via its N-

terminus and the Utp14 hook (segment c) (**Fig. 3b**). Following complete 5′ ETS removal

during the transition from State L to State M, UtpA, associated proteins, Utp18, and Rrp6

are fully released (**Fig. 3, Extended Data Fig. 6a**). During this key transition the roles of

the RNA exosome and Dhr1 are dramatically changed.

The role of the RNA exosome shifts from the active substrate engaged state with

four binding sites on the SSU processome to an overall surveillance state through the

remaining two Utp14 ERM sites (**Extended Data Fig. 6a**). Furthermore, re-engagement

of the substrate by the RNA exosome is prevented by the lack of a Utp18 binding site and

the occlusion of the Rrp6 lasso binding site by the C-terminus of ribosomal protein S4

(**Fig. 3b, Extended Data Fig. 6b and c**). Separately, the role of Dhr1 changes from a

tethered auto-inhibited state to a repositioned, substrate engaged, and activated enzyme

poised to remove U3 snoRNA from the pre-40*S* particle.

Molecular circuitry of Dhr1 activation

High-resolution crystal structures of Dhr1/DHX37 have shown that this DEAH-box helicase in isolation can exist in two states, an inactive state in which an auto-inhibitory loop prevents substrate binding, and an active state, in which single-stranded RNA acts as a substrate[11,19]. However, thus far no Utp14-mediated active state of Dhr1 has been observed. The comparison between the inactive state of Dhr1 in State H and the Utp14-mediated active state of Dhr1 in State O now rationalizes how irreversible RNA exosome-mediated 5′ ETS degradation orchestrates a cascade of events that triggers Dhr1 activation in a state-specific manner during SSU processome disassembly (**Fig. 4**).

In State H, the UtpB tetramer provides binding sites for both Dhr1 and Pno1 (elements 1 and 4), which enable additional elements of Pno1 (elements 2 and 3) to stabilize the auto-inhibited state of Dhr1. Element 3 of Pno1 stabilizes the auto-inhibitory loop of Dhr1, while element 2 of Pno1 prevents binding of segment d of Utp14, thus preventing a switch towards Dhr1 activation (**Fig. 4a**). Separately, segments of Utp14 that subsequently provide a binding site (segment b) or activate Dhr1 (segments d and e) are sequestered by Utp20/Sof1 and Utp7-Utp1 respectively (**Figs. 2c, 4a, c**).

In State O, the prior RNA exosome-mediated elimination of the UtpB complex, and the release of Utp14 segments a-e have dramatically changed the local environment for Dhr1. The new Dhr1 binding site (Utp14 segments a and b) is formed on top of its substrate U3 snoRNA, which still base-pairs with 18*S* elements that will form the central pseudoknot upon Dhr1-mediated U3 snoRNA release (**Fig. 4b**). The relative positions of Utp14 elements c and f now enable Utp14 elements d and e to stabilize a conformation of Dhr1 in which the RecA lobes are dramatically shifted to engage the substrate U3

snoRNA while displacing the auto-inhibitory loop (**Fig. 4b, d**). The architecture of the

Utp14-mediated active form of Dhr1 in State O rationalizes a wealth of genetic and

biochemical data from yeast and mammalian systems, highlighting that this mode of

activation is distinct from G-patch mediated DEAH-box helicase activation (**Extended**

Data Fig. 6e-g)[19,21].

Thus, RNA exosome-mediated remodeling of the SSU processome triggers a

series of compositional and conformational changes within the disassembling SSU

processome that collectively switch Dhr1 from an "OFF" into an "ON" state that enables

its engagement and processing of U3 snoRNA. The irreversible release of U3 snoRNA

and the formation of the central pseudoknot prepare the resulting pre-40S particle for

subsequent nuclear maturation where Utp14 is replaced by Nob1[ref2], releasing the final

link to nucleolar RNA surveillance.

Conclusions

Nuclear bodies that contain biosynthetic processing pathways such as the nucleolus,

Cajal bodies, and nuclear speckles share unified organizational principles by which RNA

synthesis and processing results in the formation and dynamics of these multi-valent

condensates[28,29]. For nucleolar SSU processome maturation and disassembly, we

present a high-resolution temporal perspective, highlighting that this highly dynamic

process is driven by the activities of two RNA helicases associated with the RNA

degradation machinery (Mtr4) and the SSU processome itself (Dhr1) (**Figs. 1 and 2**). The

irreversible unwinding and degradation of pre-ribosomal RNA (5′ ETS) changes the

composition of disassembling SSU processomes, allowing regulatory peptides of Utp14

to probe the changing landscape of these particles to state-specifically reposition and

activate Dhr1 for the terminal and irreversible release of U3 snoRNA (**Figs. 3 and 4**). The

emerging model now highlights the importance of multivalent tethering at three levels.

First, intra-SSU processome tethering enables pre-recruitment of assembly factors

required for subsequent steps including restructuring of the SSU processome and A1

cleavage. Second, the extent (valency) of tethering between large particles such as the

RNA exosome to the SSU processome changes as the exosome matures and

disassembles the SSU processome, thereby shifting the role of the RNA exosome from

initial recognition to substrate engagement and finally substrate surveillance. Third, at the

level of the organelle, overall tethering of the maturing and disassembling SSU

processomes to the nucleolus is decreased as more and more nucleolar assembly factors

are released, thereby reducing the overall valency of particles before nucleolar release

(**Fig. 5**). The scope of our study now allows us to illustrate molecular events at the

Ångstrom scale that can rationalize the organizational principles at the micrometer scale

as observed for the nucleolus[27]. This study highlights molecular transitions in large

dynamic RNA-protein complexes driven by RNA helicases, and how these changes may

be coupled to movement through biomolecular condensates in nuclear bodies.

REFERENCES

1. Dragon, F. *et al.* A large nucleolar U3 ribonucleoprotein required for 18S ribosomal RNA biogenesis. *Nature* **417**, 967–970 (2002).

2. Dörner, K., Ruggeri, C., Zemp, I. & Kutay, U. Ribosome biogenesis factors—from names to functions. *Embo J* e112699 (2023) doi:10.15252/embj.2022112699.

3. Broeck, A., Vanden & Klinge, S. Eukaryotic Ribosome Assembly. *Annual Review of Biochemistry* **Volume 93 (In press)**, (2024).

4. Kornprobst, M. *et al.* Architecture of the 90S Pre-ribosome: A Structural View on the Birth of the Eukaryotic Ribosome. *Cell* **166**, 380–393 (2016).

5. Chaker-Margot, M., Barandun, J., Hunziker, M. & Klinge, S. Architecture of the yeast small subunit processome. *Science* **355**, eaal1880 (2017).

6. Sun, Q. *et al.* Molecular architecture of the 90S small subunit pre-ribosome. *Elife* **6**, e22086 (2017).

7. Barandun, J. *et al.* The complete structure of the small-subunit processome. *Nat. Struct. Mol. Biol.* **24**, 944–953 (2017).

8. Cheng, J., Kellner, N., Berninghausen, O., Hurt, E. & Beckmann, R. 3.2-Å-resolution structure of the 90S preribosome before A1 pre-rRNA cleavage. *Nat. Struct. Mol. Biol.* **24**, 954–964 (2017).

9. Cheng, J. *et al.* 90S pre-ribosome transformation into the primordial 40S subunit. *Science* **369**, 1470–1476 (2020).

10. Du, Y. *et al.* Cryo-EM structure of 90S small ribosomal subunit precursors in transition states. *Science* **369**, 1477–1481 (2020).

11. Singh, S., Broeck, A. V., Miller, L., Chaker-Margot, M. & Klinge, S. Nucleolar maturation of the human small subunit processome. *Science* **373**, eabj5338 (2021).

12. Lau, B. *et al.* Structure of the Maturing 90S Pre-ribosome in Association with the RNA Exosome. *Mol. Cell* **81**, 293-303.e4 (2021).

13. Broeck, A. V. & Klinge, S. An emerging mechanism for the maturation of the Small Subunit Processome. *Current Opinion in Structural Biology* **73**, 102331 (2022).

14. Thoms, M. *et al.* The Exosome Is Recruited to RNA Substrates through Specific Adaptor Proteins. *Cell* **162**, 1029–1038 (2015).

15. Hunziker, M. *et al.* Conformational switches control early maturation of the
eukaryotic small ribosomal subunit. *Elife* **8**, 213 (2019).

16. Keidel, A., Long, C. L., Iwasa, J. & Conti, E. RNA-Degrading Exosome Complexes:
Molecular Mechanisms and Structural Insights. *Annu. Rev. Cell Dev. Biol.* (2025)
doi:10.1146/annurev-cellbio-111822-115115.

17. Sardana, R. *et al.* The DEAH-box helicase Dhr1 dissociates U3 from the pre-rRNA
to promote formation of the central pseudoknot. *PLoS Biol* **13**, e1002083 (2015).

18. Zhu, J., Liu, X., Anjos, M., Correll, C. C. & Johnson, A. W. Utp14 recruits and
activates the RNA helicase Dhr1 to undock U3 snoRNA from the pre-ribosome.
Molecular and Cellular Biology **36**, MCB.00773-15 (2016).

19. Boneberg, F. M. *et al.* Molecular mechanism of the RNA helicase DHX37 and its
activation by UTP14A in ribosome biogenesis. *RNA* **25**, 685–701 (2019).

20. Choudhury, P., Hackert, P., Memet, I., Sloan, K. E. & Bohnsack, M. T. The human
RNA helicase DHX37 is required for release of the U3 snoRNP from pre-ribosomal
particles. *RNA Biol* **16**, 54–68 (2018).

21. Lin, R., Elmir, E., Reynolds, M. J. & Johnson, A. W. In vitro characterization of the
yeast DEAH/RHA RNA helicase Dhr1. *J. Biol. Chem.* **301**, 108366 (2025).

22. Evans, R. *et al.* Protein complex prediction with AlphaFold-Multimer. *bioRxiv*
2021.10.04.463034 (2021) doi:10.1101/2021.10.04.463034.

23. Yu, D., Chojnowski, G., Rosenthal, M. & Kosinski, J. AlphaPulldown—a python
package for protein–protein interaction screens using AlphaFold-Multimer.
Bioinformatics **39**, btac749 (2022).

24. Schmid, E. W. & Walter, J. C. Predictomes, a classifier-curated database of
AlphaFold-modeled protein-protein interactions. *Mol. Cell* **85**, 1216-1232.e5 (2025).

25. Wasmuth, E. V., Zinder, J. C., Zattas, D., Das, M. & Lima, C. D. Structure and
reconstitution of yeast Mpp6-nuclear exosome complexes reveals that Mpp6 stimulates
RNA decay and recruits the Mtr4 helicase. *eLife* **6**, e29062 (2017).

26. Assenholt, J. *et al.* Exonucleolysis is required for nuclear mRNA quality control in
yeast THO mutants. *RNA (N. York, NY)* **14**, 2305–13 (2008).

27. Quinodoz, S. *et al.* Mapping and engineering RNA-controlled architecture of the
multiphase nucleolus. *bioRxiv* 2024.09.28.615444 (2024)
doi:10.1101/2024.09.28.615444.

28. Quinodoz, S. A. & Guttman, M. Essential Roles for RNA in Shaping Nuclear
Organization. *Cold Spring Harb. Perspect. Biol.* **14**, a039719 (2021).

29. Shan, L., Li, P., Yu, H. & Chen, L.-L. Emerging roles of nuclear bodies in genome
spatial organization. *Trends Cell Biol.* **34**, 595–605 (2024).

30. Mastronarde, D. N. Automated electron microscope tomography using robust
prediction of specimen movements. *J Struct Biol* **152**, 36–51 (2005).

31. Kimanius, D., Dong, L., Sharov, G., Nakane, T. & Scheres, S. H. W. New tools for
automated cryo-EM single-particle analysis in RELION-4.0. *Biochem J* **478**, 4169–4185
(2021).

32. Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms
for rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290–296
(2017).

33. Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for
improved cryo-electron microscopy. *Nat Methods* **14**, 331–332 (2017).

34. Zhang, K. Gctf: Real-time CTF determination and correction. *J Struct Biol* **193**, 1–12
(2016).

35. Asarnow, D., Cheng, D. A. E. P. Y. & Cheng, Y. UCSF pyem v0.5. Zenodo. (2019)
doi:10.5281/zenodo.3576630.

36. Punjani, A. & Fleet, D. J. 3D variability analysis: Resolving continuous flexibility and
discrete heterogeneity from single particle cryo-EM. *J. Struct. Biol.* **213**, 107702 (2021).

37. Wagner, T. *et al.* SPHIRE-crYOLO is a fast and accurate fully automated particle
picker for cryo-EM. *Commun Biol* **2**, 218–13 (2019).

38. Pettersen, E. F. *et al.* UCSF ChimeraX : Structure visualization for researchers,
educators, and developers. *Protein Sci* **30**, 70–82 (2020).

39. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature*
596, 583–589 (2021).

40. Casañal, A., Lohkamp, B. & Emsley, P. Current Developments in Coot for
Macromolecular Model Building of Electron Cryo-microscopy and Crystallographic Data.
Protein Sci. pro.3791 (2019) doi:10.1002/pro.3791.

41. Afonine, P. V. *et al.* Real-space refinement in PHENIX for cryo-EM and
crystallography. *Acta Crystallogr D Struct Biol* **74**, 531–544 (2018).

42. Black, J. J., Sardana, R., Elmir, E. W. & Johnson, A. W. Bud23 promotes the final
disassembly of the small subunit Processome in Saccharomyces cerevisiae. *PLoS*
Genet. **16**, e1009215 (2020).

Fig. 1. SSU processome maturation and disassembly pathway.

Simplified atomic models of the SSU processome assembly pathway with color-coded
rRNAs and ribosome assembly factors. Major changes in the rRNA between states are
in the foreground and changes in protein compositions as well as functions are noted.

449 Elements of the 5′ ETS that will be removed are highlighted in orange. Proteins that

450 become ordered or depart between different assembly intermediates are indicated. The

451 overall resolution of each cryo-EM reconstruction is noted below each state. A dashed

452 arrow indicates tethering of protein components.

Fig. 2. Interaction networks between SSU processome and the RNA exosome.

(**a**) Yeast SSU processome predictome containing assembly factors (AFs), ribosomal proteins (RPs) and RNA exosome proteins (EXO) with color-coded confidence scores. Three AlphaFold-Multimer predictions are highlighted with color-coded proteins and

confidence levels (pLDDT). (**b**) AlphaFold predictions for color-coded Utp14, Mtr4, and

Utp18 AIM mapped onto a composite model of the RNA exosome with Rrp6 (PDBs 5vzj

and 6d6q). (**c**) Cartoon representation of state H (left) and zoomed-in view (right) of the

5′ ETS 3′ end exit site (orange circles) with color-coded exosome interacting proteins.

Regulatory elements of Utp14 are highlighted in orange. (**d**) Schematic of Utp14, Rrp6,

Utp18, and Pno1 protein sequences in State H with interacting and functional elements

labeled. (**e**) Growth analysis of Rrp6 and Utp14 mutants in a Utp18 multi-Ala background

on 5-FOA plates.

Fig. 3. Helicase-mediated disassembly is coupled to RNA quality control.

(**a**) Schematic of SSU processome disassembly as a function RNA degradation. RNA and protein elements bound to the 5′ ETS RNA are displayed as colored boxes and states with corresponding 5′ ETS length are listed below. (**b**) Architecture of the changes around the 5′ ETS exit sites (orange circles) from State H to M. Protein elements are color-coded and elements of Utp14 (b-f), Pno1 (1-4) and the state of Dhr1 are highlighted.

Fig. 4. Mechanism of Utp14-mediated Dhr1 activation.

(**a**) Zoomed-in view of inactive Dhr1 in State H, with auto-inhibitory loop (light pink) and inhibitory elements of Pno1 (1-4) and elements of Utp14 highlighted. The arrow indicates the conformational change of Dhr1 RecA domains. (**b**) Zoomed-in view of activated Dhr1 in State O. Activating Utp14 elements (a, b, d, and e) are highlighted in orange. (**c**) Schematic of State H molecular circuitry between Utp14, Dhr1 and Pno1. Interacting elements are colored grey and inhibitory interactions are indicated with red lines. (**d**) Schematic of State O molecular circuitry between Utp14, Dhr1 and Pno1. Interacting elements are colored grey and activating interactions are indicated with green lines.

Fig. 5. Model of the mechanism of SSU biogenesis in the nucleolus.

(**a**) Schematic model of yeast SSU biogenesis in the nucleolus. Decreasing valency of

maturing particles is indicated below. (**b**) SSU processome maturation and disassembly

513   pathway with key transitions are indicated. Color-coded tethered proteins and complexes

514   are indicated with linkers. Arrows and numbers highlight the order of transition steps (RNA

515   degradation, conformational changes, protein release, or protein repositioning to new

516   binding sites) that lead to particles with low valency during disassembly.

METHODS

Generation of endogenously tagged DHR1, KRE33, UTP7 and UTP14 strains

Assembly intermediates of the *S. cerevisiae* SSU processomes were purified from

three genetically modified BY4741 (MATa his3Δ leu2Δ0 met15Δ0 ura3Δ0) strains. Where

genes of interest (DHR1, KRE33, UTP7 and UTP14) were endogenously tagged: the first

strain containing C-terminal tandem 3C protease-cleavable mCherry and TEV-cleavable

alfa peptide tag on DHR1 along with a 3C cleavable C-terminal GFP-tag on KRE33 (MATa

his3Δ leu2Δ0 met15Δ0 ura3Δ0 DHR1-linker-tev-alfa tag-3c-mCherry under Hygromycin

B selection and KRE33-linker-tev-GFP under Nourseothricin selection), the second strain

containing only C-terminal tandem 3C protease-cleavable mCherry and TEV-cleavable

alfa peptide tag on DHR1 (MATa his3Δ leu2Δ0 met15Δ0 ura3Δ0 DHR1-linker-tev-alfa

tag-3c-mCherry under Hygromycin B selection) and the third strain containing C-terminal

tandem 3C protease-cleavable mCherry and a TEV-cleavable alfa peptide tag on DHR1

along with TEV cleavable C-terminal GFP-tag on UTP7 and C-terminally streptavidin-

binding-peptide (sbp) tag on UTP14 (MATa his3Δ leu2Δ0 met15Δ0 ura3Δ0 DHR1-linker-

tev-alfa tag-3c-mCherry under Hygromycin B selection and UTP7-linker-tev-GFP under

G418 selection and UTP14-linker-SBP under Nourseothricin selection). The three strains

were generated with standard genomic tagging techniques using the following primers for

yeast genomic tagging:

DHR1 forward primer:

TCACCAGAAAGGGCTTCCAGACCATCACAGGTGAAGAGAAAGAAAAAAAACGCT

GCAGGTCGACGGATCC

DHR1 reverse primer:

TTAAGTGGTTGCAATTATTTGATGCCCTAGATAGGAATAATGTATTCTTCGGCAGAT

CCGCGGCCGCATAGG

Kre33 forward primer:

AAGAGATGAAAGCTATGAAAAACCAAGAAAGTCTAAAAAGGCTGCAAATACGCTG

CAGGTCGACGGATCC

KRE33 reverse primer :

TGTAAAGGTTCAAACATCAACTATGTTTCTATTCTATATTATTGTACAAAGGCAGATC

CGCGGCCGCATAGG

UTP7 forward primer:

TTTCAGAAGACCACAAGGATGTCATCGAAGAGGCATTGAGCAGATTCGGCACGCT

GCAGGTCGACGGATCC

UTP7 reverse primer:

GCTATATTAATATGCAATCGATTCTCATACTGTCAACTTTTTGAACATGAGGCAGAT

CCGCGGCCGCATAGG

UTP14 forward primer:

TCATGACTAAGCCAGGCCAAGTTATTGATCCTTTGAAGGCACCATTTAAGACGCTG

CAGGTCGACGGATCC

UTP14 reverse primer:

ATTATTCCAGTATTACTATTCTACACAATGCATAATAAATAGATATAAAAGGCAGATC

CGCGGCCGCATAGG

The three strains were grown at 30 °C in YPD media (1% yeast extract, 2%

peptone and 2% glucose) to optical density of OD 1 measured at 600 nm. Cells were

harvested at 4,000g for 5 min, washed once with 1 l of ice cold ddH$_2$O and once with a

volume of ddH$_2$O supplemented with protease inhibitors (E64, pepstatin and PMSF) equal

to the mass of the cell pellet. The final pellets of ~30-40g were flash frozen in liquid

nitrogen. Pellets were lysed by cryo-grinding using a Retsch Planetary Ball Mill PM100

and powder was stored at –80 °C.

Purification of SSU processome intermediates

Cryo-ground powder was resuspended with buffer 1 (60mM Tris pH7.5, 50mM

NaCl, 2mM MgCl2, 5% glycerol, 0.1% NP-40) with addition of PMSF, pepstatin and E64

protease inhibitors. The suspension was cleared by centrifugation at 4 °C and 40,000g

for 20 min and lysate was incubated with 800 µl of packed NHS–sepharose beads

(Cytiva) coupled to anti-mCherry nanobodies for the first capture for all purifications and

incubated for 3.5 h at 4 °C on a nutator. Beads were pelleted by centrifugation at 4 °C for

1 min at 127g. After five washes in buffer 1, complexes were eluted in buffer EB (60mM

Tris, 50mM NaCl, 5mM MgCl2, 2% glycerol, 0.01% NP-40) supplemented with 3C

protease for 1 h at 4 °C. Beads were pelleted by centrifugation at 4 °C for 1 min at 127g

and supernatant was eluted and incubated with 80 µl of packed NHS–sepharose beads

(Cytiva) coupled to anti-GFP nanobodies for second capture (Dhr1/Kre33 and Dhr1/Utp14

purifications) while anti-alfa nanobodies were used for a second Dhr1 (Dhr1 only). Beads

were incubated for 1 h at 4 °C on a nutator and pelleted by centrifugation at 4 °C for 1 min

at 127g and washed twice with 2 ml of Buffer EB. After the second wash, for Dhr1/Kre33

and Dhr1 only purifications, beads were pelleted and resuspended with 25 µl of Buffer EB

supplemented with 1mM DTT and TEV protease and incubated on ice for 1 h. Beads

were pelleted by centrifugation at 4 °C for 10 min at 21,130g and the supernatant was

collected. For Dhr1/Utp14 purification, after the second capture (here Utp7), beads were

pelleted by centrifugation at 4 °C for 1 min at 127g and flowthrough was collected, the rest

of the beads were discarded (this step was used to remove earlier contaminating

complexes). The flowthrough was incubated with 40 µl of packed NHS–sepharose beads

(Cytiva) coupled to streptavidin for 1 h at 4 °C on a nutator. Beads were pelleted by

centrifugation at 4 °C for 1 min at 127g and washed twice with 2 ml of Buffer EB. After the

second wash, beads were pelleted and resuspended with 40 µl of Buffer EB

supplemented with 1mM DTT and 5 mM of D-biotin (Amresco) and incubated on ice for

20 min. Beads were pelleted by centrifugation at 4 °C for 10 min at 21,130g and the

supernatant was collected.

Cryo-EM grid preparation and data acquisition

Cryo-EM grids were prepared using a Vitrobot Mark IV robot (FEI Company) set

to 90% humidity and 18 °C temperature. Three and a half microliters of the eluted solution

was applied to a glow-discharged Quantifoil Au R3.5/1 with a layer of 2-nm ultrathin

carbon (LFH7100AR35, Electron Microscopy Sciences). After 2.5 min incubation inside

the Vitrobot chamber, the excess solution was manually blotted and a fresh sample of 3.5

µl was reapplied inside the Vitrobot chamber and incubated for another 2.5 min. For the

Dhr1 and Kre33 dataset, a total of five applications were done on each grid. The grid was

then blotted (blot force of 8 and blot time of 9 s) and plunged into liquid ethane. For the

Dhr1dataset, a total of two applications were done for each grid. The grid was then blotted

(blot force of 8 and blot time of 7 s) and plunged into liquid ethane and for the Dhr1 and

Utp14 dataset, a total of five applications were done on each grid to achieve a good

distribution of particles. The grid was then blotted (blot force of 8 and blot time of 9 s) and

plunged into liquid ethane. Grids were imaged on a Titan Krios electron microscope (FEI)

with an energy filter (slit width of 20 eV) and a K3 Summit detector (Gatan) operating at

300 kV with a nominal magnification of ×64,000.

SerialEM[30] was used to collect four datasets. 44,272 micrographs were collected

for the Kre33/Dhr1 dataset, two datasets of Dhr1 were collected totaling to 84,463

micrographs and 85,031 micrographs were collected for Dhr1 and Utp14 dataset. All

datasets were collected with a defocus range of −1 to −2.5 μm and a super-resolution

pixel size of 0.54 Å. Micrographs contained 40 frames using a total dose of 25.3-

30.8 e−per pixel per second (specimen pixel size of 1.08 Å per pixel) with an exposure

time of 2-2.5 s and a total dose of 61.7 -63.1e− Å−2. A multi-shot strategy was used to

record nine micrographs per hole at each stage position with the same defocus range,

electron dose and frame count.

Cryo-EM data processing Dhr1 and Kre33 dataset

The Dhr1 and Kre33 dataset was processed using a combination of RELION

5beta[31] and cryoSparc v4.6[ref32]. 44,272 movies were gain corrected, dose weighted,

aligned, with each dataset having different optic groups and binned to a pixel size of 1.08

Å using RELION's implementation of a MotionCor2-like algorithm[33]. Micrograph defocus

was estimated using Gctf[34] . Particles were picked using Laplacian (autopick) in RELION

5beta and post 2D classification a total of 2,817,609 particles were re-extracted at a pixel

size of 4.32 Å per pixel (4× binning) and underwent 3D classification in RELION 5beta

with alignment using a reference map from previous datasets. Two good classes were

combined, and duplicates were removed to result in 308,351 total particles. These

particles were subjected to three rounds of CTF refinement and Bayesian polishing in

RELION 5beta. Post polishing all homogenous refinements were completed in CryoSparc

v4.6 and classifications were all done in RELION 5beta, particle positions from cryosparc

are converted into RELION using pyem software csparc2star.py[35] . The polished particles

were subjected to a homogenous refinements resulting in a reconstruction at a global

resolution of 3.3 Å. To separate the states present in the consensus reconstruction,

multiple 3D classifications without alignment were performed. This was followed by 3D

variability[36] in CryoSparc v4.6 for analysis of the type of heterogeneity present in the data

which guided the creation of a mask around the region of interest and further 3D

classification without alignment on the region of variability. Eight distinct states (sates A-

G) were isolated from the dataset that showed unique features in the progression of

maturation of the SSU processome pathway (**Supplementary Fig.1**). The global

resolution of the eight states ranged from 3-5.9 Å.

Cryo-EM data processing Dhr1 dataset

Dhr1 dataset was processed using a combination of RELION 5beta[31] and

cryoSparc v4.6[ref32] . 84,463 movies were gain corrected, dose weighted, aligned, with

each dataset having different optic groups and binned to a pixel size of 1.08 Å using

RELION's implementation of a MotionCor2-like algorithm[33] . Micrograph defocus was

estimated using Gctf[34] . Particles were picked using crYOLO 1.7.5[37] and post 2D

classification a total of 13,198,550 particles were re-extracted at a pixel size of 4.32 Å per

pixel (4× binning) and underwent heterogenous refinement in CryoSparc v4.6 using a

reference map from previous classification in RELION 5beta. One good class was

isolated resulting in 1,933,969 total particles. These particles were subjected to three

rounds of CTF refinement and Bayesian polishing in RELION 5beta. Post polishing all

homogenous refinements were completed in CryoSparc v4.6 and classifications were all

done in RELION 5beta. Particle positions from cryosparc are converted into RELION

using pyem software csparc2star.py[35] . The polished particles were subjected to a

homogenous refinement resulting in a reconstruction at a global resolution of 3 Å. To

separate the states present in the consensus reconstruction, multiple iterations of 3D

classifications without alignment was performed. Followed by 3D variability[36] in

CryoSparc v4.6 for analysis of the type of heterogeneity present in the data which guided

the creation of a mask around the region of interest and further 3D classification without

alignment on the region of variability. Seven distinct states (sates H-N) were isolated from

the dataset that showed unique features in the progression of the disassembly of the SSU

processome pathway (**Supplementary Fig.2**). The global resolution of the seven states

ranged from 2.65-4.3 Å.

Cryo-EM data processing Dhr1 and Utp14 dataset

The Dhr1 and Utp14 dataset was processed using a combination of RELION

5beta[31] and cryoSparc v4.6[ref32] . 85,031 movies were gain corrected, dose weighted,

aligned, with each dataset having different optic groups and binned to a pixel size of 1.08

Å using CryoSparc v 4.6 motion correction. Micrograph defocus was estimated using

Patch CTF and particles were picked using a template picker resulting in a total of

27,069,273 particles which underwent heterogenous refinement and subsequent global

and local CTF refinements, followed by reference motion correction. Post polishing all

homogenous refinements were completed in CryoSparc v4.6 and classifications were all

done in RELION 5beta. Particle positions from cryosparc are converted into RELION

using pyem software csparc2star.py[35] . The polished particles were subjected to a

homogenous refinement resulting in a reconstruction at a global resolution of 2.9 Å. To

separate the states present in the consensus reconstruction, multiple iterations of 3D

classifications without alignment were performed. Followed by 3D variability[36] in

CryoSparc v4.6 for analysis of the type of heterogeneity present in the data which guided

the creation of a mask around the region of interest and further 3D classification without

alignment on the region of variability. State O was isolated from the dataset with global

resolution of 3.25 Å that showed clear density for Utp14-bound Dhr1 (**Supplementary**

Fig. 3).

Generation of focused and composite maps for model buildings

Composite maps for the total of 16 states were generated from combined focused

maps to facilitate model building. Focused maps were generated using subtraction and

refinement masks generated in CryoSparc v4.6. Each focused map was made by particle

subtraction with a masked region followed by masked local refinement. Local resolution

estimation for overall and all focused maps and filtering of the overall map were performed

using CryoSparc v4.6. Focused maps were combined into a composite map using the

'vop max' command in ChimeraX[38] (**Supplementary Figs. 4-25**).

Model building and refinement

A combination of AlphaFold structure predictions[39] , existing X-ray/EM structures,

and *de novo* model building was used to build the 16 SSU processsome assembly

intermediates. A starting model (PDB: 5WLC/6KE6) that included all ribosomal proteins

and RNA was used as initial template for rigid body docking into the State H composite

map since it is of highest resolution. All template ribosomal protein models were manually

adjusted using COOT[40] . State H was then used as a template to build the proteins and

RNA into the other 15 states with manual adjustments in COOT. The final models for the

16 states were real-space refined with three cycles of refinement in PHENIX using

phenix.real_space_refine[41] using secondary structure restraints for proteins and RNA. In

regions with medium to low resolution, protein sidechains were trimmed to the C_β position

after all-atom refinement. The final model refinement statistics can be found in

Supplementary Table 1. The maps and models were analyzed and visualized in

ChimeraX[38] .

SSU processsome predictome

736 Proteins present within States A-O together with 14 exosome proteins were screened for

737 binary interactions. The resulting 3570 unique interactions were screened using the

738 default settings in the AlphaPulldown[23] implementation of Alphafold-Multimer[22]. For the

739 SSU processsome predictome (**Fig. 2a**) the median iptm_ptm score of five models was

740 plotted (**Extended Data Table 2**).

Yeast Growth Assays

The *rrp6Δ utp14 Δ* strain and *rrp6Δ utp18 Δ utp14 Δ* strain in the BY4741

background were used for all studies. These strains were transformed with two yeast

centromeric vectors, one vector under URA3 selection bearing wild-type RRP6 and

UTP14 derived from pRSII416 and second vector under LEU2 selection bearing wild-type

RRP6 or rrp6 alleles containing mutations or c-terminal truncations in conjunction with

UTP14 wild-type or utp14 alleles containing mutations derived from pRSII415. Strains

carrying both pRSII415 and pRSII416 plasmids were selected on minimal media (SD-Ura

and Leu) after transformation. Colonies grown on the selection plates were selected and

grown in minimal media (SD-Ura and Leu) liquid cultures. Loss of the *URA3* plasmid was

done on minimal media agar plates (SD–Leu + 5-FOA) plates by spotting serial 10-fold

dilutions (starting at OD_{600} of 1) of liquid cultures. Growth was monitored at 30°C over a

period of 5 days.

Data availability

Raw unaligned multi-frame movies and aligned micrographs have been deposited in the

Electron Microscopy Public Image Archive: Dhr1 and Kre33 dataset: EMPIAR-XXXX,

Dhr1 dataset: EMPIAR-XXXX, and Dhr1 and Utp14 dataset: EMPIAR-XXXX. The cryo-

EM maps and atomic models have been deposited in the Electron Microscopy Data Bank

(EMDB) and the Protein Data Bank (PDB): State A (EMDB-49075, PDB 9N6V), State A*

(EMD-49076, PDB 9N6W), State B (EMD-49077, PDB 9N6X), State C (EMD-49078, PDB

9N6Y), Sate D (EMD-49079, PDB 9N6Z), Sate E (EMD-49080, PDB 9N70), State F

(EMD-49082, PDB 9N72), State G (EMD-49083, PDB 9N73), State H (EMD-49084, PDB

9N74), State I (EMD-49085, PDB 9N75), State J (EMD-49086, PDB 9N76), State K

(EMD-49087, PDB 9N77), State L (EMD-49088, PDB 9N78), State M (EMD-49089, PDB

9N79), State N (EMD-49090, PDB 9N7A), State O (EMD-49091, PDB 9N7B).

Acknowledgements

We would like to thank M. Ebrahim, J. Sotiris, and H. Ng at the Evelyn Gruss Lipper Cryo-

Electron Microscopy Resource Center at The Rockefeller University for assistance with

grid screening and data collection. Mass spectrometry data was generated by the

Proteomics Resource Center at The Rockefeller University (RRID:SCR_017797) using

instrumentation funded by the Sohn Conferences Foundation and the Loena M. and Harry

B. Helmsley Charitable Trust, and we particularly thank H. Molina, S. Heissel, and C.

Peralta for assistance. We thank A. Vanden Broeck for advice with cryo-EM data

processing and scripts for figures. We also thank K. Ye for sharing experimental data from

a prior study[10] and members of the Klinge laboratory for critical reading of this manuscript.

Author information

Laboratory of Protein and Nucleic Acid Chemistry, The Rockefeller University, New York,

NY, USA

Sebastian Klinge, Olga Buzovetsky

Contributions

791 S.K. conceived the study. O.B. and S.K. designed the experiments and analyzed data.

792 O.B. performed all experiments, including strain generation, complex purification, cryo-

793 EM data processing, model building, and yeast genetics studies. All authors wrote and

794 edited the manuscript.

Corresponding author

797 Correspondence to Sebastian Klinge (klinge@rockefeller.edu).

Ethics declarations

800 The authors declare no competing interests.

Funding

Research reported in this publication was supported by the National Institute of General

Medical Sciences of the National Institutes of Health under Award Numbers

R01GM145950 and R35GM156426, and the Chan Zuckerberg Initiative Exploratory Cell

Network.

Supplementary Information. Supplementary Figs. 1-28

Extended Data. Extended Data Figs. 1-6, Extended Data Tables 1 and 2.

Extended Data Fig. 1. Cryo-EM maps and models of the SSU processomes.

(**a**) Composite cryo-EM maps and (**b**) atomic models colored by protein components highlighting stepwise maturation of the SSU processome.

Extended Data Fig. 2. rRNA compaction and folding events across states.

(**a**) Compaction of the 18*S* 5′ domain from States A-C side view (left) and top view (right).

Labeled RNA elements are color coded. Arrows highlight compaction between states A

818   and C. (**b**) Simplified rRNA folding events leading up to A1 cleavage. (**c**) Compaction of

819   the central and 5′ domains during states I-M.

Extended Data Fig. 3. Molecular proofreading for A1 cleavage during SSU processome maturation. Detailed views involving states prior to (States C, D, E) and after A1 cleavage (State G) with color-coded proteins. The A1 cleavage site, U3 architectural sites, and active site of Utp24 are highlighted and flexible interactions indicated by dashed lines.

Extended Data Fig. 4. Yeast SSU processome predictome.

(**a**) Color coded yeast SSU processsome predictome between assembly factors (AFs),

ribosomal proteins (RPs) and RNA exosome proteins (EXO). Key interactors of Mpp10,

Kre33, Utp14, Dhr1 and Bfr2 are color coded. (**b**) State C containing proteins listed in (a).

(**c**) State H containing proteins listed in (a). (**d**) Sas10-domain containing Rrp6/Rrp47

complex and predicted Bfr2-Lcp5, and Mpp10-Sas10 complexes. The schematic of

Mpp10 contextualizes predictions of Mpp10 interactors. (**e**) Mutually exclusive

interactions with Kre33 (human NAT10) predicted for Bms1 and Faf1 (human C1orf131).

(**f**) Predictions of Dhr1/DHX37 interactions. Mutually exclusive binding of Pno1 and Utp14

to yeast Dhr1 are highlighted. Elements of human UTP14 known to activate human

DHX37 (MIDL, MIEF)[19] are indicated. Confidence levels (pLDDT) are color coded for

each protein.

Extended Data Fig. 5. Yeast growth assays.

(**a**) Schematics of Utp18, Rrp6, Mpp6, and Utp14 with interacting and functional elements

labeled with multi-Ala mutations used in this study as well as an alignment to Mpp6

exosome recruitment motif, indicated in grey. Designed truncations of Rrp6 lasso (bottom

left) with interacting and functional elements labeled. Locations of all alanine substitutions

for Utp18, Rrp6, and Utp14 are indicated on the right. (**b**) Growth analysis of Utp18 multi-

Ala mutant in Δrrp6 and Δutp14 background (top and middle panels) and Utp18 WT in

Δrrp6 and Δutp14 background (bottom panel). Growth assay using WT, truncations and

mutant Rrp6 or Utp14 constructs in the absence of genomic Rrp6 and Utp14. Colonies

sequenced are indicated on the right of the growth assays.

Extended Data Fig. 6. Functional elements of Utp14, Rrp6, Utp18 and Dhr1.

(**a**) Schematics of Utp14, Utp18 and Rrp6 across States A-O. Utp14 functional sequence

elements structured, bound, relocated, or activating are indicated on the left and the

presence/absence of Utp18 is indicated in the middle. Visible regions of the Rrp6 C-

terminus (lasso) are indicated on the right. (**b**) Binding of Rrp6 lasso to ribosomal proteins

S24 in State L. (**c**) Rrp6 exclusion by ES6D and S4 in State M. (**d**) State C highlights the

initial binding of the Rrp6 lasso and the initial sequestration of Utp14 elements a and b

that later recruit Dhr1. (**e**) Previously characterized mutations[18,42] mapped onto Dhr1 and

Utp14 represented in State O. Functional elements of Utp14 are shown in orange. (**f**)

Mode of Dhr1 activation by Utp14 in State O. (**g**) Mode of activation of DHX15 by G-patch

protein NKRF (PDB: 6SH7).

	State A PDB 9N6V EMD-49075	State A* PDB 9N6W EMD-49076	State B PDB 9N6X EMD-49077	State C PDB 9N6Y EMD-49078	State D PDB 9N6Z EMD-49079	State E PDB 9N70 EMD-49080
Data collection and processing						
Microscope	Titan Krios	Titan Krios	Titan Krios	Titan Krios	Titan Krios	Titan Krios
Voltage (keV)	300	300	300	300	300	300
Camera	K3	K3	K3	K3	K3	K3
Magnification	64,000	64,000	64,000	64,000	64,000	64,000
Pixel size at detector (Å/pixel)	1.063	1.063	1.063	1.063	1.063	1.063
Total electron exposure (e-/Å2)	60	60	60	60	60	60
Exposure rate (e-/pixel/sec)	30	30	30	30	30	30
Number of frames collected (no.)	40	40	40	40	40	40
Defocus range (μm)	-1.0 to -2.5	-1.0 to -2.5	-1.0 to -2.5	-1.0 to -2.5	-1.0 to -2.5	-1.0 to -2.5
Energy filter slit width (V)	20	20	20	20	20	20
Automation software	SerialEM	SerialEM	SerialEM	SerialEM	SerialEM	SerialEM
Micrographs collected (no.)	44,272	44,272	44,272	44,272	44,272	44,272
Total extracted particles (no.)	2,817,609	2,817,609	2,817,609	2,817,609	2,817,609	2,817,609
Final particle images (no.)	36,965	56,168	28,151	23,899	12,826	9,125
Point-group symmetry	C1	C1	C1	C1	C1	C1
Resolution (global at FSC-0.143, Å)						
Overall map (masked)	3.23	3.05	3.71	3.65	4.74	5.17
Map sharpening B factor (Å^2)						
Overall map	-26.2	-35.8	-28.9	-23.3	-21.5	-32.6
Focused maps	-26.2 to -583.3	-35.8 to -423.3	-26.2 to -88.6	-23.3 to -80	-4.8 to -297.7	-20.6 to -241.1
Model composition						
Non-hydrogen atoms	243,904	243,904	244,461	245,176	157,426	162,545
Chains (proteins/RNA)	73/3	73/3	73/3	74/3	68/3	69/3
Protein residues	26,535	26,535	26,820	26,828	25,260	26,098
RNA residues	1847	1847	1825	1825	1430	1471
Ligands (Mg^{2+}/Zn^{2+}/ ATP/ADP/GTP)	25/2/1/0/ 1/	25/2/1/0/ 1/	34/2/1/0/ 1/	34/2/1/0/ 1/	0/0/0/0/ 0/	0/0/0/0/ 0/
Model refinement						
Refinement package (real space)	Phenix 1.19.1	Phenix 1.19.1	Phenix 1.19.1	Phenix 1.19.1	Phenix 1.19.1	Phenix 1.19.1
Initial models used (PDB code)	9N6W	9N74	9N6W/9N74	9N6W/9N74	9N6W/9N74	9N6W/9N74
Model resolution cutoff (Å)	3.4	3.3	3.9	3.9	5.0	6.5
Model-Map CC (CC$_{mask}$/CC$_{box}$/ CC$_{peaks}$/CC$_{volume}$)	0.77/0.76/ 0.68/0.77	0.79/0.76/ 0.71/0.78	0.76/0.74/ 0.65/0.76	0.77/0.76/ 0.67/0.77	0.66/0.76/ 0.50/0.66	0.69/0.76/ 0.56/0.69
Model-to-map FSC, threshold 0.50 (masked/unmasked, Å)	3.27/3.51	3.13/3.31	3.87/4.06	3.81/4.04	6.90/7.59	6.73/7.52
RMSDs						
Bond lengths (Å)	0.004	0.004	0.005	0.004	0.004	0.004
Bond angles (°)	0.890	0.912	0.955	0.923	1.021	1.028
B-factors						
Average B-factors (A^2)						
Protein	88.75	62.35	90.82	95.63	257.44	229.78
RNA	111.41	87.92	109.42	109.97	281.97	263.75
Ligand	75.43	57.49	67.50	73.12	-	-
Validation						
Clash score	3.60	3.47	5.21	4.50	1.37	1.36
MolProbity score	1.15	1.14	1.30	1.22	0.87	0.87
CaBLAM outliers	3.11	3.02	3.33	3.14	3.27	3.20
Poor rotamers (%)	0.00	0.00	0.01	0.00	0.00	0.00
C-beta deviations	0.00	0.00	0.00	0.00	0.00	0.02
EMRinger score	2.49	2.95	1.98	2.13	N/A	N/A
Ramachandran plot						
Favored (%)	98.54	98.38	97.91	98.30	98.01	98.14
Allowed (%)	1.46	1.62	2.09	1.70	1.99	1.86
Outliers (%)	0.00	0.00	0.00	0.00	0.00	0.00
RNA validation						
Average suiteness (%)	58.3	57.9	54.4	55.6	54.1	52.7
Good sugar puckers (%)	99.94	99.94	99.95	99.89	99.93	99.72

	State F PDB 9N72 EMD-49082	State G PDB 9N73 EMD-49083	State H PDB 9N74 EMD-49084	State I PDB 9N75 EMD-49085	State J PDB 9N76 EMD-49086	State K PDB 9N77 EMD-49087
Data collection and processing						
Microscope	Titan Krios	Titan Krios	Titan Krios	Titan Krios	Titan Krios	Titan Krios
Voltage (keV)	300	300	300	300	300	300
Camera	K3	K3	K3	K3	K3	K3
Magnification	64,000	64,000	64,000	64,000	64,000	64,000
Pixel size at detector (Å/pixel)	1.063	1.063	1.063	1.063	1.063	1.063
Total electron exposure (e-/Å2)	60	60	61.6	61.6	61.6	61.6
Exposure rate (e-/pixel/sec)	30	30	30.8	30.8	30.8	30.8
Number of frames collected (no.)	40	40	40	40	40	40
Defocus range (μm)	-1.0 to -2.5	-1.0 to -2.5	-1.0 to -2.5	-1.0 to -2.5	-1.0 to -2.5	-1.0 to -2.5
Energy filter slit width (V)	20	20	20	20	20	20
Automation software	SerialEM	SerialEM	SerialEM	SerialEM	SerialEM	SerialEM
Micrographs collected (no.)	44,272	44,272	84,463	84,463	84,463	84,463
Total extracted particles (no.)	2,817,609	2,817,609	13,198,550	13,198,550	13,198,550	13,198,550
Final particle images (no.)	6,227	4,987	328,554	95,512	11,394	19,571
Point-group symmetry	C1	C1	C1	C1	C1	C1
Resolution (global at FSC-0.143, Å)						
Overall map (masked)	5.94	5.96	2.69	3.01	4.23	3.94
Map sharpening B factor (Å^2)						
Overall map	-88.2	-72.7	-65.4	-45.5	-11.6	-16
Focused maps	-17.9 to -529.7	-12.7 to -309.5	-65.3 to -86.7	-51.5 to -79.5	-116 to -29.9	-16 to -40.2
Model composition						
Non-hydrogen atoms	162,433	163,806	238,225	220,521	206,690	190,271
Chains (proteins/RNA)	70/3	71/3	75/3	72/3	70/3	66/3
Protein residues	25,995	26,337	25,221	23,094	23,115	21,392
RNA residues	1491	1474	1761	1710	1722	1661
Ligands (Mg^{2+}/Zn^{2+}/ ATP/ADP/GTP)	0/0/0/0/0	0/0/0/0/0	39/3/1/1/1	59/3/1/1/1	38/2/1/1/1	48/2/0/1/1
Model refinement						
Refinement package (real space)	Phenix 1.19.1	Phenix 1.19.1	Phenix 1.19.1	Phenix 1.19.1	Phenix 1.19.1	Phenix 1.19.1
Initial models used (PDB code)	9N6W/9N74	9N6W/9N74	5WLC/6KE6	9N74	9N75	9N76
Model resolution cutoff (Å)	6.5	6.5	3.0	3.3	4.5	4.2
Model-Map CC (CC$_{mask}$/CC$_{box}$/ CC$_{peaks}$/CC$_{volume}$)	0.63/0.74/ 0.49/0.64	0.68/0.76/ 0.52/0.67	0.77/0.77/ 0.74/0.78	0.77/0.77/ 0.72/0.77	0.73/0.75/ 0.60/0.72	0.74/0.76/ 0.64/0.74
Model-to-map FSC, threshold 0.50 (masked/unmasked, Å)	7.51/8.02	7.30/7.84	2.84/2.99	3.14/3.29	4.37/6.33	4.09/4.50
RMSDs						
Bond lengths (Å)	0.005	0.004	0.005	0.004	0.004	0.004
Bond angles (°)	1.023	1.025	0.919	0.895	0.947	0.930
B-factors						
Average B-factors (A^2)						
Protein	223.98	226.01	29.43	48.61	152.23	118.18
RNA	250.68	248.95	39.81	64.27	176.40	141.33
Ligand	-	-	32.71	52.81	155.15	111.32
Validation						
Clash score	3.67	1.07	3.39	3.22	5.79	4.94
MolProbity score	1.16	0.81	1.13	1.11	1.32	1.26
CaBLAM outliers (%)	3.33	3.29	3.13	3.07	3.17	3.29
Poor rotamers (%)	0.00	0.00	0.00	0.01	0.05	0.05
C-beta deviations	0.00	0.00	0.00	0.00	0.01	0.00
EMRinger score	N/A	N/A	3.54	2.76	0.64	1.17
Ramachandran plot						
Favored (%)	98.13	98.07	98.42	98.57	98.34	98.23
Allowed (%)	1.87	1.93	1.58	1.43	1.66	1.77
Outliers (%)	0.00	0.00	0.00	0.00	0.00	0.00
RNA validation						
Average suiteness (%)	51.8	53.4	54.0	56.2	55.3	54.9
Good sugar puckers (%)	99.66	99.59	99.54	99.46	99.36	99.34

	State L	State M	State N	State O
	PDB 9N78 EMD-49088	PDB 9N79 EMD-49089	PDB 9N7A EMD-49090	PDB 9N7B EMD-49091
Data collection and processing				
Microscope	Titan Krios	Titan Krios	Titan Krios	Titan Krios
Voltage (keV)	300	300	300	300
Camera	K3	K3	K3	K3
Magnification	64,000	64,000	64,000	64,000
Pixel size at detector (Å/pixel)	1.063	1.063	1.063	1.063
Total electron exposure (e-/Å2)	61.6	61.6	61.6	61.6
Exposure rate (e-/pixel/sec)	30.8	30.8	30.8	30.8
Number of frames collected (no.)	40	40	40	40
Defocus range (μm)	-1.0 to -2.5	-1.0 to -2.5	-1.0 to -2.5	-1.0 to -2.5
Energy filter slit width (V)	20	20	20	20
Automation software	SerialEM	SerialEM	SerialEM	SerialEM
Micrographs collected (no.)	84,463	84,463	84,463	85,031
Total extracted particles (no.)	13,198,550	13,198,550	13,198,550	27,069,273
Final particle images (no.)	10,725	12,127	14,551	131,106
Point-group symmetry	C1	C1	C1	C1
Resolution (global at FSC-0.143, Å)				
Overall map (masked)	4.17	3.93	3.84	3.25
Map sharpening B factor (Å^2)				
Overall map	-12.2	-16.6	-20.2	-52.4
Focused maps	-12.2 to -30.9	-24.4 to -90.2	-27.9 to -78.8	-60.3 to -85.2
Model composition				
Non-hydrogen atoms	149,183	106,665	93,089	85,883
Chains (proteins/RNA)	60/3	46/3	40/3	38/3
Protein residues	15,942	10,667	8,701	7,602
RNA residues	1587	1493	1428	1428
Ligands	47/2/0/0/	47/2/0/0/	47/2/0/0/	47/2/0/0/
(Mg^{2+}/Zn^{2+}/ATP/ADP /GTP)	1	1	1	1
Model refinement				
Refinement package (real space)	Phenix 1.19.1	Phenix 1.19.1	Phenix 1.19.1	Phenix 1.19.1
Initial models used (PDB code)	9N77	9N78	9N79	9N7A
Model resolution cutoff (Å)	4.4	4.1	4.1	3.6
Model-Map CC ($CC_{mask}/CC_{box}/CC_{peaks}/CC_{volume}$)	0.73/0.74/ 0.59/0.72	0.78/0.77/ 0.67/0.78	0.81/0.78/ 0.70/0.81	0.74/0.78/ 0.69/0.74
Model-to-map FSC, threshold 0.50 (masked/unmasked, Å)	4.17/4.83	4.07 /4.62	3.80 /4.66	3.41 /3.73
RMSDs				
Bond lengths (Å)	0.004	0.004	0.004	0.004
Bond angles (°)	0.925	0.895	0.876	0.855
B-factors				
Average B-factors (A^2)				
Protein	144.24	190.48	149.72	61.64
RNA	150.59	158.59	131.07	85.04
Ligand	86.60	103.26	85.63	38.04
Validation				
Clash score	5.08	4.15	4.07	2.99
MolProbity score	1.27	1.20	1.19	1.09
CaBLAM outliers	3.10	2.89	3.20	3.23
Poor rotamers (%)	0.05	0.05	0.04	0.00
C-beta deviations	0.00	0.00	0.00	0.00
EMRinger score	1.08	1.46	1.79	2.33
Ramachandran plot				
Favored (%)	98.30	98.48	98.37	98.28
Allowed (%)	1.70	1.52	1.63	1.72
Outliers (%)	0.01	0.00	0.00	0.00
RNA validation				
Average suiteness (%)	55.7	56.6	56.1	57.0
Good sugar puckers (%)	99.31	99.40	99.37	99.37

Extended Data Table1. Cryo-EM data collection and refinement statistics.