1 **Aneuploidy promotes transient stress adaptation and metabolic flexibility in the human**
2 **fungal pathogen *Aspergillus fumigatus***
3
4
5

6 Anna E. Lehmann[1], Enrique Aguilar Ramírez[2], Nancy P. Keller[2], and Joseph Heitman[1*]
7
8
9
10 [1]Department of Molecular Genetics and Microbiology, Duke University School of Medicine, Durham,
11 NC, USA
12 [2]Department of Medical Microbiology and Immunology, University of Wisconsin – Madison, Madison,
13 WI, USA
14
15 *Corresponding author: Joseph Heitman
16 Email: heitm001@duke.edu

17
18
Highlights
- *A. fumigatus* rapidly gains and loses supernumerary chromosomes in response to the
calcineurin inhibitor FK506.
- Aneuploidy leads to global transcriptional and metabolic rewiring.
- Overexpression of the secondary metabolite gene cluster regulator *nscR* is sufficient to
largely recapitulate the aneuploid response to FK506 in a haploid background.
- Chr4, 6, and 7 aneuploidies reduce susceptibility to voriconazole without altering expression
of the azole target gene *cyp51A*.
27

Summary

Aneuploidy causes genome plasticity and enables adaptive responses that increase stress resistance and facilitate persistence under changing conditions in eukaryotes ranging from fungal pathogens to human cancer cells. *Aspergillus fumigatus* is a soil-resident mold and the most prevalent etiologic agent of invasive fungal infections globally. Invasive aspergillosis has an alarmingly high mortality rate, and fungal persistence in the infection microenvironment often renders treatments unsuccessful. Treatment failures can result from innate or acquired antifungal drug resistance and from fungal adaptation to other stressors encountered during infection including nutrient limitation, immune cell activity, and changes in pH and oxygen tension. Survival under changing environmental and host conditions therefore necessitates an ability to dynamically adapt to stress. We find that exposure of *A. fumigatus* to FK506, an antifungal and immunosuppressive *Streptomyces* natural product that inhibits the master regulatory phosphatase calcineurin, selects for unstable whole-chromosome aneuploidies that alleviate the polarized growth defects caused by calcineurin inhibition. Transcriptomic analysis revealed that Chr7 disomy leads to induction of the normally-silent neosartoricin biosynthetic gene cluster in response to FK506 exposure, and we demonstrate that constitutive genetic induction of this cluster is sufficient to largely recapitulate in a haploid background the response to FK506 in the aneuploid state. We further show that aneuploids undergo extensive metabolic rewiring but do not produce detectable neosartoricin, revealing global and cross-pathway effects resulting from both the aneuploid state and activation of *nscR*. We also demonstrate that the aneuploid chromosome duplications reduce susceptibility to the clinical antifungal voriconazole, underscoring the utility of aneuploidy as a flexible adaptive strategy. These findings represent a major advance in understanding how aneuploidy transiently alters stress responses and metabolism in the deadly human pathogen *A. fumigatus.*

Keywords: *Aspergillus fumigatus*, antimicrobial drug responses, aneuploidy, secondary metabolism

Introduction

Aneuploidy, an abnormal complement of whole chromosomes or chromosome segments, is a conserved adaptive strategy across eukaryotes. Whole-chromosome and segmental aneuploidies are ubiquitous in human cancers and are hypothesized to enhance tumor survival and proliferation in a dynamic and context-specific manner by altering gene dosage and activating generalized stress responses (reviewed in[1–3]). Aneuploidy is also an important persistence strategy in single-celled eukaryotes. In the model yeast *Saccharomyces cerevisiae*, unstable aneuploidy can serve as a rapidly-acquired, temporary solution to diverse stressors that potentially provides an avenue to less deleterious, stable genetic change[4–8]. Aneuploidy is also known to underlie unstable stress adaptation in multiple human and plant fungal pathogens[9]. Significantly, aneuploidy can serve as a

major cause of reduced susceptibility to triazoles, the primary antifungal class administered clinically.

In *Candida albicans*, azole resistance results from isochromosome formation of the left arm of Chr5,

which harbors the genes that encode the azole target enzyme Erg11 and the drug efflux pump-

activating transcription factor Tac1[10–12]. Whole-chromosome aneuploidy has also been reported to

reduce *in vitro* antifungal susceptibility in the pathogenic yeast *Cryptococcus neoformans*[13,14] and

the plant pathogen *Aspergillus flavus*[15]. Additionally, a recent report identified segmental duplications

of regions encoding the azole target-encoding genes *cyp51A* and *cyp51B* in azole-resistant

Aspergillus fumigatus clinical isolates, revealing a correlation between copy number variation and

clinically-relevant antifungal resistance in this globally ubiquitous pathogen[16]. These observations

support a broadly conserved role for aneuploidy in stress adaptation in pathogenic fungi. Despite the

apparent conservation of this phenomenon, however, genomic plasticity as an unstable adaptive

mechanism remains underexplored in filamentous fungi.

A. fumigatus is a globally-distributed, soil-resident fungus and the leading cause of invasive

fungal infections worldwide[17]. This organism has evolved a range of strategies that facilitate

persistence in its harsh and variable ecological niches, and this same suite of adaptive traits can

contribute to its pathogenicity. One such adaptation is the inducible production of secondary

metabolites, two of the most well-studied being 1) DHN melanin, which protects conidia from UV

radiation and reactive oxygen species and inhibits host cell apoptosis[18–20] and 2) gliotoxin, which is

toxic to mammalian host cells[21–24]. Numerous other secondary metabolites exhibit both antimicrobial

activity against soil microbes and cytotoxic activity against host immune cells, suggesting context-

dependent and evolutionarily-relevant reasons for their maintenance in the genome and selective

induction (reviewed in[25,26]). The genes involved in secondary metabolite biosynthesis are encoded in

self-contained and spatially discrete biosynthetic gene clusters (BGCs), which are often

transcriptionally inactive under standard laboratory culture conditions[27]. Difficulties in activating silent

BGCs continue to hinder efforts to elucidate the functions of these enzymes and products in fungal

physiology and adaptation. Recent efforts to activate normally-silent BGCs have employed co-

culture with soil microbes including *Streptomyces* and other Aspergilli that co-occupy a saprophytic

environmental niche[28–30]. Several such studies have resulted in the identification of new natural

products, suggesting a role for these compounds in cross-talk and interaction with other, potentially

competing microbes and further evidencing their ecological relevance[28,29,31–34].

Beyond the triggers for transcriptional activation being largely unknown, BGC induction is

further complicated by the often-observed endogenous crosstalk of metabolites within a species.

Myriad examples of this cross-talk have been observed in *A. fumigatus*, including loss of one

metabolite (e.g. fumigaclavine C or hexadehydroastechrome) resulting in increased production of

another metabolite (fumitremorgins)[35,36], evidence of an enzyme from one BGC active in synthetic

potential in another BGC[37], and changes in BGC gene expression upon deletion of genes in a

different BGC[38]. This complexity underscores the intricate and fine-tuned biology underlying *A.*

fumigatus secondary metabolism and the wealth of information that remains to be uncovered about

these processes.

Despite the apparent ubiquity of aneuploidy as a rapidly-acquired, flexible adaptive

mechanism in eukaryotes, the potential roles of duplication and loss of genetic material in *A.*

fumigatus stress responses remains a significant gap in knowledge. Here, we identify whole-

chromosome aneuploidies in the *A. fumigatus* reference strain A1163 and in several clinical and

environmental isolates. We demonstrate that Chr7 disomy induces transcription of the neosartoricin

(*nsc*) BGC, which is silent under standard laboratory culture conditions. Induction of the *nsc* gene

cluster transcription factor *nscR* in the A1163 haploid background ($nscR^{OE}$) is sufficient to alter

phenotypic responses to FK506 in a manner consistent with that observed in Chr7 aneuploids.

Neosartoricin is produced by $nscR^{OE}$, but not Chr7 aneuploids, suggesting a potential additional role

for NscR in stress responses and/or metabolism. $nscR^{OE}$ and Chr7 aneuploids also undergo broad

changes in both primary and secondary metabolism, and these expression changes do not

necessarily correspond to expression changes in synthetic pathway genes. These findings reveal

that, in addition to altering the transcriptome, aneuploidy also functions to promote metabolic

flexibility. We also find that FK506-selected aneuploid chromosome duplications reduce susceptibility

to the clinical antifungal voriconazole, illustrating the capacity for aneuploidy to confer fitness

advantages under distinct and disease-relevant conditions. These findings extend evidence of

whole-chromosome aneuploidy as a conserved adaptive strategy to the important fungal pathogen

A. fumigatus, revealing a novel avenue by which dynamic stress adaptation and metabolic rewiring

occur in this species.

Results

Unstable aneuploidy alleviates hyphal growth defects caused by FK506 in the A. fumigatus

reference strain A1163

Our previous work has utilized FK506, an antimicrobial and immunosuppressive compound

produced by *Streptomyces*, to uncover stable and unstable genetic adaptive strategies in the fungal

pathogens *Mucor*[39,40] and *Cryptococcus*[41–43]. FK506 binds to the endogenously-expressed protein

FKBP12 and forms a complex that inhibits the protein phosphatase calcineurin, a mechanism that is

broadly conserved in eukaryotes[44–46]. In *A. fumigatus*, calcineurin inhibition by pharmacologic or

genetic means impacts a host of processes including development and morphology of conidia,

septation, polarized growth, and cell wall composition[47–51]. The result of these severe growth and

morphological defects is an abolition of virulence in pathogenesis models, underscoring the

essentiality of calcineurin-mediated processes to the pathogenic capacity of this organism[48].

While several downstream effectors of calcineurin have been described, most extensively

the transcription factor CrzA[49,52,53], detailed mechanistic bases for calcineurin-dependent

morphological traits remain ill-defined. To further delineate these important processes, we employed

a suppressor screen with the goal of elucidating genetic and/or epigenetic means by which *A.*

fumigatus adapts to calcineurin inhibition. Conidia were plated on *Aspergillus* minimal medium

(AMM) containing FK506 at a concentration of 1 µg/mL, which severely impairs growth of the A1163

wild-type strain (Fig. 1*A*). Colonies that exhibited increased vegetative growth were isolated and

single-spored. A total of 52 FK506-adapted isolates were obtained, and whole-genome sequencing

revealed medium- or high-impact mutations in 14 isolates, occurring in 6 genes that we predict are

involved in bypass suppression of calcineurin (Fig 1*B*; Table S1). Surprisingly, a majority (34/52) of

these isolates had duplications of 1 or more full chromosomes, and none of these aneuploids had

high- or medium-impact sequence changes that we predict are likely to impact the observed

phenotypic changes (Fig 1*B*; Table S1). We interestingly did not identify any mutations in the genes

encoding the FK506 target, *fkbp12-1*, or the calcineurin catalytic or regulatory subunits, *cnaA* and

cnaB. One possible reason for the failure to isolate mutants in these genes could be due to

aneuploidy conferring a lower fitness cost than these genetic mutations, and/or the relatively short

timeframe and growth-permissive concentration of drug used.

152
Figure 1. Unstable aneuploidy alleviates polarized growth abnormalities caused by FK506. (*A*)
A1163 WT growth on AMM – (top) or + (bottom) 1 µg/mL FK506 for 3 days at 37°C. (*B*) Distribution
of aneuploidies, high- or medium- impact mutations, and unknown mode of change determined from
analysis of WGS data from FK506-adapted isolates. Enlarged pie chart shows frequency of

chromosome duplications in the set of 34 aneuploids. (*C*) 2 representative Chr7 aneuploids grown

on AMM + 1 μg/mL FK506 for 3 days before (top) and after (bottom) passage on drug-free AMM.

Representative images from n ≥ 3 biologically independent replicates. (*D*) Normalized read depth

averaged over 500 bp intervals, plotted relative to the closely-related CEA10 reference assembly, for

strains in (*C*). (*E*) Conidia were inoculated in AMM +/- 1 μg/mL FK506 in a microscopy growth

chamber and grown at 37°C for 24 hours. In each panel (-FK506 and +FK506), scale bars indicates

50 μm (top) and 20 μm (bottom). Representative images from n ≥ 3 biological replicates.

We elected to first focus our study on Chr7 aneuploids due to their prevalence in our screen
(9/52 isolates) and the fact that Chr7 is the smallest chromosome in the genome, and previous work
in yeast, *Drosophila*, and human cells has demonstrated that aneuploidies of smaller genomic
regions are often better tolerated than those of larger regions[3]. Chr7 disomic strains (Di 7) have a
growth defect on medium without FK506 (Fig. 1*C*), and when Di 7 conidia were plated on drug-free
medium we observed heterogeneity in colony morphology that we hypothesized could be due to
heterogeneous gain or loss of supernumerary chromosomes. To assess aneuploid chromosome
stability in the absence of FK506, strains were cultured on drug-free AMM for at least 9 days. After
drug-free passage, conidia were re-plated on FK506 medium. Passage on drug-free medium
resulted in reversion to a wild-type, haploid level of growth on FK506 (Fig. 1*C*). Sequencing of the
reverted (*REV*) isolates showed a corresponding loss of the aneuploid chromosomes, linking the
observed morphological changes directly to the gain of a second copy of Chr7 (Fig 1*D*; Table S1).

At the beginning of this study, we identified 6 isolates that originated from a single initial pool
of conidia but formed separate colonies with similar enhanced growth phenotypes on individual
FK506 selection plates. Whole-genome sequencing of these isolates revealed that all 6 carried an
extra copy of Chr7 and differed from the progenitor strain by several identical SNPs in coding
regions of the genome (Table S1). Based on the genetic and ploidy similarities, we hypothesize that
these isolates may have originated from a clonal population of Di 7 conidia in the initial pool. Due to
their apparent clonality, these 6 isolates are treated as a single independent event (Di 7-1) in all
results presented here. To avoid identifying additional clonal isolates, and to confirm that the
additional SNPs present in the aneuploids are not responsible for the aneuploid karyotype, all other
isolates described in this study were isolated from 20 different single-spore derived populations of
conidia from the original A1163 stock, and one FK506-adapted colony was isolated per replicate.

Consistent with previously-described effects of calcineurin inhibition[48,50,54], and stress
responses more generally[55,56], we observed defects in polarized growth, indicated by increased
density of hyphae, stunted extension, and aberrant branching at the apical hyphal tips, in the haploid
+FK506 (Fig. 1*F*). In contrast, Chr7 aneuploids lack the hyper-branched hyphal tips observed in the
haploid, and these phenotypes are reversed to a state that is indistinguishable from the haploid
progenitor in the reverted (*REV*), haploid strains derived from the Chr7 aneuploids (Fig. 1*F*). These
results further implicate Chr7 aneuploidy in the reversal of polarized growth abnormalities caused by
calcineurin inhibition and indicate that aneuploidy leads to an improved ability to tolerate stress
caused by FK506.

Aneuploidy enhances growth on FK506 in distinct A. fumigatus *isolates*

Unstable adaptive responses to antifungal compounds are ill-defined in *A. fumigatus*, and we
questioned whether the high prevalence of unstable aneuploidy in response to FK506 occurred due

to a specific property of A1163, or if this transient response could be generalized to other isolates.
We selected 10 diverse, non-reference strains representing the 3 clades of *A. fumigatus*[57], as well as
both clinical and environmental sources (Table S2). 3 of these strains exhibited robust growth on
FK506 at concentrations up to 50 µg/mL (Fig. S1*A*). Of the remaining 7 strains, we successfully
repeated the screen used for A1163 at 1 µg/mL FK506 for 6 strains and 0.5 µg/mL for 1 strain. 3 to 4
independent isolates with improved growth on FK506 were obtained per strain for a total of 26
isolates. All isolates were passaged on drug-free medium for at least 15 days, and re-plating conidia
on FK506 after drug-free passage showed a loss of the growth improvements initially observed in 12
of these isolates (Fig. S1*B*). In our relatively small sample size, the number of stable and unstable
responses was not evenly distributed between strains, with some giving rise to only stable isolates,
some only unstable, and some a mix of the two (Fig. 2*A*). Nevertheless, in total we obtained stable
and unstable isolates in roughly equal proportions (14 stable, 12 unstable). Sequencing of the
unstable FK506-adapted isolates revealed, similar to A1163, widespread aneuploidy involving
disomies of Chr4 and Chr7 (Fig. 2*B*). While all aneuploidies were unstable, and no stable isolates
were aneuploid, not all unstable isolates had detectable aneuploidies (Fig. 2*B*; Fig. S2). Because
aneuploids are highly unstable, one possibility is that the non-aneuploid unstable isolates were
aneuploid upon isolation and lost the duplicated genetic material during culture prior to DNA
extraction. Another possibility is that a non-aneuploidy unstable adaptive mechanism, such as
chromatin remodeling or RNA interference-mediated gene silencing, is responsible for the FK506
responses observed in these isolates[39,58]. Together, these results confirm that unstable aneuploidy is
not restricted to the A1163 reference strain, can occur frequently under the proper selective
conditions, and is conserved in strains from clinical and environmental sources representing at least
clades 1 and 2.

Figure 2. Stable and unstable FK506-adapted isolates arise in diverse strain backgrounds.
Conidia from clinical and environmental strains were plated on AMM +0.5 µg/mL FK506 (W72310) or
1 µg/mL FK506 (all other strains) and colonies with improved vegetative growth were selected.
Stability was determined by passaging isolates on drug-free AMM for at least 15 days and re-plating
on FK506. (*A*) Distribution of stable and unstable FK506-adapted isolated identified in each strain
background. (*B*) Normalized read depth for progenitor strain (P) and unstable FK506-adapted
isolates that arose in 4 strain backgrounds, plotted relative to a closely-related assembly as
indicated at the top right of each plot.

Chr7 disomy induces expression of secondary metabolite biosynthetic genes

Aneuploidy in yeasts and humans is known to alter gene expression and phenotypic traits
both through dosage-dependent transcriptional changes and by activating dosage-independent,
aneuploidy-specific responses. In *Candida albicans*, i5L isochromosome formation confers
fluconazole resistance by altering copy number and expression of *ERG11*, which encodes the azole
drug target enzyme, and *TAC1*, which encodes an efflux pump-activating transcription factor[59]. We
therefore hypothesized that changes in gene expression in Chr7 aneuploids resulting from the
changes in copy number and/or the response to altered DNA content could explain the phenotypic
changes observed in the A1163 Chr7 aneuploids. To probe a potential transcriptional basis for the
observed phenotypic changes, we conducted 2 independent RNA sequencing experiments with the
haploid progenitor strain and independent Chr7 aneuploids (Di 7-1 and Di 7-2), with and without
FK506. Total RNA was extracted from triplicate cultures and analyzed by paired-end bulk mRNA
sequencing (Materials and Methods). Parallel DNA sequencing from Di 7-1 cultured under the drug-
free RNA extraction condition confirmed that the chromosome duplication was maintained in the
drug-free condition (Fig. S3*A*).

RNA-seq reads were mapped to the A1163 reference genome[60], and PCA was performed
analyzing the top 1000 variable transcripts in each experiment, which capture the majority of
variance in the data (Fig. S3*B, C*). In both cases, the majority of the variance was explained in

principal components 1 and 2 (Fig. S3*D, E*). PCA indicates that for Di 7-1, drug condition is the
primary factor explaining gene expression differences (PC1; 52.2% of variance), followed by ploidy
(PC2; 21.6% of variance) (Fig. S3*F*). In contrast, in Di 7-2, variance is primarily driven by ploidy
(PC1; 60.5%), with drug condition explaining 19.3% of the observed variance (Fig. S3*G*). In both
experiments, analysis of differentially expressed genes (DEGs; $p_{adj} \leq 0.05$ and $|log_2FC| \geq 1$) between
aneuploids and haploid +FK506 revealed large-scale changes in gene expression, with 291
upregulated and 167 downregulated genes in experiment 1 and 1537 upregulated and 442
downregulated genes in experiment 2 (Fig. 3*A, B*; Table S3). Similar to the PCA results, these
differences in gene expression between the two experiments illustrate a much larger effect of
aneuploidy on the global transcriptome in Di 7-2 than in Di 7-1.

We predicted that phenotypic commonalities in response to FK506 observed in Di 7-1 and Di
7-2 would most likely be explainable by transcriptional change(s) shared between the two strains.
Comparison of the up- and down-regulated gene sets between the two experiments showed that Di
7-1 had 176 up-regulated and 76 down-regulated genes in common in the FK506 condition (Fig. 3*A,*
B). Interestingly, 4 genes in the *nsc* biosynthetic gene cluster (*nsc* BGC), which regulates synthesis
of the secondary metabolite neosartoricin[61], were among the most highly induced genes in both
experiments (Fig. 3*C, D*; Table S3). The *nsc* BGC, located near the left telomere of Chr7, encodes
the transcriptional regulator *nscR* and the neosartoricin biosynthetic enzymes *nscA-E* (Fig. 3*E*). This
gene cluster, with the exception of the FAD-dependent monooxygenase *nscC*, is silent or lowly
expressed in the haploid and Chr7 aneuploids -FK506, and minimally induced in the euploid +FK506
(Fig. 3*F, G*). FK506 exposure induces *nsc* gene expression specifically in the aneuploids, with *nscR*
having the highest expression (Fig. 3*G*).

273
Figure 3. Secondary metabolite transcription factor *nscR* is induced in Chr7 aneuploids and
recapitulates the Chr7 aneuploid FK506 response in the haploid. (*A* and *B*) Genes with
increased (*A*) and decreased (*B*) expression in comparison between Chr7 aneuploid and haploid in 2
RNA-seq experiments. (*C* and *D*) Differentially expressed genes in comparison between Chr7
aneuploid +FK506 and WT +FK506 in experiment 1 (*C*) and 2 (*D*). Grey colored points denote
shared up- and down-regulated genes between the two experiments. Dashed lines indicate $\log_2\text{FC}$ =
±1 and p_{adj} = 0.05. (*E*) Location of the *nsc* BGC on Chr7. (*F* and *G*) Transcript abundance (TPM) of
nsc BGC genes in haploid and Chr7 aneuploids -FK506 (*F*) and +FK506 (*G*). Points represent
technical replicates; error bars indicate standard deviation. (H) Hyphal growth phenotypes of A1163
wild-type and *nscR^{OE}*, +/- FK506, as in Fig. 1*E*.

Secondary metabolite BGCs can be transcriptionally activated by both pathway specific
transcription factors and global regulators or epigenetic modifications that themselves are induced
by various environmental stimuli[62–64]. Given the large number of transcriptional changes observed in
the Chr7 aneuploids, we assessed whether aneuploidy globally alters secondary metabolite BGC
transcription or whether the effect is specific to the *nsc* BGC. TPM quantification of the genes in the
18 additional known A1163 secondary metabolite BGCs[65] shows that these clusters are either silent
in all conditions, or do not undergo the same cluster-wide differential change as the *nsc* BGC in
response to aneuploidy and FK506 (Fig. S4). One such example is the fumagillin BGC
monooxygenase *fmaE* (AFUB_086100), which is constitutively highly expressed across strains and
conditions while the other genes in its cluster are not expressed. Several genes in the
hexadehydroastechrome and trypacidin BGCs also undergo increased transcription in response to
FK506. Unlike the *nsc* BGC, however, most genes in these clusters remained silent. These results
led us to conclude that Chr7 duplication leads to specific induction of the *nsc* BGC expression, rather
than general induction of secondary metabolic genes, in response to FK506 exposure.

nscR *overexpression recapitulates Chr7 aneuploid phenotypes in response to FK506*

To directly explore a potential role for the *nsc* BGC in the aneuploid phenotypes observed in
response to FK506, we generated a strain expressing *nscR* (AFUB_086680) under control of the *A.*
fumigatus Af293 *gpdA* promoter in the A1163 euploid background (*nscR*^{OE}). The *nscR* transcript is
constitutively overexpressed 107-fold without FK506 and 91-fold with FK506 in this strain (Fig. S5*A*).
In static submerged culture with FK506, *nscR*^{OE} exhibits decreased hyphal hyperbranching similar to
Chr7 aneuploids (Fig. 3*H*). Additionally, *nscR*^{OE} exhibits a comparable radial growth defect to Chr7
aneuploids on drug-free medium, and a comparable, minor but statistically significant, increase in
radial growth on medium containing FK506 (Fig. S5*B-D*). Taken together, these observations
indicate that *nscR* overexpression specifically alleviates FK506-dependent polarized growth defects
in a manner similar to Chr7 aneuploidy. Our findings thus reveal that overexpression of a single gene
is sufficient to largely recapitulate a particular aneuploid stress response in a haploid background.

Aneuploid secondary metabolite profiles distinguish them from haploid and nscR^OE

We hypothesized that NscR may influence the response to FK506 through the production of
neosartoricin, which could act as a signaling molecule. Alternatively, NscR might activate enzymes
involved in other synthetic processes, or an enzyme that can degrade FK506. NscR could also have
transcriptional activity outside of the *nsc* BGC that might influence the FK506 response. To
investigate a potential correlation between *nscR* expression and neosartoricin production in Chr7
aneuploids and *nscR^OE*, we profiled the metabolites produced in these strains by LC-MS.
Surprisingly, we found that only *nscR^OE* produces neosartoricin, while this compound is absent or at
undetectable levels in the Chr7 aneuploids (Fig. 4*A-C*). Considering the many instances of
metabolite cross-talk in *A. fumigatus*, we next expanded our metabolomics analysis to assess the
abundances of well-characterized *A. fumigatus* secondary metabolites in the A1163 haploid, *nscR^OE*,
and Di 7 strains. This analysis revealed clear increases in pyripyropene A, fumitremorgin C,
pseurotin A, and fumagillin biosynthesis (Fig. 4*D*; Fig. S6). Of these metabolites, only fumitremorgin
C showed a significantly increased abundance in both the Chr7 aneuploids and *nscR^OE* (Fig. S6*A*).
These changes in secondary metabolite production occurred in the absence of clear transcriptional
induction of their BGCs, indicating potential non-concordance between BGC activation and
associated secondary metabolite production in the Chr7 aneuploids (Fig. S7). Additionally, the
extensive changes in metabolite abundances that occur as a consequence of overexpressing a
single transcriptional regulator, *nscR*, underscore the complexity in regulation of these processes
and the potential for perturbation of one biosynthetic pathway to exert widespread effects on
secondary metabolism more broadly. We also examined the abundance of FK506 in haploid, Chr7
aneuploids, and *nscR^OE* and observed no significant difference in its abundance in the Chr7
aneuploids and a small, but statistically significant, reduction in abundance in *nscR^OE* (Fig. S6*E*).
Despite this reduction in FK506 abundance, the level of the compound remained relatively high in all
of the samples, which led us to conclude that elimination of FK506 alone is unlikely to explain the
shared FK506 responses.

The relative abundances of all LC-MS features associated with *A. fumigatus* metabolites
were next compared across strains by principal component analysis (PCA). The resulting chemical
profiles segregated into three distinct groups: haploid, *nscR^OE*, and the Di 7 strains, indicating that, at
a global level, the Chr7 aneuploids and *nscR^OE* display chemical profiles clearly distinct from one
another and from the haploid wild-type (Fig. 4*E*). To further elucidate how the Chr7 aneuploids differ
from their haploid progenitor, we conducted a volcano plot analysis using the LC-MS features, which
revealed that a large number of metabolites were overproduced in the aneuploids relative to haploid
(Fig. 4*F*, *G*). Interestingly, many of these metabolites were consistent with ergosterol-type
triterpenes, based on their fragmentation patterns and predicted formulas (Fig. S8). These dramatic
alterations in metabolism, coupled with the non-concordance between transcription and metabolites

produced, indicate that aneuploidy of Chr7 promotes widespread metabolic rewiring, and that these

changes are incompletely explained by changes in gene expression.

Figure 4. Chr7 aneuploids and *nscR*^{OE} undergo extensive metabolic rewiring. (*A*) Chemical structure of neosartoricin. (*B*) Abundance of neosartoricin in the different strains, determined based on the average area under the curve of the detected peak for the molecular ion [M+H]+ = 485.1802 across the four replicates analyzed, and (*C*) its associated chromatogram. (*D*) Heatmap representing the abundance of well-characterized secondary metabolites of *A. fumigatus*, based on the average area under the curve associated with their LC-MS features. (*E*) Principal component analysis (PCA) showing the variance among the different strains according to the abundance of all LC-MS–detected features in positive mode. (*F* and *G*) Volcano plots comparing the abundance of detected features (in positive mode) between the aneuploids Di 7-1 (*F*) and Di 7-2 (*G*) and the haploid. The feature corresponding to fumitremorgin C is highlighted in green. Four replicates are included in all representations.

Chr4, 6, and 7 aneuploidies reduce voriconazole susceptibility

Azole antifungals are direct inhibitors of the ergosterol biosynthetic enzyme Cyp51A, and Cyp51B is a parallel biosynthetic enzyme that is not a primary target of the drug but has been proposed to have a role in compensating for loss of Cyp51A activity in azole-resistant isolates[66]. Increased expression of *cyp51A* via promoter mutations is an established azole resistance mechanism in *A. fumigatus*[67–73], and segmental duplications of genomic regions encoding *cyp51A* (Chr4), *cyp51B* (Chr7), and the *cyp51* electron carrier protein *cyp51ec* (Chr5) were recently correlated with elevated azole resistance in *A. fumigatus* clinical isolates[16]. *cyp51A* copy number increases in these isolates correspond to increases in *cyp51A* transcript abundance under itraconazole treatment, leading to the hypothesis that the increase in gene dosage of the drug target leads to an increased concentration of drug needed to inhibit fungal growth[16]. These observations are consistent with an established azole resistance mechanism in *C. albicans*, in which duplication of Chr5L leads to fluconazole resistance by increasing the copy number of the azole target-encoding gene *ERG11*[10,59].

Given the presence of Chr4 and Chr7 disomies in our dataset, along with the observed changes in ergosterol biosynthesis, we investigated whether aneuploidy affects susceptibility to voriconazole. We measured the MICs of one aneuploid representing each karyotype observed and its corresponding reverted (*REV*) isolate (Fig. 5*A*). We found that aneuploids with duplications of Chr4, Chr6+7, and Chr4+6+7 have significantly higher MICs than haploid (Fig. 5*B*). The MICs of the *REV* isolates were not significantly different than those of the haploid progenitor. We hypothesized that the decreased voriconazole susceptibility in aneuploids may be due to an increase in *cyp51A* and/or *cyp51B* transcript abundance. We measured *cyp51A* and *cyp51B* transcript abundances by qPCR and observed no significant differences in *cyp51A* or *cyp51B* transcript abundances in any of the aneuploids relative to the haploid (Fig. 5*C-F*). To ensure that this was not due to some property of the specific culture conditions or medium used, we performed the experiment utilizing an additional culture condition (shaking flask cultures in AMM with DMSO or 1 µg/mL voriconazole) and again observed no changes in transcript abundance of the azole target-encoding genes (Fig. S9). These observations strongly suggest that, in contrast to the known aneuploidy-driven resistance mechanism in *C. albicans*, azole susceptibility in the *A. fumigatus* whole-chromosome aneuploids is not associated with an increase in copy number or expression of the drug target genes under drug treatment. Future work should investigate the potential metabolic, transcriptomic, and/or proteomic signatures in aneuploids under azole treatment to shed light on the means by which transient supernumerary chromosome acquisition can enhance tolerance of this important clinical antifungal.

Figure 5. Chr4, 6, and 7 aneuploidies reduce voriconazole susceptibility without changes in
cyp51A **or** *cyp51B* **gene expression.** (*A*) WGS coverage plots showing different karyotypes tested.
(*B*) MICs (µg/mL voriconazole) measured by CLSI broth dilution; points represent biological
replicates and statistical test is a one-way ANOVA with corrected Bartlett's test. (*C-F*) Haploid and
aneuploid strains were cultured in RPMI for 22 hours, then medium was removed and replaced with
RPMI +DMSO or RPMI +0.2 µg/mL voriconazole. Fold change of *cyp51A* (*C, D*) and *cyp51B* (*E, F*)
transcripts under DMSO (*C, E*) or voriconazole (*D, F*) treatment. Statistical significance was
determined by one-way ANOVA with Dunnett's multiple comparison test on $\Delta\Delta$Ct values. Fold
change values, relative to haploid +DMSO, are represented as linearized $\Delta\Delta$Ct values ($2^{-\Delta\Delta Ct}$).

Discussion

Aneuploidy functions as a dynamic means of generating genomic and phenotypic diversity across the kingdom Eukarya, facilitating rapid stress adaptation in a genetically transient and readily-reversible manner. Our findings extend knowledge of this phenomenon to the important fungal pathogen *Aspergillus fumigatus*, revealing a function for whole-chromosome duplications in adapting to the calcineurin inhibitor FK506 and dynamically rewiring metabolism under FK506 stress. Aneuploidy is a hallmark of cancer, and metabolic reprogramming in aneuploid cancer cells is a major avenue supporting proliferation of tumor cells[74]. The Warburg effect, in which oncogenic cells shift to primary utilization of the oxygen-independent glycolytic pathway for energy metabolism irrespective of oxygen tension, is thought to promote tumor proliferation in part by increasing the rate of ATP production and supporting growth under hypoxia[75–78]. Glycolytic shifts in tumor cells have been linked to chromosome copy number changes, supporting a potential role for the ubiquitous aneuploidies found in cancer cells in influencing oncogenic metabolic transitions[79,80]. Similarly, in *S. cerevisiae* certain chromosomal aneuploidies increase mitochondrial respiration and lead to decreased pyruvate levels[81]. In *A. fumigatus*, metabolic plasticity is necessary for survival in nutrient-limited environments encountered during infection[82,83]. Metabolic flexibility is also critical for persistence in low-oxygen microenvironments generated during pulmonary infection and in drug-resistant biofilms, which become increasingly hypoxic as they mature[84,85]. Understanding the potential parallels in aneuploidy-driven metabolic rewiring in cancer cells, yeast, and *A. fumigatus* therefore has the potential to improve our understanding of this organism's persistence in disease-relevant contexts.

Along with the A1163 reference strain, we isolated Chr4 and Chr7 aneuploids in 4 additional strain backgrounds from both clinical and environmental sources. Interestingly, 2 of the strains that readily gave rise to whole-chromosome aneuploids were isolated during a 4.5 year time course study from the sputum of a person with cystic fibrosis[86]. This finding provides direct evidence that there are isolates present in individuals with *Aspergillus* infections that are capable, under the correct selective conditions, of acquiring and maintaining unstable chromosome duplications. Another recent study, which identified multiple independent segmental duplications in azole-resistant clinical isolates[16], further supports the hypothesis that genome instability is a generalizable and clinically-relevant phenomenon in *A. fumigatus*. Because aneuploidy appears to arise frequently under stress conditions across isolates, coupled with the observation that supernumerary chromosomes are rapidly lost in the absence of drug selection, it is highly plausible that prior sequencing efforts have failed to identify naturally-occurring aneuploids in clinical and environmental samples by culturing isolates under non-selective conditions prior to DNA-sequencing. With the knowledge that aneuploids are highly unstable, future efforts to detect transient karyotype diversity in natural populations can employ isolation strategies that maintain selective pressure. Such efforts will

be important to understand the potential roles of aneuploidy in promoting clinically and agriculturally-
relevant antifungal resistance and persistence under stress.

The majority of research on aneuploidy in pathogenic fungi to this point has focused on
Candida albicans, where azole resistance arises as a consequence of increased copy number of
ERG11 and *TAC1*[10,59]. Recent findings by Khateb et al[16] similarly showed that copy number
duplication of the genomic region encoding *cyp51A* in azole-resistant *A. fumigatus* clinical isolates
corresponds to increased transcription of the drug target gene, suggesting a parallel aneuploidy-
driven resistance mechanism to that observed in *C. albicans*. In contrast to these findings, we
showed that Chr4, 6, and 7 whole-chromosome aneuploidies decrease *in vitro* susceptibility to
voriconazole, and do so without apparent changes in *cyp51* gene expression. Our findings suggest
that important differences may exist in the mechanisms by which aneuploidy can alter azole
susceptibility between different fungal pathogens and between different aneuploid isolates of the
same species.

In this study, we investigated transcriptional changes in 2 independent Chr7 aneuploids
under FK506 treatment and observed both shared and unique changes in gene expression between
the strains. Because these two aneuploids exhibited phenotypic similarities under the tested stress
condition, we focused on those differentially-expressed genes that were shared between them.
Among this shared set of genes was the transcription factor *nscR*, and we demonstrated that
overexpression of *nscR* in the haploid progenitor background largely recapitulates the FK506 stress
response shared between the Chr7 aneuploids. However, the significant number of genes
differentially expressed in one but not the other aneuploid could well underly differences in their
responses to other stresses, and the consequences of these transcriptional differences warrant
further investigation. The broad differences in transcription between two strains of the same genetic
background and karyotype, along with the observation that the transcriptional changes were not
specific to only those genes encoded on the duplicated chromosome, could indicate that there is
some degree of non-specificity in aneuploidy-driven gene expression changes. Such non-specificity
could serve as a bet-hedging strategy in which aneuploidy generates transient transcriptional
diversity in an otherwise-clonal population, potentially increasing the odds of survival upon exposure
to stress. Indeed, stochastic gene expression among clonal populations is known to underlie
differences in competence in *Bacillus subtilis*[87] and heterogeneous expression of stress response
proteins in *S. cerevisiae*[88]. In *S. cerevisiae*, aneuploidy also leads to elevated non-genetic population
heterogeneity and heterogeneous gene expression changes in response to heat shock and nutrient
stresses[89]. These prior observations indicate that transcriptional heterogeneity between genetically
identical aneuploids could have relevant consequences for fitness in environmental and infection
environments, and that studying multiple aneuploids in parallel can aid in determining the particular
aspects of the aneuploid response that are relevant to the stress condition in question.

We found that the normally-silent *nsc* BGC is specifically activated in aneuploids exposed to
FK506, and that overexpression of the *nsc* cluster transcription factor *nscR* results in reversal of the
polarized growth defects caused by FK506. These effects are independent of detectable
neosartoricin production, suggesting that NscR may have functions outside of the production of this
compound, and/or that a neosartoricin intermediate, rather than the final compound, could be the
relevant molecule in the FK506 stress response. Ongoing work seeks to characterize potential
additional functions of NscR in this context and their effects on stress responses and metabolism.
Interestingly, deletion of the gene encoding the calcineurin-activated transcription factor *crzA* is also
reported to increase *nsc* gene expression, but not neosartoricin production, further supporting a role
for this gene cluster in relation to the calcineurin pathway[90]. Several additional studies have also
shown that BGC transcript abundance does not always translate to concomitant metabolite
production[91,92]. The mechanisms underlying non-concordance between transcript and metabolite
production are understudied but can involve post-transcriptional and/or post-translational
modifications. One such example is phosphorylation regulation of AflR, the pathway specific
transcription factor required for sterigmatocystin and aflatoxin production[93]. Several studies have
also demonstrated that transcription factors encoded within one BGC can regulate the expression of
secondary metabolite genes in other BGCs, often those located on other chromosomes. For
example, in *A. fumigatus*, HasA both positively and negatively regulates multiple BGCs[94], in
Penicillium expansum, XanC specifically activates the citrinin BGC[95], and in *Aspergillus nidulans*,
ScpR drives expression of both the *inp* BGC and the asperfuranone BGC[91]. Similarly, in *Fusarium*
graminearum, TRI6 not only activates the trichothecene BGC but also influences genes involved in
the isoprenoid/mevalonate pathway[96]. Our findings that aneuploidy broadly rewires metabolism, and
that changes in secondary metabolite abundance do not necessarily correlate to transcriptional
activation of their BGCs, build on this existing body of literature and provide further evidence for the
complexity of secondary metabolism regulation. Further exploration of cross-talk and cross-
regulatory functions of BGCs in aneuploids is likely to yield valuable insight into the mechanisms
underlying metabolic flexibility and stress responses in this organism.
In the context of intra-population genetic diversity, increased copy number of full
chromosomes may increase the odds of positive selection compared to stable genetic mutations by
flexibly altering large sets of genes and rewiring metabolism in a readily-reversible manner. Given
the variability in stress conditions encountered by *A. fumigatus* both in its environment and during
host interactions, it is conceivable that aneuploidy could facilitate rapid adaptation and persistence in
both ecological and disease contexts through transcriptional changes that can be turned on and off
without the need for permanent genetic change. Overall, our results lay a foundation for future
inquiry into how flexible genomic changes may drive beneficial adaptive responses and increase the
odds of survival in response to stresses such as competing microorganisms, adverse conditions in

the environment and during host infection, and exposure to clinical and agricultural antimicrobial
compounds.
514

Resource Availability

Lead contact

Requests for resources and further information should be directed to, and will be fulfilled by, the lead
contact, Joseph Heitman (heitm001@duke.edu).

Materials availability

All unique materials generated in this study are available without restriction from the lead contact.

Data and code availability

All raw DNA- and RNA-seq data are deposited at NCBI under Bioproject PRJNA1335234,
PRJNA1335235, and PRJNA1448995. Processed sequencing data (variant calls and DESeq
results) are included in the Supplemental Material. All original code generated for this project is
available at: https://github.com/annalehmann/Afum_FK506_resistance and
https://github.com/annalehmann/Afum_Chr7_aneuploids.

Acknowledgments

We thank the Duke University Sequencing and Genomic Technologies Core Facility for
performing the sequencing and Dr. Paul Magwene, Dr. Adrianna San Roman, Dr. Marco Coelho, Dr.
Vikas Yadav, and Anna Mackey for assistance and advice with analyzing the DNA- and RNA-
sequencing data. We also thank Dr. Robert Cramer for providing the *A. fumigatus* strains and
selectable marker plasmids and for offering invaluable insights about strain heterogeneity and
antifungal drug resistance, Dr. Joe Vasselli for sharing a custom 3D-printed microscopy growth
chamber, and the members of the Heitman Lab and the Duke mycology community for constructive
discussion throughout this project. This study was supported by the NIH/NIAID grants
5R01AI172451-03 and 5R01AI170543-04 (JH) and NIH/NIGMS 1R35GM156119 (NPK). JH is also
co-director and fellow of the Canadian Institute for Advanced Research (CIFAR) program Fungal
Kingdom: Threats & Opportunities. Mass spectrometry data were obtained on equipment acquired
with support of UW-Madison and grant 2024-70410-43730 (USDA).

Author Contributions

Conceptualization: A.E.L. and J.H.; Investigation: A.E.L. and E.A.R.; Data analysis: A.E.L. and
E.A.R.; Writing – original draft: A.E.L. and E.A.R.; Writing – review & editing: J.H. and N.P.K.;
Funding acquisition: J.H. and N.P.K.

Declaration of Interests

Joseph Heitman is a member of *Current Biology*'s advisory board.

Supplemental Information

Document S1: Figures S1-S9; Tables S2 and S4.

Table S1. High- and medium-impact coding region variants in all FK506-adapted isolates.

Table S3. Gene expression changes in Chr7 aneuploids +/- FK506.

Table S5. Sequence variants between our A1163 laboratory strain and the published reference genome.

Materials and Methods

Strains and growth conditions

The A1163 reference strain served as the haploid wild type in all genetic and transcriptomic comparisons presented in this study. Additional clinical and environmental isolates described in Table S2 and previously published by Lofrgren et al[57] were also screened for aneuploidy in response to FK506. Strains were cultured on 1.5% agar Aspergillus Minimal Medium (AMM), which was prepared as described by Hill and Kafer[97]. All FK506 experiments on agar plates and in liquid culture were performed in AMM with FK506 at a concentration of 1 µg/mL or 0.5 µg/mL. Conidia were grown for experiments on AMM or AMM+FK506 for 3 days at 37°C. Unstable aneuploids were maintained on +FK506 plates to ensure retention of the ploidy differences. Conidia were collected from plates by flooding the surface of the plate with 0.01% Tween 80, manually dislodging conidia with a cell scraper, and filtering through Miracloth (Millipore Sigma). Conidia were counted with a hemacytometer.

DNA extraction and sequencing

Biomass for DNA extraction was obtained by inoculating 5 mL AMM or AMM+FK506 with conidia at a density of 5×10^5 conidia/mL and incubating at 37°C rotating at 70 rpm on a roller drum for 16-24 hours. Biomass was collected and washed 1x with diH_2O, then frozen at -80°C for >1 hour and lyophilized until dry. DNA was extracted following a modified MasterPure (Biosearch Technologies) procedure. First, lyophilized tissue was bead-beaten with 2 mm glass beads for 1 minute. The remainder of the procedure was carried out according to the manufacturer's instructions. Nucleic acid pellets were resuspended in nuclease-free water and quality was assessed by gel electrophoresis. DNA concentration was quantified with a Qubit fluorometer using the DNA broad-range kit (Thermo Fisher Scientific).

DNA sequencing was carried out by the Duke University Sequencing and Genomic
Technologies Core Facility. Libraries were prepared using the Kapa HyperPrep kit (Kapa
Bioscience). Pooled libraries were sequenced on an Illumina Novaseq X Plus instrument in 150 bp
paired-end reads.

DNA-seq data analysis

For sequence variant identification, DNA-seq reads were aligned to the A1163 reference
genome[60] with snippy[98], using the default parameters. Resulting vcf files served as input for snpEff[99],
which compares variants to a preconfigured A1163 database and predicts variant effects on the
coding transcript (low, medium, or high). For the FK506-adapted isolates, variants in coding regions
only were identified with the flags –no-downstream, –no-upstream, and –no-intergenic. Unique
variants were determined by filtering out any that were shared with our laboratory A1163 parental
strain (Table S5).

Ploidy was determined by comparing coverage depth along 500 bp windows to genome-wide
normalized read depth. Reads were aligned to the chromosome-level CEA10 reference assembly[100]
with minimap2[101], using settings for paired-end short reads. Aligned reads were filtered by a
threshold of q30 with samtools[102]. Mosdepth[103] was used to calculate read depth over 500 bp
windows. The resulting bed files were analyzed in R to compare depth in each window to genome-
wide normalized coverage. Data were visualized with ggplot2[104].

Phenotypic assays

FK506 spot growth and imaging

10^5 conidia were spotted on AMM + 1 μg/mL FK506 and incubated at 37°C for 2 days.
Images of hyphal growth at the colony edge were acquired with an AxioScop 2 fluorescence
microscope with 4x objective (Zeiss). Experiments were performed in at least 3 biologically
independent replicates.

Imaging of submerged hyphal growth

100μL conidia at a concentration of 10^3 conidia/mL in AMM ± 1 μg/mL FK506 were
inoculated in a custom 3D-printed growth chamber[105] encased between 2 transparent slides. Growth
chambers were incubated at 37°C for 24 hours. Hyphal growth images were acquired on an
AxioScop 2 fluorescence microscope with 10x and 20x objectives (Zeiss). Experiments were
performed in biological triplicate.

Transcriptomics

RNA extraction and sequencing

Biomass for RNA extraction was obtained from 10^6 conidia/mL cultures in 25 mL AMM or
AMM+FK506 grown for 16 hours at 37° with shaking at 200 RPM. Biomass was harvested, washed
1x with sterile diH$_2$O, frozen at -80°C for >1 hour, and lyophilized until dry. Lyophilized tissue was
bead-beaten for 1 minute with 2 mm glass beads and RNA was extracted with the Qiagen RNeasy
Plant Mini Kit according to the manufacturer's instructions (Qiagen). RNA concentration was
measured a Qubit fluorometer using the RNA broad-range kit (Thermo Fisher Scientific).

RNA was sequenced by the Duke University Sequencing and Genomic Technologies Core
Facility. Poly-A enriched libraries were prepared from samples with RIN ≥ 7 using the Kapa mRNA-
seq HyperPrep kit (Kapa Bioscience). Pooled libraries were sequenced on an Illumina Novaseq X
Plus instrument in 150 bp paired-end reads.

RNA-seq data analysis

Illumina universal adaptors were trimmed from paired reads with CutAdapt[106] (v4.9).
Trimmed reads were then aligned to the A1163 reference assembly[60] with HISAT2[107] (v2.2.1).
Counts per gene were quantified with featureCounts[108] (v2.0.6). Exploratory data analysis and
differential expression analysis were performed with DESeq2[109] (v1.44.0). Prior to analysis,
transcript counts were normalized by transcripts per million (TPM) within each replicate and genes
with at least 1 TPM in 1 sample were retained. Clustering and principal component analyses were
performed using vst normalized values for the top 1000 most variable genes generated by DESeq2.
Differential expression analysis was carried out in DESeq2 with alpha = 0.05 and a log$_2$FC cutoff of
+/- 1 was applied to identify DEGs from the resulting expression data. The R packages
EnhancedVolcano (v1.22.0) (https://github.com/kevinblighe/EnhancedVolcano) and pheatmap
(v1.0.13) (https://github.com/raivokolde/pheatmap) were used for data visualization.

nscR *overexpression strain construction*

The *nscR*^{OE} strain was generated in the A1163 background by ectopic integration of *nscR*
(AFUB_086680) driven by an *A. fumigatus gpdA* promoter amplified from the Af293 reference strain
(Fungal Genetics Stock Center A1100). The *gpdA* promoter (*$_p$gpdA*) was selected as the 996 bp
immediately upstream of the *gpdA* (Afu5g01970) open reading frame using primers AEL68 and
AEL69. The promoter was fused to the *nscR* open reading frame and the 504 bases immediately 3'
of the stop codon, which was amplified from A1163 gDNA with primers AEL70 and AEL71, and the
hygromycin resistance marker *hygR*. The *$_p$gpdA*^{AF}-*nscR* overexpression construct fused to *hygR*
selectable marker was amplified with primers AEL75 and AEL76. The resulting product was used to
transform A1163 protoplasts as described previously[110,111]. Transformants were selected on
osmotically-stabilized minimal medium (AMM + 1.2 M sorbitol) containing hygromycin. Successful
transformants were single spored, and genomic integration of the construct was verified by PCR and

expression was measured by RT-qPCR as described below, with RNA obtained from WT and
$nscR^{OE}$ under the same conditions ± FK506 used for RNA-seq. All primer sequences used are listed
in Table S5.

Reverse transcriptase quantitative PCR (RT-qPCR)

2.5 to 5 µg of RNA was DNase treated using the TURBO DNA-free kit (Invitrogen) following
the manufacturer's instructions. Post-DNase treatment RNA integrity was verified by running a
sample of the RNA on an agarose gel. cDNA synthesis was carried out using 500 ng of DNase-
treated RNA as described previously[104]. RT-qPCR was run on a QuantStudio 3 or QuantStudio 6 Pro
system (Thermo Fisher Scientific). Gene expression was normalized to *actA* and *tubA* in all
experiments. Fold change values were calculated by the $2^{-\Delta\Delta Ct}$ method[112].

Metabolomic studies

Four biological replicates of each strain were cultured at 10^6 conidia/mL in 25 mL of GMM
medium supplemented with 1 µg FK506/mL and incubated for two days at 37 °C with shaking at 200
rpm. After the fermentation period, each fungal culture and four medium replicates (considered as
blanks) were extracted by maceration with 25 mL of EtOAc/MeOH/AcOH (89:10:1) overnight, with
shaking at 120 rpm. The organic phase was then separated and evaporated to dryness under
reduced pressure. The resulting organic extracts were dissolved in MeOH at a concentration of 50
mg/mL, and serial dilutions (1:10 and 1:50) were prepared to obtain solutions at 0.1 mg/mL for LC-
MS analysis. Then, a 5 µL aliquot of each sample was analyzed by UPLC coupled to an Orbitrap
Exploris™ 240 mass spectrometer, acquiring both MS^1 and MS/MS spectra, using an Acquity BEH-
C18 column (2.1 mm × 100 mm, 1.7 µm).

The chromatographic method consisted of a one minute isocratic elution of 10% MeCN in
H_2O containing 0.05% formic acid, followed by a linear gradient from 10% to 100% MeCN over 20
minutes, and a 2.5-minute wash with 100% MeCN. The flow rate was maintained at 0.3 mL/min
throughout the run, with alternating positive and negative ion detection over an *m/z* range of 120–
1500 for MS/MS acquisition. Precursor ions were fragmented using higher-energy collisional
dissociation (HCD) with dynamic exclusion set to 3 s.

Once the LC-MS/MS data were acquired, the raw files generated from these analyses were
converted to .mzML format using MSConvert[113], applying the peak-picking filter for both MS^1 and
MS^2. Spectra were processed in MZmine[114] (v4.7.8). Mass detection was performed using the
lowest signal factor, with a noise threshold set at 5.0 for MS^1 and 2.5 for MS^2. A chromatogram was
constructed using the Chromatogram Builder algorithm in positive polarity mode, with parameters set
to a minimum of 4 consecutive scans, a minimum intensity of 1×10^4 for consecutive scans, a
minimum absolute height of 5×10^4, and a mass tolerance of 10 ppm.

The resulting chromatogram features were then processed using Savitzky-Golay smoothing
(5 scans retention time width), the Local Minimum Feature Resolver, a 13C isotope filter, the Join
Aligner, Feature Finder, and duplicate peak filter. From this processed feature list, features detected
in the blank replicates were manually removed, yielding a total of 4,495 features. This curated
feature list was used for principal component analysis (PCA) with Pareto scaling and missing value
imputation (set at 1/5 of the minimum value), for pairwise comparisons using unpaired Volcano Plots,
and for molecular network construction following the Feature-based Molecular Networking workflow
with default parameters[115].

The abundance of secondary metabolites was represented using the area under the curve of
the target feature (included in chromatograms, bar graphs, and heatmaps), based on LC-MS data
acquired in either negative or positive ionization mode. PCA plots, Volcano Plots, chromatograms,
bar graphs, and heatmaps were generated with Prism 10.6.0. For the heatmap, z-scores were
calculated based on the average abundance of each metabolite across the different strains.
Metabolites that were not detected in any strain were labeled as undetected. The molecular network
was visualized using Cytoscape[116] (v3.10.3). The identities of fumitremorgin C, fumagillin,
pyripyropene A, fumiquinazoline F, neosartoricin, and FK506 were confirmed by comparison of their
chromatographic and spectrometric data with authentic standards. The identity of pseurotin A was
supported by matching the fragmentation spectrum of the detected molecular ion to a high-quality
gold-standard   reference   deposited   in   the   GNPS   database
(https://gnps.ucsd.edu/ProteoSAFe/gnpslibraryspectrum.jsp?SpectrumID=CCMSLIB00011906568#
%7B%7D). The ergosterol-type triterpenoids class was supported by the predicted formula and by
the comparison of the fragmentation pattern with those reported by Lucie Ory et al[117].

Antifungal susceptibility tests

Broth dilution MIC assays were performed according to the established standard CLSI
procedure[118]. MIC tests were performed in 3 to 4 biological replicates.

Quantification of cyp51A *and* cyp51B *transcript abundances*

RNA was obtained from two distinct culture conditions to provide for potential condition-
specific, rather than drug-specific, effects on transcript abundances. In the first condition, 2.5×10^5
conidia were inoculated in 5 mL RPMI medium and incubated at 37°C rotating at 70 rpm. At 22
hours, medium was removed and replaced with 5 mL RPMI + 0.2 µg/mL voriconazole (0.1 µL of a 10
mg/mL stock in DMSO) or 0.1 µL DMSO. Cultures were incubated under the same conditions for an
additional 2 hours. For the second condition, 10^6 conidia/mL were inoculated in AMM with 0.1 µg/mL
voriconazole or DMSO and incubated at 37°C with shaking at 200 rpm. Biomass was collected at 24
hours. For both conditions, RNA was extracted and cDNA synthesized as described above. qPCR

was carried out as described for the *nscR^{OE}* strain, and *cyp51A* and *cyp51B* transcript abundances

was calculated relative to *actA*. All qPCR primers used in this study are listed in Table S4.

References

1. Knouse, K.A. et al. (2017). Aneuploidy in cancer: seq-ing answers to old questions. Annu. Rev.
Cancer Biol. 1, 335–354. DOI: 10.1146/annurev-cancerbio-042616-072231

2. Ben-David, U., and Amon, A. (2020). Context is everything: aneuploidy in cancer. Nat. Rev.
Genet. 21, 44–62. DOI: 10.1038/s41576-019-0171-x

3. Torres, E.M. et al. (2008). Aneuploidy: cells losing their balance. Genetics 179, 737–746. DOI:
10.1534/genetics.108.090878

4. Gilchrist, C., and Stelkens, R. (2019). Aneuploidy in yeast: segregation error or adaptation
mechanism? Yeast 36, 525–539. DOI: 10.1002/yea.3427

5. Sheltzer, J.M. et al. (2011). Aneuploidy drives genomic instability in yeast. Science 333, 1026–
1030. DOI: 10.1126/science.1206412

6. Sunshine, A.B. et al. (2015). The fitness consequences of aneuploidy are driven by condition-
dependent gene effects. PLOS Biol. 13, e1002155. DOI: 10.1371/journal.pbio.1002155

7. Yona, A.H. et al. (2012). Chromosomal duplication is a transient evolutionary solution to stress.
Proceed. Nat. Acad. Sci. USA 109, 21010–21015. DOI: 10.1073/pnas.1211150109

8. Gresham, D., Desai, M.M., Tucker, C.M., Jenq, H.T., Pai, D.A., Ward, A., DeSevo, C.G.,
Botstein, D., and Dunham, M.J. (2008). The repertoire and dynamics of evolutionary
adaptations to controlled nutrient-limited environments in yeast. PLOS Genet. 4, e1000303.
DOI: 10.1371/journal.pgen.1000303

9. Todd, R.T. et al. (2017). Ploidy variation in fungi: polyploidy, aneuploidy, and genome evolution.
Microbiol. Spectr. 5, DOI: 10.1128/microbiolspec.funk-0051–2016

10. Selmecki, A. et al. (2006). Aneuploidy and isochromosome formation in drug-resistant *Candida*
albicans. Science 313, 367–370. DOI: 10.1126/science.1128242

11. Sun, L. et al. (2023). Aneuploidy mediates rapid adaptation to a subinhibitory amount of
fluconazole in *Candida albicans*. Microbiol. Spectr. 11, e03016-22. DOI:
10.1128/spectrum.03016-22

12. Perepnikhatka, V. et al. (1999). Specific chromosome alterations in fluconazole-resistant
mutants of *Candida albicans*. J. Bacteriol. 181, 4041–4049. DOI: 10.1128/jb.181.13.4041-
4049.1999

13. Yang, F. et al. (2021). Adaptation to fluconazole via aneuploidy enables cross-adaptation to
Amphotericin B and Flucytosine in *Cryptococcus neoformans*. Microbiol. Spectr. 9, e00723-21.
DOI: 10.1128/Spectrum.00723-21

14. Zhang, Z. et al. (2024). Aneuploidy underlies brefeldin A-induced antifungal drug resistance in
Cryptococcus neoformans. Front. Cell. Infect. Microbiol. 14. DOI: 10.3389/fcimb.2024.1397724

15. Barda, O. et al. (2023). Aneuploidy formation in the filamentous fungus *Aspergillus flavus* in
response to azole stress. Microbiol. Spectr. 11, e04339-22. DOI: 10.1128/spectrum.04339-22

16. Khateb, A. et al. (2023). Aneuploidy Is associated with azole resistance in *Aspergillus fumigatus*.
Antimicrob. Agents Chemother. 67, e01253-22. DOI: 10.1128/aac.01253-22

17. Denning, D.W. Global incidence and mortality of severe fungal disease. Lancet Infect. Dis. 24,
428–438. DOI: 10.1016/S1473-3099(23)00692-8

18. Nguyen, K.-H. et al. (2013). UV-protectant metabolites from lichens and their symbiotic partners.
Nat. Prod. Rep. 30, 1490–1508. DOI: 10.1039/C3NP70064J

19. Brakhage, A.A., and Liebmann, B. (2005). *Aspergillus fumigatus* conidial pigment and cAMP
signal transduction: significance for virulence. Med. Mycol. 43 Suppl 1, S75-82. DOI:
10.1080/13693780400028967
20. Berkova, N. et al. (2006). *Aspergillus fumigatus* conidia inhibit tumour necrosis factor- or
staurosporine-induced apoptosis in epithelial cells. Int. Immunol. 18, 139–150. DOI:
10.1093/intimm/dxh356
21. Gardiner, D.M. et al. (2005). The epipolythiodioxopiperazine (ETP) class of fungal toxins:
distribution, mode of action, functions and biosynthesis. Microbiology 151, 1021–1032. DOI:
10.1099/mic.0.27847-0
22. Bok, J.W. et al. (2006). GliZ, a transcriptional regulator of gliotoxin biosynthesis, contributes to
Aspergillus fumigatus virulence. Infect. Immun. *74*, 6761–6768. DOI: 10.1128/iai.00780-06
23. Pardo, J. et al. (2006). The mitochondrial protein Bak is pivotal for gliotoxin-induced apoptosis
and a critical host factor of *Aspergillus fumigatus* virulence in mice. J. Cell Biol. 174, 509–519.
DOI: 10.1083/jcb.200604044
24. Spikes, S. et al. (2008). Gliotoxin production in *Aspergillus fumigatus* contributes to host-specific
differences in virulence. J. Infect. Dis. 197, 479–486. DOI: 10.1086/525044
25. Cramer, R.A. et al. (2011). Immune responses against *Aspergillus fumigatus*: what have we
learned? Curr. Opin. Infect. Dis. 24, 315–322. DOI: 10.1097/QCO.0b013e328348b159
26. Raffa, N., and Keller, N.P. (2019). A call to arms: Mustering secondary metabolites for success
and survival of an opportunistic pathogen. PLOS Pathog. 15, e1007606. DOI:
10.1371/journal.ppat.1007606
27. Lim, F.Y. et al. (2012). Chapter fifteen - toward awakening cryptic secondary metabolite gene
clusters in filamentous fungi. Methods in Enzymology Natural Product Biosynthesis by
Microorganisms and Plants, Part C., D. A. Hopwood, ed. (Academic Press), pp. 303–324. DOI:
10.1016/B978-0-12-404634-4.00015-2
28. Krespach, M.K.C. et al. (2023). *Streptomyces* polyketides mediate bacteria–fungi interactions
across soil environments. Nat. Microbiol. *8*, 1348–1361. DOI: 10.1038/s41564-023-01382-2
29. König, C.C. et al. (2013). Bacterium induces cryptic meroterpenoid pathway in the pathogenic
fungus *Aspergillus fumigatus*. ChemBioChem 14, 938–942. DOI: 10.1002/cbic.201300070
30. Nützman, H.-W. et al. (2011). Bacteria-induced natural product formation in the fungus
Aspergillus nidulans requires Saga/Ada-mediated histone acetylation. Proceed. Nat. Acad. Sci.
USA 108, 14282–14287. DOI: 10.1073/pnas.1103523108
31. Zuck, K.M. et al. (2011). Induced production of N-formyl alkaloids from *Aspergillus fumigatus* by
co-culture with *Streptomyces peucetius*. J. Nat. Prod. 74, 1653–1657. DOI: 10.1021/np200255f
32. Stroe, M.C. et al. (2020). Targeted induction of a silent fungal gene cluster encoding the
bacteria-specific germination inhibitor fumigermin. eLife 9, e52541. DOI: 10.7554/eLife.52541
33. Schroeckh, V. et al. (2009). Intimate bacterial–fungal interaction triggers biosynthesis of
archetypal polyketides in *Aspergillus nidulans*. Proceed. Nat. Acad. Sci. USA 106, 14558–
14563. DOI: 10.1073/pnas.0901870106
34. Losada, L. et al. (2009). Effect of competition on the production and activity of secondary
metabolites in *Aspergillus* species. Med. Mycol. 47, S88–S96. DOI:
10.1080/13693780802409542
35. O'Hanlon, K.A. et al. (2011). Targeted disruption of nonribosomal peptide synthetase pes3
augments the virulence of *Aspergillus fumigatus*. Infect. Immun. 79, 3978–3992. DOI:
10.1128/iai.00192-11

36. Yin, W.-B. et al. (2013). A nonribosomal peptide synthetase-derived iron(III) complex from the
pathogenic fungus *Aspergillus fumigatus*. J. Am. Chem. Soc. 135, 2064–2067. DOI:
10.1021/ja311145n
37. Tsunematsu, Y. et al. (2013). Distinct mechanisms for spiro-carbon formation reveal biosynthetic
pathway crosstalk. Nat. Chem. Biol. 9, 818–825. DOI: 10.1038/nchembio.1366
38. O'Keeffe, G. et al. (2014). RNA-seq reveals the pan-transcriptomic impact of attenuating the
gliotoxin self-protection mechanism in *Aspergillus fumigatus*. BMC Genom. 15, 894. DOI:
10.1186/1471-2164-15-894
39. Calo, S. et al. (2014). Antifungal drug resistance evoked via RNAi-dependent epimutations.
Nature 513, 555–558. DOI: 10.1038/nature13575

40. Calo, S. et al/ (2017). A non-canonical RNA degradation pathway suppresses RNAi-dependent
epimutations in the human fungal pathogen *Mucor circinelloides*. PLOS Genet. 13, e1006686.
DOI: 10.1371/journal.pgen.1006686

41. Priest, S.J. et al. (2022). Uncontrolled transposition following RNAi loss causes hypermutation
and antifungal drug resistance in clinical isolates of *Cryptococcus neoformans*. Nat. Microbiol.
7, 1239–1251. DOI: 10.1038/s41564-022-01183-z
42. Yadav, V., et al. (2025). Calcineurin controls the cytokinesis machinery during thermal stress in
Cryptococcus deneoformans. Proceed. Nat. Acad. Sci. USA 122, e2503751122.
DOI: 10.1073/pnas.2503751122
43. Huang, J. et al. (2024). Distinct evolutionary trajectories following loss of RNA interference in
Cryptococcus neoformans. Proceed. Nat. Acad. Sci. USA 121, e2416656121. DOI:
10.1073/pnas.2416656121
44. Liu, J. et al. (1991). Calcineurin is a common target of cyclophilin-cyclosporin A and FKBP-
FK506 complexes. Cell 66, 807–815. DOI: 10.1016/0092-8674(91)90124-H
45. Ho, S. et al. (1996). The mechanism of action of cyclosporin A and FK506. Clin. Immunol.
Immunopathol. 80, S40–S45. DOI: 10.1006/clin.1996.0140
46. Hemenway, C.S., and Heitman, J. (1999). Calcineurin. Cell Biochem. Biophys. 30, 115–151.
DOI: 10.1007/BF02737887
47. Juvvadi, P.R. et al. (2008). Calcineurin localizes to the hyphal septum in *Aspergillus fumigatus*:
implications for septum formation and conidiophore development. Eukaryot. Cell 7, 1606–1610.
DOI: 10.1128/ec.00200-08

48. Steinbach, W.J. et al. (2006). Calcineurin controls growth, morphology, and pathogenicity in
Aspergillus fumigatus. Eukaryot. Cell 5, 1091–1103. DOI: 10.1128/ec.00139-06

49. Cramer, R.A. et al. (2008). Calcineurin target CrzA regulates conidial germination, hyphal
growth, and pathogenesis of *Aspergillus fumigatus*. Eukaryot. Cell 7, 1085–1097. DOI:
10.1128/ec.00086-08
50. Juvvadi, P.R. et al. (2011). Localization and activity of the calcineurin catalytic and regulatory
subunit complex at the septum is essential for hyphal elongation and proper septation in
Aspergillus fumigatus. Mol. Microbiol. 82, 1235–1259. DOI: 10.1111/j.1365-2958.2011.07886.x

51. Steinbach, W.J. et al. (2004). In vitro interactions between antifungals and immunosuppressants
against *Aspergillus fumigatus*. Antimicrob. Agents Chemother. 48, 1664–1669. DOI:
10.1128/aac.48.5.1664-1669.2004

52. Soriani, F.M. et al. (2008). Functional characterization of the *Aspergillus fumigatus CRZ1*
homologue, CrzA. Mol. Microbiol. 67, 1274–1291. DOI: 10.1111/j.1365-2958.2008.06122.x

53. Ries, L.N.A. et al. (2017). The *Aspergillus fumigatus* CrzA transcription factor activates chitin
synthase gene expression during the caspofungin paradoxical effect. mBio 8. DOI:
10.1128/mbio.00705-17

54. Ferreira, M.E. da S. et al. (2007). Functional characterization of the *Aspergillus fumigatus*
calcineurin. Fungal Genet. Biol. 44, 219–230. DOI: 10.1016/j.fgb.2006.08.004

55. Ellett, F. et al. (2017). Neutrophil interactions stimulate evasive hyphal branching by *Aspergillus*
fumigatus. PLOS Pathog. 13, e1006154. DOI: 10.1371/journal.ppat.1006154

56. Schoen, T.J. et al. (2023). *Aspergillus fumigatus* transcription factor ZfpA regulates hyphal
development and alters susceptibility to antifungals and neutrophil killing during infection.
PLOS Pathog. 19, e1011152. DOI: 10.1371/journal.ppat.1011152

57. Lofgren, L.A. et al. (2022). The pan-genome of *Aspergillus fumigatus* provides a high-resolution
view of its population structure revealing high levels of lineage-specific diversity driven by
recombination. PLOS Biol. 20, e3001890. DOI: 10.1371/journal.pbio.3001890

58. Son, Y.-E. et al. (2026). Epimutations driven by RNAi or heterochromatin evoke transient
antimicrobial drug resistance in pathogenic Mucor fungi. PLOS Biol. 24, e3003598. DOI:
10.1371/journal.pbio.3003598

59. Selmecki, A. et al. (2008). An isochromosome confers drug resistance in vivo by amplification of
two genes, *ERG11* and *TAC1.* Mol. Microbiol. 68, 624–641. DOI: 10.1111/j.1365-
2958.2008.06176.x

60. Fedorova, N.D. et al. (2008). Genomic islands in the pathogenic filamentous fungus *Aspergillus*
fumigatus. PLOS Genet. 4, e1000046. DOI: 10.1371/journal.pgen.1000046

61. Chooi, Y.-H. et al. (2013). Genome mining of a prenylated and immunosuppressive polyketide
from pathogenic fungi. Org. Lett. 15, 780–783. DOI: 10.1021/ol303435y
62. Brakhage, A.A., and Schroeckh, V. (2011). Fungal secondary metabolites – strategies to activate
silent gene clusters. Fungal Genet. Biol. 48, 15–22. DOI: 10.1016/j.fgb.2010.04.004
63. Bok, J.W., and Keller, N.P. (2004). LaeA, a regulator of secondary metabolism in *Aspergillus*
spp. Eukaryot. Cell 3, 527–535. DOI: 10.1128/ec.3.2.527-535.2004
64. Mózsik, L. et al. (2022). Transcriptional activation of biosynthetic gene clusters in filamentous
fungi. Front. Bioeng. Biotechnol. 10. DOI: 10.3389/fbioe.2022.901037
65. Seo, H.-W. et al. (2024). A timeline of biosynthetic gene cluster discovery in *Aspergillus*
fumigatus: from characterization to future perspectives. J. Fungi 10, 266. DOI:
10.3390/jof10040266
66. Buied, A. et al. (2013). High-level expression of *cyp51B* in azole-resistant clinical *Aspergillus*
fumigatus isolates. J. Antimicrob. Chemother. *68*, 512–514. DOI: 10.1093/jac/dks451

67. Verweij, P.E. et al. (2007). Multiple-triazole–resistant aspergillosis. N. Engl. J. Med. 356, 1481–
1483. DOI: 10.1056/NEJMc061720

68.   Hurst, S.F. et al. (2017). Isolation of azole-resistant *Aspergillus fumigatus* from the environment
in the south-eastern USA. J. Antimicrob. Chemother. 72, 2443–2446. DOI: 10.1093/jac/dkx168

69.   Howard, S.J. et al. (2009). Frequency and evolution of azole resistance in *Aspergillus fumigatus*
associated   with   treatment   failure.   Emerg.   Infect.   Dis.   15,   1068-1076.   DOI:
10.3201/eid1507.090043

70.   Snelders, E. et al. (2009). Possible environmental origin of resistance of *Aspergillus fumigatus* to
medical triazoles. Appl. Environ. Microbiol. 75, 4053–4057. DOI: 10.1128/AEM.00231-09

71.   Snelders, E. et al. (2008). Emergence of azole resistance in *Aspergillus fumigatus* and spread of
a single resistance mechanism. PLOS Med. 5, e219. DOI: 10.1371/journal.pmed.0050219

72.   Alvarez-Moreno, C. et al. (2017). Azole-resistant *Aspergillus fumigatus* harboring TR34/L98H,
TR46/Y121F/T289A and TR53 mutations related to flower fields in Colombia. Sci. Rep. 7,
45631. DOI: 10.1038/srep45631

73.   Mellado, E. et al. (2007). A new *Aspergillus fumigatus* resistance mechanism conferring in vitro
cross-resistance to azole antifungals involves a combination of *cyp51A* alterations. Antimicrob.
Agents Chemother. 51, 1897–1904. DOI: 10.1128/aac.01092-06

74.   Hanahan, D., and Weinberg, R.A. (2011). Hallmarks of cancer: The next generation. Cell 144,
646–674. DOI: 10.1016/j.cell.2011.02.013

75.   Warburg,   O.   (1956).   On   the   origin   of   cancer   cells.   Science   123,   309–314.   DOI:
10.1126/science.123.3191.309

76.   Warburg, O. (1956). On respiratory impairment in cancer cells. Science 124, 269–270. DOI:
10.1126/science.124.3215.269

77.   Liberti, M.V., and Locasale, J.W. (2016). The Warburg Effect: How does it benefit cancer cells?
Trends Biochem. Sci. 41, 211–218. DOI: 10.1016/j.tibs.2015.12.001

78.   Bartrons, R., and Caro, J. (2007). Hypoxia, glucose metabolism and the Warburg's effect. J.
Bioenerg. Biomembr. 39, 223–229. DOI: 10.1007/s10863-007-9080-3

79.   Graham, N.A. et al. (2017). Recurrent patterns of DNA copy number alterations in tumors reflect
metabolic selection pressures. Mol. Syst. Biol. *13*, MSB167159. DOI: 10.15252/msb.20167159

80.   Williams, B.R. et al. (2008). Aneuploidy affects proliferation and spontaneous immortalization in
mammalian cells. Science 322, 703–709. DOI: 10.1126/science.1160058

81.   Kadowaki, M. et al. (2017). Chromosomal aneuploidy improves the brewing characteristics of
sake yeast. Appl. Environ. Microbiol. 83, e01620-17. DOI: 10.1128/AEM.01620-17

82.   Beattie, S.R. et al. (2017). Filamentous fungal carbon catabolite repression supports metabolic
plasticity and stress responses essential for disease progression. PLOS Pathog. 13, e1006340.
DOI: 10.1371/journal.ppat.1006340

83.   Krappmann, S. et al. (2004). The *Aspergillus fumigatus* transcriptional activator CpcA
contributes significantly to the virulence of this fungal pathogen. Mol. Microbiol. 52, 785–799.
DOI: 10.1111/j.1365-2958.2004.04015.x

84. Kowalski, C.H. et al. (2020). Fungal biofilm architecture produces hypoxic microenvironments
that drive antifungal resistance. Proceed. Nat. Acad. Sci. 117, 22473–22483. DOI:
10.1073/pnas.2003700117

85. Puerner, C. et al. (2025). Transcriptional and metabolic modeling analyses of developing
Aspergillus fumigatus biofilms reveal metabolic shifts required for biofilm maturation. mSphere
10, e00752-25. DOI: 10.1128/msphere.00752-25

86. Ross, B.S. et al. (2021). *Aspergillus fumigatus* in-host HOG pathway mutation for cystic fibrosis
lung microenvironment persistence. mBio 12. DOI: 10.1128/mbio.02153-21

87. Maamar, H. et al. (2007). Noise in gene expression determines cell fate in *Bacillus subtilis*.
Science 317, 526–529. DOI: 10.1126/science.1140818

88. Bar-Even, A. et al. (2006). Noise in protein expression scales with natural protein abundance.
Nat. Genet. 38, 636–643. DOI: 10.1038/ng1807

89. Beach, R.R. et al. (2017). Aneuploidy causes non-genetic individuality. Cell 169, 229-242.e21.
DOI: 10.1016/j.cell.2017.03.021

90. Colabardini, A.C., Wang, F., Dong, Z., Pardeshi, L., Rocha, M.C., Costa, J.H., dos Reis, T.F.,
Brown, A., Jaber, Q.Z., Fridman, M., et al. (2022). Heterogeneity in the transcriptional response
of the human pathogen *Aspergillus fumigatus* to the antifungal agent caspofungin. Genetics
220, iyab183. https://doi.org/10.1093/genetics/iyab183.

91. Bergmann, S. et al. (2010). Activation of a silent fungal polyketide biosynthesis pathway through
regulatory cross talk with a cryptic nonribosomal peptide synthetase gene cluster. Appl.
Environ. Microbiol. 76, 8143–8149. DOI: 10.1128/AEM.00683-10

92. Tannous, J. et al. (2018). Fungal attack and host defence pathways unveiled in near-avirulent
interactions of *Penicillium expansum creA* mutants on apples. Mol. Plant Pathol. 19, 2635–
2650. DOI: 10.1111/mpp.12734

93. Shimizu, K. et al. (2003). Pka, Ras and RGS protein interactions regulate activity of AflR, a
Zn(II)2Cys6 transcription factor in *Aspergillus nidulans*. Genetics 165, 1095–1104. DOI:
10.1093/genetics/165.3.1095

94. Wiemann, P. et al. (2014). Perturbations in small molecule synthesis uncovers an iron-
responsive secondary metabolite network in *Aspergillus fumigatus*. Front. Microbiol. 5. DOI:
10.3389/fmicb.2014.00530

95. Wang, W. et al. (2021). Transcription factor repurposing offers insights into evolution of
biosynthetic gene cluster regulation. mBio 12. DOI: 10.1128/mbio.01399-21

96. Shostak, K. et al. (2020). Activation of biosynthetic gene clusters by the global transcriptional
regulator TRI6 in *Fusarium graminearum*. Mol. Microbiol. 114, 664–680. DOI:
10.1111/mmi.14575

97. Hill, T., and Kafer, E. (2001). Improved protocols for *Aspergillus* minimal medium: trace element
and minimal medium salt stock solutions. Fungal Genet. Rep. 48, 20–21. DOI: 10.4148/1941-
4765.1173
98. Seemann, T. (2024). https://github.com/tseemann/snippy.
99. Cingolani, P. et al. (2012). A program for annotating and predicting the effects of single
nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain
w^{1118}□; iso-2; iso-3. Fly 6, 80–92. DOI: 10.4161/fly.19695
100. Bowyer, P. et al. (2022). Telomere-to-telomere genome sequence of the model mould pathogen
Aspergillus fumigatus. Nat. Commun. 13, 5394. DOI: 10.1038/s41467-022-32924-7
101. Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics 34, 3094–
3100. DOI: 10.1093/bioinformatics/bty191
102. Li, H. et al. (2009). The sequence alignment/map format and SAMtools. Bioinformatics 25,
2078–2079. DOI: 10.1093/bioinformatics/btp352
103. Pedersen, B.S., and Quinlan, A.R. (2018). Mosdepth: quick coverage calculation for genomes
and exomes. Bioinformatics 34, 867–868. DOI: 10.1093/bioinformatics/btx699
104. Wickham, H. (2016). ggplot2: elegant graphics for data analysis (Springer-Verlag).
105. Breuer, M.R. et al. (2022). The antidepressant sertraline induces the formation of supersized
lipid droplets in the human pathogen *Cryptococcus neoformans*. J. Fungi 8, 642. DOI:
10.3390/jof8060642
106. Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnet.journal 17, 10–12. DOI: 10.14806/ej.17.1.200

107. Kim, D. et al. (2019). Graph-based genome alignment and genotyping with HISAT2 and HISAT-
genotype. Nat. Biotechnol. *37*, 907-915. https://doi.org/10.1038/s41587-019-0201-4.
108. Liao, Y. et al. (2014). featureCounts: an efficient general purpose program for assigning
sequence   reads   to   genomic   features.   Bioinformatics   30,   923–930.   DOI:
10.1093/bioinformatics/btt656

109. Love, M.I. et al. (2014). Moderated estimation of fold change and dispersion for RNA-seq data
with DESeq2. Genome Biol. 15, 550. DOI: 10.1186/s13059-014-0550-8

110. Kerkaert, J.D. et al. (2022). An alanine aminotransferase is required for biofilm-specific
resistance of Aspergillus fumigatus to echinocandin treatment. mBio 13, e02933-21. DOI:
10.1128/mbio.02933-21

111. Zhao, C. et al. (2019). High-throughput gene replacement in *Aspergillus fumigatus*. Curr.
Protoc. Microbiol. 54, e88. DOI: 10.1002/cpmc.88

112. Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-
time   quantitative   PCR   and   the   2−ΔΔCT   method.   Methods   25,   402–408.   DOI:
10.1006/meth.2001.1262
113. Adusumilli, R., and Mallick, P. (2017). Data conversion with ProteoWizard msConvert. Methods
Mol. Biol. 1550, 339–368. DOI: 10.1007/978-1-4939-6747-6_23

114. Schmid, R. et al. (2023). Integrative analysis of multimodal mass spectrometry data in MZmine
3. Nat. Biotechnol. 41, 447–449. DOI: 10.1038/s41587-023-01690-2

115. Nothias, L.-F. et al. (2020). Feature-based molecular networking in the GNPS analysis
environment. Nat. Methods 17, 905–908. DOI: 10.1038/s41592-020-0933-6

116. Shannon, P. et al. (2003). Cytoscape: a software environment for integrated models of
biomolecular interaction networks. Genome Res. 13, 2498–2504. DOI: 10.1101/gr.1239303

117. Ory, L. et al. (2020). Detection of ergosterol using liquid chromatography/electrospray ionization
mass spectrometry: investigation of unusual in-source reactions. Rapid Commun. Mass
Spectrom. 34, e8780. DOI: doi.org/10.1002/rcm.8780

118. Rex, J. H. et al. (2008). Reference method for broth dilution antifungal susceptibility testing of
filamentous fungi, approved standard. Second edition. M38-A2. Clin. Lab. Stand. Inst. *28*, 1–
35.

1018