Chromatin landscape of endometrial tumor cells of tamoxifen users is associated with loss of progestin regulation

Luciana Ant[1], Nicolás Bellora[2], Alejandro La Greca[3], François Le Dily[4], Patricia Saragüeta[1]*

*For correspondence:
patriciasaragueta2@gmailcom (PS)

[1]Biology and Experimental Medicine Institute, IBYME-CONICET, Argentina; [2]Institute of Nuclear Technologies for Health, INTECNUS-CONICET, Argentina; [3]Laboratory of Applied Neuroscience Research (LIAN), Foundation for the Fight Against Childhood Neurological Diseases (FLENI), Institute of Neurosciences (INEU), CONICET, Argentina; [4]Centre for Genomic Regulation (CRG), Barcelona Institute for Science and Technology, Spain

Abstract   Tamoxifen is an effective and widely steroid estrogen receptor modulator (SERM) used in the treatment of breast cancer applied therapy in breast cancer. It acts through competitive inhibition of natural ligand estrogen (E2) of estrogen receptor (ER), affecting ERα interactions with other nuclear receptors, chromatin modulators, genomic structural proteins and co-regulators. Although tamoxifen inhibits the progression of breast cancer, it increases the risk of endometrial cancer, particularly in postmenopausal women. Tamoxifen could shift the DNA-binding signature of endometrial ERα upon prolonged exposure during breast cancer, resembling the signature of endometrial ERα in mammary cancer cells. Our laboratory had previously analyzed the progesterone receptor (PR) and ERα cistromes of endometrial adenocarcinoma Ishikawa cells when treated with ovarian hormones, showing that ERα binding sites are shared by PR in these endometrial adenocarcinoma cell line. This could explain, in part, progestin regulation of estrogen effects on ER/PR positive endometrial cancer cells. Here, we compared ERα cistromes of endometrial tumors of both tamoxifen users and non-users with the ERα and PR cistromes of the adenocarcinoma cell model. We found that ER cistrome of non-users is closer to transcriptionally active Ishikawa PR cistrome after R5020 treatment, while the ER cistrome of tamoxifen users is similar to Ishikawa ER cistrome after E2 treatment. These results indicate that tamoxifen treatment could change the ER chromatin landscape of endometrial tumor cells that can be occupied by PR in response to progestins. The subset of ERbs of non-users that could be shared by PR is enriched in NOTCH and inflammatory signaling pathway genes. On the other hand, the subset of ERbs of tamoxifen users that could be shared by ER in Ishikawa cells is enriched in early estrogen response pathway genes.

These results could explain a context dependent response of genomic regions critical to be taken into account to prevent a deregulation of endometrial cells under tamoxifen treatment.

Introduction

Endometrial cancer (EC) is the sixth most common malignancy worldwide (*Sung et al., 2021*). Epidemiological trends indicate a rapid rise in EC incidence (*Sung et al., 2021*; *Siegel et al., 2023*). Notably, mortality rates have increased by 1.8% annually since 2010 (*Sung et al., 2021*; *Siegel et al., 2023*). Historically, EC has been classified in two categories: Type I or Type II (*Bokhman, 1983*). Type I EC, which are more prevalent and generally associated with a favorable prognosis, are linked to unopposed estrogen exposure and typically consist of grade I or II endometrioid adenocarcinomas. In contrast, Type II ECs are considered hormone-independent and tend to be more aggressive, encompassing grade III endometrioid adenocarcinomas, serous, clear cell, undifferentiated carcinomas, and carcinosarcomas (*Makker et al., 2021*; *How et al., 2024*).

In hormone-dependent Type I EC, the interplay between estrogen and progesterone signaling -mediated by estrogen receptor alpha (ERα) and progesterone receptor (PR), respectively- plays a crucial role in tumor development and progression. Estrogens, particularly 17-estradiol (E2), are essential for the growth and differentiation of various tissues, including the endometrium, where they promote proliferation through ERα and ER. ERα modulates gene expression via several mechanisms, the most classical of which involves direct binding to estrogen response elements (EREs) within chromatin, regulating target genes such as the progesterone receptor gene (PGR) (*Hall et al., 2001*). In addition, ERs can influence transcription through ERE-independent pathways, wherein DNA binding is facilitated by interactions with other transcription factors, such as Fos and Jun (*Hall et al., 2001*).

Estrogen promotes the proliferation of the endometrial glandular epithelium -the primary cell type involved in endometrial cancer- whereas progesterone antagonizes these estrogenic effects (*Yang et al., 2011*). A disruption in the equilibrium between ERα and PR signaling has been associated with endometrial carcinogenesis (*Setiawan et al., 2013*). Specifically, prolonged exposure to unopposed estrogen can result in endometrial hyperplasia and overgrowth, a recognized precursor to malignancy (*Horn et al., 2007*; *Lydon et al., 1995*). Progesterone exerts a protective effect by inhibiting estrogen-mediated proliferation and promoting cellular differentiation (*Yang et al., 2011*). Notably, the loss of PR expression has been correlated with increased proliferation in both ER-positive and ER-negative endometrial tumors (*Tangen et al., 2014*).

Progesterone mediates its biological effects through two main isoforms of the progesterone receptors, PRA and PRB, which are transcribed from distinct promoters of the PGR gene (*Hovland et al., 1998*). In the endometrium, the relative expression of these isoforms is cell-type specific:, PRA predominates in stromal cells, whereas PRB is more abundant in epithelial cells.Upon ligand binding, progesterone receptors modulate gene expression by interacting with progesterone response elements (PREs) in the genome, thereby regulating the transcription of downstream target genes.

Selective estrogen receptor modulators (SERMs), such as tamoxifen, are commonly used in treatment of estrogen receptor-positive breast cancer. Tamoxifen exerts its effects by binding to the ligand-binding domain of ERα. While it functions predominantly as an ER antagonist in breast tissue, inhibiting estrogen-driven proliferation (*Fisher et al., 1989*; *Sutherland et al., 1983*), its activity is highly tissue-specific. In the endometrium, tamoxifen paradoxically acts as an ER agonist, promoting epithelial proliferation and increasing the risk of endometrial cancer (*Fisher et al., 1994*; *Lahti et al., 1993*; *van Leeuwen et al., 1994*). Studies using the Ishikawa endometrial cancer cell line have demonstrated that tamoxifen modulates gene expression through ERα-dependent mechanisms (*Shang and Brown, 2002*). Notably, genome-wide analyses revealed significant overlap between the ERα cistromes of tamoxifen-associated endometrial tumors and those observed in breast cancer (*Droog et al., 2016*). Moreover, tamoxifen-associated endometrial tumors exhibit a distinct ERα DNA-binding profile compared to spontaneous cases, with these altered binding patterns correlating with differential gene expression (*Droog et al., 2017*).

In breast cancer, PR modulates ERα activity by influencing its chromatin binding dynamics and

downstream gene expression (*Mohammed et al., 2015*). Studies in T47D mammary cancer cells have identified hormone control regions (HCRs), which harbor both ERα and PR binding sites and physically interact with promoters of hormone-regulated genes within hormone-responsive topologically associating domains (TADs) (*Nacht et al., 2019*; *Le Dily et al., 2019*). In contrast, the interplay between ERα and PR in EC remains poorly characterized. Recent work by *La Greca et al.* (*2022*) demonstrated that estrogen priming induces the formation of progestin-dependent PR binding sites in Ishikawa endometrial cancer cells. These sites are frequently located within 'progestin control regions' (PgCRs), a subset of HCRs situated in TADs, involving coordinated binding of ERα, PR, and the transcription factor PAX2 to regulate gene expression. Notably, PgCRs are characterized by an open chromatin conformation and are associated with the regulation of genes implicated in endometrial carcinogenesis.

While it is established that prolonged tamoxifen treatment is associated with a distinct ERα binding profile in endometrial tumors compared to those not associated with tamoxifen, the relationship between these altered ERα binding sites and the regulatory landscape of the PR remains uncharacterized. The interplay between ERα and PR is critical in hormone-dependent endometrial cancer, with PR having the capacity to modulate ERα's effects. Previous work has highlighted that ERα's positioning is a key determinant in the development of hormone-dependent endometrial cancer. Therefore, this study aims to investigate how tamoxifen-induced changes in the ERα cistrome relate to PR binding sites in endometrial cancer. We hypothesize that tamoxifen disrupts the co-regulatory relationship between ERα and PR, leading to a distinct oncogenic program. To test this, we conducted a comparative genomic analysis integrating publicly available ERα ChIP-seq data from endometrial tumors of tamoxifen users and non-users with our previously published PR and ERα cistromes from the Ishikawa endometrial cancer cell line. This approach allowed us to delineate the differential ERα and PR genomic interplay in tamoxifen-associated versus non-associated endometrial cancers, providing novel insights into the specific molecular pathways dysregulated by tamoxifen exposure.

Results

Tamoxifen-driven shifts in ER binding sites reflect estrogen-responsive, PR-independent regulation

To investigate how tamoxifen alters the estrogen receptor (ER) chromatin landscape in relation to progesterone receptor (PR) signaling, we conducted a comparative analysis of ER binding sites from endometrial tumors of tamoxifen users and non-users. Due to the absence of publicly available PR binding data from endometrial tumors, we employed our previously characterized ER and PR cistromes from the Ishikawa endometrial cancer cell line as a reference framework for interpreting the changes in ER cistrome of users and non-users. This comparative analysis supported the hypothesis that tamoxifen treatment leads to an ER binding profile less influenced by PR in tumors from tamoxifen users.

ER landscape of tamoxifen user and non-user endometrial tumors (*Droog et al., 2016*, *2017*) were compared to our previously published ER and PR cistromes generated from Ishikawa endometrial cancer cell line treated with 17-estradiol (E2) or the synthetic progestin R5020 (*La Greca et al., 2022*) (Figure 1a). In our previous work, we observed that estrogen pre-treatment leads to an increase in the total PR binding sites (PR bs), while retaining 97% of PR bs identified without E2 pre-treatment (Figure 1—Figure supplement 1a). Accordingly, we focused on the PR cistrome in estrogen-primed Ishikawa cells to characterize PR binding within the identified estrogen-responsive context (Figure 1—Figure supplement 1a).

For this comparison, raw data of ER ChIP-seqs from 6 non-users tumor and 9 tamoxifen users tumors were reprocessed under the same conditions as the ChIP-seq of Ishikawa cell line (Figure 1—Figure supplement 1b). Principal Component Analysis (PCA) of the reprocessed samples is shown in Figure 1—Figure supplement 2a. PC1 vs PC2 graph revealed two distinct clusters corre-

sponding to tamoxifen users and non-user (Figure 1—Figure supplement 2b).

To characterize the ER binding site population in the endometrial tumor samples, we classified sites into three categories: unique to tamoxifen users, unique to non-users, and common to both (Figure 1—Figure supplement 3a). In tamoxifen users, the ER ChIP-seq signal showed a loss of 2167 binding sites compared to non-users, while tamoxifen treatment contributed to the acquisition of additional 2259 binding sites in tumors (Figure 1—Figure supplement 3a). To associate these sites with enhancer activity, K3K27ac levels were analyzed, revealing that tamoxifen-treated tumors exhibited increased activity at the binding sites gained after treatment (Figure 1—Figure supplement 3b).

To understand the functional ER binding landscape in tumor samples, we quantified the overlap with the ER bs and PR bs identified in Ishikawa cells. This revealed a distinct pattern between non-users and tamoxifen users (Figure 1b). Ishikawa ER bs shared 3-fold (249/84) 165 more binding sites with ER bs from tamoxifen users than with those from non-users (Figure 1b). Conversely, Ishikawa PR bs exhibited 4.7-fold (419/89) greater overlap with ER bs from non-users compared to tamoxifen users (Figure 1b). This asymmetry was also evident considering in the percentage of binding sites uniquely shared: 5.8% of all ER bs in tamoxifen users were uniquely shared with Ishikawa ER bs (202 bs), while only 1.2% were uniquely shared with Ishikawa PR bs (20 bs). In contrast, non-users shared 0.59% of their ER bs exclusively with Ishikawa ER bs (42 bs) and 10.4% with Ishikawa PR bs (355 bs) (Figure 1c). Furthermore, the ChIP-seq signal intensity for ER at the binding sites shared with Ishikawa ER was higher in tamoxifen users than in non-users, whereas non-users showed a stronger ER signal at sites overlapping with Ishikawa PR bs (Figure 1d).

These results suggest that tamoxifen exposure in endometrial tumors is associated with an ER chromatin binding profile that more closely resembles the ER binding landscape observed in estrogen-treated Ishikawa cells. In contrast, the ER binding profile of non-tamoxifen users exhibits greater similarity to the PR binding pattern in the same cell line.

ER/PR binding in non-user links to NOTCH and inflammatory gene expression

To understand the potential regulatory roles of the 355 unique non-user ER bs that overlap with Ishikawa PR binding, we employed two complementary approaches to associate these regions with nearby genes.

First, we used the GREAT algorithm (*McLean et al., 2010*), which defines both a basal and a distal domain for each gene. The basal regulatory domain extends 5 kb upstream and 1 kb downstream of the transcription start site (TSS). Briefly, the distal regulatory domain extends in both directions from the basal domain until it reaches the basal domain of the nearest gene, but no more than 1,000 kb in any direction. Then the genes whose regulatory domains contained one of the 355 unique non-user ER bs overlapping Ishikawa PR bs were selected. Gene ontology (GO) analysis of the associated gene set revealed significant enrichment in the Mucyn type O-glycan biosynthesis pathway, Basal transcription factors, and two NOTCH-related pathways: NOTCH receptor signaling and NOTCH signaling deregulation in cancer (Figure 2a).

Given the established involvement of the NOTCH pathway in various cancers -including breast, ovarian, and cervical (*Shi et al., 2024*)- we assessed potential differential expression of these pathways in the publicly available microarray dataset of endometrial tumors from tamoxifen users and non-users (*Droog et al., 2017*). DEseq analysis of this dataset for individual differentially expressed genes failed to reveal any significant differences (p-value $\leq$ 0.05). GSEA of "NOTCH signaling deregulation in cancer" did not reveal statistically significant enrichment (p-value: 0.2) (Figure 2—Figure supplement 1a). Gene expression of NOTCH-related genes in Ishikawa cells treated with ethanol (vehicle), estradiol (E2), or tamoxifen revealed the differential expression of several (CCND1, CDKN1A, FURIN, HES1, JAG1, JAG2, MYC, NCOR2, NCSTN, NOTCH1, NOTCH3 and TWIST1) genes within both the "NOTCH receptor signaling" and "NOTCH signaling deregulation in cancer" pathways (Figure 2—Figure supplement 1b) (*Puyang et al., 2018*). These changes are variable dependent on different treatments. These findings suggest that shared PR and ER binding regions

may be involved in regulating NOTCH signaling genes, but the Ishikawa cell model may not fully
recapitulate the complex and potentially time-dependent effects of long-term tamoxifen exposure
in vivo. Further investigation is warranted to clarify the functional role of these shared binding sites
in modulating NOTCH-related gene expression.

In a second approach, we associated the shared ER bs with genes located within the same
topologically associated domains (TADs), based on published Hi-C data for Ishikawa cells (*La Greca*
et al., 2022). Ontology analysis of genes located within TADs containing these ER bs revealed
a significant enrichment in several pathways involved in inflammation processes, including the
Chemokines Receptors Bind Chemokines (R-HSA-380108), Inflammatory Response, TNF-alpha Sig-
naling via NF-kB and IL-6/JAK/STAT3 Signaling pathways (Figure 2b). We further pooled the genes
from the Chemokines Receptors Bind Chemokines and Inflammatory Response pathways and con-
ducted GSEA on the endometrial tumor microarray dataset. This analysis revealed a significant
enrichment of the "Chemokines Receptors Bind Chemokines" pathway in non-users compared to
tamoxifen users (Figure 2c), suggesting higher expression of inflammatory genes in the absence
of tamoxifen.

Among these, the proto-oncogene MYC, a key regulator of inflammation, emerged as a notable
candidate. Myc overexpression of c-Myc has been reported in over 70% of endometrial tumors
(*Zhang et al., 2018*) and its expression correlates with disease prognosis (*Geisler et al., 2004*). Prior
studies indicate that PR can act as a negative regulator of MYC transcription in endometrial can-
cer, independent of ERα status (*Kavlashvili et al., 2016*). We therefore examined the epigenetic
landscape of the MYC locus (Figure 3). A cluster of PR bs was identified upstream of the MYC
transcription start site (TSS), with a subset of overlapping ER bs in non-users. Notably, the ER
binding profiles at this locus differed between tamoxifen users and non-users. Although GSEA in-
dicated higher expression of the "Chemokines Receptors Bind Chemokines" gene set in non-users,
we observed increased H3K27ac ChIP-seq signal at the MYC locus in tamoxifen. Consistent with
this, reanalysis of RNA-seq data from Ishikawa cells (*La Greca et al., 2022*) showed significantly
increased MYC expression following E2 treatment compared to control (adjusted p-value $\leq$ 0.05).
Interestingly, MYC expression after 1-hour tamoxifen treatment did not significantly differ from E2
treatment (adjusted p-value $\leq$ 0.05)(Figure 3—Figure supplement 1). Together, these data suggest
a complex regulatory landscape involving inflammatory genes and proto-oncoges such as MYC,
with divergent regulatory mechanisms in tamoxifen users and non-users.

ER occupancy at PR-independent sites in tamoxifen treated tumors

Next, we investigated the genomic proximity of Ishikawa ER bs to the nearest PR bs (Figure 4—Fig-
ure supplement 1). Most ER bs were located within 5 kb of a PR bs (Figure 4—Figure supplement
1). To better dissect their potential interactions, we categorized ER bs into two groups: those over-
lapping PR bs by at least one base pair (ER overlapping PR) and those not overlapping PR bs (1 bp
from the nearest PR bs) (ER not overlapping PR) (Figure 4a). The former group likely represents loci
potentially co-regulated by both ER and PR, whereas the latter are probably independent of direct
PR influence. Our analysis revealed that 732 out of the 1591 Ishikawa ER bs overlapped with PR bs
(Figure 4b). To explore whether the regulatory behavior of these two categories differs between
tamoxifen users and non-users, we focused on Ishikawa ER bs (both overlapping and not overlap-
ping PR bs) that were also detected in tumors from either of the tamoxifen users or non-users.
When examining the ER ChIP-seq signal in tamoxifen users and non-users, we observed higher ER
occupancy at ER bs not co-occupied by PR in tamoxifen users than non-users (Figure 4c). In con-
trast, at ER bs co-occupied by PR, the ER signal was higher in non-users compared to tamoxifen
users (Figure 4c). Interestingly, ER bs overlapping with PR bs showed elevated H3K27ac levels in
tumors from tamoxifen users (Figure 4d), suggesting that enhancers capable of binding both ER
and PR exhibit increased activity in the context of tamoxifen.

To identify potential target genes associated with these binding sites, we applied the two ap-
proaches described previously. Ontology analysis of genes located within the same topologically

associating domains (TADs) as ER bs overlapping PR bs revealed significant enrichment for the Estrogen Response Early pathway (Figure 4e). When we instead considered genes within TADs harboring binding sites shared by both Ishikawa ER bs and PR bs, the Estrogen Response Early pathway remained the most enriched, albeit with lower significance compared to ER bs not shared with PR (Figure 4—Figure supplement 2a). Similar trends were observed with the second approach, which focused on genes proximal to the binding sites: the Estrogen Response Early pathway showed stronger enrichment for ER bs not shared with PR bs (Figure 4—Figure supplement 2b) compared to those shared with PR bs (Figure 4—Figure supplement 2c). Finally, we analyzed the ER ChIP-seq signal near the transcription start sites (TSSs) of Estrogen Response Early genes in tamoxifen users and non-users (Figure 4f). Notably, the ER signal around the TSSs of these genes was significantly higher in tamoxifen users compared to non-users.

Tamoxifen-associated Chromatin Reorganization Influences NRIP1 Expression and Estrogen Response Gene Activity

To determine whether ER binding near genes within the Estrogen Response Early pathway leads to corresponding changes in gene expression, we first performed an ontology analysis of genes differentially upregulated in Ishikawa cells treated with E2 compared to R5020 (adjusted p-value $\leq$ 0.05, log2 fold change $\geq$ 0), using data from our previous study (*La Greca et al., 2022*). As expected, genes from the Estrogen Response Early pathway were upregulated when treated with estrogen (Figure 5a). We next performed a GSEA analysis of this pathway using microarray expression profiles from tamoxifen users and non-users (*Droog et al., 2017*) (Figure 5b). This revealed higher expression of Estrogen Response Early pathway in tamoxifen users compared to non-users.

Within this pathway, the Nuclear Receptor Interacting Protein 1 (NRIP1) gene emerged as a key component (Figure 5). NRIP1 encodes a nuclear co-regulator that modulates the transcriptional activity of various transcription factors, including ER. Its differential epigenetic landscape between tamoxifen users and non-users suggests altered regulation of endometrial tumors (Figure 5c). Hi-C chromatin conformation analysis revealed a differential loop (highlighted by blue dashed lines) present in R5020-treated, but not in E2-treated, Ishikawa cells. This loop connects two CTCF binding sites. In addition, a virtual 4C analysis centered on the NRIP1 TSS indicated a higher contact frequency between the promoter and ER/PR binding sites in Ishikawa cells (5th track).

Expression profiling of NRIP1 in Ishikawa cells showed downregulation under R5020 treatment compared to E2, and upregulation when comparing E2 to untreated conditions (adjusted p-value $\leq$ 0.05) (Figure 5c, 3 last tracks). Similar epigenetic features were also observed for the genes: Growth Regulated by Estrogen in Breast Cancer 1 (GREB1), Cyclin A1 (CCNA1) and Dehydrogenase/reductase 3 (DHRS3) from the Estrogen Response Early pathway (Figure 5—Figure supplement 1). To assess NRIP1 expression in clinical samples, we analyzed RNA-seq data from 394 endometrial adenomas and adenocarcinomas obtained from The Cancer Genome Atlas Program (Figure 5d). Tumors at stage IV showed significantly lower NRIP1 expression compared to stage I (adjusted p-value $\leq$ 0.05). Since NRIP1 expression correlates with ER levels (*Flindris et al., 2021*), this downregulation may reflect the reduced ER expression typically observed in advanced-stage endometrial cancer. Finally, NRIP1 expression was significantly higher in Ishikawa cells treated with E2 (adjusted p-value $\leq$ 0.05) compared to tamoxifen or control (Figure 5e). Notably, NRIP1 has been linked to endometrial cancer and is the most frequently mutated gene in microsatellite-unstable EC (*Ferreira et al., 2014*).

Collectively, these results suggest that the Estrogen Response Early Pathway plays a pivotal role in shaping endometrial response to tamoxifen. Prolonged tamoxifen treatment may enhance ER binding at sites limited PRco-regulation, leading to upregulation of the Estrogen Response Early pathway, influencing key genes such as NRIP1, and contributing to a distinct genomic and transcriptional landscape in tamoxifen-associated endometrial cancer compared to tumors from non-users.

The findings suggest that endometrial cancer development in tamoxifen users may involve different molecular mechanisms compared to non-users, potentially reflecting a loss of PR-mediated

regulation in non-users and the deregulation of PR-independent pathways in tamoxifen users. Further research is needed to fully elucidate the long-term effects of tamoxifen on gene regulation and to identify therapeutic strategies that specifically target the unique molecular features of tamoxifen-associated endometrial cancers.

Discussion

Our analysis revealed that the ERα binding profile in tamoxifen users closely mirrors the ER landscape observed in estrogen-treated Ishikawa cells, whereas the ER profile of non-users more closely aligns with the PR binding pattern in the same cell line. This shift is consistent with previous observations (*Droog et al., 2016*, *2017*), which demonstrated a distinct ERα binding profile in tamoxifen-associated endometrial tumors. Prior studies have established that PR can modulate ERα activity and downstream gene expression in breast cancer (*Mohammed et al., 2015*) and that the coordinated binding of ERα, PR, and PAX2 regulates transcriptional programs implicated in endometrial carcinogenesis (*La Greca et al., 2022*). In this context, our findings suggest that tamoxifen disrupts PR-mediated regulation of ERα, resulting in alternative transcriptional programs that may contribute to cancer progression (Figure 6).

A major limitation is the lack of in vivo PR cistrome data in endometrial tumors, and particularly in tamoxifen-associated cases. Thus, our inference relies on Ishikawa-derived PR cistromes, which, while informative, may not fully capture the complexity of primary tumors. Specifically, Ishikawa cells cannot model the heterogeneity of tumor subtypes or the stromal–epithelial interactions that shape receptor activity in vivo. Nevertheless, the convergence of our findings with prior mechanistic studies supports the hypothesis that tamoxifen reshapes the ER–PR regulatory axis, highlighting the need for future studies using primary tumor material to define PR-dependent modulation of ERα under tamoxifen exposure.

Further characterization of the differential binding patterns revealed that ER binding sites uniquely shared with Ishikawa PR sites in non-users were enriched for pathways related to NOTCH receptor signaling and inflammatory responses, which correlated with increased expression of the associated genes. Among these, the proto-oncogene MYC was particularly notable, as its overexpression has been previously linked to endometrial tumorigenesis (*Zhang et al., 2018*). In Ishikawa cells, MYC expression was significantly upregulated following both estradiol (E2) and tamoxifen treatment, although it is important to note that short-term in vitro exposure to tamoxifen does not fully replicate the chronic treatment experienced by patients.

The anti-inflammatory effects of progesterone, mediated through PR, are well documented in the endometrium, including in endometrial adenocarcinomas (*Davies et al., 2004*). Our results are consistent with a model in which PR, acting in concert with ER, regulates inflammatory pathways in non-users, and loss of this regulatory axis may contribute to tumor development (Figure 6). By contrast, tamoxifen-associated endometrial cancer may engage alternative pathways not typically modulated by PR, such as PI3K signaling (*Kübler et al., 2025*). These observations underscore the need for further investigation to delineate how tamoxifen-induced shifts in ER cistromes reprogram transcriptional networks, particularly those governing inflammatory signaling, and how these changes influence endometrial tumor biology.

Tamoxifen exposure reshapes the epigenetic and transcriptional landscape of EC by disrupting the ERα–PR axis, leading to loss of PR-mediated control of inflammatory pathways and enhanced ER-driven activation of the Estrogen Response Early program (Figure 6). These findings define tamoxifen-associated EC as a distinct molecular subtype with important implications for prognosis and targeted therapy development.

Methods and Materials

ChIP-seq Data Processing

High-quality sequencing reads were aligned to the human reference genome (hg38, UCSC) using STAR v3.0.1 (*Dobin et al., 2013*), with alignment parameters set to a maximum intron size of 1 and end-to-end alignment enabled. For initial quality control and sample clustering, reads for each sample were aligned independently. Subsequently, samples were stratified based on tamoxifen exposure (users vs. non-users) and reads within each group were merged into a single BAM file for downstream analysis. Peak calling was performed using MACS2 v2.2.6 (*Zhang et al., 2008*) with default parameters, employing the corresponding input controls to identify regions of significant signal enrichment. Venn diagrams illustrating peak set overlaps were generated using the venn function from the Intervene package (*Khan and Mathelier, 2017*). Principal Component Analysis (PCA) and heatmap visualizations were generated using deepTools (*Ramírez et al., 2016*). Specifically, computeMatrix was used to calculate signal intensities, plotHeatmap for heatmap generation, multiBigwigSummary for region-wise signal quantification, and plotPCA for dimensionality reduction. Genomic overlap and distance between ER and PR binding sites in Ishikawa cells were assessed using the intersectBed and closest utilities from bedTools (*Quinlan and Hall, 2010*).

RNA-seq Data Processing

Differential gene expression analysis was conducted using the DESeq2 package v1.44.0. (*Love et al., 2014*). Genes exhibiting an absolute log fold change ($|log2FC|$) ≥ 1 and a Benjamini-Hochberg adjusted p-value (Padj) ≤ 0.05 were considered differentially expressed. Expression levels, quantified as FPKM (Fragment Per Kilobase of transcript, per Million mapped reads) and TPM (Transcripts Per Million) were visualized using custom Python scripts employing the seaborn library (*Waskom, 2021*).

Gene Set Enrichment Analysis (GSEA) and Gene Ontology Enrichment analysis

The association between binding sites with genes was done through two different approaches. The first one was done using GREAT algorithm (*McLean et al., 2010*; *Tanigawa et al., 2022*). This algorithm first assigns every gene a regulatory domain and then associates each genomic region with all genes whose regulatory domain it overlaps. The association was done using the Basal plus extension setting, where each gene is assigned a basal regulatory domain of a minimum distance upstream and downstream of the TSS (regardless of other nearby genes). The gene regulatory domain is extended in both directions to the nearest gene's basal domain but no more than the maximum extension in one direction. The algorithm was runned with 5kb upstream and 1kb downstream from TSS as the proximal distance and up to 1000kb as the maximum extension. The second approach to find genes associated with binding sites was performed by selecting the TADs obtained from Ishikawa Hi-C that comprised binding sites of interest and retrieving all the genes contained in that TAD.

Gene Set Enrichment Analysis (GSEA) and Gene Ontology (GO) enrichment were performed using the GSEApy package (*Fang et al., 2023*), a Python-based implementation of GSEA and a wrapper for the Enrichr database. Gene expression data obtained from GEO was analyzed using the GSEA function in GSEApy with the following parameters: permutation type set to "phenotype", 1000 permutations, and "difference of class means" as the scoring method. For GO and pathway enrichment, the enrichr function was employed to query the following libraries: "BioCarta_2016" , "KEGG_2021_Human", "Reactome_2022", "MSigDB_Hallmark_2020", and "Elsevier_Pathway_Collection". Enriched terms with an adjusted p-value (Padj) ≤ 0.05 were considered significant.

Hi-C Data Processing

Previously filtered Hi-C datasets from *La Greca et al.* (*2022*), excluding low-quality reads, were further processed using the Juicer pipeline (*Durand et al., 2016*) with the human reference genome

hg38. After individual sample contact maps were generated, contact matrices from biological repli-
cates were merged by summing the contact counts between corresponding genomic loci, resulting
in a combined contact map. Topological Associated Domains (TADs) were identified using the Ar-
rowhead algorithm, and chromatin loops (i.e., locally enriched contacts) were detected using the
HiCCUPS algorithm, both implemented within the Juicer framework.

Virtual 4C

Virtual 4C profiles were generated through custom Python scripts using Hi-C matrices of Observed/expected
contacts with a 5kb resolution and normalized via Knight-Ruiz (KR) algorithm as input. The matrices
for the region containing NRIP1 gene were obtained from hic files through Straw (*Durand et al.,
2016*).

Endometrial Cancer Samples (TCGA)

RNA-seq data (FPKM values) for human endometrial cancer samples (n = 394) were obtained from
The Cancer Genome Atlas (TCGA) project. Each sample was annotated with its corresponding FIGO
stage. Differential expression analysis of NRIP1 across FIGO stages was performed using pairwise
comparisons with Tukey's Honestly Significant Difference (HSD) post-hoc test, implemented via the
statsmodels Python package (*Seabold and Perktold, 2010*). An adjusted p-value (p-adj) ≤ 0.09 was
considered statistically significant.

Competing interests

The authors declare no competing interests.

Acknowledgments

We are grateful to members of the Saragüeta laboratory for help and suggestions and to Gabriela
Diessler for the support.

Additional information

Funding

Funder	Grant reference number	Author
Fondo para la Investigación Científica y Tecnológica	PICT 2021-1142	Patricia Saragüeta
Consejo Nacional de Investigaciones Científicas y Técnicas	PIP 2022-2024 Num 776	Patricia Saragüeta

The funders had no role in study design, data collection and interpretation, or the decision to
submit the work for publication.

Author contributions

Luciana Ant, Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Visual-
ization, Writing – original draft, Writing – original draft, Writing – review and editing; Nicolás Bellora,
Data curation, Formal analysis, Methodology, Supervision; Alejandro La Greca, Formal analysis,
Writing – review and editing; François Le Dily, Formal analysis; Patricia Saragüeta, Conceptualiza-
tion, Formal analysis, Funding acquisition, Investigation, Project administration, Supervision, Writ-
ing – original draft, Writing – review and editing.

Author ORCIDs

Luciana Ant: https://orcid.org/0009-0003-4212-8280
Nicolás Bellora: https://orcid.org/0000-0001-6637-3465

413   Alejandro La Greca: https://orcid.org/0000-0002-0309-7683

414   Francois Le Dily: https://orcid.org/0000-0002-8324-7927

415   Patricia Saragüeta: https://orcid.org/0000-0001-8222-9690

Data Availability

ER and H3K27ac ChIP-seq data and microarray data from endometrial cancers from tamoxifen
users and non-users, as reported by *Droog et al.* (*2017*, 2016) are accessible in Gene Expression
Omnibus (GEO) under accession number GSE94524. ER and PR ChIP-seq data along with RNA-seq
data from Ishikawa cells treated with E2 and R5020 (*La Greca et al., 2022*) are available under
accession number GSE139398. RNA-seq data from Ishikawa cells treated with tamoxifen and E2
(*Puyang et al., 2018*) can be accessed via GEO accession number GSE115608. CTCF binding sites
in Ishikawa cells, used in omics track figures, are available from ENCODE under accession number ENCSR000BQE. RNAseq data for human endometrial cancer samples were obtained from The
Cancer Genome Atlas (TCGA) under project TCGA. Processed data and custom scripts for virtual
4C, TCGA samples analysis and RNAseq differential expression analysis are accessible at Figshare
under reference number 28840463 (10.6084/m9.figshare.28840463).

The following previously published datasets were used:

Author(s)	Year	Dataset title	Dataset URL	Database and Identifier
Droog M, Nevedomskaya E, Dackus GM, Fles R, Kim Y, Hollema H, Mourits MJ, Nederlof PM, Van Boven HH, Linn SC, Van Leeuwen FE, Wessels LFA, Zwart W	2017	Tamoxifen-associated endometrial tumors expose differential enhancer activity for Estrogen Receptor alpha	https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE94524	NCBI Gene Expression Omnibus, GSE94524
La Greca A, Bellora N, Le Dily F, Jara R, Nacht AS, Quilez Oliete J, Villanueva JL, Vidal E, Merino G, Fresno C, Tarifa Reischle I, Vallejo G, Vicent G, Fernández E, Beato M, Saragüeta P	2022	Higher-order chromatin organization defines PR and PAX2 binding to regulate endometrial cancer cell gene expression	https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE139398	NCBI Gene Expression Omnibus, GSE139398
Puyang X, Furman C, Wu ZJ	2018	Transcriptional profiling of Ishikawa cells treated with H3B-5942, E2, or standard of care compounds	https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE115608	NCBI Gene Expression Omnibus, GSE115608
Korch C, Spillman MA, Jackson TA, Jacobsen BM, Murphy SK, Lessey BA, Jordan VC, Bradford AP	2012	DNA profiling analysis of endometrial and ovarian cell lines reveals misidentification, redundancy and contamination	https://www.encodeproject.org/experiments/ENCSR000BQE/	ENCODE, ENCSR000BQE

References

Bokhman JV. Two pathogenetic types of endometrial carcinoma. Gynecologic Oncology. 1983 Feb; 15(1):10–17. https://linkinghub.elsevier.com/retrieve/pii/0090825883901117, doi: 10.1016/0090-8258(83)90111-7.

Davies S, Dai D, Wolf DM, Leslie KK. Immunomodulatory and Transcriptional Effects of Progesterone Through Progesterone A and B Receptors in Hec50co Poorly Differentiated Endometrial Cancer Cells. Journal of the Society for Gynecologic Investigation. 2004 Oct; 11(7):494–499. http://journals.sagepub.com/doi/10.1016/j.jsgi.2004.04.003, doi: 10.1016/j.jsgi.2004.04.003.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. STAR: ultrafast universal RNA-seq aligner. Bioinformatics (Oxford, England). 2013 Jan; 29(1):15–21. doi: 10.1093/bioinformatics/bts635.

Droog M, Nevedomskaya E, Dackus GM, Fles R, Kim Y, Hollema H, Mourits MJ, Nederlof PM, Van Boven HH, Linn SC, Van Leeuwen FE, Wessels LFA, Zwart W. Estrogen receptor wields treatment-specific enhancers between morphologically similar endometrial tumors. Proceedings of the National Academy of Sciences. 2017 Feb; 114(8). https://pnas.org/doi/full/10.1073/pnas.1615233114, doi: 10.1073/pnas.1615233114.

Droog M, Nevedomskaya E, Kim Y, Severson T, Flach KD, Opdam M, Schuurman K, Gradowska P, Hauptmann M, Dackus G, Hollema H, Mourits MJ, Nederlof P, Van Boven H, Linn SC, Wessels L, Van Leeuwen FE, Zwart

W. Comparative Cistromics Reveals Genomic Cross-talk between FOXA1 and ER in Tamoxifen-Associated Endometrial Carcinomas. Cancer Research. 2016 Jul; 76(13):3773–3784. https://aacrjournals.org/cancerres/article/76/13/3773/609371/Comparative-Cistromics-Reveals-Genomic-Cross-talk, doi: 10.1158/0008-5472.CAN-14-1813.

Durand NC, Shamim MS, Machol I, Rao SSP, Huntley MH, Lander ES, Aiden EL. Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C Experiments. Cell Systems. 2016 Jul; 3(1):95–98. doi: 10.1016/j.cels.2016.07.002.

Fang Z, Liu X, Peltz G. GSEApy: a comprehensive package for performing gene set enrichment analysis in Python. Bioinformatics. 2023 Jan; 39(1):btac757. https://academic.oup.com/bioinformatics/article/doi/10.1093/bioinformatics/btac757/6847088, doi: 10.1093/bioinformatics/btac757.

Ferreira AM, Tuominen I, Sousa S, Gerbens F, Van Dijk-Bos K, Osinga J, Kooi KA, Sanjabi B, Esendam C, Oliveira C, Terpstra P, Hardonk M, Van Der Sluis T, Zazula M, Stachura J, Van Der Zee AG, Hollema H, Sijmons RH, Aaltonen LA, Seruca R, et al. New Target Genes in Endometrial Tumors Show a Role for the Estrogen-Receptor Pathway in Microsatellite-Unstable Cancers. Human Mutation. 2014 Dec; 35(12):1514–1523. https://onlinelibrary.wiley.com/doi/10.1002/humu.22700, doi: 10.1002/humu.22700.

Fisher B, Costantino JP, Redmond CK, Fisher ER, Wickerham DL, Cronin WM. Endometrial cancer in tamoxifen-treated breast cancer patients: findings from the National Surgical Adjuvant Breast and Bowel Project (NSABP) B-14. Journal of the National Cancer Institute. 1994 Apr; 86(7):527–537. doi: 10.1093/jnci/86.7.527.

Fisher B, Costantino J, Redmond C, Poisson R, Bowman D, Couture J, Dimitrov NV, Wolmark N, Wickerham DL, Fisher ER, Margolese R, Robidoux A, Shibata H, Terz J, Paterson AHG, Feldman MI, Farrar W, Evans J, Lickley HL, Ketner M, et al. A Randomized Clinical Trial Evaluating Tamoxifen in the Treatment of Patients with Node-Negative Breast Cancer Who Have Estrogen-Receptor–Positive Tumors. New England Journal of Medicine. 1989 Feb; 320(8):479–484. http://www.nejm.org/doi/abs/10.1056/NEJM198902233200802, doi: 10.1056/NEJM198902233200802.

Flindris S, Katsoulas N, Goussia A, Lazaris AC, Navrozoglou I, Paschopoulos M, Thymara I. The Expression of NRIP1 and LCOR in Endometrioid Endometrial Cancer. In Vivo. 2021; 35(5):2631–2640. http://iv.iiarjournals.org/lookup/doi/10.21873/invivo.12545, doi: 10.21873/invivo.12545.

Geisler JP, Geisler HE, Manahan KJ, Miller GA, Wiemann MC, Zhou Z, Crabtree W. Nuclear and cytoplasmic c-myc staining in endometrial carcinoma and their relationship to survival. International Journal of Gynecological Cancer. 2004 Jan; 14(1):133–137. https://ijgc.bmj.com/lookup/doi/10.1111/j.1048-891x.2004.14027.x, doi: 10.1111/j.1048-891x.2004.14027.x.

Hall JM, Couse JF, Korach KS. The Multifaceted Mechanisms of Estradiol and Estrogen Receptor Signaling. Journal of Biological Chemistry. 2001 Oct; 276(40):36869–36872. https://linkinghub.elsevier.com/retrieve/pii/S0021925819566157, doi: 10.1074/jbc.R100029200.

Horn LC, Meinel A, Handzel R, Einenkel J. Histopathology of endometrial hyperplasia and endometrial carcinoma. Annals of Diagnostic Pathology. 2007 Aug; 11(4):297–311. https://linkinghub.elsevier.com/retrieve/pii/S1092913407000858, doi: 10.1016/j.anndiagpath.2007.05.002.

Hovland AR, Powell RL, Takimoto GS, Tung L, Horwitz KB. An N-terminal Inhibitory Function, IF, Suppresses Transcription by the A-isoform but Not the B-isoform of Human Progesterone Receptors. Journal of Biological Chemistry. 1998 Mar; 273(10):5455–5460. https://linkinghub.elsevier.com/retrieve/pii/S0021925818677956, doi: 10.1074/jbc.273.10.5455.

How JA, Jazaeri AA, Westin SN, Lawson BC, Klopp AH, Soliman PT, Lu KH. Translating biological insights into improved management of endometrial cancer. Nature Reviews Clinical Oncology. 2024 Nov; 21(11):781–800. https://www.nature.com/articles/s41571-024-00934-7, doi: 10.1038/s41571-024-00934-7.

Kavlashvili T, Jia Y, Dai D, Meng X, Thiel KW, Leslie KK, Yang S. Inverse Relationship between Progesterone Receptor and Myc in Endometrial Cancer. PLOS ONE. 2016 Feb; 11(2):e0148912. https://dx.plos.org/10.1371/journal.pone.0148912, doi: 10.1371/journal.pone.0148912.

Khan A, Mathelier A. Intervene: a tool for intersection and visualization of multiple gene or genomic region sets. BMC Bioinformatics. 2017 Dec; 18(1):287. http://bmcbioinformatics.biomedcentral.com/articles/10.1186/s12859-017-1708-7, doi: 10.1186/s12859-017-1708-7.

Kübler K, Nardone A, Anand S, Gurevich D, Gao J, Droog M, Hermida-Prado F, Akhshi T, Feiglin A, Feit AS, Cohen Feit G, Dackus G, Pun M, Kuang Y, Cha J, Miller M, Gregoricchio S, Lanfermeijer M, Cornelissen S, Gibson WJ, et al. Tamoxifen induces PI3K activation in uterine cancer. Nature Genetics. 2025 Sep; 57(9):2192–2202. https://www.nature.com/articles/s41588-025-02308-w, doi: 10.1038/s41588-025-02308-w.

La Greca A, Bellora N, Le Dily F, Jara R, Nacht AS, Quilez Oliete J, Villanueva JL, Vidal E, Merino G, Fresno C, Tarifa Reischle I, Vallejo G, Vicent G, Fernández E, Beato M, Saragüeta P. Chromatin topology defines estradiol-primed progesterone receptor and PAX2 binding in endometrial cancer cells. eLife. 2022 Jan; 11:e66034. https://elifesciences.org/articles/66034, doi: 10.7554/eLife.66034.

Lahti E, Blanco G, Kauppila A, Apaja-Sarkkinen M, Taskinen PJ, Laatikainen T. Endometrial changes in post-menopausal breast cancer patients receiving tamoxifen. Obstetrics and Gynecology. 1993 May; 81(5 (Pt 1)):660–664.

Le Dily F, Vidal E, Cuartero Y, Quilez J, Nacht AS, Vicent GP, Carbonell-Caballero J, Sharma P, Villanueva-Cañas JL, Ferrari R, De Llobet LI, Verde G, Wright RHG, Beato M. Hormone-control regions mediate steroid receptor-dependent genome organization. Genome Research. 2019 Jan; 29(1):29–39. doi: 10.1101/gr.243824.118.

van Leeuwen FE, Benraadt J, Coebergh JW, Kiemeney LA, Gimbrère CH, Otter R, Schouten LJ, Damhuis RA, Bontenbal M, Diepenhorst FW. Risk of endometrial cancer after tamoxifen treatment of breast cancer. Lancet (London, England). 1994 Feb; 343(8895):448–452. doi: 10.1016/s0140-6736(94)92692-1.

Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biology. 2014 Dec; 15(12):550. https://genomebiology.biomedcentral.com/articles/10.1186/s13059-014-0550-8, doi: 10.1186/s13059-014-0550-8.

Lydon JP, DeMayo FJ, Funk CR, Mani SK, Hughes AR, Montgomery CA, Shyamala G, Conneely OM, O'Malley BW. Mice lacking progesterone receptor exhibit pleiotropic reproductive abnormalities. Genes & Development. 1995 Sep; 9(18):2266–2278. http://genesdev.cshlp.org/lookup/doi/10.1101/gad.9.18.2266, doi: 10.1101/gad.9.18.2266.

Makker V, MacKay H, Ray-Coquard I, Levine DA, Westin SN, Aoki D, Oaknin A. Endometrial cancer. Nature Reviews Disease Primers. 2021 Dec; 7(1):88. https://www.nature.com/articles/s41572-021-00324-8, doi: 10.1038/s41572-021-00324-8.

McLean CY, Bristor D, Hiller M, Clarke SL, Schaar BT, Lowe CB, Wenger AM, Bejerano G. GREAT improves functional interpretation of cis-regulatory regions. Nature Biotechnology. 2010 May; 28(5):495–501. doi: 10.1038/nbt.1630.

Mohammed H, Russell IA, Stark R, Rueda OM, Hickey TE, Tarulli GA, Serandour AA, Birrell SN, Bruna A, Saadi A, Menon S, Hadfield J, Pugh M, Raj GV, Brown GD, D'Santos C, Robinson JLL, Silva G, Launchbury R, Perou CM, et al. Progesterone receptor modulates ER action in breast cancer. Nature. 2015 Jul; 523(7560):313–317. doi: 10.1038/nature14583.

Nacht AS, Ferrari R, Zaurin R, Scabia V, Carbonell-Caballero J, Le Dily F, Quilez J, Leopoldi A, Brisken C, Beato M, Vicent GP. C/EBP mediates the growth inhibitory effect of progestins on breast cancer cells. The EMBO journal. 2019 Sep; 38(18):e101426. doi: 10.15252/embj.2018101426.

Puyang X, Furman C, Zheng GZ, Wu ZJ, Banka D, Aithal K, Agoulnik S, Bolduc DM, Buonamici S, Caleb B, Das S, Eckley S, Fekkes P, Hao MH, Hart A, Houtman R, Irwin S, Joshi JJ, Karr C, Kim A, et al. Discovery of Selective Estrogen Receptor Covalent Antagonists for the Treatment of ERWT and ERMUT Breast Cancer. Cancer Discovery. 2018 Sep; 8(9):1176–1193. https://aacrjournals.org/cancerdiscovery/article/8/9/1176/10351/Discovery-of-Selective-Estrogen-Receptor-Covalent, doi: 10.1158/2159-8290.CD-17-1229.

Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics (Oxford, England). 2010 Mar; 26(6):841–842. doi: 10.1093/bioinformatics/btq033.

Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dündar F, Manke T. deepTools2: a next generation web server for deep-sequencing data analysis. Nucleic Acids Research. 2016 Jul; 44(W1):W160–W165. https://academic.oup.com/nar/article-lookup/doi/10.1093/nar/gkw257, doi: 10.1093/nar/gkw257.

Seabold S, Perktold J. Statsmodels: Econometric and Statistical Modeling with Python. In: Austin, Texas; 2010. p. 92–96. https://doi.curvenote.com/10.25080/Majora-92bf1922-011, doi: 10.25080/Majora-92bf1922-011.

Setiawan VW, Yang HP, Pike MC, McCann SE, Yu H, Xiang YB, Wolk A, Wentzensen N, Weiss NS, Webb PM, Van Den Brandt PA, Van De Vijver K, Thompson PJ, The Australian National Endometrial Cancer Study Group, Strom BL, Spurdle AB, Soslow RA, Shu Xo, Schairer C, Sacerdote C, et al. Type I and II Endometrial Cancers: Have They Different Risk Factors? Journal of Clinical Oncology. 2013 Jul; 31(20):2607–2618. https://ascopubs.org/doi/10.1200/JCO.2012.48.2596, doi: 10.1200/JCO.2012.48.2596.

Shang Y, Brown M. Molecular determinants for the tissue specificity of SERMs. Science (New York, NY). 2002 Mar; 295(5564):2465–2468. doi: 10.1126/science.1068537.

Shi Q, Xue C, Zeng Y, Yuan X, Chu Q, Jiang S, Wang J, Zhang Y, Zhu D, Li L. Notch signaling pathway in cancer: from mechanistic insights to targeted therapies. Signal Transduction and Targeted Therapy. 2024 May; 9(1):128. https://www.nature.com/articles/s41392-024-01828-x, doi: 10.1038/s41392-024-01828-x.

Siegel RL, Miller KD, Wagle NS, Jemal A. Cancer statistics, 2023. CA: A Cancer Journal for Clinicians. 2023 Jan; 73(1):17–48. https://acsjournals.onlinelibrary.wiley.com/doi/10.3322/caac.21763, doi: 10.3322/caac.21763.

Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. CA: A Cancer Journal for Clinicians. 2021 May; 71(3):209–249. https://acsjournals.onlinelibrary.wiley.com/doi/10.3322/caac.21660, doi: 10.3322/caac.21660.

Sutherland RL, Hall RE, Taylor IW. Cell proliferation kinetics of MCF-7 human mammary carcinoma cells in culture and effects of tamoxifen on exponentially growing and plateau-phase cells. Cancer Research. 1983 Sep; 43(9):3998–4006.

Tangen IL, Werner HMJ, Berg A, Halle MK, Kusonmano K, Trovik J, Hoivik EA, Mills GB, Krakstad C, Salvesen HB. Loss of progesterone receptor links to high proliferation and increases from primary to metastatic endometrial cancer lesions. European Journal of Cancer. 2014 Nov; 50(17):3003–3010. https://linkinghub.elsevier.com/retrieve/pii/S0959804914009381, doi: 10.1016/j.ejca.2014.09.003.

Tanigawa Y, Dyer ES, Bejerano G. WhichTF is functionally important in your open chromatin data? PLoS computational biology. 2022 Aug; 18(8):e1010378. doi: 10.1371/journal.pcbi.1010378.

Waskom M. seaborn: statistical data visualization. Journal of Open Source Software. 2021 Apr; 6(60):3021. https://joss.theoj.org/papers/10.21105/joss.03021, doi: 10.21105/joss.03021.

Yang S, Thiel KW, Leslie KK. Progesterone: the ultimate endometrial tumor suppressor. Trends in Endocrinology & Metabolism. 2011 Apr; 22(4):145–152. https://linkinghub.elsevier.com/retrieve/pii/S1043276011000142, doi: 10.1016/j.tem.2011.01.005.

Zhang Q, Xu P, Lu Y, Dou H. Correlation of MACC1/c-Myc Expression in Endometrial Carcinoma with Clinical/Pathological Features or Prognosis. Medical Science Monitor: International Medical Journal of Experimental and Clinical Research. 2018 Jul; 24:4738–4744. doi: 10.12659/MSM.908812.

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM, Brown M, Li W, Liu XS. Model-based Analysis of ChIP-Seq (MACS). Genome Biology. 2008 Sep; 9(9):R137. https://doi.org/10.1186/gb-2008-9-9-r137, doi: 10.1186/gb-2008-9-9-r137.

Figure 1. Comparative analysis of ER and PR binding profiles in endometrial tumors and Ishikawa cells. a. Schematic overview of the experimental set up involving endometrial tumors from tamoxifen users and non-users, as well as Ishikawa endometrial adenocarcinoma cells. **b.** Venn diagram showing the overlap between PR binding sites (PR bs) and ER binding sites (ER bs) in Ishikawa cells, along with ER bs identified in tamoxifen users and non-users. The total number of binding sites (bs) for each group is indicated in parentheses. **c.**. Percentage of ER bs in tamoxifen users and non-users, categorized by their sharing pattern: shared between users and non-users (Between_them); shared only with Ishikawa ER (Only_with_Ishikawa_ER) or Ishikawa PR (Only_with_Ishikawa_PR); unique to each group (Unique); shared between users, non-users, and Ishikawa ER (Between_them_and_Ishikawa_ER); shared only with both Ishikawa ER and PR (Only_with_Ishikawa_ER_PR); shared across all groups (Between_them_and_Ishikawa_ER_PR); and shared between users, non-users, and Ishikawa PR (Between_them_and_Ishikawa_PR). **d.** ERα ChIP-seq signal (CPM) in tumors from tamoxifen users and non-users. Signal was quantified within a 10 kb window centered on Ishikawa ER bs and PR bs (±5 kb). Top panel: Median CPM signal across all Ishikawa ER and PR bs. Bottom panels: Signal intensity for each individual region.

a

b

c

Figure 2. Pathway enrichment analysis of genes associated with PR and non-user ER co-bound regions. a. Top enriched pathways among genes located near Ishikawa PR bs that overlap with ER bs from non-user (±5 kb from the transcription start site, TSS (left). Representation of the GREAT-based approach used to associate binding sites to nearby genes (right). The basal regulatory domain of each gene extends 5 kb upstream and 1 kb downstream of the transcription start site (TSS) and the distal regulatory domain extends in both directions from the basal domain until it reaches the basal domain of the nearest gene, but no more than 1Mb in any direction. The genes whose regulatory domains contained one of the 355 unique non-user ER bs overlapping Ishikawa PR bs were selected. **b.** Top enriched pathways in genes located within topologically associating domains (TADs) that contain at least one Ishikawa PR bs overlapping with ER bs from non-users (left). Representation of the TADs-based approach used to associate binding sites to nearby genes (right) **c.** Gene set enrichment analysis (GSEA) of gene expression profiles from tamoxifen users and non-users. The analyzed gene set includes Chemokine Receptors Bind Chemokines and Inflammatory Response genes located in TADs containing at least one Ishikawa PR bs overlapping with non-users ER bs. The enrichment profile (green) indicates the correlation of gene expression with the ranked expression dataset. Normalized enrichment score (NES) and nominal p-value (nom.p) are reported.

Figure 3. Genomic architecture, epigenetic landscape, and gene expression at MYC locus. From top to bottom, the tracks display: genome scale; contact matrices from Ishikawa cells treated with T0, E2 (E), or R5020 (R); ENCODE CTCF binding sites in Ishikawa cells; UCSC gene annotations; ER and PR ChIP-seq signal in Ishikawa cells (black lines indicate peak regions); ER ChIP-seq signal in endometrial tumors from non-users and tamoxifen users (black lines indicate peaks); H3K27ac ChIP-seq signal in non-users and tamoxifen users; and RNA-seq RPKM values in Ishikawa cells treated with T0, E2 (E), or R5020 (R).

Figure 4. ER bs overlapping PR bs and regulation of Hallmark Estrogen Response Early Pathway. a. Ishikawa ER bs were defined as overlapping PR bs if their peaks shared at least 1 bp. **b.** Number of Ishikawa ER bs that overlap or do not overlap with Ishikawa PR bs. **c.** ER signal distribution in tamoxifen users and non-users. Regions were defined within a 10kb window centered on Ishikawa ER bs (±5 kb) that were also identified as ER bs in either group (tamoxifen users and non-users). Signal intensity represents the median of accumulated reads per region. ER bs are categorized into two mutually exclusive groups: those overlapping PR bs (er_overlapping_pr) and those that do not (not_overlapping_pr). **d.** H3K27ac ChIP-seq signal distribution in tamoxifen users and non-users. Regions were defined as in (a), with signal intensity representing the median of accumulated reads per region. ER bs are categorized as in (a) **e.** Representation of the TADs-based approach used to associate binding sites to nearby genes (top). Top pathways enriched in genes located within TADs containing at least one Ishikawa ER bs that does not overlap a PR bs (bottom). **f.** Median of ER ChIP-seq signal in tamoxifen users and non-users around the transcription start sites (TSS, ±50 kb) of the 200 genes in the Hallmark Estrogen Response Early Pathway.

Figure 5. Pathway enrichment and regulatory features associated with NRIP1 and estrogen signaling. a. Top pathways enriched among genes upregulated in Ishikawa cells treated with E2 compared to R5020 (adjusted p-value $\leq$ 0.05, log_2 fold change $\geq$ 0). **b.** Gene Set Enrichment Analysis (GSEA) of the Hallmark Estrogen Response Early pathway using gene expression profiles from tamoxifen users and non-users. The enrichment profile (green) indicates the correlation with the ranked gene list. The Normalized Enrichment Score (NES) and nominal p-values (nom.p) are shown. **c.** Genomic and Epigenetic landscape, and gene expression at NRIP1 locus. Tracks, from top to bottom, display: genome scale; contact matrices from Ishikawa cells treated with R5020 or E2 (blue dashed lines indicate chromatin loops); virtual 4C centered at the NRIP1 transcription start site (TSS); ENCODE CTCF binding sites in Ishikawa cells; UCSC gene annotations; ER and PR ChIP-seq signals in Ishikawa cells (black lines indicate peaks); ER ChIP-seq signals from non-users and tamoxifen users (black lines indicate peaks); H3K27ac ChIP-seq signals from non-users and tamoxifen users; and RNA-seq RPKM values from Ishikawa cells treated with T0, E2 (E), and R5020 (R). **d.** NRIP1 expression (FPKM) from RNA-seq data in The Cancer Genome Atlas (TCGA), including 394 endometrial adenomas and adenocarcinomas. Tumors are grouped by FIGO stage. Asterisks (*) indicate statistically significant differences in expression (p $\pm$ 0.05). **e.** NRIP1 expression (TPM) in Ishikawa cells treated with DMSO, 3nM, and 30nM of estrogen or tamoxifen. Asterisks (*) indicate statistically significant differential expression between treatment groups (p $\pm$ 0.05).

Figure 6. Tamoxifen Disrupts ER–PR Crosstalk in Endometrial Cancer. Model of tamoxifen effects on ER (blue)–PR (pink) crosstalk. In non-users (Left), ER binding overlaps with PR, mediating inflammatory pathway expression (red arrow height). In tamoxifen users (Right), PR regulation is lost and ER redistributes to PR-independent sites, enhancing the Estrogen Response Early pathway.

 Supplementary figures

Figure 1—Figure supplement 1. a. Venn diagram showing the overlap of PR binding sites (PR bs) in Ishikawa cells pretreated with E2 for 12 hours and those without pretreatment. The intersection represents PR bs shared between both conditions. **b.** Distribution of ERα ChIP-seq signal in tumor samples from tamoxifen users and non-users. Regions were defined within a 10 kb window (±5 kb) centered on ER binding sites (ER bs) identified in tamoxifen users, non-users, or in both groups (common sites). Top panel: Median accumulated ERα ChIP-seq signal (CPM) across each ER bs group for tamoxifen users and non-users. Bottom panel: Heatmap showing ERα ChIP-seq signal intensity across all tamoxifen-user and non-user samples.

a

b

Figure 1—Figure supplement 2. Principal Component Analysis of ER ChIP-seq signal in tamoxifen user and non-user samples. a. Scree plot illustrating the proportion of variance in ER ChIP-seq signal explained by each principal component. **b.** Principal Component Analysis (PCA) plot showing PC1 versus PC2 based on ERα ChIP-seq signal across all tamoxifen user and non-user samples.

a

b

Figure 1—Figure supplement 3. ER and H3K27ac signal distribution in tumors from tamoxifen users and non-users. a. Regions were defined within a 10 kb window (±5 kb) centered on ER binding sites (ER bs) identified in tamoxifen users, non-users, or in both groups (common sites). Top panel: Median accumulated ERα ChIP-seq signal for each binding site group (non-users, tamoxifen users, and common). Bottom panel: Heatmap showing ERα ChIP-seq signal intensity across individual regions, grouped by ER bs category: non-users, tamoxifen users, and common sites **b.** Same analysis as in (a), showing H3K27ac ChIP-seq signal in tumors from tamoxifen users and non-users. Regions were defined as in (a).

a

NOTCH Signaling Deregulation in Cancer

NES: -1.209
Pval: 2.225e-01
FDR: 2.225e-01

non_users

Zero score at 6904   tamoxifen_user

b

Puyang *et al.* 2018

Figure 2—Figure supplement 1. Notch signaling pathway analysis in tamoxifen-treated samples and Ishikawa cells. a. Gene set enrichment analysis (GSEA) of the Notch Signaling Deregulation in Cancer pathway using gene expression profiles from tamoxifen users and non-users. The enrichment profile (green) illustrates the correlation of gene expression with the ranked gene list. Normalized enrichment scores (NES) and nominal p-values (nom. p) are shown. **b.** Expression levels (TPM) of genes from the Notch Receptor Signaling and Notch Signaling Deregulation in Cancer pathways in Ishikawa cells treated with control (DMSO), tamoxifen (3nM and 30nM), and estrogen (3nM and 30nM) (E2.3nM and E2.30nM). Asterisks (*) indicate significant differential expression between E2 (30nM) and DMSO (p ≤ 0.05).

Figure 3—Figure supplement 1. MYC expression in Ishikawa cells. MYC expression (TPM) in Ishikawa cells treated with DMSO, 3nM, and 30nM of estrogen or tamoxifen. Asterisks (*) indicate statistically significant differential expression between treatment groups ($p \leq 0.05$).

Figure 4—Figure supplement 1. Relationship between ER and PR binding sites in Ishikawa cells. Histogram displaying the distribution of ER binding sites (ER bs) based on their distance to the nearest PR bs in Ishikawa cells. The inset highlights the subset of ER bs located within 2.5 kb of a PR bs.

a

b

c

Figure 4—Figure supplement 2. Pathway enrichment associated with ER and PR binding site co-occupancy in Ishikawa cells and tumor samples. a. Top enriched pathways for genes located in TADs containing at least one Ishikawa ER binding site (ER bs) that overlaps with a PR binding site (PR bs) and is also present as an ER bs in either tamoxifen users or non-users. **b.** Top enriched pathways for genes located near Ishikawa ER bs that do not overlap with PR bs but are also present as ER bs in either tamoxifen users or non-users. **c.** Top enriched pathways for genes located near Ishikawa ER bs that overlap with PR bs and are also present as ER bs in either tamoxifen users or non-users.

Figure 5—Figure supplement 1. Epigenetic landscape and gene expression of GREB1 (a), CCNA1 (b), and DHRS3 (c), key genes in the Estrogen Early Response pathway. From top to bottom, tracks display: genome scale; contact matrices of Ishikawa cells treated with T0, E2, and R5020, along with their corresponding eigenvalues; ENCODE CTCF binding sites in Ishikawa cells (ENCSR000BQE); UCSC gene annotations; ER and PR ChIP-seq signals from Ishikawa cells; ERα ChIP-seq signals from tamoxifen users and non-users; H3K27ac ChIP-seq signals from tamoxifen users and non-users; and RNA-seq RPKM values for Ishikawa cells of T0, E2, and R5020.