1 **Cancer cells surviving cisplatin chemotherapy increase stress-induced OMA1 activity**
2 **and mitochondrial fragmentation**
3

4 Melvin Li[1,2*], Chenille A. McCullum[1], Louis T.A. Rolle[1], Qin Ni[3,4], Zhuoxu Ge[3,4], Sean X. Sun[3,4],
5 Kenneth J. Pienta[1,2], Sarah R. Amend[1,2]
6
[1]Cancer Ecology Center, The James Brady Urological Institute, Johns Hopkins School of
Medicine, Baltimore, MD 21287
[2]Pharmacology and Molecular Sciences Program, Johns Hopkins School of Medicine,
Baltimore, MD 21287
[3]Institute for NanoBio Technology, Johns Hopkins University
[4]Department of Mechanical Engineering, Johns Hopkins University
*Correspondence: Melvin Li (mli154@jh.edu)
14
Author Contact Information
Melvin Li (mli154@jh.edu)
Chenille A. McCullum (cmccul17@jh.edu)
Louis T.A. Rolle (lrolle1@jh.edu)
Qin Ni (qni4@jhu.edu)
Zhuoxu Ge (zge6@jhu.edu)
Sean X. Sun (ssun@jhu.edu)
Kenneth J. Pienta (kpienta1@jhmi.edu)
Sarah R. Amend (samend2@jhmi.edu)

51 **ABSTRACT**
52

Cancer is one of the leading causes of deaths worldwide. Once cancer acquires therapy
resistance, it becomes the main driver of cancer lethality in patients. Thus, mechanisms of
therapy resistance must be investigated to improve patient outcomes. Mitochondria are critical
organelles in the cellular stress responses, undergoing dynamic morphological and functional
changes in response to external stimuli. We and others have identified a chemotherapy-
resistant cancer cell state where cells that survive treatment exhibit a dramatic increase in cell
size and remain non-proliferative for weeks. In this study, we demonstrate that cancer cells that
enter this resistant cell state in response to cisplatin increase OMA1 activity and decrease
mitochondrial fusion and function to combat oxidative stress. These findings contribute to further
understanding the role of the mitochondrial stress responses in therapy resistance in cancer
and provide a potential therapeutic avenue to targeting cancer cells that enter this
chemotherapy-resistant cell state.
65
66 **Keywords**: cancer, mitochondrial morphology, mitochondrial dynamics, OMA1, OPA1, oxidative
67 stress

68
69 **INTRODUCTION**
70
Mitochondria are essential organelles that play multi-faceted roles in maintaining cellular
homeostasis and stress responses[1,2]. These organelles undergo dynamic morphological
changes in response to metabolic and stress stimuli[1,2]. Mitochondrial fission is induced by
oxidative stress, resulting in a fragmented phenotype to decrease oxidative capacity or to
shuttle damaged mitochondria for degradation via mitophagy[3]. Mitochondrial fusion can be
induced by increased energetic demands, creating a hyperfused branching network with
increased oxidative capacity[3]. These organelles are dysregulated in many cancers, enabling
cells to better obtain resources and adapt to their environment[3]. Recent studies have shown that
alterations in mitochondrial morphology and dynamics confer resistance to a wide range of
chemotherapies across many tumor types[4-9]. Given the evidence on the role of mitochondria in
therapy-resistant cancers, targeting mitochondria is a promising strategy to combat cancer drug
resistance.
Our group and others have previously reported that cancer cells that survive
chemotherapy enter a resistant cell state in which they increase in size over time and are
resistant to subsequent treatments[10-14]. This phenomenon has been reported in mouse models
and in human tumors across various tumor types and is induced by multiple classes of
chemotherapy and environmental conditions[13-19]. In addition, the presence of this cell state in
patient primary tumors predict progression to metastatic disease, further strengthening its
significance *in vivo*[13,20,21]. Our previous work has identified that these cells exhibit altered
nuclear morphology, a dramatic increase and displacement of intracellular labile iron and
lysosomes, and increased antioxidant responses through NRF2 signaling[10,22]. Mitochondria
interact with all these processes, including communicating with the nucleus through retrograde
signaling, storing labile iron in iron-sulfur clusters, colocalizing with lysosomes to degrade
damaged portions of mitochondria, and undergoing turnover in response to NRF2 signaling[23-28].
Mitochondrial mass has been reported to increase in cancer cells entering this resistant
state from paclitaxel treatment in breast cancer, while another study observed increased
mitophagy in cancer cells surviving chemotherapy treatment in a head and neck cancer
model[29,30]. Beyond these observations, the molecular mechanisms of the mitochondrial
phenotypes in this chemotherapy-resistant cancer cell state have been largely unexplored. In
this study, we investigated mitochondrial morphology, dynamics, and stress responses in
prostate cancer cells surviving cisplatin chemotherapy. Through various imaging modalities and

protein expression analyses, we identified that the cells surviving chemotherapy exhibit
increased OMA1 activity, aberrant cristae morphology, and reduced mitochondrial fusion,
leading to decreased mitochondrial function. Our findings suggest that targeting OMA1 activity
could be a potential therapeutic vulnerability specific to this resistant phenotype.
106
107 **RESULTS**
108
Cancer cells surviving chemotherapy exhibit increased levels of reactive oxygen species
and increased mitochondrial fragmentation. The PC3 prostate cancer cell line was treated
with an LD50 dose of cisplatin (6 µM) for 72 hours[10,31]. At the end of treatment, cisplatin-
containing media was removed, and the surviving cells were then cultured in fresh media for up
to 10 days (**Fig. 1A**). During the 10-day period after cisplatin removal, surviving cells were not
proliferative and increased in cell size over time, resulting in a 40-fold increase in cell volume at
10 Days Post-Treatment Removal (10 Days PTR) (**Fig. 1B-C**). We have previously published
that cells continue to die after treatment removal, and the death rate plateaus at 10 Days PTR[10].
We initially hypothesized that these surviving cells have recovered from the cellular damage
induced by chemotherapy, including the increased oxidative stress and DNA damage induced
by chemotherapy treatment[32]. We found that the cells 10 Days PTR exhibited increased levels
of reactive oxygen species (ROS) per cell area when compared to untreated cells (**Fig. 1D-E**).
To validate this finding, both groups were treated with antioxidant N-acetyl cystine (NAC) for 24
hours. While the PC3 cells did not show any change in DCF-DA staining, cells 10 Days PTR
exhibited a decrease in DCF-DA staining under NAC-treated conditions, (**Supplementary Fig.**
S1).
Mitochondrial morphology is altered under oxidative stress[33]. We observed that cells 10
Days PTR have more mitochondria on a per cell basis, although this difference is non-significant
when normalizing to cell area (**Fig. 1F-H**). Quantification of mitochondrial morphology showed
that cells 10 Days PTR exhibited mitochondrial fragmentation, with a 1.5-fold decrease in
branches per mitochondrion and a 1.5-fold decrease in branch length per mitochondrion (**Fig.**
1F, I-J).
131
Surviving cells increase DRP1 localization to mitochondria. Mitochondrial fragmentation
can be the result of an increase in fission or a decrease in fusion[34]. We probed for the
expression of dynamin-related GTPases that are known to remodel mitochondrial structure.
Dynamin-related protein 1 (DRP1) is the main driver of mitochondrial fission at the outer
mitochondrial membrane (OMM) upon translocation from the cytosol to the mitochondria.
Phosphorylation of DRP1 at Ser616 promotes its translocation to the OMM, while
phosphorylation of DRP1 at Ser637 prevents the translocation from occurring[1]. Total DRP1
expression was unchanged in cells 10 days PTR compared to control and Ser616
phosphorylation was not detected in either group (**Fig. 2A**). Cells 10 Days PTR had decreased
DRP1 Ser637 phosphorylation compared to untreated cells, suggesting a decrease in DRP1
inhibition and an increase in DRP1 localization to mitochondria (**Fig. 2A**). DRP1 binds to
receptors mitochondrial fission factor (MFF), MiD49, or MiD51 on the OMM that allow DRP1 to
assemble on the mitochondrial surface for its fission activity[1,2,35]. Compared to control, cells 10
Days PTR showed increased expression of MFF and MiD49, but a decrease in MiD51
expression (**Fig. 2A**). We next assessed DRP1 mitochondrial localization via
immunofluorescence. Cells 10 Days PTR had increased DRP1-mitochondria localization when
compared to PC3 control cells (**Fig. 2B-F**). These results suggest that DRP1 may play a role in
the fragmentation phenotype that we observe in the mitochondria of surviving cells.
150
Mitochondria in surviving cells exhibit aberrant cristae morphology. In addition to DRP1-
mediated fission, mitochondrial fragmentation can occur through a decrease in fusion, which is

mainly regulated by Mitofusin 1 (MFN1), Mitofusin 2 (MFN2), and optic atrophy 1 protein (OPA1)
[1,35]. MFN1 and MFN2 are responsible for the fusion of the OMM [1]. There were no differences in
MFN1 and MFN2 expression between surviving cells and untreated cells (**Fig. 3A**). OPA1 is an
inner membrane remodeler that maintains cristae structure[1,35]. The long isoforms of OPA1
mediate fusion of the inner mitochondrial membrane and promote tight stacking of lamellar
cristae[36,37]. Under oxidative stress or mitochondrial depolarization, OPA1 is cleaved by OMA1
into short isoforms (L1 to S3, and L2 to S5), leading to decreases in mitochondrial fusion and
energetic capacity[37-39].
Cells that survived cisplatin treatment showed a decrease in the L2 isoform of OPA1 and
an increase of the S5 isoform, suggesting that L-OPA1 is cleaved to S-OPA1 by OMA1 (**Fig.
3A-B**). Upon stress conditions, OMA1 cleaves itself when activated to act on L-OPA1[39,40]. We
observed decreased expression of OMA1 in cells 10 Days PTR when compared to PC3 control
(**Fig. 3A, C**). Taken together, the altered levels of L2 and S5 isoforms of OPA1 and decreased
OMA1 levels in cells 10 Days PTR suggest OMA1 activity is higher in the cells that survive
cisplatin than control.
OPA1 is responsible for maintaining mitochondrial cristae structure. To evaluate this
directly, we performed transmission electron microscopy (TEM) to assess mitochondrial cristae
morphology. Mitochondria in untreated cells had tight lamellar cristae, while the mitochondria in
cells 10 Days PTR exhibited amorphous and wider cristae shape (**Fig. 3D**). Quantification of
cristae morphology showed that mitochondria in cells 10 Days PTR had fewer cristae, higher
mean cristae area, and higher cristae width (**Fig. 3E-G**). In addition, mitochondria in in cells 10
Days PTR were 3.5 times smaller by area than the mitochondria in untreated cells, supporting
the fragmentation phenotype that we observed in the immunofluorescence imaging in **Figure
1F, I-J** (**Fig. 3H**).
Overall, these data indicate that mitochondria in cells 10 Days PTR have impaired
cristae organization. The complexes of the electron transport chain (ETC) reside on the cristae
structure. The ETC complexes generate a negative membrane potential ($\Delta\Psi_m$) across the inner
mitochondrial membrane (IMM) as electrons are shuttled through[41]. To assess the functional
implications of this altered mitochondrial cristae structure, we measured the mitochondrial
membrane potential via live-cell staining with MitoTracker Red CMXRos. We observed that the
mitochondria in cells 10 Days PTR had decreased MitoTracker Red CMXRos signal when
compared to untreated cells, indicating that mitochondria are less active in cells 10 Days PTR
(**Fig. 3I-J**).
186
**Cells that survive cisplatin treatment decrease both mitochondrial fission and fusion
dynamics**. Given the changes in both DRP1 phosphorylation and OPA1 processing in cells 10
Days PTR, we next directly evaluated mitochondrial fission and fusion dynamics. We performed
time lapse imaging to capture mitochondrial fission and fusion events on a per-mitochondrion
basis, taking an image every 30 seconds for a total of 15 minutes. Overall, we observed that the
fission and fusion rate scaled with increasing mitochondrial area (**Fig. 4A-C**). When normalized
to mitochondrial area, cells 10 Days PTR had a 50% decrease in fission rate and a 50%
decrease in fusion rate than untreated cells (**Fig. 4A, D-E, Supplementary Video 1-4**).
195
196 **DISCUSSION**
197
Given the critical roles of mitochondria in cellular stress responses and their implications
in therapy-resistant cancer cells, recent advancements in identifying molecular regulators that
drive morphological and functional adaptations provide potential therapeutic avenues for
cancer[3,4,6,35,42]. Our group and others have reported the emergence of an enlarged and non-
proliferative cancer cell state in response to various chemotherapy treatments [10-14,22]. These
cells are resistant to subsequent chemotherapy and are associated with worse prognosis in

mouse models and in patients[13-20]. In this study, we evaluated the mitochondrial stress
responses in cancer cells that survive in the days following chemotherapy.
These cells 10 Days PTR had increased levels of ROS (**Fig. 1D-E**), consistent with
previous reports in bladder cancer where giant cells that emerge after cisplatin treatment
survive with increased levels of ROS[43]. The accumulation of ROS can induce oxidation of
macromolecules that are essential for cellular homeostasis, leading to DNA damage, misfolding
of proteins, and lipid peroxidation[44-47]. We have previously published that cells surviving
chemotherapy have continued DNA damage response (DDR), even at 10- and 15-Days PTR[10].
and that cells 10 Days PTR exhibit increased lipid peroxidation[22]. Moreover, this work also
showed that these surviving cells have an increase in nuclear factor erythroid 2-related factor
(NRF2)-mediated antioxidant response, further demonstrating that they are in a state of
oxidative stress[22].
Mitochondrial biogenesis, morphology, and function are altered under oxidative
stress[33,48-51]. While we observe that surviving cells have more mitochondria on a per cell basis,
when normalizing to cell size, there was no difference in mitochondrial number (**Fig. 1F-H**).
When analyzing the structure of the mitochondria, we found that the mitochondria in cells 10
Days PTR were more fragmented and had fewer branches per mitochondrion when compared
to the mitochondria of the untreated cells (**Fig. 1F, I-J**). We hypothesize that mitochondria in
surviving cells underwent fragmentation in response to the increase in ROS levels when
compared to untreated cells. Building on observations of this fragmentation phenotype in other
in cancer cells subjected to oxidative stress[48,50,52], our findings align with recent work by Song et
al. showing that cancer cells with a similar morphological phenotype that survive paclitaxel
treatment had more mitochondria per cell but exhibited a smaller and more fragmented
phenotype when compared to the untreated controls[53]. The mitochondria in those surviving cells
also moved more slowly and had fewer fission and fusion events than the mitochondria in the
control cells[53]. This suggests that these treatment-induced mitochondrial phenotypes could span
across therapies that act through different mechanisms, as paclitaxel is a microtubule stabilizer,
while cisplatin creates adducts in purine bases and ultimately causes DNA double strand
breaks[54,55].
When probing for the expression of the molecular regulators of mitochondrial fusion, we
identified differential expression of multiple isoforms of OPA1 when comparing cells 10 Days
PTR and untreated cells (**Fig. 3A-B**). The decrease in L2-OPA1 and increase in S5-OPA1
expression suggested that the stress-induced protease OMA1 was active in cells 10 Days PTR
(**Fig. 3A-B**). This is further supported by the decreased level of OMA1 itself, likely due to its
autocatalytic turnover. The S3 and S5 isoforms of OPA1 are fusion incompetent, and have been
shown to induce mitochondrial fission through interactions with DRP1 and other fission
machinery and were shown to be protective in events of oxidative stress[38,56]. The accumulation
of these OPA1 isoforms may be a mechanism to reduce mitochondrial fusion and decrease
OXPHOS to prevent further production of ROS, as mitochondria are major sources of
intracellular ROS when electrons are shuttled through the ETC[49].
We observed that the mitochondria in cells 10 Days PTR had larger and more
amorphous cristae than the tight, lamellar cristae in the mitochondria of untreated cells (**Fig. 3D-
H**). This aberrant morphology can be attributed to the decrease in L-OPA1 and accumulation of
S-OPA1 via OMA1 cleavage (**Fig. 3A-C**), as the short isoforms are not tethered to the IMM to
promote cristae folding[36,39,40]. Cristae folding increases the surface area of the IMM for oxidative
phosphorylation by bringing electron transport chain (ETC) complexes closer together and
concentrating the proton gradient[36,57,58]. Cells 10 Days PTR had decreased ETC efficiency of
(**Fig. 3I-J**). It has been reported that cells engineered to express S-OPA1 exclusively have
higher survival under oxidative stress than cells engineered to express only L-OPA1[56]. Cells
expressing solely L-OPA1 produced more superoxide radicals and therefore were sensitized to
oxidative stress[56]. Thus, the decrease in cristae folding in surviving cells further supports that

surviving cells may reduce OXPHOS and subsequent endogenous ROS production in response
to oxidative stress.
Mitochondrial morphology is mediated by dynamic fission and fusion processes. We
observed a surprising decrease in the mitochondrial fission rate in cells 10 Days PTR (**Fig. 4A,**
D). This was unexpected given the parallel findings of increased DRP1 localization to
mitochondria (via TOM20-DRP1 colocalization analysis), increased expression of MFF and
MiD49, and decreased expression of phospho-DRP1 Ser637 (**Fig. 2A-F**). This may suggest an
increase in the upstream signaling to promote mitochondrial fission, but a mechanism
downstream of DRP1 leading to the decreased fission rate we observed in the time lapse
imaging. There are several factors that affect mitochondrial fission after DRP1 docks onto the
various receptors on the OMM. The endoplasmic reticulum (ER) is a significant player in
mitochondrial fission. The molecular regulators of mitochondrial fission and fusion colocalize
with ER-mitochondrial contact sites to regulate mitochondrial dynamics[59,60]. The polymerization
of actin around the ER and the mitochondria and the transfer of calcium from the ER to
mitochondria mediate the constriction of the outer mitochondrial membrane that is required for
mitochondrial fission to occur[61-64]. Cells 10 Days PTR may have decreased ER-mitochondrial
contacts, which could explain the paradoxical increase in DRP1 localization to mitochondria and
a decrease in mitochondrial fission rate, but this has yet to be tested. Based on the TEM images
in **Figure 3D**, the ER appear more fragmented and further from mitochondria in cells 10 Days
PTR when compared to the ER in the PC3 control cells. Cytoskeletal filaments also play
important roles in regulating mitochondrial fission, although their effects may be cell type or
context specific. In a model of *X. laevis* melanocytes, depolymerizing microtubules with
nocodazole decreased mitochondrial distribution, motility, and shape dynamics across the entire
cell[65], while mitochondrial associations with microtubules inhibited mitochondrial fission in a
model of fission yeast[66]. How the cytoskeleton filaments affect mitochondrial dynamics in cancer
cells in this resistant state is unknown and warrants further investigation. In addition to
decreased mitochondrial fission rate, cells 10 Days PTR also exhibited a decrease in the rate of
mitochondrial fusion (**Fig. 4A, E**). This data coincides with the increased OMA1 activity,
decreased expression of the fusion-promoting L-OPA1 isoforms, and increased expression of
the fusion-incompetent S-OPA1 isoforms we observed from **Fig. 3A-C**. The decrease in
mitochondrial fusion could also lead to the fragmentation phenotype we observed in the
mitochondria of the cells 10 Days PTR (**Fig. 1F, I-J**).
In summary, our findings suggest that cancer cells surviving chemotherapy increase
OMA1 activity in response to oxidative stress. The stress-induced OMA1 activity cleaves OPA1
into its shorter isoforms, leading to a decrease in mitochondrial fusion, aberrant mitochondrial
cristae morphology, and decreased efficiency of oxidative phosphorylation, ultimately promoting
cell survival under increased ROS conditions (**Fig. 5**). Although there are currently no specific
OMA1 inhibitors developed, this work contributes to the need for the screening of compounds
against OMA1 activity, as other groups have solely used genetic mechanisms to inhibit OMA1 in
various cancer models[42,67-69]. Further work involves knocking out OMA1 in PC3 cells to assess
its role in the entrance to this cancer cell state. We hypothesize that these surviving cancer cells
are critical mediator of cancer lethality in patients. Additional characterization of the cellular
stress responses and molecular programs of cells in this resistant state will shine light on
potential vulnerabilities that can be leveraged to target them for destruction and improve patient
prognosis.
300
301 **MATERIALS AND METHODS**
302
Cell culture. The PC3 cell line was purchased from American Type Culture Collection (ATCC).
Cells were cultured in RPMI 1640 medium (ThermoScientific; Cat# 11875119) with 10% fetal

bovine serum (FBS) (Avantar; Cat# 97068-085) and 1% penicillin and streptomycin (P/S)
(Gibco; Cat#15140-122). Cells were incubated at 37℃ and 5% CO2.

Induction of resistant cancer cell state. PC3 cells were plated in 150 mm dishes (Corning;
Cat#: 353025) at a density of 1,250,000 cells per dish. 24 hours after seeding, cells were
treated with an IC50 dose (6µM) of cisplatin (Millipore Sigma; Cat#: 232120) in RPMI media with
10% FBS and 1% P/S for 72 hours. Drug-containing media was then removed, and fresh cell
culture media was added onto surviving cells. The surviving cells were monitored for 10 days,
with media changes every 3 days. Cells were analyzed 10-days post-treatment removal (PTR)
for all experiments.

Phase contrast microscopy. Cells were cultured to their desired timepoint in T75 flasks.
Phase contrast microscopy was performed on an EVOS M7000 (ThermoScientific) with a 10X
objective.

Cell volume measurement with Fluorescence eXclusion method. The microfluidic
Fluorescence Exclusion method (FXm) was used to measure the volume of PC3 untreated cells
and cells 10 Days PTR [101, 102]. In this method, microfluidic channels with known heights
were fabricated. Cells were seeded into the device in the presence of a membrane-
impermeable fluorescent dextran and imaged using an epifluorescence microscope. The
reduction in fluorescence intensity due to cell volume exclusion is proportional to the ratio
between the cell height and the channel height, allowing us to calculate cell volume. FXm
channel fabrication and volume calculation are described in previously published work by Ni et
al. and Rochman et al. [102, 103].

Immunofluorescence and Airyscan super-resolution imaging. Cells were plated onto poly-l-
lysine coated coverslips (Thomas Scientific; Cat#: 1217N79). Cells were fixed with methanol-
free 4% paraformaldehyde (ThermoFisher Scientific; Cat#28908) in 1X PBS for 15 minutes at
room temperature followed by permeabilization with 0.2% Triton-X 100 (ThermoFisher
Scientific; Cat#28314) in 1X PBS for 10 minutes at room temperature. Blocking was performed
with 10% Normal Goat Serum (abcam; Cat# ab7481) in 0.1% PBS-Tween20 for 30 minutes at
room temperature. Cells were incubated with primary antibody diluted in 10% Normal Goat
Serum in 0.1% PBS-Tween20 overnight at 4℃: anti-Tom20 (Santa Cruz; Cat# sc-17764) diluted
1:100 and anti-DRP1 (Cell Signaling Technology; Cat#: 8570S) diluted 1:100. Secondary
antibody incubation was performed using Goat anti-mouse IgG2a Cross-Absorbed Alexa Fluor
647 (ThermoFisher Scientific; Cat# A-21131) and Goat anti-rabbit IgG (H+L) Highly Cross-
Absorbed Secondary Alexa Fluor Plus 555 diluted 1:1000 in 10% Normal Goat Serum in 0.1%
PBS-Tween20 for 1 hour. Cells were also stained for F-actin during the secondary antibody
incubation with Alexa Fluor 488 Phalloidin (ThermoFisher Scientific; Cat# A22287). Following
secondary antibody and phalloidin incubation, cells were washed with 0.1% PBS-Tween20
three times and subsequently washed twice with 1X PBS. Coverslips were washed once in
dH2O prior to mounting. Coverslips were mounted onto slides with 10 µL of Prolong Glass
Antifade Mountant (ThermoFisher Scientific; Cat#: P36984) and samples were cured for at least
50 hours prior to imaging. All slides were imaged on a Zeiss LSM 880 microscope (Carl Zeiss)
in Airyscan SR mode with an 63X, numerical aperture 1.4 PlanApo oil objective. Images were
taken in a 3X3 tile z-stack at a resolution of 1768X1768 pixels per tile and an axial step size of
187 nm. 3D Airyscan Joint Deconvolution was performed on Zen software for image processing
and downstream image analyses were performed in ImageJ[70] and Imaris (Bitplane).

Mitochondrial morphology analysis. Images were imported into ImageJ[70] and converted to 8-
bit images. The ImageJ pipeline, Mitochondria Analyzer, was used to analyze mitochondrial

morphology[71]. In short, 3D thresholding was performed on each image within the Mitochondria
Analyzer plugin[71]. The settings used were Rolling = 1.0, Max Radius = 1.25, Block Size = 3, C-
Value = 3-5, Post-Processing Outlier Radius = 8.0 – 9.823 pixels. 2D slices were then obtained
from the 3D thresholded images for 2D mitochondrial morphology analysis. Features such as
branches per mitochondrion and branch length per mitochondrion were extracted in the
Mitochondria Analyzer plugin with the 2D thresholded slices as input images.

Colocalization analysis. In 2D, images were imported into ImageJ[70] and split into individual
channels and maximum Z-projections were generated for each channel. A line drawing was
used to generate intensity profiles along the length of the line. Intensities were normalized within
the DRP1 and Tom20 channel and overlayed together into a graph using RStudio[72] with
ggplot2[73]. For 3D analysis, images were analyzed in Imaris (Bitplane) for Tom20-DRP1
colocalization. Background subtraction was applied using a 10 µm filter, then surface renderings
were generated of the Tom20 and DRP1 channels. Using the object-to-object statistics
parameter, only DRP1 that was touching the Tom20 surface (0.0 µm away) was filtered for, and
a separate surface channel was created to only include those DRP1 surfaces. Percent
colocalization per cell was determined by taking the number of colocalized DRP1 surfaces
divided by the total number of DRP1 surfaces in the cell, multiplied by 100.

Mitochondrial membrane potential imaging and analysis. Cells were plated into a black 24-
well plate (ibidi; Cat# 82426) and allowed to adhere overnight. Cells were stained with 100 nM
MitoTracker Red CMXRos (ThermoScientific; Cat#: M7512) for 30 minutes at 37℃ and 5% CO2
and washed twice with 1X PBS. RPMI phenol red-free media containing 10% FBS and 1% P/S
was then added into each well of the 24-well plate and cells were imaged on an EVOS M7000
high content imager (ThermoScientific) with 37℃ and 5% CO2 incubation. Images were
analyzed in CellProfiler[74] for mean fluorescence intensity of MitoTracker Red CMXRos staining.

Transmission electron microscopy and mitochondrial cristae morphology analysis.
Transmission electron microscopy sample prep was performed with a previously published
protocol from Kostecka et al.[75]. Briefly, cells were cultured to the desired timepoints and were
fixed overnight at 4℃ in a solution of 2.5% glutaraldehyde, 3mM MgCl2, in 0.1 M sodium
cacodylate buffer, pH 7.2. Samples were rinsed with buffer followed by additional fixation in the
dark and on ice for 1 hour with 1% osmium tetroxide reduced with 1.5% potassium ferrocyanide
in 0.1 M sodium cacodylate buffer. Samples were rinsed with dH2O rinse and en bloc staining
was performed for one hour in 2% uranyl acetate (aq). A graded series of ethanol was used to
dehydrate the samples and samples were then embedded in Eponate resin overnight at 60℃. A
diamond knife on the Reichert-Jung Ultracut E ultramicrotome was used to cut 60 to 90 nm
sections. 2x1 mm formvar coated copper slot grids were used to pick up the sections. 2% uranyl
acetate (aq.) and 0.4% lead citrate were used to stain the grids before imaging on a
ThermoFisher Talos L120C at 120 kV with a ThermoFisher Ceta CCD (16 megapixel CMOS,
16-bit).

Time lapse microscopy for mitochondrial dynamics. Cells were plated into 8-well ibiTreat µ-
slides (ibidi; Cat#: 80826) and allowed to adhere overnight. Cells were stained with 150 nM
MitoTracker Green (ThermoScientific; Cat#: M7514) for 30 minutes at 37℃ and 5% CO2 and
washed twice with 1X PBS. RPMI phenol red-free media containing 10% FBS and 1% P/S was
then added into each well of the 8-well ibiTreat µ-slide. Cells were imaged on a Zeiss LSM 880
microscope (Carl Zeiss) in Airyscan SR mode with an 63X, numerical aperture 1.4 PlanApo oil
objective, at 37℃ and humidified 5% CO2. Cells were excited with a 488 nm laser and 2D
images were taken every 30 seconds for 15 minutes at a resolution of 1280X1280 pixels to
observe mitochondrial fission and fusion events. 2D Airyscan Processing was performed in Zen

software, and images were then subjected to background subtraction in ImageJ[70] using the
rolling ball method with a radius of 50 pixels. Mitochondrial fission and fusion events were
identified through Mitometer, a MATLAB-based software to track mitochondria over time[76].
410
Western blot. PC3 control cells and cells 10 Days PTR were lifted from T150 flasks with
TrypLETM Express (GibcoTM; Ref# 12604-013) in lysate form using a RIPA buffer (Sigma-
Aldrich; Cat#: R0278-500ML) with Halt Protease & Phosphatase cocktail (ThermoScientific;
Cat#: 78440) and 0.5M EDTA (ThermoScientific; Cat#: 1861274). Lysates were incubated at
4°C on a rotator for 30 minutes and then were subsequently spun in a 4°C centrifuge at 21,000
rcf and stored at -20°C. The Pierce BCA Protein Assay was used to equally allocate 50 µg
protein from each lysate for each lane being run in the gel. Each sample was made with the
appropriate amount of lysate, 5 µL 4X Loading Buffer (4X Laemmli Buffer (Bio-Rad #1610747),
beta-Mercaptoethanol (Bio-Rad; Cat#: 1610710XTU), and completed with UltraPure distilled
water (ThermoScientific; Cat#: 10977023). Samples were boiled in a thermocycler set to 99°C
with a lid temperature of 105°C for 10 minutes. A 4-20% Mini-PROTEAN TGX gel (Bio-Rad
#4561094) was loaded with each sample and Chameleon Duo Pre-stained Protein Ladder (LI-
COR Biosciences; Cat#: 928-60000). The gel run was set at 140 V for 60 minutes. For OPA1
probing, a 7.5% Mini-PROTEAN TGX gel (Bio-Rad #4561023) was used, and gels were run at
50 V for 300 minutes. Gels were transferred onto a 0.2 µm nitrocellulose membrane in the
Trans-Blot Turbo Transfer Pack (Bio-Rad; Cat#: 1704158) using the BioRad Trans-Blot Turbo
Transfer System. The membrane was blocked in 1X Casein Buffer (10X Casein Buffer (Sigma-
Aldrich; Cat#: B6429-500ML) diluted with UltraPure distilled water on a room temperature
shaker for 1 hour. The membrane was then incubated in a primary antibody specific to the
target protein at a 1:1000 dilution in 1X Casein Buffer with mouse β-actin primary antibody
(Sigma Aldrich; Cat#: A5441-0.5ML) at 1:5000 dilution overnight at 4°C. The primary antibodies
used were Phospho-DRP1 Ser616 (Cell Signaling Technology; Cat# 3455S), Phospho-DRP1
Ser637 (Cell Signaling Technology; Cat#: 4867S), DRP1 (Cell Signaling Technology; Cat#:
8570S), MFF (Cell Signaling Technology; Cat#: 84580S), MiD49 (Proteintech; Cat# 28718-1-
AP), MiD51 (Proteintech; Cat#: 20164-1-AP), OPA1 (Cell Signaling Technology; Cat#: 80471S),
OMA1 (Cell Signaling Technology; Cat#: 95473S), MFN1 (Cell Signaling Technology; Cat#:
14739S), MFN2 (Cell Signaling Technology; Cat#: 9482S). For the OPA1 blots, Vinculin was
used as the loading control (Cell Signaling; Cat#: 13901S) at a 1:1000 dilution. The following
day, the membrane was washed four times for 5 minutes each in 0.1% TBS-Tween 20
(UltraPure distilled water, 10X TBS (Quality Biological #351-086-101CS), 0.1% Tween 20
(Sigma-Aldrich #9005645)). Then, the membrane was incubated in a secondary antibody
diluted in 1X Casein Buffer for 1 hour on a room temperature shaker. The secondary antibodies
used were IRdye® 800CW goat anti-rabbit IgG at a 1:15000 dilution, and IRdye® 680RD goat
anti-mouse IgG at a 1:20000 dilution (LI-COR Biosciences; Cat# 926-32211 and 926-68070,
respectively). The membrane was washed another four times for 5 minutes each in 0.1% TBS-
Tween 20. Imaging was performed on the Azure Sapphire. (Azure Biosystems).
447
448 ***Statistics***. Statistical analyses were performed in GraphPad Prism version 9.1 (GraphPad
449 Software, LLC) and RStudio[72]. Statistical tests performed are reported in respective figure
450 legends. An alpha value of 0.05 was used for all tests. ns, *, **, ***, and **** indicated $p > 0.05$,
451 $p < 0.05$, $p < 0.01$, $p < 0.001$, and $p < 0.0001$, respectively. In the violin plots, the solid line
452 represents the median and the dotted lines represent the lower and upper quartiles.
453
454 **DATA AVAILABILITY**
455 Data is provided within the manuscript or in supplementary information files.
456
REFERENCES

1   Giacomello, M., Pyakurel, A., Glytsou, C. & Scorrano, L. A.-O. The cell biology of
mitochondrial membrane dynamics.

2   Wai, T. Is mitochondrial morphology important for cellular physiology? LID - S1043-
2760(24)00123-1 [pii] LID - 10.1016/j.tem.2024.05.005 [doi].

3   Jin, P. *et al.* Mitochondrial adaptation in cancer drug resistance: prevalence,
mechanisms, and management. *Journal of Hematology & Oncology* **15**, 97 (2022).
https://doi.org:10.1186/s13045-022-01313-4

4   Huang, Q. *et al.* Increased mitochondrial fission promotes autophagy and hepatocellular
carcinoma cell survival through the ROS-modulated coordinated regulation of the NFKB
and TP53 pathways.

5   Tomková, V., Sandoval-Acuña, C., Torrealba, N. & Truksa, J. Mitochondrial
fragmentation, elevated mitochondrial superoxide and respiratory supercomplexes
disassembly is connected with the tamoxifen-resistant phenotype of breast cancer cells.
Free Radical Biology and Medicine **143**, 510-521 (2019).
https://doi.org:https://doi.org/10.1016/j.freeradbiomed.2019.09.004

6   Decker, C. W. *et al.* Mitofusin-2 mediates doxorubicin sensitivity and acute resistance in
Jurkat leukemia cells.

7   Li, S. A.-O., Wu, Y., Ding, Y., Yu, M. & Ai, Z. CerS6 regulates cisplatin resistance in oral
squamous cell carcinoma by altering mitochondrial fission and autophagy.

8   Han, Y. *et al.* Mitochondrial fission causes cisplatin resistance under hypoxic conditions
via ROS in ovarian cancer cells. *Oncogene* **38**, 7089-7105 (2019).
https://doi.org:10.1038/s41388-019-0949-5

9   Chen, X. *et al.* Targeting Mitochondrial Structure Sensitizes Acute Myeloid Leukemia to
Venetoclax Treatment. *Cancer Discovery* **9**, 890-909 (2019).
https://doi.org:10.1158/2159-8290.CD-19-0117

10   Kim, C. J. *et al.* Nuclear morphology predicts cell survival to cisplatin chemotherapy.

11   Niu, N. *et al.* Linking genomic reorganization to tumor initiation via the giant cell cycle.
Oncogenesis **5**, e281-e281 (2016). https://doi.org:10.1038/oncsis.2016.75

12   Pienta, K. A.-O., Hammarlund, E. U., Brown, J. S., Amend, S. A.-O. & Axelrod, R. A.-O.
Cancer recurrence and lethality are enabled by enhanced survival and reversible cell
cycle arrest of polyaneuploid cells. LID - 10.1073/pnas.2020838118 [doi] LID -
e2020838118.

13   Fei, F. *et al.* The number of polyploid giant cancer cells and epithelial-mesenchymal
transition-related proteins are associated with invasion and metastasis in human breast
cancer. *Journal of Experimental & Clinical Cancer Research* **34**, 158 (2015).
https://doi.org:10.1186/s13046-015-0277-8

14   Schmidt, M. J. *et al.* Polyploid cancer cells reveal signatures of chemotherapy resistance.
Oncogene **44**, 439-449 (2025). https://doi.org:10.1038/s41388-024-03212-z

15   Zhang, X. *et al.* Targeting polyploid giant cancer cells potentiates a therapeutic response
and overcomes resistance to PARP inhibitors in ovarian cancer. *Science Advances* **9**,
eadf7195   https://doi.org:10.1126/sciadv.adf7195

16   Zhang, S. *et al.* Generation of cancer stem-like cells through the formation of polyploid
giant cancer cells. *Oncogene* **33**, 116-128 (2014). https://doi.org:10.1038/onc.2013.96

17   Mannan, R. *et al.* Polypoidal giant cancer cells in metastatic castration-resistant prostate
cancer: observations from the Michigan Legacy Tissue Program. *Medical Oncology* **37**,
16 (2020). https://doi.org:10.1007/s12032-020-1341-6

18   Amend, S. A.-O. *et al.* Polyploid giant cancer cells: Unrecognized actuators of
tumorigenesis, metastasis, and resistance.

19   Lopez-Sánchez, L. M. *et al.* CoCl2, a mimic of hypoxia, induces formation of polyploid
giant cells with stem characteristics in colon cancer.

20   Trabzonlu, L., Pienta, K. J., Trock, B. J., De Marzo, A. A.-O. & Amend, S. A.-O. Presence of
cells in the polyaneuploid cancer cell (PACC) state predicts the risk of recurrence in
prostate cancer.

21   Alharbi, A. M., De Marzo Am Fau - Hicks, J. L., Hicks Jl Fau - Lotan, T. L., Lotan Tl Fau -
Epstein, J. I. & Epstein, J. I. Prostatic Adenocarcinoma With Focal Pleomorphic Giant Cell
Features: A Series of 30 Cases.

22   Loftus, L. V., Rolle, L. T. A., Wang, B., Pienta, K. J. & Amend, S. R. Dysregulation of Labile
Iron Predisposes Chemotherapy Resistant Cancer Cells to Ferroptosis. *International*
Journal of Molecular Sciences **26** (2025).

23   Yang, D. & Kim, J. A.-O. Mitochondrial Retrograde Signalling and Metabolic Alterations in
the Tumour Microenvironment. LID - 10.3390/cells8030275 [doi] LID - 275.

24   Braymer, J. J. & Lill, R. Iron-sulfur cluster biogenesis and trafficking in mitochondria.

25   Wong, Y. C., Ysselstein, D. & Krainc, D. Mitochondria–lysosome contacts regulate
mitochondrial fission via RAB7 GTP hydrolysis. *Nature* **554**, 382-386 (2018).
https://doi.org:10.1038/nature25486

26   Chen, Q. *et al.* Nanoscale monitoring of mitochondria and lysosome interactions for
drug screening and discovery. *Nano Research* **12**, 1009-1015 (2019).
https://doi.org:10.1007/s12274-019-2331-x

27   Wong, Y. C., Kim, S., Peng, W. & Krainc, D. Regulation and Function of Mitochondria–
Lysosome Membrane Contact Sites in Cellular Homeostasis. *Trends in Cell Biology* **29**,
500-513 (2019). https://doi.org:https://doi.org/10.1016/j.tcb.2019.02.004

28   Luchkova, A., Mata, A. & Cadenas, S. A.-O. Nrf2 as a regulator of energy metabolism and
mitochondrial function.

29   Sirois, I. *et al.* A Unique Morphological Phenotype in Chemoresistant Triple-Negative
Breast Cancer Reveals Metabolic Reprogramming and PLIN4 Expression as a Molecular
Vulnerability.

30   You, B. *et al.* AMPK-mTOR-Mediated Activation of Autophagy Promotes Formation of
Dormant Polyploid Giant Cancer Cells.

31   Mallin, M. M. *et al.* Cells in the polyaneuploid cancer cell (PACC) state have increased
metastatic potential.

32   Yang, H. *et al.* The role of cellular reactive oxygen species in cancer chemotherapy.

33   Xu, X., Pang, Y. & Fan, X. Mitochondria in oxidative stress, inflammation and aging: from
mechanisms to therapeutic advances. *Signal Transduction and Targeted Therapy* **10**, 190
(2025). https://doi.org:10.1038/s41392-025-02253-4

34   Berman, S. B., Pineda Fj Fau - Hardwick, J. M. & Hardwick, J. M. Mitochondrial fission
and fusion dynamics: the long and short of it.

Tilokani, L., Nagashima, S., Paupe, V. & Prudent, J. Mitochondrial dynamics: overview of molecular mechanisms.

Fry, M. Y. *et al.* In situ architecture of Opa1-dependent mitochondrial cristae remodeling.

Gilkerson, R., De La Torre, P. & St Vallier, S. Mitochondrial OMA1 and OPA1 as Gatekeepers of Organellar Structure/Function and Cellular Stress Response.

Anand, R. *et al.* The i-AAA protease YME1L and OMA1 cleave OPA1 to balance mitochondrial fusion and fission.

MacVicar, T. & Langer, T. A.-O. OPA1 processing in cell death and disease - the long and short of it.

Baker, M. J. *et al.* Stress-induced OMA1 activation and autocatalytic turnover regulate OPA1-dependent mitochondrial dynamics.

Zorova, L. D. *et al.* Mitochondrial membrane potential.

Rivera-Mejías, P. *et al.* The mitochondrial protease OMA1 acts as a metabolic safeguard upon nuclear DNA damage.

Adibi, R., Moein, S. & Gheisari, Y. Zoledronic acid targets chemo-resistant polyploid giant cancer cells. *Scientific Reports* **13**, 419 (2023). https://doi.org:10.1038/s41598-022-27090-1

Schieber, M. & Chandel, N. S. ROS function in redox signaling and oxidative stress.

Srinivas, U. S., Tan, B. W. Q., Vellayappan, B. A. & Jeyasekharan, A. D. ROS and the DNA damage response in cancer.

Malhotra, J. D. *et al.* Antioxidants reduce endoplasmic reticulum stress and improve protein secretion.

Wang, B. *et al.* ROS-induced lipid peroxidation modulates cell death outcome: mechanisms behind apoptosis, autophagy, and ferroptosis. *Archives of Toxicology* **97**, 1439-1451 (2023). https://doi.org:10.1007/s00204-023-03476-6

Iqbal, S. & Hood, D. A. Oxidative stress-induced mitochondrial fragmentation and movement in skeletal muscle myoblasts.

Ježek, J. A.-O. X., Cooper, K. F. & Strich, R. Reactive Oxygen Species and Mitochondrial Dynamics: The Yin and Yang of Mitochondrial Dysfunction and Cancer Progression. LID - 10.3390/antiox7010013 [doi] LID - 13.

Wu, S., Zhou F Fau - Zhang, Z., Zhang Z Fau - Xing, D. & Xing, D. Mitochondrial oxidative stress causes mitochondrial fragmentation via differential modulation of mitochondrial fission-fusion proteins.

Lee, H. C. & Wei, Y. H. Mitochondrial biogenesis and mitochondrial DNA maintenance of mammalian cells under oxidative stress.

Chang, X. *et al.* ROS-Drp1-mediated mitochondria fission contributes to hippocampal HT22 cell apoptosis induced by silver nanoparticles. *Redox Biology* **63**, 102739 (2023). https://doi.org:https://doi.org/10.1016/j.redox.2023.102739

Song, Q. *et al.* Evolutionary adaptation and asymmetric inheritance of polyploid giant cancer cells in esophageal squamous cell carcinoma. *Cancer Letters*, 217818 (2025). https://doi.org:https://doi.org/10.1016/j.canlet.2025.217818

Díaz, J. F., Valpuesta, J. M., Chacón, P., Diakun, G. & Andreu, J. M. Changes in Microtubule Protofilament Number Induced by Taxol Binding to an Easily Accessible

Site: INTERNAL MICROTUBULE DYNAMICS*. *Journal of Biological Chemistry* **273**, 33803-
33810 (1998). https://doi.org:https://doi.org/10.1074/jbc.273.50.33803

55   Dasari, S. & Tchounwou, P. B. Cisplatin in cancer therapy: molecular mechanisms of
action.

56   Lee, H. A.-O., Smith, S. B., Sheu, S. A.-O. & Yoon, Y. A.-O. The short variant of optic
atrophy 1 (OPA1) improves cell survival under oxidative stress.

57   Cogliati, S., Enriquez, J. A. & Scorrano, L. Mitochondrial Cristae: Where Beauty Meets
Functionality.

58   Decker, S. T. & Funai, K. Mitochondrial membrane lipids in the regulation of bioenergetic
flux.

59   Abrisch, R. G., Gumbin, S. C., Wisniewski, B. T., Lackner, L. L. & Voeltz, G. K. Fission and
fusion machineries converge at ER contact sites to regulate mitochondrial morphology.
LID - 10.1083/jcb.201911122 [doi] LID - e201911122.

60   Sonn, S. K. *et al.* ER-associated CTRP1 regulates mitochondrial fission via interaction with
DRP1. *Experimental & Molecular Medicine* **53**, 1769-1780 (2021).
https://doi.org:10.1038/s12276-021-00701-z

61   Chakrabarti, R. A.-O. *et al.* INF2-mediated actin polymerization at the ER stimulates
mitochondrial calcium uptake, inner membrane constriction, and division.

62   Ji, W.-k., Hatch, A. L., Merrill, R. A., Strack, S. & Higgs, H. N. Actin filaments target the
oligomeric maturation of the dynamin GTPase Drp1 to mitochondrial fission sites. *eLife*
4, e11553 (2015). https://doi.org:10.7554/eLife.11553

63   Gatti, P., Schiavon, C., Cicero, J., Manor, U. & Germain, M. Mitochondria- and ER-
associated actin are required for mitochondrial fusion. *Nature Communications* **16**, 451
(2025). https://doi.org:10.1038/s41467-024-55758-x

64   Moltedo, O., Remondelli, P. & Amodio, G. The Mitochondria-Endoplasmic Reticulum
Contacts and Their Critical Role in Aging and Age-Associated Diseases.

65   Fernández Casafuz, A. B., De Rossi, M. C. & Bruno, L. Mitochondrial cellular organization
and shape fluctuations are differentially modulated by cytoskeletal networks. *Scientific*
Reports **13**, 4065 (2023). https://doi.org:10.1038/s41598-023-31121-w

66   Choudhury, S. A.-O., Ananthanarayanan, V. A.-O. & Ayappa, K. A.-O. X. Coupling of
mitochondrial population evolution to microtubule dynamics in fission yeast cells: a
kinetic Monte Carlo study.

67   Daverey, A. *et al.* Depletion of mitochondrial protease OMA1 alters proliferative
properties and promotes metastatic growth of breast cancer cells. *Scientific Reports* **9**,
14746 (2019). https://doi.org:10.1038/s41598-019-49327-2

68   Chen, L. *et al.* Inhibition of mitochondrial OMA1 ameliorates osteosarcoma
tumorigenesis. *Cell Death & Disease* **15**, 786 (2024). https://doi.org:10.1038/s41419-
024-07127-1

69   Wu, Z. A.-O. *et al.* OMA1 reprograms metabolism under hypoxia to promote colorectal
cancer development.

70   Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image
analysis. *Nature Methods* **9**, 671-675 (2012). https://doi.org:10.1038/nmeth.2089

71   Chaudhry, A., Shi, R. & Luciani, D. A.-O. A pipeline for multidimensional confocal analysis
of mitochondrial morphology, function, and dynamics in pancreatic β-cells.

72   RStudio: Integrated Development Environment for R (Posit Software, PBC, Boston, MA,
2025).
73   Valero-Mora, P. M. ggplot2: Elegant Graphics for Data Analysis. *Journal of Statistical*
Software, Book Reviews **35**, 1 - 3 (2010). https://doi.org:10.18637/jss.v035.b01
74   Stirling, D. R. *et al.* CellProfiler 4: improvements in speed, utility and usability. *BMC*
Bioinformatics **22**, 433 (2021). https://doi.org:10.1186/s12859-021-04344-9
75   Kostecka, L. A.-O. *et al.* Cancer cells employ lipid droplets to survive toxic stress.
76   Lefebvre, A., Ma, D., Kessenbrock, K., Lawson, D. A.-O. & Digman, M. A.-O. Automated
segmentation and tracking of mitochondria in live-cell time-lapse images.
642

ACKNOWLEDGEMENTS

This work was also supported by the William and Carolyn Stutt Research Fund, Ronald Rose, MC Dean, Inc., William and Marjorie Springer, Mary and Dave Stevens, Louis Dorfman, and the Jones Family Foundation. We thank the Johns Hopkins School of Medicine Microscope Facility for the acquisition of transmission electron microscopy images with the ThermoFisher Talos L120C.

FUNDING

This work was supported by the US Department of Defense CDMRP/PCRP 367 (W81XWH-20-10353), the Prostate Cancer Foundation, and the Patrick C. Walsh Prostate Cancer Research Fund to SRA; and NCI grants U54CA143803, CA163124, CA093900, and CA143055, and the Prostate Cancer Foundation to KJP. This work was also supported by the Office of the Director and the National Institute of General Medical Sciences of the National Institutes of Health under award number S10OD023548.

AUTHOR CONTRIBUTIONS

Conceptualization: ML, KJP, SRA; Experimental work: ML, CAM, LTAR, QN, ZG; Data analysis and figure generation: ML; Writing – original draft preparation: ML; Writing – review and editing: ML, KJP, SRA. Funding acquisition: SXS, KJP, SRA.

COMPETING INTERESTS

KJP is a consultant to Cue Biopharma, Inc., an equity holder in PEEL Therapeutics, and a founder and equity holder in Keystone Biopharma, Inc. SRA is an equity holder in Keystone Biopharma, Inc.

ETHICS DECLARATIONS

Approval for animal experiments: Not applicable

Approval for human experiments: Not applicable

Consent to participate/consent to publish: Not applicable

SUPPLEMENTARY INFORMATION

Supplementary Figure S1. N-acetyl cysteine decreases levels of reactive oxygen species in cells 10 Days Post-Treatment Removal. **(a)** Representative DCF-DA fluorescence images of cells treated with N-acetyl cysteine and vehicle. (**b**) Quantification of mean fluorescence intensity of DCF-DA between indicated groups.

Supplementary Figure S2. Uncropped blot images from Figure 2a. Boxed areas delineate the cropped portion displayed in Figure 2a.

Supplementary Figure S3. Uncropped blot images from Figure 3a. Boxed areas delineate the
cropped portion displayed in Figure 3a.
Supplementary Video 1. Representative time lapse of mitochondrial dynamics in untreated
PC3 cells stained with MitoTracker Green.
Supplementary Video 2. Second representative time lapse of mitochondrial dynamics in
untreated PC3 cells stained with MitoTracker Green.
Supplementary Video 3. Representative time lapse of mitochondrial dynamics in cells 10 Days
Post-Treatment Removal stained with MitoTracker Green.
Supplementary Video 4. Second representative time lapse of mitochondrial dynamics in cells
10 Days Post-Treatment Removal stained with MitoTracker Green.
692
FIGURE LEGENDS
Figure 1. Cancer cells surviving chemotherapy have increased levels of reactive oxygen
species and exhibit mitochondrial fragmentation. (**a**) Timeline for the induction of the
chemotherapy-resistant cancer cell state. Created in BioRender. Li, M. (2025)
https://BioRender.com/vkhgx05 (**b**) Phase contrast images of untreated cells and cells 1- and
10-Days Post-Treatment Removal (PTR). (**c**) Cell volume measurements of untreated cells and
cells 1- and 10-Days PTR. (**d**) Phase contrast images and DCF-DA fluorescence images in
untreated cells and cells 10 Days PTR. (**e**) Quantification of mean fluorescence intensity of
DCF-DA staining. (**f**) Representative images of TOM20 and Phalloidin in untreated cells and
cells 10 Days PTR. Segmentation of TOM20 signal was performed to quantify mitochondrial
morphology features. (**g**) Quantification of number of mitochondria per cell. (**h**) Quantification of
mitochondria normalized to cell area. (**i**) Quantification of number of branches per
mitochondrion. (**j**) Quantification of branch length per mitochondrion. n = 679, 257, and 249 cells
for untreated, cells 1 Day PTR, and cells 10 Days PTR, respectively (**c**); n = 1141 and 362 cells
for untreated and cells 10 Days PTR, respectively (**e**); n = 37 and 22 cells for untreated and
cells 10 Days PTR, respectively (**g-j**). Data are presented as mean ± s.e.m. (**c**, **g-j**), while data
in violin plots were presented as median and corresponding interquartile ranges (**e**). p values
were calculated with a one-way ANOVA followed by a post-hoc Tukey's multiple comparisons
test (**c**), a two-tailed Mann Whitney test (**e**), and an unpaired Student's two-tailed t-test (**g-j**). ns
not significant, **** $p<0.0001$.
713
Figure 2. Cells 10 Days PTR increase DRP1 localization to mitochondria. (**a**)
Representative western blots of Phospho-DRP1 Ser637, Phospho-DRP1 Ser616, DRP1, MFF,
MiD49, MiD51, and β-actin loading control. n = 3 biological replicates. We note that due to
similarities in molecular weight and identical primary antibody species, each protein was probed
independently on separate blots with their own separate β-actin loading control. Original,
uncropped blots (including the β-actin loading control for each blot) are presented in
Supplementary Figure S2. (**b**) Max intensity projection images of TOM20 and DRP1 channels in
untreated cells and cells 10 Days PTR. (**c**) Normalized profile intensity plot of TOM20 and DRP1
signal in untreated cells. (**d**) Normalized profile intensity plot of TOM20 and DRP1 signal in cells
10 Days PTR. (**e**) 3-dimensional (3D) rendering of TOM20, DRP1, and Phalloidin signal in
untreated cells and cells 10 Days PTR. (**f**) Quantification of TOM20-DRP1 colocalization in 3D.
n = 11 cells each for untreated and cells 10 Days PTR (**f**). Data are presented as mean ± s.e.m.
in (**f**), and p values were calculated with an unpaired Student's two-tailed t-test (**f**). ** $p<0.01$.
727
Figure 3. Mitochondria in cells 10 Days PTR have aberrant cristae morphology. (**a**)
Representative western blots of OPA1, OMA1, Vinculin, and β-actin loading control. We note
that due to similarities in molecular weight and identical primary antibody species, each protein
was probed independently on separate blots with their own separate β-actin or vinculin loading
control. Original, uncropped blots (including the β-actin or vinculin loading control for each blot)

are presented in Supplementary Figure S3. **b**) Quantifications of OPA1 isoform protein
expression in untreated cells and cells 10 Days PTR. Samples derive from the same experiment
in each blot that was quantified. (**c**) Quantification of OMA1 expression in untreated cells and
cells 10 Days PTR. Samples derive from the same experiment in each blot that was quantified.
(**d**) Representative transmission electron microscopy (TEM) images of mitochondria in
untreated cells and cells 10 Days PTR. (**e**) Quantification of number of cristae per
mitochondrion. (**f**) Quantification of mean cristae area per mitochondrion. (**g**) Quantification of
cristae width. (**h**) Quantification of mitochondrial area between untreated cells and cells 10 Days
PTR. (**i**) Representative fluorescence images of MitoTracker Red CMXRos staining in untreated
cells and cells 10 Days PTR. (**j**) Quantification of mean fluorescence intensity of MitoTracker
Red CMXRos staining between indicated groups. n = 4 biological replicates (**b**); n = 3 biological
replicates (**c**); n = 42 and 59 mitochondria for untreated and cells 10 Days PTR, respectively (**e-**
f, h); n = 229 and 177 cristae for untreated and cells 10 Days PTR, respectively (**g**); n = 12,147
and 2502 cells for untreated and cells 10 Days PTR, respectively (**j**). Data are presented as
mean ± s.e.m. (**b-c**), while data in violin plots are presented as median and corresponding
interquartile ranges (**e-h, j**). p values were calculated with a two-tailed Kolmogorov-Smirnov test
(**e-h**), and a two-tailed Mann-Whitney test (**j**). * $p<0.05$, ** $p<0.01$, **** $p<0.0001$.
750
Figure 4. Cells 10 Days PTR decrease both mitochondrial fission and fusion dynamics.
(**a**) Representative fluorescence images of mitochondria every 60 seconds for 6 minutes in
untreated cells and cells 10 Days PTR. Yellow arrow and box indicate a fusion event, while a
blue arrow and box indicate a fission event. (**b**) Linear regression analysis of fission rate vs
mitochondrial area. (**c**) Linear regression analysis of fusion rate vs mitochondrial area. (**d**)
Quantification of fission rate in untreated cells and cells 10 Days PTR when normalized to
mitochondrial area. (**e**) Quantification of fusion rate in untreated cells and cells 10 Days PTR
when normalized to mitochondrial area. n = 37 and 17 cells for untreated and cells 10 Days
PTR, respectively (**b-c, d-e**). Data are presented as scatter plots (**b-c**) and mean ± s.e.m. (**d-e**).
R-squared values were calculated with the lm() and summary() functions in RStudio[72] (**b-c**). p
values were calculated with the lm() and summary() functions in RStudio[72] (**b-c**) and a two-tailed
Mann-Whitney test (**d-e**). * $p<0.05$, ** $p<0.01$.
763
Figure 5. Cancer cells surviving chemotherapy increase OMA1 activity and decrease
mitochondrial fusion and function to combat oxidative stress. Cancer cells that survive
chemotherapy enter a resistant cell state that increases their cell size and remain non-
proliferative. Cells in this state have increased levels of reactive oxygen species (ROS) and
exhibit fragmented mitochondria. Cells in this state increase OMA1 activity, cleaving L-OPA1 to
S-OPA1, ultimately disrupting the mitochondrial cristae structure and reducing mitochondrial
fusion. This altered cristae structure generates a diffuse proton gradient and decreases
mitochondrial membrane potential, which lowers oxidative capacity and endogenous ROS
production as a survival mechanism against oxidative stress in these cells. Created in
BioRender. Li, M. (2025) https://BioRender.com/41jh49x.

Fig. 1

a

0 hr | 6μM Cisplatin | 72 hr | Remove Cisplatin treatment | Days post-Cisplatin treatment removal | End

PC3 Cells (Untreated) — Day 1 — Day 10

b Days Post-Treatment Removal

Untreated | 1 | 10

c Cell volume (μm³) — Untreated, 1 Day PTR, 10 Days PTR ****

d Phase | DCF-DA — Untreated, 10 Days PTR

e DCF-DA — Normalized DCF-DA Mean Intensity — Untreated, 10 Days PTR ****

f Composite | Phalloidin | TOM20 | TOM20 Segmentation | TOM20 Segmentation (Zoom) — Untreated, 10 Days PTR

g Mitochondria per Cell — Mitochondria per cell — Untreated, 10 Days PTR ****

h Mitochondria per Cell Area — Mitochondria per per μm² — Untreated, 10 Days PTR ns

i Branches per mitochondrion — Branches per mitochondrion — Untreated, 10 Days PTR ****

j Branch Length — Branch length per mitochondrion (μm) — Untreated, 10 Days PTR ****

Fig. 2

a

	Untreated	10 Days PTR
Phospho-DRP1 Ser637 (78 kDa)		
Phospho-DRP1 Ser616 (78 kDa)		
DRP1 (78 kDa)		
MFF (25-35 kDa)		
MiD49 (49 kDa)		
MiD51 (51 kDa)		
β-actin (42 kDa)		

b

TOM20 | DRP1 | Composite | Composite (Zoom)

Untreated

10 Days PTR

50 μm | 50 μm | 50 μm | 5 μm

c Untreated Normalized Profile

Relative Intensity vs Distance (μm)

Protein
DRP1
TOM20

d 10 Days Post–Treatment Removal Normalized Profile

Relative Intensity vs Distance (μm)

Protein
DRP1
TOM20

e

3D Composite | 3D Composite (Zoom)

Untreated

10 Days PTR

20 μm | 2 μm
20 μm | 2 μm

f TOM20-DRP1 3D Colocalization

% Colocalization

**

Untreated | 10 Days PTR

TOM20 | Mitochondrial DRP1 | Cytoplasmic DRP1 | Phalloidin

Fig. 3

a

OPA1 (70-100 kDa): L1, L2, S3, S4, S5
Vinculin (124 kDa)
OMA1 (37 kDa)
MFN1 (82 kDa)
MFN2 (82 kDa)
β-actin (42 kDa)

Untreated / 10 Days PTR

○ Untreated
○ 10 Days PTR

b OPA1

Relative Densitometry vs OPA1 Isoforms (L1, L2, S3, S4, S5)
**** * *

c OMA1

Relative Densitometry: Untreated vs 10 Days PTR
****

d Mitochondria / Mitochondria (Zoom)

Untreated — 0.5 μm / 0.1 μm
10 Days PTR — 0.5 μm / 0.1 μm

e Cristae per Mitochondrion
Cristae per Mitochondrion: Untreated vs 10 Days PTR
**

f Mean Cristae Area per Mitochondrion
Mean Cristae Area per Mitochondrion (μm²): Untreated vs 10 Days PTR
*

g Cristae Width
Cristae Width (μm): Untreated vs 10 Days PTR
****

h Mitochondrial Area
Mitochondrial Area (μm²): Untreated vs 10 Days PTR
**

i MitoTracker Red CMXRos / MitoTracker Red CMXRos (Zoom)

Untreated — 100 μm / 50 μm
10 Days PTR — 100 μm / 50 μm

j Mitochondrial Membrane Potential ($\Delta\psi_m$)
Normalized MitoTracker Red Mean Intensity: Untreated vs 10 Days PTR
****

Fig. 4

a

Time | 0" | 60" | 120" | 180" | 240" | 300" | 360"

Untreated (2 μm scale bars)

10 Days PTR (2 μm scale bars)

Fission Fusion

b

Fission Rate vs Mitochondrial Area

- Untreated
- 10 Days PTR

$R^2 = 0.599$
p-value = 6.96e−12

Fission Rate (min^{-1})

Mitochondrial Area (μm^2)

c

Fusion Rate vs Mitochondrial Area

- Untreated
- 10 Days PTR

$R^2 = 0.66$
p-value = 8.67e−14

Fusion Rate (min^{-1})

Mitochondrial Area (μm^2)

d

Fission

*

Fission Rate (min^{-1} μm^{-2})

Untreated   10 Days PTR

e

Fusion

**

Fusion Rate (min^{-1} μm^{-2})

Untreated   10 Days PTR

Fig. 5