1 **Soybean *Rhg1* resistance to *Heterodera glycines* impedes the**

2 **parasitism of *Pratylenchus penetrans***

Deepak Haarith[1,2], Ann E. MacGuidwin[1], and Andrew F. Bent[1,*]

[1]Department of Plant Pathology, University of Wisconsin-Madison, Madison, Wisconsin,

USA 53706

[2]Present address: Entomology and Nematology Department, University of Florida,

Gainesville, Florida, USA 32611

*Corresponding author: Andrew F. Bent, E-mail: afbent@wisc.edu

Keywords: Root lesion nematode, soybean cyst nematode, SCN, *rhg1*, host resistance

Funding: Wisconsin Soybean Marketing Board award #MSN274578 2023-2024

Abstract

Rhg1 has been the most effective QTL deployed in soybeans to control soybean cyst

nematode (*Heterodera glycines*; SCN). However, the resistance mechanisms and

specificity of *Rhg1* towards SCN as opposed to other plant parasitic nematodes are not

well known. In this study we report that *Rhg1* can impede the parasitism of the root lesion

nematode *Pratylenchus penetrans* (Pp), which is commonly found along with SCN in

northern U.S. soybean fields. Two elite public soybean varieties, carrying either *rhg1-a* +

Rhg4 or *rhg1-b*, negatively impacted Pp populations measured at 30 dpi. This contribution

of *Rhg1* was more rigorously demonstrated using near isogenic lines for *rhg1-b*.

Additionally, in three of the four soybean SCN-susceptible genotypes tested we observed

significantly more Pp when Pp were co-inoculated with SCN, consistent with a previous

study. No significant changes in SCN cyst numbers upon co-inoculation with Pp were

observed for any of the *rhg1* resistant lines. No nematode-induced changes in *Rhg1*

transcript abundances were observed when sampling infected root sections in any of the

plant genotypes, and no changes in a jasmonate response indicator were observed.

However, the 3 dpi salicylate-mediated response to SCN or Pp became more robust

during SCN + Pp co-infection. Multiple hypotheses for further dissection and manipulation

of soybean resistance to *Pratylenchus* root lesion nematodes are suggested based on the

present findings.

Introduction

Root lesion nematodes (RLN; *Pratylenchus* spp.) are important agricultural pests (Jones et

al. 2013; Rosa de Araújo et al. 2025). They are migratory nematodes with a wide host

range including soybean. *Pratylenchus brachyurus* is often considered to be the most

damaging nematode pest of soybeans in South America (Castanheira et al. 2020;

Favoreto et al. 2019; Nomura et al. 2024). *Pratylenchus* populations survive cold winters in

dead roots through anhydrobiosis or slow desiccation (Townshend 1984). *Pratylenchus*

are some of the most common plant parasitic nematodes found in the soybean producing

midwestern U.S. states like Wisconsin, Minnesota, Iowa, and Illinois, with *Pratylenchus*

penetrans (Pp) being among the most common species, and a species for which soybean

yield losses have been documented (Chen et al. 2012; MacGuidwin and Bender 2012;

MacGuidwin et al. 2023; Saikai and MacGuidwin 2022; Schumacher and Bird 2025)

(Schmitt and Barker 1981). Unlike sedentary endoparasitic nematodes such as the

soybean cyst nematode (SCN, *Heterodera glycines*), Pp feed on soybean roots at multiple

sites as ectoparasites or endoparasites (Endo 1967; Saikai and MacGuidwin 2020). SCN

spend limited time exposed to environmental disturbances as they shelter inside host roots

or in cyst-encased eggs for much of their life cycle. In contrast, the migratory lifestyle of Pp

makes them more directly exposed to, and to some extent adapted to, changes in soil pH

and nutrients, soil moisture, soil type, own population density, microbial communities, and

competition from other nematodes.

SCN is the most economically important pathogen or pest on soybeans in the United

States (Bradley et al. 2021). Across all of the North Central U.S. and Ontario for ten recent

years, SCN is estimated to have caused approximately 4% soybean yield loss (about 1/4

of the total for all soybean pathogens) (Allen et al. 2017; Bradley et al. 2021).

Understandably, most efforts in U.S. soybean systems have been directed at managing

SCN and not *Pratylenchus* spp. Estimates of the impact of RLN on soybean production

are rare, but RLN and SCN frequently co-occur (MacGuidwin et al. 2023; Saikai and

MacGuidwin 2022; Schumacher and Bird 2025). Northern U.S. soybean fields are typically

rotated with maize, an excellent host for RLN. In one deeply studied set of Wisconsin

soybean fields, RLN was found to cause up to 4.5% yield loss by itself (Saikai and

MacGuidwin 2022), and about 3.79% yield loss on corn (MacGuidwin and Bender 2016).

The farm gate values of corn and soybean in Wisconsin alone was over USD 2.4 billion

and USD 1.3 billion respectively (USDA), which illustrates the economic impact of even a

1% yield loss to RLN.

Along with crop rotation, the most important tool for managing SCN over the past

three decades has been host resistance provided by the major-effect quantitative trait

locus (QTL) *Rhg1* (Resistance to *Heterodera glycines* 1) (Niblack et al. 2006) (Bent 2022;

Niblack et al. 2006). There are two widely known resistance-conferring haplotype variants

of *Rhg1* in soybean. The haplotype *rhg1-a* is commonly derived from "Peking" while *rhg1-b*

is often derived from "PI 88788", with the latter being used in over 95% of all commercial

nematode resistant varieties (McCarville et al. 2017; Niblack et al. 2006). The Peking-type

rhg1-a haplotype is less effective if it is not coupled with resistance-associated alleles at

the unlinked *Rhg4* or *rhg2* loci (Basnet et al. 2022; Brucker et al. 2005; Yu et al. 2016).

SCN populations that at least partially overcome three reference sources of *rhg1-b*

resistance are designated as HG type 2.5.7. Throughout soybean-producing areas of the

U.S., HG type 2.5.7 and other *rhg1-b*-virulent SCN populations are increasingly common

and are gradually increasing their virulence on *rhg1-b* (Kleczewski et al. 2023; McCarville

et al. 2017). In response, expanded use of Peking-type resistance and introduction of

varieties that carry PI 88788-type *rhg1-b* resistance stacked with additional minor-effect

resistance QTLs is anticipated (Bent 2022; Brzostowski and Diers 2017; Tylka et al. 2012)

The *rhg1-a* and *rhg1-b* haplotypes both carry three tightly linked genes that

contribute to SCN resistance (Cook et al. 2014; Cook et al. 2012a). *Glyma.18g022400*,

Glyma.18g022500 and *Glyma.18g022700* respectively encode AAT_{Rhg1} (putative amino

acid transporter), $\alpha\text{-}SNAP_{Rhg1}$ (vesicle trafficking mediator, also called GmSNAP18) and

$WI12_{Rhg1}$ (unknown function, carries a WI12 wound inducible protein domain). While the

causal *Rhg1* genes were identified a decade ago and their mechanism(s) of action are

being investigated, the resistance has been deployed for multiple decades. The $\alpha\text{-}$

$SNAP_{Rhg1}$ proteins differ between the *rhg1-a* and *rhg1-b* haplotypes, as does the copy

number of tandem repeats of the entire locus (Cook et al. 2014). The abundance of

resistance-associated $\alpha\text{-}SNAP_{Rhg1}$ protein is known to increase 10-15-fold in the

syncytium, the SCN feeding site (Bayless et al. 2016; Bayless et al. 2019). AAT_{Rhg1} protein

increases an order of magnitude along the SCN root penetration path, where a substantial

pool of AAT_{Rhg1} is associated with macrovesicles (Han et al. 2023). Less is known about

how the $WI12_{Rhg1}$ interacts with SCN, although impacts on root gibberellin levels have

been discovered (Dong and Hudson 2022). Silencing or knocking out any one of the three

genes increases susceptibility to SCN (Cook et al. 2012b; Dong and Hudson 2022; Liu et

al. 2017; Schumacher and Bird 2025). Soybean *Rhg1* genes also conferred resistance to

other cyst nematodes when moved into Arabidopsis and potato (Butler et al. 2019; Butler

et al. 2021).

We hypothesized that *Rhg1* could also provide some resistance to Pp and that the

commercially popular *rhg1-b* may have been beneficial against Pp. Melakeberhan and

Dey previously reported that the infection rate of *H. glycines* in some instances decreased

with increasing *P. penetrans* proportions, and that the rate of *P. penetrans* infection

increased when co-inoculated at a 1:1 ratio with *H. glycines* but did not increase at higher

or lower co-inoculation ratios (2003). That work was done using an SCN-susceptible

soybean cultivar. Schumacher and Bird recently reported some instances where the

presence of Pp increased SCN population densities on *rhg1-a* resistance sources, but with

no overall consistent trend (Schumacher and Bird 2025). Interactions between *Rhg1* and

other pathogens are only beginning to receive study, although most soybeans grown in

recent years across the U.S. carry this genetic resistance (Lopez-Nicora et al. 2023). On

soybean, Niblack and colleagues previously found that resistance to SCN was not reduced

during co-infections with the root knot nematode *M. incongnita* (Niblack et al. 1986).

Significantly, the *rhg1-a + rhg2 (GmSNAP11)* combination has recently been shown to

enhance quantitative resistance against reniform nematodes (Usovsky et al. 2021;

Usovsky et al. 2022)

In this study we assessed (i) the success of Pp and an SCN HG 2.5.7 population on

the soybean lines IL3025N (*rhg1-b*), IL3849N (*rhg1-a/Rhg4*) and Williams 82 (SCN-

susceptible *rhg1-c*), in terms of total nematodes recovered per plant at the end of 30 days,

(ii) the impact on nematode numbers when SCN and Pp were co-inoculated at 2:1 or 1:1

SCN:Pp ratios, and, using a set of near isogenic lines (NILs) for *rhg1-b*, (iii) how *rhg1-b*

impacts Pp and SCN nematode numbers and the transcript abundance response of the

three *rhg1-b* genes as well as salicylic acid (SA) and jasmonic acid (JA) response indicator

genes.

Materials and Methods

Sand cup preparation

135 Torpedo sand was sieved to remove residual gravel and the resulting sand was

136 moistened, placed in heat-tolerant bins at a maximum of 7.5 cm depth, and heated in a 65

137 °C oven for 48 h. The sand was cooled to room temperature and filled in 5 cm x 5 cm cups

138 (plastic pots with drainage holes) and saturated with 50 mL of tap water.

Plant genotypes

Seeds of Williams 82 (*rhg1-c*), IL3025N (*rhg1-b*) and IL3849N (*rhg1-a,* and *Rhg4*)

were used in the first set of cup assays. Seeds of Williams 82 and near-isogenic lines

Resistant NIL (*rhg1-b*; LD09-15604) and Susceptible NIL (*rhg1-c*; LD09-15630) were used

in the second set of cup assays and differential gene expression studies. The NILs were

developed by Brian Diers, University of Illinois at Urbana-Champaign, through four

backcrosses using LD00-3309 (*rhg1-b*) as the recurrent parent and IA3023 as donor of the

susceptible *rhg1-c* allele (resulting NIL accessions predicted to be >96% LD00-3309

genotype). All seeds were surface sterilized using chlorine vapor as described by Clough

and Bent (1998). Seeds were germinated in paper towels at 27°C for 4-5 days in the dark

until a 5 cm radicle had formed, and were then planted in sand cups. For nematode

infection cup assays, five replicates (individual plants in individual cups) for each plant

genotype and each nematode treatment were transplanted. Replicate blocks were

composed of one replicate of all treatments arranged in a tray. Plants within each replicate

block were randomized and covered with a humidity dome for the first 2 days of growth.

For expression analyses, three replicate blocks each were set up for 3 dpi and 7 dpi

sampling points. Each experiment was repeated twice (two separate dates).

Nematode inoculum

Eggs of an HG 2.5.7 Illinois field population of SCN maintained on IL3025N (FI

~70), termed "SCN HG 2.5.7 IL", were hatched in 4 mM $ZnCl_2$ solution at 27°C and dark

conditions. Freshly hatched J2 stage SCN were rinsed and re-suspended in de-ionised

water at 1000 J2/mL or 500 J2/mL. The *Pratylenchus penetrans* population was collected

in Wisconsin and maintained on 'IO Chief' sweet corn root explants. All life stages of *P.*

penetrans were extracted from corn root explants grown on Gamborg B5 medium using

the Baermann funnel method (Baermann 1917). The nematodes were rinsed and

resuspended in de-ionised water. *P. penetrans* inoculum batches contained approximately

35% adult females, 15% adult males, 40% third and fourth stage juveniles and 10% first

and second stage juveniles. The nematodes were inoculated into the sand cups 24 h after

transplanting the germinated soybean seedlings (VE) for most experiments, or for

experiments that used V1 plants, when the first set of trifoliate leaves had emerged.

Nematode suspensions were inoculated into the sand via a small hole made using a 200

uL pipette tip, parallel to and 1 cm away from the soybean seedling, which was then re-

covered with sand. The plants were covered with humidity domes for 12 h post inoculation.

Nematode count cup assays

Inoculated plants were maintained in a growth room with 14h daylight, day

temperature 27°C, night temperature 22°C, for 30 days. The plants were watered with 10

mL of tap water each day for the first week, 15 mL a day the second week, alternating

between 15 mL and 30 mL a day the third week and 30 mL a day thereafter. The cups

were fertilized once during week 3 and again a week later using MiracleGrow as per

manufacturer recommendations instead of tap water.

Nematodes were extracted from the sand using water flotation and decanted into a

nested sieve system consisting of a 2380 μm, 177 μm and 38 μm sieve (US standard

sieves 7, 80, and 400 respectively). Plant materials, coarse sand particles and other debris

retained on the 2380 μm sieve after vigorous rinsing with tap water were collected, then

the roots of all samples were chopped into ~1 cm pieces and incubated on Baermann

funnels with de-ionised water for 48 h at room temperature (25°C). 50 mL of water was

eluted from the bottom of the funnels and evaluated for Pp (all life stages) and SCN J2s

and males. Contents retained by the 177 μm sieve were retained for SCN cyst

enumeration under a stereoscope while contents from the 38 μm sieve were enumerated

for *Pratylenchus penetrans* of all life stages and SCN males and juveniles. Nematode

samples were further clarified using 70% (70 g into 100ml H_2O) sucrose solution prior to

microscopic enumeration. In NIL samples co-inoculated with both nematodes, contents

from 38 μm sieve were also evaluated for SCN J2 and male numbers.

NIL *rhg1* differential gene expression assay

For RNA extraction, roots from plants (Williams 82, Resistant NIL and Susceptible

NIL) inoculated with either SCN (500 J2) or Pp (500 worms – all life stages) or both (500

J2 + 500 Pp worms) as well as non-inoculated controls were rinsed in water and observed

under a stereoscope at 3 dpi or 7 dpi. For each sample, about 100 mg of root tissue from

regions with visible lesions from a single plant was placed in ice-cold cryovials with 5

ceramic beads and immediately frozen in liquid nitrogen. These samples were stored at -

80°C. Frozen tissue was processed at 2500 rpm for 30 s in a powerlyzer and returned to

liquid nitrogen for a minute. This process was repeated 2-3 times until tissue was

pulverized. RNA was then extracted from pulverized root tissue with Trizol (Zymo) using

manufacturer's protocol. cDNA was synthesized using Powerscript reagent and

degenerate primers from 1 ug of RNA for each sample. Real-time quantification and Ct

values were obtained for transcripts encoding AAT_{Rhg1}, $WI12_{Rhg1}$, $\alpha\text{-SNAP}_{Rhg1}$ (detects all

haplotypes), $\alpha\text{-SNAP}_{Rhg1}HC$ (detects only the HC/ *rhg1-b* haplotype), as well as for *AAT*

(*Glyma.10G109500*; JA response indicator), *NIMIN1* (*Glyma.10G010100*; SA response

indicator) and the internal standard gene *NREG1* (*Glyma.11G079600*) (Beyer et al.

2021).To determine the fold-change in transcript abundance in response to nematode

inoculations, the dCt values obtained using NREG1 as internal standard were normalized

to the average dCt values of all no-nematode samples for the same plant genotype within

the same experiment date to obtain ddCt values. Alternatively, to evaluate relative

transcript abundances across plant genotypes, the dCt values for all genotypes and

treatments were normalized to the average dCt values of no-nematode inoculated Sus.

NIL plants to obtain ddCt values.

Statistical analyses

All data were analyzed and visualized using R (version 4.3.0). Graphs show mean

and standard error of mean for the original non-transformed data. For statistical tests,

count data were square root transformed or log transformed to achieve normality,

multivariate analyses of variance were performed using the aov() function, and the means

were separated using TukeyHSD() in the package stats() with an α=0.05. Normality was

confirmed using Shapiro.test() function for all variables.

The SCN cyst data for the experiments were analyzed using the formula:

$$\sqrt[2]{Cysts} \sim [(experiments|replicates) + Plant\ genotype * Nematode\ inoculation]$$

Similarly, Pp count data was analyzed using the formula:

$$\sqrt[2]{P.penetrans}\ or \log(P.penetrans + 1)\ \sim [(experiments|replicates) + Plant\ genotype$$

$$* Nematode\ inoculation]$$

For the NIL nematode count evaluations, we used:

$$\log(SCN\ cysts) \sim [(experiments|replicates) + Plant\ genotype * Nematode\ inoculation]$$

$$\log(Total\ P.penetrans)\ \sim [(experiments|replicates) + Plant\ genotype * Nematode\ inoculation]$$

For the transcript abundance evaluations, statistical analyses were performed on the ddCt

values since the fold-change (2^-ddCt) values did not pass the Shapiro-wilk normality test.

The formula we used to evaluate transcript induction within a plant genotype in response

to nematodes:

$$ddCt \sim \frac{Experiments}{root\ replicates} + Plant\ genotype * Nematode\ treatment$$

We then visualized the data and analyses using Microsoft Office and Adobe Illustrator.

Results

Pp parasitism of soybean is enhanced by co-infection with SCN

Controlled environment single-species inoculation and co-inoculation experiments with Pp and SCN were carried out (Pp-only, SCN-only or Pp+SCN all within the same experiment). Experiments were first conducted using soybean genotypes that differ at hundreds of loci, including different SCN resistance haplotypes. Williams 82 (*rhg1-c*) is SCN-susceptible while IL3025N (also known as LD11-2170) carries *rhg1-b* and IL3849N (also known as LD07-3395bf) carries *rhg1-a* and *Rhg4*. Four independent experiments were conducted, two using a 1:1 ratio and two using a 2:1 SCN:Pp ratio for co-inoculations, with five replicates of each treatment within each experiment.

Upon co-inoculation with virulent SCN HG 2.5.7 IL, more Pp were recovered compared to the Pp-only inoculations (($p < 0.001$), orange vs. blue bars in Figure 1A and 1B; ANOVA tables associated with all figures are provided as Supplementary Tables). The experiments utilized an SCN HG 2.5.7 population that, as expected, reproduced well on the susceptible *rhg1-c* (Williams 82) and substantially overcame *rhg1-b* resistance (IL3025N) while exhibiting poor reproduction on a plant line carrying the *rhg1-a* and *Rhg4* combination (IL3849N) (green bars in Figure 1C and 1D; all differences between SCN-only inoculations (green bars) were significant at $p<0.001$). Within single soybean genotypes, the increase in Pp populations upon co-inoculation with SCN (orange vs. blue bars) was statistically significant only on the Williams 82 and IL3025N soybean genotypes that support substantial SCN growth and reproduction (Figure 1A and 1B). On IL3849N, which largely prevented growth of the HG 2.5.7 SCN population, the increases in Pp abundance in the co-inoculations as opposed to the inoculations only with Pp were not statistically

significant (Figure 1A and 1B). Fewer Pp were recovered from IL3849N than from

IL3025N in both single and co-inoculations (Figure 1A and 1B). SCN:Pp inoculation ratios

of 2:1 and 1:1 were tested and gave similar results (Figure 1A and 1B). In all of the

experiments in Figure 1, about twice as many Pp were recovered 30 days after inoculation

when Pp were co-inoculated with SCN HG 2.5.7, compared to Pp-only inoculations (the

ratio on the same plant genotype in the same experiment ranged from 1.7 to 2.3, across

all plant genotypes and both ratios of co-inoculation).

Within the same experiments as above, there were no significant differences in SCN

HG 2.5.7 IL cyst numbers between SCN-only inoculations and co-inoculations with Pp on

the same plant genotype (green vs. blue bars, Figure 1C and 1D). Nearly 10-fold more

cysts were recovered across treatments when 1000 (Figure 1D) rather than 500 (Figure

1C) SCN J2s were inoculated, for both solo inoculations and co-inoculations with Pp.

rhg1-b impedes the parasitism of Pp

A separate phenomenon was observed in the experiments of Figure 1, suggesting

that soybean SCN resistance might also impact RLN. When inoculated by itself in these

experiments, significantly fewer Pp were recovered after 30 days from the SCN-resistant

variety IL3849N than from Williams 82 (blue bars in Figure 1A and 1B). Pp abundance on

IL3025N was intermediate, being significantly lower only in the 1:1 SCN:Pp experiments

(Figure 1A and 1B).

A more rigorous test for an effect of the SCN resistance *rhg1-b* haplotype on Pp

was conducted using near-isogenic soybean lines (NIL) expected to carry the same

genotype at >96% of all loci but carrying either *rhg1-c* (Sus. NIL) or *rhg1-b* (Res. NIL) at

the *Rhg1* locus. The results supported the conclusion that *rhg1-b* can negatively impact Pp

(Figure 2). Mirroring the experiments of Figure 1, significantly fewer Pp were recovered

from the resistant NIL compared to the susceptible NIL when singly inoculated with Pp

(Figure 2A blue bars). When co-inoculated with both nematodes (Figure 2A orange bars),

fewer Pp were again recovered from Res. NIL plants than Sus. NIL plants but with a

difference that was at the margin of statistical significance (p = 0.056).

For SCN, multiple studies have shown that the proportion of female to male

nematodes formed is often lower on soybean plants with effective SCN resistance than on

susceptible plants (Colgrove and Niblack 2005). Intriguingly, with migratory Pp we also

observed that the ratios of females to males recovered from the *rhg1-b* Res. NIL were

significantly lower than those for *rhg1-c* Sus. NIL plants (Figure 2B). This was true in both

Pp-only and SCN co-inoculated treatments. When the proportion of Pp found in roots vs.

the surrounding sand at 30 dpi was enumerated, the *rhg1-b* Res. NIL had a higher

proportion of the Pp in the sand than the Sus. NIL, but intriguingly, that was only in the Pp

+ SCN co-inoculated samples (Figure 2C). For the Res. NIL line, a ~5-fold higher

proportion of Pp were in the sand than in the root upon co-inoculation with SCN, compared

to the Pp-only samples (Figure 2C). The relative presence of juvenile Pp 30 days after

inoculation was examined as another parasitism/resistance metric, because those

juveniles are predominantly progeny produced from the initial inoculum. At 30 days after

inoculation the number of juvenile-stage Pp (Figure 2D) and the proportion of juvenile-

stage vs. adult Pp (Figure 2E) again showed that presence of SCN shifted Pp population

demographics. Juvenile Pp were significantly less common on *rhg1-b* Res. NIL plants

compared to the *rhg1-c* Sus. NIL. For the juvenile:adult ratio they were less common on

Res. NIL plants only when Pp and SCN were co-inoculated (Figure 2E). The findings of

Figure 2 together indicate that *rhg1-b* can exert a negative impact on Pp, in the presence

of SCN but also in the absence of SCN.

Pp co-inoculation reduces SCN cyst numbers only in the susceptible *rhg1-c* lines

313   As expected, in SCN-only inoculations fewer SCN HG 2.5.7 IL cysts were

recovered from the resistant *rhg1-b* NIL plants compared to the susceptible *rhg1-c* NIL

plants or Wm82 (Figure 3A; data are from same experiments at Figure 2). Co-inoculation

of Pp with SCN reduced the number of cysts recovered on the SCN-susceptible *rhg1-c*

Williams 82 and Sus. NIL lines (Figure 3A). The difference was statistically significant only

for Williams 82 (Figure 3A), but a similar trend was also observed on Williams 82 in the

experiments of Figure 1A ($p = .094$) and on Sus. NIL in the experiments of Figure 3A ($p =$

$.070$). When the plants had partial resistance to SCN that reduced cyst production (Res.

NIL), co-inoculation with Pp did not further reduce cyst numbers (Figure 3A). In the same

experiments, the J2 and male SCN populations in roots and sand at 30 dpi were also

enumerated. The results were not consistent across the two SCN-susceptible plant

genotypes, as fewer total SCN J2s were recovered after co-inoculation with Pp only in the

case of the Sus. NIL plants (Figure 3B). Co-inoculation with Pp did not reduce J2 SCN

numbers for Williams 82 and Res. NIL plants (Figure 3B). The number of SCN males

recovered exhibited substantial variation, and no statistically significant differences were

observed between plant genotypes or between plants inoculated with only SCN and those

inoculated with both nematodes (Figure 3C).

No significant changes in *rhg1-b* transcript abundance detected in response to

nematode infection

In light of the observed interactions between SCN, Pp and *rhg1-b*, we investigated

nematode induction of increases in transcript abundance of the three *rhg1-b* genes as well

as indicator genes of SA-response and JA-response activity in soybeans, at 3dpi and 7dpi

(Figures 4 and 5). As has previously been observed (Cook et al. 2014), *Rhg1* transcripts

were approximately 5- to 20-fold more abundant for the high-copy *rhg1-b* genotype than

for the single-copy *rhg1-c* genotype (Supplemental Figure 1). Transcript abundances are

reported in Figures 4 and 5 as fold-change relative to the mean of the abundance for the

non-inoculated control of the same plant genotype within the same experiment (ddCt).

Figure 4 A-F show that even with two independent experiments that each sampled

infected zones from three independent root systems, we did not detect any significant

nematode-induced transcript abundance of any of the *rhg1-b* genes when plants were

inoculated at VE stage (emerging seedlings, cotyledons not yet separated). We also

observed no significant differences with plants inoculated at the V1 stage (fully expanded

trifoliate; Supplemental Figure 2).

Combined infection by SCN and Pp induces a stronger salicylic acid response

At either 3 dpi or 7 dpi, neither SCN, Pp nor their combination induced a significant

change in transcript abundance (ddCt) for the JA-response indicator gene '*AAT*' (Figure 5A

and 5B). Note that this established soybean JA-response indicator gene *AAT* (Beyer et al.

2021) is *Glyma.10G109500* and is not the *Rhg1* gene for AAT$_{Rhg1}$. In contrast to the JA

result, transcript abundances for the soybean SA-response indicator gene *NIMIN1*

(*Glyma.10G010100*) did exhibit statistically significant increases at 3 dpi in many of the

nematode-inoculated samples (Figure 5C). Only dual inoculation with SCN and Pp

consistently caused significant elevation of the SA indicator in all three soybean genotypes

(Figure 5C). The elevated SA response indicated by *NIMIN1* transcript levels was gone by

7 dpi (Figure 5D).

Discussion

Although SCN is the most economically important nematode of soybeans in the US,

in more tropical soybean producing states such as Brazil, RLN (especially *Pratylenchus*

brachyurus) has been reported to be more consequential (Castanheira et al. 2020;

Nomura et al. 2024). No soybean varieties labeled as having strong resistance to RLN

have been deployed but searches are underway (e.g., Chowdhury et al. 2022; Hamawaki

et al. 2019; Machado and Araúujo 2016). SCN and RLN frequently co-occur. Given the

importance of the soybean crop, it is crucial to understand if there are interactions

between the two nematodes on soybean and if that could lead to a change in their relative

abundance over time. Only limited surveys and studies are available on the tripartite

interaction of Soybean-SCN-RLN (Melakeberhan and Dey 2003; Schumacher and Bird

2025). At the time of writing this manuscript, no published reports are available on how the

deployed *Rhg1* resistance against SCN factors into this tripartite interaction.

Many soybean fields are infested with both SCN and RLN and the limited research

reported suggests that their interaction is complicated. MacGuidwin et al. suggested a

competitive relationship (population densities of both SCN and RLN suppressed by co-

occurence) based on an analysis of field survey data (MacGuidwin et al. 2023).

Melakeberhan and Dey tested four 1000-nematode inocula with varying proportions of *H.*

glycines and *P. penetrans* and found evidence of competition when SCN outnumbered Pp

3:1. When both species were equally represented the interaction appeared to be

antagonistic, as numbers of Pp increased and SCN numbers decreased as compared to

the single-species case (Melakeberhan and Dey 2003). Those experiments used HG 0

SCN on an SCN-susceptible soybean host. In the present study we also observed that

presence of SCN increased Pp numbers on susceptible hosts. This was true both on

Williams 82, which lacks SCN resistance, and on IL3025N, which carries *rhg1-b* but is

susceptible to the HG 2.5.7 IL (FI ~70) population that was used. Hypotheses for how SCN

assists Pp parasitism include effector-mediated suppression host defenses, and/or simply

by providing established entry points into the root.

We used HG 2.5.7 SCN both because it is informative with respect to *Rhg1* (it

overcomes *rhg1-b* but not *rhg1-a*), and because it is becoming the most common type of

SCN found in North Central U.S. soybean fields. The positive effect of SCN on Pp

population numbers was not observed on IL3849N, which carries *rhg1-a + Rhg4* SCN

resistance that is effective against HG 2.5.7 IL. The failure of SCN to increase Pp

numbers on IL3849N is a related topic for future study - it may for example be due to the

reduced number of SCN present on the resistant host after the first few days of infection,

and/or to the plant defense responses induced by SCN on the resistant host.

In our previous studies with *Rhg1* genes we had shown their capacity to increase

resistance to other cyst nematode species and genera (*Heterodera schactii, Globodera*

pallida, *Globodera rostochiensis*) when transgenically expressed in Arabidopsis or potato

(Butler et al. 2019). Similarly, the combined presence of SCN resistance loci *rhg1-a* and

rhg2 was shown to also provide quantitative resistance against reniform nematodes

(*Rotylenchulus reniformis*) (Usovsky et al. 2021; Usovsky et al. 2022). For the present

study we hypothesized that SCN-resistant soybeans deployed in fields may have been

partially controlling RLN. Our data support that early (30 dpi) establishment of Pp is

impeded by *rhg1-a*, as evidenced by the reduction of Pp population sizes in IL3849N

relative to Williams 82. Early Pp establishment was also impeded on IL3025N (*rhg1-b*)

when initial SCN and Pp population densities were similar, but was not significant when

SCN outnumbered Pp. We further hypothesized that the interaction between SCN and Pp

could reduce the efficacy of *Rhg1*. In support of that, relative to inoculation of the same

plants with just Pp, we found an increase in Pp recovery when simultaneous SCN

pressure was applied on plants that are susceptible to the SCN population (Pp + HG 2.5.7

SCN on Williams 82 and IL3025N). Interestingly, for the virulent SCN HG 2.5.7 IL

population that we tested, counts of SCN cysts were not reduced by co-inoculation with Pp

except at 1:1 ratio when *rhg1-a* or *rhg1-b* were absent. A similar dynamic between SCN

and Pp was observed by Melakeberhan and Dey on multiple soybean cultivars that lacked

the *rhg1-a* or *rhg1-b* resistance (Melakeberhan and Dey 2003).   We have not investigated

full-season resistance or yield protection (see also Pedro et al. 2024), but the reductions in

parasitism observed in the present study and that of Melakeberhan and Dey suggest a

potential contribution to full-season disease resistance against Pp. The field studies of

Saikai and MacGuidwin (Saikai and MacGuidwin 2022) were done using varieties with

rhg1-b resistance, so a yield loss impact of Pp on SCN-resistant soybean varieties has

been demonstrated. The present study suggests that Pp damage could be even worse if

use of *rhg1-b* soybeans was not prevalent, but further field-level and full-season or multi-

season studies are needed to investigate this point.

As noted above, in our experiments and those of Melakeberhan and Dey (2003) co-

inoculation with SCN helped Pp or was neutral, in all cases tested. Even on resistant

hosts, we observed no reduction of Pp at 30 dpi when co-inoculated with SCN. In

contrast, with wheat, Laserre et al. 1994 and Mokrini et al. 2018 both observed that on

susceptible hosts co-inoculation with cereal cyst nematode tended to reduce *Pratylenchus*

population sizes (Lasserre et al. 1994; Mokrini et al. 2018). Intriguingly Laserre and

colleagues also showed, on a susceptible host, a reduction in eggs and vermiform *P.*

neglectus upon inoculation of *H. avenae* onto a separate part of the root system. However,

on wheat accessions with resistance to *H. avenae* they observed that co-inoculation with

H. avenae led in some cases to higher numbers of *Pratylenchus*, similar to the trend in our

experiments (Lasserre et al. 1994).

The *rhg1-b* haplotype SCN resistance, unlike the *rhg1-a* haplotype resistance, is not

co-dependent on the *Rhg4* locus on chromosome 08 (Patil et al. 2019). We were able to

obtain near-isogenic lines for *rhg1-b* that, after four backcrosses, should be genetically

>96% identical, for a more well-controlled test of the contribution of *rhg1-b* to resistance

against Pp. As with IL3025N and IL3849N, we observed fewer Pp on Res. NIL plants

compared to the Sus. NIL plants when only Pp was inoculated, strongly linking *rhg1-b*

SCN resistance to partial Pp resistance as well. In addition to the overall reduction in Pp

numbers in the Res. NIL, we also observed reductions in the ratios of females to males

and juveniles to adults. RLN males and females differ in how they feed on the roots and

how much damage they can subsequently cause (Saikai and MacGuidwin 2020). Females

make larger lesions and burrow more frequently into the roots than the males. Hence

having overall fewer Pp and a population that is skewed towards adults, especially males,

may be less damaging to the crop. When considered together, these observations indicate

that *rhg1-b* imposes stress on Pp populations and further support the indication that *rhg1-b*

may be reducing the severity of damage to soybean caused by Pp.

In an earlier pair of studies, we had observed AAT_{Rhg1} protein abundance increasing

along SCN penetration path (Han et al. 2023), unlike the $\alpha\text{-SNAP}_{Rhg1}$ that exhibited a

strong abundance increase in the syncytium (Bayless et al. 2016). Because Pp do not

form syncytia, in the present study we hypothesized that *Rhg1-GmAAT* could be one of the

early response genes that functions against diverse nematodes. However, for the genes

encoding AAT_{Rhg1}, $\alpha\text{-SNAP}_{Rhg1}$ or $WI12_{Rhg1}$ we did not observe any statistically significant

transcript induction at 3 dpi for plants inoculated with Pp at the VE or the V1 stage, or at 7

dpi after VE stage inoculation. It is possible that any transcript induction is sufficiently local

to infected cells that the vast majority of non-infected cells in the root samples diluted out

any signal from cells at or immediately adjacent to the site of nematode infection. It is also

possible that expression of the *Rhg1* gene products is regulated in part at the post-

transcriptional level. This absence of an observed transcript induction is consistent with a

previous study where we saw no significant differences in transcript induction for α -

$\text{SNAP}_{Rhg1}\text{HC}$ at 7 dpi but did observe significant changes in protein abundance (Haarith et

al. 2025).

Transcript abundances for *NIMIN1*, a preferred soybean salicylic acid response

pathway indicator (Beyer et al. 2021), were upregulated significantly at 3 dpi in response

to SCN, and were more reliably upregulated when Pp co-infected alongside SCN. Perhaps

of greater interest, Pp alone did not reproducibly elicit this SA response while Pp+SCN did

(Figure 5), and this Pp co-infection with SCN enhanced both SA responses and Pp

population sizes (Figure 1). Antagonism between SA- and JA-mediated defense

responses has been widely reported (Glazebrook et al. 2003; Verma et al. 2025). There

are multiple instances of a "host mis-direction" virulence strategy in which pathogens or

pests that are more effectively defended against by the one set of defense responses

cause the plant to instead activate the other defense responses (Schweiger et al. 2014).

The non-sedentary (migratory) parasitism exhibited by Pp may be more similar to plant

attack by herbivorous insects than it is to a haustorium-forming biotrophic fungus or

oomycete, or a syncytium-forming cyst nematode. Our results suggest that Pp may lose

more to JA-mediated plant defenses than to SA-mediated plant defenses; future studies

could investigate if Pp in fact benefits from stronger elicitation of SA-mediated defenses.

The findings of (Usovsky et al. 2021) correlating α-SNAP-encoding genetic loci with

resistance to both SCN and reniform nematode support speculation that α-SNAP protein

variants may also contribute to resistance against Pp. A large body of evidence has

implicated SNAP proteins in SCN resistance, including major roles for the α-SNAP$_{Rhg1}$HC

protein encoded by *rhg1-b* and the α-SNAP$_{Rhg1}$LC/GmSNAP18 protein encoded by *rhg1-a*,

and more subtle contributions from a γ-SNAP (gamma-SNAP) encoded at *cqSCN-006*

(*Glyma.15G191200*) and loss of the α-SNAPs encoded at *rhg2*/*GmSNAP02*

(*Glyma.02G260400*), and *GmSNAP11* (*Glyma.11g234500*) (Bayless et al. 2016; Bent

2022; Butler et al. 2021; Cook et al. 2014; Cook et al. 2012a; Haarith et al. 2025;

Lakhssassi et al. 2017; Liu et al. 2017; Matsye et al. 2012; Usovsky et al. 2023) SNAP

proteins interact with each other as they form multimers on SNARE protein bundles that

mediate subcellular vesicle trafficking (Zhao et al. 2015). Future research could reveal that

combinations of SCN resistance-associated α-SNAP-encoding loci may provide stronger

Pp resistance than *rhg1-b*, and/or provide diversified resistance mediators that impose

distinct selection pressures on Pp populations. However, a leading hypothesis for the

impact on cyst nematodes of the non-canonical SCN resistance-associated α-SNAPs is

their cytotoxicity and elevated abundance in syncytia (Bayless et al. 2016; Bayless et al.

2018). Pp are not sedentary feeders that induce syncytia formation in the host. The *Rhg1*-

encoded AAT_{Rhg1} and $WI12_{Rhg1}$ proteins also contribute to SCN resistance and are equally

strong candidates for contributing to Pp resistance.

Taken together, the present results indicate that the *rhg1-b* haplotype impedes the

parasitism of Pp as well as SCN – a pair of plant parasitic nematode species that are

highly distinct taxonomically and behaviorally. The present controlled environment results

also show, in agreement with a past study, that on SCN-susceptible soybean genotypes

SCN can enhance Pp population sizes. There is measurable interaction between the two

nematodes regarding the efficacy of *rhg1-b* resistance, and in their induction of salicylic

acid responses within the plant. The findings suggest multiple avenues for future study and

manipulation of soybean resistance to *Pratylenchus* root lesion nematodes.

Author Contributions

DH conceptualized, designed, and performed the experiments, analyzed the data, and authored the manuscript. AEM provided Pp inoculum, methods and expertise in *Pratylenchus penetrans*, and edited the manuscript. AFB co-designed the experiments, and analyzed and co-authored the manuscript. AFB and DH are co-PIs on the grant that funded this study.

Acknowledgements

Funding for this study was provided primarily by a grant to AFB and DH from the Wisconsin Soybean Marketing Board, and also by grants to AFB from the United Soybean Board. We thank undergraduate John Stuntebeck for providing experiment support.

Figure Captions

Figure 1. Nematode counts on Wm82, IL3849N (*rhg1*-a) and IL3025N (*rhg1*-b) plants at

30 dpi with either or both SCN HG 2.5.7 IL and Pp. (A) Pp counts at 30 dpi at 1:1 SCN: Pp

inoculation (B) Pp counts at 30 dpi at 2:1 SCN:Pp inoculation. Blue bars correspond to Pp-

only inoculation and orange bars represent combined inoculation of both SCN and Pp. (C)

Numbers of SCN cysts recovered at 30 dpi at 1:1 SCN:Pp inoculation. (D) Numbers of

SCN cysts recovered at 30 dpi at 2:1 SCN:Pp inoculation. Green bars represent SCN-only

inoculation and the light-blue bars represent combined inoculation of both SCN and Pp.

Values shown are mean +/- standard error of mean. Statistical significance for differences

between means are displayed as p-values from ANOVA Tukey HSD tests performed on

data transformed to achieve normality (see Methods). The SCN HG 2.5.7 IL used partially

overcomes *rhg1-b* resistance and represents the most prevalent HG type in the USA. All

data are for two independent experiments with five replicates within each experiment,

using distinct batches of seedlings and fresh nematode inoculum.

Figure 2. Pp counts on Wm82, Sus. NIL and Res. NIL plants at 30 dpi in either Pp-only

inoculations (blue bars) or a combination of SCN and Pp at 1:1 ratio (orange bars). (A)

Counts of all Pp. (B) Ratio of Pp females to males. (C) Ratio of Pp in sand fraction to root

fraction. (D) Ratio of Pp juveniles (J1 through J4) to adults (male + female). (E) Counts of

Pp juveniles (J1 through J4). Values shown are mean +/- standard error of mean.

Instances of possible significant differences of means are marked with p-values from

ANOVA Tukey HSD tests performed on data transformed to achieve normality (see

Methods; p > 0.2 if no p value is shown). The SCN HG 2.5.7 IL used partially overcomes

rhg1-b resistance and represents the most prevalent HG type in the USA. All data are for

two independent experiments with five replicates within each experiment, using distinct

batches of seedlings and fresh nematode inoculum.

Figure 3. SCN cyst counts on Wm82, Sus. NIL and Res. NIL plants at 30 dpi in either

SCN-only inoculation (green bars) or a combination of SCN and Pp at 1:1 ratio (light-blue

bars). (A) SCN cyst counts. (B) SCN J2 counts (sand + root). (C) SCN male counts (sand

+ root). Values are mean +/- standard error of mean. Instances of possible significant

differences of means are marked with p-values from ANOVA Tukey HSD tests performed

on data transformed to achieve normality (see Methods) (p > 0.2 if no p value is shown).

The SCN HG 2.5.7 IL used partially overcomes *rhg1-b* resistance and represents the most

prevalent HG type in the USA. All data are for two independent experiments with five

replicates within each experiment, using distinct batches of seedlings and fresh nematode

inoculum.

Figure 4. *Rhg1* transcript induction (fold change) relative to (w.r.t.) no-nematode controls

within each plant genotype. The protein encoded by the analyzed transcript is noted in the

upper-left corner of each graph. Wm82, Sus. NIL and Res. NIL plants were inoculated at

VE stage and root samples from infection areas taken at: (A, C and E) 3 dpi, or (B, D and

F) 7 dpi. No statistically significant differences between samples were observed using

ANOVA and Tukey HSD test. The SCN HG 2.5.7 IL used partially overcomes *rhg1-b*

resistance and represents the most prevalent HG type in the USA. All data are for two

independent experiments with four replicate roots assessed within each experiment, using

distinct batches of seedlings and fresh nematode inoculum.

Figure 5. Induction (fold change) of JA-response indicator gene *Glyma.10G109500* and

SA-response indicator gene *Glyma.10G010100* transcripts at 3 dpi and 7dpi, relative to

(w.r.t.) no-nematode controls within each plant genotype within the same experiment.

Wm82, Sus. NIL and Res. NIL soybean accessions inoculated at VE stage; transcript

measured and sample dpi indicated in upper left corner of graph. Significant differences of

means are marked with p-values from ANOVA Tukey HSD tests ($p > 0.2$ if no p value is

shown). The SCN HG 2.5.7 IL used partially overcomes *rhg1-b* resistance and represents

the most prevalent HG type in the USA. All data are for two independent experiments with

four replicate roots assessed within each experiment, using distinct batches of seedlings

and fresh nematode inoculum.

Supplemental Figure 1. Same data as in Figure 4 but with data for all genotypes

normalized to mean Sus. NIL no nematode control within same experiment, to show

relative abundances between genotypes. *Rhg1* transcript induction (fold change)

compared to no-nematode controls within each plant genotype. The protein encoded by

the analyzed transcript is noted in the upper-left corner of each graph. Wm82, Sus. NIL

and Res. NIL plants were inoculated at VE stage and root samples from infection areas

taken at: (A, C, E and G) 3 dpi, or (B, D, F and H) 7 dpi. No statistically significant

differences between samples were observed using ANOVA and Tukey HSD test. The SCN

HG 2.5.7 IL used partially overcomes *rhg1-b* resistance and represents the most prevalent

HG type in the USA. All data are for two independent experiments with four replicate roots

assessed within each experiment, using distinct batches of seedlings and fresh nematode

inoculum.

Supplemental Figure 2. *Rhg1* transcript induction (fold change) for V1 plants at 3 dpi

compared to no-nematode controls within each plant genotype. The protein encoded by

the analyzed transcript is noted in the upper-left corner of each graph. Wm82, Sus. NIL

and Res. NIL plants were inoculated at V1 stage. (A, C, E) Data normalized to mean for

no-nematode control of same genotype in same experiment. (B, D, F) Same data as A, C,

E with data for all genotypes normalized to mean Sus. NIL no nematode control within

same experiment, to show relative abundances between genotypes. Instances of possible

significant differences of means are marked with p-values from ANOVA Tukey HSD tests

(p > 0.2 if no p value is shown). The SCN HG 2.5.7 IL used partially overcomes *rhg1-b*

resistance and represents the most prevalent HG type in the USA. All experiments were

repeated twice with three replicate roots assessed within each experiment, using distinct

batches of seedlings and fresh nematode inoculum.

Literature Cited

Allen, T. W., Bradley, C. A., Sisson, A. J., Byamukama, E., Chilvers, M. I., Coker, C. M., Collins, A. A., Damicone, J. P., Dorrance, A. E., Dufault, N. S., Esker, P. D., Faske, T. R., Giesler, L. J., Grybauskas, A. P., Hershman, D. E., Hollier, C. A., Isakeit, T., Jardine, D. J., Kelly, H. M., Kemerait, R. C., Kleczewski, N. M., Koenning, S. R., Kurle, J. E., Malvick, D. K., Markell, S. G., Mehl, H. L., Mueller, D. S., Mueller, J. D., Mulrooney, R. P., Nelson, B. D., Newman, M. A., Osborne, L., Overstreet, C., Padgett, G. B., Phipps, P. M., Price, P. P., Sikora, E. J., Smith, D. L., Spurlock, T. N., Tande, C. A., Tenuta, A. U., Wise, K. A., and Wrather, J. A. 2017. Soybean Yield Loss Estimates Due to Diseases in the United States and Ontario, Canada, from 2010 to 2014. Plant Health Progress 18:19-27.

Baermann, G. 1917. Eine einfache Methode zur Auffindung von Anklostomum (Nematoden) Larven in Erdproben. Geneeskundig tijdschrift voor Nederlandsch-Indië 57:131-137.

Basnet, P., Meinhardt, C. G., Usovsky, M., Gillman, J. D., Joshi, T., Song, Q., Diers, B., Mitchum, M. G., and Scaboo, A. M. 2022. Epistatic interaction between Rhg1-a and Rhg2 in PI 90763 confers resistance to virulent soybean cyst nematode populations. Theor Appl Genet 135:2025-2039.

Bayless, A. M., Zapotocny, R. W., Grunwald, D. J., Amundson, K. K., Diers, B. W., and Bent, A. F. 2018. An atypical N-ethylmaleimide sensitive factor enables the viability of nematode-resistant Rhg1 soybeans. Proc Natl Acad Sci U S A 115:E4512-E4521.

Bayless, A. M., Zapotocny, R. W., Han, S., Grunwald, D. J., Amundson, K. K., and Bent, A. F. 2019. The rhg1-a (Rhg1 low-copy) nematode resistance source harbors a copia-family retrotransposon within the Rhg1-encoded alpha-SNAP gene. Plant Direct 3:e00164.

Bayless, A. M., Smith, J. M., Song, J., McMinn, P. H., Teillet, A., August, B. K., and Bent, A. F. 2016. Disease resistance through impairment of alpha-SNAP-NSF interaction and vesicular trafficking by soybean Rhg1. Proc Natl Acad Sci U S A 113:E7375-E7382.

Bent, A. 2022. Exploring Soybean Resistance to Soybean Cyst Nematode. Annu Rev Phytopathol 60:379-409.

Beyer, S. F., Bel, P. S., Flors, V., Schultheiss, H., Conrath, U., and Langenbach, C. J. G. 2021. Disclosure of salicylic acid and jasmonic acid-responsive genes provides a molecular tool for deciphering stress responses in soybean. Sci Rep 11:20600.

Bradley, C. A., Allen, T. W., Sisson, A. J., Bergstrom, G. C., Bissonnette, K. M., Bond, J., Byamukama, E., Chilvers, M. I., Collins, A. A., Damicone, J. P., Dorrance, A. E., Dufault, N. S., Esker, P. D., Faske, T. R., Fiorellino, N. M., Giesler, L. J., Hartman, G. L., Hollier, C. A., Isakeit, T., Jackson-Ziems, T. A., Jardine, D. J., Kelly, H. M., Kemerait, R. C., Kleczewski, N. M., Koehler, A. M., Kratochvil, R. J., Kurle, J. E., Malvick, D. K., Markell, S. G., Mathew, F. M., Mehl, H. L., Mehl, K. M., Mueller, D. S., Mueller, J. D., Nelson, B. D., Overstreet, C., Padgett, G. B., Price, P. P., Sikora, E. J., Small, I., Smith, D. L., Spurlock, T. N., Tande, C. A., Telenko, D. E. P., Tenuta, A. U., Thiessen, L. D., Warner, F., Wiebold, W. J., and Wise, K. A. 2021. Soybean Yield Loss Estimates Due to Diseases in the United States and Ontario, Canada, from 2015 to 2019. Plant Health Progress 22:483-495.

Brucker, E., Carlson, S., Wright, E., Niblack, T., and Diers, B. 2005. Rhg1 alleles from
soybean PI 437654 and PI 88788 respond differentially to isolates of Heterodera
glycines in the greenhouse. Theor Appl Genet 111:44-49.
Brzostowski, L. F., and Diers, B. W. 2017. Pyramiding of Alleles from Multiple Sources
Increases the Resistance of Soybean to Highly Virulent Soybean Cyst Nematode
Isolates. Crop Science 57:2932-2941.
Butler, K. J., Chen, S., Smith, J. M., Wang, X., and Bent, A. F. 2019. Soybean Resistance
Locus Rhg1 Confers Resistance to Multiple Cyst Nematodes in Diverse Plant
Species. Phytopathology 109:2107-2115.
Butler, K. J., Fliege, C., Zapotocny, R., Diers, B., Hudson, M., and Bent, A. F. 2021.
Soybean Cyst Nematode Resistance Quantitative Trait Locus cqSCN-006 Alters the
Expression of a gamma-SNAP Protein. Mol Plant Microbe Interact 34:1433-1445.
Castanheira, C. M., Falcão, H. G., Ida, E. I., Dias-Arieira, C. R., and Bolanho Barros, B. C.
2020. Pratylenchus brachyurus parasitism on soybean: effects on productivity,
vegetative and nematological parameters and chemical properties. European
Journal of Plant Pathology 157:651-661.
Chen, S., Shaeffer, C. C., Wyse, D. L., Nickel, P., and Kandel, H. 2012. Plant-parasitic
Nematode Communities and Their Associations with Soil Factors in Organically
Farmed Fields in Minnesota. Journal of Nematology 44:361-369.
Chowdhury, I. A., Yan, G. P., Kandel, H., and Plaisance, A. 2022. Population Development
of the Root-Lesion Nematode on Soybean Cultivars. Plant Disease 106:2117-2126.
Colgrove, A. L., and Niblack, T. L. 2005. The effect of resistant soybean on male and
female development and adult sex ratios of. Journal of Nematology 37:161-167.
Cook, D. E., Bayless, A. M., Wang, K., Guo, X., Song, Q., Jiang, J., and Bent, A. F. 2014.
Distinct Copy Number, Coding Sequence, and Locus Methylation Patterns Underlie
Rhg1-Mediated Soybean Resistance to Soybean Cyst Nematode. Plant Physiol
165:630-647.
Cook, D. E., Lee, T. G., Guo, X., Melito, S., Wang, K., Bayless, A. M., Wang, J., Hughes, T.
J., Willis, D. K., Clemente, T. E., Diers, B., Jiang, J., Hudson, M., and Bent, A. F.
2012a. Copy Number Variation of Multiple Genes at Rhg1 Mediates Nematode
Resistance in Soybean. Science 338:1206-1209.
Cook, D. E., Lee, T. G., Guo, X., Melito, S., Wang, K., Bayless, A. M., Wang, J., Hughes, T.
J., Willis, D. K., Clemente, T. E., Diers, B. W., Jiang, J., Hudson, M. E., and Bent, A.
F. 2012b. Copy number variation of multiple genes at Rhg1 mediates nematode
resistance in soybean. Science 338:1206-1209.
Dong, J., and Hudson, M. E. 2022. WI12(Rhg1) interacts with DELLAs and mediates
soybean cyst nematode resistance through hormone pathways. Plant Biotechnol J
20:283-296.
Endo, B. Y. 1967. Comparitive Population Increase of Pratylenchus brachyurus and P.
zeae in Corn and in Soybean Varieties Lee and Peking. Phytopathology 57:118-
120.
Favoreto, L., Meyer, M. C., Dias-Arieira, C. R., Machado, A. C. Z., Santiago, D. C., and
Riberiro, N. R. 2019. Diagnosis and Management of Nematodes In Soybean Crop.
Informe Agropecuario 40:18-29.
Glazebrook, J., Chen, W. J., Estes, B., Chang, H. S., Nawrath, C., Metraux, J. P., Zhu, T.,
and Katagiri, F. 2003. Topology of the network integrating salicylate and jasmonate

signal transduction derived from global expression phenotyping. Plant Journal
34:217-228.
Haarith, D., Das, S., Nelson, E., Zapotocny, R., and Bent, A. F. 2025. Overexpression of α-
SNAPRhg1 Can Improve rhg1-a Mediated Soybean Resistance to Soybean Cyst
Nematode. Phytopathology.
Hamawaki, O. T., Hamawaki, R. L., Nogueira, A. P. O., Glasenapp, J. S., Hamawaki, C. D.
L., and da Silva, C. O. 2019. Evaluation of soybean breeding lineages to new
sources of root-knot nematode resistance. Cienc Agrotec 43.
Han, S., Smith, J. M., Du, Y., and Bent, A. F. 2023. Soybean transporter AAT Rhg1
abundance increases along the nematode migration path and impacts vesiculation
and ROS. Plant Physiol 192:133-153.
Jones, J. T., Haegeman, A., Danchin, E. G., Gaur, H. S., Helder, J., Jones, M. G., Kikuchi,
T., Manzanilla-Lopez, R., Palomares-Rius, J. E., Wesemael, W. M., and Perry, R. N.
2013. Top 10 plant-parasitic nematodes in molecular plant pathology. Mol Plant
Pathol 14:946-961.
Kleczewski, N., Colsgrove, A., Bowman, D., Harbach, C., and Plewa, D. 2023. A Survey of
Soybean Cyst Nematode Population Densities and Phenotypes in Illinois: 2018 and
2020. Plant Health Progress 24:221-225.
Lakhssassi, N., Liu, S., Bekal, S., Zhou, Z., Colantonio, V., Lambert, K., Barakat, A., and
Meksem, K. 2017. Characterization of the Soluble NSF Attachment Protein gene
family identifies two members involved in additive resistance to a plant pathogen.
Sci Rep 7:45226.
Lasserre, F., Rivoal, R., and Cook, R. 1994. Interactions between Heterodera-Avenae and
Pratylenchus-Neglectus on Wheat. Journal of Nematology 26:336-344.
Liu, S., Kandoth, P. K., Lakhssassi, N., Kang, J., Colantonio, V., Heinz, R., Yeckel, G.,
Zhou, Z., Bekal, S., Dapprich, J., Rotter, B., Cianzio, S., Mitchum, M. G., and
Meksem, K. 2017. The soybean GmSNAP18 gene underlies two types of resistance
to soybean cyst nematode. Nat Commun 8:14822.
Lopez-Nicora, H. D., Ralston, T. I., Diers, B. W., Dorrance, A. E., and Niblack, T. L. 2023.
Interactions Among Heterodera glycines, Macrophomina phaseolina, and Soybean
Genotype. Plant Dis 107:401-412.
MacGuidwin, A. E., and Bender, B. E. 2012. Estimating Population Densities of Root
Lesion Nematodes, Pratylenchus spp., from Soil Samples Using Dual Active and
Passive Assays. Plant Health Progress 13.
MacGuidwin, A. E., and Bender, B. E. 2016. Development of a Damage Function Model for
Pratylenchus penetrans on Corn. Plant Dis 100:764-769.
MacGuidwin, A. E., Smith, D. L., Conley, S. P., and Saikai, K. A. 2023. Prevalence of Pest
Nematodes Associated with Soybean (*Glycine max*) in Wisconsin from 1998 to
2021. J Nematol 55:20230053.
Machado, A. C. Z., and Araúujo, J. V. 2016. Broad-sense heritability and variance
component estimates for resistance in Brazilian soybean genotypes. Tropical Plant
Pathology 41:390-396.
Matsye, P. D., Lawrence, G. W., Youssef, R. M., Kim, K. H., Lawrence, K. S., Matthews, B.
F., and Klink, V. P. 2012. The expression of a naturally occurring, truncated allele of
an alpha-SNAP gene suppresses plant parasitic nematode infection. Plant Mol Biol
80:131-155.

McCarville, M. T., Marett, C. C., Mullaney, M. P., Gebhart, G. D., and Tylka, G. L. 2017.
Increase in Soybean Cyst Nematode Virulence and Reproduction on Resistant
Soybean Varieties in Iowa From 2001 to 2015 and the Effects on Soybean Yields.
Plant Health Progress 18:146-155.
Melakeberhan, H., and Dey, J. 2003. Competition between *Heterodera glycines* and
Meloidogyne incognita or *Pratylenchus penetrans*: Independent Infection Rate
Measurements. Journal of Nematology 35:1-6.
Mokrini, F., Viaene, N., Waeyenberge, L., Dababat, A. A., and Moens, M. 2018.
Investigation of resistance to *Pratylenchus penetrans* and *P. thornei* in international
wheat lines and its durability when inoculated together with the cereal cyst
nematode *Heterodera avenae*, using qPCR for nematode quantification. European
Journal of Plant Pathology 151:875-889.
Niblack, T. L., Hussey, R. S., and Boerma, H. R. 1986. Effects of Interactions among
Heterodera-Glycines, Meloidogyne-Incognita, and Host Genotype on Soybean Yield
and Nematode Population-Densities. Journal of Nematology 18:436-443.
Niblack, T. L., Lambert, K. N., and Tylka, G. L. 2006. A model plant pathogen from the
kingdom animalia: Heterodera glycines, the soybean cyst nematode. Annual
Review of Phytopathology 44:283-303.
Nomura, R. B. G., Lopes-Caitar, V. S., Hishinuma-Silva, S. M., Machado, A. C. Z., Meyer,
M. C., and Marcelino-Guimarães, F. C. 2024. Pratylenchus brachyurus: status and
perspectives in Brazilian agriculture. Tropical Plant Pathology 49:573-589.
Patil, G. B., Lakhssassi, N., Wan, J., Song, L., Zhou, Z., Klepadlo, M., Vuong, T. D., Stec,
A. O., Kahil, S. S., Colantonio, V., Valliyodan, B., Rice, J. H., Piya, S., Hewezi, T.,
Stupar, R. M., Meksem, K., and Nguyen, H. T. 2019. Whole-genome re-sequencing
reveals the impact of the interaction of copy number variants of the rhg1 and Rhg4
genes on broad-based resistance to soybean cyst nematode. Plant Biotechnol J
17:1595-1611.
Pedro, M. S., da Silva, S. A., Picoli, L. H., da Cunha, L. S., and Machado, A. C. Z. 2024.
Experimental and statistical approaches to evaluate the reaction of soybean to.
Tropical Plant Pathology 49:612-621.
Rosa de Araújo, T., de Assis dos Santos Diniz, F., Mendes, B. L., Almeida, L. C., and
Santiago, T. R. 2025. Pratylenchus zeae is the main root lesion nematode affecting
sugarcane in the world's most important producing regions. Physiological and
Molecular Plant Pathology 136.
Saikai, K., and MacGuidwin, A. E. 2020. Difference in Lesion Formation by Male and
Female *Pratylenchus penetrans*. Journal of Nematology 52.
Saikai, K., and MacGuidwin, A. E. 2022. Impact of Pratylenchus penetrans on Soybean
Grown in Wisconsin, U.S.A. Plant Dis 106:2904-2910.
Schmitt, D. P., and Barker, K. R. 1981. Damage and Reproductive Potentials of
Pratylenchus-Brachyurus and Pratylenchus-Penetrans on Soybean. Journal of
Nematology 13:327-332.
Schumacher, L. A., and Bird, G. W. 2025. Distribution of *Pratylenchus* spp. and *Heterodera*
glycines in Michigan and Combined Reproduction on Peking and PI 437654
Soybean. Plant Health Progress.

Schweiger, R., Heise, A. M., Persicke, M., and Muller, C. 2014. Interactions between the
jasmonic and salicylic acid pathway modulate the plant metabolome and affect
herbivores of different feeding types. Plant Cell Environ 37:1574-1585.
Townshend, J. L. 1984. Anhydrobiosis in *Pratylenchus penetrans*. Journal of Nematology
16:282-289.
Tylka, G. L., Gebhart, G. D., Marrett, C. C., and Mullaney, M. P. 2012. Evaluation of
soybean varieties resistant to soybean cyst nematode in Iowa in 2012. Iowa State
University Extension and Outreach.
Usovsky, M., Robert T. Robbins, Juliet Fultz Wilkes, Devany Crippen, Vijay Shankar, Tri D.
Vuong, Paula Agudelo, and Nguyen, H. T. 2022. Classification methods and
identification of reniform nematode resistance in known soybean cyst nematode
resistant soybean genotypes. Plant Disease 106:382-389.
Usovsky, M., Lakhssassi, N., Patil, G. B., Vuong, T. D., Piya, S., Hewezi, T., Robbins, R. T.,
Stupar, R. M., Meksem, K., and Nguyen, H. T. 2021. Dissecting nematode
resistance regions in soybean revealed pleiotropic effect of soybean cyst and
reniform nematode resistance genes. Plant Genome 14:e20083.
Usovsky, M., Gamage, V. A., Meinhardt, C. G., Dietz, N., Triller, M., Basnet, P., Gillman, J.
D., Bilyeu, K. D., Song, Q., Dhital, B., Nguyen, A., Mitchum, M. G., and Scaboo, A.
M. 2023. Loss-of-function of an alpha-SNAP gene confers resistance to soybean
cyst nematode. Nat Commun 14:7629.
Verma, K., Kumari, K., Rawat, M., Devi, K., and Joshi, R. 2025. Crosstalk of Jasmonic acid
and Salicylic acid with other Phytohormones Alleviates Abiotic and Biotic Stresses
in Plants. Journal of Soil Science and Plant Nutrition 25:4997-5019.
Yu, N., Lee, T. G., Rosa, D. P., Hudson, M., and Diers, B. W. 2016. Impact of Rhg1 copy
number, type, and interaction with Rhg4 on resistance to Heterodera glycines in
soybean. Theor Appl Genet 129:2403-2412.
Zhao, M., Wu, S., Zhou, Q., Vivona, S., Cipriano, D. J., Cheng, Y., and Brunger, A. T. 2015.
Mechanistic insights into the recycling machine of the SNARE complex. Nature
518:61-67.

Total RLN counts

SCN cyst counts

Figure 1. Nematode counts on Wm82, IL3849N (*rhg1-a*) and IL3025N (*rhg1-b*) plants at 30 dpi with either or both SCN HG 2.5.7 IL and Pp. (A) Pp counts at 30 dpi at 1:1 SCN: Pp inoculation (B) Pp counts at 30 dpi at 2:1 SCN:Pp inoculation. Blue bars correspond to Pp-only inoculation and orange bars represent combined inoculation of both SCN and Pp. (C) Numbers of SCN cysts recovered at 30 dpi at 1:1 SCN:Pp inoculation. (D) Numbers of SCN cysts recovered at 30 dpi at 2:1 SCN:Pp inoculation. Green bars represent SCN-only inoculation and the light-blue bars represent combined inoculation of both SCN and Pp. Values shown are mean +/- standard error of mean. Statistical significance for differences between means are displayed as p-values from ANOVA Tukey HSD tests performed on data transformed to achieve normality (see Methods). The SCN HG 2.5.7 IL used partially overcomes *rhg1-b* resistance and represents the most prevalent HG type in the USA. All data are for two independent experiments with five replicates within each experiment, using distinct batches of seedlings and fresh nematode inoculum.

Figure 2. Pp counts on Wm82, Sus. NIL and Res. NIL plants at 30 dpi in either Pp-only inoculations (blue bars) or a combination of SCN and Pp at 1:1 ratio (orange bars). (A) Counts of all Pp. (B) Ratio of Pp females to males. (C) Ratio of Pp in sand fraction to root fraction. (D) Ratio of Pp juveniles (J1 through J4) to adults (male + female). (E) Counts of Pp juveniles (J1 through J4). Values shown are mean +/- standard error of mean. Instances of possible significant differences of means are marked with p-values from ANOVA Tukey HSD tests performed on data transformed to achieve normality (see Methods; p > 0.2 if no p value is shown). The SCN HG 2.5.7 IL used partially overcomes *rhg1-b* resistance and represents the most prevalent HG type in the USA. All data are for two independent experiments with five replicates within each experiment, using distinct batches of seedlings and fresh nematode inoculum.

Figure 3. SCN cyst counts on Wm82, Sus. NIL and Res. NIL plants at 30 dpi in either SCN-only inoculation (green bars) or a combination of SCN and Pp at 1:1 ratio (light-blue bars). (A) SCN cyst counts. (B) SCN J2 counts (sand + root). (C) SCN male counts (sand + root). Values are mean +/- standard error of mean. Instances of possible significant differences of means are marked with p-values from ANOVA Tukey HSD tests performed on data transformed to achieve normality (see Methods) (p > 0.2 if no p value is shown). The SCN HG 2.5.7 IL used partially overcomes *rhg1-b* resistance and represents the most prevalent HG type in the USA. All data are for two independent experiments with five replicates within each experiment, using distinct batches of seedlings and fresh nematode inoculum.

Figure 4. *Rhg1* transcript induction (fold change) relative to (w.r.t.) no-nematode controls within each plant genotype. The protein encoded by the analyzed transcript is noted in the upper-left corner of each graph. Wm82, Sus. NIL and Res. NIL plants were inoculated at VE stage and root samples from infection areas taken at: (A, C and E) 3 dpi, or (B, D and F) 7 dpi. No statistically significant differences between samples were observed using ANOVA and Tukey HSD test. The SCN HG 2.5.7 IL used partially overcomes *rhg1-b* resistance and represents the most prevalent HG type in the USA. All data are for two independent experiments with four replicate roots assessed within each experiment, using distinct batches of seedlings and fresh nematode inoculum.

Figure 5. Induction (fold change) of JA-response indicator gene *Glyma.10G109500* and SA-response indicator gene *Glyma.10G010100* transcripts at 3 dpi and 7dpi, relative to (w.r.t.) no-nematode controls within each plant genotype within the same experiment. Wm82, Sus. NIL and Res. NIL soybean accessions inoculated at VE stage; transcript measured and sample dpi indicated in upper left corner of graph. Significant differences of means are marked with p-values from ANOVA Tukey HSD tests (p > 0.2 if no p value is shown). The SCN HG 2.5.7 IL used partially overcomes *rhg1-b* resistance and represents the most prevalent HG type in the USA. All data are for two independent experiments with four replicate roots assessed within each experiment, using distinct batches of seedlings and fresh nematode inoculum.