Soybean *Rhg1* resistance to *Heterodera glycines* hinders the parasitism of *Pratylenchus penetrans*

Journal:	*Plant Disease*
Manuscript ID	PDIS-12-25-2530-RE.R1
Manuscript Type:	Research
Date Submitted by the Author:	n/a
Complete List of Authors:	Haarith, Deepak; University of Florida, Entomology and Nematology MacGuidwin, Ann; University of Wisconsin-Madison, Plant Pathology Bent, Andrew; University of Wisconsin-Madison, Plant Pathology
Keywords:	Nematodes < Causal Agent, oilseeds and legumes < Field crops < Crop Type, cultivar/resistance < Disease management < Subject Areas

Soybean *Rhg1* resistance to *Heterodera glycines* ~~impedes~~

<u>hinders</u> the parasitism of *Pratylenchus penetrans*

Deepak Haarith[1,2], Ann E. MacGuidwin[1], and Andrew F. Bent[1,*]

[1]Department of Plant Pathology, University of Wisconsin-Madison, Madison, Wisconsin,

USA 53706

[2]Present address: Entomology and Nematology Department, University of Florida,

Gainesville, Florida, USA 32611

*Corresponding author: Andrew F. Bent, E-mail: afbent@wisc.edu

Keywords: host resistance, *rhg1*, root lesion nematode, SCN, soybean cyst nematode

Funding: Wisconsin Soybean Marketing Board award #MSN274578 2023-2024 and

United Soybean Board award #25-209-S-C-3-A

Deepak Haarith
Plant Disease

Abstract

Rhg1 has been the most effective QTL deployed in soybeans (*Glycine max*) to control

soybean cyst nematode (*Heterodera glycines*; SCN). However, the resistance

mechanisms and specificity of *Rhg1* towards SCN as opposed to other plant parasitic

nematodes are not well known. In this study we report that *Rhg1* can ~~impede~~ hinder the

parasitism of the root lesion nematode *Pratylenchus penetrans* (Pp), which is commonly

found along with SCN in northern U.S. soybean fields. Two elite public soybean varieties,

carrying either *rhg1-a* + *Rhg4* or *rhg1-b*, negatively impacted Pp populations measured at

30 dpi. This contribution of *Rhg1* was more rigorously demonstrated using near isogenic

lines for *rhg1-b*. Additionally, in three of the four soybean genetic backgrounds tested we

observed significantly more Pp at 30 dpi when Pp were co-inoculated with SCN, consistent

with a previous study. No significant changes in SCN cyst numbers upon co-inoculation

with Pp were observed for any of the SCN-resistant *rhg1* lines. No nematode-induced

changes in *Rhg1* transcript abundances were observed when sampling infected root

sections in any of the plant genotypes, and no changes in a jasmonate response indicator

were observed. However, the 3 dpi salicylate-mediated response to SCN or Pp became

more robust during SCN + Pp co-infection. Multiple hypotheses for further dissection and

manipulation of soybean resistance to *Pratylenchus* root lesion nematodes are suggested

based on the present findings.

Introduction

Root lesion nematodes (RLN[a]; *Pratylenchus* spp.) are important agricultural pests

(Jones et al. 2013; Rosa de Araújo et al. 2025). They are migratory ectoparasitic and

endoparasitic nematodes (Zunke 1990) with a wide host range including soybean.

Pratylenchus brachyurus is often considered to be the most damaging nematode pest of

soybeans in South America (Castanheira et al. 2020; Favoreto et al. 2019; Nomura et al.

2024). *Pratylenchus* populations survive cold winters in dead roots, including through

anhydrobiosis or slow desiccation (Townshend 1984). *Pratylenchus* are some of the most

common plant parasitic nematodes found in the soybean producing midwestern U.S.

states like Wisconsin, Minnesota, Iowa, and Illinois, with *Pratylenchus penetrans* (Pp)

being among the most common species, and a species for which soybean yield losses

have been documented (Chen et al. 2012; MacGuidwin and Bender 2012; MacGuidwin et

al. 2023; Saikai and MacGuidwin 2022; Schmitt and Barker 1981; Schumacher and Bird

2025). Unlike sedentary endoparasitic nematodes such as the soybean cyst nematode

(SCN, *Heterodera glycines*), Pp feed on soybean roots at multiple sites as ectoparasites or

endoparasites (Endo 1967; Saikai and MacGuidwin 2020). SCN spend limited time

exposed to environmental disturbances as they shelter inside host roots or in cyst-

encased eggs for much of their life cycle. In contrast, the migratory lifestyle of Pp (Zunke

1990) may make them more directly exposed to, and to some extent adapted to, changes

in soil pH and nutrients, soil moisture, soil type, own population density, microbial

communities, and competition from other nematodes. SCN is the most economically

important pathogen or pest on soybeans in the United States (Bradley et al. 2021). Across

all of the North Central U.S. and Ontario for ten recent years, SCN is estimated to have

caused approximately 4% soybean yield loss (about 1/4 of the total for all soybean

pathogens) (Allen et al. 2017; Bradley et al. 2021). Understandably, most efforts in U.S.

soybean systems have been directed at managing SCN and not *Pratylenchus* spp.

Estimates of the impact of RLN on soybean production are rare, but RLN and SCN

frequently co-occur (MacGuidwin et al. 2023; Saikai and MacGuidwin 2022; Schumacher

and Bird 2025). Northern U.S. soybean fields are typically rotated with maize, an excellent

host for RLN. In one deeply studied set of Wisconsin soybean fields, RLN was found to

cause up to 4.5% yield loss by itself (Saikai and MacGuidwin 2022), and about 3.79%

yield loss on corn (MacGuidwin and Bender 2016). In 2024 the farm gate values of corn

and soybean in Wisconsin alone were over USD $2.26 billion and USD $1.01 billion

respectively

(https://www.nass.usda.gov/Quick_Stats/Ag_Overview/stateOverview.php?state=WISCON

SIN), which illustrates the economic impact of even a 1% yield loss to RLN.

Along with crop rotation, the most important tool for managing SCN over the past

three decades has been host resistance provided by the major-effect quantitative trait

locus (QTL) *Rhg1* (Resistance to *Heterodera glycines* 1) (Bent 2022; Niblack et al. 2006).

There are two widely known resistance-conferring haplotype variants of *Rhg1* in soybean.

The haplotype *rhg1-a* is commonly derived from "Peking" while *rhg1-b* is often derived

from "PI 88788", with the latter being used in over 95% of all commercial nematode

resistant varieties (McCarville et al. 2017; Niblack et al. 2006). The Peking-type *rhg1-a*

haplotype is less effective if it is not coupled with resistance-associated alleles at the

unlinked *Rhg4* or *rhg2* loci (Basnet et al. 2022; Brucker et al. 2005; Yu et al. 2016). SCN

populations that at least partially overcome three reference sources of *rhg1-b* resistance

are designated as HG type 2.5.7. Throughout soybean-producing areas of the U.S., HG

type 2.5.7 and other *rhg1-b*-virulent SCN populations are increasingly common and are

gradually increasing their virulence on *rhg1-b* (Kleczewski et al. 2023; McCarville et al.

2017). In response, expanded use of Peking-type resistance and introduction of varieties

that carry PI 88788-type *rhg1-b* resistance stacked with additional minor-effect resistance

QTLs is anticipated (Bent 2022; Brzostowski and Diers 2017; Tylka et al. 2012)

The *rhg1-a* and *rhg1-b* haplotypes both carry three tightly linked genes that

contribute to SCN resistance (Cook et al. 2014; Cook et al. 2012a). *Glyma.18g022400*,

Glyma.18g022500 and *Glyma.18g022700* respectively encode AAT_{Rhg1} (putative amino

acid transporter), $\alpha\text{-}SNAP_{Rhg1}$ (vesicle trafficking mediator, also called GmSNAP18) and

$WI12_{Rhg1}$ (unknown function, carries a WI12 wound inducible protein domain). While the

causal *Rhg1* genes were identified a decade ago and their mechanism(s) of action are

being investigated, the resistance has been deployed for multiple decades. The $\alpha\text{-}$

$SNAP_{Rhg1}$ proteins differ between the *rhg1-a* and *rhg1-b* haplotypes, as does the copy

number of tandem repeats of the entire locus (Cook et al. 2014). The abundance of

resistance-associated $\alpha\text{-}SNAP_{Rhg1}$ protein is known to increase 10-15-fold in the

syncytium, the SCN feeding site (Bayless et al. 2016; Bayless et al. 2019). AAT_{Rhg1} protein

increases an order of magnitude along the SCN root penetration path, where a substantial

pool of AAT_{Rhg1} is associated with macrovesicles (Han et al. 2023). Less is known about

how the $WI12_{Rhg1}$ interacts with SCN, although impacts on root gibberellin levels have

been discovered (Dong and Hudson 2022). Silencing or knocking out any one of the three

genes increases susceptibility to SCN (Cook et al. 2012b; Dong and Hudson 2022; Liu et

al. 2017; Schumacher and Bird 2025). In transgenic potato, expression of the set of three

Deepak Haarith
Plant Disease

soybean *Rhg1* genes from their native soybean promoters or from strong constitutive

promoters elevated resistance to potato cyst nematodes (Butler et al. 2019).

We hypothesized that *Rhg1* could also provide some resistance to Pp and that the

commercially popular *rhg1-b* may have been beneficial against Pp. Melakeberhan and

Dey previously reported that the infection rate of SCN in some instances decreased with

increasing Pp proportions, and that the rate of Pp infection increased when co-inoculated

at a 1:1 ratio with *H. glycines* but did not increase at higher or lower co-inoculation ratios

(Melakeberhan and Dey 2003). That work was done using an SCN-susceptible soybean

cultivar. Schumacher and Bird recently reported some instances where the presence of

Pp increased SCN population densities on *rhg1-a* resistance sources, but with no overall

consistent trend (Schumacher and Bird 2025). Interactions between *Rhg1* and other

pathogens are only beginning to receive study, although most soybeans grown in recent

years across the U.S. carry this genetic resistance (Lopez-Nicora et al. 2023). On

soybean, Niblack and colleagues previously found that resistance to SCN was not reduced

during co-infections with the root knot nematode *M. incognita* (Niblack et al. 1986).

Significantly, the *rhg1-a* + *rhg2 (GmSNAP11)* combination has recently been shown to

enhance quantitative resistance against reniform nematodes (Usovsky et al. 2021;

Usovsky et al. 2022)

In this study we assessed (i) the success of Pp and an SCN HG 2.5.7 population on

the soybean lines IL3025N (*rhg1-b*), IL3849N (*rhg1-a/Rhg4*) and Williams 82 (SCN-

susceptible *rhg1-c*), in terms of total nematodes recovered per plant at the end of 30 days,

(ii) the impact on nematode numbers when SCN and Pp were co-inoculated at 2:1 or 1:1

SCN:Pp ratios, and, using a set of near isogenic lines (NILs) for *rhg1-b*, (iii) how *rhg1-b*

Deepak Haarith
Plant Disease

132 impacts Pp and SCN nematode numbers and the transcript abundance response of the

133 three *rhg1-b* genes as well as salicylic acid (SA) and jasmonic acid (JA) response indicator

134 genes.

Materials and Methods

Sand cup preparation

Quarried sand from Dane County, Wisconsin was sieved with a 0.25mm screen to

remove larger gravel or sand particles and the resulting fine sand was moistened, placed

in Nalgene polypropylene heat-tolerant autoclave bins at a maximum of 7.5 cm depth, and

heated in a 65 °C oven for 48 h. The sand was cooled to room temperature, filled in plastic

pots (9 cm deep, 6x6 cm at top tapering to 4x4 cm with drainage holes, filled with 190 +/-

10 cc of sand) and each of these sand cups (pots) was saturated with 50 mL of tap water.

Plant genotypes

Seeds of Williams 82 (Wm82; *rhg1-c*), IL3025N (*rhg1-b*) and IL3849N (*rhg1-a* and

Rhg4, susceptible allele at *Rhg2*) were used in the first set of cup assays. Wm82 is a

standard SCN-susceptible check; soybean lines that carry *rhg1-b* exhibit strong resistance

to HG 0 SCN but only quantitative/partial resistance to HG 2.5.7 SCN; lines with *rhg1-a* +

Rhg4 exhibit strong resistance to HG 0 SCN and HG 2.5.7 SCN (Bent 2022; Niblack et al.

2006). Seeds of Wm82 and near-isogenic lines Resistant NIL (Res. NIL; *rhg1-b*; LD09-

15604) and Susceptible NIL (Sus. NIL; *rhg1-c*; LD09-15630) were used in the second set

of cup assays and differential gene expression studies. IL3025N (synonyms: Illini 3025N,

LD11-2170) and IL3849N (synonyms: Illini 3849N, LD07-3395bf) were developed by Brian

Deepak Haarith
Plant Disease

Diers, University of Illinois at Urbana-Champaign, and are available through Baird Seed

Co. (Williamsfield, Illinois). The pedigree for IL3025N is Syngenta_03JR313108 x LD05-

3171. IL3849N is an SCN-resistant reselection from LD07-3395 whose pedigree is

Syngenta_WW115926 x LD00-2817. The NILs were developed by Brian Diers, University

of Illinois at Urbana-Champaign, through four backcrosses using LD00-3309 (from

Maverick x Dwight, carries *rhg1-b*) as the recurrent parent and IA3023 as donor of the

susceptible *rhg1-c* allele, hence the resulting NIL accessions are predicted to be >96%

LD00-3309 genotype. All seeds were surface sterilized using chlorine vapor as described

by Clough and Bent (1998).

Nematode inoculum

The Pp population was collected in Wisconsin and maintained on 'IO Chief' sweet

corn (*Zea mays*) root explants by inoculation onto 1-week old roots from freshly

germinated seed grown on 1X Gamborg's B-5 medium with 2% sucrose and 1.5% agar

(Rebois and Huettel, 1986). Pp subcultures were started 3 months prior to use in

experiments. The "SCN HG 2.5.7 IL" field population, obtained courtesy of Alison

Colgrove, was maintained on IL3025N soybean plants and exhibited an FI ~70 on that

rhg1-b soybean host relative to reproduction on Williams 82.

SCN eggs from crushed cysts were hatched in 4 mM $ZnCl_2$ solution at 27°C and

dark conditions. Freshly hatched J2 stage SCN were rinsed and re-suspended in de-

ionised water immediately prior to inoculation. Pp (all life stages other than eggs/J1) were

collected together from corn root explants using the Baermann funnel method (Baermann

1917). The Pp were then rinsed and resuspended in de-ionised water. Pp inoculum

batches had inherent variation but did consistently exhibit a demographic representation

Deepak Haarith
Plant Disease

of, very approximately, 35% adult females, 15% adult males, 40% third and fourth stage

juveniles and 10% second stage juveniles. Within each experiment the Pp aliquots applied

to each of the Pp-inoculated plants were highly similar because all aliquots were drawn

from the same Pp nematode preparation. Nematode inoculum density was adjusted so

that 2 ml of liquid (1 ml containing Pp or water + 1 ml containing SCN or water) was

inoculated into the sand of each pot via a small hole made using a 200 uL pipette tip,

parallel to and 1 cm away from the soybean seedling, which was then re-covered with

sand. The plants remained covered with humidity domes for one day post inoculation.

Nematode count cup assays

Unless otherwise noted, three nematode treatments were included in each

experiment: SCN alone, Pp alone, or SCN + Pp. For the experiments presented in Figure

1, four independent experiments were conducted, two using a 1:1 ratio of SCN:Pp (500 of

each) and two using a 2:1 SCN:Pp ratio (approximately 1500 SCN:750 Pp) for co-

inoculations. Other pots in these experiments received only one nematode species (500 of

either SCN or Pp in the 1:1 experiments; 1500 SCN or 750 Pp in the 2:1 experiments)The

other pots within the 1:1 and 2:1 experiments received the noted number of only one or

the other nematode species. Each pot contained ~190 cc of sand (described above). The

2:1 experiments were conducted one week apart using separate SCN hatches but same

batch of Ppused Pp from the same Baermann funnel extraction. The two 1:1 experiments

each used independent preparations of SCN and Pp and were conducted {X} months

apart from the eachother and separate from the 2:1 experiments. Within each experiment

there were five replicates of each treatment (3 plant genotypes x 3 nematode treatments =

9 treatments per block), arranged in a randomized complete block design. The three plant

Deepak Haarith
Plant Disease

genotypes (SCN-susceptible Wm82, SCN-resistant IL3029N and partially SCN-resistant

IL3025N) are described above. Each block of 9 pots was in a separate tray with a different

randomized arrangement, with trays adjacent to each other in a large controlled

environment growth room, and harvest\processing was done by block. Seeds were

germinated in paper towels at 27°C for 4-5 days in the dark until a 5 cm radicle had formed

and were then transplanted with one plant per sand pot. Nematode inoculum (see

previous section) was applied one day after transplanting these VE stage seedlings

(Vegetative Emergence) (Pedersen 2009). Plants were covered with a humidity dome for

the first 2 days after transplanting. Each experiment was done twice (two separate dates).

Inoculated plants were maintained in a growth room with 14h daylight, day temperature

27°C, night temperature 22°C, for 30 days. The plants were watered minimally (liquid did

not flow from holes into the tray) by applying 10 mL of tap water to each pot each day for

the first week, 15 mL a day the second week, alternating between 15 mL and 30 mL a day

the third week and 30 mL a day thereafter. The pots were fertilized once after 15 days and

again 7 days later using MiracleGro (Scotts, Marysville OH) as per manufacturer

recommendations instead of tap water.

For the experiments presented in Figures 2 and 3, a similar experimental design was

used (3 plant genotypes x 3 nematode treatments = 9 treatments per block, arranged in a

randomized complete block design; (two independent experiments with five replicates

within each experiment, using distinct batches of seedlings and fresh nematode inoculum

in each experiment),). The three plant genotypes in this case were standard SCN-

susceptible check Wm82, SCN-susceptible Sus. NIL and partially SCN-resistant (*rhg1-b*)

Deepak Haarith
Plant Disease

Res. NIL, and treated with approximately 500 SCN, 500 Pp or 500 SCN + 500 Pp.

nematodes of the noted type (Pp and/or SCN) were used.

After 30 days, nematodes were extracted from the sand using water flotation and

decanted into a nested sieve system consisting of a 2800 μm, 710 μm, 177 μm and 38 μm

sieve (ASTM E11 US Alternative sieves 7, 25, 80, and 400 respectively). Plant roots and

other material retained on the 2800 μm sieve after vigorous rinsing with tap water were

collected, then the roots of all samples were chopped into ~1 cm pieces and incubated on

Baermann funnels with de-ionised water for 48 h at room temperature (25°C). 50 mL of

water was eluted from the bottom of the funnels and evaluated for Pp (all life stages other

than eggs/J1) and SCN J2s and males. Contents retained by the 177 μm sieve were

enumeratedretained for SCN cyst enumerations under a stereoscope while contents from

the 38 μm sieve were enumerated for Pp of all life stages (juveniles and adults) and for

SCN males and juveniles. Nematode samples were further clarified by dilution using 70%

(70 g into 100ml H_2O) sucrose solution prior to microscopic enumeration. In pilot tests with

no plants, average Pp extraction efficiency 24 h after inoculation was 54% (std.dev. 1.1%)

using the above protocol. In NIL samples co-inoculated with both nematodes, contents

from 38 μm sieve were also evaluated for SCN J2 and male numbers.

NIL *rhg1* differential mRNA abundance assay

Each mRNA abundance result reports data from two independent experiments that used

distinct batches of seedlings and nematode inoculum. Within each experiment that used

roots from VE seedlings, three replicate blocks were set up in a randomized complete

block design as described above for nematode count cup assays except with two pots of

each treatment within each block, to allow sampling at the 3 dpi (days post inoculation)and

7 dpi time points. Note that infection can be asynchronous and these time points can

include infections initiated 0-3 days or 0-7 days prior to sampling. For gene expression

experiments that used V1 (Vegetative 1st stage) plants (Pedersen, 2009), inoculation

occurred when the first set of trifoliate leaves had emerged and two blocks were utilized.

For RNA extraction, roots from plants (Wm82, Res. NIL and Sus. NIL) inoculated with

either SCN (500 J2) or Pp (500 J2-to-adult worms) or both (500 SCN + 500 Pp) as well as

non-inoculated controls were rinsed in water and observed under a stereoscope at 3 dpi or

7 dpi. Each sample was from a separate plant. For each sample, about 100 mg of root

tissues from a single plant, from ~1 cm regions segments with multiple visible

lesionswater-soaked foci (putative infection sites), all from a single plant, was were placed

in ice-cold cryovials with 5 ceramic beads and immediately frozen in liquid nitrogen. These

samples were stored at -80°C. Infection sites were in the zone of maturation on lateral

roots and similar root sites were collected from non-inoculated control root systems.

Frozen tissue was processed at 2500 rpm for 30 s in a powerlyzer and returned to liquid

nitrogen for a minute. This process was repeated 2-3 times until tissue was pulverized.

RNA was then extracted from pulverized root tissue using Zymo (Irvine, CA) Direct-Zol

TriZol-based miniprep kit using manufacturer protocol. cDNA was synthesized from 1µg of

RNA for each sample using Solis BioDyne (Tartu, Estonia) FireScriptTM RT cDNA

Synthesis mix with included Olido dT random degenerate hexamer primers (Catalog #06-

20-00500), according to manufacturer protocol.

Real-time quantification and Ct values were obtained for transcripts encoding

AAT_{Rhg1}, $WI12_{Rhg1}$, α-SNAP$_{Rhg1}$ (detects all haplotypes), α-SNAP$_{Rhg1}$HC (detects only the

HC/ *rhg1-b* haplotype), as well as for *AAT* (*Glyma.10G109500*; JA response indicator),

NIMIN1 (*Glyma.10G010100*; SA response indicator) and the internal standard gene

NREG1 (*Glyma.11G079600*) (Beyer et al. 2021). ΔCt values were the average of two

technical replicates from the same sample. To determine the fold-change in transcript

abundance in response to nematode inoculations the $2^{-\Delta\Delta Ct}$ method was used (Livak

and Schmittgen 2001). For comparisons within plant genotypes, ΔCt values obtained using

NREG1 as internal standard were normalized to the average ΔCt values of all no-

nematode samples for the same plant genotype within the same experiment date to obtain

ΔΔCt values. ~~Alternatively, to evaluate relative transcript abundances across~~For

comparisons between plant genotypes, the ΔCt values for all genotypes and treatments

were normalized to the average ΔCt values of no-nematode inoculated Sus. NIL plants to

obtain ΔΔCt values. A full list of the oligonucleotide primers used in this study is available

in Supplementary Table 1.

Statistical analyses

All data were analyzed and visualized using R (version 4.3.0). Graphs show mean

and standard error of mean for the original non-transformed data. For statistical tests,

nematode count data were square root transformed or log transformed to achieve

normality (residuals checked graphically to confirm equal variance and normality). ANOVA

was performed using the aov() function and the means were separated using TukeyHSD()

in the package stats(). Plant genotype, nematode treatment, experiment, and blocks within

experiment were all fixed effects and the interaction between genotype and nematode

treatment was included in the model. The specific R model used is provided for each

analysis in the Supplemental Tables. A typical analysis assessed one nematode type

(e.g., Pp numbers in Pp-only and Pp+SCN treatments) and had 60 entries (2 experiments,

Deepak Haarith
Plant Disease

291   5 replicate blocks per experiment, 3 plant genotypes, 2 treatments). For the transcript

292   abundance evaluations, experiments were set up similarly to the above and the same

293   model was fit as for nematode counts (models stated in Supplementary Tables). Transcript

294   statistical analyses were performed on the ∆∆Ct values (mRNA abundance).

Results

Pp parasitism of soybean is enhanced by co-infection with SCN

Controlled environment single-species inoculation and co-inoculation experiments

with Pp and SCN were carried out (Pp-only, SCN-only or Pp+SCN all within the same

experiment). Experiments were first conducted using soybean genotypes that differ at

hundreds of loci, including different SCN resistance haplotypes. Wm82 (*rhg1-c*) is SCN-

susceptible while IL3025N (also known as LD11-2170) carries *rhg1-b* and IL3849N (also

known as LD07-3395bf) carries *rhg1-a* and *Rhg4*. Four independent experiments were

conducted, two using a 1:1 ratio and two using a 2:1 SCN:Pp ratio for co-inoculations, with

five replicates of each treatment within each experiment. The experiments utilized an SCN

HG 2.5.7 population that, as expected, reproduced well on the susceptible *rhg1-c* (Wm82)

and substantially overcame *rhg1-b* resistance (IL3025N) while exhibiting poor reproduction

on a plant line carrying the *rhg1-a* and *Rhg4* combination (IL3849N) (green bars in Figure

1C and 1D; all differences between SCN-only inoculations (green bars) were significant at

p<0.001). ANOVA tables associated with this and subsequent figures are provided as

Supplementary Tables.

Deepak Haarith
Plant Disease

Figure 1A and 1B show that more Pp were recovered upon co-inoculation with

virulent SCN HG 2.5.7 IL compared to the Pp-only inoculations, in experiments on SCN-

susceptible Wm82 or moderately SCN-susceptible IL3025N soybean plants (orange vs.

blue bars for same soybean genotype, p < 0.001). More Pp were also recovered upon co-

inoculation with SCN on IL3849N (Figure 1A and 1B), which is a genotype that largely

prevents SCN cyst production (Figure 1C and 1D). The above observations were true for

SCN:Pp inoculation ratios of 2:1 and 1:1. In all of the experiments in Figure 1, about twice

as many Pp were recovered 30 days after inoculation when Pp were co-inoculated with

SCN HG 2.5.7, compared to Pp-only inoculations (the ratio on the same plant genotype in

the same experiment ranged from 1.7 to 2.3, across all plant genotypes and both ratios of

co-inoculation).

Within the same experiments as above, there were no significant differences in SCN

HG 2.5.7 IL cyst numbers between SCN-only inoculations and co-inoculations with Pp on

the same plant genotype (green vs. blue bars, Figure 1C and 1D). Nearly 10-fold more

cysts were recovered across treatments when ~1500 (Figure 1D) rather than 500 (Figure

1C) SCN J2s were inoculated, for both solo inoculations and co-inoculations with Pp.

rhg1-b impedes the parasitism of Pp

330   A separate phenomenon was observed in the experiments of Figure 1, suggesting

331   that soybean SCN resistance might also impact RLN. When inoculated by itself or together

332   with SCN, significantly fewer Pp were recovered after 30 days from the SCN-resistant

333   variety IL3849N than from Wm82 (compare bars of same color in Figure 1A and 1B). Pp

334   abundance on IL3025N was lower than on Wm82 in three of the four comparisons (Figure

Deepak Haarith
Plant Disease

1A and 1B). Fewer Pp were recovered from IL3849N than from IL3025N only for the 2:1

SCN:Pp inoculations (Figure 1B vs. 1A). However, the genetic backgrounds of these

soybean genotypes differ at thousands of loci.

A more rigorous test for an effect of the SCN resistance *rhg1-b* haplotype on Pp

was conducted by using near-isogenic soybean lines (NIL) expected to carry the same

genotype at >96% of all loci but carrying either *rhg1-c* (Sus. NIL) or *rhg1-b* (Res. NIL) at

the *Rhg1* locus. The results supported the conclusion that *rhg1-b* can negatively impact Pp

(Figure 2). Significantly fewer Pp were recovered from the Res. NIL compared to the Sus.

NIL when singly inoculated with Pp (Figure 2A blue bars). When co-inoculated with both

nematodes (Figure 2A orange bars), fewer Pp were again recovered from Res. NIL plants

than Sus. NIL plants but with a difference that was at the margin of statistical significance

(p = 0.056).

For SCN, multiple studies have shown that the proportion of female to male

nematodes formed is often lower on soybean plants with effective SCN resistance than on

susceptible plants (Colgrove and Niblack 2005). Intriguingly, with migratory Pp we also

observed that the ratios of females to males recovered from the *rhg1-b* Res. NIL were

significantly lower than those for *rhg1-c* Sus. NIL plants (Figure 2B). This was true in both

Pp-only and SCN co-inoculated treatments. When the proportion of Pp found in roots vs.

the surrounding sand at 30 dpi was enumerated, the *rhg1-b* Res. NIL had a higher

proportion of the Pp in the sand than the Sus. NIL, but intriguingly, that was only in the Pp

+ SCN co-inoculated samples (Figure 2C). For the Res. NIL line, a ~5-fold higher

proportion of Pp were in the sand than in the root upon co-inoculation with SCN, compared

to the Pp-only samples (Figure 2C). The relative presence of juvenile Pp 30 days after

Deepak Haarith
Plant Disease

inoculation was examined as another parasitism/resistance metric, because those

juveniles are predominantly progeny produced from the initial inoculum. At 30 days after

inoculation the number of juvenile-stage Pp (Figure 2D) and the proportion of juvenile-

stage vs. adult Pp (Figure 2E) again showed that presence of SCN shifted Pp population

demographics. Juvenile Pp were significantly less common on *rhg1-b* Res. NIL plants

compared to the *rhg1-c* Sus. NIL. For the juvenile:adult ratio they were less common on

Res. NIL plants only when Pp and SCN were co-inoculated (Figure 2E). The findings of

Figure 2 together indicate that *rhg1-b* can exert a negative impact on Pp, in the presence

of SCN but also in the absence of SCN.

Pp co-inoculation reduces SCN cyst numbers only in the susceptible *rhg1-c* lines

As expected, in SCN-only inoculations fewer SCN HG 2.5.7 IL cysts were

recovered from the resistant *rhg1-b* NIL plants compared to the susceptible *rhg1-c* NIL

plants or Wm82 (Figure 3A; data are from same experiments as Figure 2). Co-inoculation

of Pp with SCN reduced the number of cysts recovered on the SCN-susceptible *rhg1-c*

Wm82 and Sus. NIL lines (Figure 3A). The difference was statistically significant only for

Wm82 (Figure 3A), but a similar trend was also observed on Wm82 in the experiments of

Figure 1A (p = 0.094) and on Sus. NIL in the experiments of Figure 3A (p = 0.070). When

the plants had partial resistance to SCN that reduced cyst production (Res. NIL), co-

inoculation with Pp did not further reduce cyst numbers (Figure 3A). In the same

experiments, the J2 and male SCN populations in roots and sand at 30 dpi were also

enumerated. The results were not consistent across the two SCN-susceptible plant

genotypes, as fewer total SCN J2s were recovered after co-inoculation with Pp only in the

case of the Sus. NIL plants (Figure 3B). Co-inoculation with Pp did not reduce J2 SCN

numbers for Wm82 and Res. NIL plants (Figure 3B). The number of SCN males recovered

exhibited substantial variation, and no statistically significant differences were observed

between plant genotypes or between plants inoculated with only SCN and those

inoculated with both nematodes (Figure 3C).

387 No significant changes in *rhg1-b* transcript abundance detected in response to

388 nematode infection

In light of the observed interactions between SCN, Pp and *rhg1-b*, we investigated

nematode induction of increases in transcript abundance of the three *rhg1-b* genes as well

as indicator genes of SA-response and JA-response activity in soybeans, at 3dpi and 7dpi

(Figures 4 and 5). As has previously been observed (Cook et al. 2014), *Rhg1* transcripts

were approximately 5- to 20-fold more abundant for the high-copy *rhg1-b* genotype than

for the single-copy *rhg1-c* genotype (Supplemental Figure 1). Transcript abundances are

reported in Figures 4 and 5 as fold-change relative to the mean of the abundance for the

non-inoculated control of the same plant genotype within the same experiment (ΔΔCt).

Figure 4 A-F show that even with two independent experiments that each sampled

infected zones from three independent root systems, we did not detect any significant

nematode-induced transcript abundance of any of the *rhg1-b* genes when plants were

inoculated at VE stage (emerging seedlings, cotyledons not yet separated). We also

observed no significant differences with plants inoculated at the V1 stage (fully expanded

trifoliate; two independent roots in each of two experiments; Supplemental Figure 2).

Combined infection by SCN and Pp induces a stronger salicylic acid response

At either 3 dpi or 7 dpi, neither SCN, Pp nor their combination induced a significant

change in transcript abundance ($\Delta\Delta$Ct) for the JA-response indicator gene '*AAT*' (Figure

5A and 5B). Note that this established soybean JA-response indicator gene *AAT* (Beyer

et al. 2021) is *Glyma.10G109500* and is not the *Rhg1* gene for AAT$_{Rhg1}$. In contrast to the

JA result, transcript abundances for the soybean SA-response indicator gene *NIMIN1*

(*Glyma.10G010100*) did exhibit statistically significant increases at 3 dpi in many of the

nematode-inoculated samples (Figure 5C). Only dual inoculation with SCN and Pp

consistently caused significant elevation of the SA indicator in all three soybean genotypes

(Figure 5C). The elevated SA response indicated by *NIMIN1* transcript levels was no

longer observable at 7 dpi (Figure 5D).

Discussion

Although SCN is the most economically important nematode of soybeans in the US,

in more tropical soybean producing states such as Brazil, RLN (especially *Pratylenchus*

brachyurus) has been reported to be more consequential (Castanheira et al. 2020;

Nomura et al. 2024). No soybean varieties labeled as having strong resistance to RLN

have been deployed but searches are underway (e.g., Chowdhury et al. 2022; Hamawaki

et al. 2019; Machado and Araúujo 2016). SCN and RLN frequently co-occur. Given the

importance of the soybean crop, it is crucial to understand if there are interactions

between the two nematodes on soybean and if that could lead to a change in their relative

abundance over time. Only limited surveys and studies are available on the tripartite

Deepak Haarith
Plant Disease

interaction of Soybean-SCN-RLN (Melakeberhan and Dey 2003). At the time of writing this

manuscript, no published reports are available on how the deployed *Rhg1* resistance

against SCN factors into this tripartite interaction.

Many soybean fields are infested with both SCN and RLN and the limited research

reported suggests that their interaction is complicated. MacGuidwin et al. suggested a

competitive relationship (population densities of both SCN and RLN suppressed by co-

occurrence) based on an analysis of field survey data (MacGuidwin et al. 2023).

Melakeberhan and Dey tested four 1000-nematode inocula with varying proportions of

SCN and Pp and found evidence of competition when SCN outnumbered Pp 3:1. When

both species were equally represented the interaction appeared to be antagonistic, as

numbers of Pp increased and SCN numbers decreased as compared to the single-species

case (Melakeberhan and Dey 2003). Those experiments used HG 0 SCN on an SCN-

susceptible soybean host. Taking a cue from that prior study (Melakeberhan and Dey

2003), we co-inoculated equal numbers of SCN and Pp, and then also examined how

twice as many SCN as Pp altered the result. As in the study of Melakeberhan and Dey

wWe also observed that presence of SCN increased Pp numbers on soybean roots, but

also observed the increase on an SCN-resistant host (IL3849N). In contrast to the prior

study (Melakeberhan and Dey 2003), we observed more not fewer Pp 30 days after co-

inoculation with twice as many SCN. Hypotheses for how SCN assists Pp parasitism

include effector-mediated suppression host defenses, and/or simply by providing

established entry points into the root.

We used HG 2.5.7 SCN both because it is informative with respect to *Rhg1* (it largely

overcomes *rhg1-b* but not *rhg1-a + Rhg4*), and because it is becoming the most common

Deepak Haarith
Plant Disease

type of SCN found in North Central U.S. soybean fields. The capacity of SCN to increase

Pp numbers on IL3849N is a related topic for future study - Pp may for example still be

quantitatively impacted by the reduced number of SCN present on the resistant host after

the first few days of infection, and/or by the plant defense responses induced by SCN on

the resistant host.

The Pp life cycle can require longer than 30 days. However, we were testing Pp

together with SCN and hence performed 30-day experiments. 30 days is the established

assay time for SCN resistance assays because longer time points introduce the issue of

multiple generations of SCN reproduction. A priori, impacts on any plant-parasitic

nematode can be documented without taking the study through a full nematode life cycle.

However, future studies could incorporate a full life cycle assessment of Pp during co-

infection with SCN.

The overall infection efficiency of Pp on soybean as opposed to other hosts also

merits future study. Pp did infect soybean in the present study setting, and infected SCN-

susceptible soybean lines more when there were more SCN present (Figure 1). At the 30-

day time point it was often the case (see Figure 2C) that in the absence of SCN over half

of the Pp were in the soybean roots and not in the sand media. We also were able to

readily find water-soaked infection sites on the roots for our RNA extraction and qPCR

experiments. Some attrition (nematodes that do not find and feed on the root and hence

do not survive) is common in any nematode inoculation assay and attrition will increase as

inoculum size per root increases. However, out of an initial inoculum of 500, the number

of Pp recovered from roots and sand at 30 dpi was often below 10%. We speculate that,

despite the care taken to water pots consistently and with small volumes, sub-optimal

Deepak Haarith
Plant Disease

watering (that washes nematodes from the root zone and/or imposes low-moisture stress

on the nematodes) may have been responsible for this low Pp recovery rate.

In our previous studies with *Rhg1* genes we had shown their capacity to increase

resistance to other cyst nematode species and genera (*Heterodera schactii*, *Globodera*

pallida, *Globodera rostochiensis*) when transgenically expressed in *Arabidopsis* or potato

(Butler et al. 2019). Similarly, the combined presence of SCN resistance loci *rhg1-a* and

rhg2 was shown to also provide quantitative resistance against reniform nematodes

(*Rotylenchulus reniformis*) (Usovsky et al. 2021; Usovsky et al. 2022). The present study

was designed as an initial investigation of the broader hypothesis that SCN-resistant

soybeans deployed in fields may have been partially controlling RLN. Our data are not

inconsistent with the hypothesis that early (30 dpi) establishment of Pp can be impeded by

rhg1-a + Rhg4, as evidenced by our observed reduction of Pp population sizes in IL3849N

relative to Wm82. Early Pp establishment was also impeded on IL3025N (*rhg1-b*) when

initial SCN and Pp population densities were similar, but was not significant when SCN

outnumbered Pp. We further hypothesized that the interaction between SCN and Pp

could reduce the efficacy of *Rhg1*. In support of that, as noted above, relative to

inoculation of the same plants with just Pp we found an increase in Pp recovery when

simultaneous SCN pressure was applied ~~on plants that are susceptible to the SCN~~

~~population (Pp + HG 2.5.7 SCN on Wm82 and IL3025N)~~. Interestingly, for the virulent SCN

HG 2.5.7 IL population that we tested, counts of SCN cysts were not reduced by co-

inoculation with Pp except at 1:1 ratio when *rhg1-a* or *rhg1-b* were absent. A similar

dynamic between SCN and Pp was observed by Melakeberhan and Dey on multiple

soybean cultivars that lacked the *rhg1-a* or *rhg1-b* resistance (Melakeberhan and Dey

2003).   We have not investigated full-season resistance or yield protection (see also

Pedro et al. 2024), but the reductions in parasitism observed in the present study and that

of Melakeberhan and Dey suggest a potential contribution to full-season disease

resistance against Pp. The field studies of Saikai and MacGuidwin (Saikai and

MacGuidwin 2022) were done using varieties with *rhg1-b* resistance, so a yield loss impact

of Pp on SCN-resistant soybean varieties has been demonstrated. The present study

suggests that Pp damage could be even worse if use of *rhg1-b* soybeans was not

prevalent, but further field-level and full-season or multi-season studies are needed to

investigate this point.

As noted above, in our experiments and those of Melakeberhan and Dey (2003) co-

inoculation with SCN helped Pp or was neutral, in all cases tested. Even on resistant

hosts, we observed no reduction of Pp at 30 dpi when co-inoculated with SCN.   In

contrast, with wheat, Laserre et al. 1994 and Mokrini et al. 2018 both observed that on

susceptible hosts co-inoculation with cereal cyst nematode tended to reduce *Pratylenchus*

population sizes (Lasserre et al. 1994; Mokrini et al. 2018).   Intriguingly Laserre and

colleagues also showed, on a susceptible host, a reduction in eggs and vermiform *P.*

neglectus upon inoculation of *H. avenae* onto a separate part of the root system. However,

on wheat accessions with resistance to *H. avenae* they observed that co-inoculation with

H. avenae led in some cases to higher numbers of *Pratylenchus*, similar to the trend in our

experiments (Lasserre et al. 1994).

The *rhg1-b* haplotype SCN resistance, unlike the *rhg1-a* haplotype resistance, is not

co-dependent on the *Rhg4* locus on chromosome 08 (Patil et al. 2019). We were able to

obtain near-isogenic lines for *rhg1-b* that, after four backcrosses, should be genetically

Deepak Haarith
Plant Disease

>96% identical, for a more well-controlled test of the contribution of *rhg1-b* to resistance

against Pp. As with IL3025N and IL3849N, we observed fewer Pp on Res. NIL plants

compared to the Sus. NIL plants when only Pp was inoculated, strongly linking *rhg1-b*

SCN resistance to partial Pp resistance as well. In addition to the overall reduction in Pp

numbers in the Res. NIL, we also observed reductions in the ratios of females to males

and juveniles to adults. RLN males and females differ in how they feed on the roots and

how much damage they can subsequently cause (Saikai and MacGuidwin 2020). Females

make larger lesions and burrow more frequently into the roots than the males. Hence

having overall fewer Pp and a population that is skewed towards adults, especially males,

may be less damaging to the crop. When considered together, these observations indicate

that *rhg1-b* imposes stress on Pp populations and further support the indication that *rhg1-b*

may be reducing the severity of damage to soybean caused by Pp.

In an earlier pair of studies, we had observed AAT_{Rhg1} protein abundance increasing

along SCN penetration path (Han et al. 2023), unlike the α-$SNAP_{Rhg1}$ that exhibited a

strong abundance increase in the syncytium (Bayless et al. 2016). Because Pp do not

form syncytia, in the present study we hypothesized that *Rhg1-GmAAT* could be one of

the early response genes that functions against diverse nematodes. However, for the

genes encoding AAT_{Rhg1}, α-$SNAP_{Rhg1}$ or $WI12_{Rhg1}$ we did not observe any statistically

significant transcript induction at 3 dpi for plants inoculated with Pp at the VE or the V1

stage, or at 7 dpi after VE stage inoculation. It is possible that any transcript induction is

sufficiently local to infected cells that the vast majority of non-infected cells in the root

samples diluted out any signal from cells at or immediately adjacent to the site of

nematode infection. It is also possible that expression of the *Rhg1* gene products is

Deepak Haarith
Plant Disease

regulated in part at the post-transcriptional level. This absence of an observed transcript

induction is consistent with a previous study where we saw no significant differences in

transcript induction for α -SNAP$_{Rhg1}$HC at 7 dpi but did observe significant changes in

protein abundance (Haarith et al. 2025).

Transcript abundances for *NIMIN1*, a preferred soybean salicylic acid response

pathway indicator (Beyer et al. 2021), were upregulated significantly at 3 dpi in response

to SCN, and were more reliably upregulated when Pp co-infected alongside SCN. Perhaps

of greater interest, Pp alone did not reproducibly elicit this SA response while Pp+SCN did

(Figure 5), and this Pp co-infection with SCN enhanced both SA responses and Pp

population sizes (Figure 1). Antagonism between SA- and JA-mediated defense

responses has been widely reported (Glazebrook et al. 2003; Verma et al. 2025). There

are multiple instances of a "host mis-direction" virulence strategy in which pathogens or

pests that are more effectively defended against by the one set of defense responses

cause the plant to instead activate the other defense responses (Schweiger et al. 2014).

The non-sedentary (migratory) parasitism exhibited by Pp may be more similar to plant

attack by herbivorous insects than it is to a haustorium-forming biotrophic fungus or

oomycete, or a syncytium-forming cyst nematode. Our results suggest that Pp may lose

more to JA-mediated plant defenses than to SA-mediated plant defenses; future studies

could investigate if Pp in fact benefits from stronger elicitation of SA-mediated defenses.

The findings of (Usovsky et al. 2021) correlating α-SNAP-encoding genetic loci with

resistance to both SCN and reniform nematode support speculation that α-SNAP protein

variants may also contribute to resistance against Pp. A large body of evidence has

implicated SNAP proteins in SCN resistance, including major roles for the α-

SNAP$_{Rhg1}$HC/GmSNAP18 protein encoded by *rhg1-b* and the α-SNAP$_{Rhg1}$LC/GmSNAP18

protein encoded by *rhg1-a*, and more subtle contributions from a γ-SNAP (gamma-SNAP)

encoded at *cqSCN-006* (*Glyma.15G191200*) and loss of the α-SNAPs encoded at

GmSNAP02 (*Glyma.02G260400*), and *rhg2/GmSNAP11* (*Glyma.11g234500*) (Bayless et

al. 2016; Bent 2022; Butler et al. 2021; Cook et al. 2014; Cook et al. 2012a; Haarith et al.

2025; Lakhssassi et al. 2017; Liu et al. 2017; Matsye et al. 2012; Usovsky et al. 2023).

Four separate α-SNAP proteins bind to a single SNARE bundle as they mediate NSF

binding and SNARE bundle disassembly (Zhao et al. 2015), and presence of multiple

types of SNAP proteins is likely to impact their function. Future research could reveal that

certain combinations of SCN resistance-associated α-SNAP-encoding loci may provide

stronger Pp resistance than *rhg1-b*, and/or provide diversified resistance mediators that

impose distinct selection pressures on Pp populations. However, a leading hypothesis for

the impact on cyst nematodes of the non-canonical SCN resistance-associated α-SNAPs

is their cytotoxicity and elevated abundance in syncytia (Bayless et al. 2016; Bayless et al.

2018), and Pp are not sedentary feeders that induce syncytia. Some Pp individuals do

feed on a single host cell for several hours (Zunke 1990), which may be enough time for

SCN resistance-associated SNAPs to perturb the susceptible status of the host cell. But

the *Rhg1*-encoded AAT$_{Rhg1}$ and WI12$_{Rhg1}$ proteins also contribute to SCN resistance (Cook

et al. 2012). An equally strong hypothesis for future research is that AAT$_{Rhg1}$ and/or

WI12$_{Rhg1}$, moreso than α-SNAP$_{Rhg1}$, are the primary mediators of the impeded Pp

parasitism caused by *Rhg1*.

Taken together, the present results indicate that the *rhg1-b* haplotype impedes the

parasitism of Pp as well as SCN – a pair of plant parasitic nematode species that are

Deepak Haarith
Plant Disease

highly distinct taxonomically and behaviorally. The present controlled environment results

also show, in agreement with a past study, that on SCN-susceptible soybean genotypes

SCN can enhance Pp population sizes. There is measurable interaction between the two

nematodes regarding the efficacy of *rhg1-b* resistance, and in their induction of salicylic

acid responses within the plant. The findings suggest multiple avenues for future study

and manipulation of soybean resistance to *Pratylenchus* root lesion nematodes.

Author Contributions

598 DH conceptualized, designed, and performed the experiments, analyzed the data, and

599 authored the manuscript. AEM provided Pp inoculum, methods and expertise in

600 *Pratylenchus* biology and edited the manuscript. AFB co-designed the experiments, and

601 analyzed and co-authored the manuscript. AFB and DH are co-PIs on the grant that

602 funded this study.

Acknowledgements

Funding for this study was provided primarily by a grant to AFB and DH from the

Wisconsin Soybean Marketing Board, and also by grants to AFB from the United Soybean

Board. We thank Brian Diers of University of Illinois at Urbana-Champaign for providing

characterized soybean accessions, undergraduate John Stuntebeck for providing

experiment support, and Nicholas Keuler of the UW-Madison Statistical Consulting Group

for advice on statistical analyses.

Deepak Haarith
Plant Disease

Figure Captions

Figure 1. Nematode counts on Wm82, IL3849N (*rhg1*-a) and IL3025N (*rhg1*-b) plants at

30 dpi with either or both SCN HG 2.5.7 IL and Pp. (A) Pp counts at 30 dpi at 1:1 SCN: Pp

inoculation (B) Pp counts at 30 dpi at 2:1 SCN:Pp inoculation. Blue bars correspond to Pp-

only inoculation and orange bars represent combined inoculation of both SCN and Pp. (C)

Numbers of SCN cysts recovered at 30 dpi at 1:1 SCN:Pp inoculation. (D) Numbers of

SCN cysts recovered at 30 dpi at 2:1 SCN:Pp inoculation. Green bars represent SCN-only

inoculation and the light-blue bars represent combined inoculation of both SCN and Pp.

Values shown are mean +/- standard error of mean for raw (non-transformed) nematode

counts. For the comparisons of greatest interest, statistical significance for differences

between transformed means are displayed as p-values from ANOVA Tukey HSD tests

(see Methods). The SCN HG 2.5.7 IL used partially overcomes *rhg1-b* resistance and

represents the most prevalent HG type in the USA. Data are for two independent

experiments with five replicates within each experiment (n = 10 for each bar).

Figure 2. Pp counts on Wm82, Sus. NIL and Res. NIL plants at 30 dpi in either Pp-only

inoculations (blue bars) or a combination of SCN and Pp at 1:1 ratio (orange bars). (A)

Counts of all Pp. (B) Ratio of Pp females to males. (C) Ratio of Pp in sand fraction to root

fraction. (D) Ratio of Pp juveniles (J1 through J4) to adults (male + female). (E) Counts of

Pp juveniles (J1 through J4). Values shown are mean +/- standard error of mean for non-

transformed data. Instances of possible significant differences of means are marked with

p-values from ANOVA Tukey HSD tests performed on transformed data (see Methods). p

Deepak Haarith
Plant Disease

> 0.2 if no p value is shown. The SCN HG 2.5.7 IL used partially overcomes *rhg1-b*

resistance and represents the most prevalent HG type in the USA. Data are for two

independent experiments with five replicates within each experiment (n = 10 for each bar).

Figure 3. SCN cyst counts on Wm82, Sus. NIL and Res. NIL plants at 30 dpi in either

SCN-only inoculation (green bars) or a combination of SCN and Pp at 1:1 ratio (light-blue

bars). (A) SCN cyst counts. (B) SCN J2 counts (sand + root). (C) SCN male counts (sand

+ root). Values are mean +/- standard error of mean for non-transformed data. Instances

of possible significant differences of means are marked with p-values from ANOVA Tukey

HSD tests performed on transformed data (see Methods). p > 0.2 if no p value is shown.

The SCN HG 2.5.7 IL used partially overcomes *rhg1-b* resistance and represents the most

prevalent HG type in the USA. Data are for two independent experiments with five

replicates within each experiment (n = 10 for each bar).

Figure 4. *Rhg1* transcript induction (fold change) relative to (w.r.t.) no-nematode controls

within each plant genotype. The protein encoded by the analyzed transcript is noted in the

upper-left corner of each graph. Wm82, Sus. NIL and Res. NIL plants were inoculated at

VE stage and root samples from infection areas taken at: (A, C and E) 3 dpi, or (B, D and

F) 7 dpi. No statistically significant differences between samples were observed using

ANOVA and Tukey HSD test. The SCN HG 2.5.7 IL used partially overcomes *rhg1-b*

resistance and represents the most prevalent HG type in the USA. Data are from two

Deepak Haarith
Plant Disease

656 independent experiments with three replicate roots assessed within each experiment (n =

657 6 for each bar).

Figure 5. Induction (fold change) of JA-response indicator gene *Glyma.10G109500* and

SA-response indicator gene *Glyma.10G010100* transcripts at 3 dpi and 7dpi, relative to

(w.r.t.) no-nematode controls within each plant genotype within the same experiment.

Wm82, Sus. NIL and Res. NIL soybean accessions inoculated at VE stage; transcript

measured and sample dpi indicated in upper left corner of graph. Significant differences of

means are marked with p-values from ANOVA Tukey HSD tests (p > 0.2 if no p value is

shown). The SCN HG 2.5.7 IL used partially overcomes *rhg1-b* resistance and represents

the most prevalent HG type in the USA. Data are from two independent experiments with

three replicate roots assessed within each experiment (n = 6 for each bar).

Supplemental Figure 1. Same data as in Figure 4 but with data for all genotypes

normalized to mean Sus. NIL no nematode control within same experiment, to show

relative abundances between genotypes. *Rhg1* transcript induction (fold change)

compared to no-nematode controls within each plant genotype. The protein encoded by

the analyzed transcript is noted in the upper-left corner of each graph. Wm82, Sus. NIL

and Res. NIL plants were inoculated at VE stage and root samples from infection areas

taken at: (A, C, E and G) 3 dpi, or (B, D, F and H) 7 dpi. No statistically significant

differences between samples were observed using ANOVA and Tukey HSD test. The

SCN HG 2.5.7 IL used partially overcomes *rhg1-b* resistance and represents the most

Deepak Haarith
Plant Disease

prevalent HG type in the USA. Data are from two independent experiments with three

replicate roots assessed within each experiment (n = 6 for each bar).

Supplemental Figure 2. *Rhg1* transcript induction (fold change) for V1 plants at 3 dpi

compared to no-nematode controls within each plant genotype. The protein encoded by

the analyzed transcript is noted in the upper-left corner of each graph. Wm82, Sus. NIL

and Res. NIL plants were inoculated at V1 stage. (A, C, E) Data normalized to mean for

no-nematode control of same genotype in same experiment. (B, D, F) Same data as A, C,

E with data for all genotypes normalized to mean Sus. NIL no nematode control within

same experiment, to show relative abundances between genotypes. Instances of possible

significant differences of means are marked with p-values from ANOVA Tukey HSD tests

(p > 0.2 if no p value is shown). The SCN HG 2.5.7 IL used partially overcomes *rhg1-b*

resistance and represents the most prevalent HG type in the USA. All experiments were

repeated twice with two replicate roots assessed within each experiment (n = 4 for each

bar).

Deepak Haarith
Plant Disease

Literature Cited

Allen, T. W., Bradley, C. A., Sisson, A. J., Byamukama, E., Chilvers, M. I., Coker, C. M.,

Collins, A. A., Damicone, J. P., Dorrance, A. E., Dufault, N. S., Esker, P. D., Faske,

T. R., Giesler, L. J., Grybauskas, A. P., Hershman, D. E., Hollier, C. A., Isakeit, T.,

Jardine, D. J., Kelly, H. M., Kemerait, R. C., Kleczewski, N. M., Koenning, S. R.,

Kurle, J. E., Malvick, D. K., Markell, S. G., Mehl, H. L., Mueller, D. S., Mueller, J.

D., Mulrooney, R. P., Nelson, B. D., Newman, M. A., Osborne, L., Overstreet, C.,

Padgett, G. B., Phipps, P. M., Price, P. P., Sikora, E. J., Smith, D. L., Spurlock, T.

N., Tande, C. A., Tenuta, A. U., Wise, K. A., and Wrather, J. A. 2017. Soybean

Yield Loss Estimates Due to Diseases in the United States and Ontario, Canada,

from 2010 to 2014. Plant Health Prog. 18:19-27.

Baermann, G. 1917. Eine einfache Methode zur Auffindung von Anklostomum

(Nematoden) Larven in Erdproben. Geneeskundig tijdschrift voor Nederlandsch-

Indië 57:131-137.

Basnet, P., Meinhardt, C. G., Usovsky, M., Gillman, J. D., Joshi, T., Song, Q., Diers, B.,

Mitchum, M. G., and Scaboo, A. M. 2022. Epistatic interaction between *Rhg1-a*

and *Rhg2* in PI 90763 confers resistance to virulent soybean cyst nematode

populations. Theor. Appl. Genet. 135:2025-2039.

Bayless, A. M., Zapotocny, R. W., Grunwald, D. J., Amundson, K. K., Diers, B. W., and

Bent, A. F. 2018. An atypical N-ethylmaleimide sensitive factor enables the viability

of nematode-resistant *Rhg1* soybeans. Proc. Natl. Acad. Sci. U S A 115:E4512-

E4521.

Deepak Haarith
Plant Disease

Bayless, A. M., Zapotocny, R. W., Han, S., Grunwald, D. J., Amundson, K. K., and Bent,

A. F. 2019. The *rhg1-a* (*Rhg1* low-copy) nematode resistance source harbors a

copia-family retrotransposon within the *Rhg1*-encoded alpha-SNAP gene. Plant

Direct 3:e00164.

Bayless, A. M., Smith, J. M., Song, J., McMinn, P. H., Teillet, A., August, B. K., and

Bent, A. F. 2016. Disease resistance through impairment of alpha-SNAP-NSF

interaction and vesicular trafficking by soybean *Rhg1*. Proc. Natl. Acad. Sci. U S A

113:E7375-E7382.

Bent, A. 2022. Exploring Soybean Resistance to Soybean Cyst Nematode. Annu. Rev.

Phytopathol. 60:379-409.

Beyer, S. F., Bel, P. S., Flors, V., Schultheiss, H., Conrath, U., and Langenbach, C. J.

G. 2021. Disclosure of salicylic acid and jasmonic acid-responsive genes provides

a molecular tool for deciphering stress responses in soybean. Sci. Rep. 11:20600.

Bradley, C. A., Allen, T. W., Sisson, A. J., Bergstrom, G. C., Bissonnette, K. M., Bond,

J., Byamukama, E., Chilvers, M. I., Collins, A. A., Damicone, J. P., Dorrance, A. E.,

Dufault, N. S., Esker, P. D., Faske, T. R., Fiorellino, N. M., Giesler, L. J., Hartman,

G. L., Hollier, C. A., Isakeit, T., Jackson-Ziems, T. A., Jardine, D. J., Kelly, H. M.,

Kemerait, R. C., Kleczewski, N. M., Koehler, A. M., Kratochvil, R. J., Kurle, J. E.,

Malvick, D. K., Markell, S. G., Mathew, F. M., Mehl, H. L., Mehl, K. M., Mueller, D.

S., Mueller, J. D., Nelson, B. D., Overstreet, C., Padgett, G. B., Price, P. P., Sikora,

E. J., Small, I., Smith, D. L., Spurlock, T. N., Tande, C. A., Telenko, D. E. P.,

Tenuta, A. U., Thiessen, L. D., Warner, F., Wiebold, W. J., and Wise, K. A. 2021.

Deepak Haarith
Plant Disease

Soybean Yield Loss Estimates Due to Diseases in the United States and Ontario,

Canada, from 2015 to 2019. Plant Health Prog. 22:483-495.

Brucker, E., Carlson, S., Wright, E., Niblack, T., and Diers, B. 2005. *Rhg1* alleles from

soybean PI 437654 and PI 88788 respond differentially to isolates of Heterodera

glycines in the greenhouse. Theor. Appl. Genet. 111:44-49.

Brzostowski, L. F., and Diers, B. W. 2017. Pyramiding of Alleles from Multiple Sources

Increases the Resistance of Soybean to Highly Virulent Soybean Cyst Nematode

Isolates. Crop. Sci. 57:2932-2941.

Butler, K. J., Chen, S., Smith, J. M., Wang, X., and Bent, A. F. 2019. Soybean

Resistance Locus *Rhg1* Confers Resistance to Multiple Cyst Nematodes in

Diverse Plant Species. Phytopathol. 109:2107-2115.

Butler, K. J., Fliege, C., Zapotocny, R., Diers, B., Hudson, M., and Bent, A. F. 2021.

Soybean Cyst Nematode Resistance Quantitative Trait Locus *cqSCN-006* Alters

the Expression of a gamma-SNAP Protein. Mol. Plant Microbe Interact. 34:1433-

1445.

Castanheira, C. M., Falcão, H. G., Ida, E. I., Dias-Arieira, C. R., and Bolanho Barros, B.

C. 2020. *Pratylenchus brachyurus* parasitism on soybean: effects on productivity,

vegetative and nematological parameters and chemical properties. Eur. J. Plant

Pathol. 157:651-661.

Chen, S., Shaeffer, C. C., Wyse, D. L., Nickel, P., and Kandel, H. 2012. Plant-parasitic

Nematode Communities and Their Associations with Soil Factors in Organically

Farmed Fields in Minnesota. J. Nematol. 44:361-369.

Deepak Haarith
Plant Disease

Chowdhury, I. A., Yan, G. P., Kandel, H., and Plaisance, A. 2022. Population

Development of the Root-Lesion Nematode on Soybean Cultivars. Plant Dis.

106:2117-2126.

Colgrove, A. L., and Niblack, T. L. 2005. The effect of resistant soybean on male and

female development and adult sex ratios of. J. Nematol. 37:161-167.

Cook, D. E., Bayless, A. M., Wang, K., Guo, X., Song, Q., Jiang, J., and Bent, A. F.

2014. Distinct Copy Number, Coding Sequence, and Locus Methylation Patterns

Underlie *Rhg1*-Mediated Soybean Resistance to Soybean Cyst Nematode. Plant

Physiol. 165:630-647.

Cook, D. E., Lee, T. G., Guo, X., Melito, S., Wang, K., Bayless, A. M., Wang, J.,

Hughes, T. J., Willis, D. K., Clemente, T. E., Diers, B., Jiang, J., Hudson, M., and

Bent, A. F. 2012. Copy Number Variation of Multiple Genes at *Rhg1* Mediates

Nematode Resistance in Soybean. Science 338:1206-1209.

Dong, J., and Hudson, M. E. 2022. WI12(Rhg1) interacts with DELLAs and mediates

soybean cyst nematode resistance through hormone pathways. Plant Biotechnol.

J. 20:283-296.

Endo, B. Y. 1967. Comparative Population Increase of *Pratylenchus brachyurus* and *P.*

zeae in Corn and in Soybean Varieties Lee and Peking. Phytopathol. 57:118-120.

Favoreto, L., Meyer, M. C., Dias-Arieira, C. R., Machado, A. C. Z., Santiago, D. C., and

Riberiro, N. R. 2019. Diagnosis and Management of Nematodes In Soybean Crop.

Informe Agropecuario 40:18-29.

Glazebrook, J., Chen, W. J., Estes, B., Chang, H. S., Nawrath, C., Metraux, J. P., Zhu,

T., and Katagiri, F. 2003. Topology of the network integrating salicylate and

jasmonate signal transduction derived from global expression phenotyping. Plant J.

34:217-228.

Haarith, D., Das, S., Nelson, E., Zapotocny, R., and Bent, A. F. 2025. Overexpression of

α-SNAP$_{Rhg1}$ Can Improve *rhg1-a* Mediated Soybean Resistance to Soybean Cyst

Nematode. Phytopathol.115(9): https://doi.org/10.1094/PHYTO-02-25-0077-R

Hamawaki, O. T., Hamawaki, R. L., Nogueira, A. P. O., Glasenapp, J. S., Hamawaki, C.

D. L., and da Silva, C. O. 2019. Evaluation of soybean breeding lineages to new

sources of root-knot nematode resistance. Cienc. Agrotec. 43.

https://doi.org/10.1590/1413-7054201943009519

Han, S., Smith, J. M., Du, Y., and Bent, A. F. 2023. Soybean transporter AAT$_{Rhg1}$

abundance increases along the nematode migration path and impacts vesiculation

and ROS. Plant. Physiol. 192:133-153.

Jones, J. T., Haegeman, A., Danchin, E. G., Gaur, H. S., Helder, J., Jones, M. G.,

Kikuchi, T., Manzanilla-Lopez, R., Palomares-Rius, J. E., Wesemael, W. M., and

Perry, R. N. 2013. Top 10 plant-parasitic nematodes in molecular plant pathology.

Mol. Plant Pathol. 14:946-961.

Kleczewski, N., Colsgrove, A., Bowman, D., Harbach, C., and Plewa, D. 2023. A Survey

of Soybean Cyst Nematode Population Densities and Phenotypes in Illinois: 2018

and 2020. Plant Health Prog. 24:221-225.

Lakhssassi, N., Liu, S., Bekal, S., Zhou, Z., Colantonio, V., Lambert, K., Barakat, A.,

and Meksem, K. 2017. Characterization of the Soluble NSF Attachment Protein

gene family identifies two members involved in additive resistance to a plant

pathogen. Sci. Rep. 7:45226.

Lasserre, F., Rivoal, R., and Cook, R. 1994. Interactions between Heterodera avenae

and Pratylenchus neglectus on Wheat. J. Nematol. 26:336-344.

Liu, S., Kandoth, P. K., Lakhssassi, N., Kang, J., Colantonio, V., Heinz, R., Yeckel, G.,

Zhou, Z., Bekal, S., Dapprich, J., Rotter, B., Cianzio, S., Mitchum, M. G., and

Meksem, K. 2017. The soybean *GmSNAP18* gene underlies two types of

resistance to soybean cyst nematode. Nat. Commun. 8:14822.

Livak, K. J. and Schmittgen, T. D. 2001. Analysis of relative gene expression data using

real-time quantitative PCR and the $2^{-\Delta\Delta Ct}$ method. Methods 25:402-8.

Lopez-Nicora, H. D., Ralston, T. I., Diers, B. W., Dorrance, A. E., and Niblack, T. L.

2023. Interactions Among *Heterodera glycines*, *Macrophomina phaseolina*, and

Soybean Genotype. Plant Dis. 107:401-412.

MacGuidwin, A. E., and Bender, B. E. 2012. Estimating Population Densities of Root

Lesion Nematodes, *Pratylenchus* spp., from Soil Samples Using Dual Active and

Passive Assays. Plant Health Prog. 13.E1535-1025.

MacGuidwin, A. E., and Bender, B. E. 2016. Development of a Damage Function Model

for *Pratylenchus penetrans* on Corn. Plant Dis. 100:764-769.

Deepak Haarith
Plant Disease

MacGuidwin, A. E., Smith, D. L., Conley, S. P., and Saikai, K. A. 2023. Prevalence of

Pest Nematodes Associated with Soybean (*Glycine max*) in Wisconsin from 1998

to 2021. J. Nematol. 55:20230053.

Machado, A. C. Z., and Araúujo, J. V. 2016. Broad-sense heritability and variance

component estimates for resistance in Brazilian soybean genotypes. Tropical

Plant Pathology 41:390-396.

Matsye, P. D., Lawrence, G. W., Youssef, R. M., Kim, K. H., Lawrence, K. S., Matthews,

B. F., and Klink, V. P. 2012. The expression of a naturally occurring, truncated

allele of an alpha-SNAP gene suppresses plant parasitic nematode infection. Plant

Mol Biol 80:131-155.

McCarville, M. T., Marett, C. C., Mullaney, M. P., Gebhart, G. D., and Tylka, G. L. 2017.

Increase in Soybean Cyst Nematode Virulence and Reproduction on Resistant

Soybean Varieties in Iowa From 2001 to 2015 and the Effects on Soybean Yields.

Plant Health Prog. 18:146-155.

Melakeberhan, H., and Dey, J. 2003. Competition between *Heterodera glycines* and

Meloidogyne incognita or *Pratylenchus penetrans*: Independent Infection Rate

Measurements. J. Nematol. 35:1-6.

Mokrini, F., Viaene, N., Waeyenberge, L., Dababat, A. A., and Moens, M. 2018.

Investigation of resistance to *Pratylenchus penetrans* and *P. thornei* in international

wheat lines and its durability when inoculated together with the cereal cyst

nematode *Heterodera avenae*, using qPCR for nematode quantification. Eur. J.

Plant Pathol.151:875-889.

Deepak Haarith
Plant Disease

Niblack, T. L., Hussey, R. S., and Boerma, H. R. 1986. Effects of Interactions among

Heterodera glycines, *Meloidogyne incognita*, and Host Genotype on Soybean Yield

and Nematode Population-Densities. J. Nematol. 18:436-443.

Niblack, T. L., Lambert, K. N., and Tylka, G. L. 2006. A model plant pathogen from the

kingdom animalia: Heterodera glycines, the soybean cyst nematode. Annu. Rev.

Phytopathol. 44:283-303.

Nomura, R. B. G., Lopes-Caitar, V. S., Hishinuma-Silva, S. M., Machado, A. C. Z.,

Meyer, M. C., and Marcelino-Guimarães, F. C. 2024. *Pratylenchus brachyurus*:

status and perspectives in Brazilian agriculture. Trop. Plant Pathol. 49:573-589.

Patil, G. B., Lakhssassi, N., Wan, J., Song, L., Zhou, Z., Klepadlo, M., Vuong, T. D.,

Stec, A. O., Kahil, S. S., Colantonio, V., Valliyodan, B., Rice, J. H., Piya, S.,

Hewezi, T., Stupar, R. M., Meksem, K., and Nguyen, H. T. 2019. Whole-genome

re-sequencing reveals the impact of the interaction of copy number variants of the

rhg1 and *Rhg4* genes on broad-based resistance to soybean cyst nematode. Plant

Biotechnol. J. 17:1595-1611.

Pedro, M. S., da Silva, S. A., Picoli, L. H., da Cunha, L. S., and Machado, A. C. Z. 2024.

Experimental and statistical approaches to evaluate the reaction of soybean to

Pratylenchus brachyurus. Trop. Plant Pathol. 49:612-621.

Pedersen, P. 2009. Soybean Growth and Development. Iowa State University

Extension publication PM1945.

Rebois, R.V. and Huettel, R. N. 1986. Population-Dynamics, Root Penetration, and

Feeding-Behavior of Pratylenchus-Agilis in Monoxenic Root Cultures of Corn,

Tomato, and Soybean. J. Nematology 18:392-397.

Rosa de Araújo, T., de Assis dos Santos Diniz, F., Mendes, B. L., Almeida, L. C., and

Santiago, T. R. 2025. *Pratylenchus zeae* is the main root lesion nematode affecting

sugarcane in the world's most important producing regions. Physiol. Mol. Plant

Pathol.136: 102576.

Saikai, K., and MacGuidwin, A. E. 2020. Difference in Lesion Formation by Male and

Female *Pratylenchus penetrans*. J. Nematol. 52.E2020-90.

Saikai, K., and MacGuidwin, A. E. 2022. Impact of *Pratylenchus penetrans* on Soybean

Grown in Wisconsin, U.S.A. Plant Dis. 106:2904-2910.

Schmitt, D. P., and Barker, K. R. 1981. Damage and Reproductive Potentials of

Pratylenchus brachyurus and *Pratylenchus penetrans* on Soybean. J. Nematol.

13:327-332.

Schumacher, L. A., and Bird, G. W. 2025. Distribution of *Pratylenchus* spp. and

Heterodera glycines in Michigan and Combined Reproduction on Peking and PI

437654 Soybean. Plant Health Prog. https://doi.org/10.1094/PHP-10-24-0101-RS

Schweiger, R., Heise, A. M., Persicke, M., and Muller, C. 2014. Interactions between

the jasmonic and salicylic acid pathway modulate the plant metabolome and affect

herbivores of different feeding types. Plant Cell Environ. 37:1574-1585.

Deepak Haarith
Plant Disease

Townshend, J. L. 1984. Anhydrobiosis in *Pratylenchus penetrans*. J. Nematol. 16:282-

289.

Tylka, G. L., Gebhart, G. D., Marrett, C. C., and Mullaney, M. P. 2012. Evaluation of

soybean varieties resistant to soybean cyst nematode in Iowa in 2012. Iowa State

University Extension and Outreach.

https://faculty.sites.iastate.edu/gltylka/files/inline-files/evaluation_of_scn-

resistant_soybean_varieties_in_iowa_-_2018_ipm_52.pdf

Usovsky, M., Robert T. Robbins, Juliet Fultz Wilkes, Devany Crippen, Vijay Shankar, Tri

D. Vuong, Paula Agudelo, and Nguyen, H. T. 2022. Classification methods and

identification of reniform nematode resistance in known soybean cyst nematode

resistant soybean genotypes. Plant Dis. 106:382-389.

Usovsky, M., Lakhssassi, N., Patil, G. B., Vuong, T. D., Piya, S., Hewezi, T., Robbins,

R. T., Stupar, R. M., Meksem, K., and Nguyen, H. T. 2021. Dissecting nematode

resistance regions in soybean revealed pleiotropic effect of soybean cyst and

reniform nematode resistance genes. Plant Genome 14:e20083.

Usovsky, M., Gamage, V. A., Meinhardt, C. G., Dietz, N., Triller, M., Basnet, P., Gillman,

J. D., Bilyeu, K. D., Song, Q., Dhital, B., Nguyen, A., Mitchum, M. G., and Scaboo,

A. M. 2023. Loss-of-function of an alpha-SNAP gene confers resistance to

soybean cyst nematode. Nat. Commun. 14:7629.

Verma, K., Kumari, K., Rawat, M., Devi, K., and Joshi, R. 2025. Crosstalk of Jasmonic

acid and Salicylic acid with other Phytohormones Alleviates Abiotic and Biotic

Stresses in Plants. J. Soil Sci. Plant Nutr. 25:4997-5019.

Yu, N., Lee, T. G., Rosa, D. P., Hudson, M., and Diers, B. W. 2016. Impact of *Rhg1*

copy number, type, and interaction with *Rhg4* on resistance to Heterodera glycines

in soybean. Theor. Appl. Genet. 129:2403-2412.

Zhao, M., Wu, S., Zhou, Q., Vivona, S., Cipriano, D. J., Cheng, Y., and Brunger, A. T.

2015. Mechanistic insights into the recycling machine of the SNARE complex.

Nature 518:61-67.

Zunke, U. 1990. Observations on the Invasion and Endoparasitic Behavior of the Root

Lesion Nematode *Pratylenchus-Penetrans*. J. Nematol. 22:309-320.

Figure 1. Nematode counts on Wm82, IL3849N (*rhg1-a*) and IL3025N (*rhg1-b*) plants at 30 dpi with either or both SCN HG 2.5.7 IL and Pp. (A) Pp counts at 30 dpi at 1:1 SCN: Pp inoculation (B) Pp counts at 30 dpi at 2:1 SCN:Pp inoculation. Blue bars correspond to Pp-only inoculation and orange bars represent combined inoculation of both SCN and Pp. (C) Numbers of SCN cysts recovered at 30 dpi at 1:1 SCN:Pp inoculation. (D) Numbers of SCN cysts recovered at 30 dpi at 2:1 SCN:Pp inoculation. Green bars represent SCN-only inoculation and the light-blue bars represent combined inoculation of both SCN and Pp. Values shown are mean +/- standard error of mean. For the comparisons of greatest interest, statistical significance for differences between means are displayed as p-values from ANOVA Tukey HSD tests performed on data transformed to achieve normality (see Methods). The SCN HG 2.5.7 IL used partially overcomes *rhg1-b* resistance and represents the most prevalent HG type in the USA. Data are for two independent experiments with five replicates within each experiment (n=10 for each bar), using distinct batches of seedlings and fresh nematode inoculum.

161x232mm (300 x 300 DPI)

Figure 2. Pp counts on Wm82, Sus. NIL and Res. NIL plants at 30 dpi in either Pp-only inoculations (blue bars) or a combination of SCN and Pp at 1:1 ratio (orange bars). (A) Counts of all Pp. (B) Ratio of Pp females to males. (C) Ratio of Pp in sand fraction to root fraction. (D) Ratio of Pp juveniles (J2 through J4) to adults (male + female). (E) Counts of Pp juveniles (J2 through J4). Values shown are mean +/- standard error of mean. Instances of possible significant differences of means are marked with p-values from ANOVA Tukey HSD tests performed on data transformed to achieve normality (see Methods; p > 0.2 if no p value is shown). The SCN HG 2.5.7 IL used partially overcomes *rhg1-b* resistance and represents the most prevalent HG type in the USA. Data are for two independent experiments with five replicates within each experiment (n=10 for each bar).

159x241mm (300 x 300 DPI)

Figure 3. SCN cyst counts on Wm82, Sus. NIL and Res. NIL plants at 30 dpi in either SCN-only inoculation (green bars) or a combination of SCN and Pp at 1:1 ratio (light-blue bars). (A) SCN cyst counts. (B) SCN J2 counts (sand + root). (C) SCN male counts (sand + root). Values are mean +/- standard error of mean for non-transformed data. Instances of possible significant differences of means are marked with p-values from ANOVA Tukey HSD tests performed on data transformed to achieve normality (see Methods) (p > 0.2 if no p value is shown). The SCN HG 2.5.7 IL used partially overcomes *rhg1-b* resistance and represents the most prevalent HG type in the USA. Data are for two independent experiments with five replicates within each experiment (n=10 for each bar).

153x231mm (300 x 300 DPI)

Figure 4. *Rhg1* transcript induction (fold change) relative to (w.r.t.) no-nematode controls within each plant genotype. The protein encoded by the analyzed transcript is noted in the upper-left corner of each graph. Wm82, Sus. NIL and Res. NIL plants were inoculated at VE stage and root samples from infection areas taken at: (A, C and E) 3 dpi, or (B, D and F) 7 dpi. No statistically significant differences between samples were observed using ANOVA and Tukey HSD test. The SCN HG 2.5.7 IL used partially overcomes *rhg1-b* resistance and represents the most prevalent HG type in the USA. Data are for two independent experiments with three replicate roots assessed within each experiment (n=6 for each bar).

155x235mm (300 x 300 DPI)

**Figure 5. Induction (fold change) of JA-response indicator gene *Glyma.10G109500*
and SA-response indicator gene *Glyma.10G010100* transcripts at 3 dpi and 7dpi,
relative to (w.r.t.) no-nematode controls within each plant genotype within the
same experiment.** Wm82, Sus. NIL and Res. NIL soybean accessions inoculated at VE
stage; transcript measured and sample dpi indicated in upper left corner of graph.
Significant differences of means are marked with p-values from ANOVA Tukey HSD tests
($p > 0.2$ if no p value is shown). The SCN HG 2.5.7 IL used partially overcomes *rhg1-b*
resistance and represents the most prevalent HG type in the USA. Data are for two
independent experiments with three replicate roots assessed within each experiment
(n=6 for each bar).

156x209mm (300 x 300 DPI)

Supplemental Figure 1. Same data as in Figure 4 but with data for all genotypes normalized to mean Sus. NIL no nematode control within same experiment, to show relative abundances between genotypes. *Rhg1* transcript induction (fold change) compared to no-nematode controls within each plant genotype. The protein encoded by the analyzed transcript is noted in the upper-left corner of each graph. Wm82, Sus. NIL and Res. NIL plants were inoculated at VE stage and root samples from infection areas taken at: (A, C, E and G) 3 dpi, or (B, D, F and H) 7 dpi. No statistically significant differences between samples were observed using ANOVA and Tukey HSD test. The SCN HG 2.5.7 IL used partially overcomes *rhg1-b* resistance and represents the most prevalent HG type in the USA. Data are for two independent experiments with three replicate roots assessed within each experiment (n=6 for each bar).

Supplemental Figure 2. *Rhg1* **transcript induction (fold change) for V1 plants at 3 dpi compared to no-nematode controls within each plant genotype.** The protein encoded by the analyzed transcript is noted in the upper-left corner of each graph. Wm82, Sus. NIL and Res. NIL plants were inoculated at V1 stage. (A, C, E) Data normalized to mean for no-nematode control of same genotype in same experiment. (B, D, F) Same data as A, C, E with data for all genotypes normalized to mean Sus. NIL no nematode control within same experiment, to show relative abundances between genotypes. Instances of possible significant differences of means are marked with p-values from ANOVA Tukey HSD tests (p > 0.2 if no p value is shown). The SCN HG 2.5.7 IL used partially overcomes *rhg1-b* resistance and represents the most prevalent HG type in the USA. All experiments were repeated twice with two replicate roots assessed within each experiment (n=4 for each bar).

Supplementary Table 1. List of all qPCR primers used in this study to quantify mRNA/transcript abundance

Gene name	Primer Direction	5' to 3' Primer Sequence	Reference
Glyma.18G022500 α-SNAP$_{Rhg1}$	Forward	GCAATTGATGAAGAAGATGTTG	Haarith et al., 2025
	Reverse	TCAAGTAATAGCCTCATGCTG	
Glyma.18G022700 WI12$_{Rhg1}$	Forward	AGGTCACGTGTTGCCGTTG	Cook et al., 2012
	Reverse	AAACCACACCAATAACAACAAAGCTCT	
Glyma.18G022400 AAT$_{Rhg1}$	Forward	CGTGTAGAGTCCTTGAAGTACAGC	
	Reverse	ACCAGAGCTGTGATAGCCAACC	
Glyma.10G109500 AAT (JA)	Forward	GAGTTGATACCAGCCAACAG	Beyer et al., 2021
	Reverse	GAGACCAACAAGCAGTGTAG	
Glyma.10G010100 NIMIN1 (SA)	Forward	ATGTTGAACACGGCATTCTC	
	Reverse	GGTACGGTGTGACTTTCTTG	
Glyma.11G079600 NREG1 (Reference gene)	Forward	TCCTGGCTGCTAACTACTTG	
	Reverse	GGTCTTGCGAATTTCCTCTG	

Supplementary Table 2. ANOVA for Pp numbers in 1:1 SCN:Pp inoculation (Wm82, IL3045N and IL3849N)

Formula: aov(Sqrt(Pp count) ~ (Experiment/replicate)+Plant genotypes*Nematode inoculation)

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	4.476	4.4759	12.1744	0.001113	**
Plant genotypes	2	33.379	16.6896	45.3959	2.011e-11	***
Nematode inoculation	1	29.596	29.5964	80.5025	1.696e-11	***
Experiment/replicate	8	9.173	1.1466	3.1189	0.007046	**
Plant genotypes x Nematode inoculation	2	0.037	0.0185	0.0502	0.0951098	
Residuals	44	16.176	0.3676			

Significance : 0 '*' 0.001 '**' 0.01 '*'**

Tukey HSD (p-values)		
Wm82+Pp	IL3849N+Pp	**0.0000004**
Wm82+Pp	IL3025N+Pp	**0.0000718**
Wm82+Both	Wm82+Pp	**0.0001694**
Wm82+Both	IL3849N+Pp	**0**
Wm82+Both	IL3849N+Both	**0.0000008**
Wm82+Both	IL3025N+Pp	**0**
Wm82+Both	IL3025N+Both	**0.0004221**
IL3849N+Pp	IL3025N+Pp	0.6180058
IL3849N+Both	Wm82+Pp	0.5687882
IL3849N+Both	IL3849N+Pp	**0.0000925**
IL3849N+Both	IL3025N+Pp	**0.0116435**
IL3849N+Both	IL3025N+Both	0.3892712
IL3025N+Both	Wm82+Pp	0.9996929
IL3025N+Both	IL3849N+Pp	**0.0000001**
IL3025N+Both	IL3025N+Pp	**0.0000274**

Supplementary Table 3. ANOVA for SCN cyst numbers in 1:1 SCN:Pp inoculation (Wm82, IL3045N and IL3849N)

Formula: aov(log(Cyst count) ~ (Experiment/replicate)+Plant genotypes*Nematode inoculation)

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	14.217	14.217	52.4515	4.367e-09	***
Plant genotypes	2	92.931	46.465	171.7184	<2.2e-16	***
Nematode inoculation	1	0.001	0.001	0.0045	0.94687	
Experiment/replicate	8	1.438	0.180	0.6641	0.71986	
Plant genotypes x Nematode inoculation	2	2.516	1.258	4.6485	0.01462	*
Residuals	45	12.177	0.271			

Significance : 0 '*' 0.001 '**' 0.01 '*'**

Tukey HSD (p-values)		
Wm82+SCN	IL3849N+SCN	**0.000000**
Wm82+SCN	IL3025N+SCN	**0.003122**
Wm82+Both	Wm82+SCN	0.168464
Wm82+Both	IL3849N+SCN	**0.000000**
Wm82+Both	IL3849N+Both	**0.000000**
Wm82+Both	IL3025N+SCN	0.628442
Wm82+Both	IL3025N+Both	0.912637
IL3849N+SCN	IL3025N+SCN	**0.000000**
IL3849N+Both	Wm82+SCN	**0.000000**
IL3849N+Both	IL3849N+SCN	0.500285
IL3849N+Both	IL3025N+SCN	**0.000000**
IL3849N+Both	IL3025N+Both	**0.000000**
IL3025N+Both	Wm82+Pp	0.050215
IL3025N+Both	IL3849N+Pp	**0**
IL3025N+Both	IL3025N+Pp	**0.0002341**

Supplementary Table 4. ANOVA for Pp numbers in 2:1 SCN:Pp inoculation(Wm82, IL3045N and IL3849N)

Formula: aov(log(Pp count) ~ (Experiment/replicate)+Plant genotypes*Nematode inoculation)

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	0.8422	0.8422	5.9161	0.01904	*
Plant genotypes	2	10.9284	5.4642	38.3814	1.876e-10	***
Nematode inoculation	1	7.9374	7.9374	55.7533	2.081e-09	***
Experiment/replicate	8	0.4230	0.0529	0.3714	0.93021	
Plant genotypes x Nematode inoculation	2	0.1748	0.0874	0.6140	0.54565	
Residuals	45	6.4065	0.1424			

Significance : 0 '*' 0.001 '**' 0.01 '*'**

Tukey HSD (p-values)		
Wm82+Pp	IL3849N+Pp	**0.0000364**
Wm82+Pp	IL3025N+Pp	0.4505769
Wm82+Both	Wm82+Pp	**0.0003304**
Wm82+Both	IL3025N+Pp	**0.0000007**
Wm82+Both	IL3849N+Pp	**0**
Wm82+Both	IL3849N+Both	**0.0000004**
Wm82+Both	IL3025N+Both	0.5168828
IL3849N+Pp	IL3025N+Pp	**0.0114356**
IL3849N+Both	Wm82+Pp	0.3737783
IL3849N+Both	IL3849N+Pp	**0.0163912**
IL3849N+Both	IL3025N+Pp	0.9999942
IL3849N+Both	IL3025N+Both	**0.0001528**
IL3025N+Both	Wm82+Pp	0.050215
IL3025N+Both	IL3849N+Pp	**0**
IL3025N+Both	IL3025N+Pp	**0.0002341**

Supplementary Table 5. ANOVA for SCN cyst numbers in 2:1 SCN:Pp inoculation (Wm82, IL3045N and IL3849N)

Formula: aov(Sqrt(Cyst count) ~ (Experiment/replicate)+Plant genotypes*Nematode inoculation)

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	221.4	221.44	21.4969	3.053e-05	***
Plant genotypes	2	5954.2	2977.08	289.0036	<2.2e-16	***
Nematode inoculation	1	2.4	2.42	0.2350	0.6301709	
Experiment/replicate	8	404.6	50.58	4.9097	0.0002125	***
Plant genotypes x Nematode inoculation	2	5.0	2.50	0.2430	0.7852932	
Residuals	45	463.6	10.30			
Significance : 0 '*' 0.001 '**' 0.01 '*'**						

Tukey HSD (p-values)		
Wm82+SCN	IL3849N+SCN	**0.0000000**
Wm82+SCN	IL3025N+SCN	**0.0000632**
Wm82+Both	Wm82+SCN	1.0000000
Wm82+Both	IL3849N+SCN	**0.0000000**
Wm82+Both	IL3849N+Both	0.0000000
Wm82+Both	IL3025N+SCN	**0.0000668**
Wm82+Both	IL3025N+Both	**0.0010353**
IL3849N+SCN	IL3025N+SCN	**0.0000000**
IL3849N+Both	Wm82+SCN	0.0000000
IL3849N+Both	IL3849N+SCN	1.0000000
IL3849N+Both	IL3025N+SCN	**0.0000000**
IL3849N+Both	IL3025N+Both	0.0000000
IL3025N+Both	Wm82+SCN	**0.0009817**
IL3025N+Both	IL3849N+SCN	**0.0000000**
IL3025N+Both	IL3025N+SCN	0.9563879

Supplementary Table 6. ANOVA for Total Pp numbers on NILs

Formula: aov(sqrt(Pp count) ~ (Experiment/replicate)+Plant genotypes*Nematode inoculation)

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	103.184	103.184	122.6661	1.926e-14	***
Plant genotypes	2	26.913	13.456	15.9970	5.649e-06	***
Nematode inoculation	1	14.541	14.541	17.2866	0.0001423	***
Experiment/replicate	8	5.670	0.709	0.8426	0.5706470	
Plant genotypes x Nematode inoculation	2	4.094	2.047	2.4333	0.0992143	
Residuals	45	37.853	0.841			

Significance : 0 '*' 0.001 '**' 0.01 '*'**

Tukey HSD (p-values)		
Wm82+Pp	Wm82+Both	**0.0018532**
Wm82+Pp	Sus. NIL+Pp	0.9932719
Wm82+Pp	Sus. NIL+Both	0.5104347
Wm82+Pp	Res. NIL+Pp	0.0994669
Wm82+Pp	Res. NIL+Both	0.9609094
Wm82+Both	Sus. NIL+Both	0.172806
Wm82+Both	Res. NIL+Both	**0.0001337**
Sus. NIL+Pp	Wm82+Both	**0.0095477**
Sus. NIL+Pp	Sus. NIL+Both	0.8400984
Sus. NIL+Pp	Res. NIL+Pp	**0.0258896**
Sus. NIL+Pp	Res. NIL+Both	0.7362778
Sus. NIL+Both	Res. NIL+Both	0.1255234
Res. NIL+Pp	Wm82+Both	**0.0000003**
Res. NIL+Pp	Sus. NIL+Both	**0.0008159**
Res. NIL+Pp	Res. NIL+Both	0.4436716

Supplementary Table 7. ANOVA for Total Pp Sand:Root ratio on NILs

Formula: aov(sqrt(Pp Sand:Root count ratio) ~ (Experiment/replicate)+Plant genotypes*Nematode inoculation)

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	2.5698	2.5698	8.2737	0.006128	**
Plant genotypes	2	0.7019	0.3509	1.1298	0.332080	
Nematode inoculation	1	9.8983	9.8983	31.8680	1.047e-06	***
Experiment/replicate	8	2.2440	0.2805	0.9031	0.522299	
Plant genotypes x Nematode inoculation	2	7.9391	3.9695	12.7801	4.024e-05	***
Residuals	45	13.9771	0.3106			
Significance : 0 '*' 0.001 '**' 0.01 '*'**						

Tukey HSD (p-values)		
Wm82+Pp	Wm82+Both	0.9992374
Wm82+Pp	Sus. NIL+Pp	0.5181065
Wm82+Pp	Sus. NIL+Both	0.9977317
Wm82+Pp	Res. NIL+Pp	**0.0007679**
Wm82+Pp	Res. NIL+Both	0.0747088
Wm82+Both	Sus. NIL+Both	0.9999991
Wm82+Both	Res. NIL+Both	0.1598771
Sus. NIL+Pp	Wm82+Both	0.3133064
Sus. NIL+Pp	Sus. NIL+Both	0.2696013
Sus. NIL+Pp	Res. NIL+Pp	0.0927678
Sus. NIL+Pp	Res. NIL+Both	**0.0005679**
Sus. NIL+Both	Res. NIL+Both	0.90591
Res. NIL+Pp	Wm82+Both	**0.0002492**
Res. NIL+Pp	Sus. NIL+Both	**0.0001862**
Res. NIL+Pp	Res. NIL+Both	**0.0000001**

Supplementary Table 8. ANOVA for Total Pp Female:Male ratio on NILs

aov(sqrt(Female:Male count ratio) ~ (Experiment/replicate)+Plant genotypes*Nematode inoculation)

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	0.2124	0.21241	1.4121	0.2402	
Plant genotypes	2	4.7374	2.36870	15.7469	4.724e-06	***
Nematode inoculation	1	0.3330	0.33301	2.2139	0.1429	
Experiment/replicate	2	0.1788	0.08940	0.5943	0.5557	
Plant genotypes x Nematode inoculation	2	0.1189	0.05945	0.3952	0.6756	
Residuals	51	7.6716	0.15042			

Significance : 0 '*' 0.001 '**' 0.01 '*'**

Tukey HSD (p-value)		
Wm82+Pp	Wm82+Both	0.9589550
Wm82+Pp	Sus. NIL+Pp	0.9999986
Wm82+Pp	Sus. NIL+Both	0.8754047
Wm82+Pp	Res. NIL+Pp	**0.0286405**
Wm82+Pp	Res. NIL+Both	0.0712761
Wm82+Both	Sus. NIL+Both	0.9997857
Wm82+Both	Res. NIL+Both	**0.0080514**
Sus. NIL+Pp	Wm82+Both	0.9759452
Sus. NIL+Pp	Sus. NIL+Both	0.9121446
Sus. NIL+Pp	Res. NIL+Pp	**0.0221532**
Sus. NIL+Pp	Res. NIL+Both	0.0565343
Sus. NIL+Both	Res. NIL+Both	**0.0036522**
Res. NIL+Pp	Wm82+Both	**0.0027149**
Res. NIL+Pp	Sus. NIL+Both	**0.0011828**
Res. NIL+Pp	Res. NIL+Both	0.9990226

Supplementary Table 9. ANOVA for Total Pp Juveniles (J2+J3+J4) on NILs

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	49.310	49.310	103.5823	2.988e-13	***
Plant genotypes	2	23.238	11.619	24.4071	6.626e-08	***
Nematode inoculation	1	11.532	11.532	24.2241	1.193e-05	***
Experiment/replicate	8	4.080	0.510	1.0712	0.39996	
Plant genotypes x Nematode inoculation	2	4.297	2.149	4.5136	0.01635	*
Residuals	45	21.422	0.476			

Significance : 0 '***' 0.001 '**' 0.01 '*'

Tukey HSD (p-value)		
Wm82+Pp	Wm82+Both	**0.0000487**
Wm82+Pp	Sus. NIL+Pp	0.8636626
Wm82+Pp	Sus. NIL+Both	0.1015930
Wm82+Pp	Res. NIL+Pp	0.0752190
Wm82+Pp	Res. NIL+Both	0.8762907
Wm82+Both	Sus. NIL+Both	0.1126620
Wm82+Both	Res. NIL+Both	**0.0000012**
Sus. NIL+Pp	Wm82+Both	**0.0018727**
Sus. NIL+Pp	Sus. NIL+Both	0.6445899
Sus. NIL+Pp	Res. NIL+Pp	**0.0036033**
Sus. NIL+Pp	Res. NIL+Both	0.2393802
Sus. NIL+Both	Res. NIL+Both	**0.0058241**
Res. NIL+Pp	Wm82+Both	**0.0000000**
Res. NIL+Pp	Sus. NIL+Both	**0.0000000**
Res. NIL+Pp	Res. NIL+Both	0.5404546

Supplementary Table 10. ANOVA for Total Pp Juveniles (J2+J3+J4):Adults (Male+Female) on NILs

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	0.1004	0.10043	1.1352	0.292359	
Plant genotypes	2	1.4132	0.70662	7.9874	0.001075	**
Nematode inoculation	1	0.4680	0.46803	5.2903	0.026135	*
Experiment/replicate	8	0.7269	0.09086	1.0271	0.430151	
Plant genotypes x Nematode inoculation	2	0.3017	0.15083	1.7049	0.193321	
Residuals	51	3.9810	0.08847			

Significance : 0 '***' 0.001 '**' 0.01 '*'

Tukey HSD (p-value)		
Wm82+Pp	Wm82+Both	0.1379612
Wm82+Pp	Sus. NIL+Pp	0.7765115
Wm82+Pp	Sus. NIL+Both	0.0702945
Wm82+Pp	Res. NIL+Pp	0.9867943
Wm82+Pp	Res. NIL+Both	0.9799576
Wm82+Both	Sus. NIL+Both	0.9996147
Wm82+Both	Res. NIL+Both	**0.0259743**
Sus. NIL+Pp	Wm82+Both	0.827706
Sus. NIL+Pp	Sus. NIL+Both	0.653358
Sus. NIL+Pp	Res. NIL+Pp	0.3818849
Sus. NIL+Pp	Res. NIL+Both	0.3468304
Sus. NIL+Both	Res. NIL+Both	**0.0113626**
Res. NIL+Pp	Wm82+Both	**0.0305982**
Res. NIL+Pp	Sus. NIL+Both	**0.013538**
Res. NIL+Pp	Res. NIL+Both	0.9999998

Supplementary Table 11. ANOVA for Total SCN cysts on NILs

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	13.0096	13.0096	115.6445	5.082e-14	***
Plant genotypes	2	1.0089	0.5045	4.4843	0.01675	*
Nematode inoculation	1	2.4579	2.4579	21.8488	2.698e-05	***
Experiment/replicate	8	0.2749	0.0344	0.3054	0.96010	
Plant genotypes x Nematode inoculation	2	0.3164	0.1582	1.4064	0.25560	
Residuals	45	5.0624	0.1125			

Significance : 0 '*' 0.001 '**' 0.01 '*'**

Tukey HSD (p-values)		
Wm82+SCN	Wm82+Both	**0.0039955**
Wm82+SCN	Sus. NIL+SCN	0.9978039
Wm82+SCN	Sus. NIL+Both	**0.0371747**
Wm82+SCN	Res. NIL+SCN	0.0572952
Wm82+SCN	Res. NIL+Both	**0.0007447**
Wm82+Both	Sus. NIL+Both	0.9643226
Wm82+Both	Res. NIL+Both	0.9937006
Sus. NIL+SCN	Wm82+Both	**0.0139943**
Sus. NIL+SCN	Sus. NIL+Both	0.1046342
Sus. NIL+SCN	Res. NIL+SCN	0.1516324
Sus. NIL+SCN	Res. NIL+Both	**0.0028780**
Sus. NIL+Both	Res. NIL+Both	0.7510163
Res. NIL+SCN	Wm82+Both	0.9202250
Res. NIL+SCN	Sus. NIL+Both	0.9999748
Res. NIL+SCN	Res. NIL+Both	0.6445139

Supplementary Table 12. ANOVA for RT q-PCR transcript abundance for 3dpi VE inoculation 1:1 SCN:Pp. Analyses were done on ddCt values = (Ct sample – CtNREG1) - avg(Ct No nematode sample of the same plant genotype – CtNREG1).

AAT$_{Rhg1}$ 3dpi VE inoculation

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	5.456	5.4560	4.8545	0.03178	*
Plant x Nematode Treatment	11	9.183	0.8348	0.7428	0.69323	
Experiment/replicate (root)	4	41.001	10.2504	9.1203	1.003e-05	***
Residuals	55	61.815	1.1239			
Significance : 0 '*' 0.001 '**' 0.01 '*'**						

α-SNAP$_{Rhg1}$ 3dpi VE inoculation

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	6.559	6.5590	5.7852	0.01955	*
Plant x Nematode Treatment	11	19.880	1.8073	1.5941	0.12644	
Experiment/replicate (root)	4	47.520	11.8800	10.4784	2.212e-06	***
Residuals	55	62.357	1.1338			
Significance : 0 '*' 0.001 '**' 0.01 '*'**						

WI12$_{Rhg1}$ 3dpi VE inoculation

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	0.9014	0.9014	2.1180	0.1513	
Plant x Nematode Treatment	11	4.8492	0.4408	1.0358	0.4289	
Experiment/replicate (root)	4	24.2835	6.0709	14.2642	4.749e-08	***
Residuals	55	23.4081	0.4256			
Significance : 0 '*' 0.001 '**' 0.01 '*'**						

α-SNAP$_{Rhg1}$HC 3dpi VE inoculation

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	1.0645	1.0645	5.7852	0.01955	*
Plant x Nematode Treatment	3	0.7744	0.7744	1.5941	0.12644	
Experiment/replicate (root)	4	6.6203	6.6203	10.4784	2.212e-06	***
Residuals	15	0.2607	0.2607			
Significance : 0 '*' 0.001 '**' 0.01 '*'**						

AAT (JA response) 3dpi VE inoculation

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	0.210	0.2097	0.3154	0.5767	
Plant x Nematode Treatment	11	5.761	0.5238	0.7876	0.6511	
Experiment/replicate (root)	4	62.289	15.5722	23.4174	2.438e-11	***
Residuals	55	36.574	0.6650			
Significance : 0 '*' 0.001 '**' 0.01 '*'**						

NIMIN1 (SA response) 3dpi VE inoculation

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	19.621	19.6207	28.1187	2.087e-06	***
Plant x Nematode Treatment	11	53.984	4.9076	7.0331	2.848e-07	***
Experiment/replicate (root)	4	37.738	9.4345	13.5207	9.706e-08	***
Residuals	55	38.378	0.6978			
Significance : 0 '*' 0.001 '**' 0.01 '*'**						

Supplementary Table 13. ANOVA for RT q-PCR transcript abundance for 7dpi VE inoculation 1:1 SCN:Pp. Analyses were done on ddCt values = (Ct sample – CtNREG1) - avg(Ct No nematode sample of the same plant genotype – CtNREG1).

AAT$_{Rhg1}$ 3dpi VE inoculation

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	1.711	1.7713	1.8180	0.1830820	
Plant x Nematode Treatment	11	4.747	0.4316	0.4585	0.9205702	
Experiment/replicate (root)	4	25.400	6.3499	6.7458	0.0001717	***
Residuals	55	51.772	0.9413			
Significance : 0 '*' 0.001 '**' 0.01 '*'**						

α-SNAP$_{Rhg1}$ 3dpi VE inoculation

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	0.5753	0.57525	1.0549	0.30889	
Plant x Nematode Treatment	11	4.2437	0.38579	0.7074	0.72612	
Experiment/replicate (root)	4	6.7825	1.69564	3.1093	0.02231	*
Residuals	55	29.9937	0.54534			
Significance : 0 '*' 0.001 '**' 0.01 '*'**						

WI12$_{Rhg1}$ 3dpi VE inoculation

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	0.0001	0.00007	0.0001	0.991382	
Plant x Nematode Treatment	11	2.2066	0.20060	0.3510	0.969036	
Experiment/replicate (root)	4	10.4876	2.62191	4.5881	0.002868	**
Residuals	55	31.4299	0.57145			
Significance : 0 '*' 0.001 '**' 0.01 '*'**						

α-SNAP$_{Rhg1}$HC 3dpi VE inoculation

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	0.154	0.1535	0.4299	0.5220	
Plant x Nematode Treatment	3	0.261	0.0871	0.2440	0.8643	
Experiment/replicate (root)	4	92.762	23.1906	64.9315	2.735e-09	***
Residuals	15	5.357	0.3572			
Significance : 0 '*' 0.001 '**' 0.01 '*'**						

AAT (JA response) 3dpi VE inoculation

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	9.498	9.498	19.3388	5.061e-05	***
Plant x Nematode Treatment	11	4.119	0.374	0.7625	0.6747	
Experiment/replicate (root)	4	170.788	42.697	86.9339	<2.2e-16	***
Residuals	55	27.013	0.491			
Significance : 0 '*' 0.001 '**' 0.01 '*'**						

NIMIN1 (SA response) 3dpi VE inoculation

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	7.284	7.284	7.0647	0.01027	*
Plant x Nematode Treatment	11	16.375	1.489	1.4438	0.18027	
Experiment/replicate (root)	4	151.174	37.793	36.6539	6.423e-15	***
Residuals	55	56.710	1.031			
Significance : 0 '*' 0.001 '**' 0.01 '*'**						

Supplementary Table 14. ANOVA for RT q-PCR transcript abundance for 3dpi V1 inoculation 1:1 SCN:Pp. Analyses were done on ddCt values = (Ct sample – CtNREG1) - avg(Ct No nematode sample of the same plant genotype – CtNREG1).

AAT$_{Rhg1}$ 3dpi VE inoculation

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	11.146	11.1458	4.7505	0.03653	*
Plant x Nematode Treatment	11	23.710	2.1554	0.9187	0.53433	
Experiment/replicate (root)	2	4.453	2.2265	0.9490	0.39745	
Residuals	33	77.425	2.3462			

Significance : 0 '***' 0.001 '**' 0.01 '*'

α-SNAP$_{Rhg1}$ 3dpi VE inoculation

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	10.949	10.9491	5.5854	0.02415	*
Plant x Nematode Treatment	11	25.812	2.3466	1.1970	0.32688	
Experiment/replicate (root)	2	12.521	6.2607	3.1938	0.05396	*
Residuals	33	64.690	1.9603			

Significance : 0 '***' 0.001 '**' 0.01 '*'

WI12$_{Rhg1}$ 3dpi VE inoculation

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	1.150	1.1501	1.0715	0.3081	
Plant x Nematode Treatment	11	19.808	1.8008	1.6777	0.1227	
Experiment/replicate (root)	2	2.368	1.1840	1.1031	0.3438	
Residuals	33	35.420	1.0733			

Significance : 0 '***' 0.001 '**' 0.01 '*'

α-SNAP$_{Rhg1}$HC 3dpi VE inoculation

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	1.69	1.69	0.0175	0.8963569	
Plant x Nematode Treatment	11	70.75	6.43	0.0667	0.9999709	
Experiment/replicate (root)	2	2834.42	1417.21	14.6904	0.0001972	***
Residuals	17	1640.03	96.47			

Significance : 0 '***' 0.001 '**' 0.01 '*'

Plant Disease

AAT (JA response) 3dpi VE inoculation

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	1.272	1.2724	0.6445	0.42785	
Plant x Nematode Treatment	11	24.857	2.2598	1.1445	0.36080	
Experiment/replicate (root)	2	15.736	7.8679	3.9850	0.02817	*
Residuals	33	65.154	1.9744			
Significance : 0 '*' 0.001 '**' 0.01 '*'**						

NIMIN1 (SA response) 3dpi VE inoculation

	Df	Sum. Sq	Mean Sq.	F value	Pr > F	Significance
Experiment	1	0.015	0.0152	0.0080	0.929478	
Plant x Nematode Treatment	11	74.317	6.7561	3.5360	0.002533	**
Experiment/replicate (root)	2	14.637	7.3183	3.8303	0.032259	*
Residuals	32	61.140	1.9106			
Significance : 0 '*' 0.001 '**' 0.01 '*'**						