The Connectome Interpreter Toolkit

Yijie Yin[1], Judith Hoeller[2], Alexander Mathiasen[3], Joanna Tsang[3], Monique Estelle Charrier[3], and Albert Cardona[4,1]

[1]*Department of Physiology, Development and Neuroscience, University of Cambridge*
[2]*Howard Hughes Medical Institute at Janelia Research Campus*
[3]*Unaffiliated*
[4]*MRC Laboratory of Molecular Biology*

September 30, 2025

Abstract

Complete synaptic wiring diagrams, or connectomes, of whole brains open new opportunities for studying the structure-function relationship of neural circuits. However, the large number of nodes and edges in the graphs makes the analysis challenging. Here, we present the **Connectome Interpreter**, an open-source software toolkit for efficient graph traversal to find polysynaptic pathways, compute the effective connectivity and receptive fields for arbitrarily deep neurons, slice out subcircuits, and non-linear but differentiable circuit modeling, implemented using efficient approaches tailored to the high density and size of connectomes such as that of the fruit fly *Drosophila melanogaster*. Our approach delivers results orders of magnitude faster than conventional methods in consumer computer hardware. We demonstrate the capabilities of our toolkit with select applications, including quantifying the density of polysynaptic connections in the whole adult fruit fly brain, exploring the necessity for non-linearities in circuit modeling, and combining known function of neurons with the connectome to aid in formulating hypotheses of circuit function.

1 Introduction

Neural circuit architecture constrains the functional capabilities of brains, with mapped neuronal wiring diagrams, or connectomes, serving as a basis for disambiguating competing hypotheses of neural function [1]. Recent large-scale research projects reconstructed neurons and labeled their synapses from nanometer-resolution volumes of the brains of fruit flies and humans [2–10]. These large, directed, weighted, signed (excitation/inhibition) connectomes establish a strong foundation for studying brain function: all circuits are now accessible for analysis.

Upon completion of connectome mapping, research efforts can then proceed into circuit analysis. However, connectomic datasets reveal unexpected complexity: an average neuron connects to ∼130 other neurons (median ∼90) and

~80 cell types (median ~55) in the central brain of the fly, indicating that the synaptic connectivity graph is dense enough to render circuit analysis a challenge. While a common approach is to focus on the stronger synaptic weights, the continuous distribution of connection weights makes defining a well-justified threshold challenging. Moreover, because most connections are weak, thresholding effectively removes most of the connections, inadvertently discarding potentially crucial signal integration patterns.

The observed dense synaptic connectivity of brain connectomes indicates that many neurons simultaneously participate in multiple sensory-motor pathways underlying natural behaviors. Yet, knowledge of neuronal function typically advances by experimentally manipulating and monitoring specific neurons with controlled, limited stimuli. Such an approach poses a fundamental scaling problem: exhaustively testing the relationships among combinations of sensory inputs, neuronal activations, and behavioral outputs is infeasible.

The study of the relationship between circuit structure and function fundamentally involves linking the known to the unknown (Figure 1A) — a central theme of neuroscience since its inception. Whether mapping sensory receptive fields or assigning functional roles based on circuit interactions, neuroscientists leverage prior data (e.g., characterized sensory neurons or behaviors) to interpret unexamined neural circuits. The advent of multiple connectomic datasets amplify both the need and opportunity for new methods that systematically bring functional data into circuit analysis.

To address these challenges, we developed an approach to study neuronal function in which the connectome serves as not just a lookup table but an effective basis for translating structural complexity into functional insights.

Here, we introduce *Connectome Interpreter*, an open-source toolbox (`https://github.com/YijieYin/connectome_interpreter`) specifically developed to analyze circuit function on the basis of combining connectomes with known but potentially sparsely-annotated neuronal function. The toolbox accelerates common network analysis operations such as effective connectivity calculation [11–14] and polysynaptic path finding, while enabling exploration of previously inaccessible questions, including the identification of the stimulus optimal for activating neurons of interest. We demonstrate the utility of our toolbox through highlighting neural circuits of potential functional significance, and examining the role of non-linearity in signal propagation.

2 Results

2.1 Table of neurons with known functions

Knowledge of the function of at least some cell types is essential to infer the function of any particular cell type, by propagating functional information across the circuits comprised in a connectome. While connectomes are generally only now starting to be available, literature on the function of cell types has been accumulating for decades in multiple animal models.

We have collected information on the function of known cell types from the fly neuroscience literature into a table (`https://tinyurl.com/known-neuron-function`). By mapping the known function of some neurons onto the circuits described by the connectome, we open up the opportunity to efficiently generate connectome-based hypotheses of neural function in the fly.

The table shows that there is partial knowledge on at least 30% of the cells, comprising 5% of all cell types. In particular, the table includes 2.4% (174/7175) of the central brain cell types, 15% (71/481) of the descending cell types, 17% (60/352) of visual projection, and 4% (15/423) of the sensory neurons, indicating that more is known closer to the sensory/motor periphery of the central nervous system (as illustrated by Figure 1A).

We recognize that literature coverage of cell types continuously evolves, and therefore we opened up the table to the neuroscience community to further contribute by reviewing, correcting, and adding new entries via: `https://tinyurl.com/known-neuron-function`, to ensure all relevant work is represented. For maximal accessibility, we integrated the table into the Codex data curation initiative (`https://codex.flywire.ai`, 15), and include a snapshot in the supplementary material 1.

2.2 Connectome-driven target neuron selection

Our goal is to deliver effective tools for, given sparse functional data and dense connectomic data, identifying specific, understudied neurons for subsequent and targeted experimentation. Here, we provide both linear and non-linear approaches to tackle this analysis.

2.2.1 Linear approach: effective connectivity

The effective connectivity approximates the amount of influence a source neuron could have on a target neuron, across all polysynaptic pathways of length n. One popular approach to estimating the effective connectivity is to multiply connection weights along each path from a source to a target neuron, and then add up this product across all parallel pathways between the same source and target [11–14].

However, multiplying and summing connection weights, which is equivalent to taking the power of the connectivity matrix (Figure 1B-C), is in practice limited to networks with a small number of nodes (i.e., neurons in the connectome), because the computational cost of multiplying two $n \times n$ connectivity matrices scales to the third power of the number of nodes (n), and linearly with the number of hops (h) in the longest polysynaptic path to consider ($O(n^3 * h)$) (Figure 1F, Methods 5.0.1).

Our method avoids the n^3 scaling by first leveraging the sparsity of the fly connectome: for its 140,000 neurons, on average any one neuron receives synapses from 1/1,000 of all neurons, i.e., about 140, when considering all mapped synapses. The sparse matrix representation brings huge savings in

Figure 1: Linear approaches of target neuron selection with examples. **A.** Illustration of connectome-based hypothesis generation. Green fill indicates existing research on that neuron. **B.** Illustration of the equivalence of two representations of a network M. **C.** Calculation of all-to-all two-hop effective connectivity is equivalent to multiplying the connectivity matrix with itself once. The blue dashed boxes highlight the numbers multiplied to yield 0.46, and the fill colors show the "column picture of matrix multiplication": in this case, the first column of the result matrix is the sum of: $0\times$ the first column, $0.2\times$ the second column, and $0.8\times$ the last column. **D-E**. We combine sparsity with chunking, by dividing the last matrix into one dense column chunk at a time, and multiplying it with all the sparse matrices before (E), resulting in a dense matrix the size of the chunk. We then combine the (thresholded) dense column chunks together. **F.** Comparison of our sparse chunked method with dense matrix multiplication on GPU and CPU, for different connectivity matrix sizes. **G.** To calculate signed effective connectivity, we divide the connectivity matrix into excitatory rows and inhibitory rows, and multiply each separately with excitatory and inhibitory rows, for all combinations of ee, ei, ie,ii. We then add ee and ii together for the "new" excitatory rows; and ei and ie together for the "new" inhibitory rows. **H.** HRN_VP5 connects to the ipsilateral $DNb05$. The edge labels indicate normalized input for $DNb05$. **I.** $JO-D$ connects to the contralateral $CB0916$. **J.** $JO-A$ and $JO-B$ connect to the ipsilateral $DNp01$ (giant fiber), $DNp02$ and $DNp11$. **K.** oviLN connects bilaterally to $DNc01$ and $DNc02$ through indirect inhibition. **L.** Direct and effective signed connectivity between 100 randomly-selected cell types, colored by the path length. **M.** $LPLC1$ connects bilaterally, indirectly, to $DNa05$, through excitatory connections. The connections are stronger contralaterally. **N.** Neuroglancer scene snapshot, visualizing two-hop effective connectivity from individual $LPLC1$ neurons, each in its own color, to $DNa05$ (cyan). Yellow means stronger connectivity. All wiring diagrams are thresholded at 1% normalized input.

computer memory usage and number of operations to perform, skipping all the zeros during matrix multiplication.

However, the divergence and convergence of the connectivity (Figure 4A) entail that the matrix products quickly become denser the more matrices are multiplied, rendering sparse representation no longer useful for longer polysynaptic hops. We address this challenge by dividing the dense products into chunks with a customizable size (Figure 1D-E), such that only one small dense chunk has to be held in memory at any one time, drastically reducing the memory requirement for such large-scale operations.

Our approach makes the same computation orders of magnitude faster, taking minutes instead of hours or days to compute the effective connectivity between all neurons (Figure 1F), for a connectome the size of the brain of the adult fly brain, at ~140,000 [2, 3]. Further, we cater to a diversity of available resources in typical neuroscience labs, by making our implementation customizable in memory consumption, facilitating execution on widely-accessible computational resources, from a laptop to Google Colab, when conventional approaches (such as dense matrix multiplication; see Methods 5.0.1) otherwise require high performance computing resources for these datasets. The results can then be saved and used for instant queries of effective connectivity between any groups of neurons.

As a proof of principle, we combine the (effective) connectivity matrix with our table of functionally characterized neurons (Supplement 2 and 3). This yields, for every cell type, a "contextualized receptive field" (RFc), which we define as the receptive field in terms of not only peripheral sensory neurons (the 'conventional' receptive field), but also the functionally characterized neurons anywhere in the connectome. This approach was previously illustrated for the receptive field of the lobula plate tangential cells (LPTCs) via their direct T4/T5 inputs [16]. To illustrate the utility of the contextualized receptive field, we look for unreported (direct and indirect) putative functional relationships between reported cell types in the literature.

We consider the following monosynaptic (direct) connections:

1. [17] showed that *Drosophila melanogaster* prefer humidity. Correspondingly, we find that the HP5 glomerulus of the hygrosensory receptor neurons, sensitive to humidity [18], synapse onto the ipsilateral side of DNb05 which is responsible for ipsiversive turning (turning towards the ipsilateral side; [19]) (Figure 1H). Through effectively combining the existing knowledge of cell types with the connectivity matrix, we identify the HP5-to-DNb05 connection as potentially the neural circuit basis for turning towards the side with higher humidity, a hypothesis that could now be tested experimentally.

2. Johnston's organ neurons, JO-D, shown to signal static deflection of the antenna [20], are connected to the contralateral CB0916 which is involved in neck motor control [21] (Figure 1I). The direct synapse from JO-D to CB0916 could therefore drive neck movement upon antennal deflection.

3. Johnston's organ neurons JO-A and JO-B detect sound [22, 23]. We find that they synapse directly not only to the Giant Fiber neuron as shown before [24], but also onto additional descending neurons (DNp02 and DNp11) implicated in directional escaping in response to visual input from different angles [25] (Figure 1J). We thus hypothesize that these descending neurons could also drive directional escaping in response to sound.

We considered as well the following polysynaptic (indirect) connection: [26] showed that oviINs inhibit oviDNs responsible for oviposition. We show that they also indirectly inhibit DNc01 and DNc02 (Figure 1K), two cell types known to release SIFamide [27, 28]. Activation of SIFamide-releasing neurons enhances appetitive behavior and food intake [27], while silencing them results in promiscuous flies [28]. The indirect inhibition from oviINs to DNc01 and DNc02 thus indicates that when oviINs are inhibited during oviposition [26], DNc01 and DNc02 can be disinhibited, putatively enhancing appetitive behavior and increasing food intake, while making flies less susceptible to mating. Correspondingly, [29] showed that females become unreceptive immediately after mating until the end of egg laying.

2.2.2 Linear approach: effective connectivity with excitation/inhibition

Distinguishing excitation from inhibition is crucial for determining how the signals flow through the neural circuits. We developed a method for computing the signed effective connectivity without having to enumerate all combinations of excitation and inhibition in polysynaptic pathways. Our implementation again profitably uses a chunked sparse multiplication approach, retaining computational accessibility.

We demonstrate the capabilities of our method for computing the all-to-all signed effective connectivity for the FAFB/FlyWire connectome. We find that the effective excitation and inhibition between two cell types become similar as the path length between them increases (up to 5 hops) (Figure 1L): an analysis that would be computationally impractical without our chunked sparse matrix approach.

We computed the RFc for all cell types in the adult fly brain, considering the excitatory and inhibitory connections (Supplement 4). As an example, we consider here a circuit for visual looming stimulus. LPLC1 is a visual projection neuron type present in repeated instances tiling the visual field and that detects visual loom [30]. Downstream, our analysis surfaced that DNa05 integrates indirect excitatory input from LPLC1 (Figure 1M). DNa05 is a neuron type that drives flight saccades towards the ipsilateral side [31, 32]. We show further that LPLC1s provide stronger polysynaptic excitation to the contralateral DNa05, with ventral LPLC1 neurons showing stronger connections (Figure 1N). This asymmetry in the strength of the contralateral polysynaptic excitatory pathways suggests that a looming stimulus on one side would activate DNa05 more strongly on the contralateral side, causing the fly to saccade away from the stimulus. And indeed, a looming stimulus on the left activates the right DNa05,

leading the fly to turn right [31]. Furthermore, the observed dorsal-ventral gradient of connectivity between LPLC1 and DNa05 (Figure 1N) may make the flies more sensitive to ventral stimuli when flying, since similar synaptic gradients are functional in directional escape responses and visual object pursuit [25, 33].

2.2.3 Non-linear target neuron selection

The linear approaches described above are useful for measuring the general influence from one group of neurons to another averaged across time, but inadequate for modeling the non-linear interactions in real time, where, for the same increment in input, the neuron's increment in activation depends on its current activation state (Figure 2A). For example, if a neuron's activity is just below the threshold potential, then minimal additional excitation results in an action potential. To incorporate the non-linearity, we build a simple model based on the following assumptions (Figure 2B-C):

1. A neuron cannot be 'negatively active', and its activation saturates;

2. A signal takes about twice the time to propagate through two synaptic hops than one;

3. Excitatory and inhibitory signals propagate at identical speeds;

4. Each neuron is modeled as a single unit.

We further require that the model be differentiable, so that experimental data can be fit via backpropagation, and for the application of activation-maximization for computing the optimal stimulus (see section 2.2.4). With these assumptions and requirements, we build a model with the following activation function:

$$act_{i,t+1} = \tanh(ReLU(excitability \times \sum_{j=1}^{j=n} W_{ji} \times act_{j,t} + bias_i + tau \times act_{i,t}))$$

where W_{ji} (a scalar, i.e., a number) is the connection weight from neurons j to i, and $act_{i,t+1}$ (a scalar) is the activation of neuron i at time point $t+1$. $bias_i$ is the bias (a non-negative scalar) for neuron i, reflecting the baseline activity observed in experimental studies. tau controls the extent to which neuron i's activity persists across time.

In other words, the activation of neuron i at the next timestep is the weighted sum of the upstream partners' activation at the current timestep, multiplied by an *excitability* parameter (a scalar), added to the baseline activity and the remaining activity from neuron i itself, and then passed through a rectified linear unit (*ReLU*) and a tanh function. The *ReLU* guarantees that a neuron is not negatively active, and the *tanh* guarantees that the activation saturates. We chose tanh over a *sigmoid* for a sharper slope when the input is weak.

Figure 2: Non-linear modeling and optimal stimulus search. **A.** Non-linearity introduces context-dependence. The blue and red colors are based on the applying the function to x and y, indicated on the top of each column. Each black line in the last column indicates when the color is the same for all points on that line. When the line is straight, there is no context-dependence, i.e., the same change in x always results in the same change in color. This is the case for the linear plot, the ReLU plot when $x > 0$ and $y > 0$; and approximately true for the tanh plot when x and y are approximately 0. There is context-dependence when the lines are curved: the curvature indicates that the same change in x could result in difference changes in color, depending on the current value of x and y (the context). **B.** The input to the model is two-dimensional: the activation of input neurons across time. **C.** The model has a layered structure, where each layer represents a timepoint, and contains all neurons. The activity of all neurons is also two-dimensional (number of neurons, number of timepoints). The connectome with signed connectivity is between any two adjacent timepoints. Some neurons can be both externally stimulated (pink) and activated by upstream connectivity (blue). Neurons' activations are illustrated with fill colors. In activation maximization, some neurons' activation values at some timepoints are set as the target, and gradient descent looks for the two-dimensional input neurons' activity across time that reproduces the target. **D.** The activation function of the model. **E.** Subtractive inhibition, as long as it scales with the input, is divisive in effect, i.e., it changes the slope/excitability of the target neuron.

We further implemented the possibility of modeling a subset of inhibitory connections as contributing to divisive normalization, where inhibitory input scales down the gain of excitatory input ('divisive') [34], instead of subtracting from the excitatory input ('subtractive'). Divisive normalization is implemented as a change in the *excitability* of the target neurons, scaled by the connection weight with the input, the input activation, and a user-defined divisive normalization strength parameter.

We show that the model is capable of divisive normalization without this explicit implementation (Figure 2E). For instance, suppose neuron A is excitatory, and neuron B provides feedback inhibition through $A \rightarrow B \rightarrow A$ connections, then *the inhibition from B to A scales with A's activity* (due to the $A \rightarrow B$ connection), rendering this inhibitory connection divisive in effect. The scaling of B's activity according to A's is necessary for this subtractive inhibition to be "divisive in effect", suggesting that an explicit implementation of divisive normalization is still required otherwise.

We model temporal summation of inputs in a way that preserves differentiability, which is necessary for parameter tuning and activation maximization. In our model, a neuron only integrates presynaptic input from the timestep before. However, its presynaptic partners can accumulate activation across time through tau, the activity decay parameter; when tau is set to zero there isn't any accummulation. When presynaptic neurons have $tau > 0$ then the postsynaptic neuron receives inputs that are representative of a temporal integration, and therefore in this way we implement temporal summation without spiking and hence maintaining differentiability.

Our model supports targeted specification of experimentally observed data by declaring simulated activity patterns, defining activation profiles over time, and customizing excitability and bias parameters, per neuron. Furthermore, our implementation supports fitting select parameters based on training data.

2.2.4 Explainable-AI (XAI)-assisted interpretation

Building a model answers the question of "what would happen following a given input", but not "what input should be given in the first place". In the analysis of the connectome, determining which stimulus pattern best activates a select group of neurons is not straightforward given the complexity of the synaptic connectivity, which includes many divergent and convergent polysynaptic pathways with multiple lateral, feedback and recurrent connections. Furthermore, attempting to try all possible combinations of input patterns is impractical, given the high-dimensional sensory input space in the fly's ecological niche.

We approach the interpretation of the connectome by determining optimal input patterns for select neurons anywhere in the brain with a technique from the field of explainable-AI: activation maximization [35, 36]. This technique manipulates inputs to optimize for the activation of target neuron(s) of interest, without searching through the entire space of possible stimuli, by formulating the search as an optimization problem that can be solved with gradient descent (Figure 2B-C). To obtain the most parsimonious stimulus and polysynaptic

pathways to the target neurons possible, we define a loss function that minimizes the average activation across all neurons during the optimization.

The activation maximization approach merely delivers the optimal pattern of input neurons' activity to reach the specified target. Therefore, the same approach supports addressing the case of setting an arbitrary, potentially time-varying, activation pattern as the target, for a select group of neurons (for example descending neurons), and then searching for the optimal activity pattern for the input neurons (for example olfactory receptor neurons).

In addition, the flexibility in specifying which neurons can be manipulated (similar to how some neurons are activated by optogenetics in a lab experiment) supports the use case of determining not only the conventional receptive field, but also the contextualised receptive field (RFc). This is useful because linking concepts, such as looming stimulus represented by LC4 [30, 37], with behaivor, can sometimes be more informative than linking individual photoreceptor activations with behaviour.

2.3 Pathfinding: retrieving polysynaptic pathways from inputs to targets

While the complete connectome of a whole brain can be vast, understanding of circuit function can advance by studying one subcircuit at a time. Experimental neuroscience excels at targeting select neurons in the brain, and connectomics datasets provide the basis for recovering the subcircuit between those neurons and, together with neurotransmitter signatures [10], formulating mechanistic hypotheses of circuit function.

However, identifying multi-hop polysynaptic pathways from sources to targets is computationally demanding, even impractical, in dense, highly interconnected networks like a connectome (Figure 3C). Restricting the analysis to the shortest polysynaptic pathways or thresholding graph edges to retain only the strongest connections trades off completeness for practicality, yielding potentially limited insight.

We successfully retrieve complete circuits relating source and target neurons by introducing an efficient implementation of an algorithm that rapidly retrieves all possible circuit pathways between two neuron groups, limited only by defining the number of hops and an optional threshold on the connection weight. Our approach yields results in seconds for connectomes such as the adult fruit fly brain in consumer-grade laptops (Figure 3C).

2.3.1   An efficient search algorithm

First we represent the connectome as a table (a Python Pandas dataframe with indexed columns) with n rows, where each row is a connection with 3 properties: the presynaptic neuron ID, the postsynaptic neuron ID, and the connection weight, with an $O(n)$ cost. Second, perform a bidirectional breadth-first search, by subsetting the table once per hop in each direction, using the set of IDs of the neurons of each group (source and target) as keys to retain, and proceed

Figure 3: Path-finding. **A.** We identify intermediate neurons on paths of length n between source and target neurons through bidirectional breadth-first search: we propagate selection downstream from the source neurons (through the orange edges), and upstream from the target neurons (through the blue edges), until they meet at the middle, where we take the intersection of the neurons reached, those indirectly connected to both the source neurons and the target neurons (green ticks). We then propagate this selection backwards upstream (through the blue ticks) and downstream (orange ticks) to sources and targets. **B.** Alternatively, instead of propagating the selection bidirectionally, we use the pre-computed (thresholded) effective connectivity from the source, to intersect with those connected with the targets, starting from the last layer. **C.** We benchmark the speed of our algorithms ("oflength" refers to panel **A**, "iterative" refers to panel **B**) against Python packages networkx [38] and networkit [39], for different path lengths (x axis) and source node count. **D.** Direct connections from visual projection neurons to DNa10, where connection strength $> 1\%$ of DNa10's total input. **E.** The three-hop visual receptive field through the individual neurons in each type that connects to DNa10 in **D**, shown in hexagonal heatmaps, based on the hexagonal coordinates from [40]. Equivalent three-hop and four-hop receptive fields are attached in Supplement S1.

iteratively. The termination condition is reaching half the specified number of hops (or half minus one for one end when odd) from each side. The result is the complete set of edges or connections, relating all neurons in the circuit between the source and target neurons and spanning the specified polysynaptic pathway depth.

We also implemented a second method, consisting in propagating the neuron selection upstream from the target neurons only, and intersect those connected to the target neurons with those indirectly connected to the source neurons, using the precomputed effective connectivity (Figure 3B). Though slightly faster, this algorithm relies on the (thresholded) effective connectivity matrices, which by design (for performance tuning and dramatic data storage savings) can be constructed to leave out weak connections via thresholding.

Both methods retrieve the synaptic graph of the subcircuit of interest relating all source and target neurons. If the complete list of polysynaptic pathways was required (where each pathway contains n edges from a source to a target for a path of length n), depth-first search can proceed based on the extracted subcircuit only, so that every run returns a hit, bypassing the unsuccessful searches therefore gaining in performance.

Our approach significantly reduces unnecessary operations, enabling interactive path-finding in mere seconds on widely accessible consumer hardware or Google Colab, improving performance substantially over conventional methods that can take hours.

2.3.2 Example search: a visual receptive field

In the fruit fly brain, the optic lobe alone contains about 50,000 neurons on one side [9]. The circuits for visual processing are organised in repeated columnar units corresponding to ommatidia that tile the surface of the eye, capturing a two-dimensional slice of the visual field in a retinotopic map [9, 41, 42]. Deeper in the brain, neural circuits for input convergence across the visual field results in individual neurons responding to specific visual features [16].

To showcase our approach to swiftly retrieve neural circuits between source and target neurons separated by polysynaptic pathways, we investigate the visual receptive field of DNa10, a descending neuron cell type hypothesized to control wing thrust, lift, and pitch on the basis of connectomics data [32]. Our preparatory analysis revealed that DNa10 receives over 1% of its direct synaptic input from visual projection neuron types LLPC3, LPLC4, LC10d, LC10c, LTe64, LC22, each of which is composed of individual neurons tiling the visual field (Figure 3D).

Assuming different VPNs detect different visual features, we ask whether the visual features DNa10 detects are location-dependent. We retrieved 3-hop polysynaptic pathways, from the large monopolar cells $L1$, $L2$ and $L3$ (directly postsynaptic to photoreceptors), and photoreceptors $R7$ and $R8$, to DNa10, separately through each VPN type. Based on the columnar assignments of the large monopolar cells and photoreceptors in [40], we group the pathways by hexagonal coordinates tiling the visual field, and calculate the effective connec-

Figure 4: Density and effective connectivity of polysynaptic pathways. We randomly select 100 cell types as sources, and 100 as targets from the FAFB/FlyWire connectome. We look for poly-synaptic pathways between them one pair at a time, and calculate effective connectivity when such pathways exist, for all 10,000 pairs, for path lengths between 2 and 5. **A.** Histogram of the number of cell type pairs with paths of the tested lengths, given a threshold on all the direct connections in the pathways. **B.** Histogram of effective connectivity between source and target cell types for the paths found. The x axis are square-root-transformed, and the y axis log-transformed for visualization.

tivity from these coordinates to DNa10, to generate the hexagonal heatmaps describing the visual-feature-specific receptive field (Figure 3E). The equivalent plots for 4-hop and 5-hop visualizations are in Supplement S1. The visual features that DNa10 can respond to could thus vary depending upon the area of the visual field, which is plausible for a descending neuron involved in flight control.

2.3.3   Structuring circuits into layers

Structuring a circuit into layers facilitates the identification of motifs of known computational properties such as feedforward, feedback and lateral inhibition, as well as winner-take-all architectures and recurrent motifs where prior processed signals interact with newly incoming signals. To this end, we integrated a metric of information flow [43] which labels neurons into layers as a probabilistic function of the connection weight (see Methods 5.1).

2.4   Depth and density of brain circuits

With the ability to swiftly retrieve polysynaptic pathways across the whole fly brain connectome, we set to measure how many hops separate any one cell type from another. We sample 100 random cell types as sources, and another 100 random cell types as targets, and then search for polysynaptic pathways from 1 single source cell type to 1 single target cell type at a time . Across all 10,000 possible pairs of cell types, we found that, for a connection threshold of 0, ∼70% of tested pairs yielded pathway lengths of 2 (i.e., one single neuron in between), a result that changes dramatically for a connection threshold of 1%, yielding only

2% of the pairs with a pathway of 2 hops (Figure 4A). For polysynaptic pathways of 5 hops, the difference is much smaller between a threshold of 0 (100%) and 1% (∼84%). These findings indicate that the fly connectome is very dense when considering weak connections or polysynaptic pathways of longer length, with significant implications for reductionist approaches to studying brain function.

2.5   Impact of circuit density on effective connectivity

Using the pathways between the selected cell types, we compute the effective connectivity locally, to compare the effective connectivity value across path lengths. Effective connectivity calculation typically uses connection weights normalized by the total number of post-synapses for the postsynaptic neuron, and is therefore a number between 0 and 1. Taking the product of such numbers intuitively results in a smaller number the more times the multiplication takes place. However, we show here that effective connectivity is similar in magnitude across path lengths. This is likely related to the density of the connectivity, i.e., the abundant divergence and convergence in the nervous system, such that there are more pathways between two cell types the longer the path length is. The accumulation of these pathways likely counteracts the diminishing product of connection weights.

2.6   Effect of non-linearity

Non-linearity introduces context dependence with important consequences for neuronal activation when integrating different types (e.g., excitatory or inhibitory) of upstream signals with different weights. Here, we consider two toy circuits to exemplify when a non-linear model produces neural dynamics different from a linear model. The first example illustrates the role of *tanh* (saturation) and the second the role of *ReLU* (non-negativity).

2.6.1   Modeling convergent feedforward excitation and inhibition to explore the role of *tanh*

In the first example, we explore how the difference in weights for feedforward excitation and feedforward inhibition changes the activity of the output neuron across varying strengths of input (Figure 5A).

To demonstrate the differences between the linear and non-linear neuron activation responses to increasing inputs, we constructed the following network model with parallel feedforward excitation and feedforward inhibition converging onto a single output neuron. Hence, a single input unit transmits signals to two units (the interneurons), one excitatory, with a stronger weight, and one inhibitory, with a weaker weight, and both are connected to the single output neuron. The strength of the weights ($w1$ and $w2$) represent here the aggregate convergent input from many neurons, which can co-activate in ecologically-realistic environments. Here we model this convergence with one unit for simplicity.

With the non-linear model, we find that the output neuron's activity is not always the highest when the input neuron's activity is the highest (Figure 5B). In contrast, higher input entails higher output for the linear model (Figure 5C).

Our reasoning is the following. The up-and-down pattern of output neuron responses in Figure 5B arises from the differences in the excitatory drive onto the excitatory ($w1$) and inhibitory ($w2$) interneurons. When $w1 > w2$, at a high input value, the excitatory interneuron saturates first. At even higher inputs, the excitatory interneuron can no longer increase the excitatory drive to the target, therefore limiting the feedforward excitation, whereas the inhibitory drive can be higher still. This effect stems from the saturating non-linear region of the $tanh$ function (Figure 2D), supporting its use in our neuron model.

We further explored this phenomenon by varying the difference in connection weights from the input neuron onto the excitatory and inhibitory interneurons (when $w1 + w2 = 1$) (Figure 5B). When $w1 = w2 = 0.5$, the excitation and inhibition onto the target neuron cancel out; when $w1 - w2 = 1$, $w2 = 0$, the output neuron only receives excitation, so it quickly saturates; the nonlinearity in response manifests the most when $1 > w1 > w2 > 0$, where the excitatory interneuron approaches saturation, and the inhibitory interneuron, while providing inhibition, does not approach saturation yet.

This example highlights the context-dependence in non-linear models: the excitatory interneuron is approaching saturation, so further excitation makes little difference in its activation; in contrast, the inhibitory interneuron does not saturate yet, so little excitation leads to substantial increase in activity. Together, they cause the non-linear response in the output neuron.

2.6.2 Differential transmission of inhibition and excitation: role of the non-negativity constraint with $ReLU$

In the second example, we explore how the existence of inhibitory neurons, while enforcing a non-negativity constraint on neuron activity, favors the use of a non-linear model over a linear one. Here, the non-linearity of $ReLU$ plays a key role by introducing a non-negativity constraint: when an interneuron receives more inhibition than excitation (such as that in Figure 5D), the net activity is zero, instead of negative, and therefore does not pass on any of such excess inhibition – just like in biological neurons. The target neuron thus receives less inhibition than the connection weight alone would suggest if a linear model was used.

Together, these two examples highlight how simple non-linearities - saturation via $tanh$ and thresholding via $ReLU$ - can reshape circuit responses in ways not predictable from linear connectivity alone.

3 Discussion

We have introduced the Connectome Interpreter (`https://github.com/YijieYin/connectome_interpreter`), an open-source computational toolbox that translates the complex structural wiring of a brain into testable functional hypothe-

Figure 5: Toy circuits for probing non-linearity. **A.** A simple circuit with both feed-forward excitation ($w1 \rightarrow w3$) and feed-forward inhibition ($w2 \rightarrow w4$). We assume $w3 = w4$. **B.** Higher input activation may not lead to higher target activation in a non-linear model, depending on the difference in saturation (here due to the difference between $w1$ and $w2$) between feed-forward excitation and feed-forward inhibition. **C.** Higher input activation leads to higher target activation in a linear model, as long as effective excitation $>$ effective inhibition. **D.** A simple circuit where the effective inhibition (through $w9 \rightarrow w10 \rightarrow w11$) outweighs the effective excitation (through $w6 \rightarrow w7 \rightarrow w8$). However, activation of the inputs in the non-linear model activates, instead of inhibits, the target, because the neuron downstream of $w10$ is silenced and does not pass on the inhibition.

ses. We demonstrated that out approach for computing effective connectivity, with or without excitation/inhibition, scales efficiently to large connectomic datasets such as the adult fruit fly brain with ∼140,000 neurons and millions of synaptic connections [2, 3] with predicted neurotransmitter signatures [10]. By integrating the fly connectome with a curated and community-expandable compendium of known neuronal functions, our methods move beyond a static anatomical map to enable the dynamic exploration of signal propagation, both linearly and non-linearly, across the entire brain. Our toolkit facilitates combining experimentally acquired neuronal activity data with the connectome and the synaptic signatures to infer *in silico* the receptive field of any neuron in the brain, and to slice out relevant subnetworks between select sets of source and target neurons, aiding in bridging the gap between circuit architecture and neural computation.

3.1 The need for the whole connectome to analyze individual neurons and circuits

The fruit fly connectome is very dense (Figure 4A; [44]). By this we mean that the activity of any one neuron will depend on the activity of many other neurons. Therefore, our intuition falls short in formulating hypotheses of when or whether a neuron would be active as a function of the activity of some other neurons.

With our toolkit, we measured the density of 5-hop polysynaptic connectivity between random cell types, at 1% threshold for normalized input and not considering the sign of the connections. We found that for 84% of all possible pairs of cell types there was at least 1 polysynaptic path (Figure 4A), which strikes us as very high. Our finding highlights the abundance of convergent and divergent connections in the nervous system of the fly, and has implications in the analysis of circuit function.

For example, let's consider action selection in the fruit fly. The observed dense connectivity fully supports how the fruit fly, in response to a specific sensory stimulus, selects one course of action while suppressing others, because all other inputs must be integrated towards taking the decision, which requires strong convergence. Furthermore, all other possible courses of action have to be suppressed, which requires both divergence (lateral inhibition) and feedback inhibition to setup expectations via corollary discharge. With the Connectome Interpreter toolkit, dense connectomes can be analyzed with ease to assist in formulating hypotheses of neural circuit function that can be tested experimentally in the laboratory by monitoring and manipulating neuronal activity.

3.2   Necessity of non-linear models

Biological neurons are non-linear in nature, given a dynamically adjustable temporal integration window, a spike that resets input accumulation and transmits signals to postsynaptic neurons, and a refractory period that limits maximal activity, by limiting the maximum number of spikes per unit of time (the firing rate). A consequence of spiking is that neurons are either silent (zero) or active (positive), and cannot be negatively active. While graded potential neurons exist, for example in the optic lobe of the fly [45], their activity dynamics is also constrained to be non-negative and are subject to thresholds for synaptic transmission, both of which are better represented by non-linear functions.

A good model captures enough from the system under study that insight can be derived from the study of the model that would not be possible otherwise, often because of the impracticality of observing the whole system with enough spatial and temporal resolution. What makes a good model then depends very much on the goals of the desired analysis. Here, we set to capture fundamental features of neuronal activity such as the non-negativity constraint and the saturation of activity, features that all reflect different aspects of a neuron's non-linear activation function, while retaining differentiability for using experimental data to fit model parameters and for computing the optimal stimulus with activation maximization.

We modeled the saturation of neural activity with a *tanh* and the non-negativity constraint with a *ReLu* centered at zero. In biological neurons, *tanh*-like saturation has been shown experimentally [46], and the prevalence of such saturation in neural responses likely depends on the overall activity level of the system: as network activity increases, more neurons approach their firing-rate ceiling and saturate. The *ReLU* function trivially implements the non-negativity constraint by discarding all inputs below a defined threshold,

here being zero. The non-negativity constraint models an effective deficit in inhibitory transmission: the fact that inhibition does not propagate beyond the postsynaptic neuron as negative signals would when not prevented to do so. The effective deficit in inhibitory transmission requires only that a neuron receives more net inhibition than the amount that would completely silence it by canceling out excitatory input. Given that essentially all brain circuits are composed of both excitatory and inhibitory neurons, the condition of effective inhibitory deficit is presumably widespread, indicating that the non-negativity is fundamental to capture in the model for the propagation of excitatory and inhibitory signals across the connectome.

We tested our model and found that nonlinearities such as saturation and thresholding can have non-intuitive effects, where higher input activation may not entail higher target activation (Figure 5A-B), due to $tanh$-based saturation of neurons; and where effective net inhibition may not entail the inhibition of the target (Figure 5D), due to the silencing of the inhibitory path through $ReLU$.

3.3 The Connectome Interpreter toolkit in practice

We show the utility of combining existing literature on neuron function with effective connectivity, by highlighting previously unreported links between neurons with known functions (Figure 1H-K, Figure 1M-N, Figure 3D-E). The interpretability of these links underscores the promise of contextualized receptive fields (RFc) for hypothesizing on the functions of currently understudied neurons. Together, effective connectivity and contextualized receptive field provide a principled way to move beyond direct synapses, link indirectly-connected neurons representing high-level concepts, discover putative functional relationships between neurons, and generate testable hypotheses of neural function.

Our model provides flexible capabilities to incorporate experimentally-tested features of the nervous system, such as divisive normalization, baseline activity, excitability, and persistence of activity across time. The interpretability of the parameters makes incorporating and testing assumptions straightforward, where the model serves as a conduit for propagating activity, and a hypothesis-generation tool by answering "what-if" scenarios under plausible assumptions. The library of functions we provide facilitates the interpretation and visualization of the model results, guiding the analysis of the model's predictions.

The toolkit includes library utility functions for calculating effective connectivity and path-finding, that operate orders of magnitude faster than other methods. The performance improvement we achieved, together with the accessibility afforded by lowering the minimum hardware requirements, democratizes data analysis and therefore supports a qualitative change in approaches to connectomic analyzes. We offer a range of examples in the form of annotated programs as online notebooks, alongside detailed documentation, for easy adoption. Our software, implemented as open source and thoroughly documented, can be used as a standalone or incorporated into existing data-query platforms such as neuPrint [47], Codex [15] and CATMAID [48], to further lower the barriers of access and facilitate the discovery of neurons and circuits relevant to any and

all experimental paradigms inquiring into brain function.

4   Acknowledgements

We thank Mateo Espinosa Zarlenga for his helpful advice on activation maximization. We thank Katherine Nagel, Kavin Nuñez and Shamik Dasgupta for their feedback and suggestions on the package. We thank Gregory S.X.E. Jefferis and Michael Reiser for their support. We thank the LMB High Performance Computing facilities for the computational resources. This project was funded by a Wellcome Trust Investigator Award to Albert Cardona (Ref: 205038/Z/16/Z) and the MRC LMB core funding.

5   Methods

5.0.1   Calculating effective connectivity

A standard approach is to multiply the weights in a multi-hop path between two neurons, and add the products across parallel pathways [11, 13, 14, 49], without any non-linearity such as thresholding or saturation. This can be implemented as multiplication of the connectivity matrix with itself (see explained in Figure 1C). Calculating matrix power scales to the third power of the number of nodes because: for matrix multiplication $A \times B = C$, to calculate each element in C, n multiplications need to take place (n elements in a row of A, each multiplied with the corresponding element in a column of B). There are $n \times n$ elements in C.

While theoretically straightforward, this approach rapidly encounters practical limits: Storing a 140,000 x 140,000 connectivity matrix in computer random access memory (RAM), where each element takes 32 bits (float32), requires:

$$\frac{140000 \times 140000 \text{ elements } \times 32 \text{ bits per element } \div 8 \text{ bits per byte}}{10^9 \text{ bytes per gigabyte}} = 78.4 \text{ gigabyte}$$

, exceeding the capacity of common laptops. Matrix multiplication requires at least three times this amount (result = one matrix x another matrix). Matrix multiplication is faster on GPUs than CPUs (Figure 1F). However, the maximum RAM an A100 chip has is 80GB, indicating that additional parallelization is necessary, even on high performance computing clusters, to take powers of the dense matrix the size of an adult fly brain.

One inherent aspect of calculating polysynaptic strengths by multiplication is the rapid diminishing (for normalized weights) or amplification (for synapse counts) of values as the pathway length increases. We believe this accurately reflects divergence and convergence in the nervous system: the signal in one neuron can be traced to many in a small number of hops, corresponding to the increase in connection density (Figure 4A). Our default option therefore does not process the products further.

Nonetheless, users have the flexibility to normalize these pathway strengths using an n-th root transformation, to "control for the path lengths". However, due to the divergence and convergence, the rooted effective connectivity values tend to be bigger with longer path lengths. This limitation of diminishing or amplification of values is better addressed using the non-linear model (Section 2.2.3).

5.1 Inferring Hierarchy from Connectivity to Sensory Neurons

To facilitate interpretation of large connectivity matrices, we define a hierarchy of neurons roughly based on how many hops away a neuron is from sensory neurons. This hierarchy approximates the feedforward processing of information from sensory neurons toward (motor-proxy) descending neurons, while revealing deviations due to lateral and feedback connectivity.

In an idealized feedforward network, neurons are organized into discrete layers based on the number of hops that separate them from inputs. However, neurons in real brains are more richly interconnected. For example, a sensory neuron may strongly connect to neuron A and only weakly connect to neuron B, and A strongly connects to B. In this case, we have to decide if neuron B is one hop or two hops away from sensory neurons. An additional complication is that downstream neurons may connect back to sensory neurons, obfuscating a simple layering.

We adopt a nonlinear signal propagation model previously developed for the Drosophila olfactory system [43], which simulates the hierarchical spread of activity from an initial set of active neurons. The model treats each neuron as binary–active or inactive–and updates activity in discrete steps. At each step, inactive neurons are probabilistically activated based on their total weighted input from already active neurons. This probability follows a sigmoid function, capturing both integration and saturation. Once active, a neuron remains active for all subsequent steps.

The layer of a neuron is defined as the average number of steps it takes to become active in this simulation, given a starting set of input neurons. These layers capture a continuous (non-integer-valued) position in a hierarchy that starts at the input neurons. The model is implemented using the Navis toolbox [50] and allows users to visualize pathways together with the hierarchical position of neurons. The hierarchy can also be used to identify feedforward, lateral, and feedback neurons, i.e., neurons which predominantly connect to other neurons of later, similar, or earlier layers, respectively.

5.2 Modeling

For user friendliness, we implement parameter-sharing: for parameters such as bias, slope, and divisive normalization, users need to first specify how the

neurons are to be groups (by e.g. cell type), and then simply use dictionaries that map from the group names to the values of the corresponding group.

We also provide training capability. Given that data from the experimental literature are not always time series, but sometimes a number representing average activity across time, we support training using either the average activation value for each neuron, or time series data. Parameter sharing also applies to training, such that each neuron group have the same parameter values after training.

5.2.1 Saliency

We implement another technique from explainable AI, saliency, which calculates the gradients of the target with respect to the input. Compared to activation maximization, which looks for the best stimulus to reach the target activation, saliency addresses the question of "which part of the current stimulus (i.e. which input neurons, among all activated) matters more for the current target neuron activation?". Note that due to non-linearity, the importance only refers to the current activation value for both the input and the target. The important part of the stimulus may change for a different target neuron activation value.

5.3 User-centric design

We aim for user-centric design in writing the package, and therefore included a number of convenience functions for visualization of pathways in (interactive) wiring diagrams, with options to sort nodes by connection weight, or user-specified order. Effective connectivity can not only be visualized as colored tables, but also in neuroglancer, or for visual information, in hexagonal heatmaps or eyemaps [51]. Pathways can also be plotted through layers in neuroglancer.

To make the non-linear model interpretable, we implement a function that extracts the "relevant neurons" - those that are connected (poly-synaptically) to the target with a connectivity and activity threshold. Conversely, users can start from a pathway and query the activity of all neurons in a pathway. In addition, the activity evolution of groups of neurons can be visualized in an interactive plot, or in neuroglancer.

To facilitate research on vision and navigation, we add functions that construct looming stimuli and sine-shaped stimuli.

To further reduce the barrier of linking stimulus space to neuron activation space, we collate published datasets of neruon responses to odour presentations [46, 52? –55], so that the community can easily combine experimental literature with connectome-based modeling.

Last but not least, we illustrate the functionalities of the software through example Google Colab notebooks (`https://github.com/YijieYin/connectome_interpreter`), using public datasets, to enable browser-based analysis without local installation.

6   Supplementary materials

Figure S1: Four-hop (**A**) and five-hop (**B**) visual receptive field through the individual neurons in each visual projection type that connects to DNa10 in Figure 3D

.

1. Supplementary table 1. A snapshot of the table of neurons with known functions.

2. Supplementary table 2. Direct connectivity combined with table of neurons with known functions.

3. Supplementary table 3. Effective connectivity combined with table of neurons with known functions.

4. Supplementary table 4. Signed effective connectivity combined with table of neurons with known functions.

References

[1] Winfried Denk, Kevin L. Briggman, and Moritz Helmstaedter. Structural neurobiology: missing link to a mechanistic understanding of neural computation. *Nature Reviews Neuroscience*, 13(5):351–358, May 2012. Publisher: Nature Publishing Group.

[2] Philipp Schlegel, Yijie Yin, Alexander S. Bates, Sven Dorkenwald, Katharina Eichler, Paul Brooks, Daniel S. Han, Marina Gkantia, Marcia dos Santos, Eva J. Munnelly, Griffin Badalamente, Laia Serratosa Capdevila, Varun A. Sane, Alexandra M. C. Fragniere, Ladann Kiassat, Markus W. Pleijzier, Tomke Stürner, Imaan F. M. Tamimi, Christopher R. Dunne,

Irene Salgarella, Alexandre Javier, Siqi Fang, Eric Perlman, Tom Kazimiers, Sridhar R. Jagannathan, Arie Matsliah, Amy R. Sterling, Szi-chieh Yu, Claire E. McKellar, Marta Costa, H. Sebastian Seung, Mala Murthy, Volker Hartenstein, Davi D. Bock, and Gregory S. X. E. Jefferis. Whole-brain annotation and multi-connectome cell typing of Drosophila. *Nature*, 634(8032):139–152, October 2024. Publisher: Nature Publishing Group.

[3] Sven Dorkenwald, Arie Matsliah, Amy R. Sterling, Philipp Schlegel, Szi-chieh Yu, Claire E. McKellar, Albert Lin, Marta Costa, Katharina Eichler, Yijie Yin, Will Silversmith, Casey Schneider-Mizell, Chris S. Jordan, Derrick Brittain, Akhilesh Halageri, Kai Kuehner, Oluwaseun Ogedengbe, Ryan Morey, Jay Gager, Krzysztof Kruk, Eric Perlman, Runzhe Yang, David Deutsch, Doug Bland, Marissa Sorek, Ran Lu, Thomas Macrina, Kisuk Lee, J. Alexander Bae, Shang Mu, Barak Nehoran, Eric Mitchell, Sergiy Popovych, Jingpeng Wu, Zhen Jia, Manuel A. Castro, Nico Kemnitz, Dodam Ih, Alexander Shakeel Bates, Nils Eckstein, Jan Funke, Forrest Collman, Davi D. Bock, Gregory S. X. E. Jefferis, H. Sebastian Seung, and Mala Murthy. Neuronal wiring diagram of an adult brain. *Nature*, 634(8032):124–138, October 2024. Publisher: Nature Publishing Group.

[4] Julia Buhmann, Arlo Sheridan, Caroline Malin-Mayor, Philipp Schlegel, Stephan Gerhard, Tom Kazimiers, Renate Krause, Tri M. Nguyen, Larissa Heinrich, Wei-Chung Allen Lee, Rachel Wilson, Stephan Saalfeld, Gregory S. X. E. Jefferis, Davi D. Bock, Srinivas C. Turaga, Matthew Cook, and Jan Funke. Automatic detection of synaptic partners in a whole-brain Drosophila electron microscopy data set. *Nature Methods*, 18(7):771–774, July 2021.

[5] Louis K Scheffer, C Shan Xu, Michal Januszewski, Zhiyuan Lu, Shin-ya Takemura, Kenneth J Hayworth, Gary B Huang, Kazunori Shinomiya, Jeremy Maitlin-Shepard, Stuart Berg, Jody Clements, Philip M Hubbard, William T Katz, Lowell Umayam, Ting Zhao, David Ackerman, Tim Blakely, John Bogovic, Tom Dolafi, Dagmar Kainmueller, Takashi Kawase, Khaled A Khairy, Laramie Leavitt, Peter H Li, Larry Lindsey, Nicole Neubarth, Donald J Olbris, Hideo Otsuna, Eric T Trautman, Masayoshi Ito, Alexander S Bates, Jens Goldammer, Tanya Wolff, Robert Svirskas, Philipp Schlegel, Erika Neace, Christopher J Knecht, Chelsea X Alvarado, Dennis A Bailey, Samantha Ballinger, Jolanta A Borycz, Brandon S Canino, Natasha Cheatham, Michael Cook, Marisa Dreher, Octave Duclos, Bryon Eubanks, Kelli Fairbanks, Samantha Finley, Nora Forknall, Audrey Francis, Gary Patrick Hopkins, Emily M Joyce, SungJin Kim, Nicole A Kirk, Julie Kovalyak, Shirley A Lauchie, Alanna Lohff, Charli Maldonado, Emily A Manley, Sari McLin, Caroline Mooney, Miatta Ndama, Omotara Ogundeyi, Nneoma Okeoma, Christopher Ordish, Nicholas Padilla, Christopher M Patrick, Tyler Paterson, Elliott E Phillips, Emily M Phillips, Neha Rampally, Caitlin Ribeiro, Madelaine K Robertson, Jon Thomson Rymer, Sean M Ryan, Megan Sammons, Anne K Scott, Ash-

ley L Scott, Aya Shinomiya, Claire Smith, Kelsey Smith, Natalie L Smith, Margaret A Sobeski, Alia Suleiman, Jackie Swift, Satoko Takemura, Iris Talebi, Dorota Tarnogorska, Emily Tenshaw, Temour Tokhi, John J Walsh, Tansy Yang, Jane Anne Horne, Feng Li, Ruchi Parekh, Patricia K Rivlin, Vivek Jayaraman, Marta Costa, Gregory SXE Jefferis, Kei Ito, Stephan Saalfeld, Reed George, Ian A Meinertzhagen, Gerald M Rubin, Harald F Hess, Viren Jain, and Stephen M Plaza. A connectome and analysis of the adult Drosophila central brain. *eLife*, 9:e57443, September 2020. Publisher: eLife Sciences Publications, Ltd.

[6] Michael Winding, Benjamin D. Pedigo, Christopher L. Barnes, Heather G. Patsolic, Youngser Park, Tom Kazimiers, Akira Fushiki, Ingrid V. Andrade, Avinash Khandelwal, Javier Valdes-Aleman, Feng Li, Nadine Randel, Elizabeth Barsotti, Ana Correia, Richard D. Fetter, Volker Hartenstein, Carey E. Priebe, Joshua T. Vogelstein, Albert Cardona, and Marta Zlatic. The connectome of an insect brain. *Science*, 379(6636):eadd9330, March 2023. Publisher: American Association for the Advancement of Science.

[7] Meike Sievers, Alessandro Motta, Martin Schmidt, Yagmur Yener, Sahil Loomba, Kun Song, Johannes Bruett, and Moritz Helmstaedter. Connectomic reconstruction of a cortical column, March 2024. Pages: 2024.03.22.586254 Section: New Results.

[8] Alexander Shapson-Coe, Michał Januszewski, Daniel R. Berger, Art Pope, Yuelong Wu, Tim Blakely, Richard L. Schalek, Peter H. Li, Shuohong Wang, Jeremy Maitin-Shepard, Neha Karlupia, Sven Dorkenwald, Evelina Sjostedt, Laramie Leavitt, Dongil Lee, Jakob Troidl, Forrest Collman, Luke Bailey, Angerica Fitzmaurice, Rohin Kar, Benjamin Field, Hank Wu, Julian Wagner-Carena, David Aley, Joanna Lau, Zudi Lin, Donglai Wei, Hanspeter Pfister, Adi Peleg, Viren Jain, and Jeff W. Lichtman. A petavoxel fragment of human cerebral cortex reconstructed at nanoscale resolution. *Science*, 384(6696):eadk4858, May 2024. Publisher: American Association for the Advancement of Science.

[9] Aljoscha Nern, Frank Loesche, Shin-ya Takemura, Laura E. Burnett, Marisa Dreher, Eyal Gruntman, Judith Hoeller, Gary B. Huang, Michał Januszewski, Nathan C. Klapoetke, Sanna Koskela, Kit D. Longden, Zhiyuan Lu, Stephan Preibisch, Wei Qiu, Edward M. Rogers, Pavithraa Seenivasan, Arthur Zhao, John Bogovic, Brandon S. Canino, Jody Clements, Michael Cook, Samantha Finley-May, Miriam A. Flynn, Imran Hameed, Alexandra M. C. Fragniere, Kenneth J. Hayworth, Gary Patrick Hopkins, Philip M. Hubbard, William T. Katz, Julie Kovalyak, Shirley A. Lauchie, Meghan Leonard, Alanna Lohff, Charli A. Maldonado, Caroline Mooney, Nneoma Okeoma, Donald J. Olbris, Christopher Ordish, Tyler Paterson, Emily M. Phillips, Tobias Pietzsch, Jennifer Rivas Salinas, Patricia K. Rivlin, Philipp Schlegel, Ashley L. Scott, Louis A. Scuderi, Satoko

Takemura, Iris Talebi, Alexander Thomson, Eric T. Trautman, Lowell Umayam, Claire Walsh, John J. Walsh, C. Shan Xu, Emily A. Yakal, Tansy Yang, Ting Zhao, Jan Funke, Reed George, Harald F. Hess, Gregory S. X. E. Jefferis, Christopher Knecht, Wyatt Korff, Stephen M. Plaza, Sandro Romani, Stephan Saalfeld, Louis K. Scheffer, Stuart Berg, Gerald M. Rubin, and Michael B. Reiser. Connectome-driven neural inventory of a complete visual system. *Nature*, 641(8065):1225–1237, May 2025. Publisher: Nature Publishing Group.

[10] Nils Eckstein, Alexander Shakeel Bates, Andrew Champion, Michelle Du, Yijie Yin, Philipp Schlegel, Alicia Kun-Yang Lu, Thomson Rymer, Samantha Finley-May, Tyler Paterson, Ruchi Parekh, Sven Dorkenwald, Arie Matsliah, Szi-Chieh Yu, Claire McKellar, Amy Sterling, Katharina Eichler, Marta Costa, Sebastian Seung, Mala Murthy, Volker Hartenstein, Gregory S. X. E. Jefferis, and Jan Funke. Neurotransmitter classification from electron microscopy images at synaptic sites in Drosophila melanogaster. *Cell*, 187(10):2574–2594.e23, May 2024. Publisher: Elsevier.

[11] Feng Li, Jack W Lindsey, Elizabeth C Marin, Nils Otto, Marisa Dreher, Georgia Dempsey, Ildiko Stark, Alexander S Bates, Markus William Pleijzier, Philipp Schlegel, Aljoscha Nern, Shin-ya Takemura, Nils Eckstein, Tansy Yang, Audrey Francis, Amalia Braun, Ruchi Parekh, Marta Costa, Louis K Scheffer, Yoshinori Aso, Gregory SXE Jefferis, Larry F Abbott, Ashok Litwin-Kumar, Scott Waddell, and Gerald M Rubin. The connectome of the adult Drosophila mushroom body provides insights into function. *eLife*, 9:e62576, December 2020. Publisher: eLife Sciences Publications, Ltd.

[12] H. Sebastian Seung. Predicting visual function by interpreting a neuronal wiring diagram. *Nature*, 634(8032):113–123, October 2024. Publisher: Nature Publishing Group.

[13] Brad K Hulse, Hannah Haberkern, Romain Franconville, Daniel Turner-Evans, Shin-ya Takemura, Tanya Wolff, Marcella Noorman, Marisa Dreher, Chuntao Dan, Ruchi Parekh, Ann M Hermundstad, Gerald M Rubin, and Vivek Jayaraman. A connectome of the Drosophila central complex reveals network motifs suitable for flexible navigation and context-dependent action selection. *eLife*, 10:e66039, October 2021. Publisher: eLife Sciences Publications, Ltd.

[14] Claire Eschbach, Akira Fushiki, Michael Winding, Casey M. Schneider-Mizell, Mei Shao, Rebecca Arruda, Katharina Eichler, Javier Valdes-Aleman, Tomoko Ohyama, Andreas S. Thum, Bertram Gerber, Richard D. Fetter, James W. Truman, Ashok Litwin-Kumar, Albert Cardona, and Marta Zlatic. Recurrent architecture for adaptive regulation of learning in the insect brain. *Nature Neuroscience*, 23(4):544–555, April 2020. Number: 4 Publisher: Nature Publishing Group.

[15] Arie Matsliah, Amy R Sterling, Sven Dorkenwald, Kai Kuehner, Ryan Morey, H Sebastian Seung, and Mala Murthy. Codex: Connectome Data Explorer. 2023. Publisher: Unpublished.

[16] Arthur Zhao, Aljoscha Nern, Sanna Koskela, Marisa Dreher, Mert Erginkaya, Connor W. Laughland, Henrique Ludwigh, Alex Thomson, Judith Hoeller, Ruchi Parekh, Sandro Romani, Davi D. Bock, Eugenia Chiappe, and Michael B. Reiser. A comprehensive neuroanatomical survey of the Drosophila Lobula Plate Tangential Neurons with predictions for their optic flow sensitivity. *eLife*, 13, January 2024. Publisher: eLife Sciences Publications Limited.

[17] Anders Enjin, Emanuela E. Zaharieva, Dominic D. Frank, Suzan Mansourian, Greg S. B. Suh, Marco Gallio, and Marcus C. Stensmyr. Humidity Sensing in *Drosophila*. *Current Biology*, 26(10):1352–1358, May 2016.

[18] Dominic D. Frank, Anders Enjin, Genevieve C. Jouandet, Emanuela E. Zaharieva, Alessia Para, Marcus C. Stensmyr, and Marco Gallio. Early Integration of Temperature and Humidity Stimuli in the *Drosophila* Brain. *Current Biology*, 27(15):2381–2388.e4, August 2017.

[19] Helen H. Yang, Bella E. Brezovec, Laia Serratosa Capdevila, Quinn X. Vanderbeck, Atsuko Adachi, Richard S. Mann, and Rachel I. Wilson. Fine-grained descending control of steering in walking *Drosophila*. *Cell*, 187(22):6290–6308.e27, October 2024.

[20] Eriko Matsuo, Daichi Yamada, Yuki Ishikawa, Tomonori Asai, Hiroshi Ishimoto, and Azusa Kamikouchi. Identification of novel vibration- and deflection-sensitive neuronal subgroups in Johnston's organ of the fruit fly. *Frontiers in Physiology*, 5, May 2014. Publisher: Frontiers.

[21] Benjamin Gorko, Igor Siwanowicz, Kari Close, Christina Christoforou, Karen L. Hibbard, Mayank Kabra, Allen Lee, Jin-Yong Park, Si Ying Li, Alex B. Chen, Shigehiro Namiki, Chenghao Chen, John C. Tuthill, Davi D. Bock, Hervé Rouault, Kristin Branson, Gudrun Ihrke, and Stephen J. Huston. Motor neurons generate pose-targeted movements via proprioceptive sculpting. *Nature*, 628(8008):596–603, April 2024. Publisher: Nature Publishing Group.

[22] Azusa Kamikouchi, Hidehiko K. Inagaki, Thomas Effertz, Oliver Hendrich, André Fiala, Martin C. Göpfert, and Kei Ito. The neural basis of Drosophila gravity-sensing and hearing. *Nature*, 458(7235):165–171, March 2009. Publisher: Nature Publishing Group.

[23] Akira Mamiya and Michael H. Dickinson. Antennal Mechanosensory Neurons Mediate Wing Motor Reflexes in Flying Drosophila. *Journal of Neuroscience*, 35(20):7977–7991, May 2015. Publisher: Society for Neuroscience Section: Articles.

[24] Adeline Pézier, Sami H. Jezzini, Bruno Marie, and Jonathan M. Blagburn. Engrailed Alters the Specificity of Synaptic Connections of Drosophila Auditory Neurons with the Giant Fiber. *Journal of Neuroscience*, 34(35):11691–11704, August 2014. Publisher: Society for Neuroscience Section: Articles.

[25] Mark Dombrovski, Martin Y. Peek, Jin-Yong Park, Andrea Vaccari, Marissa Sumathipala, Carmen Morrow, Patrick Breads, Arthur Zhao, Yerbol Z. Kurmangaliyev, Piero Sanfilippo, Aadil Rehan, Jason Polsky, Shada Alghailani, Emily Tenshaw, Shigehiro Namiki, S. Lawrence Zipursky, and Gwyneth M. Card. Synaptic gradients transform object location to action. *Nature*, 613(7944):534–542, January 2023. Publisher: Nature Publishing Group.

[26] Fei Wang, Kaiyu Wang, Nora Forknall, Christopher Patrick, Tansy Yang, Ruchi Parekh, Davi Bock, and Barry J. Dickson. Neural circuitry linking mating and egg laying in Drosophila females. *Nature*, 579(7797):101–105, March 2020. Publisher: Nature Publishing Group.

[27] Carlotta Martelli, Ulrike Pech, Simon Kobbenbring, Dennis Pauls, Britta Bahl, Mirjam Vanessa Sommer, Atefeh Pooryasin, Jonas Barth, Carmina Warth Perez Arias, Chrystalleni Vassiliou, Abud Jose Farca Luna, Haiko Poppinga, Florian Gerhard Richter, Christian Wegener, André Fiala, and Thomas Riemensperger. SIFamide Translates Hunger Signals into Appetitive and Feeding Behavior in Drosophila. *Cell Reports*, 20(2):464–478, July 2017. Publisher: Elsevier.

[28] Selim Terhzaz, Philippe Rosay, Stephen F. Goodwin, and Jan A. Veenstra. The neuropeptide SIFamide modulates sexual behavior in *Drosophila*. *Biochemical and Biophysical Research Communications*, 352(2):305–310, January 2007.

[29] Aubrey Manning. The control of sexual receptivity in female *Drosophila*. *Animal Behaviour*, 15(2):239–250, April 1967.

[30] Nathan C. Klapoetke, Aljoscha Nern, Edward M. Rogers, Gerald M. Rubin, Michael B. Reiser, and Gwyneth M. Card. A functionally ordered visual feature map in the Drosophila brain. *Neuron*, 110(10):1700–1711.e6, May 2022. Publisher: Elsevier.

[31] Bettina Schnell, Ivo G. Ros, and Michael H. Dickinson. A Descending Neuron Correlated with the Rapid Steering Maneuvers of Flying Drosophila. *Current Biology*, 27(8):1200–1205, April 2017. Publisher: Elsevier.

[32] H. S. J. Cheong, K. Eichler, T. Stürner, S. K. Asinof, A. S. Champion, E. C. Marin, T. B. Oram, M. Sumathipala, L. Venkatasubramanian, S. Namiki, I. Siwanowicz, M. Costa, S. Berg, Janelia FlyEM Project Team, G. S. X. E. Jefferis, and G. M. Card. Transforming descending input into motor

output: An analysis of the Drosophila Male Adult Nerve Cord connectome, June 2025. Pages: 2023.06.07.543976 Section: New Results.

[33] Matthew F. Collie, Chennan Jin, Emily Kellogg, Quinn X. Vanderbeck, Alexandra K. Hartman, Stephen L. Holtz, and Rachel I. Wilson. Specialized parallel pathways for adaptive control of visual object pursuit, April 2025. Pages: 2025.04.23.650240 Section: New Results.

[34] Shawn R. Olsen, Vikas Bhandawat, and Rachel I. Wilson. Divisive Normalization in Olfactory Population Codes. *Neuron*, 66(2):287–299, April 2010. Publisher: Elsevier.

[35] Dumitru Erhan, Y. Bengio, Aaron Courville, and Pascal Vincent. Visualizing Higher-Layer Features of a Deep Network. *Technical Report, Univeristé de Montréal*, January 2009.

[36] Chris Olah, Alexander Mordvintsev, and Ludwig Schubert. Feature Visualization. *Distill*, 2(11):e7, November 2017.

[37] Jan M. Ache, Jason Polsky, Shada Alghailani, Ruchi Parekh, Patrick Breads, Martin Y. Peek, Davi D. Bock, Catherine R. von Reyn, and Gwyneth M. Card. Neural Basis for Looming Size and Velocity Encoding in the *Drosophila* Giant Fiber Escape Pathway. *Current Biology*, 29(6):1073–1081.e4, March 2019.

[38] Aric A. Hagberg, Daniel A. Schult, and Pieter J. Swart. Exploring Network Structure, Dynamics, and Function using NetworkX. pages 11–15, Pasadena, California, June 2008.

[39] Christian L. Staudt, Aleksejs Sazonovs, and Henning Meyerhenke. NetworKit: A Tool Suite for Large-scale Complex Network Analysis, November 2015. arXiv:1403.3005 [cs].

[40] Arie Matsliah, Szi-chieh Yu, Krzysztof Kruk, Doug Bland, Austin T. Burke, Jay Gager, James Hebditch, Ben Silverman, Kyle Patrick Willie, Ryan Willie, Marissa Sorek, Amy R. Sterling, Emil Kind, Dustin Garner, Gizem Sancer, Mathias F. Wernet, Sung Soo Kim, Mala Murthy, and H. Sebastian Seung. Neuronal parts list and wiring diagram for a visual system. *Nature*, 634(8032):166–180, October 2024. Publisher: Nature Publishing Group.

[41] I. A. Meinertzhagen and S. D. O'Neil. Synaptic organization of columnar elements in the lamina of the wild type in Drosophila melanogaster. *Journal of Comparative Neurology*, 305(2):232–263, 1991. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/cne.903050206.

[42] Marta Rivera-Alba, Shiv N. Vitaladevuni, Yuriy Mischenko, Zhiyuan Lu, Shin-ya Takemura, Lou Scheffer, Ian A. Meinertzhagen, Dmitri B. Chklovskii, and Gonzalo G. de Polavieja. Wiring economy and volume exclusion determine neuronal placement in the Drosophila brain. *Current biology : CB*, 21(23):2000–2005, December 2011.

[43] Philipp Schlegel, Alexander Shakeel Bates, Tomke Stürner, Sridhar R Jagannathan, Nikolas Drummond, Joseph Hsu, Laia Serratosa Capdevila, Alexandre Javier, Elizabeth C Marin, Asa Barth-Maron, Imaan FM Tamimi, Feng Li, Gerald M Rubin, Stephen M Plaza, Marta Costa, and Gregory S X E Jefferis. Information flow, cell types and stereotypy in a full olfactory connectome. *eLife*, 10:e66018, May 2021. Publisher: eLife Sciences Publications, Ltd.

[44] Albert Lin, Runzhe Yang, Sven Dorkenwald, Arie Matsliah, Amy R. Sterling, Philipp Schlegel, Szi-chieh Yu, Claire E. McKellar, Marta Costa, Katharina Eichler, Alexander Shakeel Bates, Nils Eckstein, Jan Funke, Gregory S. X. E. Jefferis, and Mala Murthy. Network statistics of the whole-brain connectome of Drosophila. *Nature*, 634(8032):153–165, October 2024. Publisher: Nature Publishing Group.

[45] Juergen Haag and Alexander Borst. Encoding of Visual Motion Information and Reliability in Spiking and Graded Potential Neurons. *Journal of Neuroscience*, 17(12):4809–4819, June 1997. Publisher: Society for Neuroscience Section: Articles.

[46] Vikas Bhandawat, Shawn R. Olsen, Nathan W. Gouwens, Michelle L. Schlief, and Rachel I. Wilson. Sensory Processing in the Drosophila Antennal Lobe Increases the Reliability and Separability of Ensemble Odor Representations. *Nature neuroscience*, 10(11):1474–1482, November 2007.

[47] Stephen M. Plaza, Jody Clements, Tom Dolafi, Lowell Umayam, Nicole N. Neubarth, Louis K. Scheffer, and Stuart Berg. neuPrint: An open access tool for EM connectomics. *Frontiers in Neuroinformatics*, 16, July 2022. Publisher: Frontiers.

[48] Stephan Saalfeld, Albert Cardona, Volker Hartenstein, and Pavel Tomančák. CATMAID: collaborative annotation toolkit for massive amounts of image data. *Bioinformatics*, 25(15):1984–1986, August 2009.

[49] H. Sebastian Seung. Interneuron diversity and normalization specificity in a visual system, April 2024. Pages: 2024.04.03.587837 Section: New Results.

[50] Philipp Schlegel, Chris Barnes, Frank Loesche, Andrew Champion, dokato, Sridhar Jagannathan, Rob Court, Jinhan Choi, Forrest Collman, Stuart Berg, Ben Pedigo, Guspan Tanadi, Peter Newstein, Yervand Azatian, and Antonio. navis-org/navis: Version 1.10.0, February 2025.

[51] Arthur Zhao, Eyal Gruntman, Aljoscha Nern, Nirmala Iyer, Edward M. Rogers, Sanna Koskela, Igor Siwanowicz, Marisa Dreher, Miriam A. Flynn, Connor Laughland, Henrique Ludwig, Alexander Thomson, Cullen Moran, Bruck Gezahegn, Davi D. Bock, and Michael B. Reiser. Eye structure shapes neuron function in Drosophila motion vision. *Nature*, pages 1–8, July 2025. Publisher: Nature Publishing Group.

[52] Elissa A. Hallem and John R. Carlson. Coding of Odors by a Receptor Repertoire. *Cell*, 125(1):143–160, April 2006. Publisher: Elsevier.

[53] Daniel Münch and C. Giovanni Galizia. DoOR 2.0 - Comprehensive Mapping of Drosophila melanogaster Odorant Responses. *Scientific Reports*, 6(1):21841, February 2016. Number: 1 Publisher: Nature Publishing Group.

[54] Hany K. M. Dweck, Shimaa A. M. Ebrahim, Tom Retzke, Veit Grabe, Jerrit Weißflog, Ales Svatoš, Bill S. Hansson, and Markus Knaden. The Olfactory Logic behind Fruit Odor Preferences in Larval and Adult Drosophila. *Cell Reports*, 23(8):2524–2531, May 2018. Publisher: Elsevier.

[55] Laurent Badel, Kazumi Ohta, Yoshiko Tsuchimoto, and Hokto Kazama. Decoding of Context-Dependent Olfactory Behavior in Drosophila. *Neuron*, 91(1):155–167, July 2016. Publisher: Elsevier.