1

Covert spatial attention is uniform across cardinal meridians despite differential adaptation

Hsing-Hao Lee[1*] & Marisa Carrasco[1,2]

[1] Department of Psychology, New York University, New York, NY, USA
[2] Center for Neural Sciences, New York University, New York, NY, USA

* Hsing-Hao Lee
Email: hsinghaolee@nyu.edu

Competing Interest Statement: The authors declare no competing interests.

Abstract

Visual adaptation and attention are two processes that help manage the brain's limited bioenergetic resources for perception. Visual perception is heterogeneous around the visual field: it is better along the horizontal than the vertical meridian (horizontal-vertical anisotropy, HVA), and better along the lower than the upper vertical meridian (vertical meridian asymmetry, VMA). Recently, we showed that visual adaptation is more pronounced at the horizontal than the vertical meridian, but whether and how this differential adaptation modulates the effects of covert spatial attention remains unknown. In this study, we investigated whether and how the effects of endogenous (voluntary) and exogenous (involuntary) covert attention on an orientation discrimination task vary at the cardinal meridians, with and without adaptation. We manipulated endogenous (Experiment 1) or exogenous (Experiment 2) attention via an informative central or uninformative peripheral cue, respectively. Results showed that (1) in the non-adapted condition, the typical HVA and VMA emerged in contrast thresholds; (2) the adaptation effect was stronger at the horizontal than the vertical meridian; and (3) regardless of adaptation, both endogenous and exogenous attention enhanced and impaired performance at the attended and unattended locations, respectively, to a similar degree at both cardinal meridians. Together, these findings reveal that, despite differences between endogenous and exogenous attention, their effects remain uniform across cardinal meridians–even under differential adaptation that reduces intrinsic asymmetries of visual field representations.

Keywords: visual adaptation, endogenous attention, exogenous attention, visual performance asymmetries, contrast sensitivity

Introduction

Visual adaptation and attention are two processes that optimize performance and help manage the brain's limited bioenergetic resources by allocating them according to task demands (Carrasco, 2011; Lee, Fernández, & Carrasco, 2024; Lennie, 2003; Pestilli, Viera, & Carrasco, 2007). Although both processes modulate sensory responses, they have opposite effects on the contrast response function. Visual adaptation helps manage bioenergetic resources by increasing metabolic efficiency–it reduces sensitivity to repeated features and enhances sensitivity to novel ones. For example, contrast adaptation can adjust the gain of the neural response so that its dynamic range is matched to the range of levels in the stimulus (Boynton & Finney, 2003; Gardner et al., 2005; Kohn, 2007; Perini, Cattaneo, Carrasco, & Schwarzbach, 2012; Vergeer, Mesik, Baek, Wilmerding, & Engel, 2018; Webster, 2011, 2015). In contrast, visual attention selectively improves information processing at an attended location while impairing processing elsewhere – a ubiquitous performance tradeoff considered a push-pull mechanism (e.g., Dosher & Lu, 2000b; Ling & Carrasco, 2006a; Pestilli & Carrasco, 2005; Pestilli, Ling, & Carrasco, 2009; Pestilli et al., 2007; for reviews, Carrasco, 2011, 2014; Desimone & Duncan, 1995; Olivers, 2025).

There are two types of covert spatial attention: endogenous and exogenous. Endogenous attention is voluntary, goal-driven, and flexible; exogenous attention is involuntary, stimulus-

driven, and automatic. Endogenous attention takes ~300 ms to be deployed and can be sustained
for many seconds, whereas exogenous attention peaks at ~120 ms and is transient (reviews:
Carrasco, 2011, 2014). Despite these differences, both types of attention improve performance
in many visual tasks, e.g., contrast sensitivity (e.g., Herrmann, Montaser-Kouhsari, Carrasco, &
Heeger, 2010; Pestilli et al., 2009), appearance (review: Carrasco & Barbot, 2019), and
orientation discrimination (e.g., Fernández, Okun, & Carrasco, 2022). However, they have distinct
effects in other tasks, e.g., texture segregation (e.g., Barbot & Carrasco, 2017; Jigo, Heeger, &
Carrasco, 2021; Yeshurun & Carrasco, 1998), and alter sensitivity across a different spatial
frequency range (Fernández et al., 2022; Jigo & Carrasco, 2020).

Exogenous attention restores contrast sensitivity after adaptation; although adaptation reduces
sensitivity, the magnitude of the exogenous attentional benefit at the attended location and its
concurrent cost at the unattended location remain comparable to those observed without
adaptation (Lee et al., 2024; Pestilli et al., 2007). However, whether and how endogenous
attention operates after adaptation is unknown. Thus, our first goal was to examine whether
endogenous attention restores contrast sensitivity after adaptation. It is possible that after
adaptation endogenous attention (1) enhances contrast sensitivity to a similar extent as without
adaptation, assuming that similar to exogenous attention (Lee et al., 2024; Pestilli et al., 2007),
endogenous attention and adaptation yield independent effects on contrast sensitivity (**Figure 1A**;
Hypothesis 1); (2) enhances sensitivity more than before adaptation, reflecting a compensatory
process given the flexible nature of endogenous attention, which optimizes performance as a
function of task demands (Barbot & Carrasco, 2017; Barbot, Landy, & Carrasco, 2012; Giordano,
McElree, & Carrasco, 2009; Hein, Rolke, & Ulrich, 2006; Yeshurun, Montagna, & Carrasco, 2008);
it may help more than without adaptation, as after decreases there is more room for improvement
(**Figure 1B**; Hypothesis 2); or (3) enhances sensitivity less than without adaptation; if reduced
baseline sensitivity limits the push-pull effects of endogenous attention (**Figure 1C**; Hypothesis
3).

Because endogenous attention is flexible (Barbot & Carrasco, 2017; Barbot et al., 2012; Giordano
et al., 2009; Hein et al., 2006; Yeshurun et al., 2008), but exogenous attention is not (Barbot et
al., 2012; Carrasco, Loula, & Ho, 2006; Crotty, Massa, Tellez, White, & Grubb, 2025; Giordano et
al., 2009; Hein et al., 2006; Luck & Thomas, 1999; Yantis & Jonides, 1996; Yeshurun & Carrasco,
1998), and because distinct brain regions are critical for their effect –right frontal eye fields for
endogenous attention (Fernández, Hanning, & Carrasco, 2023), and early visual cortex for
exogenous attention (Fernández & Carrasco, 2020; Lee et al., 2024), where it interacts with
adaptation (Lee et al., 2024) –it is possible that they exert different effects on contrast sensitivity
after adaptation. Therefore, our second goal was to determine whether endogenous and
exogenous attention have similar or different effects on contrast sensitivity following adaptation.

Finally, we investigated whether target location matters. In adult humans, visual performance is
better at the horizontal than the vertical meridian (horizontal-vertical anisotropy, HVA), and better
at the lower than the upper vertical meridian (vertical meridian asymmetry, VMA). These visual
field asymmetries, known as performance fields, are present in many fundamental visual tasks,
including contrast sensitivity (Abrams, Nizam, & Carrasco, 2012; Baldwin, Meese, & Baker, 2012;
Cameron, Tai, & Carrasco, 2002; Carrasco, Talgar, & Cameron, 2001; Corbett & Carrasco, 2011;
Fuller, Rodriguez, & Carrasco, 2008; Himmelberg, Winawer, & Carrasco, 2020; Lee & Carrasco,

2025; Purokayastha, Roberts, & Carrasco, 2021), visual acuity (Kwak, Hanning, & Carrasco, 2023;
Montaser-Kouhsari & Carrasco, 2009), spatial resolution (Altpeter, Mackeben, & Trauzettel-
Klosinski, 2000; Carrasco, Williams, & Yeshurun, 2002; Greenwood, Szinte, Sayim, & Cavanagh,
2017; Talgar & Carrasco, 2002), and motion (Fuller & Carrasco, 2009; Tünçok, Kiorpes, &
Carrasco, 2025), as well as mid-level visual processes, such as texture segregation (Barbot, Xue,
& Carrasco, 2021; Greenwood et al., 2017; Talgar & Carrasco, 2002; Z. Wang, Murai, & Whitney,
2020) and crowding (Greenwood et al., 2017; Kurzawski et al., 2023; Petrov & Meleshkevich,
2011), and high-level tasks, such as numerosity perception (Chakravarthi, Papadaki, & Krajnik,
2022), face perception (Afraz, Pashkam, & Cavanagh, 2010; Kim & Chong, 2024), word
identification (Tsai, Liao, Hou, Jang, & Chen, 2024), and visual short-term memory (Montaser-
Kouhsari & Carrasco, 2009).

These visual field asymmetries are resistant to endogenous (Purokayastha et al., 2021; Tünçok,
Carrasco, & Winawer, 2025) and exogenous attention (Cameron et al., 2002; Carrasco et al.,
2001; Roberts, Ashinoff, Castellanos, & Carrasco, 2018; Roberts, Cymerman, Smith, Kiorpes, &
Carrasco, 2016), as well as to temporal (Fernández, Denison, & Carrasco, 2019) attention. Thus,
performance fields are not easily reshaped. On the contrary, presaccadic attention, which
enhances the processing at the location of the impending saccade target, exacerbates
performance asymmetries at the cardinal locations by enhancing contrast sensitivity the most at
the horizontal meridian and the least at the upper vertical meridian (Hanning, Himmelberg, &
Carrasco, 2022, 2024; Kwak, Hanning, & Carrasco, in press; Kwak, Zhao, Lu, Hanning, &
Carrasco, 2024).

A recent study showed that visual adaptation is stronger at the horizontal than the vertical
meridian, leading to more homogeneous perception by mitigating the HVA (Lee & Carrasco, 2025).
It remains unknown, however, whether and how endogenous and exogenous attention reshape
performance fields after such differential adaptation. Thus, our third goal was to investigate
whether, following adaptation, covert spatial attention enhances contrast sensitivity (1) to the
same extent at the cardinal meridians around polar angle, similar to without adaptation (e.g.,
Carrasco et al., 2001; Purokayastha et al., 2021; Roberts et al., 2018; Roberts et al., 2016; Tünçok,
Carrasco, & Winawer, 2025) (**Figure 1D**; Hypothesis 4), (2) more at the vertical than the horizontal
meridian, and more at the upper than the lower vertical meridian, acting as a compensatory
mechanism to reduces asymmetries (**Figure 1E**; Hypothesis 5), or (3) more where baseline
performance is already better (i.e., the horizontal meridian) than where it is worse (i.e., vertical
meridian, especially the upper vertical meridian), thereby exaggerating asymmetries (**Figure 1F**;
Hypothesis 6).

Both adaptation (Altan, Morgan, Dakin, & Schwarzkopf, 2025; Dao, Lu, & Dosher, 2006; Gardner
et al., 2005; Perini et al., 2012; Pestilli et al., 2007) and endogenous attention (Dosher & Lu,
2000a; Ling & Carrasco, 2006a; Lu, Lesmes, & Dosher, 2002; Pestilli et al., 2009) primarily affect
the contrast gain of the contrast response function (**Figure 1A**), i.e., a shift in threshold, whereas
exogenous attention primarily affects response gain (Fernández & Carrasco, 2020; Pestilli et al.,
2009), i.e., a shift in asymptote. Additionally, according to a prominent normalization model of
attention (Reynolds & Heeger, 2009), exogenous attention can also affect contrast gain when the
attentional window is wider than the stimulus size, and endogenous attention can also affect
response gain when the attentional window is narrower than the stimulus size (Herrmann et al.,

2010). In this study, to directly compare the two types of attention before and after adaptation at
the cardinal meridians, we induced a larger attentional window in the exogenous attention
experiment, enabling contrast gain effects predicted by Reynolds and Heeger's (2009)
normalization model of attention.

Figure 1. (A-C) Hypotheses regarding effects on contrast sensitivity. (A) Hypothesis 1: Attentional effect is
comparable with and without adaptation. The c_{50} and c_{50}-adapted indicate the contrast threshold derived
from the titration procedures in the non-adapted and adapted conditions, respectively. (B) Hypothesis 2:
Attentional effect is larger with than without adaptation. (C) Hypothesis 3: Attentional effect is smaller than
without adaptation. (D-F) Hypotheses regarding effects on contrast sensitivity as a function of location at
the cardinal locations. (D) Hypothesis 4: Attentional effect is comparable around polar angle after adaptation.
(E) Hypothesis 5: Attentional effect is stronger at the vertical than horizontal meridian. (F) Hypothesis 6:
Attentional effect is smaller at the vertical than horizontal meridian.

In summary, we asked (1) whether and how endogenous attention restores contrast sensitivity
following adaptation, (2) whether endogenous and exogenous attention have similar or distinct
effects on contrast sensitivity before and after adaptation, and (3) whether these effects uniformly
or differentially across the cardinal meridians around the visual field. These findings are essential
for elucidating how the visual system engages adaptation and attention—two fundamental visual
processes that manage limited bioenergetic resources—to optimize performance across locations
that differ in intrinsic discriminability and in their corresponding representation in cortical surface
area.

Experiment 1 – Endogenous attention

Methods

Participants

Twelve adults (5 females, age range: 24-36 years old), including author HHL, participated in the
experiment. All of them had normal or corrected-to-normal vision. Sample size was based on
previous studies on adaptation (Lee et al., 2024), with an effect size of d=1.3, and on performance
fields (Lee & Carrasco, 2025), with an effect size of d=1.41 for performance in the neutral trials.
According to G*Power 3.0 (Faul, Erdfelder, Lang, & Buchner, 2007), we would need 9 participants
for adaptation and 8 participants for performance fields to reach a power=0.9. We also estimated
the required sample size for the interaction between adaptation and location, based on a recent
study between adaptation and performance fields (Lee & Carrasco, 2025) (η_p^2=0.34), by
assuming SD=1, we would need 10 subjects to reach a power=0.9 according to the Monte-Carlo
simulation (1,000 iterations per possible subject number). The Institutional Review Board at New
York University approved the experimental procedures, and all participants provided informed
consent before they started the experiment.

Stimuli and apparatus

The target Gabor (diameter = 4°, 5 cpd, 1.25° full-width at half maximum) was presented on the
left, right, upper and lower cardinal meridian locations (8° from the center to center). There were
four placeholders (length = 0.16°, width = 0.06°) 0.5° away from the Gabor's edge. The fixation
cross consisted of a plus sign (length = 0.25°; width = 0.06°) at the center of the screen. The
endogenous attentional cue (length = 0.75°; width = 0.2°) was presented at the center.

Participants were in a dimly lit, sound-attenuated room, with their head placed on a chinrest 57
cm away from the monitor. All stimuli were generated using MATLAB (MathWorks, MA, USA) and
the Psychophysics Toolbox (Brainard, 1997; Pelli, 1997) on a gamma-corrected 20-inch
ViewSonic G220fb CRT monitor with a spatial resolution of 1,280 x 960 pixels and a refresh rate
of 100 Hz. To ensure fixation, participants' eye movements were recorded using EYELINK 1000
(SR Research, Osgoode, Ontario, Canada) with a sample rate of 1,000 Hz.

Experimental design and procedures

Figure 2 shows the procedure of titration and the endogenous attention task. In the adapted
condition, at the beginning of each block, participants adapted to a vertical 5-cpd Gabor patch

flickering at 7.5 Hz in a counterphase manner, presented at the target location for 60 seconds.
Each trial started with a 2s top-up phase to ensure a continuous adaptation effect throughout the
block. In the non-adaptation condition, participants maintained fixation at the center for 4s (without
Gabor) at the beginning of each block and for 2s at the beginning of each trial.

221
222

223
Figure 2. Experimental procedure: Participants performed either adaptation or nonadaptation blocks, each
in separate experimental sessions. The target Gabor stimulus was always presented within the black
placeholder, and target meridians were blocked. The target, two vertical Gabor stimuli were presented 8°
away from the center (e.g., at the vertical meridian here; at the horizontal meridian in a different block).
Participants were instructed to respond whether the Gabor was tilted clockwise or counterclockwise from
vertical. The pre-cue either matches (valid condition), mismatches (invalid condition) the response cue, or
does not provide location information (neutral condition). For illustration purposes, the stimulus size and
spatial frequency shown here are not to scale.
232
After the top-up, there was a 200 ms ISI before an endogenous pre-cue was presented for 100
ms. Following a 200-ms ISI the tilted Gabor was then presented for 67 ms, followed by another
200-ms ISI and then the response cue. In a valid trial, the location indicated by the response cue
matches the precue; in an invalid trial, they mismatch; in a neutral cue condition, the pre-cue
points at both locations. Participants had to judge whether the target Gabor was tilted clockwise
or counterclockwise off vertical. The tilt angle was 2.5°, based on pilot data and our previous study
(Lee & Carrasco, 2025), to ensure an adaptation effect while avoiding floor or ceiling performance.
240
A feedback tone was presented when participants gave an incorrect response. The target
locations were blocked in a horizontal block or a vertical block, where the target locations were
presented at the horizontal or vertical meridians, respectively. Participants were asked to respond

as accurately as possible while fixating at the center of the screen throughout the trial. A trial
would be interrupted and repeated at the end of the block if participants' eyes position deviated
≥1.5° from the center, from the pre-cue onset until the response cue onset.
247
Participants completed the adapted and non-adapted attentional task on the vertical and the
horizontal meridian on different days, with a counterbalanced order. The order of horizontal and
vertical meridian blocks was randomized, and the adaptation and non-adaptation titration were
implemented on different days, with a counterbalanced order. There were 4 independent
staircases for each adaptation condition and location, varying Gabor contrast from 2% to 85% to
reach ~75% accuracy for the orientation discrimination task. Each staircase started from 4
different points (85%, 2%, the median contrast of 43.5%, and a random point between 2% and
85%) and contained 48 trials. Four blocks (192 trials per location for each adaptation and non-
adaptation conditions) were conducted consecutively for the horizontal meridian block or the
vertical meridian block. The contrast threshold was derived using an adaptive staircase procedure
using the Palamedes toolbox (Prins & Kingdom, 2018), as in previous studies (e.g., Fernández &
Carrasco, 2020; Hanning et al., 2022; Jigo & Carrasco, 2018; Lee & Carrasco, 2025; Lee et al.,
2024) and averaging the last 8 trials. The Gabors were always preceded by a neutral pre-cue,
which, as in many studies (e.g., Dosher & Lu, 2000b; Fernández et al., 2022; Huang, Liao, Chen,
& Chen, 2025; Jigo & Carrasco, 2020; Li, Pan, & Carrasco, 2021; Luzardo & Yeshurun, 2025;
Palmieri & Carrasco, 2024; Ramamurthy, White, & Yeatman, 2024; Tünçok, Carrasco, & Winawer,
2025), provided the same temporal information as the valid and invalid cues, but no information
about the spatial location.
266
267 In this endogenous attention task, for each adapted and non-adapted condition, 20% of the trials
268 had a neutral cue, which pointed at both locations; 80% of the trials had an attentional cue pointing
269 toward a location, 75% among them were valid cues, and the other 25% were invalid cues. All
270 participants completed a practice session to familiarize themselves with the task procedure.
271
Psychometric function fitting
We fitted a Weibull function for the accuracy as a function of contrast threshold. For each location
and adaptation condition, a logistic function was fit to the data using maximum likelihood
estimation using the fmincon function in MATLAB. The results derived from the psychometric
function estimation positively correlated (*ps*<.01) with the staircase results in all experiments,
verifying our procedure in all conditions.
278
Behavioral data analyses
Behavioral data analyses were performed using R (Team, 2000). A three-way repeated-measures
analysis of variance (ANOVA) on d' was conducted on the factors of location (horizontal meridian,
upper, lower), adaptation (adapted, non-adapted), and attention (valid, neutral, invalid) conditions
to assess statistical significance. Repeated-measures ANOVA along with effect size (η^2) were
computed in R and used to assess statistical significance.
285
286
287

288 **Results**
289
290 **Adaptation effect varied around polar angle**
291
After deriving the c_{50} contrast for the horizontal meridian (HM), upper, and lower vertical meridians
for both the adapted and non-adapted conditions, we conducted a two-way ANOVA on contrast
thresholds (**Figure 3**). This analysis showed a main effect of location [$F(2,22)=7.89$, $p=.003$,
$\eta_p^2=0.42$] and a higher threshold in the adapted than non-adapted conditions [$F(1,11)=18.44$,
$p=.001$, $\eta_p^2=0.63$], and an interaction [$F(2,22)=3.58$, $p=.045$, $\eta_p^2=0.25$], indicating that the
adaptation effect varied across locations.
298
299 We confirmed that the HVA and VMA emerged in the non-adaptation condition (**Figure 3**):
300 Contrast thresholds were lower along the horizontal than the vertical meridian [$t(11)=5.87$, $p<.001$,
301 $d=1.69$) and lower at the lower than upper vertical meridian [$t(11)=2.37$, $p=.037$, $d=0.68$].
302
Next, we assessed the adaptation effect at the horizontal and vertical meridians. The normalized
adaptation effect (calculated as the difference between adapted and non-adapted thresholds
divided by the sum of the thresholds, as in Lee and Carrasco (2025) was stronger at the horizontal
than the vertical meridian [$t(11)=3.39$, $p=.006$, $d=0.98$] (**Figure 3,** see gaps between adapt and
non-adapt conditions for different locations), but no significant difference between the upper and
lower vertical meridian [$t(11)<1$].
309

310
311 **Figure 3**. The contrast thresholds for different locations and adaptation conditions. The thresholds were
312 higher in the vertical than horizontal meridian (HM), and higher in the upper than lower vertical meridian.
313 The thresholds were also higher in the adapted than non-adapted conditions. Critically, the adaptation effect
314 was stronger in the horizontal than vertical meridian. The error bars indicate ±1 SEM.

315 **Endogenous attentional effect**
316
Figure 4 shows the results. We compared the endogenous attentional effect on d′ by conducting
a three-way ANOVA on the factors of location (HM, upper, lower), attentional validity (valid,
neutral, invalid), and adaptation (adaptation, non-adaptation). Given that we titrated the contrast
thresholds across locations and adaptation conditions, we expected no main effects of either
adaptation or location. Indeed, there was a main effect of attention [$F(2,22)=53.18$, $p<.001$,
$\eta_p^2=0.83$], but neither of location [$F(2,22)<1$], nor of adaptation [$F(1,11)<1$]. There was neither a
3-way interaction nor 2-way interactions [all $ps>.1$].
324
325

326
Figure 4. Performance in Experiment 1. (A) d′ was higher in the valid followed by neutral and invalid
conditions in both non-adapted and adapted conditions. There was no difference between the adapted and
non-adapted conditions. (B) The attentional effects were similar around polar angle –horizontal meridian
(HM), and upper and lower vertical meridian– and were comparable in the adapted and non-adapted
conditions. The error bars above the bar plots indicate ±1 SEM of the difference between conditions. ***
$p<.001$, n.s. $p>.05$.

The results were further confirmed by separating the adapted and non-adapted conditions into
two 2-way ANOVAs on attention and location. For the non-adapted condition, we observed a main
effect of attention [$F(2,22)=46.74$, $p<.001$, $\eta_p^2=0.81$] but not of location [$F(2,22)<1$] or an
interaction [$F(4,44)=1.68$, $p>.1$]. The same pattern emerged for the adapted condition: a main
effect of attention [$F(2,22)=38.59$, $p<.001$, $\eta_p^2=0.78$] but not of location [$F(2,22)<1$] or an
interaction [$F(4,44)=1.48$, $p>.1$]. Thus, neither adaptation state nor location modulated the
pronounced overall effect of attention.
340
341 We plot the individual data for the endogenous attentional effect (valid d' − invalid d') in the
342 adapted and non-adapted conditions (**Figure 5**). There was no difference between the two
343 conditions [$t(11)=1.27$, $p>.1$].
344
345 In sum, the endogenous attentional effect was comparable across locations and adaptation
346 conditions.
347

348
349 **Figure 5.** Comparison of the endogenous attentional effects (valid d' − invalid d') in Experiment 1. The
350 attentional effects were comparable in the adapted and non-adapted conditions. The red circle indicates
351 the mean of all participants, and the error bars indicate ±1 SEM.
352
353

Experiment 2 — Exogenous attention

355
Experiment 1 shows that endogenous attention does not reshape the performance fields, even
after differential adaptation effects across meridians. In Experiment 2, we examined whether
exogenous attention exhibits a similar or distinct pattern as endogenous attention, given their well-
established differences in temporal dynamics: Whereas endogenous attention takes about 300
ms to deploy and its effects can be sustained for many seconds, exogenous attention effects peak

at about 120 ms and its effects are transient (reviews: Carrasco, 2011, 2014; Carrasco & Barbot,
2014). Moreover, endogenous attention is flexible whereas exogenous attention is not (e.g.,
Barbot & Carrasco, 2017; Barbot et al., 2012; Giordano et al., 2009; Hein et al., 2006; Nakayama
& Mackeben, 1989; Yantis & Jonides, 1996; Yeshurun & Carrasco, 1998, 2008), and the effects
of endogenous attention scale with cue validity, whereas those of exogenous attention do not
(e.g., Giordano et al., 2009; Kinchla, 1980; Mangun & Hillyard, 1990; Sperling & Melchner, 1978).
367
To manipulate exogenous attention, we used a peripheral cue (a bolded placeholder) presented
before the target onset. According to a normalization model of attention, exogenous attention can
also affect contrast gain when the attentional window is large enough (Reynolds & Heeger, 2009;
Herrmann et al., 2010). To induce a large attentional window while maintaining overlap between
the target and adaptors and ensure the adaptation effect, we randomly presented the target in
one of the five locations within the placeholders (**Figure 6**), and participants were explicitly
instructed to attend to the whole space encompassed by the placeholder, as the target could
appear anywhere within the placeholder. This procedure has been successfully used to
manipulate the size of the attentional window in both exogenous and endogenous covert spatial
attention, as well as in presaccadic attention (e.g., Binda & Murray, 2015; Cutrone, Heeger, &
Carrasco, 2018; Feng & Spence, 2017; Grubb et al., 2013; Herrmann et al., 2010; Li et al., 2021).
379
380   **Methods**
381
382   **Participants**
383   Eleven out of 12 participants[1] who participated in Experiment 1, including author HHL, also
384   participated in Experiment 2. We tested the same group of participants to compare the results
385   from endogenous and exogenous attentional effects after adaptation.
386
Stimuli and apparatus
Figure 6 shows an experimental trial. The target stimuli and the apparatus were the same as
Experiment 1. The placeholders in Experiment 2 (length = 0.256° for placeholders that were
further away from the center, length = 0.192° for placeholders that were closer to the center, all
width = 0.06°) were larger, given that there were 5 possible target locations: center and 2° on the
upper, lower, left, or right of the central Gabor. During the cue presentation, the placeholders
became thicker (6 pixels bigger for the frame elements closer to the center and 8 pixels bigger for
the frame elements further away from the center) to capture participants' exogenous attention.
395
396   **Experimental design and procedures**
397
398   The same c_{50} contrast derived from Experiment 1 was used in Experiment 2 for the adapted and
399   non-adapted conditions across locations. The experimental design and procedures were the
400   same as in Experiment 1, except for the following: After the top-up, there was a 200-ms ISI before
401   the exogenous pre-cue appeared for 60 ms, followed by 40-ms ISI. The tilted Gabor was then
402   presented for 67 ms followed by another 200-ms ISI and the response cue. Participants were

---

[1] One participant was no longer around NYC to participate

403 explicitly told that the exogenous cues were not informative; i.e., they were equally likely to be
404 valid, neutral and invalid (33% each). Participants were instructed to enlarge their attentional
405 window during the task, as they were explicitly told that the target could appear anywhere within
406 the placeholders.
407

408
409
Figure 6. Experimental procedure: The procedure was the same as Experiment 1 except for the pre-cue
and ISI timings. The pre-cue (bolded placeholders) either matches (valid condition), mismatches (invalid
condition) the response cue, or does not provide location information (neutral condition). The placeholders
were wider (8°) than Experiment 1. The target, two vertical Gabor stimuli were presented on average 8°
away from the center (e.g., at the horizontal meridian for example here; at the vertical meridian in a different
block). There were 5 possible target locations, which were 2° away from the central Gabor. For illustration
purposes, the stimulus size and spatial frequency shown here are not to scale.
417
418

Results

420
Figure 7 shows our results. As in Experiment 1, we compared the exogenous attentional effect
on d′ by conducting a three-way ANOVA on the factors of location (HM, upper, lower), attentional
validity (valid, neutral, invalid), and adaptation (adaptation, non-adaptation). There was a main
effect of attention [$F(2,20)=20.7$, $p<.001$, $\eta_p^2=0.67$], but neither of location [$F(2,20)=2.61$, $p=.099$],
nor of adaptation [$F(1,10)=1.08$, $p>.1$]. There was neither a 3-way interaction [$F(4,40)=2.36$,
$p=.069$] nor 2-way interactions [all $ps>.1$].
427

428
Figure 7. Performance in Experiment 2. (A) d' was higher in the valid followed by neutral and invalid
conditions in both non-adapted and adapted conditions. There was no difference between the adapted and
non-adapted conditions. (B) The attentional effects were similar around polar angle –horizontal meridian
(HM), and upper and lower vertical meridian– and were comparable in the adapted and non-adapted
conditions. The error bars above the bar plots indicate ±1 SEM of the difference between conditions. ***
$p<.001$, ** $p<.01$, * $p<.05$, n.s. $p>.05$.

435
436

The results here were further confirmed by separating the adapted and non-adapted conditions
into two 2-way ANOVAs on attention and location. For the non-adapted condition, we observed a
main effect of attention [$F(2,20)=13.33$, $p<.001$, $\eta_p^2=0.57$] but neither of location [$F(2,20)<1$], nor
an interaction [$F(4,40)=1.33$, $p>.1$]. The same pattern emerged for the adapted condition: a main
effect of attention [$F(2,20)=21.19$, $p<.001$, $\eta_p^2=0.68$] but neither of location [$F(2,20)=2.33$, $p>.1$]
nor an interaction [$F(4,40)=2.01$, $p>.1$].

444 We plot the individual data for the exogenous attentional effect (valid d' − invalid d') in the adapted
445 and non-adapted conditions (**Figure 8**). There was no difference between the two conditions
446 [$t(10)<1$].

448 In sum, the exogenous attentional effect was comparable across locations and adaptation
449 conditions.

451
452 **Figure 8**. Comparison of the exogenous attentional effects (valid d' − invalid d') in Experiment 2. The
453 attentional effects were comparable in the adapted and non-adapted conditions. The red circle indicates
454 the mean of all participants, and the error bars indicate ±1 SEM.

457 **Comparing Experiments 1 and 2**

459 Given that we had 11 common participants in Experiments 1 and 2, we conducted a 4-way
460 repeated-measures within-subject ANOVA on the factor of type of attention (endogenous,
461 exogenous), attentional validity (valid, neutral, invalid), adaptation (adapted, non-adapted), and
462 location (HM, upper, lower). There was a main effect of attentional validity [$F(2,20)=51.72$, $p<.001$,
463 $\eta_p^2=0.84$], and an interaction between attentional validity and type of attention [$F(2,20)=7.38$,
464 $p=.004$, $\eta_p^2=0.42$]. Post-hoc analyses indicated that valid condition had the highest d' followed by
465 neutral [valid − neutral: $t(10)=6.78$, $p<.001$, $d=2.04$] and invalid conditions [neutral − invalid:
466 $t(10)=6.17$, $p<.001$, $d=1.86$]. The attentional effect (valid d' − invalid d') was stronger for
467 endogenous than exogenous attention [$t(10)=2.95$, $p=.015$, $d=0.89$]. Importantly, there were no
468 4-way interaction [$F(4,40)<1$] nor any other significant effects [all $ps>.05$], indicating that the effect
469 for both types of attention did not vary across locations nor across adaptation conditions.

470
471 Furthermore, we found a positive Pearson correlation [r=.39, p=.025] between the exogenous and
472 endogenous overall attentional effect (collapsing across adaptation conditions and locations),
473 which indicates that those observers who had a stronger effect of one type of attention also had
474 a stronger effect for the other type.

475
476

Discussion

478
In this study, we investigated whether attention interacts with adaptation around polar angle. Our
results are consistent with separate studies showing: (1) without adaptation, the typical
performance fields emerged, with lower contrast thresholds at the horizontal than the vertical
meridian (HVA) and at the lower than the upper vertical meridian (VMA) (e.g., Abrams et al., 2012;
Baldwin et al., 2012; Cameron et al., 2002; Carrasco et al., 2001; Corbett & Carrasco, 2011; Fuller
et al., 2008; Himmelberg et al., 2020; Lee & Carrasco, 2025); (2) adaptation effects were stronger
at the horizontal than the vertical meridian (Lee & Carrasco, 2025); and (3) both endogenous
attention (Purokayastha et al., 2021; Tünçok, Carrasco, & Winawer, 2025) and exogenous
attention (Cameron et al., 2002; Carrasco et al., 2001; Roberts et al., 2018; Roberts et al., 2016)
enhanced contrast sensitivity similarly across all tested locations. Furthermore, our results
revealed that: (1) endogenous attention restored contrast sensitivity following adaptation, (2)
endogenous and exogenous attention had similar effects on contrast sensitivity before and after
adaptation–both enhanced contrast sensitivity at the attended location, with concomitant costs at
unattended locations, and (3) they did so uniformly at the cardinal meridians around the visual
field–despite differential adaptation effects.

494
The finding that endogenous attention enhances contrast sensitivity to a similar extent in adapted
and non-adapted conditions indicates that visual adaptation does not modulate the attentional
effect. This novel finding is consistent with corresponding findings for exogenous attention on
contrast sensitivity after adaptation (Lee et al., 2024; Pestilli et al., 2007). Despite its flexible
nature (reviews: Carrasco, 2011, 2014; Carrasco & Barbot, 2014; Olivers, 2025), endogenous
attention neither increased nor decreased contrast sensitivity differentially before and after
adaptation, indicating that these two processes, which help manage limited bioenergetic
resources, play independent roles in shaping performance.

503
Typically, the effect of exogenous attention manifests as response gain and the effect of
endogenous attention as contrast gain (Ling & Carrasco, 2006a; Pestilli et al., 2009). In the
exogenous attention experiment, we induced contrast gain by manipulating the size of the
attentional window, presenting the target Gabor at one of 5 different locations within a larger
stimulus placeholder. According to Reynolds and Heeger's normalization model of attention,
attention produces contrast gain rather than response gain when the attentional window is large
relative to stimulus size (Reynolds & Heeger, 2009), a prediction confirmed psychophysically and
with functional magnetic resonance imaging (fMRI) (Herrmann et al., 2010). By contrast,
endogenous attention can induce response gain when deployed over a relatively smaller
attentional window than the stimulus size (Fernández et al., 2023; Herrmann et al., 2010; Morrone,

Denti, & Spinelli, 2004). Consistent with previous findings (Lee et al., 2024; Pestilli et al., 2007),
exogenous attention modulated contrast sensitivity to a similar extent in adapted and non-adapted
conditions, indicating that adaptation did not modulate its effect. These results support Hypothesis
1: after adaptation, covert spatial attention modulates contrast sensitivity to the same extent as
without adaptation (**Figure 1A**).
519

In this exogenous attention experiment, to induce a larger attentional window, participants were
explicitly told that the target could appear anywhere within the placeholders. This manipulation
should not affect the effects of exogenous attention as it cannot induce endogenous attention.
The effects of endogenous attention scale with cue validity (e.g., Giordano et al., 2009; Kinchla,
1980; Mangun & Hillyard, 1990; Sperling & Melchner, 1978), and in the exogenous attention
experiment, the cue was uninformative: each of the valid, invalid, and neutral cues was presented
on 33% of the trials, so when a cue indicated one location out of two, its validity was 50%. Thus,
had observers deployed endogenous attention in Experiment 2, performance would have been
similar for valid and invalid conditions. Instead, we found significant benefits at the attended
location and significant costs at unattended locations, consistent with an exogenous attention
effect. Moreover, given the timing of the exogenous cue (~120 ms)–and that endogenous
attention takes ~300 ms to be deployed (e.g., Cheal, Lyon, & Hubbard, 1991; Geweke, Pokta, &
Störmer, 2021; Liu, Stevens, & Carrasco, 2007; Nakayama & Mackeben, 1989; Remington,
Johnston, & Yantis, 1992, for reviews, see Carrasco 2011, 2014, Carrasco & Barbot, 2015)–
endogenous attention could not contribute.
535

Adaptation was more pronounced at the horizontal than the vertical meridian. Unlike our previous
study (Lee & Carrasco, 2025), which blocked each target location, here we introduced greater
target uncertainty by using two possible target locations per trial (**Figure 2**). The replication of the
adaptation pattern across studies shows that that the previous findings are robust to target
uncertainty and generalize across participants. Most adaptation studies have examined only the
horizontal meridian (e.g., Beaton & Blakemore, 1981; Carrasco et al., 2006; Gao, Webster, &
Jiang, 2019; Greenlee, Georgeson, Magnussen, & Harris, 1991; Pestilli et al., 2007; Schieting &
Spillmann, 1987), only the vertical meridian (e.g., Bell, Gheorghiu, Hess, & Kingdom, 2011; Bell,
Gheorghiu, & Kingdom, 2009; Montaser-Kouhsari & Rajimehr, 2004), or did not analyze target
locations separately (e.g., Bao, Fast, Mesik, & Engel, 2013; Lin, Zhou, Naya, Gardner, & Sun,
2021; Ling & Carrasco, 2006b; Zimmermann, Weidner, Abdollahi, & Fink, 2016). Our results add
further evidence that adaptation differs across meridians, an important finding to consider in future
studies and models of vision.
549

Endogenous and exogenous attention enhanced contrast sensitivity similarly around polar angle,
despite the differential effects of adaptation. Consistent with previous studies (Cameron et al.,
2002; Carrasco et al., 2001; Roberts et al., 2018; Roberts et al., 2016), asymmetries at the
cardinal locations were resistant to both endogenous and exogenous attention, indicating their
resilient nature and that they cannot be easily reshaped. In contrast, consistent with a recent
finding (Lee & Carrasco, 2025), visual adaptation reduced contrast sensitivity more at the
horizontal than the vertical meridian, yet neither type of covert spatial attention modulated the
extent of the asymmetries altering the shape of the performance fields, notwithstanding the

differential adaptation effect. This similar effect is notable given that endogenous attention is flexible and exogenous attention automatic (e.g., Carrasco, 2011, 2014; Carrasco & Barbot, 2014; Olivers, 2025), yet neither compensated for poor performance. These findings provide further evidence regarding the resilience of polar angle asymmetries and support Hypothesis 4 (**Figure 1D**): visual adaptation does not modulate the effects of covert spatial attention, even at the location of poorest performance.

What contributes to performance asymmetries in the HVA and VMA? These asymmetries arise from both retinal and cortical factors. Retinally, cone density is higher at the horizontal than the vertical meridian (Curcio, Sloan Jr, Packer, Hendrickson, & Kalina, 1987; Curcio, Sloan, Kalina, & Hendrickson, 1990), and midget-RGC density is higher at the lower than the upper vertical meridian (Curcio et al., 1990; Song, Chui, Zhong, Elsner, & Burns, 2011). Cortically, V1 surface area is larger for the horizontal than the vertical meridian, and for the lower than the upper vertical meridian (Benson, Kupers, Barbot, Carrasco, & Winawer, 2021; Himmelberg et al., 2021; Himmelberg, Kwak, Carrasco, & Winawer, 2025; Himmelberg, Tünçok, et al., 2023; Himmelberg, Winawer, & Carrasco, 2022, 2023; Lee & Carrasco, 2025; Silva et al., 2018). Moreover, cortical factors account for more variance in these asymmetries than retinal factors (Kupers, Benson, Carrasco, & Winawer, 2022). Still, these factors cannot fully explain behavioral differences observed in psychophysical tasks, which are diminished but still present once stimulus size is cortically magnified (Jigo, Tavdy, Himmelberg, & Carrasco, 2023), suggesting that additional factors –such as sensory tuning and neuronal computations– also contribute to the HVA and VMA (Himmelberg, Winawer, & Carrasco, 2023; Jigo et al., 2023; Xue, Barbot, Abrams, Chen, & Carrasco, 2025).

Endogenous and exogenous attention rely on different neural substrates. fMRI studies show differential activity modulation across the frontoparietal network (Beck & Kastner, 2014; Buschman & Miller, 2009; Chica, Bartolomeo, & Lupiáñez, 2013; Fiebelkorn & Kastner, 2020; Kastner & Buschman, 2017; Meyer, Du, Parks, & Hopfinger, 2018), temporoparietal junction (Dugué, Merriam, Heeger, & Carrasco, 2018), and visual cortex (Dugué, Merriam, Heeger, & Carrasco, 2020; Hopfinger & West, 2006; Ling, Jehee, & Pestilli, 2015). Transcranial magnetic stimulation (TMS) studies, which disrupt the neuronal balance between excitation and inhibition (Bradley, Nydam, Dux, & Mattingley, 2022; Kobayashi & Pascual-Leone, 2003; Valero-Cabré, Pascual-Leone, & Coubard, 2011), revealed that early visual cortex plays a critical role for adaptation (Lee et al., 2024; Perini et al., 2012) and exogenous attention (Fernández & Carrasco, 2020; Lee et al., 2024), whereas the human homologue of the right frontal eye fields (rFEF+) plays a critical role for endogenous attention (Fernández et al., 2023). Critically, disrupting rFEF+ does not affect exogenous attention (Chen et al., 2025), and disrupting early visual cortex does not affect endogenous attention (Fernández et al., 2023), indicating a double dissociation. Despite these distinct neural underpinnings, both types of covert spatial attention affected contrast sensitivity uniformly at the cardinal meridians around polar angle and did not interact with location or adaptation. These findings suggest that distinct neuronal populations underlie polar angle asymmetries, adaptation, and attentional modulation.

We found stronger attentional effects for endogenous than exogenous attention. Given that
adaptation is more effective when the adaptor and the target spatially overlap (Kovács, Zimmer,
Harza, & Vidnyánszky, 2007; Larsson & Harrison, 2015; Webster, 2011, 2015), we introduced
target uncertainty with 5 possible target locations and allowed 2° overlap between adaptor and
target to elicit adaptation while allowing exogenous attention to operate via contrast gain. This
manipulation may have yielded a slightly narrower exogenous attentional window than for
endogenous attention in our design, as well as compared with previous studies. For example,
Herrmann et al. (2010) used five possible target locations with no overlap, whereas in our current
study the target Gabors could overlap by 2° within placeholders. According to Reynolds and
Heeger's normalization model of attention (Reynolds & Heeger, 2009), attention multiplies
stimulus-evoked activity before divisive normalization. In our task, normalization may have pooled
a broader suppressive drive than in typical exogenous attention tasks, but not as broad as in
typical endogenous attention tasks —leading to less pronounced contrast gain and thus weaker
exogenous than endogenous attention effects.

Why do type of spatial covert attention, adaptation, and polar angle asymmetries not interact?
The visual cortex plays a crucial role in all three processes. fMRI studies have shown that covert
endogenous spatial attention modulates activity in visual cortex via feedback from frontoparietal
cortex (Buschman & Miller, 2009; Chica et al., 2013; Corbetta, Patel, & Shulman, 2008; Corbetta
& Shulman, 2002; Dugué et al., 2020; Lauritzen, D'Esposito, Heeger, & Silver, 2009; Pestilli,
Carrasco, Heeger, & Gardner, 2011) and increasingly modulates activity in the occipital visual
areas (Dugué et al., 2020), with V1/V2, its early visual areas, being not critical for endogenous
attention, as TMS on these areas does not alter its effect on visual perception (Fernández et al.,
2023). In contrast, exogenous attention modulates visual cortex via feedforward activation
(Dassanayake, Michie, & Fulham, 2016; Dugué et al., 2020; Hopfinger, Luck, & Hillyard, 2004;
Liu, Pestilli, & Carrasco, 2005; F. Wang, Chen, Yan, Zhaoping, & Li, 2015; Westerberg, Schall,
Woodman, & Maier, 2023), and V1/V2 are critical for its effect, as TMS on these areas eliminates
the effect of exogenous attention on visual perception (Fernández & Carrasco, 2020; Lee et al.,
2024). Moreover, these two attention types also differentially modulate visual subregions of the
temporoparietal-junction (Dugué et al., 2018). All these differences underscore the distinct
contributions of endogenous and exogenous in modulating visual perception.

Early visual cortex also plays a critical role in visual adaptation. TMS over V1/V2 decreases
contrast adaptation (Perini et al., 2012), and adaptation modulates contrast response functions in
V1/V2 (Altan et al., 2025; Gardner et al., 2005; Vinke, Bloem, & Ling, 2022). A TMS study revealed
that adaptation and exogenous attention interact in early visual cortex (Lee et al., 2024), but it is
unknown whether they do so systematically around polar angle, as several factors shape
asymmetries. Moreover, it is presently unknown whether endogenous attention and adaptation
interact either in early occipital or in frontal areas.

Both adaptation (Lee & Carrasco, 2025) and polar angle asymmetries (Benson et al., 2021;
Himmelberg et al., 2022; Himmelberg, Winawer, & Carrasco, 2023; Lee & Carrasco, 2025)
correlate with V1 surface area, but surface area alone cannot fully account for these asymmetries
(Jigo et al., 2023). Additional factors such as neural gain also contribute to these asymmetries

(Xue et al., 2025). Future research integrating computational modeling, neuroimaging,
neurostimulation, and psychophysics will be essential to assess the relative contributions of
cortical and computational factors to attention, adaptation and polar angle asymmetries.
648
In conclusion, this study reveals that performance asymmetries are resistant to the effects of both
endogenous and exogenous covert spatial attention, despite their distinct temporal dynamics and
differences in flexibility–even after adaptation induces differential effects across meridians.
Although both adaptation and attention help allocate limited resources according to task demands,
neither type of covert spatial attention differentially enhance target processing at locations that
differ in intrinsic discriminability and their corresponding representation in cortical surface area.
655
656

657 **Acknowledgments**
658 This study was supported by NIH NEI R01-EY027401 to M.C. and the Ministry of Education in
659 Taiwan and 2025 National Science and Technology Council (NSTC) Taiwanese Overseas
660 Pioneers Grants (TOP Grants) for PhD Candidates to H.-H.L. We thank Nick Crotty, David Tu
661 and Shutian Xue for their feedback on a previous version of the manuscript.

662
663

References
Abrams, J., Nizam, A., & Carrasco, M. (2012). Isoeccentric locations are not equivalent: The
extent of the vertical meridian asymmetry. *Vision Research, 52*(1), 70-78.
Afraz, A., Pashkam, M. V., & Cavanagh, P. (2010). Spatial heterogeneity in the perception of face
and form attributes. *Current Biology, 20*(23), 2112-2116.
Altan, E., Morgan, C. A., Dakin, S. C., & Schwarzkopf, D. S. (2025). Spatial frequency adaptation
modulates population receptive field sizes. *elife, 13*, RP100734.
Altpeter, E., Mackeben, M., & Trauzettel-Klosinski, S. (2000). The importance of sustained
attention for patients with maculopathies. *Vision Research, 40*(10-12), 1539-1547.
Baldwin, A. S., Meese, T. S., & Baker, D. H. (2012). The attenuation surface for contrast sensitivity
has the form of a witch's hat within the central visual field. *Journal of Vision, 12*(11), 23.
Bao, M., Fast, E., Mesik, J., & Engel, S. (2013). Distinct mechanisms control contrast adaptation
over different timescales. *Journal of Vision, 13*(10), 14.
Barbot, A., & Carrasco, M. (2017). Attention modifies spatial resolution according to task demands.
Psychological Science, 28(3), 285-296.
Barbot, A., Landy, M. S., & Carrasco, M. (2012). Differential effects of exogenous and
endogenous attention on second-order texture contrast sensitivity. *Journal of Vision, 12*(8),
6.
Barbot, A., Xue, S., & Carrasco, M. (2021). Asymmetries in visual acuity around the visual field.
Journal of Vision, 21(1), 2.
Beaton, A., & Blakemore, C. (1981). Orientation selectivity of the human visual system as a
function of retinal eccentricity and visual hemifield. *Perception, 10*(3), 273-282.
Beck, D. M., & Kastner, S. (2014). Neural systems for spatial attention in the human brain:
Evidence from neuroimaging in the framework of biased competition. In A. C. Nobre & S.
Kastner (Eds.), *Oxford Handbook of Attention*: Oxford University Press.
Bell, J., Gheorghiu, E., Hess, R. F., & Kingdom, F. A. (2011). Global shape processing involves
a hierarchy of integration stages. *Vision Research, 51*(15), 1760-1766.

Bell, J., Gheorghiu, E., & Kingdom, F. A. A. (2009). Orientation tuning of curvature adaptation
reveals both curvature-polarity-selective and non-selective mechanisms. *Journal of Vision,*
9(12), 3.
Benson, N. C., Kupers, E. R., Barbot, A., Carrasco, M., & Winawer, J. (2021). Cortical
magnification in human visual cortex parallels task performance around the visual field.
Elife, 10, e67685.
Binda, P., & Murray, S. O. (2015). Spatial attention increases the pupillary response to light
changes. *Journal of Vision, 15*(2), 1.
Boynton, G. M., & Finney, E. M. (2003). Orientation-specific adaptation in human visual cortex.
Journal of Neuroscience, 23(25), 8781-8787.
Bradley, C., Nydam, A. S., Dux, P. E., & Mattingley, J. B. (2022). State-dependent effects of neural
stimulation on brain function and cognition. *Nature Reviews Neuroscience, 23*(8), 459-
475.
Brainard, D. H. (1997). The psychophysics toolbox. *Spatial Vision, 10*(4), 433-436.
Buschman, T. J., & Miller, E. K. (2009). Serial, covert shifts of attention during visual search are
reflected by the frontal eye fields and correlated with population oscillations. *Neuron, 63*(3),
386-396.
Cameron, E. L., Tai, J. C., & Carrasco, M. (2002). Covert attention affects the psychometric
function of contrast sensitivity. *Vision Research, 42*(8), 949-967.
Carrasco, M. (2011). Visual attention: The past 25 years. *Vision Research, 51*(13), 1484-1525.
Carrasco, M. (2014). Spatial covert attention: Perceptual modulation. *The Oxford Handbook of*
Attention, 183, 230.
Carrasco, M., & Barbot, A. (2014). *How attention affects spatial resolution.* Paper presented at
the Cold Spring Harbor Symposia on Quantitative Biology.
Carrasco, M., & Barbot, A. (2019). Spatial attention alters visual appearance. *Current Opinion in*
Psychology, 29, 56-64.
Carrasco, M., Loula, F., & Ho, Y.-X. (2006). How attention enhances spatial resolution: Evidence
from selective adaptation to spatial frequency. *Perception & Psychophysics, 68*(6), 1004-
1012.
Carrasco, M., Talgar, C. P., & Cameron, E. L. (2001). Characterizing visual performance fields:
Effects of transient covert attention, spatial frequency, eccentricity, task and set size.
Spatial Vision, 15(1), 61.
Carrasco, M., Williams, P. E., & Yeshurun, Y. (2002). Covert attention increases spatial resolution
with or without masks: Support for signal enhancement. *Journal of Vision, 2*(6), 4.
Chakravarthi, R., Papadaki, D., & Krajnik, J. (2022). Visual field asymmetries in numerosity
processing. *Attention, Perception, & Psychophysics, 84*(8), 2607-2622.
Cheal, M., Lyon, D. R., & Hubbard, D. C. (1991). Does attention have different effects on line
orientation and line arrangement discrimination? *The Quarterly Journal of Experimental*
Psychology, 43(4), 825-857.
Chen, Q., Lee, H.-H., Hoxha, K., Fernández, A., Hanning, N. M., & Carrasco, M. (2025). Does
human right frontal eye field (rFEF+) play a critical role in exogenous attention? A
Transcranial Magnetic Stimulation (TMS) study. *Journal of Vision, 25*(9), 2612-2612.
Chica, A. B., Bartolomeo, P., & Lupiáñez, J. (2013). Two cognitive and neural systems for
endogenous and exogenous spatial attention. *Behavioural Brain Research, 237*, 107-123.
Corbett, J. E., & Carrasco, M. (2011). Visual performance fields: Frames of reference. *PLoS One,*
6(9), e24470.
Corbetta, M., Patel, G., & Shulman, G. L. (2008). The reorienting system of the human brain:
From environment to theory of mind. *Neuron, 58*(3), 306-324.
Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven attention in
the brain. *Nature Reviews Neuroscience, 3*(3), 201-215.

Crotty, N., Massa, N., Tellez, D., White, A. L., & Grubb, M. A. (2025). A preliminary investigation of the interaction between expectation and the reflexive allocation of covert spatial attention. *Scientific Reports, 15*(1), 5778.

Curcio, C. A., Sloan Jr, K. R., Packer, O., Hendrickson, A. E., & Kalina, R. E. (1987). Distribution of cones in human and monkey retina: Individual variability and radial asymmetry. *Science, 236*(4801), 579-582.

Curcio, C. A., Sloan, K. R., Kalina, R. E., & Hendrickson, A. E. (1990). Human photoreceptor topography. *Journal of Comparative Neurology, 292*(4), 497-523.

Cutrone, E. K., Heeger, D. J., & Carrasco, M. (2018). On spatial attention and its field size on the repulsion effect. *Journal of Vision, 18*(6), 8.

Dao, D. Y., Lu, Z.-L., & Dosher, B. A. (2006). Adaptation to sine-wave gratings selectively reduces the contrast gain of the adapted stimuli. *Journal of Vision, 6*(7), 6.

Dassanayake, T. L., Michie, P. T., & Fulham, R. (2016). Effect of temporal predictability on exogenous attentional modulation of feedforward processing in the striate cortex. *International Journal of Psychophysiology, 105*, 9-16.

Desimone, R., & Duncan, J. (1995). Neural mechanisms of selective visual attention. *Annual Review of Neuroscience, 18*(1), 193-222.

Dosher, B. A., & Lu, Z.-L. (2000a). Mechanisms of perceptual attention in precuing of location. *Vision Research, 40*(10), 1269-1292.

Dosher, B. A., & Lu, Z.-L. (2000b). Noise exclusion in spatial attention. *Psychological Science, 11*(2), 139-146.

Dugué, L., Merriam, E. P., Heeger, D. J., & Carrasco, M. (2018). Specific visual subregions of TPJ mediate reorienting of spatial attention. *Cerebral Cortex, 28*(7), 2375-2390.

Dugué, L., Merriam, E. P., Heeger, D. J., & Carrasco, M. (2020). Differential impact of endogenous and exogenous attention on activity in human visual cortex. *Scientific Reports, 10*(1), 21274.

Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G* Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods, 39*(2), 175-191.

Feng, J., & Spence, I. (2017). The effects of spatial endogenous pre-cueing across eccentricities. *Frontiers in Psychology, 8*, 888.

Fernández, A., & Carrasco, M. (2020). Extinguishing exogenous attention via transcranial magnetic stimulation. *Current Biology, 30*(20), 4078-4084. e4073.

Fernández, A., Denison, R. N., & Carrasco, M. (2019). Temporal attention improves perception similarly at foveal and parafoveal locations. *Journal of Vision, 19*(1), 12.

Fernández, A., Hanning, N. M., & Carrasco, M. (2023). Transcranial magnetic stimulation to frontal but not occipital cortex disrupts endogenous attention. *Proceedings of the National Academy of Sciences, 120*(10), e2219635120.

Fernández, A., Okun, S., & Carrasco, M. (2022). Differential effects of endogenous and exogenous attention on sensory tuning. *Journal of Neuroscience, 42*(7), 1316-1327.

Fiebelkorn, I. C., & Kastner, S. (2020). Functional specialization in the attention network. *Annual Review of Psychology, 71*(1), 221-249.

Fuller, S., & Carrasco, M. (2009). Perceptual consequences of visual performance fields: The case of the line motion illusion. *Journal of Vision, 9*(4), 13.

Fuller, S., Rodriguez, R. Z., & Carrasco, M. (2008). Apparent contrast differs across the vertical meridian: Visual and attentional factors. *Journal of Vision, 8*(1), 16.

Gao, Y., Webster, M. A., & Jiang, F. (2019). Dynamics of contrast adaptation in central and peripheral vision. *Journal of Vision, 19*(6), 23.

Gardner, J. L., Sun, P., Waggoner, R. A., Ueno, K., Tanaka, K., & Cheng, K. (2005). Contrast adaptation and representation in human early visual cortex. *Neuron, 47*(4), 607-620.

Geweke, F., Pokta, E., & Störmer, V. S. (2021). Spatial distance of target locations affects the time course of both endogenous and exogenous attentional deployment. *Journal of Experimental Psychology: Human Perception and Performance, 47*(6), 774.

Giordano, A. M., McElree, B., & Carrasco, M. (2009). On the automaticity and flexibility of covert attention: A speed-accuracy trade-off analysis. *Journal of Vision, 9*(3), 30.

Greenlee, M. W., Georgeson, M. A., Magnussen, S., & Harris, J. P. (1991). The time course of adaptation to spatial contrast. *Vision Research, 31*(2), 223-236.

Greenwood, J. A., Szinte, M., Sayim, B., & Cavanagh, P. (2017). Variations in crowding, saccadic precision, and spatial localization reveal the shared topology of spatial vision. *Proceedings of the National Academy of Sciences, 114*(17), E3573-E3582.

Grubb, M. A., Behrmann, M., Egan, R., Minshew, N. J., Carrasco, M., & Heeger, D. J. (2013). Endogenous spatial attention: Evidence for intact functioning in adults with autism. *Autism Research, 6*(2), 108-118.

Hanning, N. M., Himmelberg, M. M., & Carrasco, M. (2022). Presaccadic attention enhances contrast sensitivity, but not at the upper vertical meridian. *Iscience, 25*(2).

Hanning, N. M., Himmelberg, M. M., & Carrasco, M. (2024). Presaccadic attention depends on eye movement direction and is related to V1 cortical magnification. *Journal of Neuroscience, 44*(12).

Hein, E., Rolke, B., & Ulrich, R. (2006). Visual attention and temporal discrimination: Differential effects of automatic and voluntary cueing. *Visual Cognition, 13*(1), 29-50.

Herrmann, K., Montaser-Kouhsari, L., Carrasco, M., & Heeger, D. J. (2010). When size matters: Attention affects performance by contrast or response gain. *Nature Neuroscience, 13*(12), 1554-1559.

Himmelberg, M. M., Kurzawski, J. W., Benson, N. C., Pelli, D. G., Carrasco, M., & Winawer, J. (2021). Cross-dataset reproducibility of human retinotopic maps. *Neuroimage, 244*, 118609.

Himmelberg, M. M., Kwak, Y., Carrasco, M., & Winawer, J. (2025). Unpacking the V1 map: Differential covariation of preferred spatial frequency and cortical magnification across spatial dimensions. *PLOS Computational Biology, 21*(10), e1013599.

Himmelberg, M. M., Tünçok, E., Gomez, J., Grill-Spector, K., Carrasco, M., & Winawer, J. (2023). Comparing retinotopic maps of children and adults reveals a late-stage change in how V1 samples the visual field. *Nature Communications, 14*(1), 1561.

Himmelberg, M. M., Winawer, J., & Carrasco, M. (2020). Stimulus-dependent contrast sensitivity asymmetries around the visual field. *Journal of Vision, 20*(9), 18.

Himmelberg, M. M., Winawer, J., & Carrasco, M. (2022). Linking individual differences in human primary visual cortex to contrast sensitivity around the visual field. *Nature Communications, 13*(1), 3309.

Himmelberg, M. M., Winawer, J., & Carrasco, M. (2023). Polar angle asymmetries in visual perception and neural architecture. *Trends in Neurosciences, 46*(6), 445-458.

Hopfinger, J. B., Luck, S. J., & Hillyard, S. A. (2004). Selective attention: Electrophysiological and neuromagnetic studies. *The Cognitive Neurosciences, 3*, 561-574.

Hopfinger, J. B., & West, V. M. (2006). Interactions between endogenous and exogenous attention on cortical visual processing. *NeuroImage, 31*(2), 774-789.

Huang, D., Liao, K., Chen, F., & Chen, Y. (2025). The effects of acute alcohol intake on endogenous vs. exogenous attention in visual perception. *Addictive Behaviors*, 108465.

Jigo, M., & Carrasco, M. (2018). Attention alters spatial resolution by modulating second-order processing. *Journal of Vision, 18*(7), 2.

Jigo, M., & Carrasco, M. (2020). Differential impact of exogenous and endogenous attention on the contrast sensitivity function across eccentricity. *Journal of Vision, 20*(6), 11.

Jigo, M., Heeger, D. J., & Carrasco, M. (2021). An image-computable model of how endogenous and exogenous attention differentially alter visual perception. *Proceedings of the National Academy of Sciences, 118*(33), e2106436118.

Jigo, M., Tavdy, D., Himmelberg, M. M., & Carrasco, M. (2023). Cortical magnification eliminates differences in contrast sensitivity across but not around the visual field. *eLife, 12*, e84205.

Kastner, S., & Buschman, T. J. (2017). Visual attention. In *Oxford Research Encyclopedia of Neuroscience*.

Kim, C., & Chong, S. C. (2024). Metacognition of perceptual resolution across and around the visual field. *Cognition, 253*, 105938.

Kinchla, R. (1980). The measurement of attention. In R. S. Nikerson (Ed.), *Attention and performance VIII* (pp. 213-238): Psychology Press.

Kobayashi, M., & Pascual-Leone, A. (2003). Transcranial magnetic stimulation in neurology. *The Lancet Neurology, 2*(3), 145-156.

Kohn, A. (2007). Visual adaptation: Physiology, mechanisms, and functional benefits. *Journal of Neurophysiology, 97*(5), 3155-3164.

Kovács, G., Zimmer, M., Harza, I., & Vidnyánszky, Z. (2007). Adaptation duration affects the spatial selectivity of facial aftereffects. *Vision Research, 47*(25), 3141-3149.

Kupers, E. R., Benson, N. C., Carrasco, M., & Winawer, J. (2022). Asymmetries around the visual field: From retina to cortex to behavior. *PLoS Computational Biology, 18*(1), e1009771.

Kurzawski, J. W., Burchell, A., Thapa, D., Winawer, J., Majaj, N. J., & Pelli, D. G. (2023). The Bouma law accounts for crowding in 50 observers. *Journal of Vision, 23*(8), 6.

Kwak, Y., Hanning, N. M., & Carrasco, M. (2023). Presaccadic attention sharpens visual acuity. *Scientific Reports, 13*(1), 2981.

Kwak, Y., Hanning, N. M., & Carrasco, M. (in press). Saccade direction modulates the temporal dynamics of presaccadic attention. *Journal of Vision*.

Kwak, Y., Zhao, Y., Lu, Z.-L., Hanning, N. M., & Carrasco, M. (2024). Presaccadic attention enhances and reshapes the Contrast Sensitivity Function differentially around the visual field. *eNeuro, 11*(9).

Larsson, J., & Harrison, S. J. (2015). Spatial specificity and inheritance of adaptation in human visual cortex. *Journal of Neurophysiology, 114*(2), 1211-1226.

Lauritzen, T. Z., D'Esposito, M., Heeger, D. J., & Silver, M. A. (2009). Top–down flow of visual spatial attention signals from parietal to occipital cortex. *Journal of Vision, 9*(13), 18.

Lee, H.-H., & Carrasco, M. (2025). Visual adaptation stronger at the horizontal than the vertical meridian: Linking performance with V1 cortical surface area. *Proceedings of the National Academy of Sciences, 122*(29), e2507810122.

Lee, H.-H., Fernández, A., & Carrasco, M. (2024). Adaptation and exogenous attention interact in the early visual cortex: A TMS study. *Iscience, 27*(11).

Lennie, P. (2003). The cost of cortical computation. *Current Biology, 13*(6), 493-497.

Li, H.-H., Pan, J., & Carrasco, M. (2021). Different computations underlie overt presaccadic and covert spatial attention. *Nature Human Behaviour, 5*(10), 1418-1431.

Lin, Y., Zhou, X., Naya, Y., Gardner, J. L., & Sun, P. (2021). Voxel-wise linearity analysis of increments and decrements in BOLD responses in human visual cortex using a contrast adaptation paradigm. *Frontiers in Human Neuroscience, 15*, 541314.

Ling, S., & Carrasco, M. (2006a). Sustained and transient covert attention enhance the signal via different contrast response functions. *Vision Research, 46*(8-9), 1210-1220.

Ling, S., & Carrasco, M. (2006b). When sustained attention impairs perception. *Nature Neuroscience, 9*(10), 1243-1245.

Ling, S., Jehee, J. F., & Pestilli, F. (2015). A review of the mechanisms by which attentional feedback shapes visual selectivity. *Brain Structure and Function, 220*(3), 1237-1250.

Liu, T., Pestilli, F., & Carrasco, M. (2005). Transient attention enhances perceptual performance and fMRI response in human visual cortex. *Neuron, 45*(3), 469-477.

891 Liu, T., Stevens, S. T., & Carrasco, M. (2007). Comparing the time course and efficacy of spatial
892    and feature-based attention. *Vision Research, 47*(1), 108-113.
893 Lu, Z.-L., Lesmes, L. A., & Dosher, B. A. (2002). Spatial attention excludes external noise at the
894    target location. *Journal of Vision, 2*(4), 4.
895 Luck, S. J., & Thomas, S. J. (1999). What variety of attention is automatically captured by
896    peripheral cues? *Perception & Psychophysics, 61*(7), 1424-1435.
897 Luzardo, F., & Yeshurun, Y. (2025). Large-scale examination of the benefit and cost of spatial
898    attention and their individual variability. *Cognition, 264*, 106242.
899 Mangun, G., & Hillyard, S. (1990). Allocation of visual attention to spatial locations: Tradeoff
900    functions for event-related brain potentials and detection performance. *Perception &*
901    *Psychophysics, 47*(6), 532-550.
902 Meyer, K. N., Du, F., Parks, E., & Hopfinger, J. B. (2018). Exogenous vs. endogenous attention:
903    Shifting the balance of fronto-parietal activity. *Neuropsychologia, 111*, 307-316.
904 Montaser-Kouhsari, L., & Carrasco, M. (2009). Perceptual asymmetries are preserved in short-
905    term memory tasks. *Attention, Perception, & Psychophysics, 71*(8), 1782-1792.
906 Montaser-Kouhsari, L., & Rajimehr, R. (2004). Attentional modulation of adaptation to illusory
907    lines. *Journal of Vision, 4*(6), 3.
908 Morrone, M. C., Denti, V., & Spinelli, D. (2004). Different attentional resources modulate the gain
909    mechanisms for color and luminance contrast. *Vision Research, 44*(12), 1389-1401.
910 Nakayama, K., & Mackeben, M. (1989). Sustained and transient components of focal visual
911    attention. *Vision Research, 29*(11), 1631-1647.
912 Olivers, C. N. (2025). Selective attention and eccentricity: A comprehensive review. *Neuroscience*
913    *& Biobehavioral Reviews*, 106368.
914 Palmieri, H., & Carrasco, M. (2024). Task demand mediates the interaction of spatial and temporal
915    attention. *Scientific Reports, 14*(1), 9228.
916 Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics: transforming numbers
917    into movies. *Spatial Vision, 10*(4), 437-442.
918 Perini, F., Cattaneo, L., Carrasco, M., & Schwarzbach, J. V. (2012). Occipital transcranial
919    magnetic stimulation has an activity-dependent suppressive effect. *Journal of*
920    *Neuroscience, 32*(36), 12361-12365.
921 Pestilli, F., & Carrasco, M. (2005). Attention enhances contrast sensitivity at cued and impairs it
922    at uncued locations. *Vision Research, 45*(14), 1867-1875.
923 Pestilli, F., Carrasco, M., Heeger, D. J., & Gardner, J. L. (2011). Attentional enhancement via
924    selection and pooling of early sensory responses in human visual cortex. *Neuron, 72*(5),
925    832-846.
926 Pestilli, F., Ling, S., & Carrasco, M. (2009). A population-coding model of attention's influence on
927    contrast response: Estimating neural effects from psychophysical data. *Vision Research,*
928    *49*(10), 1144-1153.
929 Pestilli, F., Viera, G., & Carrasco, M. (2007). How do attention and adaptation affect contrast
930    sensitivity? *Journal of Vision, 7*(7), 9.
931 Petrov, Y., & Meleshkevich, O. (2011). Asymmetries and idiosyncratic hot spots in crowding.
932    *Vision Research, 51*(10), 1117-1123.
933 Prins, N., & Kingdom, F. A. (2018). Applying the model-comparison approach to test specific
934    research hypotheses in psychophysical research using the Palamedes toolbox. *Frontiers*
935    *in Psychology, 9*, 1250.
936 Purokayastha, S., Roberts, M., & Carrasco, M. (2021). Voluntary attention improves performance
937    similarly around the visual field. *Attention, Perception, & Psychophysics, 83*(7), 2784-2794.
938 Ramamurthy, M., White, A. L., & Yeatman, J. D. (2024). Children with dyslexia show no deficit in
939    exogenous spatial attention but show differences in visual encoding. *Developmental*
940    *Science, 27*(3), e13458.

Remington, R. W., Johnston, J. C., & Yantis, S. (1992). Involuntary attentional capture by abrupt
onsets. *Perception & Psychophysics, 51*(3), 279-290.
Reynolds, J. H., & Heeger, D. J. (2009). The normalization model of attention. *Neuron, 61*(2),
168-185.
Roberts, M., Ashinoff, B. K., Castellanos, F. X., & Carrasco, M. (2018). When attention is intact in
adults with ADHD. *Psychonomic Bulletin & Review, 25*(4), 1423-1434.
Roberts, M., Cymerman, R., Smith, R. T., Kiorpes, L., & Carrasco, M. (2016). Covert spatial
attention is functionally intact in amblyopic human adults. *Journal of Vision, 16*(15), 30.
Schieting, S., & Spillmann, L. (1987). Flicker adaptation in the peripheral retina. *Vision Research,*
27(2), 277-284.
Silva, M. F., Brascamp, J. W., Ferreira, S., Castelo-Branco, M., Dumoulin, S. O., & Harvey, B. M.
(2018). Radial asymmetries in population receptive field size and cortical magnification
factor in early visual cortex. *NeuroImage, 167*, 41-52.
Song, H., Chui, T. Y. P., Zhong, Z., Elsner, A. E., & Burns, S. A. (2011). Variation of cone
photoreceptor packing density with retinal eccentricity and age. *Investigative*
Ophthalmology & Visual Science, 52(10), 7376-7384.
Sperling, G., & Melchner, M. J. (1978). The attention operating characteristic: Examples from
visual search. *Science, 202*(4365), 315-318.
Talgar, C. P., & Carrasco, M. (2002). Vertical meridian asymmetry in spatial resolution: Visual
and attentional factors. *Psychonomic Bulletin & Review, 9*(4), 714-722.
Team, R. C. (2000). R language definition. *Vienna, Austria: R foundation for statistical computing,*
3(1), 116.
Tsai, L.-T., Liao, K.-M., Hou, C.-H., Jang, Y., & Chen, C.-C. (2024). Visual field asymmetries in
visual word form identification. *Vision Research, 220*, 108413.
Tünçok, E., Carrasco, M., & Winawer, J. (2025). How spatial attention alters visual cortical
representation during target anticipation. *Nature Communications, 16*, 8746.
Tünçok, E., Kiorpes, L., & Carrasco, M. (2025). Opposite asymmetry in visual perception of
humans and macaques. *Current Biology, 35*(3), 681-687. e684.
Valero-Cabré, A., Pascual-Leone, A., & Coubard, O. A. (2011). Transcranial magnetic stimulation
(TMS) in basic and clinical neuroscience research. *Revue Neurologique, 167*(4), 291-316.
Vergeer, M., Mesik, J., Baek, Y., Wilmerding, K., & Engel, S. A. (2018). Orientation-selective
contrast adaptation measured with SSVEP. *Journal of Vision, 18*(5), 2.
Vinke, L. N., Bloem, I. M., & Ling, S. (2022). Saturating nonlinearities of contrast response in
human visual cortex. *Journal of Neuroscience, 42*(7), 1292-1302.
Wang, F., Chen, M., Yan, Y., Zhaoping, L., & Li, W. (2015). Modulation of neuronal responses by
exogenous attention in macaque primary visual cortex. *Journal of Neuroscience, 35*(39),
13419-13429.
Wang, Z., Murai, Y., & Whitney, D. (2020). Idiosyncratic perception: A link between acuity,
perceived position and apparent size. *Proceedings of the Royal Society B, 287*(1930),
20200825.
Webster, M. A. (2011). Adaptation and visual coding. *Journal of Vision, 11*(5), 3.
Webster, M. A. (2015). Visual adaptation. *Annual Review of Vision Science, 1*(1), 547-567.
Westerberg, J. A., Schall, J. D., Woodman, G. F., & Maier, A. (2023). Feedforward attentional
selection in sensory cortex. *Nature Communications, 14*(1), 5993.
Xue, S., Barbot, A., Abrams, J., Chen, Q., & Carrasco, M. (2025). System-Level computations
underlie visual field heterogeneity. *bioRxiv*. doi.org/10.1101/2025.09.19.677418
Yantis, S., & Jonides, J. (1996). Attentional capture by abrupt onsets: new perceptual objects or
visual masking? *Journal of Experimental Psychology: Human Perception and*
Performance, 22(6), 1505-1513.
Yeshurun, Y., & Carrasco, M. (1998). Attention improves or impairs visual performance by
enhancing spatial resolution. *Nature, 396*(6706), 72-75.

Yeshurun, Y., Montagna, B., & Carrasco, M. (2008). On the flexibility of sustained attention and
its effects on a texture segmentation task. *Vision Research, 48*(1), 80-95.
Zimmermann, E., Weidner, R., Abdollahi, R. O., & Fink, G. R. (2016). Spatiotopic adaptation in
visual areas. *Journal of Neuroscience, 36*(37), 9526-9534.
996