Both terminal misfolding and polymerisation contribute to disease-relevant phenotypic changes in cell models of α₁-antitrypsin deficiency-associated liver disease

Panayiota Stylianou[1,2]*, Kish Adoni[3]*, Mun Peak Nyon[3,4]*, Adam Cryar[3]*, Natacha O. Lee[3], Imran Haq[3,4], Jennifer A. Dickens[5], Charis-Patricia Segeritz[6], Graeme C. Smith[3], Adriana Ordonez[5], James A. Irving[3,4], David A. Lomas[3,4], Kyra-Mae Leighton[1], Jessica Beasley[1], Aleck W.E. Jones[7], S. Tamir Rashid[8], Ludovic Vallier[5], Konstantinos Thalassinos[3]§, Bibek Gooptu[1,2]§

* Joint first authors
§ Joint corresponding authors:
KT
email: k.thalassinos@ucl.ac.uk.
BG
email: bg129@leicester.ac.uk

[1] University of Leicester, Institute for Structural & Chemical Biology (LISCB) and Department of Respiratory Sciences, Henry Wellcome Building, University Road, Leicester, LE1 5HB, UK;

[2] NIHR Leicester BRC – Respiratory Theme, Glenfield Hospital, Groby Road, Leicester, LE3 9QP, UK;

[3] Institute of Structural and Molecular Biology (ISMB), Division of Biosciences, University College London, Darwin Building, Gower Street, London, WC1E 6BT, UK and School of Crystallography, Department of Biological Sciences, Birkbeck, University of London, Malet Street, London, WC1E 7HX, UK;

[4] Lungs for Living, Respiratory Medicine, Rayne Building,

[5] National School of Tropical Medicine, Department of Pediatrics, Baylor College of Medicine, One Baylor Plaza, BCM113, Houston, TX, 77030, USA; Texas Children's Hospital Center for Vaccine Development, Baylor College of Medicine, 1102 Bates Street, Houston, TX, 77030, USA.;

[6] Cambridge Institute for Medical Research, MRC/Wellcome Trust Building, Hills Road, Cambridge, CB2 0XY, UK;

[7] Department of Women & Children's Medicine, King's College London, 10th Floor, North Wing, St Thomas' Hospital, Westminster Bridge Road, London, SE1 7EH, UK;

[8] Department of Metabolism, Digestion and Reproduction, Faculty of Medicine, Imperial College, Commonwealth Building, Du Cane Road, London, W12 0NN, UK

Abstract

Polymerisation of α_1-antitrypsin within hepatocytes is considered central to pathogenesis in α_1-antitrypsin deficiency-associated liver fibrosis, most commonly seen in homozygotes for the Z (p.Glu342Lys) allele. Polymerisation proceeds via self-association of monomeric intermediate states. In parallel, >50% of synthesised Z α_1-antitrypsin is instead recognized as terminally-misfolded and degraded. It is unclear whether this contributes to Z α_1-antitrypsin deficiency-associated liver disease. We characterised the relationships between polymer formation, terminal misfolding and their cellular consequences, using ion-mobility and label-free proteomics mass spectrometry, light and electron microscopy, and cellular assays. IM-MS findings supported the relationship between polymerisation propensity of different variants and expansion of the protein consistent with opening of its β-sheet A. Proteomic analyses of well-established CHO cell models of hepatocyte handling of α_1-antitrypsin variants indicated that cellular responses to the Z mutation were surprisingly similar to those seen with the $Null_{HongKong}$ variant (NHK), which can only misfold terminally and cannot polymerise. A minor set of proteins showed increases associated with Z and not NHK α_1-antitrypsin expression, consistent with a polymer-specific response, characterized by association with increased organellar organization and vesicle-mediated transport. Conversely, proteostatic and pro-fibrotic integrin-associated pathways increased with terminal misfolding tendency of the expressed α_1-antitrypsin variant. Bioenergetic pathway changes indicated concomitant switching from oxidative to glycolytic metabolism. Cell studies further correlated fibrosis-associated behaviours with terminal misfolding rather than polymerisation. Terminal misfolding, as well as polymerisation behaviour, may therefore be important for pro-fibrotic responses including metabolic reprogramming and senescence in Z α_1-antitrypsin deficiency. Molecular therapies may prove most efficacious for associated liver disease if they address terminal misfolding as well as polymerisation consequences.

Introduction

The Z (Glu342Lys) variant of α_1-antitrypsin confers a high risk of liver fibrosis (1) leading to prevalence of clinically-significant liver disease (e.g. cirrhosis, hepatocellular carcinoma) in 10% of homozygotes (2). The heterozygote state (PiMZ) is also associated with ~8-fold increased risk relative to wild-type (M) homozygotes (PiMM) (3). The risk arises due to toxic conformational behaviour of the mutant α_1-antitrypsin polypeptide chain after translation, during and/or after folding within the hepatocyte endoplasmic reticulum (ER) (4-7). Only about 15% of synthesized Z α_1-antitrypsin is secreted and this has impaired antiprotease function (8). The loss-of-function consequences within the lung, the organ where α_1-antitrypsin function is critical, predispose to early-onset emphysema (9).

Hepatocytes from PiZZ individuals with liver disease are characterized by ER inclusion bodies composed of α_1-antitrypsin molecules, non-covalently linked into polymeric chains that entangle and precipitate (7). Other α_1-antitrypsin variants associated with deficiency are also polymerogenic (10-14). Where assessed, polymerisation propensity generally correlates with severity of disease phenotype, and inversely with ability to fold to the native state which is associated with inhibitory function. However, in the case of the Baghdad variant, refolding studies indicated it both polymerizes more readily yet folds better than Z α_1-antitrypsin (15). Polymerisation and terminal misfolding, while related processes, may therefore give rise to distinct pools of conformational behaviour (Fig. 1a).

In PiZZ individuals, polymer levels within liver cells and extracellularly within the circulation, appear to correlate with key disease features (16, 17). Moreover, polymerogenic mutants of related proteins within the serine protease inhibitor (serpin) superfamily cause a range of diseases (serpinopathies) through analogous combinations of gain- and loss-of-function effects (18). ~15% of synthesized α_1-antitrypsin is retained within the polymeric inclusions and >50% is recognized as terminally misfolded and targeted for ER associated degradation (ERAD) (4, 6). Polymer accumulation, however, does not trigger detectable unfolded protein (UPR) and ER stress responses *per se* in cell models of disease-mutant α_1-antitrypsin expression that recapitulate *in vivo* hepatocyte phenotypes, though it is associated with sensitisation to this (19). A separate 'ordered protein response' has been observed in cell models of α_1-antitrypsin deficiency (20). Polymers are also secreted extracellularly and have pro-inflammatory effects (21-23). Together these findings have led to a consensus model of α_1-antitrypsin deficiency pathogenesis where polymerisation is regarded as a central event (24). On this basis we and others have worked to develop small molecules specifically designed to block the conformational changes that lead to polymerisation (25-30). Null variants of α_1-antitrypsin, where no α_1-antitrypsin

is detectable by standard biochemical assays, is retained within the hepatocyte or secreted into the circulation, are believed to contribute to substantial risk of emphysema but not to liver disease (31).

There is a need to develop a better understanding of the mechanistic steps by which the conformational events associated with folding of the Z and other common, polymerogenic variants of α_1-antitrypsin within the ER trigger hepatic fibrosis. We previously used transcriptomic and proteomic techniques to study the effects of the Z mutation in the inducible pluripotent stem cell (iPSC)-derived hepatocyte-like (iHep) cell model of disease derived from a patient and validated our findings in *ex vivo* PiZZ liver disease tissue (32). However, this was not cross-referenced with a model of terminal misfolding alone. Changes associated with polymerisation *per se* could therefore not be distinguished from those related to terminal misfolding and high activity of the ERAD system.

This study aimed to better characterize correlations between the effects of deficiency mutations upon folding to native ensembles and polymerisation propensity of α_1-antitrypsin, and to dissect the cellular consequences of terminal misfolding of Z α_1-antitrypsin from those of its polymerisation. For the former we undertook ion mobility mass spectrometry (IM-MS). For the latter, we performed proteomic studies in the well-characterised and validated CHO cell model of hepatocyte behaviour in α_1-antitrypsin deficiency. This allowed us to compare findings in the contexts of stable transfection and inducible expression of M and Z α_1-antitrypsin but also with expression of the rarer Null$_{HongKong}$ (NHK) variant. This arises due to a frameshift mutation affecting residue 318 that introduces 15 new residues and then a premature stop codon at position 333, resulting in a 61 amino-acid C-terminal truncation (33). These changes result in terminal misfolding of 100% of synthesized α_1-antitrypsin polypeptide chains, saturating the ERAD response and triggering the UPR and ER stress. Indeed, NHK α_1-antitrypsin expression is widely used as an experimental trigger in studies characterising these responses (33-38). Expression of the Z variant of α_1-antitrypsin does not cause ER stress in the CHO cell model and similar findings have been reported in human hepatocytes. However, increased susceptibility to ER stress is observed in these cells in the context of further experimental challenge. We have proposed a "two hit" model for ER stress in response to Z α_1-antitrypsin expression, and that this susceptibility relates to the terminally-misfolded polypeptide fraction reducing reserve proteostatic capacity within the ER. We hypothesized that the toxic gain-of-function pathway attributable to Z α_1-antitrypsin polymer fraction would be defined by a proteomic signature distinct from those seen with either expression of the wild-type (M) or the NHK variant of α_1-antitrypsin.

Materials and Methods

Preparation and proteomics of CHO cell ER samples

Chinese hamster ovary (CHO)-K1 cells stably transfected with Tet-On expression systems for M, Z, or Null$_{HongKong}$ (NHK) variants of α_1-antitrypsin were cultured, and expression induced with doxycycline, as described previously (19). Cells were harvested and lysed after 5 days post-seeding (2 days post-induction). Lysates were fractionated and processed for GeLC-MS^E as described previously (39) LC-MS^E data were processed using ProteinLynx Global Server (PLGS) v2.5.2. The data processing prior to database searching included ion detection, alignment and clustering methods detailed elsewhere (40). Thresholds used for peak selection were as follows, 150 counts for low energy ion detection, 40 counts for high energy ion detection and 500 counts for precursor exact mass retention time (EMRT) integrated intensity. Samples were run on a Waters Synapt G2 with an Waters Acquity LC system.

Database searches

Data were searched using PLGS v2.5.2 against a Chinese Hamster Ovary (CHO) database (www.chogenome.org/files/CHO_refseq_protein.fasta) appended with human α_1-antitrypsin and common contaminant proteins. Carbamidomethyl (C) was set as a fixed modification while oxidation (M), acetylation (K & N-term) and deamidation (N &Q) were set as variable modifications. Precursor and fragment ion tolerances were determined automatically by PLGS. Protein identification criteria were set as the detection of at least 2 fragments ion per peptide, at least 2 fragments per protein and at least 1 peptide identification per protein. A maximum of 2 missed cleavages were allowed. The protein level false discovery rate was maintained at < 5%, estimated based upon the number of proteins identified from a randomized database. Post-processing the false discovery rate was minimized further by applying the additional criterion that for a successful identification a protein must be identified in a minimum of two biological repeats. The mass spectrometry data and associated database search results have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD002219 and 10.6019/PXD002219.

Protein quantification

Protein quantification was achieved through the Hi3 technique. Hi3 is based upon the observation that the summed intensity of the three most ionizing peptides per protein correlates with high accuracy and precision to protein concentration (43). As label-free quantification directly compares samples analysed in concurrent runs, quantification errors can be introduced by run-to-run instrument

variability. To reduce this; within-condition normalization was applied prior to quantification. This was undertaken at the peptide level using a script written in-house in R to implement locally-weighted scatter point smoothing (LOESS, Fig. 1D). Post LOESS normalization, further global normalization was applied based upon the total detected protein intensity per run, in order to account for sample preparation and loading variability across biological replicates and across conditions.

Differential expression

Pairwise comparisons of relative peptide abundances were performed for the three α_1-antitrypsin expression contexts. Differential abundance was calculated based on fold change and statistical analysis. Fold change was calculated for each comparison using the mean of Hi3 protein intensity across replicates. P values were calculated using a two-sided Student's t-test, with equal variance and performed on $\log_2$ transformed intensity data. Benjamini-Hochberg correction was applied for multiple comparisons. Proteins that were calculated to have a $\log_2$ fold change larger than 1.3 and a p value < 0.05 were defined as differentially expressed. MS^E quantitative technical variation has been previously demonstrated to be reproducible at 10-15% (39). A 1.3x $\log_2$ fold threshold therefore represents a value typically 2-3 times larger than estimated technical measurement error and compares well with more established label-based methods.

Data Processing

For quantitative analysis of proteomics data across the three conditions including control (M), Z and NHK variant CHO cells; data were filtered such that each protein must be identified in at least 1 biological replicate (and two technical replicates) of each of the three conditions. As such, proteins that were not detected in all three conditions were excluded from further analysis. IBAQ values were imputed for missing values as an average of the identified IBAQ values of that condition. Principle component analysis (PCA) was performed using PC1 (59.0%) and PC2 (9.8%) with Origin v2024b (OriginLab Corporation, Northampton, MA, USA) using the parameter settings analyze: Correlation Matrix, Number of Components to Extract: 2. Hierarchical Clustering Analysis was also performed using Origin v2024b with the mean IBAQ value of the protein for each condition (for columns and rows; Cluster Method: Group Average, Distance Type: Pearson Correlation, Number of Clusters: 1). Next, proteins were grouped according to the following IBAQ value criteria: NHK>Z>M, M>Z>NHK, Z<M<NHK and Z<NHK<M. For each criterion, the protein list was analysed using STRING v12.0 (41) with standard settings (network edges: confidence, all active interaction scores, confidence: 0.4). Networks were then exported to Cytoscape v3.10.3 (42) where functional analysis was performed for

biological process ontology and network annotations. For investigation of specific pathways, all validated proteins were input into STRING and the list of proteins associated with specific Gene Ontology Biological Processes were extracted. The $\log_2$ IBAQ of the proteins, in conditions M, Z and NHK were then visualised via a heatmap using Prism GraphPad v10.4.2.

Cell phenotyping

To assess mitochondrial morphology, cultured CHO-K1 cells expressing M, Z or NHK variants of α_1-antitrypsin mutation were incubated with the MitoTracker Red stain M7512 Red CMXRos (Life Technologies, ThermoFisher, MA, USA) as per manufacturer's instructions. Live cells were imaged on an incubated LSM710 laser point scanning confocal microscope (Carl Zeiss Ltd, Cambridge UK) and acquired with the Zen 2010 software using a 63x 1.4NA Plan Apochromat objective lens. The experiments were repeated 3 times for each cell line. For quantitative analysis of mitochondrial morphology, mitochondria were selected in an automated manner and their sphericity scores calculated (Volocity, PerkinElmer, MA, USA; Imaris, Bitplane, Zurich, Switzerland).

Senescence behaviour was evaluated by assays to quantify senescence associated-β-galactosidase expression, proliferation by BrdU uptake, and p16 expression as described previously (44, 45). Immunofluorescence studies were undertaken using antibodies against KDEL motif (ENZ-ABS679, Enzo Life Sciences), Drp1 and Mfn-1 (respectively D6C7 #8570 amd D6E2S #14739, Cell Signaling Technology), galectin-3 (sc-23938 produced by Santa Cruz Biotechnology, TX, USA), β1-integrin (ab24693, Abcam, Cambridge, UK) and p16 (ab241543, Abcam, Cambridge UK). Cells were immunostained (45) and imaging was undertaken on a Leica TCS SP5 microscope.

Results

We hypothesised that we could use mass spectrometry modalities to interrogate the disease mechanism of the Z mutation. Specifically, we aimed to study its effects upon conformational behaviour of the native state in comparison with other polymerogenic variants, and the consequences of Z α_1-antitrypsin polymerisation in cell models of disease.

Native state expansion reports polymerogenic behaviour of α_1-antitrypsin variants by ion mobility mass spectrometry

We confirmed the polymerisation tendencies of different α_1-antitrypsin deficiency variants correlated with subtle but highly-reproducible expansion of the native state (Supp. Fig. 1). Even the mild S variant showed native state expansion to a similar degree to that seen with the ~5 Å lateral expansion of β-sheet A that arises from the insertion of an extra strand (as seen in the cleaved and latent species). Z and Baghdad variants demonstrated even greater CCS expansion, such that native states were expanded in inverse relation to activation energies previously determined for transitions to the polymerogenic M* state (CCS M<Z<Baghdad, activation energies M>Z>Baghdad (15)).

Global proteomic profiles indicate high similarity in cell responses to Z and NHK variant relative to M α_1-antitrypsin expression

We went on to explore the proteomic profiles of existing cell models for the expression of M, Z and NHK α_1-antitrypsin that recapitulate their expression and subcellular behaviour in hepatocytes. Contrary to our initial hypothesis, both Z and NHK α_1-antitrypsin expression were associated with similar global proteomic profiles relative to the M protein whether assessed by principal component analysis (Fig. 1a), Volcano plot (Fig. 1b) or hierarchical cluster analysis (Fig. 1c). The data were interrogated to better understand the relationships between the observations, polymerisation and terminal misfolding.

Cellular pathways most associated with polymerisation

To first explore the set of proteins whose relative abundance was most up- or down-regulated with polymerisation, we identified proteins whose abundance was greatest in cells expressing Z α_1-antitrypsin and least in those expressing the NHK variant, and vice versa. These were termed NHK<M<Z and NHK>M>Z respectively (Fig. 2a). This strategy should down-prioritise responses associated with terminal misfolding. Relatively small subsets of proteins were identified in each case.

Interestingly, the profile associated with the order NHK<M<Z was enriched for organelle organisation and vesicle mediated transport. This may reflect changes to ER organisation described in association with polymer accumulation. The profile defined NHK>M>Z represented very few proteins.

Proteins whose expression levels followed the order M<NHK<Z, or M>NHK>Z are shown in Supp. Fig. 2a and represent changes that are qualitatively similar for both variants relative to wild-type expression. Pathway analyses for all identified protein profiles are shown in Supp. Fig. 2b.

Cellular pathways varying with terminal misfolding tendency

We next defined the sets of proteins whose expression followed the order of terminal misfolding events of α_1-antitrypsin, i.e. positively correlated: M<Z<NHK (network analysis Fig. 2b); inversely correlated: M>Z>NHK (Fig. 2c). As the representation within these profiles was unexpectedly high relative to our initial hypothesis, we validated key findings using heatmap representations and cell biology studies (Figs 3-6).

Translation, cytoskeleton and protein folding

Increases in terminal misfolding tendency of the expressed α_1-antitrypsin variant (M<Z<NHK) were associated with bi-directional changes in expression of translation-associated (e.g. ribosomal proteins) and cytoskeletal proteins (Figs. 2b). Proteins associated with protein folding and re-folding were also increased in this order. The data on these were compared with heatmap colouring (Fig. 3a) and validated by Western blot data confirming altered expression of ER chaperone proteins in equally-loaded samples of cell lysate fractions (Fig. 3b). The classically-defined ER stress response pathways are not activated in CHO cell models of Z α_1-antitrypsin expression. However, interestingly the subset of proteins associated with protein misfolding that were identified as 'ER stress response' do change similarly in cells expressing this and those expressing the NHK variant. Together these data suggest that the classic ER stress response is triggered when a threshold is reached along a spectrum of misfolding-related behaviour.

Metabolic reprogramming and mitochondrial behaviour

Proteins associated with anaerobic metabolism increased with terminal misfolding tendency (M<Z<NHK) whilst those representing aerobic respiration processes (tricarboxylic acid cycle, TCA, and oxidative phosphorylation) reduced (NHK<Z<M) as shown in Figs. 2b-c and 4a. These findings are consistent with metabolic reprogramming observed in fibrosis (46-48). Across the larger set of

mitochondrial proteins in general, hierarchical cluster analysis (lower panel) confirmed that cells expressing either misfolding Z or NHK α_1-antitrypsin resembled each other more than those expressing M (Fig. 4a). The findings were evaluated at the subcellular level in the CHO cell models. Observations were compared between the CHO cell M, Z and NHK α_1-antitrypsin overexpression models. The functional (reactive oxidant species-producing) mitochondria in these states showed different morphological phenotypes, with elongation (reducing sphericity) increasing step-wise from CHO cell models expressing M to Z to NHK α_1-antitrypsin (Fig. 4b). These changes were not associated with changes in Drp1 or mitofusin-1 levels (Supp. Fig. 3).

We validated these findings in inducible pluripotent stem cell (iPSC)-derived hepatocyte-like (iHep) cell models of PiMM and PiZZ behaviour derived from the same PiZZ donor with and without gene editing (49). The observed, functional mitochondria were visualised in iHeps, they again demonstrated elongation in the Z homozygous state, relative to the corrected M homozygous state. Such behaviour is seen as a response of healthy mitochondria to mitophagy, as a potential mechanism to evade incorporation into the phagolysosomal pathway (50-52). The activation of mitophagy by the deleterious effects of the Z mutation on α_1-antitrypsin behaviour was supported by EM studies of iHep cells. These demonstrated appearances consistent with mitophagy of grossly abnormal mitochondria in PiZZ but not PiMM cells (Fig. 4b). Together our findings indicate that α_1-antitrypsin misfolding within the ER results in mitophagy of damaged mitochondria on the one hand, and selection for functional mitochondria in more elongated species.

Conserved and α_1-antitrypsin deficiency-associated fibrotic pathways

Proteins are not annotated for specifically pro-fibrotic roles by gene ontology or network analysis descriptors. Moreover, CHO paralogs for many proteins identified in human disease pathways are not well-defined. These factors may have contributed to the low number of pro-fibrotic pathway proteins identified following a bespoke search. One notable exception was serpinH1, also known as heat shock protein (HSP)47. This protein chaperones formation of the functionally-critical triple helical structure of collagen (53), a defining component of scar tissue in all fibrosis. Such extracellular matrix proteins are generally synthesised by mesenchymal rather than cells of an epithelial lineage such as CHO cells and hepatocytes. However in fibrotic contexts epithelial cells develop mesenchymal characteristics (54). SerpinH1/HSP47 expression increased with terminal misfolding tendency of the expressed α_1-antitrypsin variant (M<Z<NHK).

Another important set of proteins whose co-expression was associated with misfolding tendency was the integrins. α- and β-integrin expression levels increased in line with misfolding tendency (M<Z<NHK). Stepwise changes were also identified in the same direction with two important proteins that interact functionally with integrins. CD98 heavy chain (CD98hc, gene identifier SLC3A2) is constitutively complexed with integrins (55). Such complexes may stabilise the pro-fibrotic TGF-β1 signalling pathway via interactions with galectin-3 and the TGF-β receptor (TGF-βR)2 (56). Interestingly galectin-3 levels were increased in the context of NHK but not Z variant α_1-antitrypsin expression. Levels of G-protein peptides (small monomeric GTPase protein) also increased with misfolding tendency (M<Z<NHK α_1-antitrypsin expression models). G-proteins such as Rho, Ras and Rab proteins activate integrin functions, with Rho-mediated pathways identified as key elements of pro-fibrotic signalling (57). Levels of the alternative integrin interactors intracellular adhesion molecule (ICAM)-1 (58), and tetraspanin (59), a human paralog of which protects against fibrosis (60) showed reciprocal decreases (M>Z>NHK). Levels of talin, that links integrins with the actin cytoskeleton and hence can connect extracellular and intracellular stiffening processes (61), increased in the context of NHK though not Z α_1-antitrypsin expression relative to expression of the wild-type M protein. To confirm pro-fibrotic behaviours suggested by the proteomic data, galectin-3:β1-integrin interactions at the cell surface were assessed in CHO cells (Fig 6). These demonstrated increased co-localisation in the order M<Z<NHK variant expression.

The increasing prevalence of protein disulphide isomerase (PDI)A4 with misfolding tendency is also of specific interest as its expression levels are correlated with liver fibrosis associated with circulating α_1-antitrypsin deficiency in Z variant homozygotes (62). This was elevated in the context of the CHO model of Z α_1-antitrypsin expression relative to the M expression model, but levels were higher still in the context of NHK α_1-antitrypsin expression. Together the proteomic findings suggest that expression of Z α_1-antitrypsin triggers intracellular responses within the ER, cytosol and cell membrane that are significantly related to terminal misfolding rather than polymerisation *per se.* The profiles of the changes between α_1-antitrypsin variant expression models supported a role of terminal misfolding as a direct stimulator of cellular responses on-pathway for profibrotic behaviour. To test these implications arising from analyses of the proteomic data, we went on to assess a range of cellular phenotypic behaviours generally associated with fibrotic disease

Senescence

To test these implications arising from analyses of the proteomic data, we next assessed senescence as a range of cellular behaviours associated with fibrotic disease (63). Based upon the proteomic

findings, we hypothesised that senescence markers would track with terminal misfolding rather than polymerisation tendencies. We therefore assessed senescence behaviours in the CHO cells expressing the different α_1-antitrypsin variants (Fig. 7). Again, these demonstrated stepwise changes supporting the hypothesis. Senescence-associated-β-galactosidase (SA-β-gal) staining increased in cells expressing Z α_1-antitrypsin, relative to those expressing M α_1-antitrypsin, and increased still further in cells expressing the NHK variant (Fig. 7a). Consistent with this, proliferation assessed by BrdU assay reduced stepwise with expression of M to Z to NHK α_1-antitrypsin (Fig. 7b). Interestingly, however, p16 staining was unchanged between cells expressing the three variants (Fig. 7c). This suggested that in these cell models, senescence had not progressed to its late phase within the timeframe of the experiment.

Discussion

The Z (Glu342Lys) mutation renders α_1-antitrypsin protein less competent at adopting a stable native fold. Instead, it is vulnerable to terminal misfolding and ERAD, and to polymerisation. The associated hepatic fibrosis has been attributed to the observed accumulation of α_1-antitrypsin polymers within hepatocyte inclusion bodies. However the findings in this paper indicate that terminally misfolded α_1-antitrypsin can drive phenotypic changes in cell models of α_1-antitrypsin that are on-pathway for fibrosis. Specifically we have identified metabolic reprogramming, senescence and expression of galectin-3, a cell surface mediator of profibrotic behaviour. This is surprising as expression of the Z variant in CHO cells, iHeps and hepatocytes is not associated with the unfolded protein response (UPR) or ER stress observed with expression of the NHK variant in cell models (20).

We and others have previously reported changes in metabolic pathways in iHeps expressing Z rather than M α_1-antitrypsin (32, 64). Metabolic reprogramming is a hallmark of fibrotic behaviour in pathological fibrosis in the lung and other organs (46-48). It is characterised by a shift from oxidative metabolic pathways to glycolytic behaviour. However, our studies are the first to suggest that in α_1-antitrypsin deficiency these may correlate with the terminally-misfolded protein load rather than solely from the polymer burden.

In our studies, this was associated with a change in mitochondrial morphology. Changes in mitochondrial sphericity are described in other protein misfolding disorders and even in models of dementia caused by misfolding and polymerisation of neuroserpin, a protein closely related to α_1-antitrypsin (65). Commonly, the pathological state is associated with fragmentation of mitochondria to smaller and hence more spherical states as a precursor to apoptosis. Recent studies of other Z homozygote iHep models of α_1-antitrypsin deficiency report similar findings to this. Our EM cellular studies presented here and previously (32) also demonstrated this behaviour. Conversely the findings from our fluorescence studies report increased elongation of observed species across large numbers of mitochondria. These differences are explained by the choice of fluorescent dye used in the studies, and hence what is reported. The studies by Kaserman *et al* used a dye that stains the mitochondrial membrane and will therefore identify all mitochondrial species (64). For our fluorescence work following up the proteomic data on bioenergetic pathways we were particularly interested in the appearances of functional mitochondria. We therefore used a marker for this, i.e. a dye that recognises non-depolarised species (preserved $\Delta\psi$m). The morphologies of these mitochondria visualised in M and Z CHO cells were validated by the observation of similar phenotypes in the

corresponding iHep models. However the EM appearances in the iHep models indicate severe co-existent mitochondrial degeneration with closely-associated evidence of early mitophagy in PiZZ compared with PiMM iHeps. These characteristics define depolarised, dysfunctional mitochondria that will not be reported with our fluorescent marker. Taken together, these findings very closely recapitulate observations of mitochondrial injury, mitophagy and elongation observed in PiZZ liver tissue with established liver disease (52). Mitochondrial dysfunction (66) and megamitochondria (67) are also observed as common features in a range of chronic liver diseases. The findings in this study demonstrate that in α_1-antitrypsin deficiency such changes are early events, closely coupled with protein misfolding, but not specifically to polymerisation, within the ER.

We note that translocation of aggregated Z α_1-antitrypsin into mitochondria has been observed in a liver-derived cell line (68). Our studies were not designed to look for this. If this were indeed polymeric material, it should not be centrally relevant to the process that we are observing not only with Z α_1-antitrypsin but also to a greater degree with expression of the NHK variant which cannot polymerise.

To explore further the possibility that terminal misfolding of α_1-antitrypsin rather than polymerisation alone correlates with cellular processes that are on-pathway for liver fibrosis we went on to evaluate markers of senescence in these cell lines. The findings indicated increased senescence behaviour with increasing misfolding, again following the pattern M<Z<NHK variant expression. Lastly, we looked at the cell surface co-localisation of galectin-3 and β1-integrin. Both proteins are common mediators of fibrosis across multiple organs including lung and liver (56, 69, 70). We have identified an increase in their close (<40 nm) co-localisation at the cell surface in epithelial cell and tissue models of early lung fibrosis in response to stimulation by the key pro-fibrotic cytokine transforming growth factor (TGF-)β1 (56). It is therefore interesting to observe similar close co-localisation in response to expression of the NHK variant compared to M α_1-antitrypsin in the CHO cell models of liver fibrosis, and that this is also observed to a lesser extent with expression of the Z mutant.

We have therefore identified a range of responses to expression of pathogenic α_1-antitrypsin mutants that can polymerise and/or undergo terminal misfolding, that increase as the latter increases, even when polymerisation cannot be achieved. From the proteomic to subcellular and functional levels these are generally on-pathway for fibrosis. Figure 8 outlines two ways in which these could arise within the different models. Firstly it is possible that the common patterns of cellular changes reported here at multiple scales with both misfolding variants could have different triggers in each

case. With the Z α_1-antitrypsin expression models, this may be the formation of polymers but not the UPR and ER stress response, since these models do not manifest this in the absence of a further proteostatic challenge ("second hit"). Conversely, in the NHK α_1-antitrypsin expression model where polymerisation cannot occur, but the UPR and ER stress response are activated, the latter would be the likeliest triggers.

The second hypothesis is that the common responses described here arise from a common primary trigger, the terminally-misfolded fraction of α_1-antitrypsin that is recognised and targeted for degradation by ERAD. This has been quantified as the fate of ~70% of all Z α_1-antitrypsin polypeptides synthesised and will account for 100% of synthesised NHK variant polypeptides (4). This is the more parsimonious explanation. It complicates previous understanding of polymerisation as the key toxic gain-of-function event in α_1-antitrypsin deficiency-associated liver disease (24), and the significance of some identified polymer-response pathways (71). On the other hand it is highly consistent with the priming of PiZZ disease models to induction of the UPR and ER stress in response to a second hit, relative to the PiMM state (19, 72). This hypothesis is also consistent with findings from hypothesis-free (exomic and proteomic) studies of the effects of the Z mutation in cell models and human subjects (62, 73). These most strongly implicated ER proteins involved in the general misfolding (protein disulphide isomerase (PDI)A4) and ERAD responses as mediators of liver toxicity and fibrosis in PiZZ α_1-antitrypsin deficiency.

The two mechanistic hypotheses need not be entirely mutually exclusive, and the contribution of terminal misfolding relative to polymerisation loads in liver disease associated with PiZZ α_1-antitrypsin deficiency deserves further clarification. The findings further indicate that the homozygote states for the NHK and other terminally misfolding Null variants of α_1-antitrypsin should not be assumed benign in terms of liver disease risk. The absence of an observed clinical association may relate to their rarity, rather than representing evidence of absence of cellular pathology.

Acknowledgements

PS received support from the National Institute for Health Research (NIHR) Leicester Biomedical Research Centre (BRC) Respiratory Theme, British Lung Foundation (BLF)/Asthma+Lung (A+L)UK and the Wellcome Trust (WT). KA and KT have received support from the WT and Alpha-1 Foundation (A1F). MPN was supported by A1F and a WT VIP award. AC was supported by a … studentship. NOL was supported by an MRC studentship. IH, JAI, DAL were supported by MRC, WT, GlaxoSmithKline, the Rosetrees Trust, the Engineering and Physical Sciences Research Council (EPSRC) and NIHR UCLH BRC and Papworth Hospital. C-PS was supported by the Children's Liver Disease Foundation. GS was supported by a Wellcome Trust ISMB DTP studentship. AO was supported by an eALTA grant. JB was supported by an MRC AIM PhD studentship. AWEJ was supported by the MRC, the Australian National Health and Medical Research Council, the Australian Mitochondrial Disease Foundation and Victorian Government Operational Infrastructure Support. STR was supported by WT, MRC, NIHR Imperial BRC, A1F, and NC3Rs. LV was supported by European Research Council (ERC), NIHR CUH BRC, and the WT/MRC Cambridge Stem Cell Institute. BG received support from NIHR Leicester BRC Respiratory Theme, BLF/A+LUK, MRC and A1F.

We are grateful to Prof Stefan Marciniak (University of Cambridge) for very helpful discussions in the development of this work. We thank the Advanced Imaging Facility (RRID:SCR_020967) at the University of Leicester for support.

References

1.   Fromme M, Schneider V, Pereira V, Hamesch K, Pons M, Reichert MC, et al. Hepatobiliary phenotypes of adults with alpha-1 antitrypsin deficiency. Gut. 2021;71:415-23.

2.   Tanash HA, Piitulainen E. Liver disease in adults with severe alpha-1-antitrypsin deficiency. J Gastroenterol. 2019;54(6):541-8.

3.   Nakanishi T, Forgetta V, Handa T, Hirai T, Mooser V, Lathrop GM, et al. The undiagnosed disease burden associated with alpha-1 antitrypsin deficiency genotypes. Eur Respir J. 2020;56:2001441.

4.   Kroeger H, Miranda E, MacLeod I, Perez J, Crowther DC, Marciniak SJ, et al. Endoplasmic reticulum-associated degradation (ERAD) and autophagy cooperate to degrade polymerogenic mutant serpins. J Biol Chem. 2009;284:22793-802.

5.   Gooptu B, Dickens JA, Lomas DA. The molecular and cellular pathology of α1-antitrypsin deficiency. Trends in Molecular Medicine. 2014;20(2):116-27.

6.   Teckman JH, Burrows J, Hidvegi T, Schmidt B, Hale PD, Perlmutter DH. The proteasome participates in degradation of mutant α_1-antitrypsin Z in the endoplasmic reticulum of hepatoma-derived hepatocytes. J Biol Chem. 2001;276:44865-72.

7.   Lomas DA, Evans DL, Finch JT, Carrell RW. The mechanism of Z α_1-antitrypsin accumulation in the liver. Nature. 1992;357:605-7.

8.   Lomas DA, Evans DL, Stone SR, Chang W-SW, Carrell RW. Effect of the Z mutation on the physical and inhibitory properties of α1-antitrypsin Biochemistry. 1993;32:500-8.

9.   Gooptu B, Ekeowa UI, Lomas DA. Mechanisms of emphysema in α1-antitrypsin deficiency: molecular and cellular insights. Eur Respir J. 2009;34(2):475-88.

10.   Mahadeva R, Chang W-SW, Dafforn T, Oakley DJ, Foreman RC, Calvin J, et al. Heteropolymerisation of S, I and Z α1-antitrypsin and liver cirrhosis. J Clin Invest. 1999;103:999-1006.

11.   Dafforn TR, Mahadeva R, Elliott PR, Sivasothy P, Lomas DA. A kinetic mechanism for the polymerisation of α1-antitrypsin. J Biol Chem. 1999;274:9548-55.

12.   Giacopuzzi E, Laffranchi M, Berardelli R, Ravasio V, Ferrarotti I, Gooptu B, et al. Real-world clinical applicability of pathogenicity predictors assessed on SERPINA1 mutations in alpha-1-antitrypsin deficiency. Hum Mutat. 2018;39(9):1203-13.

13.   Fra AM, Gooptu B, Ferrarrotti I, Miranda E, Scabini R, Ronzoni R, et al. Three new alpha$_1$-antitrypsin deficiency variants help to define a C-terminal region regulating conformational change and polymerization. PLoS One. 2012;7(6):e38405, doi:10.1371/journal.pone.0038405.

14.   Miranda E, Perez J, Ekeowa UI, Hadzic N, Kalsheker N, Gooptu B, et al. A novel monoclonal antibody to characterize pathogenic polymers in liver disease associated with α1-antitrypsin deficiency. Hepatology. 2010;52:1078-88.

15.   Haq I, Irving JA, Saleh AD, Dron L, Regan-Mochrie GL, Motamedi-Shad N, et al. Deficiency mutations of alpha1-antitrypsin. Effects upon folding, function and polymerization. Am J Respir Cell Mol Biol. 2016;54:71-80.

16.   Mela M, Smeeton W, Davies SE, Miranda E, Scarpini C, Coleman N, et al. Alpha-1 antitrypsin polymer load correlates with hepatocyte senescence, fibrosis stage and liver-related mortality. Chronic Obstructive Pulmonary Disease. 2020;7:151-62.

17.   Tan L, Dickens JA, DeMeo DL, Miranda E, Perez J, Rashid ST, et al. Circulating polymers in α1-antitrypsin deficiency. Eur Respir J. 2014;43:1501-4.

18.   Gooptu B, Lomas DA. Conformational pathology of the serpins: themes, variations and therapeutic strategies. Annu Rev Biochem. 2009;78:147-76.

19.   Ordóñez A, Snapp EL, Tan L, Miranda E, Marciniak SJ, Lomas DA. Endoplasmic reticulum polymers impair luminal protein mobility and sensitise to cellular stress in α_1-antitrypsin deficiency. Hepatology. 2013;57:2049-60.

20.   Marciniak SJ, Ordonez A, Dickens JA, Chambers JE, Patel V, Dominicus CS, et al. New Concepts in Alpha-1 Antitrypsin Deficiency Disease Mechanisms. Annals of the American Thoracic Society. 2016;13 Suppl 4:S289-96.

21.   DeMeo DL, Banos M, Perez J, Tan L, Barker AF, Campbell E, et al. Circulating polymers of alpha-1 antitrypsin in PiZZ subjects. Am J Resp Crit Care Med. 2012;185:A4361.

22.   Mahadeva R, Atkinson C, Li J, Stewart S, Janciauskiene S, Kelley DG, et al. Polymers of Z $\alpha1$-antitrypsin co-localise with neutrophils in emphysematous alveoli and are chemotactic *in vivo*. Am J Pathol. 2005;166(2):377-86.

23.   Parmar JS, Mahadeva R, Reed BJ, Farahi N, Cadwallader K, Bilton D, et al. Polymers of $\alpha1$-antitrypsin are chemotactic for human neutrophils: a new paradigm for the pathogenesis of emphysema. Am J Respir Cell Mol Biol. 2002;26:723-30.

24.   Gooptu B, Lomas DA. Polymers and inflammation: disease mechanisms of the serpinopathies. J Exp Med. 2008;205(7):1529-34.

25.   Lomas DA, Hurst JR, Gooptu B. Update on alpha-1 antitrypsin deficiency: New therapies. J Hepatol. 2016;65(2):413-24.

26.   Nyon MP, Gooptu B. Therapeutic targeting of misfolding and conformational change in α_1-antitrypsin deficiency. Future Medicinal Chemistry. 2014;6:1047-65.

27.   Patschull AOM, Gooptu B, Ashford P, Daviter T, Nobeli I. *In silico* assessment of potential druggable pockets on the surface of α_1-antitrypsin conformers. PLoS One. 2012;7(5):e36612. doi:10.1371/journal.pone.0036612.

28.   Chang Y-P, Mahadeva R, Patschull AOM, Nobeli I, Ekeowa UI, McKay AR, et al. Targeting serpins in high-throughput and structure-based drug design. Methods in Enzymology. 2011;501:139-75.

29.   Nyon MP, Prentice T, Day J, Kirkpatrick J, Sivalingam G, Levy GR, et al. An integrative approach combining ion mobility mass spectrometry, X-ray crystallography, and nuclear magnetic resonance spectroscopy to study the conformational dynamics of $\alpha1$-antitrypsin upon ligand binding. Protein Science. 2015;24:1301-12.

30.   Mallya M, Phillips, R,L., Saldanha, S.A., Gooptu, B., Leigh Brown, S.C., Termine, D.J., Shirvani, A.M., Wu, Y., Sifers, R.N., Abagyan, R., Lomas, D.A. Small molecules block the polymerization of Z alpha1-antitrypsin and increase the clearance of intracellular aggregates. J Med Chem. 2007;50(22):5357-63.

31.   Lee JH, Brantly M. Molecular mechanisms of alpha 1-antitrypsin null alleles. Respir Med. 2000;94Suppl C:S7-11.

32.   Segeritz CP, Rashid ST, de Brito MC, Serra MP, Ordonez A, Morell CM, et al. hiPSC hepatocyte model demonstrates the role of unfolded protein response and inflammatory networks in alpha1-antitrypsin deficiency. J Hepatol. 2018;69(4):851-60.

33.   Sifers RN, Brashears-Macatee S, Kidd VJ, Muensch H, Woo SLC. A frameshift mutation results in a truncated α_1-antitrypsin that is retained within the rough endoplasmic reticulum. J Biol Chem. 1988;263(15):7330-5.

34.   Pan S, Wang S, Utama B, Huang L, Blok N, Estes MK, et al. Golgi localization of ERManI defines spatial separation of the mammalian glycoprotein quality control system. Mol Biol Cell. 2011;22(16):2810-22.

35.   Wu Y, Swulius MT, Moremen KW, Sifers RN. Elucidation of the molecular logic by which misfolded α1-antitrypsin is preferentially selected for degradation. Proc Natl Acad Sci USA. 2003;100(14):8229-34.

36.   Cabral CM, Liu Y, Moremen KW, Sifers RN. Organizational diversity among distinct glycoprotein ER-associated degradation programs. Molec Biol Cell. 2002;13:2639-50.

37.   Zhong Y, Shen H, Wang Y, Yang Y, Yang P, Fang S. Identification of ERAD components essential for dislocation of the null Hong Kong variant of alpha-1-antitrypsin (NHK). Biochemical and biophysical research communications. 2015;458(2):424-8.

38.   Greenblatt EJ, Olzmann JA, Kopito RR. Derlin-1 is a rhomboid pseudoprotease required for the dislocation of mutant alpha-1 antitrypsin from the endoplasmic reticulum. Nat Struct Mol Biol. 2011;18(10):1147-52.

39.   Patel VJ, Thalassinos K, Slade SE, Connolly JB, Crombie A, Murrell JC, et al. A comparison of labeling and label-free mass spectrometry-based proteomics approaches. J Proteome Res. 2009;8(7):3752-9.

40.   Geromanos SJ, Vissers JP, Silva JC, Dorschel CA, Li GZ, Gorenstein MV, et al. The detection, correlation, and comparison of peptide precursor and product ions from data independent LC-MS with data dependant LC-MS/MS. Proteomics. 2009;9(6):1683-95.

41.   Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, et al. The STRING database in 2023: protein-protein association networks and functional enrichment analyses for any sequenced genome of interest. Nucleic Acids Res. 2023;51(D1):D638-D46.

42.   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a software environment for integrated models of biomolecular interaction networks. Genome Res. 2003;13(11):2498-504.

43.   Silva JC, Gorenstein MV, Li GZ, Vissers JP, Geromanos SJ. Absolute quantification of proteins by LCMSE: a virtue of parallel MS acquisition. Mol Cell Proteomics. 2006;5(1):144-56.

44.   Gonzalez-Gualda E, Baker AG, Fruk L, Munoz-Espin D. A guide to assessing cellular senescence in vitro and in vivo. FEBS J. 2021;288(1):56-80.

45.   Stylianou P, Clark K, Gooptu B, Smallwood D, Brightling CE, Amrani Y, et al. Tensin1 expression and function in chronic obstructive pulmonary disease. Sci Rep. 2019;9(1):18942.

46.   Horn P, Tacke F. Metabolic reprogramming in liver fibrosis. Cell Metab. 2024;36(7):1439-55.

47.   Zhao X, Kwan JYY, Yip K, Liu PP, Liu FF. Targeting metabolic dysregulation for fibrosis therapy. Nat Rev Drug Discov. 2020;19(1):57-75.

48.   Bueno M, Calyeca J, Rojas M, Mora AL. Mitochondria dysfunction and metabolic reprogramming as drivers of idiopathic pulmonary fibrosis. Redox Biol. 2020;33:101509.

49.   Rashid ST, Corbineau S, Hannan N, Marciniak SJ, Miranda E, Alexander GJ, et al. Modeling inherited metabolic disorders of the liver using human induced pluripotent stem cells. J Clin Invest. 2010;120:3127-36.

50.   Rambold AS, Kostelecky B, Lippincott-Schwartz J. Together we are stronger: fusion protects mitochondria from autophagosomal degradation. Autophagy. 2011;7(12):1568-9.

51.   Gomes LC, Di Benedetto G, Scorrano L. During autophagy mitochondria elongate, are spared from degradation and sustain cell viability. Nat Cell Biol. 2011;13(5):589-98.

52.   Teckman JH, An JK, Blomenkamp K, Schmidt B, Perlmutter D. Mitochondrial autophagy and injury in the liver in alpha 1-antitrypsin deficiency. Am J Physiol Gastrointest Liver Physiol. 2004;286:G851-62.

53.   Dafforn TR, Della M, Miller AD. The molecular interactions of heat shock protein 47 (hsp 47) and their implications for collagen biosynthesis. J Biol Chem. 2001;276:49310-9.

54.   Wynn TA. Cellular and molecular mechanisms of fibrosis. J Pathol. 2008;214(2):199-210.

55.   Prager GW, Feral CC, Kim C, Han J, Ginsberg MH. CD98hc (SLC3A2) interaction with the integrin beta subunit cytoplasmic domain mediates adhesive signaling. J Biol Chem. 2007;282(33):24477-84.

56.   Calver JF, Parmar NR, Harris G, Lithgo RM, Stylianou P, Zetterberg FR, et al. Defining the mechanism of galectin-3-mediated TGF-beta1 activation and its role in lung fibrosis. J Biol Chem. 2024;300(6):107300.

57.   Tsou PS, Haak AJ, Khanna D, Neubig RR. Cellular mechanisms of tissue fibrosis. 8. Current and future drug targets in fibrosis: focus on Rho GTPase-regulated gene transcription. Am J Physiol Cell Physiol. 2014;307(1):C2-13.

58.   Longhurst CM, Jennings LK. Integrin-mediated signal transduction. Cell Mol Life Sci. 1998;54(6):514-26.

59.   Termini CM, Gillette JM. Tetraspanins Function as Regulators of Cellular Signaling. Front Cell Dev Biol. 2017;5:34.

60.   Tsujino K, Takeda Y, Arai T, Shintani Y, Inagaki R, Saiga H, et al. Tetraspanin CD151 protects against pulmonary fibrosis by maintaining epithelial integrity. Am J Respir Crit Care Med. 2012;186(2):170-80.

61.   Goult BT, Yan J, Schwartz MA. Talin as a mechanosensitive signaling hub. The Journal of cell biology. 2018;217(11):3776-84.

62.   Karatas E, Raymond A-A, Leon C, Dupuy J-W, Di-Tommaso S, Senant N, et al. Hepatocyte proteomes reveal the role of protein disulfide isomerase 4 in alpha 1-antitrypsin deficiency. JHEP Rep. 2021;3:100297.

63.   O'Reilly S, Tsou PS, Varga J. Senescence and tissue fibrosis: opportunities for therapeutic targeting. Trends Mol Med. 2024;30(12):1113-25.

64.   Kaserman JE, Werder RB, Wang F, Matte T, Higgins MI, Dodge M, et al. Human iPSC-hepatocyte modeling of alpha-1 antitrypsin heterozygosity reveals metabolic dysregulation and cellular heterogeneity. Cell Rep. 2022;41(10):111775.

65.   D'Acunto E, Gianfrancesco L, Serangeli I, D'Orsi M, Sabato V, Guadagno NA, et al. Polymerogenic neuroserpin causes mitochondrial alterations and activates NFkappaB but not the UPR in a neuronal model of neurodegeneration FENIB. Cell Mol Life Sci. 2022;79(8):437.

66.   Chu Q, Gu X, Zheng Q, Wang J, Zhu H. Mitochondrial Mechanisms of Apoptosis and Necroptosis in Liver Diseases. Anal Cell Pathol (Amst). 2021;2021:8900122.

67.   Shami GJ, Samarska IV, Koek GH, Li A, Palma E, Chokshi S, et al. Giant mitochondria in human liver disease. Liver Int. 2023;43(11):2365-78.

68.   Khodayari N, Wang RL, Oshins R, Lu Y, Millett M, Aranyos AM, et al. The Mechanism of Mitochondrial Injury in Alpha-1 Antitrypsin Deficiency Mediated Liver Disease. Int J Mol Sci. 2021;22(24).

69.   Reed NI, Jo H, Chen C, Tsujino K, Arnold TD, DeGrado WF, et al. The alphavbeta1 integrin plays a critical in vivo role in tissue fibrosis. Sci Transl Med. 2015;7(288):288ra79.

70.   Li LC, Li J, Gao J. Functions of galectin-3 and its role in fibrotic diseases. J Pharmacol Exp Ther. 2014;351(2):336-43.

71.   Davies MJ, Miranda E, Roussel BD, Kaufman RJ, Marciniak SJ, Lomas DA. Neuroserpin polymers activate NF-kappaB by a calcium signaling pathway that is independent of the unfolded protein response. J Biol Chem. 2009;284(27):18202-9.

72.   Lawless MW, Greene CM, Mulgrew A, Taggert CC, O'Neill SJ, McElvaney NG. Activation of endoplasmic reticulum-specific stress responses associated with the conformational disease Z α1-antitrypsin deficiency. J Immunol. 2004;172:5722-6.

73.   Joly P, Vignaud H, Di Martino J, Ruiz M, Garin R, Restier L, et al. ERAD defects and the HFE-H63D variant are associated with increased risk of liver damages in Alpha 1-Antitrypsin Deficiency. PLoS One. 2017;12:e0179369.

Figure Legends

Figure 1

Alpha$_1$-antitrypsin misfolding and polymerisation in deficiency variants

a) **Model of effects of pathogenic mutations on α_1-antitrypsin folding behaviour**

Energy diagram illustrating folding trajectories for (left to right) wild-type (M) α_1-antitrypsin, Z, S, Baghdad, and NHK variants. Polymerogenic mutations lower the activation energy for the native-to-polymer transition and may favour the population of near-native, polymerogenic intermediate states during or after folding within the ER. For the NHK variant, folding to native or ordered polymer states is not possible due to C-terminal deletion, and so it populates a disordered, terminally-misfolded state that is prone to ER-associated degradation (ERAD). >50% of synthesised Z α_1-antitrypsin polypeptides also undergo ERAD, yet the Baghdad variant which is more polymerogenic refolds more efficiently. These observations suggest that partitioning can arise between terminally-misfolded and polymerogenic states during folding of the Z variant.

b-d) **Proteomic analysis of CHO cells expressing M, Z or NHK α_1-antitrypsin demonstrates high global similarity for expression of Z and NHK variants relative to M** by: b) principal component (PC), c) Volcano plot, and d) hierarchical cluster analyses. In c) vertical dashed lines indicate threshold for confident discrimination of change (log$_2$-fold change=1.3), and horizontal dashes delineate the p=0.05 probability threshold following Benjamini-Hochberg correction for multiple comparisons. Each point plotted represents an identified protein, with log$_2$ mapping of the number of unique peptides used for its assignment (spot radius), and of signal intensity (colour heatmapping). In d) low-to-high detected levels are depicted in purple-to-orange spectrum between … and …

Figure 2

Network analyses of functional clusters identify distinct response repertoires to polymerisation and misfolding.

Medium stringency STRING analyses of proteins differentially co-expressed with Z and NHK variants, relative to observations with expression of M α_1-antitrypsin in CHO cells.

a) Changes observed with increasing (left, NHK<M<Z) and decreasing (right, Z<M<NHK) ratios of polymerisation:ERAD behaviours of α_1-antitrypsin variants.

b) Protein increases observed with increasing terminal misfolding behaviours (M<Z<NHK)

c) Protein increases observed with decreasing terminal misfolding behaviours

Supp. Fig. 2

a) Qualitatively similar changes observed in CHO cells expressing NHK and Z relative to findings with M α_1-antitrypsin expression but not following order of terminal misfolding or polymerisation tendencies. Increasing expression with M<NHK<Z (left) and Z<NHK<M (right).

b) Pathway analyses for all profiles shown in Fig. 2.

Figure 3

Increases in proteostatic response M<Z<NHK

a) Heatmap depiction of proteins associated with protein folding processes and ER stress response responses. Proteins are listed in order of increasing levels in cells expressing M α_1-antitrypsin. Purple depicts the lowest levels detected and orange the highest. Thus in this case, both sets of proteins show generally increased co-expression with the α_1-antitrypsin variants in the order M<Z<NHK.

b) Changes in ER chaperone (detected by anti-KDEL western blot) repertoire co- expressed in responses to variants confirm the expected increase in these proteins M<Z<NHK, across a subcellular fractionation gradient.

Supp. Fig. 3

SDS-PAGE of cell lysates for CHO cell models expressing M, Z and NHK α_1-antitrypsin probed with western blots using antibodies to recognize Drp1 and Mfn-1. Sample loading demonstrated by β-actin loading shown below.

Figure 4

Bio-energetic and mitochondrial behaviour in CHO models

a) Heatmap depiction (format as outlined for Fig. 3) of co-expression of "Anaerobic" (e.g. glycolytic) and "Aerobic" (tricarboxylic acid, oxidative phosphorylation) metabolic pathway proteins with different α_1-antitrypsin variants. Abundance of anaerobic pathway proteins generally increased (M<Z<NHK) whilst aerobic pathway proteins generally decreased (M>Z>NHK) with terminal-misfolding tendency. More complex patterns were seen with the global set of mitochondrial proteins ("Mitochondrial", upper). HCA confirmed the co-expression profiles with the misfolding variants were more similar to each other than to the profile observed with M α_1-antitrypsin expression ("Mitochondrial", lower).

b) Observed changes in morphology of functional mitochondria in association with metabolic shifts observed in Fig. 4a in representative views (confocal fluorescence microscopy), and quantified by

sphericity score. Upper panel shows illustrative examples of the observed morphologies of functional mitochondria associated most closely with expression of each α_1-antitrypsin variant. Such mitochondria are shown with red fluorescent signal in the upper panel, and false coloured by sphericity score in the lower panel (red high, to blue, low) with overall sphericity data from field views quantified in box and whisker plot.

Figure 5

Grossly-disordered mitochondrial morphology in iHep models of Z relative to M α_1-antitrypsin expression and *ex vivo*.

a) Representative views of cellular mitochondria in iHep cells derived from the same donor with (MM) and without (ZZ) correction of the pathogenic α_1-antitrypsin mutation, false coloured by sphericity score as in Fig. 4. Box and whisker plot quantitation

b) Mitochondrial changes observed by SEM, with double membrane structures (white arrows) wrapping around grossly abnormal mitochondria, consistent with mitophagy.

Figure 6

a) Heatmap representation of co-expression levels of proteins involved in fibrosis pathways and liver disease risk in PiZZ individuals in cells expressing M, Z or NHK α_1-antitrypsin. Formatted as in Figure 3.

b) CHO cell co-expression of galectin-3 and β1-integrin with M, Z or NHK α_1-antitrypsin with overlapping co-immunofluorescence correlated by Mander's and Pearson's co-efficients.

Figure 7

Senescence readouts assessed for CHO cells expressing M, Z or NHK α_1-antitrypsin, n=3:

a) SA-β-galactosidase stain intensity quantitation (above) and representative micrographs (lower panel). Increases indicate increased senescence behaviour.

b) BrdU proliferation assay, reductions indicate increased senescence behaviour.

c) p16 marker of late senescence.

Figure 8

Different models for the activation of pro-fibrotic behaviour. Model A includes the combination of polymer accumulation and UPR/ER stress to drive a two-trigger pathway. Model B involves the

stimulation of pro-fibrotic activity from a common primary trigger, the terminally-misfolded fraction of α_1-antitrypsin that is recognised and targeted for degradation by ERAD.