Ecophylogenetic patterns of rhizosphere bacterial community assembly in *Pisum* spp. (Fabaceae, Fabeae) reveal strong plant-mediated ecological filtering

Victor Angot[1*], Vincent Pailler[2], Abdelhamid Kebieche[3], Elodie Belmonte[2], Virginie Bourion[1], Yanis Bouchenak-Khelladi[1]

[1] Université Bourgogne Europe, Institut Agro, INRAE, UMR Agroécologie, 21000 Dijon, France

[2] INRAE, UMR GDEC, Gentyane, Clermont-Ferrand, France

[3] Institut Agro, Université d'Angers, INRAE, IRHS, SFR 4207 QUASAV, Angers, France

[*] Corresponding author: victor.angot@inrae.fr

Abstract

Plant-microorganisms interactions are among the oldest biotic relationships and play a fundamental role in shaping biological systems. These associations involve several and diverse species, each evolving on different timescales. Understanding these interactions requires approaches that integrate both ecological community dynamics and evolutionary processes, which drive the adaptation of plants and microorganisms. We investigated bacterial community assembly dynamics associated with four diverse *Pisum* spp. accessions grown in greenhouse conditions on soil. Bacterial DNA was extracted from bulk soil, rhizosphere, rhizoplane, and endosphere microhabitats across three plant growth stages, followed by full-length 16S rRNA gene sequencing. Bacterial communities varied in diversity, composition and structure across microhabitats and growth stages. Ecophylogenetic analyses, that is integrating community ecology dynamics into a phylogenetic framework, indicated strong host-filtering, with community assembly across space and time being structured by phylogenetic constraints. This illustrates the role of the plant in creating and shaping distinct ecological niches, where selective recruitment favors specific and closely related lineages. Our findings suggest that an ecophylogenetic approach provides valuable insights into plant-microbiota dynamics by integrating ecological and evolutionary processes, thereby offering a powerful perspective to investigate the co-adaptation of plants and their associated microbiota.

Key words: plant-microorganisms interactions, *Pisum* spp., plant development, ecophylogenetic, ecological filtering

Acknowledgments: The authors thank the 4PMI greenhouse platform team at INRAe Dijon for help in designing the experiment and the ECP team (LEGae, INRAE Dijon) for valuable assistance in the laboratory. The authors wish to thank Nadine Praeg and Mathieu Barret for critically evaluating and providing very valuable comments that improved significantly the manuscript. This research was funded by INRAE and the Agence National de la Recherche (ANR, France).

Author contributions: Conceptualization: Y.B.K.; Methodology: V.A., Y.B.K., V.B.; Investigation: A.K., E.B, V.B., Y.B.K.; Data curation: V.A., V.P.; Formal analysis: V.A., V.P.; Visualization: V.A.; Writing – original draft: V.A.; Writing – review & editing: V.A., Y.B.K.; Supervision: Y.B.K.; Funding acquisition: Y.B.K.

Declaration of competing interest: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Introduction

Biotic interactions are responsible for major evolutionary changes, such as increased genetic diversity within populations or triggering adaptive changes leading to ecological speciation, across multiple spatial and temporal scales. These complex interactions, that can involve two or thousands of species and occur within or across trophic levels, contribute to crucial ecological patterns that structure biological systems (Schemske *et al.*, 2009; Spicer & Woods, 2022). One of the most intricate and complex biotic interaction is the association between plants and microorganisms (Lyu *et al.*, 2021), that is among the oldest relationship that may have originated around 460-475 Mya (Brundrett, 2002), and represents an ideal model for studying host-microbiota association dynamics.

Legumes (Fabaceae) is a highly diverse family of angiosperms, many of which form symbiotic associations with nitrogen-fixing bacteria and play an important role in fixing atmospheric nitrogen. Globally, grain legumes (tribe Fabeae) were estimated to fix a total of 35.5 million tons of nitrogen every year (Herridge et al., 2022). Among these, *Pisum* spp. include one of the first domesticated plant species (Zohary & Hopf, 1973). *Pisum* is divided into two species : *P. fulvum* and *P. sativum* which is subdivided in 3 sub-species : *P. sativum* subsp. *sativum*, *P. sativum* subsp. *elatius* and *P. sativum* subsp. *abyssinicum* (Smýkal et al., 2011). While the symbiotic relationship of *Pisum* spp. with *Rhizobia* is well-documented (Bourion *et al.*, 2018; Hawkins & Oresnik, 2022), the complex interactions with non-symbiotic microorganisms remain largely unknown, whereas the rhizosphere microbiota play a crucial role in shaping *Rhizobia*-legumes interactions (Han *et al.*, 2020). The non-symbiotic bacteria could also improve the effectiveness of *Rhizobia*-inoculants by enhancing their ability to colonize, survive, and persist in the soil, as well as their capacity to interact with other microorganisms (O'Callaghan et al., 2022; Kong & Liu, 2022). Therefore, it is critical to understand the assembly rules of the whole bacterial community, beyond *Rhizobium* symbiosis. Plant-microorganisms interactions are governed by a complex 'chemical dialogue' through the production of exudates by the plant (primary and secondary metabolites) (Duc *et al.*, 2022). Some exudates, such as organic acids (Rekha *et al.*, 2018), amino-acids (Feng *et al.*, 2018) and glucose (Zhang *et al.*, 2015), have chemo-attractive properties that facilitate bacterial colonization (Chaparro *et al.*, 2013). Other compounds, such as antimicrobial peptides, have antibiotic properties that prevent the colonization of pathogenic microorganisms (Pacheco-Cano *et al.*, 2018). In addition, root architecture and root hairs structure further modulate the assembly of the rhizosphere microbial communities (Kawasaki *et al.*, 2016; Robertson-Albertyn *et al.*, 2017) by altering the physical microhabitat properties. Therefore, through these chemical and physical modifications, plants can strongly influence the composition of rhizosphere microbial communities, exceeding the influence of the original soil properties (Tkacz *et al.*, 2020). In return, these shaped communities can enhance plant fitness through nitrogen fixation, nutrient acquisition, and increased resistance to biotic and abiotic stresses (Bhattacharyya & Jha, 2012; Peix *et al.*, 2015; de la Fuente Cantó *et al.*, 2020; Yang *et al.*, 2021).

The assembly dynamics of microbial communities are shaped by multiple factors, requiring multifactorial analyses that account for both spatial and temporal dimensions. Different plant compartments (referred hereafter as plant-associated microhabitats), such as the rhizosphere, rhizoplane and endosphere contain distinct microbial communities (Cregger et al., 2018; Gupta et al., 2021). Therefore, the identification of microbial associations and enrichment or impoverishment dynamics through an ecological gradient (i.e., from the rhizosphere to the endosphere) is crucial in order to characterize the host-mediated ecological filters associated with different plant-associated microhabitats (Zhang et al., 2017; Iven et al., 2022). Studies have shown that plant-associated microbial communities are specific to their host species and can shift through plant growth (Chaparro *et al.*, 2014; Edwards *et al.*, 2015; Shade *et al.*, 2017), suggesting both spatial and temporal filtering. While it is demonstrated that assembly of soil microbial communities is determined by biotic interactions such

as mutualism (Goberna et al., 2014; Romdhane et al., 2022), facilitation or competition (Faust & Raes,
2012), the association and assembly dynamics throughout plant-associated microhabitats and plant
growth remain largely unknown (Moroenyane *et al.*, 2021). To decipher these spatio-temporal
dynamics, an ecophylogenetic framework is particularly effective because it computes ecological
dynamics while considering phylogenetic relatedness to infer underlying biotic interactions. For
instance, insights from human microbiomes indicate that microbial recruitment can follow an
underdispersion assembly pattern (Darcy *et al.*, 2020) (i.e. taxa are more likely to be recruited if a
phylogenetically close relative was previously established). Whether similar processes govern the
mode and tempo of microbial recruitment in plant-associated microhabitats remains largely
speculative. The use of ecophylogenetic frameworks is therefore needed to reveal association patterns
with host plants (Wang & Sugiyama, 2020). The assembly of these community may be attributed to
phylogenetic niche conservatism (Losos, 2008; Pyron et al., 2015), when closely related species inhabit
similar ecological niches. However, instances of competitive exclusion have been shown, such as in
floral nectar (Peay *et al.*, 2012), and the balance between these processes may be influenced by
resource availability overtime (Pekkonen et al., 2013). Similarly, the intensity of plant rhizodeposition
and its ability to shape ecological niches for bacteria could affect the pattern of phylogenetic
association. The specialization of microbiota on the plant's physiological needs and on its
microhabitats suggest that plant-microbiota interactions prioritize specific functions over particular
microbial taxa (Castellano-Hinojosa & Strauss, 2021). Since phylogenetic relatedness often predicts
functional similarity (Burns & Strauss, 2011; Martiny *et al.*, 2015; Anacker & Strauss, 2016), community
assembly is expected to be phylogenetically structured (Aguirre de Cárcer, 2019). Therefore, an
ecophylogenetic approach can help identify whether plant-mediated ecological filtering shapes
microbial assembly across spatial and temporal scales.
111
To explicitly link spatial and temporal host-mediated ecological filters with phylogenetic community
assembly, we conducted a greenhouse experiment with four *Pisum spp.* accessions to study the
composition and structure of plant-associated belowground bacterial communities at different growth
stages using both ecological and phylogenetic approaches. Our hypotheses were (i) that bacterial
community diversity, composition and phylogenetic diversity differ across the plant-associated
microhabitat and reflect a gradient of host-filtering, (ii) that bacterial community composition and
phylogenetic structure shift across plant growth stages, reflecting dynamic changes in host-filtering,
and (iii) that host-filtering results in non-random phylogenetic assembly patterns.

Material and Methods

Experimental design and data collection

Four *Pisum* accessions from the Pea Genetic Resource Centre at INRAE Dijon were selected: (i) one
Pisum fulvum (IPIP10023) and (ii) one *Pisum sativum* subsp*. elatius* (IPI200733) as two wild species
accessions, one *Pisum sativum* subsp*. abyssinicum* 'ATAR' (IPI200596) and one *Pisum sativum* subsp*.
sativum* 'CAMEOR' (IPI200293) as two accessions representative of domestication or selection events.
The seeds were stored in a cold room prior to sowing. Seeds from *Pisum fulvum* and *Pisum sativum*
subsp*. elatius* were scarified using a scalpel to enable germination. Soil was collected at Auxonne, Côte
d'Or, France (47.185N, 5.403E) from an organic farm, air-dried (Han *et al.*, 2026), and sieved to 4 mm.
Soil nitrate, ammonium, and total nitrogen, as well as water content and bulk density, were assessed
at the laboratory "SADEF Agronomy & Environment" (Appendix 1). The experiment was conducted as
a completely randomized design and was carried out in a climate-controlled glasshouse located at
INRAE Dijon from February to May 2023. The greenhouse was equipped with a temperature control
system that maintained a temperature of 20 °C during the day and 15 °C at night, and a 16 hours

photoperiod. Seeds were sown into sterilized 2 L pots filled with sieved soil. Two seeds were sown in each pot to ensure the successful germination of at least one. Pots were top-watered with osmotic water with an automatic dripper system until germination. Afterwards, pots were watered daily with 100 mL of a nutritive solution (MgSO4 + 7H2O 1 mL/L, CaCl2 0.63 mL/L, K2SO4 1.39 mL/L, NaCl 0.1 mL/L, Oligo 1 mL/L, and iron versenate 1 mL/L; pH 6.5). Plants were harvested at three successive growth stages: the first stage (S1) was the vegetative stage (5 to 6 leaf nodes, 16-24 days), the second stage (S2) was the beginning of flowering (appearance of the first flower, 31-56 days); and the third stage (S3) was the beginning of seed filling (the first pod containing internal seeds at least 8mm in length, 36-64 days). The aim was to have nine plants for each growth stage and for each accession (nine replicates x three growth stages x four accessions) resulting in a total of 108 plants, with 27 plants per accession. However, not all seeds germinated, with only 25/27 for *Pisum sativum* subsp*. abyssinicum,* 21/27 *for P. sativum* subsp*. sativum, 15/27 for P. fulvum and 15/27 for P. sativum* subsp*. elatius*, which led to a total of 76 plants. Eight pots were left unsown to provide bulk soil samples without any association with *Pisum*, serving as controls. After retrieving the plant, the roots were transferred to a sterile centrifuge tube with 35 mL of 1X PBS and kept at 4 °C. Roots were initially sonicated for one minute and 30 seconds to collect rhizosphere microorganisms (R), then transferred to a new tube containing 35 mL of 1X PBS and subjected to a second round of sonication for 10 minutes to isolate rhizoplane microorganisms (Rp). The R and Rp tubes were then centrifuged at 5000 x g for 10 minutes, and 250 mg of the resulting pellet was sampled for DNA extraction. Additionally, 400 mg of the central root pivot (the first 3 cm below seed insertion) were sampled and grinded to retrieve the endosphere (E) microbial communities. At the start of the experiment, and at each stage, bulk soil samples were collected as controls and analyzed using the same procedure as for R and Rp. Some plant nodules were also sampled, however, due to the low number of plants that formed nodules, the results concerning this microhabitat could not be included in this study. In total, we collected 260 samples that encompassed all plant-associated microhabitats at three growth stages and 32 bulk soils.

Bacterial DNA extraction, amplification and sequencing

DNA was extracted from all samples using DNeasy® PowerSoil® Pro Kits (QIAGEN®, Hilden, Germany) following the manufacturer's protocol and stored at -20 °C. Concentrations and purity of the extractions were assessed using a Nanodrop2000™ spectrophotometer (Thermo Fischer Scientific™, Waltham, USA). DNA extraction yielded insufficient concentration and purity in 11 samples, which were therefore excluded from amplification. The amplification of full-length 16S rRNA gene (V1-V9 regions) was performed according to the PacBio® procedure and general best practices recommendations. Single-molecule Real-time long reads sequencing was performed at Gentyane Sequencing Platform (Clermont-Ferrand, France) with a PacBio Sequel II Sequencer (Pacific Biosciences®, Menlo Park, CA, USA). Amplicons were normalized at 5 ng/µL, pooled and purified with 1.3X SMRTbell cleanup beads. The SMRTbell library was prepared using a SMRTbell prep kit 3.0, following the "Preparing multiplexed amplicon libraries using SMRTbell prep kit 3.0" protocol. A Femto Pulse (Agilent Technologies®, Santa Clara, CA, USA) assay was used to assess the fragments size distribution. Pools were carried into the enzymatic reactions to remove the single-strand overhangs and to repair any damage that may be present on the DNA backbone. An A-tailing reaction followed by the overhang adapter ligation was conducted to generate the SMRTbell template. After nuclease treatment, pools were cleanup with 1.3X SMRTbell cleanup beads. The SMRTbell libraries were quality inspected and quantified on a Femto Pulse (Agilent Technologies®) and a Qubit fluorimeter with Qubit dsDNA HS reagent Assay kit (Life Technologies®). A ready-to-sequence SMRTbell Polymerase Complex was created using a Binding Kit 3.1 (PacBio®). The PacBio Sequel II instrument was programmed to load a 150pM library and sequenced in CCS mode on a PacBio SMRTcell 8M, with the Sequencing Plate 2.0

(Pacific Biosciences®, Menlo Park, CA, USA) with 0.5 hour of pre-extension time and acquiring one
movie of 10 hours.

Four SMRTcells were used and the CCS reads were generated from raw data (i.e., subreads) with "ccs"
and each sample was demultiplexed from each pool using "lima". These were performed using "SMRT-
Link v13.0" software (Pacific Biosciences®, Menlo Park, CA, USA). Two hundred and forty-nine samples
were analyzed in a fastq file format. The DADA2 package (Callahan *et al.*, 2016) in R was used to
perform taxonomic affiliations of the samples (see supplementary material). Primers were removed
and strands orientation was conducted. Then, reads were filtered according to their sizes (between
1,000 and 1,600bp), followed by a dereplication and denoising steps to obtain the amplicon sequence
variants (ASV, mean length = 1,449 bp). Taxonomic affiliations were done using the "silva_nr99_v138"
database (Yilmaz *et al.*, 2014), and ASVs present in less than 2 samples were removed. The total
number of ASVs retrieved was 19,747 (ccs reads data available in Genbank; SAMN45805597-
SAMN45805845). Chloroplast, mitochondrial or eukaryote 16S rRNA sequences were removed, which
resulted in 18,335 ASVs for 249 samples (see supplementary material,
https://doi.org/10.5281/zenodo.18400087).

Community ecology and statistical analyses

The statistical analyses were performed using RStudio software (version 4.3.0). To ensure data quality
and that sampling of the different microhabitats was properly conducted, outlier samples from the
initial set of 249 were identified using two approaches. Firstly, richness of the samples was calculated,
and 62 outliers with aberrantly low or high richness were removed (likely due to technical issues that
do not reflect true biological variation). Secondly, a Bray-Curtis dissimilarity matrix was constructed,
and Nonmetric Multidimensional Scaling (NMDS) analysis was performed for each microbial
community compartment using the "vegan" package (Oksanen *et al.*, 2022). The NMDS coordinates
were used to calculate Mahalanobis distances between samples within each microhabitat. Outliers
were defined as points with Mahalanobis distances exceeding the 99% Chi-squared threshold
(Appendix 2), resulting in the removal of seven additional outliers and a final dataset of 180 samples.
Finally, we removed ASVs representing less than 0.01% of total sequences to reduce computational
load for phylogenetic inference, leading to a set of 1,590 ASVs. The same filtered dataset was then
used for ecological and ecophylogenetic analyses, with minimal impact on results and biological
interpretation (Appendix 3). The completeness of community coverage was confirmed using
rarefaction curves (Schöler *et al.*, 2017). Shannon index (H) and species richness (S) were calculated
using the *diversity* function in the "vegan" package. Analyses of the alpha diversity indexes were
conducted using multiple Kruskal-Wallis test from the package "agricolae" (de Mendiburu, 2023).
Community structure was analyzed and visualized using distance-based redundancy analysis with Bray-
Curtis distances (db-RDA, capscale, "vegan", 10,000 permutations; Oksanen *et al.*, 2022). This
technique has proven to be useful when measuring multispecies responses to structured multifactorial
designs (Legendre & Anderson, 1999). We assessed the homogeneity of multivariate dispersions by
applying permutation tests to the residuals of the db-RDA model (Appendix 4). Differential abundance
analyses were performed using "edgeR" (Robinson *et al.*, 2010). ASV count data were filtered to retain
taxa with counts per million normalization > 1 in at least two samples and normalized by scaling each
sample to the median library size. Differential abundance was assessed by contrasting bulk soil with
each plant-associated microhabitat, and S1 to S3 within each plant-associated microhabitats
($|log2FC| > 1$, $p < 0.05$). Data and scripts required to reproduce the analyses are available at
https://doi.org/10.5281/zenodo.18400087.

Bacterial phylogenetic inference

In order to infer a bacterial phylogeny including ASVs we retrieved from our samples, we aligned 1,590 16S rRNA sequences using MUSCLE 3.6 (Edgar, 2004) and used Gblocks 0.91 (Castresana, 2000) to eliminate poorly aligned positions. The resulting matrix included 1,590 taxa and 1,195 characters (16S rRNA sequence matrix is available with the online version of the manuscript). Phylogenetic inferences were performed using BEAST 2.6.4 (Bouckaert *et al.*, 2019). The substitution model used was GTR +I+G (Yang, 1994); three runs of 100 million chains were performed, sampling every 5,000 generations. We checked for convergence and viewed statistics for each run using TRACER 1.7.2 (Rambaut *et al.*, 2018). After combining the three runs and removing 40% of burn-in samples, we summarized the results using a maximum clade credibility (MCC) tree.

We computed the Rao's quadratic entropy (FD_Q Rao, 1982, "SYNCA", Debastiani & Pillar, 2012), a weighted measure of diversity integrating both phylogenetic distances and relative abundances of bacterial taxa, to assess the effects of host-filtering on phylogenetic diversity. Temporal phylogenetic dispersion model of bacterial communities was estimated by calculating the empirical dispersion parameter (*D*) for each plant-associated microhabitats, following the method developed by Darcy *et al.* (2020). The dispersion parameter tracks how newly recruited taxa relate phylogenetically to previously observed taxa over time, allowing to test whether new recruits are more closely or more distantly related to earlier community members than expected by chance (Darcy *et al.*, 2020). All above analyses were performed using the MCC tree.

Results

Bacterial community composition and structure

Bacterial communities were characterized from a total of 180 samples. To ensure a good coverage of the bacterial communities, we generated rarefaction curves using the full set of 18,335 ASVs, illustrating the increase in observed ASVs as a function of sequencing depth (Appendix 5). All samples reached an asymptote, indicating that the bacterial communities were fully sampled in all microhabitats and that rarefaction was unnecessary. Community diversity did not differ significantly among the four *Pisum* accessions, and although community structure varied significantly, the explained variance was low (< 5%, Appendix 6). Therefore, all genotypes were combined for subsequent analyses. Bacterial community diversity varied across the different plant-associated microhabitats (Figure 1A, B). Specific richness was the highest in the bulk soil (bulk, median S = 547), followed by the rhizosphere (R, S = 399) and rhizoplane (Rp, S = 389), with the lowest richness observed in the endosphere (E, S = 70; Figure 1A, $p < 0.001$). The Shannon diversity index significantly decreased across the microhabitats, from 5.73 in bulk to 5.42 in R, 5.14 in Rp, and 3.62 in E (Figure 1B, $p < 0.001$). Although specific richness and Shannon diversity were not significantly influenced by growth stages, it is noteworthy that the initial community exhibited the highest diversity, with S = 660 and H = 6.05 in the initial bulk soil (i.e., soil sampled before sowing) (Appendix 7).

Bacterial communities' structure differs across plant-associated microhabitats, growth stages and host *Pisum* genotypes ($R^2 = 0.50$, $p < 0.001$, Figure 1C). Changes in structure were best explained by microhabitats (26.60%, $p < 0.001$, Figure 1C). There was no overlap of 95% CI ellipses between microhabitats that were not in close physical proximity (Figure 1C). The highest overlap in 95% CI was observed between R and Rp, while the overlaps with the other microhabitats were small. Growth stages had a lower but significant effect on bacterial communities' structure (4.21%, $p < 0.001$; Figure 1C). The host genotype had a significant but marginal influence on the associated bacterial communities (3.95%, $p < 0.001$; Figure 1C). Time had no significant effect on the bacterial communities in the bulk soil ($p = 0.15$; Appendix 8A). In contrast, growth stages had a significant and increasingly

pronounced effect on the bacterial communities in the plant-associated microhabitats, explaining
respectively 9.31%, 10.92%, and 13.27% of the variance in R, Rp and E ($p < 0.001$; Appendix 8B, C, D).

The taxonomic composition of bacterial communities varied across plant-associated microhabitats
(Figure 1D, Appendix 9A, B, C). Bulk soil communities were the most heterogeneous, with low-
abundance orders ("others") accounting for 39.6% of relative abundance, alongside Bacillales (27.2%),
Rhizobiales (12.2%), Vicinamibacterales (6.7%), Micrococcales (4.8%) and Burkholderiales (4.7%)
(Figure 1D). In the rhizosphere, community composition remained broadly similar to bulk soil but
showed an enrichment in Burkholderiales (15.6%), mainly driven by ASVs affiliated to *Massilia*,
Rhizobacter, *Ramlibacter* and *Noviherbaspirillum* (Appendix 10A), together with a decrease in
Bacillales (19.5%). In the rhizoplane, communities shifted toward a dominance of Burkholderiales
(28.9%), Micrococcales (11%) and Pseudomonadales (5.9%), driven by *Delftia*, *Massilia*,
Paeniglutamicibacter and several *Pseudomonas* species (Appendix 10B), while Bacillales (9.8%) and
Vicinamibacterales (2%) decreased. In the endosphere, communities were dominated by
Burkholderiales (43.8%) and Rhizobiales (26.8%), with Bacillales, Micrococcales and Vicinamibacterales
nearly absent. The highest increase in the endosphere is for *Rhizobium leguminosarum* (Appendix 10C).
Notably, several orders that were nearly absent in bulk soil (<1%), including Pseudomonadales,
Flavobacteriales, Xanthomonadales, Sphingomonadales and Cellvibrionales, progressively increased in
relative abundance across plant-associated microhabitats (Figure 1D).

Community composition within plant-associated microhabitats also varied across growth stages (S1 to
S3), whereas no significant differences were observed in bulk soil (Figure 1D, Appendix 9D, E, F, G). In
the rhizosphere, Bacillales (14.3 to 26.5%), Rhizobiales (11.7 to 14.7%) and Micrococcales (4.9 to 8.7%)
increased, driven mainly by *Bacillus* species, *Paeniglutamicibacter sulfureus* and *Paenarthrobacter*
aurescens (Appendix 10D), while Burkholderiales (17.6 to 10.6%), Flavobacteriales (7.8 to 0.9%) and
Cellvibrionales (4.0 to 0.1%) declined. In the rhizoplane, Micrococcales (2.9 to 14.2%) and
Pseudomonadales (5.8 to 6.7%) increased, driven primarily by *Paeniglutamicibacter sulfureus* and
Pseudomonas species (Appendix 10E), whereas Burkholderiales (38.8 to 25.5%), Cellvibrionales (3.9 to
0.4%), Flavobacteriales (4.7 to 2.1%) and Xanthomonadales (4.1 to 1.7%) decreased. In the
endosphere, the only marked increase was observed for Rhizobiales (22.8 to 31.8%), mainly driven by
Rhizobium leguminosarum (Appendix 10F), while Burkholderiales (54.6 to 35.2%), Cellvibrionales (3.7
to 1.2%) and Pseudomonadales (4.7 to 2.9%) declined.

Temporal and spatial recruitment in a phylogenetic context

Using our phylogenetic tree of 1,590 bacterial ASVs, we examined the phylogenetic diversity of
bacterial communities across the different microhabitats and calculated the phylogenetic dispersion
parameter (D) to assess the phylogenetic patterns of bacterial recruitment throughout growth stages
in each plant-associated microhabitat. Negative D values indicate underdispersion, corresponding to
the preferential recruitment of phylogenetically close taxa from the established community over time,
whereas positive values indicate overdispersion, reflecting the recruitment of phylogenetically more
distant taxa. Phylogenetic diversity was the highest in the bulk with $FD_Q = 0.67$ ($p < 0.001$, Figure 2A)
and progressively decreased across the plant-associated compartments with $FD_Q = 0.65$ in R, $FD_Q = 0.60$
in Rp and $FD_Q = 0.44$ in E ($p < 0.001$, Figure 2A). The phylogenetic diversity significantly varied across
growth stages only within the endosphere, rising from $FD_Q = 0.40$ at S1 to $FD_Q = 0.44$ at S2 and
$FD_Q = 0.46$ at S3 ($p < 0.01$, data not shown). The phylogenetic dispersion parameter differs across plant-
associated microhabitats (Figure 2B). Bacterial communities in R exhibited overdispersion pattern with
$D > 0$ ($D = 0.34$, $p < 0.05$, Figure 2B). In Rp, D was lower than in R ($p < 0.1$) and not different to $D = 0$,
indicating a neutral phylogenetic dispersion pattern ($D = 0.03$, $p > 0.05$, Figure 2B). Communities in E

showed underdispersion, with D = -0.40, significantly lower than in Rp and different from the neutral
model ($p < 0.05$, Figure 2B), indicating that closely related ASVs have more probabilities of being
recruited throughout growth stages.

Discussion

Microbial community assembly

Bacterial communities were specific to the different microhabitats. These offer distinct ecological
niches, which is reflected by the significant changes in abundance, composition, and structure of
bacterial communities. The highest diversity was observed in the bulk soil and gradually decreased
from the rhizosphere to the endosphere. Plant-associated microhabitats act as ecological filters along
the spatial gradient, where communities from the bulk soil are shaped in the rhizosphere, which are
further influenced by the rhizoplane, ultimately resulting in only a small part of these communities
being present in the endosphere. Community composition exhibited a clear successional shift from
bulk soil to the endosphere, marked by a progressive enrichment of several orders nearly absent in
bulk soil, including Pseudomonadales, Flavobacteriales, Xanthomonadales, Sphingomonadales, and
Cellvibrionales. Notably, Burkholderiales became dominant along the gradient, while orders such as
Bacillales and Vicinamibacterales decreased. Within the Burkholderiales, potential PGP genera such as
Massilia (Dai *et al.*, 2020; Angot *et al.*, 2025) and *Noviherbaspirillum* (Barros *et al.*, 2022) drove these
shifts. Within Pseudomonadales, *Pseudomonas* species, which are common in the endosphere and
nodules of legumes (Mayhood & Mirza, 2021; Hossain *et al.*, 2023) and may influence the outcome of
root-nodule symbiosis (Crosbie *et al.*, 2022) drove the signal. The increase in Rhizobiales in the
endosphere was mainly driven by *Rhizobium leguminosarum*, likely due to the spatial proximity with
root nodules.

Host growth stages also influenced the structure and composition of bacterial communities within
plant-associated microhabitats, while bulk soil communities remained stable. In the rhizosphere, we
observed an increase in Bacillales driven by potential PGP *Bacillus species* (El-Esawi *et al.*, 2018; Blake
et al., 2021), and Micrococcales, mainly driven by *Paeniglutamicibacter sulfureus*, alongside a decline
in Burkholderiales, Flavobacteriales, and Cellvibrionales. The rhizoplane exhibited a pronounced
increase in Micrococcales, primarily due to *Paeniglutamicibacter sulfureus*, and a slight increase in
Pseudomonadales (*Pseudomonas spp.*), while Burkholderiales and several other orders decreased. This
marked enrichment of *Paeniglutamicibacter* at later stages is notable. Although its direct role in plant
interactions is unknown, its capacity for hydrocarbon degradation in soils (Brzeszcz *et al.*, 2024)
suggests a potential response to root metabolites, given the structural similarities between
hydrocarbons and many plant secondary metabolites (Singer, 2006; Rohrbacher & St-Arnaud, 2016).
In the endosphere, the most significant temporal change was a further increase in Rhizobiales,
dominated by *Rhizobium leguminosarum*, consistent with the hypothesis of dispersion from root
nodules, whose abundance rises with plant development (Mortier *et al.*, 2012).

The stability of bulk soil communities, compared to plant-associated microhabitats, highlights the
importance of host's influence on the temporal dynamics of bacterial communities. Even if further
work is necessary to understand the mechanistic processes driving plant-bacterial recruitment
throughout plant development, numerous studies have illustrated the influence of plant growth stages
on the non-symbiotic microbial communities (Qiao *et al.*, 2017; Dai *et al.*, 2019; Larrouy *et al.*, 2023).
These variations seem to be tied to shifts in the plant's requirements throughout its development, as
well as to changes in the quality and quantity of rhizodeposits produced by the host (Hütsch et al.,
2002; Houlden et al., 2008; Chaparro et al., 2013). Indeed, rhizodeposition is dependent on plant
carbon allocation (Shamoot et al., 1968) which varies through growth stages (Morin et al., 2022).

Carbon allocation in the roots is important during the vegetative stage (Voisin et al., 2003b). At the flowering stage, the concentration of carbon increases to supply shoot apical meristem and induces floral transition (Cho et al., 2018). During the reproductive stage, carbon is remobilized from the leaves, stem and roots to fill the seeds (Weber et al., 1997). Because the microorganisms associated with *Pisum* spp*.* can enhance its nitrogen acquisition, the dynamics of the rhizosphere microbiota are linked to the plant's nitrogen needs, which vary at different growth stages (Du et al., 2023). During the vegetative phase, nitrogen requirements are significant to ensure plant growth and to build reserves in the above-ground parts (Voisin et al., 2007). During seed filling stages, nitrogen acquisition is reduced and the plant uses its nitrogen stocks to fill the seeds (Voisin et al., 2003a). Both carbon and nitrogen fluxes variations impact the rhizosphere microbiota; however, the influence of developmental stages on the recruitment of microbial communities remains uncertain (Biget et al., 2024), especially given the potential confounding effect of the time since sowing (Dibner et al., 2021).

Our findings reveal that plants host diverse bacterial communities that vary across the microhabitats they provide and throughout their growth stages. These shifts in community composition and structure reflect the plant's capacity to act as an ecological filter, creating distinct niches that select for particular bacterial taxa. At the same time, microorganisms also contribute to shaping their environments, feeding back into the construction and modification of plant-associated microhabitats. Given the complexity of integrating spatial and temporal factors in host-microbiota systems, moving toward an eco-evolutionary framework appears particularly valuable. Such an approach would help disentangle the mechanisms driving community recruitment, persistence, and changes across plant development.

Phylogenetic constraints on bacterial community assembly

Our bacterial phylogeny, based on full-length 16S rRNA gene sequencing of 1,590 ASVs, is largely congruent with recent genomic references (Zhu *et al.*, (2019), Appendix 11). All Bacterial clades were retrieved (Appendix 11). Proteobacteria (with Gamma- and Alphaproteobacteria as sister groups), Desulfobacterota (sister to Proteobacteria), and Bacteroidota. Firmicutes, Actinobacteria, Chloroflexi, and Cyanobacteria are also monophyletic. The placement of Acidobacteriota differs, possibly due to our denser sampling (128/1,590 vs 56/10,575 taxa in Zhu et al. (2019)). Planctomycetota, not sampled in Zhu et al. (2019), are placed as sister to Thermotogae, which is sister to Cyanobacteria in Zhu et al. (2019), supporting our topology (Rensink *et al.*, 2020). These results suggest that, although recombination and horizontal gene transfer processes affecting the 16S rRNA gene can weaken the concordance of 16S rRNA phylogenies (Hassler *et al.*, 2022), using full-length sequencing enables the construction of phylogenies that are congruent with phylogenomic inferences. In addition, 16S rRNA gene phylogenies were shown to reflect the ecological and functional traits of soil bacterial communities (Borenstein *et al.*, 2008). The phylogenetic diversity (PD) varied across microhabitats, with the highest PD being observed in the bulk soil, followed by a gradual decrease across the rhizosphere, rhizoplane, and endosphere. Host filtering does not randomly reduce bacterial diversity but rather selects for specific, phylogenetically related lineages, likely by intensifying niche constraints. Goldford *et al.* (2018), using ex situ cultivation of microbial communities from different habitats, showed that communities were assembled at family-level, suggesting that this pattern may reflect functional selection linked to competitive advantages. Our results support the prediction that phylogenetically conserved set of traits allow specific bacteria lineages to pass through ecological constraints imposed by the host plants (Aguirre de Cárcer, 2019). This suggests that selective recruitment is correlated with bacterial functions required for plant development, as previously reported by Mougel *et al.* (2006). In addition, the temporal phylogenetic pattern of community assembly varied across the plant-associated microhabitats. The model we used considers the likelihood that newly recruited taxa are similar to closely related taxa previously recruited (Darcy *et al.*, 2020). Overdispersion observed in the rhizosphere suggests that interspecific competition among

closely related bacteria plays a crucial role in shaping the community (Nemergut *et al.*, 2013). Closely related taxa, which share similar niches, are competitively excluded, leading to phylogenetic overdispersion of the bacterial community. The rhizosphere, as an open and nutrient-rich microhabitat, could provide a diverse array of ecological niches. These niches can support the coexistence of distantly related bacteria, probably with distinct ecological functions. Conversely, the underdispersion pattern observed in the endosphere indicates the opposite process: the bacterial community is not mainly structured by competition among closely related bacteria but rather by strong host-filtering (Mayfield & Levine, 2010), as previously observed for mycorrhizal fungal community assembly (Frew *et al.*, 2023). The endosphere, as a highly selective and closed microhabitat, imposes significant host-mediated environmental constraints that can be physical (barriers of root cells) or chemical (plant-derived chemical compounds; Lumibao *et al.* (2020)). It shows that closely related bacteria are able to coexist in a community (Webb *et al.*, 2002), with specific functional traits, often conserved among phylogenetically related taxa. The pattern of phylogenetic underdispersion in the endosphere is consistent with ecological selection acting on phylogenetically conserved traits (Webb *et al.*, 2002; Gerhold *et al.*, 2015). If the traits being selected for are horizontally transferred between bacteria lineages with no connection with their relatedness, we would not observe any phylogenetic signature (Darcy *et al.*, 2020). Several studies have demonstrated that plant hosts actively select for specific bacterial communities or strains. (Bertin *et al.*, 2003; Bulgarelli *et al.*, 2013; Sasse *et al.*, 2018; Bourion *et al.*, 2018; Xiong *et al.*, 2021). Over evolutionary timescales, this persistent host-mediated ecological filtering could have driven the evolutionary history of plant-associated bacterial communities. Such reciprocal interactions, which impact the fitness of both partners, could be a keystone in the evolutionary trajectory of land plants and a mark of coevolution (Fitzpatrick et al., 2018).

Altogether, our results illustrate the strong influence of plants on their associated soil bacterial communities, shaping both their ecological and evolutionary dynamics across spatial and temporal scales. Bacterial communities were highly specific to underground plant-associated microhabitats, and their assembly was phylogenetically structured, with the host filtering taxa according to their evolutionary relatedness. However, investigating the heritability of bacterial communities is crucial for understanding whether these associations are shaped by (i) horizontal transfer from the environment to plant-associated microhabitats, (ii) plant-specific traits, or (iii) vertical transmission across plant generations. A limitation of this study is the absence of direct measurements of plant physiology and root exudates. Consequently, while we document clear spatio-temporal patterns in community assembly, it cannot be directly linked to bacterial plant growth promotion or plant recruitment mechanisms. Future work integrating host phenotyping, exudate profiling, and microbial functional potential is essential for a mechanistic understanding of the plant-bacteria interactions and their evolutionary outcomes.

References

Aguirre de Cárcer D. **2019**. A conceptual framework for the phylogenetically constrained assembly of microbial communities. *Microbiome* **7**: 142.

Anacker BL, Strauss SY. **2016**. Ecological similarity is related to phylogenetic distance between species in a cross-niche field transplant experiment. *Ecology* **97**: 1807–1818.

Angot V, Thiour-Mauprivez C, Schellenberger R, Lamboeuf M, Crépin O, Garmyn D, Blouin M, Jacquiod S. **2025**. Alleviation of water-deficit inhibition of plant growth by rhizosphere microbiota conditioning. *New Phytologist* **247**: 2914–2926.

Barros FM do R, Pedrinho A, Mendes LW, Freitas CCG, Andreote FD. **2022**. Interactions between Soil Bacterial Diversity and Plant-Parasitic Nematodes in Soybean Plants. *Applied and Environmental Microbiology* **88**: e00963-22.

Bertin C, Yang X, Weston LA. **2003**. The role of root exudates and allelochemicals in the rhizosphere. *Plant and Soil* **256**: 67–83.

Bhattacharyya PN, Jha DK. **2012**. Plant growth-promoting rhizobacteria (PGPR): emergence in agriculture. *World Journal of Microbiology & Biotechnology* **28**: 1327–1350.

Biget M, Mony C, Wang T, Ling N, Miteul A, Jambon O, Causse-Védrines R, Michon-Coudouel S, Hervé M, Chable V, *et al.* **2024**. Intra- and inter-annual changes in root endospheric microbial communities of grapevine are mainly deterministic. *Plant and Soil* **494**: 217–233.

Blake C, Christensen MN, Kovács ÁT. **2021**. Molecular Aspects of Plant Growth Promotion and Protection by Bacillus subtilis. *Molecular plant-microbe interactions: MPMI* **34**: 15–25.

Borenstein E, Kupiec M, Feldman MW, Ruppin E. **2008**. Large-scale reconstruction and phylogenetic analysis of metabolic environments. *Proceedings of the National Academy of Sciences* **105**: 14482–14487.

Bouckaert R, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M, Gavryushkina A, Heled J, Jones G, Kühnert D, De Maio N, *et al.* **2019**. BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. *PLoS computational biology* **15**: e1006650.

Bourion V, Heulin-Gotty K, Aubert V, Tisseyre P, Chabert-Martinello M, Pervent M, Delaitre C, Vile D, Siol M, Duc G, *et al.* **2018**. Co-inoculation of a Pea Core-Collection with Diverse Rhizobial Strains Shows Competitiveness for Nodulation and Efficiency of Nitrogen Fixation Are Distinct traits in the Interaction. *Frontiers in Plant Science* **8**: 2249.

Brundrett MC. **2002**. Coevolution of roots and mycorrhizas of land plants. *New Phytologist* **154**: 275–304.

Brzeszcz J, Steliga T, Ryszka P, Kaszycki P, Kapusta P. **2024**. Bacteria degrading both n-alkanes and aromatic hydrocarbons are prevalent in soils. *Environmental Science and Pollution Research International* **31**: 5668–5683.

Bulgarelli D, Schlaeppi K, Spaepen S, Ver Loren van Themaat E, Schulze-Lefert P. **2013**. Structure and functions of the bacterial microbiota of plants. *Annual Review of Plant Biology* **64**: 807–838.

Burns JH, Strauss SY. **2011**. More closely related species are more ecologically similar in an experimental test. *Proceedings of the National Academy of Sciences* **108**: 5302–5307.

Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. **2016**. DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods* **13**: 581–583.

Castellano-Hinojosa A, Strauss SL. **2021**. Insights into the taxonomic and functional characterization of agricultural crop core rhizobiomes and their potential microbial drivers. *Scientific Reports* **11**.

Castresana J. **2000**. Selection of Conserved Blocks from Multiple Alignments for Their Use in Phylogenetic Analysis. *Molecular Biology and Evolution* **17**: 540–552.

Chaparro JM, Badri DV, Bakker MG, Sugiyama A, Manter DK, Vivanco JM. **2013**. Root exudation of phytochemicals in Arabidopsis follows specific patterns that are developmentally programmed and correlate with soil microbial functions. *PloS One* **8**: e55731.

Chaparro JM, Badri DV, Vivanco JM. **2014**. Rhizosphere microbiome assemblage is affected by plant development. *The ISME Journal* **8**: 790–803.

Cho L-H, Pasriga R, Yoon J, Jeon J-S, An G. **2018**. Roles of Sugars in Controlling Flowering Time. *Journal of Plant Biology* **61**: 121–130.

Cregger MA, Veach AM, Yang ZK, Crouch MJ, Vilgalys R, Tuskan GA, Schadt CW. **2018**. The Populus holobiont: dissecting the effects of plant niches and genotype on the microbiome. *Microbiome* **6**: 31.

Crosbie DB, Mahmoudi M, Radl V, Brachmann A, Schloter M, Kemen E, Marín M. **2022**. Microbiome profiling reveals that Pseudomonas antagonises parasitic nodule colonisation of cheater rhizobia in Lotus. *The New Phytologist* **234**: 242–255.

Dai Y, Yang F, Zhang L, Xu Z, Fan X, Tian Y, Wang T. **2020**. Wheat-associated microbiota and their correlation with stripe rust reaction. *Journal of Applied Microbiology* **128**: 544–555.

Dai L, Zhang G, Yu Z, Ding H, Xu Y, Zhang Z. **2019**. Effect of Drought Stress and Developmental Stages on Microbial Community Structure and Diversity in Peanut Rhizosphere Soil. *International Journal of Molecular Sciences* **20**: 2265.

Darcy JL, Washburne AD, Robeson MS, Prest T, Schmidt SK, Lozupone CA. **2020**. A phylogenetic model for the recruitment of species into microbial communities and application to studies of the human microbiome. *The ISME journal* **14**: 1359–1368.

Dibner RR, Weaver AM, Brock MT, Custer GF, Morrison HG, Maignien L, Weinig C. **2021**. Time outweighs the effect of host developmental stage on microbial community composition. *FEMS Microbiology Ecology* **97**: fiab102.

Du C, Xu R, Zhao X, Liu Y, Zhou X, Zhang W, Zhou X, Hu N, Zhang Y, Sun Z, *et al.* **2023**. Association between host nitrogen absorption and root-associated microbial community in field-grown wheat. *Applied Microbiology and Biotechnology* **107**: 7347–7364.

Duc NH, Vo HTN, van Doan C, Hamow KÁ, Le KH, Posta K. **2022**. Volatile organic compounds shape belowground plant-fungi interactions. *Frontiers in Plant Science* **13**: 1046685.

Edgar RC. **2004**. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* **32**: 1792–1797.

Edwards J, Johnson C, Santos-Medellín C, Lurie E, Podishetty NK, Bhatnagar S, Eisen JA, Sundaresan V. **2015**. Structure, variation, and assembly of the root-associated microbiomes of rice. *Proceedings of the National Academy of Sciences* **112**: E911–E920.

El-Esawi MA, Alaraidh IA, Alsahli AA, Alamri SA, Ali HM, Alayafi AA. **2018**. Bacillus firmus (SW5) augments salt tolerance in soybean (Glycine max L.) by modulating root system architecture, antioxidant defense systems and stress-responsive genes expression. *Plant physiology and biochemistry: PPB* **132**: 375–384.

Faust K, Raes J. **2012**. Microbial interactions: from networks to models. *Nature Reviews Microbiology* **10**: 538–550.

Feng H, Zhang N, Du W, Zhang H, Liu Y, Fu R, Shao J, Zhang G, Shen Q, Zhang R. **2018**. Identification of Chemotaxis Compounds in Root Exudates and Their Sensing Chemoreceptors in Plant-Growth-Promoting Rhizobacteria Bacillus amyloliquefaciens SQR9. *Molecular Plant-Microbe Interactions®* **31**: 995–1005.

Fitzpatrick CR, Copeland J, Wang PW, Guttman DS, Kotanen PM, Johnson MTJ. **2018**. Assembly and ecological function of the root microbiome across angiosperm plant species. *Proceedings of the National Academy of Sciences of the United States of America* **115**: E1157–E1165.

Frew A, Heuck MK, Aguilar-Trigueros CA. **2023**. Host filtering, not competitive exclusion, may be the main driver of arbuscular mycorrhizal fungal community assembly under high phosphorus. *Functional Ecology* **37**: 1856–1869.

de la Fuente Cantó C, Simonin M, King E, Moulin L, Bennett MJ, Castrillo G, Laplaze L. **2020**. An extended root phenotype: the rhizosphere, its formation and impacts on plant fitness. *The Plant Journal* **103**: 951–964.

Gerhold P, Cahill Jr JF, Winter M, Bartish IV, Prinzing A. **2015**. Phylogenetic patterns are not proxies of community assembly mechanisms (they are far better). *Functional Ecology* **29**: 600–614.

Goberna M, García C, Verdú M. **2014**. A role for biotic filtering in driving phylogenetic clustering in soil bacterial communities. *Global Ecology and Biogeography* **23**: 1346–1355.

Goldford JE, Lu N, Bajić D, Estrela S, Tikhonov M, Sanchez-Gorostiaga A, Segrè D, Mehta P, Sanchez A. **2018**. Emergent simplicity in microbial community assembly. *Science (New York, N.Y.)* **361**: 469–474.

Gupta R, Anand G, Gaur R, Yadav D. **2021**. Plant-microbiome interactions for sustainable agriculture: a review. *Physiology and Molecular Biology of Plants: An International Journal of Functional Plant Biology* **27**: 165–179.

Han X, Cuartero J, Koppe V, Nohl S, Sneyders A, Vancampenhout K, Frey B, Frossard A. **2026**. Air-drying of soil preserves microbial and faunal eDNA abundance and diversity regardless of land-use type or management intensity. *Soil Biology and Biochemistry* **215**: 110082.

Han Q, Ma Q, Chen Y, Tian B, Xu L, Bai Y, Chen W, Li X. **2020**. Variation in rhizosphere microbial communities and its
association with the symbiotic efficiency of rhizobia in soybean. *The ISME journal* **14**: 1915–1928.

Hassler HB, Probert B, Moore C, Lawson E, Jackson RW, Russell BT, Richards VP. **2022**. Phylogenies of the 16S rRNA gene
and its hypervariable regions lack concordance with core genome phylogenies. *Microbiome* **10**: 104.

Hawkins JP, Oresnik IJ. **2022**. The Rhizobium-Legume Symbiosis: Co-opting Successful Stress Management. *Frontiers in Plant*
Science **12**: 796045.

Herridge DF, Giller KE, Jensen ES, Peoples MB. **2022**. Quantifying country-to-global scale nitrogen fixation for grain legumes
II. Coefficients, templates and estimates for soybean, groundnut and pulses. *Plant and Soil* **474**: 1–15.

Hossain MS, Frith C, Bhattacharyya SS, DeLaune PB, Gentry TJ. **2023**. Isolation and Characterization of Bacterial Endophytes
from Small Nodules of Field-Grown Peanut. *Microorganisms* **11**: 1941.

Houlden A, Timms-Wilson TM, Day MJ, Bailey MJ. **2008**. Influence of plant developmental stage on microbial community
structure and activity in the rhizosphere of three field crops. *FEMS Microbiology Ecology* **65**: 193–201.

Hütsch B, Augustin J, Merbach W. **2002**. Plant rhizodeposition - An important source for carbon turnover in soils. *Journal of*
Plant Nutrition and Soil Science **165**: 397–407.

Iven H, Walker TWN, Anthony M. **2022**. Biotic Interactions in Soil are Underestimated Drivers of Microbial Carbon Use
Efficiency. *Current Microbiology* **80**: 13.

Kawasaki A, Donn S, Ryan PR, Mathesius U, Devilla R, Jones A, Watt M. **2016**. Microbiome and Exudates of the Root and
Rhizosphere of Brachypodium distachyon, a Model for Wheat. *PLOS ONE* **11**: e0164533.

Kong Z, Liu H. **2022**. Modification of Rhizosphere Microbial Communities: A Possible Mechanism of Plant Growth Promoting
Rhizobacteria Enhancing Plant Growth and Fitness. *Frontiers in Plant Science* **13**: 920813.

Larrouy JL, Dhami MK, Jones EE, Ridgway HJ. **2023**. Physiological stage drives fungal community dynamics and diversity in
Leptospermum scoparium (mānuka) flowers. *Environmental Microbiology* **25**: 766–771.

Legendre P, Anderson MJ. **1999**. Distance-Based Redundancy Analysis: Testing Multispecies Responses in Multifactorial
Ecological Experiments. *Ecological Monographs* **69**: 1–24.

Losos JB. **2008**. Phylogenetic niche conservatism, phylogenetic signal and the relationship between phylogenetic relatedness
and ecological similarity among species. *Ecology Letters* **11**: 995–1003.

Lumibao CY, Kimbrough ER, Day RH, Conner WH, Krauss KW, Van Bael SA. **2020**. Divergent biotic and abiotic filtering of root
endosphere and rhizosphere soil fungal communities along ecological gradients. *FEMS Microbiology Ecology* **96**: fiaa124.

Lyu D, Msimbira LA, Nazari M, Antar M, Pagé A, Shah A, Monjezi N, Zajonc J, Tanney CAS, Backer R, *et al.* **2021**. The
Coevolution of Plants and Microbes Underpins Sustainable Agriculture. *Microorganisms* **9**: 1036.

Martiny JBH, Jones SE, Lennon JT, Martiny AC. **2015**. Microbiomes in light of traits: A phylogenetic perspective. *Science* **350**:
aac9323.

Mayfield MM, Levine JM. **2010**. Opposing effects of competitive exclusion on the phylogenetic structure of communities.
Ecology Letters **13**: 1085–1093.

Mayhood P, Mirza BS. **2021**. Soybean Root Nodule and Rhizosphere Microbiome: Distribution of Rhizobial and Nonrhizobial
Endophytes. *Applied and Environmental Microbiology* **87**: e02884-20.

de Mendiburu F. **2023**. agricolae: Statistical Procedures for Agricultural Research.

Morin A, Maurousset L, Vriet C, Lemoine R, Doidy J, Pourtau N. **2022**. Carbon fluxes and environmental interactions during
legume development, with a specific focus on Pisum sativum. *Physiologia Plantarum* **174**: e13729.

Moroenyane I, Mendes L, Tremblay J, Tripathi B, Yergeau É. **2021**. Plant Compartments and Developmental Stages Modulate
the Balance between Niche-Based and Neutral Processes in Soybean Microbiome. *Microbial Ecology* **82**: 416–428.

Mortier V, Holsters M, Goormachtig S. **2012**. Never too many? How legumes control nodule numbers. *Plant, Cell &*
Environment **35**: 245–258.

Mougel C, Offre P, Ranjard L, Corberand T, Gamalero E, Robin C, Lemanceau P. **2006**. Dynamic of the genetic structure of
bacterial and fungal communities at different developmental stages of Medicago truncatula Gaertn. cv. Jemalong line J5. *New*
Phytologist **170**: 165–175.

Nemergut DR, Schmidt SK, Fukami T, O'Neill SP, Bilinski TM, Stanish LF, Knelman JE, Darcy JL, Lynch RC, Wickey P, *et al.*
2013. Patterns and processes of microbial community assembly. *Microbiology and molecular biology reviews: MMBR* **77**: 342–
356.

O'Callaghan M, Ballard RA, Wright D. **2022**. Soil microbial inoculants for sustainable agriculture: Limitations and
opportunities. *Soil Use and Management* **38**: 1340–1369.

Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Solymos P, Stevens MHH, Szoecs E, *et al.*
2022. vegan: Community Ecology Package.

Pacheco-Cano RD, Salcedo-Hernández R, López-Meza JE, Bideshi DK, Barboza-Corona JE. **2018**. Antimicrobial activity of
broccoli (Brassica oleracea var. italica) cultivar Avenger against pathogenic bacteria, phytopathogenic filamentous fungi and
yeast. *Journal of Applied Microbiology* **124**: 126–135.

Peay KG, Belisle M, Fukami T. **2012**. Phylogenetic relatedness predicts priority effects in nectar yeast communities.
Proceedings. Biological Sciences **279**: 749–758.

Peix A, Ramírez-Bahena MH, Velázquez E, Bedmar EJ. **2015**. Bacterial Associations with Legumes. *Critical Reviews in Plant*
Sciences **34**: 17–42.

Pekkonen M, Ketola T, Laakso JT. **2013**. Resource Availability and Competition Shape the Evolution of Survival and Growth
Ability in a Bacterial Community. *PLoS ONE* **8**: e76471.

Pyron RA, Costa GC, Patten MA, Burbrink FT. **2015**. Phylogenetic niche conservatism and the evolutionary basis of ecological
speciation. *Biological Reviews* **90**: 1248–1262.

Qiao Q, Wang F, Zhang J, Chen Y, Zhang C, Liu G, Zhang H, Ma C, Zhang J. **2017**. The Variation in the Rhizosphere Microbiome
of Cotton with Soil Type, Genotype and Developmental Stage. *Scientific Reports* **7**: 3940.

Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. **2018**. Posterior Summarization in Bayesian Phylogenetics Using
Tracer 1.7. *Systematic Biology* **67**: 901–904.

Rao CR. **1982**. Diversity and dissimilarity coefficients: A unified approach. *Theoretical Population Biology* **21**: 24–43.

Rekha K, Baskar B, Srinath S, Usha B. **2018**. Plant-growth-promoting rhizobacteria Bacillus subtilis RR4 isolated from rice
rhizosphere induces malic acid biosynthesis in rice roots. *Canadian Journal of Microbiology* **64**: 20–27.

Rensink S, Wiegand S, Kallscheuer N, Rast P, Peeters SH, Heuer A, Boedeker C, Jetten MSM, Rohde M, Jogler M, *et al.* **2020**.
Description of the novel planctomycetal genus Bremerella, containing Bremerella volcania sp. nov., isolated from an active
volcanic site, and reclassification of Blastopirellula cremea as Bremerella cremea comb. nov. *Antonie van Leeuwenhoek* **113**:
1823–1837.

Robinson MD, McCarthy DJ, Smyth GK. **2010**. edgeR: a Bioconductor package for differential expression analysis of digital
gene expression data. *Bioinformatics (Oxford, England)* **26**: 139–140.

Rohrbacher F, St-Arnaud M. **2016**. Root Exudation: The Ecological Driver of Hydrocarbon Rhizoremediation. *Agronomy* **6**: 19.

Romdhane S, Spor A, Aubert J, Bru D, Breuil M-C, Hallin S, Mounier A, Ouadah S, Tsiknia M, Philippot L. **2022**. Unraveling
negative biotic interactions determining soil microbial community assembly and functioning. *The ISME journal* **16**: 296–306.

Sasse J, Martinoia E, Northen T. **2018**. Feed Your Friends: Do Plant Exudates Shape the Root Microbiome? *Trends in Plant*
Science **23**: 25–41.

Schemske DW, Mittelbach GG, Cornell HV, Sobel JM, Roy K. **2009**. Is There a Latitudinal Gradient in the Importance of Biotic
Interactions? *Annual Review of Ecology, Evolution, and Systematics* **40**: 245–269.

Schöler A, Jacquiod S, Vestergaard G, Schulz S, Schloter M. **2017**. Analysis of soil microbial communities based on amplicon
sequencing of marker genes. *Biology and Fertility of Soils* **53**: 485–489.

Shade A, Jacques M-A, Barret M. **2017**. Ecological patterns of seed microbiome diversity, transmission, and assembly. *Current*
Opinion in Microbiology **37**: 15–22.

Shamoot S, McDonald L, Bartholomew WV. **1968**. Rhizo-Deposition of Organic Debris in Soil. *Soil Science Society of America*
Journal **32**: 817–820.

Singer AC. **2006**. The Chemical Ecology of Pollutant Biodegradation: Bioremediation and Phytoremediation from Mechanistic
and Ecological Perspectives. In: Mackova M, Dowling D, Macek T, eds. Phytoremediation Rhizoremediation. Dordrecht:
Springer Netherlands, 5–21.

Smýkal P, Kenicer G, Flavell AJ, Corander J, Kosterin O, Redden RJ, Ford R, Coyne CJ, Maxted N, Ambrose MJ, *et al.* **2011**.
Phylogeny, phylogeography and genetic diversity of the Pisum genus. *Plant Genetic Resources* **9**: 4–18.

Spicer ME, Woods CL. **2022**. A case for studying biotic interactions in epiphyte ecology and evolution. *Perspectives in Plant*
Ecology, Evolution and Systematics **54**: 125658.

Tkacz A, Bestion E, Bo Z, Hortala M, Poole PS. **2020**. Influence of Plant Fraction, Soil, and Plant Species on Microbiota: a
Multikingdom Comparison. *mBio* **11**: e02785-19.

Voisin A-S, Bourion V, Duc G, Salon C. **2007**. Using an Ecophysiological Analysis to Dissect Genetic Variability and to Propose
an Ideotype for Nitrogen Nutrition in Pea. *Annals of Botany* **100**: 1525–1536.

Voisin AS, Salon C, Jeudy C, WAREMBOURG FR. **2003a**. Seasonal Patterns of 13C Partitioning Between Shoots and Nodulated
Roots of N2- or Nitrate-fed Pisum sativum L. *Annals of Botany* **91**: 539–546.

Voisin AS, Salon C, Jeudy C, Warembourg FR. **2003b**. Symbiotic N2 fixation activity in relation to C economy of Pisum sativum
L. as a function of plant phenology. *Journal of Experimental Botany* **54**: 2733–2744.

Wang B, Sugiyama S. **2020**. Phylogenetic signal of host plants in the bacterial and fungal root microbiomes of cultivated
angiosperms. *The Plant Journal* **104**: 522–531.

Webb CO, Ackerly DD, McPeek MA, Donoghue MJ. **2002**. Phylogenies and Community Ecology. *Annual Review of Ecology*
and Systematics **33**: 475–505.

Weber H, Borisjuk L, Wobus U. **1997**. Sugar import and metabolism during seed development. *Trends in Plant Science* **2**: 169–
174.

Xiong C, Zhu Y-G, Wang J-T, Singh B, Han L-L, Shen J-P, Li P-P, Wang G-B, Wu C-F, Ge A-H, *et al.* **2021**. Host selection shapes
crop microbiome assembly and network complexity. *The New Phytologist* **229**: 1091–1104.

Yang Z. **1994**. Maximum likelihood phylogenetic estimation from DNA sequences with variable rates over sites: Approximate
methods. *Journal of Molecular Evolution* **39**: 306–314.

Yang N, Nesme J, Røder HL, Li X, Zuo Z, Petersen M, Burmølle M, Sørensen SJ. **2021**. Emergent bacterial community
properties induce enhanced drought tolerance in Arabidopsis. *NPJ biofilms and microbiomes* **7**: 82.

Yilmaz P, Parfrey LW, Yarza P, Gerken J, Pruesse E, Quast C, Schweer T, Peplies J, Ludwig W, Glöckner FO. **2014**. The SILVA
and 'All-species Living Tree Project (LTP)' taxonomic frameworks. *Nucleic Acids Research* **42**: D643-648.

Zhang Y, Xu J, Riera N, Jin T, Li J, Wang N. **2017**. Huanglongbing impairs the rhizosphere-to-rhizoplane enrichment process
of the citrus root-associated microbiome. *Microbiome* **5**: 97.

Zhang N, Yang D, Wang D, Miao Y, Shao J, Zhou X, Xu Z, Li Q, Feng H, Li S, *et al.* **2015**. Whole transcriptomic analysis of the
plant-beneficial rhizobacterium Bacillus amyloliquefaciens SQR9 during enhanced biofilm formation regulated by maize root
exudates. *BMC Genomics* **16**: 685.

Zhu Q, Mai U, Pfeiffer W, Janssen S, Asnicar F, Sanders JG, Belda-Ferre P, Al-Ghalith GA, Kopylova E, McDonald D, *et al.*
2019. Phylogenomics of 10,575 genomes reveals evolutionary proximity between domains Bacteria and Archaea. *Nature*
Communications **10**: 5477.

Zohary D, Hopf M. **1973**. Domestication of Pulses in the Old World: Legumes were companions of wheat and barley when
agriculture began in the Near East. *Science (New York, N.Y.)* **182**: 887–894.

Figure 1: Bacterial community diversity, structure and composition. **Panel A-B:** Specific richness (A) and Shannon diversity index (B) of bacterial communities across bulk soil and the different plant-associated microhabitats. Multiple Kruskal-Wallis comparisons. **Panel C:** Bray-Curtis distance-based redundancy analysis (similar to Constrained Analysis of Principal Coordinates). The model was tested using PERMANOVA with 10,000 permutations. The R-squared value indicates the percentage of variance explained by the model, including the three variables (microhabitat, growth stages and host genotype) and their interactions. Note that for figure clarity, host genotypes are not displayed. **Panel D:** Average taxonomic composition of bacterial communities based on relative abundance at the Order level across bulk soil, plant-associated microhabitats and growth stages. Only the eleven most abundant Orders are shown, with the remaining Orders grouped together and labeled as "others". Note that data from all host genotypes were compiled. (N_{total} = 180, N_{Bulk} = 21, N_R = 46, N_{Rp} = 56, N_E = 57)

Figure 2: Phylogenetic diversity and dispersion parameter of bacterial communities across the different microhabitats. **Panel A:** *Abundance weighted phylogenetic diversity (Rao's quadratic entropy index, FD_Q) (Rao, 1982) of bacterial communities across the different microhabitats. Multiple Kruskal-Wallis test, $p < 0.001$, N = 180.* **Panel B:** *Phylogenetic dispersion parameter (D) (Darcy et al., 2020) of bacterial communities across the different plant-associated microhabitats. D > 0 indicates overdispersion, D = 0 is the neutral model and D < 0 indicates underdispersion. N = 159. The p-values indicate significant differences in D values to neutral model, rather than differences between microhabitats.*