Title

Notch expression during ctenophore development gives insight into its ancestral function

Authors

*Brent Foster[1], Fredrik Hugosson[1], Cezar Borba[1], James Strother[1], Mark Q. Martindale[1]

Affiliations

1. The Whitney Laboratory for Marine Bioscience, Department of Biology, University of Florida, St.

Augustine, FL 32080, USA

* Corresponding author: bf400@exeter.ac.uk

Summary

The canonical *Notch* pathway is a juxtacrine signaling module with widely conserved roles maintaining

progenitor cell populations, promoting binary cell fate decisions, and establishing tissue boundaries. This

pathway emerged in Metazoan lineages, although many molecular components and regulators existed

prior to that divergence. Given that the vast majority of *Notch* studies focus on bilaterians, it is unclear

when or how *Notch* gained its developmental signaling functions. To clarify the ancestral function of

Notch, we turned to an early branching Metazoan — the ctenophore *Mnemiopsis leidyi* — and conducted

structural analyses of putative *Notch* components to evaluate predicted signaling potential. We

characterized gene expression of these components with *in situ* hybridization and show they are expressed

during late embryogenesis. Using hybridization chain reaction (HCR[TM]), we examine *MlNotch* expression

relative to canonical transcriptional targets and putative stem cell markers and identify differential

expression patterns at a cellular resolution. Pharmacological inhibition reveals zones of active and

inactive *MlNotch* during late embryonic development. Our results suggest that *Notch* evolved its signaling

potential by the Metazoan divergence and is likely involved in regulating progenitor cell populations and

coordinating developmental fate decisions.

Keywords

Introduction

The canonical *Notch* pathway is a juxtacrine signaling module involved in various aspects of development across Metazoa. It is best known for its role in promoting binary cell fate decisions via lateral inhibition and/or induction, maintaining stem and progenitor cell populations, and regulating terminal differentiation (for reviews of the role of *Notch* during development, see [1,2]). Many of the key regulatory enzymes and transcriptional targets for *Notch* signaling predate the divergence of Metazoan lineages, suggesting that this pathway evolved by co-opting molecules from pre-existing signaling modules[3,4] (see also Figure 1A).

Canonical *Notch* signaling consists of a transmembrane protein that undergoes several distinct proteolytic cleavages for its activation. In the Golgi apparatus, Furin-like convertase forms a *Notch* heterodimer (S1 cleavage) that translocates to the cell membrane, where it acts as a signal receptor. The *Notch* receptor can then interact with a ligand expressed in a neighboring cell, exposing a second extracellular cleavage site recognized by a membrane-bound metalloprotease (S2 cleavage). This cleavage releases the *Notch* extracellular domain (NECD) from the membrane, allowing it to be taken up by the neighboring signaling cell. Meanwhile, γ-secretase in the signal-receiving cell recognizes a third cleavage site and releases the *Notch* intracellular domain (NICD) from the membrane (S3 cleavage), whereupon it translocates to the nucleus and acts as a co-factor to regulate transcription (see Figure 1B for a detailed cell diagram depicting canonical *Notch* signaling in bilaterians).

The canonical *Notch* pathway is likely a Metazoan innovation and is present in virtually all bilaterian model systems, although when and how this signaling module emerged during animal evolution remains a mystery. Genomic evidence in single-celled Eukaryotes suggests that domain shuffling of separate proteins could have led to the formation of Metazoan *Notch*[5,6]. A more recent report has identified a *Notch-like* gene with conserved PFAM domains in their canonical order in the choanoflagellate species *Mylnosiga fluctuans*, although to date no study has identified a non-Metazoan species that encodes both *Notch* receptor and a recognizable activating ligand[7]. The vast majority of *Notch* studies are conducted in bilaterians and have demonstrated a broad conservation of components and functions[4,8–10]. Comparative genomics has shown that *Notch* orthologs and putative pathway

components have been identified in each non-bilaterian clades, however, to our knowledge *in situ*

characterizations of *Notch* components in non-bilaterians have only been conducted in a handful of

sponges[11,12] and cnidarians[13–17]. To better understand how the *Notch* pathway and its various

developmental functions evolved, it is necessary to understand the developmental context in which these

components are expressed in other early branching Metazoan representatives.

Ctenophores, as the likely sister group to all other Metazoans[18–20], can offer key evolutionary

insight into how *Notch* signaling and its developmental functions emerged in Metazoans, regardless of

whether a canonical pathway is intact or not. *Mnemiopsis leidyi,* the primary subject of this report,

encodes at least 16 out of 22 canonical *Notch* pathway components in its genome, with several other

components reported to be missing so-called diagnostic domains, raising questions concerning their

orthology and role in a functional pathway[18]. Existing data only provide presence or absence information

which has led subsequent reports to equate the absence of these canonical *Notch* components and PFAM

domains as evidence for the absence of a functional signaling pathway[9,10,18]. To date, no study has

analyzed the cellular and developmental context in which putative canonical components are expressed

during ctenophore development. We therefore have examined the predicted signaling capacity of *M. leidyi*

Notch (*MlNotch*) and the spatiotemporal expression of putative pathway regulators and targets during

early ctenophore development to gain insight into how these components may have evolved to form a

functional developmental signaling module.

Results

Figure 1. The canonical *Notch* signaling pathway emerged in Metazoan lineages. A) Simple phylogeny
depicting when molecular components of the canonical *Notch* pathway evolved. B) Cell diagram of
canonical *Notch* pathway in bilaterians, with components color-matched to A) when they are believed to
have first emerged. C) Hypothetical cell diagram of components of the putative canonical Notch pathway
encoded in the *Mnemiopsis leidyi* genome. D) Developmental RNA-seq of putative canonical *Notch*
pathway genes[37]. Bolded genes in the key represent genes expressed during the developmental peak
shaded in gray. * = previously unreported genes.

M. leidyi genome encodes many canonical Notch components with characteristic PFAM domains

Many components of the canonical *Notch* pathway are definitively encoded in the genome of the

ctenophore *Mnemiopsis leidyi*[18], although these characterizations have been limited to an absent/present

dichotomy that assumed that missing components or diagnostic domains of putative *Notch* genes equated

non-functionality. However, if these putative canonical components are expressed in the same or adjacent

cells at the same time, they could ostensibly behave as a rudimentary pathway (compare Figure 1B to

Figure 1C). We therefore examined the cellular and developmental context in which these components are

expressed to test the hypothesis that canonical *Notch* components are expressed in shared domains during

early ctenophore development and function as a primitive pathway.

Figure S1. Domain architectures of proteins implicated in the canonical *Notch* pathway. A) Candidate
ligands, *MlDelta-like* and *MlCad-like*. B) Putative S2 proteases, *MlFurin-like1* and *MlFurin-like2*. C)
Hairy-Enhancer of Split orthologs, *MlHES1* and *MlHES2*. D) E3 ubiquitin ligase, *MlMindbomb-like.*

We first confirmed that these ctenophore *Notch* components contain all of their characteristic

PFAM domains. We conducted independent BLAST searches of canonical *Notch* components against the

M. leidyi genome and confirmed hits with reciprocal BLAST and PFAM analysis. Upon examination of

the reported *MlDelta-like* (ML21438a) gene, we were able to identify the delta-serrate ligand (DSL)

PFAM domains previously reported missing (see Figure S1A and File S1). However, reciprocal BLAST

searches did not result in any hit with matching PFAM domains, leading us to conclude that this hit was

an error in the genome sequencing. Furthermore, BLAST searches of the *Drosophila Delta-like* sequence

against the *M. leidyi* genome yielded the gene identified as *MlNotch* (ML128617a) and a gene rich in both

EGF repeats and LamininG domains, characteristic of proto-cadherins (ML24144a; see Figure S1A and

File S1). Given that these genes lack both DSL and MNNL domains, we do not consider them true *Delta-*

like proteins. We therefore refer to ML24144a as *MlCadherin-like* (*MlCad-like*). Our searches also

revealed that the Furin-like gene previously reported[18] (ML07022a) retains several Furin-like repeats and

subtilase domains characteristic of Furin convertases; we therefore refer to this gene as *MlFurin-like1*. We

also identified previously unreported orthologs for a second Furin-like convertase (ML279823a; Figure

S1B and File S1), a second HES transcription factor (ML065739a; Figure S1C and File S1), and an

ortholog of the *Delta-like* regulator *Mindbomb* (MleFL.S3.22491; Figure S1D and File S1). We refer to

these genes as *MlFurin-like2*, *MlHES2*, and *MlMindbomb*, respectively.

Using existing developmental RNA-seq data from *M. leidyi*[21], we plotted the median transcripts

per million (TPMs) of the putative *Notch* components encoded in the genome identified in our

independent searches. We noted three apparent peaks of gene expression at approximately 4 hours post

fertilization (hpf), 8hpf, and 12hpf (Figure 1D). We opted to focus our gene expression studies on the

8hpf timepoint to avoid potential confounds with the maternal-to-zygotic transcriptional transition[22]. The

genes expressed in this 8hpf peak correspond to putative components of the core canonical *Notch*

pathway: *MlNotch* (receptor), *MlPCad-like* (candidate ligand), *MlFurin-like2* (S1 cleavage), *MlADAM10*

(S2 cleavage), *MlPresenilin* (S3 cleavage), *MlSuH* (co-factor), and *MlHES2* (transcriptional target).

Figure 2 *Notch* domains predate Metazoans. A) Maximum likelihood/Bayesian phylogeny of *Notch* across
Metazoa showing conserved *Notch* Intracellular Domains (NICDs). Node labels are Maximum Likelihood
/ Posterior Probabilities values. B) AlphaFold 3D prediction of *MlNotch* Negative Regulatory Region
(NRR) and NICD. C) Basic alignments of *Notch* juxta-transmembrane domains highlighting conserved
S1, S2, and S3 cleavage recognition sites. Species in panel C) are color-matched to phyla from panel A).

Notch negative regulatory regions and intracellular domains are highly conserved

We next wanted to evaluate the conserved signaling capacity of the *Notch* receptor by conducting a

phylogenetic analysis of *Notch* orthologs across Metaozoa and analyzing their respective PFAM domain

architectures and juxta-transmembrane domains (J/TM). Lin-*Notch* Repeats (LNR) and Ankyrin repeats

characteristic of *Notch* evolved before the Metazoan lineages. In fact, the LNR and *Notch* Intracellular

Domain (NICD) are remarkably conserved across Metazoans, whereas the extracellular Epidermal

Growth Factor (EGF) repeats are highly variable (Figure 2A). Our phylogram of *Notch* has strong support

placing ctenophore *Notch* proteins at the base of Metazoa, supporting previous supposition that NICD

architecture was present in the last common animal ancestor[7]. While each of the predicted ctenophore

Notch proteins contains conserved nuclear localization signals, we were unable to identify these domains

in all of the representative holozoan *Notch-like* proteins.

Given that canonical *Notch* signaling activity requires three discrete cleavage events, we wanted

to evaluate whether *MlNotch* has conserved regulatory structures and cleavage recognition sites. Our

AlphaFold prediction shows that *MlNotch* LNRs cover the extracellular J/TM region in *MlNotch* (Figure

2B). Our multiple sequence alignment shows that residues recognized by Furin-like and γ-secretase are

highly conserved, whereas the putative S2 cleavage site is poorly conserved (Figure 2C). The minimal

Furin-like recognition motif (RXXR) is only observed in bilaterian lineages, although we found a highly

conserved Arginine residue aligned in nearly all of the examined sequences. W and P residues of the

ΦWΦP motif are also highly conserved. Taken together, these analyses suggest that *MlNotch* may retain

signal-transducing capacity independent of S1 or S2 cleavage via Furin-like convertase and ADAM

metalloproteases, respectively.

Figure 3. Genes implicated in canonical *Notch* signaling are expressed in developing pharynx and tentacle bulbs at 8hpf. A–A') *MlNotch*. B–B') *MlCad-like*. C–C') *MlFurin-like1*. D–D') *MlFurin-like2*. E–E') *MlADAM10*. F–F') *MlPresenillin*. G–G') *Suppressor-of-Hairless (MlSuH)*. H–H') *MlHairy-Enhancer-of-Split2 (MlHES2)*. * = oral pole. All scalebars = 50µm. White arrowheads indicate distinct patches of expression in aboral cells.

Figure S2. *MlDelta-like* is not detectable 8hpf. A) Lateral view. B) Aboral view. * = oral pole. All scalebars = 50µm.

Putative Notch components share expression domains during late ctenophore embryogenesis

We examined gene expression of *MlNotch* and *MlCad-like* during late embryonic development. *MlNotch*

transcripts are broadly expressed at 8hpf in developing pharynx, tentacle bulbs, and aboral organ and in

ectodermal cells invaginating at the oral pole (Figure 3A). We also identified a peculiar symmetrical

MlNotch expression pattern in groups of ectodermal cells at the aboral end of developing embryos (Figure

3A'). *MlCad-like* is expressed in the developing pharynx and tentacle bulbs, similar to *MlNotch* (Figure

3B and Figure 3B'), although we failed to detect clear expression in the aboral end (Figure 3B'). We did

not detect expression of the previously identified *MlDelta-like* 8hpf (Figure S2). All four putative protease

genes share broad expression domains at 8hpf in the developing pharynx and tentacle bulbs, although we

failed to detect expression of these genes in the aboral end (Figure 3C–3F).

We next examined gene expression of a putative *MlNotch* co-factor (*MlSuH*) and transcriptional

target (*MlHES2*). Colorimetric *in situs* of *MlSuH* illustrate a similar expression pattern to *MlNotch* with

broad expression domains in the developing pharynx and tentacle bulbs (Figure 3G). We also note patches

of *MlSuH* expression in the aboral end of the developing embryo, similar to our observations for *MlNotch*

(Figure 3G'). *MlHES2* is expressed in similar domains (Figure 3H and Figure 3H').

Figure 4. *MlNotch* and *MlCad-like* transcripts are both differentially and co-expressed 8hpf. A–C) Lateral
views of *MlNotch* (A), MlCad-like (B), and Merged channels (C). D–F) Aboral views of *MlNotch* (D),
MlCad-like (E), and Merged channels (F). * = oral pole. All scalebars = 50µm. White arrows indicate
cells co-expressing *MlNotch* and *MlCad-like*. Yellow arrows indicate *MlNotch*-positive cells. Magenta
arrows indicate *MlCad-like*-positive cells.
189

MlNotch and a candidate ligand are expressed in distinct cell populations

To obtain a more cellular resolution of these gene expression patterns, we modified a hybridization chain

reaction (HCR^{TM}) protocol[23]. We observed weak expression of *MlNotch* in the developing pharynx and

tentacle bulbs and identified a strong *MlNotch* signal in distinct patches of cells at the aboral end of 8hpf

embryos (Figure 4A). *MlCad-like* signal is most pronounced at the oral pole and developing tentacle

bulbs (Figure 4B). While we note apparent regions where *MlNotch* and *MlCad-like* are co-expressed

(Figure 4C; see insert 4C'), we also identify differential expression in the developing pharynx and

tentacle bulbs (Figure 4C; see inserts 4C''–4C'''). Aboral views confirm the symmetrical nature of

MlNotch expression in distinct patches of cells, similar to what we observed in our colorimetric *in situs*

(Figure 4D; see insert 4D'). *MlCad-like* does not appear to be expressed in the same or neighboring cells

in the aboral end of developing embryos (Figure 4E; see insert 4E'). We note regions where *MlNotch* and

MlCad-like appear to be co-expressed in the ectoderm of the developing tentacle bulbs (Figure 4D–4F;

see inserts 4D'', 4E'', and 4F'').

Figure 5. *MlNotch*-positive cells neighbor cells expressing canonical transcriptional targets. A–C) Lateral views of *MlNotch* (A), *MlHES2* (B), and Merged channels (C). D–F) Aboral views of *MlNotch* (D), *MlHES2* (E), and Merged channels (F). G–I) Lateral views of *MlNotch* (G), *MlHES2* (H), and Merged channels (I). Aboral views of *MlNotch* (J), *MlHES2* (K), and Merged channels (L). M) Cartoon schematic representing expression of *MlNotch* relative to *MlSox1* from the panels in J–L. * = oral pole. All scalebars = 50µm. Magenta arrowheads indicate cells expressing *MlHES2* or *MlSox1*, but not *MlNotch*. White arrowheads indicate cells that co-express *MlNotch* and either *MlHES2* or *MlSox1,* respectively.

MlNotch-positive cells co-express canonical transcriptional targets or are found adjacent to cells

expressing target transcripts

We next considered expression of *MlNotch* relative to *MlHES2* and *MlSox1*, both canonical

transcriptional targets. *MlHES2* shares similar expression domains as *MlCad-like* in the developing

tentacle bulbs and pharynx 8hpf, with several cells differentially expressed between *MlNotch*-positive

domains in the aboral end (Figure 5A–F). Cells in the developing tentacle bulb appear to co-express

MlNotch and *MlHES2* transcripts (Figure 5D–F), suggestive of two distinct cell populations or perhaps of

a transcriptional transition in cell identity during development.

MlSox1 is a member of the SoxB gene family and has been shown to be expressed in distinctive

domains in the aboral end of developing *M. leidyi* embryos with colorimetric *in situ* hybridization[24]. We

corroborate that expression pattern here with HCR[TM] (Figure 5G–L) and show *MlNotch*-positive cells

neighboring *MlSox1*-positive cells in the aboral end and developing tentacle bulbs of embryos 8hpf

(Figure 5J–L). We also show a previously unidentified ring of *MlSox1*-positive cells medial to an outer

ring of *MlNotch*-positive cells near where the apical organ will develop (Figure 5J–L; see inserts 5J",

5K", 5L"). These expression patterns are highly reminiscent of classical models of *Notch*-induced lateral

inhibition and lateral induction[1,25,26] (see Figure 5M).

Figure S3. *MlNotch* expression 24hpf. A–A") Colorimetric ISH of *MlNotch* 24hpf. A) Lateral view showing *MlNotch* expression in mesogleal cells, tentacle bulbs, and floor of the apical organ. A') Aboral view at the comb row plane showing *MlNotch* expression in presumptive endodermal cells surrounding the comb rows. A") Aboral view at the plane of the apical organ. B–C') HCR of *MlNotch* 24hpf. B) Lateral view showing *MlNotch* expression in the floor of the apical organ and in mesogleal cells (see arrowheads). C) Closeup lateral view of *MlNotch*-expressing cells that line the endodermal canals and are sub-adjacent to the comb rows (see arrowhead). C') Closeup lateral view of *MlNotch*-expressing cells in the floor of the apical organ. * = oral pole. All scalebars = 50µm. P = pharynx, CR = comb rows, TB = tentacle bulb, T = tentacle, AO = apical organ.

MlNotch is expressed in the floor of the apical organ, comb rows, and endodermal canals 24hpf

We were curious to see how *MlNotch* expression and its implied function might change during ctenophore development. Colorimetric *in situ* hybridization shows *MlNotch* expression in gut regions, tentacles, and the floor of the fully developed apical organ of 24h cydippids (Figure S3A–A"). We also note what appears to be cells in the tissue lining endodermal canals (see arrowheads in Figure S3A and 3A'). Aboral views reveal *MlNotch* expression in cells lining the ciliary comb rows (Figure S3A') and cells that make up the floor of the apical organ (Figure S3A"). HCR^TM experiments corroborate our colorimetric *in situs* with strong *MlNotch* signal in neighboring cells of the apical organ (Figure S3B), comb rows and endodermal canals (Figure S3C and Figure S3C').

Figure S4. Whole-mount colorimetric *in situ* hybridization of putative genes involved in S1–S3 cleavage. A–A''') *MlFurin-like1* expression show in in lateral view (A), plane of the apical organ (A'), plane of the comb rows (A''), and plane of the tentacle bulbs (A'''). B–B''') *MlFurin-like2* expression show in in lateral view (B), plane of the apical organ (B'), plane of the comb rows (B''), and plane of the tentacle bulbs (B'''). C–C''') *MlADAM10* expression show in in lateral view (C), plane of the apical organ (C'), plane of the comb rows (C''), and plane of the tentacle bulbs (C'''). D–D''') *MlPresenillin* expression show in in lateral view (D), plane of the apical organ (D'), plane of the comb rows (D''), and plane of the tentacle bulbs (D'''). * = oral pole. All scalebars = 50µm.

Putative protease genes share expression domains in the apical organ, tentacle bulbs, and pharynx

24hpf

We evaluated expression patterns of putative S1–S3 protease genes at 24hpf via whole-mount

colorimetric *in situ* hybridization. Both *MlFurin-like1* and *MlFurin-like2* are exclusively expressed in

tentacle bulbs in 24h cydippids (Figure S4A–4B; see also Figure S4A''' and 4B'''). *MlADAM10* is also

expressed in the tentacle bulbs and pharynx (Figure S4C) with additional expression in the apical organ

(Figure S4C'), around the comb rows (Figure 4SC'') and tentacle bulb (Figure S4C'''). *MlPresenillin*, the

active subunit of γ-secretase responsible for S3 cleavage, shares similar expression domains in the

pharynx, apical organ, and cells surrounding the comb rows (Figure S4D–D'''). These results support

conditions in which these genes may be co-expressed to interact in the tentacle bulbs, although they also

hint at differential expression — and therefore function — in cells of the apical organ, comb rows, and

pharynx.

Figure 6. Traditional *Notch* inhibitors reduce expression of *MlNICD* in developing ctenophores. A)
Experimental setup of pharmacological inhibition experiments. B–B'') Representative ctenophore embryo
incubated in 1% DMSO. B = lateral view, B' = aboral view. B'' shows a lateral view. White arrowheads in
i. and ii. point to mesenchymal cells harboring *MlNICD*. C–C') Representative ctenophore embryo
incubated in 10μM DAPT. C = lateral view, C' = aboral view. D–D') Representative ctenophore embryo
incubated in 25μM LY411575. D = lateral view, D' = aboral view. * = oral pole. All scalebars = 50μm.

Pharmacological inhibition reveals regions of active and inactive MlNotch

To test whether *MlNotch* is cleaved during late ctenophore embryogenesis, we utilized DAPT and

LY411575 — both potent γ-secretase inhibitors[27] — combined with an antibody that recognizes cleaved

Notch (see Figure 6A for experimental setup). We note NICD signal localized to the invaginating pharynx

and developing tentacle bulbs (Figure 6B), suggesting that these regions are rich in activated *MlNotch*.

However, we failed to detect signal in the distinctive aboral patches of cells characterized with our

colorimetric and HCR *in situs* (Figure 6B'), suggesting that these cell populations are most likely loaded

with unprocessed *MlNotch* transcripts or inactive *MlNotch* protein. Mesenchymal cells within the

mesoglea also seem to harbor activated *MlNotch* (Figure 6B''; see white arrow heads). Both DAPT- and

LY411575-treated embryos show a noticeable decrease in signal within the pharynx and tentacle bulb

regions (Figure 6C & Figure 6D). We did not detect any obvious phenotype or loss of cell types in the

embryonic or cydippid stages. Furthermore, drug-treated embryos appear to develop into phenotypically

normal cydippids. Together, our results are suggestive of a mechanism where "inactive" *MlNotch*

regulates a distinct sub-population of cells that, upon cleavage-induced activation, are expressed in

mesenchymal cells and cells that contribute to the future tentacle bulb sheath and pharyngeal epithelium.

Discussion

A Notch receptor existed early in the Metazoan divergence and likely retained signaling capacity

independent of S1 cleavage

Our bioinformatic structural analysis of the *Notch* juxta-transmembrane domains suggests that *MlNotch*

likely does not undergo S1 cleavage. Previous work has identified a Furin-independent function for *Notch*

and concluded that this particular role is derived[28,29], although these studies were limited to cell cultures

and *Drosophila*. Here, we show that the minimum Furin-like recognition motif (RXXR) in *Notch* was

either lost in ctenophore lineages or a bilaterian innovation. Given that our analysis shows that other non-

bilaterian Metazoan *Notch* proteins also lack this exact motif, we favor the latter interpretation. This

suggests that S1 cleavage and the heterodimerization of the *Notch* receptor could in fact be derived, which

in turn would indicate that the Furin-independent function of *Notch* was ancestral.

Similarly, our analysis shows that the S2 cleavage site seems to be highly diverse in non-

bilaterian lineages, although *MlNotch* has an LNR domain architecture that may retain a negative

regulatory function by covering an as-of-yet unidentified site recognized by another membrane-bound

protease (Figure 2B). Choanoflagellates offer a twist in that their *Notch-like* genes also contain structures

reminiscent of the negative regulatory regions that, in canonical *Notch* signaling, functions as a cap that

covers the S2 sites recognized by ADAM proteinases. In the choanoflagellate *Monosiga brevicolis*,

Notch-like genes contain an intracellular LNR domain, suggesting that pre-Metazoan *Notch-like* proteins

may have been capable of intracellular S2-like cleavages, whereas *M. fluctuans* retains the canonical

order of PFAM domains and therefore may have been capable of a more canonical extracellular cleavage.

Given that *M. brevicolis* is a more recent lineage, this would suggest that the shuffled *Notch* domains is a

derived choanoflagellate characteristic and that the pre-Metazoan *Notch* looked — and may have even

functioned — similar to the canonical Metazoan *Notch*.

Developmental and cellular context of MlNotch expression is consistent with models of cell-cell

signaling

The putative *Notch* pathway transcripts characterized here all share similar expression domains at the

same developmental stage, consistent with what we would expect from components interacting with each

other in a functional signaling pathway. The intriguing aboral patches of *MlNotch*-positive cells combined

with the data from our pharmacological *Notch* inhibition suggest that these populations are loaded with

MlNotch transcripts that have not been processed. We hypothesize that these cells are a self-regulating

progenitor population characterized by uncleaved — or "inactive" — *MlNotch*. *MlNotch*-positive

mesenchymal cells and cells within the developing tentacle bulb and pharynx appear to harbor cleaved —

or "active" — *MlNotch*. It is unclear if these active *MlNotch*-positive cells are migrating, but we cannot

rule out this possibility. *Notch* signaling has been implicated in coordinating the epithelial-to-

mesenchymal transition in cell culture[30] and regulating migratory peripheral glia in *Drosophila*[31], motile

cells in zebrafish trunk neural crest[32], and migrating neurons in developing mouse cerebral cortex[33]. To

our knowledge, no study has identified a role for *Notch* signaling in regulating migratory cells outside of

bilateria. Our results are suggestive of a mechanism where "inactive" *MlNotch* regulates a distinct sub-

population of cells that, upon cleavage-induced activation, may be involved in the epithelial-to-

mesenchymal transition and the formation of the tentacle bulb sheath and pharyngeal epithelium, pointing

to an ancient role for *Notch* regulation of migratory cells. It is unclear what the functions of the other

canonical *Notch* pathway genes are in these *Notch*-expressing cells that apparently lack stable cell-cell

interactions.

As noted previously, the expression of *MlNotch* in relation to putative transcriptional targets,

MlHES2 and *MlSox1*, is highly reminiscent of classical models of lateral inhibition and induction. This

result is intriguing, given that *MlSox1* belongs to the SoxB group[34] and *Notch* has been shown to have a

conserved role in regulating *Sox* genes. Pharmacological inhibition of *Notch* in *Nematostella vectensis*

upregulates *SoxB2* genes[35] while downregulating HES genes[14]. A *Sox*-related gene regulatory network is

also moderated by *Notch* in *Hydra vulgaris*[36,37]. *Notch* has also been shown to regulate *Sox* in cephalopod

and mouse retinal development[38,39]. *Drosophila* neuroblast formation appears to be gated by a *Notch* and

SoxB interplay[40]. *SoxB1* coordinates with *NICD* to maintain neural cells in an undifferentiated state in

chick embryos[41]. We do not show any clear evidence of *MlNotch* or *MlSox1* promoting neuronal or

sensory cell fate in ctenophores, although we do not preclude that possibility. *M. leidyi* embryos develop

recognizable neurons by 24hpf, with condensations around the floor of the apical organ by 24hpf[42],

precisely where we see *MlNotch* expressed (Figure S3A''). Given that *MlNotch* is expressed well before

the first presumptive neuron is differentiated, we hypothesize that this embryonic expression is not

promoting terminal neuronal fate, although it could very well be involved in regulating early progenitor

cell populations. The localization of *MlNotch* expression at 24hpf in endodermal cells subjacent to comb

row cells — where the future gonads will develop — hints at a possible role in promoting primordial

germ cell fate.

Pharmacological inhibition of MlNotch points to an as-of-yet unknown mechanism of activation in

ctenophores

It is still unclear what activates *MlNotch*. The inability to identify an *MlNotch*-activating ligand has been

one of the primary reasons previous reports have concluded that ctenophores do not have a functional

Notch pathway[9,10,18]. However, the genome of the choanoflagellate species *S. dolichothecata* encodes a

protein with MNNL and DSL domains characteristic of *Delta-like*, contrary to previous work suggesting

that the *Delta-like* ligand was a Metazoan innovation[7]. Taken together with our ctenophore data, we see at

least two possibilities for the emergence of a bonafide *Delta-like* ligand: *Delta-like* could have emerged

prior to the Metaozan divergence and was subsequently lost in the ctenophore lineage, or we simply are

unable to identify *MlDelta-like* with current methods available to ctenophore researchers.

If *Delta-like* was lost in the ctenophore lineage, then *MlNotch* could either function independent

of a ligand or it could recognize an as-of-yet unidentified activating ligand. *Notch* activity can occur

during *Drosophila* oogenesis independent of ligand activation[43,44]. Perhaps *MlNotch* is capable of such a

ligand-independent function. It may even be possible that the ancestral *Notch* could have functioned as a

homodimer, acting as both signal and receiver, and *Delta-like* emerged after the ctenophore divergence. It

has been hypothesized that the DSL domains characteristic of *Delta-like* proteins shares ancestry with

LNR domains of the *Notch* receptor[3]. Furthermore, the DSL domain is extremely similar to EGF domains

— perhaps genes rich in EGF architecture could be viable candidates for identifying non-canonical

binding partners. The gene hit from our BLAST searches — ML24144a — is an intriguing candidate

given its EGF-rich and LamininG-rich extracellular domains, generally characteristic of cadherins.

Furthermore, upon re-examination of the genomic architecture surrounding ML24144a, we identified a

neighboring gene (ML24143a) that appears to harbor the intracellular domain of a proto-cadherin. It is

possible that these two genes are misannotated and are in fact a single proto-cadherin that is co-expressed

in cells presumably loaded with active *MlNotch*. This raises the intriguing possibility that the ancestral

Metazoan *Notch* interacted with cadherin-like proteins. N-cadherins have been shown to interact with

Wnt and *Notch* pathways to regulate neural progenitor cells[45]. Our results are consistent with previous

hypotheses that *Notch* signaling could have emerged from a cell adhesion module that predates

Metazoa[46]. Our HCR™ data show that *MlNotch* and *MlCad-like* transcripts are co-expressed in distinct

cell populations. *Delta-like* and *Jagged-like*, both classical *Notch* ligands, have been shown to be *cis*-

activators in synthetic cell cultures[47]. It may be possible that *Notch* communication originated as an

autocrine signaling module in single-celled eukaryotes that, due to its association with primitive

cadherins, evolved its juxtacrine functions at the base of animal multicellularity.

Alternatively, ctenophores could still have a *Delta-like* gene that has evolved such that traditional

sequence-based methods are not sensitive enough to identify homologs. Recent work has shown that

genes identified as "lineage-specific" may in fact be false-negatives due to the rapid rate in which some

genes have evolved such that homologs are unable to be detected[48]. Reasoning that bases homology on

sequence similarity or the presence/absence of diagnostic PFAM domains may be similarly flawed. We

therefore caution researchers from concluding that specific genes are missing or lineage-specific based on

bioinformatic inferences alone, particularly in species where existing genomic and transcriptomic data are

prone to errors or mis-annotations.

Conclusion

This study paints a more complex picture for *Notch* evolution and its activity during ctenophore

development than was previously supposed, laying the groundwork to enrich and expand future studies on

the evolution of signaling pathways. We have identified and characterized unreported *Notch* pathway

components encoded in the *M. leidyi* genome and clarified the developmental context in which these

components are expressed. Our results demonstrate that *MlNotch* expression during ctenophore

development is consistent with classical models of cell-cell communication, suggesting that its role as a

juxtacrine signal transducer may have evolved at the base of the Metazoan divergence, perhaps through a

co-option of a pre-existing cell adhesion module. Furthermore, our results highlight the importance of 1)

validating bioinformatic data with *in situ* characterizations before making generalized conclusions and 2)

expanding the repertoire of model systems to understand the evolution of developmental signaling

pathways. While a relative handful of model systems are powerful to study the basic characteristics of

signaling pathways, they offer an extremely limited scope for understanding how these pathways evolved.

Despite the unique phylogenetic position and intriguing biology of ctenophores, they remain relatively

understudied. *In situ* characterizations of signaling pathways in early-branching Metazoans are crucial for

understanding how they evolved their developmental functions.

Acknowledgments

419   This work was supported by the National Science Foundation (NSF grant IOS-1755364), the National
420   Aeronautics and Space Agency (NASA grant 80NSSC18K1067), and The Paul G. Allen Frontiers Group
421   (grant #12970). The authors thank Dr. Alicia Boyd for her valuable comments while drafting this
422   manuscript.

Author contributions

425   Conceptualization, B.F., F.H., and M.Q.M; methodology, B.F., F.H., C.B., and M.Q.M.; validation, B.F.,

426   F.H., and C.B.; formal analysis, B.F., F.H., and C.B.; investigation, B.F., F.H., and C.B.; resources, B.F.,

427   F.H., C.B., J.S., and M.Q.M; data curation, B.F., F.H., and C.B.; writing – original draft, B.F; writing –

428   review and editing, B.F., F.H., C.B., J.S., and M.Q.M.; visualization, B.F., F.H., and C.B.; supervision,

429   F.H. and M.Q.M.; project administration, J.S. and M.Q.M.; funding acquisition, J.S and M.Q.M.

Declaration of interests

The authors declare no competing interests.

STAR Methods

Resource availability

Lead contact

Further information and requests for resources and reagents will be fulfilled by the lead contact, Brent

Foster (bf400@exeter.ac.uk), or Mark Q. Martindale (mqmartin@ufl.edu).

Materials availability

Animals, reagents, and probes generated by this study are available from Mark Q. Martindale

(mqmartin@ufl.edu).

Data and code availability

This study did not generate any new code.

Key resources table

Reagent or Resource	Source	Identifier
Animal husbandry		
Kreisel		
Grade A Brine Shrimp Eggs	Brine Shrimp Direct	NA
S. Presso, Selco®	Artemia International	NA
Forceps		
Kits		
PGEM-T vector	Promega	Cat #A3600
GeneJET Plasmid Miniprep kit	ThermoFisher	Cat #K0503
MEGAscript SP6	ThermoFisher	Cat #AM1330
MEGAscript T7	ThermoFisher	Cat #AM1334

Fixatives

ACME	PBS	
	Methanol	
	Glacial Acetic Acid	
	Glycerol	
Rain-X®	Amazon, Inc.	
PFA	Electron Microscopy Sciences	Cat #15710
Gluteraldehyde	Electron Microscopy Sciences	Cat # 16216-10
Methanol	Fisher Scientific	Cat #A4121

In situ **and** *HCR probes*

In situ **hybridization solutions**

Proteinase K	Invitrogen	Cat# 25530049
Glycine	Sigma Aldrich	Cat# 410225
Triethanolamine	Sigma Aldrich	Cat #90279
Acetic anhydride	Sigma Aldrich	Cat# 320102
Hybridization buffer (with salmon sperm DNA)	Formamide (Sigma Aldrich)	Cat #FX0420
	20X SSC (pH4.5)	NA
	20mg/ml heparin	NA
	100% Tween-20 (ThermoFisher)	Cat # P2287
	20% SDS (ThermoFisher)	Cat# BP166-500
	10mg/ml salmon sperm DNA (Invitrogen)	Cat #AM9680
Hybridization buffer (without salmon sperm DNA)	Formamide (Sigma Aldrich)	Cat #FX0420
	20X SSC (pH4.5)	NA
	20mg/ml heparin	NA
	100% Tween-20 (ThermoFisher)	Cat # P2287
	20% SDS (ThermoFisher)	Cat# BP166-500
2X Saline Sodium-Citrate (SSC), ph7	Sigma Aldrich	Cat# S6639
1X PBS-Tween	100ml 10X PBS	NA
	899ml Milli-Q H_2O	
	1ml Tween-20	

Blocking Buffer	Roche	Cat# 11096176001
Maleic Acid Buffer	6.905g maleic acid	NA
	4.38g NaCl	
	Fill to 1L Milli-Q H_2O	
AP Buffer	100mM NaCl	NA
	100mM Tris-Cl	
	0.5% Tween-20	
	DEPC-H_2O	
AP Buffer with $MgCl_2$	100mM NaCl	NA
	50mM $MgCl_2$	
	100mM Tris-Cl	
	0.5% Tween-20	
	DEPC-H_2O	
NBT (100mg/ml)	Roche	Cat #11383213001
BCIP (50mg/ml)	Roche	Cat #11383223001
AP Developing Buffer	AP Buffer with $MgCl_2$	NA
	NBT (0.33mg/ml)	
	BCIP (0.165mg/ml)	

Antibodies

Anti-DIG	Roche	Cat #11093274910
Cleaved Notch1 antibody	Cell Signaling Technology	Cat #4147
goat-α-Rabbit secondary antibody, AP-conjugated	Invitrogen	Cat #31346

HCR solutions

HCR™ Probe Hybridization Buffer v3.0	Molecular Instruments, Inc.	NA
HCR™ Probe Wash Buffer v3.0	Molecular Instruments, Inc.	NA
HCR™ Amplifier Buffer v3.0	Molecular Instruments, Inc.	NA
HCR™ Amplifier (3.0): B2-488	Molecular Instruments, Inc.	NA
HCR™ Amplifier (3.0): B1-647	Molecular Instruments, Inc.	NA
HCR™ Amplifier (3.0): B3-647	Molecular Instruments, Inc.	NA

Animal collection and husbandry

Adult *Mnemiopsis leidyi* were collected in the Matanzas River estuarine system near St. Augustine and

Beverly Beach, FL, and transported to the Whitney Laboratory. Animals were transitioned into 2/3X

filtered sea water (FSW) (approximately 25 ppt) and kept in a self-circulating Kreisel system in constant

light. Adults were fed several times per day with artemia brine shrimp enriched with S. Presso (Seclo®).

Prior to spawning, adult animals were placed in the dark for 3 hours and then transitioned to glass finger

bowls. When spawning began, roughly one hour later, embryos were collected in a 30-minute window

and allowed to develop until 4 hours post fertilization (hpf), after which the vitelline membrane was

manually removed with sharp forceps. Embryos were then allowed to grow up until 8hpf or 24hpf when

they were fixed for *in situ* hybridization or HCR[TM].

Gene Identification and Domain Analysis

M. leidyi genes were identified in the *Mnemiopsis* Genome Portal from gene identification numbers[18].

Orthology was confirmed via reciprocal BLAST against the NCBI database[49] and PFAM analysis. PFAM

architectures were evaluated using a combination of SMART[50] and Prosite[51–53]. Domain significance was

determined by E-value (SMART) or score threshold (Prosite) (see Supplementary File 1 for PFAM

scores). Juxta-transmembrane domains were confirmed using TMHMM 2.0[54] and the Conserved Domain

Search (CDS) from NCBI[55–57] (E-value ≤ 0.01). Nuclear localization signals (NLS) were identified with

NL Stradamus[58] (significant posterior probability ≥ 0.6). PEST domains were identified with EPestfind

(http://emboss.bioinformatics.nl/cgi-bin/emboss/epestfind; threshold ≥ +5.0 score). Putative S1, S2, and

S3 recognition sites were identified from previous publications[59]. *MlNotch* juxta-transmembrane domain

(JTMD) was aligned with reported *TaNotch*, *NvNotch*, *HsNotch1, HsNotch4, HsNotch2, HsNotch3,*

DmNotch JTMDs (see Supplementary File 2).

Phylogeny

Sequences for ctenophores (*Boroe ovata*, *Mnemiopsis leidyi*, and *Hormiphora Califoniensis*) and

choanoflagellates (*Capsaspora owczarzaki* and *Mylnosiga fluctuans*) were added to existing

alignments[3,16] using MAFFT[60,61] (see Supplementary File 3 for alignments). EGF repeat domains were

excluded from the alignments due to their variability across species[3]. Bootstrapped maximum likelihood

phylogenies were prepared with RAxMLGUI2.0 using a WAG substitution matrix and executing 500 ML

tree inferences and thereafter a thorough ML search. All free model parameters were estimated by

RAxML using a gamma model of rate heterogeneity. Bayesian analysis was performed using MrBayes

3.2.6 in Geneious[62] using the WAG fixed model and gamma rate variation. Markov Chain Monte Carlo

runs of 1,100,000 generations were calculated with trees sampled every 200 generations and with a prior

burn-in of 110,000 generations. The best-scoring tree was visualized using FigTree (v1.4.2;

http://tree.bio.ed.ac.uk/software/figtree) and annotated in Adobe Illustrator 2022. Juxtatransmembrane

alignments were analyzed and visualized with Jalview[63]. AlphaFold predictions were made with

ColabFold v1.5.5: AlphaFold2 using MMseqs2[64] (see Supplementary File 3).

Developmental RNAseq for M. leidyi

Developmental RNA-seq data[21] was accessed via the *Mnemiopsis* Genome Portal[18] and plotted in Excel

(see Supplementary File 5).

Cloning of in situ probe templates

Probe preparation and colorimetric *in situ* hybridization were conducted as previously described[65], with

slight modifications. Gene fragments were ordered with flanking SP7 and T7 promoters from Twist

Biosciences, Inc. (see Supplementary File 6 for fragment sequences) and directly PCR amplified for

probe synthesis using SP6 and T7 primers. Some genes were amplified from cDNA using gene specific

primers. These were then ligated into Promega PGEM-T vector (A3600), then transformed and verified

via colony PCR. Plasmids were purified using the ThermoFisher GeneJET Plasmid Miniprep kit

(#K0503) and analyzed by Sanger sequencing. DIG-labled RNA probes were transcribed using Invitrogen

MEGAscript SP6 (#AM1330) or T7 (#AM1334) kits according to manufacturer directions.

Colorimetric whole-mount in situ hybridization

Whole-mount *in situ* hybridizations were conducted as previously described[65] with slight modifications.

Animals were fixed in ACME solution[66] for one hour at room temperature, then subsequently washed in

PBS-Tween (see Supplementary File 7 for the detailed protocol). In brief, samples underwent a secondary

fixation in 4% PFA and 0.8% glutaraldehyde for five minutes at room temperature, then transitioned into

4% PFA solution overnight at 4ºC. The following day, samples were transitioned to 100% methanol and

stored at -20ºC. Samples were subjected to Proteinase K digestion and washes in glycine, triethanolamine,

and acetic anhydride before proceeding with an overnight pre-hybridization at 63ºC. Probes were

hybridized using 1ng/µl concentrations at 63ºC for 24–72 hours. Samples were transitioned into SSC and

PBS-Tween solutions and then blocked in Roche Blocking Buffer (Roche, #11096176001) overnight at

4ºC. The following day, samples were incubated with DIG-antibody (Sigma, #11093274910) diluted in

Roche Blocking Buffer (Roche, #11096176001, 1:5000) overnight at 4ºC. Samples were then washed in

PBS-Tween and washed in alkaline phosphatase (AP) developing buffer (1M NaCl, 2M MgCl2, 1M Tris,

Tween-20, NBT, BCIP) to develop DIG-labeled probes.

Hybridization chain reaction RNA fluorescence in situ hybridization (HCR RNA-FISH)

HCR probes were made using a customized Python code[67] and ordered through IDT, Inc., and diluted to

100nM stock concentration. Animals were fixed at 8hpf using Rain-X® primary fixation[68] for between

20–60 minutes at room temperature, transitioned into PBS-Tween and then subjected to an overnight

secondary fixation in 4% PFA at 4ºC. Samples were transitioned into 100% methanol and stored at -20ºC

until ready for use, after which they were transitioned back into PBS-Tween. We then followed a standard

HCR[TM] RNA-FISH protocol as described previously[23,69,70], with slight modifications (see Supplementary

File 8 for the detailed protocol). Samples were washed in HCR™ Probe Hybridization Buffer v3.0

(Molecular Instruments, Inc.) and incubated at 37ºC. Probes were diluted to 20nM in HCR™ Probe

Hybridization Buffer v3.0 (Molecular Instruments, Inc.) and were allowed to hybridize overnight at 37ºC,

then washed with HCR™ Probe Wash Buffer v3.0 (Molecular Instruments, Inc.). Samples were then

washed in HCR™ Amplifier Buffer v3.0 (Molecular Instruments, Inc). Hairpins were heated at 95ºC for

90 seconds, then allowed to cool to room temperature in the dark before being added to samples and

allowed to amplify overnight at room temperature in the dark. Samples were then washed in 5X SSC and

counterstained with Hoechst (1:1000).

Imaging

Colorimetric *in situ* samples were mounted in 80% glycerol and imaged on a Zeiss Imager M2. Images

were post-processed in Adobe Photoshop 2022. HCR[TM] samples were mounted in 5X SSC and imaged on

a Zeiss 710 laser scanning confocal microscope. Confocal slices and Z-stacks were analyzed in FIJI[71].

Pharmacological Inhibition

Embryos were de-membranated between 3–4hpf, then treated in 10µM DAPT (Sigma, D5942), 25µM

LY411575 (Selleck Chemicals, S2714), and 1% DMSO (Sigma, D4540) in 2/3X FSW for approximately

4 hours. We then adapted a standard protocol for antibody staining of *Mnemiopsis leidyi* specimens (see

Supplementary File 9 for the detailed antibody staining protocol). In brief, embryos were fixed in Rain-

X® (200µl per 1ml) for 20 minutes, then washed into PBS-Tween and stored in 4% paraformaldehyde at

4ºC overnight. The following day, embryos were washed 4 times in PBS-Triton, then permeabilized with

4 washes in PBT and blocked in 5% Normal Goat Serum (NGS) for 1 hour at room temperature. Cleaved

Notch1 antibody (Cell Signaling Technology, #4147, 1:500) was diluted in 5% NGS and incubated with

samples overnight at 4ºC. Embryos were washed 7 times in PBT, then incubated with goat-α-Rabbit

secondary antibody conjugated with AP (Invitrogen, #31346, 1:3500) and incubated overnight at 4ºC. The

specimens were then washed twice in AP developing buffer without $MgCl_2$ (1M NaCl, 1M Tris, Tween-

20), then washed twice in AP buffer with $MgCl_2$ (1M NaCl, 2M $MgCl_2$, 1M Tris, Tween-20), then

developed in AP buffer with NBT and BCIP.

Supplemental information

References

1.   Borggrefe, T., and Oswald, F. (2009). The Notch signaling pathway: Transcriptional regulation at
559   Notch target genes. Cell. Mol. Life Sci. *66*, 1631–1646. https://doi.org/10.1007/s00018-009-8668-7.

560   2.   Sachan, N., Sharma, V., Mutsuddi, M., and Mukherjee, A. (2024). Notch signalling: multifaceted role
561   in development and disease. The FEBS Journal *291*, 3030–3059. https://doi.org/10.1111/febs.16815.

562   3.   Gazave, E., Lapébie, P., Richards, G.S., Brunet, F., Ereskovsky, A.V., Degnan, B.M., Borchiellini, C.,
563   Vervoort, M., and Renard, E. (2009). Origin and evolution of the Notch signalling pathway: An
564   overview from eukaryotic genomes. BMC Evolutionary Biology *9*, 1–27.
565   https://doi.org/10.1186/1471-2148-9-249/FIGURES/12.

566   4.   Babonis, L.S., and Martindale, M.Q. (2017). Phylogenetic evidence for the modular evolution of
567   metazoan signalling pathways. Philosophical Transactions of the Royal Society B: Biological
568   Sciences *372*. https://doi.org/10.1098/RSTB.2015.0477.

569   5.   King, N., Westbrook, M.J., Young, S.L., Kuo, A., Abedin, M., Chapman, J., Fairclough, S., Hellsten,
570   U., Isogai, Y., Letunic, I., et al. (2008). The genome of the choanoflagellate Monosiga brevicollis and
571   the origin of metazoans. Nature *451*, 783–788. https://doi.org/10.1038/nature06617.

572   6.   Suga, H., and Ruiz-Trillo, I. (2013). Development of ichthyosporeans sheds light on the origin of
573   metazoan multicellularity. Developmental Biology *377*, 284–292.
574   https://doi.org/10.1016/j.ydbio.2013.01.009.

575   7.   Richter, D.J., Fozouni, P., Eisen, M.B., and King, N. (2018). Gene family innovation, conservation
576   and loss on the animal stem lineage. eLife *7*, e34226. https://doi.org/10.7554/eLife.34226.

8.   Lv, Y., Pang, X., Cao, Z., Song, C., Liu, B., Wu, W., and Pang, Q. (2024). Evolution and Function of
the Notch Signaling Pathway: An Invertebrate Perspective. IJMS *25*, 3322.
https://doi.org/10.3390/ijms25063322.

9.   Lövy, A., Kyslík, J., Picard-Sánchez, A., Holzer, A.S., Krejčí, A., and Fiala, I. (2025). The Notch
pathway in Metazoa: a comparative analysis across cnidarians and beyond. EvoDevo *16*, 8.
https://doi.org/10.1186/s13227-025-00246-6.

10.   He, X., Wu, F., Zhang, L., Li, L., and Zhang, G. (2021). Comparative and evolutionary analyses
reveal conservation and divergence of the notch pathway in lophotrochozoa. Scientific Reports *11*.
https://doi.org/10.1038/s41598-021-90800-8.

11.   Richards, G.S., and Degnan, B.M. (2012). The expression of Delta ligands in the sponge
Amphimedon queenslandica suggests an ancient role for Notch signaling in metazoan development.
EvoDevo *3*, 15. https://doi.org/10.1186/2041-9139-3-15.

12.   Richards, G.S., Simionato, E., Perron, M., Adamska, M., Vervoort, M., and Degnan, B.M. (2008).
Sponge Genes Provide New Insight into the Evolutionary Origin of the Neurogenic Circuit. Current
Biology *18*, 1156–1161. https://doi.org/10.1016/j.cub.2008.06.074.

13.   Marlow, H., Roettinger, E., Boekhout, M., and Martindale, M.Q. (2012). Functional roles of Notch
signaling in the cnidarian Nematostella vectensis. Developmental Biology *362*, 295–308.
https://doi.org/10.1016/J.YDBIO.2011.11.012.

14.   Layden, M.J., and Martindale, M.Q. (2014). Non-canonical Notch signaling represents an ancestral
mechanism to regulate neural differentiation. EvoDevo *5*, 30. https://doi.org/10.1186/2041-9139-5-
30.

15.   Haillot, E., Lebedeva, T., Steger, J., Genikhovich, G., Montenegro, J.D., Cole, A.G., and Technau, U.
(2024). Notch, β-catenin and MAPK signaling segregate endoderm and mesoderm in the diploblast
Nematostella vectensis. Preprint, https://doi.org/10.1101/2024.10.29.620801
https://doi.org/10.1101/2024.10.29.620801.

16.   Gahan, J.M., Schnitzler, C.E., DuBuc, T.Q., Doonan, L.B., Kanska, J., Gornik, S.G., Barreira, S.,
Thompson, K., Schiffer, P., Baxevanis, A.D., et al. (2017). Functional studies on the role of Notch
signaling in Hydractinia development. Developmental Biology *428*, 224–231.
https://doi.org/10.1016/J.YDBIO.2017.06.006.

17.   Käsbauer, T., Towb, P., Alexandrova, O., David, C.N., Dall'Armi, E., Staudigl, A., Stiening, B., and
Böttger, A. (2007). The Notch signaling pathway in the cnidarian Hydra. Developmental biology *303*,
376–390. https://doi.org/10.1016/J.YDBIO.2006.11.022.

18.   Ryan, J.F., Pang, K., Schnitzler, C.E., Nguyen, A.D., Moreland, R.T., Simmons, D.K., Koch, B.J.,
Francis, W.R., Havlak, P., Smith, S.A., et al. (2013). The genome of the ctenophore Mnemiopsis
leidyi and its implications for cell type evolution. Science *342*.
https://doi.org/10.1126/SCIENCE.1242592.

19.   Moroz, L.L., Kocot, K.M., Citarella, M.R., Dosung, S., Norekian, T.P., Povolotskaya, I.S.,
Grigorenko, A.P., Dailey, C., Berezikov, E., Buckley, K.M., et al. (2014). The ctenophore genome and
the evolutionary origins of neural systems. Nature *510*, 109–114.
https://doi.org/10.1038/NATURE13400.

20. Schultz, D.T., Haddock, S.H.D., Bredeson, J.V., Green, R.E., Simakov, O., and Rokhsar, D.S. (2023).
Ancient gene linkages support ctenophores as sister to other animals. Nature *618*, 110–117.
https://doi.org/10.1038/s41586-023-05936-6.

21. Levin, M., Anavy, L., Cole, A.G., Winter, E., Mostov, N., Khair, S., Senderovich, N., Kovalev, E.,
Silver, D.H., Feder, M., et al. (2016). The mid-developmental transition and the evolution of animal
body plans. Nature *531*, 637–641. https://doi.org/10.1038/nature16994.

22. Davidson, P.L., Koch, B.J., Schnitzler, C.E., Henry, J.Q., Martindale, M.Q., Baxevanis, A.D., and
Browne, W.E. (2017). The maternal-zygotic transition and zygotic activation of the *Mnemiopsis*
leidyi genome occurs within the first three cleavage cycles. Molecular Reproduction Devel *84*, 1218–
1229. https://doi.org/10.1002/mrd.22926.

23. Choi, H.M.T., Schwarzkopf, M., Fornace, M.E., Acharya, A., Artavanis, G., Stegmaier, J., Cunha, A.,
and Pierce, N.A. (2018). Third-generation *in situ* hybridization chain reaction: multiplexed,
quantitative, sensitive, versatile, robust. Development *145*, dev165753.
https://doi.org/10.1242/dev.165753.

24. Schnitzler, C.E., Simmons, D.K., Pang, K., Martindale, M.Q., and Baxevanis, A.D. (2014).
Expression of multiple Sox genes through embryonic development in the ctenophore Mnemiopsis
leidyi is spatially restricted to zones of cell proliferation. EvoDevo *5*. https://doi.org/10.1186/2041-
9139-5-15.

25. Sjöqvist, M., and Andersson, E.R. (2019). Do as I say, Not(ch) as I do: Lateral control of cell fate.
Developmental Biology *447*, 58–70. https://doi.org/10.1016/j.ydbio.2017.09.032.

26. Bray, S.J., and Bigas, A. (2025). Modes of Notch signalling in development and disease. Nat Rev
Mol Cell Biol *26*, 522–537. https://doi.org/10.1038/s41580-025-00835-2.

27. Morohashi, Y., Kan, T., Tominari, Y., Fuwa, H., Okamura, Y., Watanabe, N., Sato, C., Natsugari, H.,
Fukuyama, T., Iwatsubo, T., et al. (2006). C-terminal Fragment of Presenilin Is the Molecular Target
of a Dipeptidic γ-Secretase-specific Inhibitor DAPT (N-[N-(3,5-Difluorophenacetyl)-L-alanyl]-S-
phenylglycine t-Butyl Ester). Journal of Biological Chemistry *281*, 14670–14676.
https://doi.org/10.1074/jbc.M513012200.

28. Kidd, S., and Lieber, T. (2002). Furin cleavage is not a requirement for Drosophila Notch function.
Mechanisms of Development *115*, 41–51. https://doi.org/10.1016/S0925-4773(02)00120-X.

29. Bush, G., diSibio, G., Miyamoto, A., Denault, J.-B., Leduc, R., and Weinmaster, G. (2001). Ligand-
Induced Signaling in the Absence of Furin Processing of Notch1. Developmental Biology *229*, 494–
502. https://doi.org/10.1006/dbio.2000.9992.

30. Zavadil, J., Cermak, L., Soto-Nieves, N., and Böttinger, E.P. (2004). Integration of TGF-β/Smad and
Jagged1/Notch signalling in epithelial-to-mesenchymal transition. EMBO J *23*, 1155–1165.
https://doi.org/10.1038/sj.emboj.7600069.

31. Edenfeld, G., Altenhein, B., Zierau, A., Cleppien, D., Krukkert, K., Technau, G., and Klämbt, C.
(2007). Notch and Numb are required for normal migration of peripheral glia in Drosophila.
Developmental Biology *301*, 27–37. https://doi.org/10.1016/j.ydbio.2006.11.013.

32. Alhashem, Z., Feldner-Busztin, D., Revell, C., Alvarez-Garcillan Portillo, M., Camargo-Sosa, K.,
Richardson, J., Rocha, M., Gauert, A., Corbeaux, T., Milanetto, M., et al. (2022). Notch controls the
cell cycle to define leader versus follower identities during collective cell migration. eLife *11*,
e73550. https://doi.org/10.7554/eLife.73550.

33. Hashimoto-Torii, K., Torii, M., Sarkisian, M.R., Bartley, C.M., Shen, J., Radtke, F., Gridley, T.,
Šestan, N., and Rakic, P. (2008). Interaction between Reelin and Notch Signaling Regulates Neuronal
Migration in the Cerebral Cortex. Neuron *60*, 273–284. https://doi.org/10.1016/j.neuron.2008.09.026.

34. Reitzel, A.M., Pang, K., and Martindale, M.Q. (2016). Developmental expression of "germline"- and
"sex determination"-related genes in the ctenophore Mnemiopsis leidyi. EvoDevo *7*.
https://doi.org/10.1186/S13227-016-0051-9.

35. Richards, G.S., and Rentzsch, F. (2015). Regulation of Nematostella neural progenitors by SoxB,
Notch and bHLH genes. Development (Cambridge, England) *142*, 3332.
https://doi.org/10.1242/DEV.123745.

36. Moneer, J., Siebert, S., Krebs, S., Cazet, J., Prexl, A., Pan, Q., Juliano, C., and Böttger, A. (2021).
Differential gene regulation in DAPT-treated Hydra reveals candidate direct Notch signalling targets.
Journal of cell science *134*. https://doi.org/10.1242/jcs.258768.

37. Steichele, M., Sauermann, L., Pan, Q., Moneer, J., De La Porte, A., Heß, M., Mercker, M., Strube, C.,
Flaswinkel, H., Jenewein, M., et al. (2025). Notch signaling mediates between two pattern-forming
processes during head regeneration in *Hydra*. Life Sci. Alliance *8*, e202403054.
https://doi.org/10.26508/lsa.202403054.

38. Koenig, K.M., Sun, P., Meyer, E., and Gross, J.M. (2016). Eye development and photoreceptor
differentiation in the cephalopod *Doryteuthis pealeii*. Development, dev.134254.
https://doi.org/10.1242/dev.134254.

39. Napoli, F.R., Daly, C.M., Neal, S., McCulloch, K.J., Zaloga, A.R., Liu, A., and Koenig, K.M. (2022).
Cephalopod retinal development shows vertebrate-like mechanisms of neurogenesis. Current Biology
32, 5045-5056.e3. https://doi.org/10.1016/j.cub.2022.10.027.

40. Arefin, B., Parvin, F., Bahrampour, S., Stadler, C.B., and Thor, S. (2019). Drosophila Neuroblast
Selection Is Gated by Notch, Snail, SoxB, and EMT Gene Interplay. Cell Reports *29*, 3636-3651.e3.
https://doi.org/10.1016/j.celrep.2019.11.038.

41. Holmberg, J., Hansson, E., Malewicz, M., Sandberg, M., Perlmann, T., Lendahl, U., and Muhr, J.
(2008). SoxB1 transcription factors and Notch signaling use distinct mechanisms to regulate
proneural gene function and neural progenitor differentiation. Development *135*, 1843–1851.
https://doi.org/10.1242/dev.020180.

42. Norekian, T.P., and Moroz, L.L. (2021). Development of the nervous system in the early hatching
larvae of the ctenophore Mnemiopsis leidyi. Journal of Morphology *282*, 1466–1477.
https://doi.org/10.1002/JMOR.21398.

43. Palmer, W.H., and Deng, W.-M. (2015). Ligand-Independent Mechanisms of Notch Activity. Trends
in Cell Biology *25*, 697–707. https://doi.org/10.1016/j.tcb.2015.07.010.

44. Palmer, W.H., Jia, D., and Deng, W.-M. (2014). Cis-interactions between Notch and its ligands block
ligand-independent Notch activity. eLife *3*, e04415. https://doi.org/10.7554/eLife.04415.

45. Miyamoto, Y., Sakane, F., and Hashimoto, K. (2015). N-cadherin-based adherens junction regulates
the maintenance, proliferation, and differentiation of neural progenitor cells during development. Cell
Adhesion & Migration *9*, 183–192. https://doi.org/10.1080/19336918.2015.1005466.

46. Murata, A., and Hayashi, S.-I. (2016). Notch-Mediated Cell Adhesion. Biology.
https://doi.org/10.3390/biology5010005.

47. Nandagopal, N., Santat, L.A., and Elowitz, M.B. (2019). Cis-activation in the Notch signaling
pathway. eLife *8*. https://doi.org/10.7554/ELIFE.37880.

48. Weisman, C.M., Murray, A.W., and Eddy, S.R. (2020). Many, but not all, lineage-specific genes can
be explained by homology detection failure. PLoS Biol *18*, e3000862.
https://doi.org/10.1371/journal.pbio.3000862.

49. Sayers, E.W., Beck, J., Bolton, E.E., Brister, J.R., Chan, J., Connor, R., Feldgarden, M., Fine, A.M.,
Funk, K., Hoffman, J., et al. (2025). Database resources of the National Center for Biotechnology
Information in 2025. Nucleic Acids Research *53*, D20–D29. https://doi.org/10.1093/nar/gkae979.

50. Letunic, I., Khedkar, S., and Bork, P. (2021). SMART: recent updates, new developments and status
in 2020. Nucleic Acids Research *49*, D458–D460. https://doi.org/10.1093/nar/gkaa937.

51. Sigrist, C.J.A., De Castro, E., Cerutti, L., Cuche, B.A., Hulo, N., Bridge, A., Bougueleret, L., and
Xenarios, I. (2012). New and continuing developments at PROSITE. Nucleic Acids Research *41*,
D344–D347. https://doi.org/10.1093/nar/gks1067.

52. Sigrist, C.J.A. (2002). PROSITE: A documented database using patterns and profiles as motif
descriptors. Briefings in Bioinformatics *3*, 265–274. https://doi.org/10.1093/bib/3.3.265.

53. De Castro, E., Sigrist, C.J.A., Gattiker, A., Bulliard, V., Langendijk-Genevaux, P.S., Gasteiger, E.,
Bairoch, A., and Hulo, N. (2006). ScanProsite: detection of PROSITE signature matches and
ProRule-associated functional and structural residues in proteins. Nucleic Acids Research *34*, W362–
W365. https://doi.org/10.1093/nar/gkl124.

54. Krogh, A., Larsson, B., Von Heijne, G., and Sonnhammer, E.L.L. (2001). Predicting transmembrane
protein topology with a hidden markov model: application to complete genomes11Edited by F.
Cohen. Journal of Molecular Biology *305*, 567–580. https://doi.org/10.1006/jmbi.2000.4315.

55. Wang, J., Chitsaz, F., Derbyshire, M.K., Gonzales, N.R., Gwadz, M., Lu, S., Marchler, G.H., Song,
J.S., Thanki, N., Yamashita, R.A., et al. (2023). The conserved domain database in 2023. Nucleic
Acids Research *51*, D384–D388. https://doi.org/10.1093/nar/gkac1096.

56. Lu, S., Wang, J., Chitsaz, F., Derbyshire, M.K., Geer, R.C., Gonzales, N.R., Gwadz, M., Hurwitz,
D.I., Marchler, G.H., Song, J.S., et al. (2020). CDD/SPARCLE: the conserved domain database in
2020. Nucleic Acids Research *48*, D265–D268. https://doi.org/10.1093/nar/gkz991.

57. Marchler-Bauer, A., Bo, Y., Han, L., He, J., Lanczycki, C.J., Lu, S., Chitsaz, F., Derbyshire, M.K.,
Geer, R.C., Gonzales, N.R., et al. (2017). CDD/SPARCLE: functional classification of proteins via

subfamily domain architectures. Nucleic Acids Res *45*, D200–D203.
https://doi.org/10.1093/nar/gkw1129.

58. Nguyen Ba, A.N., Pogoutse, A., Provart, N., and Moses, A.M. (2009). NLStradamus: a simple
Hidden Markov Model for nuclear localization signal prediction. BMC Bioinformatics *10*, 202.
https://doi.org/10.1186/1471-2105-10-202.

59. Van Tetering, G., and Vooijs, M. (2011). Proteolytic Cleavage of Notch: “HIT and
RUN” CMM *11*, 255–269. https://doi.org/10.2174/156652411795677972.

60. Katoh, K. (2002). MAFFT: a novel method for rapid multiple sequence alignment based on fast
Fourier transform. Nucleic Acids Research *30*, 3059–3066. https://doi.org/10.1093/nar/gkf436.

61. Katoh, K., and Standley, D.M. (2013). MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. Molecular Biology and Evolution *30*, 772–780.
https://doi.org/10.1093/molbev/mst010.

62. Huelsenbeck, J.P., and Ronquist, F. (2001). MRBAYES: Bayesian inference of phylogenetic trees.
Bioinformatics *17*, 754–755. https://doi.org/10.1093/bioinformatics/17.8.754.

63. Waterhouse, A.M., Procter, J.B., Martin, D.M.A., Clamp, M., and Barton, G.J. (2009). Jalview
Version 2—a multiple sequence alignment editor and analysis workbench. Bioinformatics *25*, 1189–
1191. https://doi.org/10.1093/bioinformatics/btp033.

64. Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., and Steinegger, M. (2022).
ColabFold: making protein folding accessible to all. Nat Methods *19*, 679–682.
https://doi.org/10.1038/s41592-022-01488-1.

65. Mitchell, D.G., Edgar, A., Mateu, J.R., Ryan, J.F., and Martindale, M.Q. (2024). The ctenophore
Mnemiopsis leidyi deploys a rapid injury response dating back to the last common animal ancestor.
Commun Biol *7*, 203. https://doi.org/10.1038/s42003-024-05901-7.

66. García-Castro, H., Kenny, N.J., Iglesias, M., Álvarez-Campos, P., Mason, V., Elek, A., Schönauer, A.,
Sleight, V.A., Neiro, J., Aboobaker, A., et al. (2021). ACME dissociation: a versatile cell fixation-
dissociation method for single-cell transcriptomics. Genome Biol *22*, 89.
https://doi.org/10.1186/s13059-021-02302-5.

67. Kuehn, E., Clausen, D.S., Null, R.W., Metzger, B.M., Willis, A.D., and Özpolat, B.D. (2022).
Segment number threshold determines juvenile onset of germline cluster expansion in *Platynereis*
dumerilii. J Exp Zool Pt B *338*, 225–240. https://doi.org/10.1002/jez.b.23100.

68. Mitchell, D.G., Edgar, A., and Martindale, M.Q. (2021). Improved histological fixation of gelatinous
marine invertebrates. Front Zool *18*, 29. https://doi.org/10.1186/s12983-021-00414-z.

69. Wang, Y., Zeng, Y., Saka, S.K., Xie, W., Goldaracena, I., Kohman, R.E., Yin, P., and Church, G.M.
(2020). Multiplexed in situ protein imaging using DNA-barcoded antibodies with
extended hybridization chain reactions. bioRxiv, 274456. https://doi.org/10.1101/274456.

70. Tsuneoka, Y., and Funato, H. (2020). Modified in situ Hybridization Chain Reaction Using Short
Hairpin DNAs. Front. Mol. Neurosci. *13*, 75. https://doi.org/10.3389/fnmol.2020.00075.

71. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image
analysis. Nature Methods 2012 9:7 *9*, 676–682. https://doi.org/10.1038/nmeth.2019.

A

Bilateria

Cnidaria

Porifera

Ctenophora

?

Choanoflagellata

Fungi

Amoebozoa

B

Notch OFF

Signal-sending Cell

Delta-like

Signal-receiving Cell

NECD Notch
NICD

SMRT
SuH ⊣ HES

Canonical

Notch ON

Signal-sending Cell

Neuralized
Mindbomb

Signal-receiving Cell

NICD NECD
ADAM10 S2
ADAM17

NECD

NICD NICD S3
γ-Secretase

Furin-like
O-Fut S1 NECD
NICD Sudx Deltex → ⊣ Numb
Notchless
Fringe NICD
SuH → HES
MAM

C

Notch OFF?

Signal-sending Cell

Delta-like/ EGF-rich?

Signal-receiving Cell

NECD Notch
NICD

SuH → HES

Hypothesized *M. leidyi*

Notch ON?

Signal-sending Cell

Mindbomb

Signal-receiving Cell

NICD NECD
S2
ADAM10

NICD NICD S3
γ-Secretase

Furin-like?
S1 NECD
NICD

NICD
SuH → HES

D

Median TPMs

0   1   2   3   4   4.5   5   5.5   6   6.5   7   7.5   8   8.5   9   10   11   12   13   14   15   16   17   18   19   20

Hours Post Fertilization (hpf)

MlNotch ∗
MlDelta-like1
EGF-rich1 ∗
MlFurin-like1
MlFurin-like2 ∗
MlADAM17
MlADAM10
MlPresenillin
MlNicastrin
MlAPH1
MlPen2
MlSuH
MlNumb
MlHES1
MlHES2 ∗
MlMindbomb ∗
MlDeltex
MlNedd4/Sudx
MlNotchless

A

EGF repeats | LNR repeats | NODP | J/TM | Ankyrin repeats | NLS | PEST

B

Mle NRR + NICD

C

Juxta-transmembrane Domains

Receptor & candidate ligand

MlNotch *MlCad-like*

Putative regulatory proteases

S1 S1 S2 S3

MlFurin-like1 *MlFurin-like2* *MlADAM10* *MlPresenillin*

Canonical co-factor & transcriptional target

MlSuH *MlHES2*

A — 8hpf — N = 43/43
B — 8hpf — N = 49/49
C — 8hpf — N = 31/36
D — 8hpf — N = 23/27
E — 8hpf — N = 38/42
F — 8hpf — N = 60/63
G — 8hpf — N = 57/59
H — 8hpf — N = 31/31

A' — 8hpf
B' — 8hpf
C' — 8hpf
D' — 8hpf
E' — 8hpf
F' — 8hpf
G' — 8hpf
H' — 8hpf

A

Signal-receiving Cell

Furin-like

S1

NICD NECD

NICD NECD

S2

ADAM10
ADAM17

NICD

NICD

S3

γ-Secretase

DAPT
LY411575

X

NICD

AP

α-NICD

Shuck

DMSO
LY411575
DAPT

0   1   2   3   4   5   6   7   8

Hours Post Fertilization (hpf)

1% DMSO

10µM DAPT

25µM LY411575

Anti-NICD

B   8hpf

N = 16/16

C   8hpf

N = 12/18

D   8hpf

N = 12/13

B'

C'

D'

B''   *

i.

ii.

i.   ii.