March 12, 2026

Evidence for holocentric centromeres in the early branching apicomplexan

parasite *Cryptosporidium parvum*

Running title: Holocentricity in *Cryptosporidium*

Abigail Kimball[1][#], Wanyi Huang[1][#], Rui Xu[1,a], Melanie Key[1], Lisa Funkhouser-Jones[1,b], and L.

David Sibley[1*]

[#] Contributed equally

Department of Molecular Microbiology, Washington University School of Medicine, St. Louis,

MO 63110, USA

Present addresses:

[a] State Key Laboratory for Animal Disease Control and Prevention, South China Agricultural

University, Guangzhou 510642, China

[b] Department of Microbial Pathogenesis & Immunology, Naresh K. Vashisht School of

Medicine, Texas A&M University, Bryan, TX 77807, USA

*Address correspondence to Dr. L. David Sibley, Department of Molecular Microbiology,

Washington University School of Medicine, 660 S. Euclid, CB 8230 St. Louis MO 63110. E-

mail address: sibley@wustl.edu

Keywords: apicomplexan, mitosis, centromeres, holocentric, *Cryptosporidium*

ABSTRACT (150 limit)

During its asexual cycle, *Cryptosporidium parvum* (*C. parvum*) amplifies through three rounds of nuclear division before undergoing cytokinesis to form infectious merozoites. However, chromosome organization and segregation during nuclear division remain unstudied. Here, we visualized H3-histones, including the centromeric histone H3 (CENH3), telomeres, and centrosomes during mitosis. Nuclear division was accompanied by centrosome duplication and elongated microtubules that spanned the dividing nuclei. Surprisingly, *C. parvum* centromeres detected by CENH3 showed diffuse staining throughout nuclear division, that overlapped with telomeres in the apical nucleus. Analysis of centromeres based on CENH3 capture and DNA sequencing revealed numerous distinct binding sites scattered across all eight chromosomes, including both coding regions and intergenic regions, that were typified by GA rich repeat sequences. The unique holocentric chromosome structure of *C. parvum* is unlike the single regional centromeres found in related apicomplexans, suggesting it arose independently from other known examples in plants, insects and worms.

INTRODUCTION

Cryptosporidium parvum (C. parvum), an obligate intracellular parasite, remains a serious global health and agricultural threat [1-4]. Transmission occurs through the ingestion of infectious oocysts and causes severe diarrheal illness in infants, young animals, and immunocompromised adults [5-7]. Once ingested, sporozoites exit the oocyst, invade host cells, and undergo three rounds of nuclear division within a single shared cytoplasm (the process of merogony), followed by cytokinesis to produce eight identical merozoites. Newly formed merozoites are released into the lumen, where they invade new host cells, and repeat the

invasion-egress cycle three times before transitioning to the sexual phase of the life cycle [8, 9].

Despite its significance as a global health problem, and the novelty of apicomplexan nuclear

division processes, *C. parvum* mitosis remains largely uncharacterized, with our current

understanding limited to basic microscopy studies [8, 10-15].

To divide rapidly, apicomplexans appear to rely on several features that are thought to

streamline nuclear division. Comparative cell biology suggests that maintaining chromosomes in

a minimally condensed state may reduce the need for extensive chromatin remodeling during

division, although the extent to which this directly accelerates mitosis remains to be fully

resolved [16, 17]. This adaptation necessitates strict nuclear organization during mitosis, where

centromeres are typically maintained on the apical end of the nucleus attached to the

centrosomes, and telomeres persistently cluster at the basal end of the nuclear periphery [18-20].

Additionally, apicomplexans maintain their nuclear envelope throughout mitosis, minimizing the

energy and time required for breakdown and reformation [17, 21]. A consequence of this closed

mitosis is restricted access of spindle fibers to chromosomes, necessitating alternative

segregation mechanisms based on intranuclear spindles. Rather than centrosomes orchestrating

microtubule organization from the cytoplasm, apicomplexan centrosomes (spindle pole bodies,

microtubule-organizing centers) are embedded within the nuclear envelope and serve as spindle

assembly hubs that elongate the microtubules so they can attach to the centromeres and

ultimately coordinate chromosome movement during division [22-29]. It has been postulated that

nuclear pores are essential for nucleocytoplasmic transport through the intact envelope and

directly support the process of spindle assembly [30, 31]. Electron microscopy studies suggest

that *C. parvum* undergoes schizogony, with multiple rounds of nuclear replication preceding a

cytoplasmic division, followed by the egress of daughter parasites from the host cell [13].

During mitosis and meiosis, centromeres function to orchestrate chromosome segregation

by serving as sites for microtubule anchoring to assure faithful separation at metaphase [32].

Centromeres have variable architecture in different organisms consisting of point centromeres

(small 250 bp single site per chromosome) in yeast [33], regional centromeres (gene poor islands

of heterochromatin ranging from several to hundreds of kb) in *T. gondii* and *P. falciparum* [26,

27, 34], and holocentric chromosomes (multiple centromere sites along the chromosome) found

in multiple species of plants, insects and worms [35-37]. Although centromere function is

essential and highly conserved across eukaryotes, centromeric DNA sequences and size are

subject to rapid evolution [38-40]. Variation in centromere architecture affects nuclear

organization, karyotype changes, and ultimately drives speciation [41, 42].

The nuclear structures, chromosome features, and mitotic mechanisms, including the role of

centromeres, have not been examined previously in *C. parvum*. Here, we delineate the

fundamental biology of *C. parvum* mitosis, revealing significant deviations from previously

assumed mechanisms. Using high-resolution imaging we examined nuclear division during the

first round of merogony, leveraging multiple transgenic parasite strains and mitosis-specific

antibodies. Additionally, we used Cleavage Under Targets and Nuclease Release (CUT&RUN)

[43] to identify the location of the centromeres based on CENH3 and patterns of centromere-

associated histone modifications in *C. parvum* and the related parasite *Toxoplasma gondii* (*T.*

gondii). In contrast to the monocentric centromeric structure of *T. gondii*, we found that *C.*

parvum has a dispersed centromeric chromosome structure with CENH3 binding sites scattered

across multiple small regions on all 8 chromosomes, resulting in diffuse CENH3 centromere

staining throughout mitosis. Overall, these results suggest that *C. parvum* employs a divergent

mitotic mechanism compared to other well-studied apicomplexans and has convergently evolved

holocentric architecture.

RESULTS

Characterization of centromeric histone H3 (CENH3) and histone H3 candidates in C.

parvum

The genome annotation of *C. parvum* contains two genes encoding Histone H3 *cgd3_2540* and

cgd4_3220 (referred to here as H3.1 and H3.2, respectively) and a third gene *cgd4_2030*

annotated as a centromere-specific Histone H3 variant (CENH3). To validate these annotations,

we performed multiple BLASTP analyses using the protein products of each of these three *C.*

parvum genes to identify top hits in the following groups: 1) Apicomplexa (excluding all

Cryptosporidium species), 2) all eukaryotic organisms (excluding all Apicomplexa), 3) *Homo*

sapiens, and 4) *Saccharomyces cerevisiae*. Collectively, these analyses provide strong support

for cgd4_2030 as the CENH3 ortholog in *C. parvum* (**Table 1**). To examine the relationship

between the protein sequences for the *C. parvum* genes of interest and the top hits of BLASTP

analyses, a phylogenetic tree was generated (**Fig 1A**). We found that *cgd3_2540* (H3.1) and

cgd4_3220 (H3.2) closely cluster with histone H3 sequences from other organisms, whereas

cgd4_2030 (CENH3) clustered with other centromeric histone H3 proteins. We then directly

compared the amino acid sequences of H3.1 and H3.2 with a global sequence alignment and

found that the two sequences were highly similar (99.26%), but less similar compared to the

CENH3 (65.10% and 64.43%, respectively) (**Table 2 and Fig 1B**).

Table 1: The results of BLASTP analyses using the *C. parvum* centromeric histone H3-like protein (*cgd4_2030*), histone H3-like

proteins (*cgd3_2540* and *cgd4_3220).*

Query	Top hits in Apicomplexa*	E value	Top hits in all Eukaryotes#	E value	Top hits in *Saccharomyces cerevisiae*	E value	Top hits in *Homo sapiens*	E value
cgd4_2030 (CENH3)	histone H3-like centromeric protein CSE4 [*C. cionae*]	5e-46	centromeric DNA-binding histone H3-like protein cse4 [*N. ribis*]	3e-39	Chain A, Cse4	2e-30	Chain A, Histone H3.1, Histone H3-like centromeric protein A, Histone H3.1	5e-39
cgd3_2540 (H3.1)	histone H3 [*C. cionae*]	2e-86	histone H3-like [*P. hypophthalmus*]	3e-86	histone H3	4e-81	histone H3.2	1e-88
cgd4_3220 (H3.2)	histone H3 [*C. cionae*]	2e-90	histone H3-like [*P. hypophthalmus*]	2e-87	histone H3	2e-81	histone H3.2	1e-89

118   *parvum* proteins were compared against all organisms in the Apicomplexa phylum (except for *Cryptosporidium* spp.),

119   # *C. parvum* proteins were compared against all organisms (except for Apicomplexa)

122   **Table 2:** The identity and similarity results from multiple global sequence alignments between

123   H3.1, H3.2, and CENH3 proteins using a BLOSUM62 scoring matrix.

Cp H3.1 and *Cp* H3.2	*Cp* CENH3 and *Cp* H3.1	*Cp* CENH3 and *Cp* H3.2
	Identity: 46.31% Similarity: 65.10% Gaps: 14.09%	
Identity: 97.78% Similarity: 99.26% Gaps: 0.00%		Identity: 46.31% Similarity: 64.43% Gaps: 14.09%

To further explore the distribution of H3.1 and H3.2 H3 variants, we generated two

transgenic lines of parasites that replaced the dispensable thymidine kinase (*tk*) gene with a

second copy of *cgd3_2540* (referred to as H3.1-3HA) or *cgd4_3220* (referred to as H3.2-3HA)

fused with a triple hemagglutinin (3HA) epitope tag using CRISPR/Cas9 (**Fig 1C, S1 A-B Fig,**

Table S1). Transgenic parasites were used to infect a human adenocarcinoma cell line (HCT-8)

monolayer, cultures were fixed at 18 hours post infection (hpi) during merogony, stained, and the

localization of the 3HA tag was observed by immunofluorescence (IFA) high-resolution

confocal laser scanning microscopy equipped with Airyscan detection (LSCM-A). We found that

H3.1-3HA (**Fig 1D**) and H3.2-3HA (**Fig 1E**) had a diffuse staining pattern, which was expected

based on histone H3 staining in other apicomplexan parasites [27, 44, 45]

The staining pattern of C. parvum centromeric histone H3 (CENH3) is distinct.

To visualize *C. parvum* centromeres, we used CRISPR-Cas9 gene editing to add a 3HA tag to

the C-terminus of the centromere-specific histone H3 variant protein (referred to as CENH3-

3HA) (**Fig 2A, S3 A-B Fig, Table S1**). After successfully selecting and amplifying these

transgenic parasites in immunocompromised mice, we infected HCT-8 cells for 22 h and

examined the staining pattern by LSCM-A microscopy. For these studies, we chose mature

meronts that were oriented with the basal side of the vacuole at the surface of the host cell where

it attaches at the feeder organelle and where the majority of meronts were projecting upward, as

ascertained by fluorescence staining and Z-stack images running perpendicular from bottom to

top of the parasite (see cartoon on **Fig 3C**). The CENH3 staining pattern observed in *C. parvum*

was unexpectedly highly diffuse and occupied the apical half of the nucleus (**Fig 2B-C**) in

contrast to the more dispersed pattern found in the H3.1-3HA and H3.2-3HA tagged parasites

(**Fig 1 D,E**). The diffuse pattern of CENH3 staining in *C. parvum* contrasts sharply with the

focal pattern of regional centromeres previously described in *T. gondii* [27, 46]. Hence, we

utilized a CENH3-3HA-tagged *T. gondii*, gifted by Dr. Kami Kim, as a staining control for our

experiments. These parasites were used for the infection of HCT-8 cells, fixed at 24 hpi, stained,

and examined by high-resolution LSCM-A immunofluorescence (IFA) microscopy (**Fig 2D-E**).

We found that the staining pattern of the centromeric histone H3 in *T. gondii* is punctate, similar

to what has been found previously in *T. gondii* by other investigators [27, 46].

Distribution of centrosomes in C. parvum.

A vital step in mitosis is centrosome duplication, and previous studies in apicomplexans have

shown that it can be traced using antibodies that recognize the centrin 1 protein [27, 47]. A single

point adjacent to the nucleus occurs in non-replicating cells, while two centrin 1 positive points

per nucleus indicates the parasite is undergoing replication [27, 47]. In *Plasmodium*, during

schizogony, asynchronous duplications of centriolar plaques, centrosomes, have been observed

across multiple nuclei within a single cytoplasm, in this study we sought to identify analogous

centrosomes in *C. parvum*. [28]. To identify parasites undergoing early mitosis, we infected

HCT-8 cells with excysted wild type *C. parvum* sporozoites, washed off uninvaded sporozoites

at 2 h, fixed the cultures at various timepoints between 6.5-8.5 hpi, and stained with Hoechst and

an anti-centrin-1 antibody. Initially, trophozoites contained a single centrin-1 focus associated

with the single nucleus (**S4A**). However, in other examples of trophozoites, we often observed

that each nucleus consistently exhibited two centrin-1 positive points (**Fig 2F**). Furthermore, by

visualizing the number of nuclei based on Hoechst staining, and centrin-1 to monitor dividing

centrosomes, we observed that all nuclei found in a common cytoplasm shared the same centrin-

1 staining pattern. For example, when 2N or 4N meronts were observed to have duplicated the

centriole as shown by centrin-1 staining, the duplicated pattern was consistently observed for all

nuclei within that cell (**Fig 2F**). To determine if the diffuse centromere staining pattern persisted

during active mitosis in *C. parvum*, we infected cultures with CENH3-3HA parasites and

repeated the experiment previously described. We found that nuclei undergoing mitosis also

displayed a diffuse CENH3-3HA staining pattern by LSCM-A (**Fig 2G**). We also attempted to

localize CENH3 using ultrastructure expansion or immuno-EM staining methods. Despite

extensive troubleshooting, we could not identify positive signals in N- or C-terminally tagged

CENH3-3HA lines, or in separate line expressing a spaghetti monster (sm)HA tag (i.e. CENH3-

smHA), by either method, which may be due to the low abundance of CENH3 or difficulty

preserving the epitope during these procedures.

The telomere (G-strand binding protein, GBP) staining pattern is apical in C. parvum.

Previous studies in *Plasmodium* have found that the telomeres are consistently found on the

opposite pole to the centromeres [48-50]. Since the centromeres were located toward the apical

end of the nuclei in meronts, we expected that *C. parvum* telomeres would be more basal. To

examine the distribution of telomere in *C. parvum*, we utilized a surrogate protein marker. G-

quadruplex structures readily form at repetitive telomeric sequences and require G-quadruplex

binding proteins (GBPs) for stabilization (**Fig 3A**). A previous study identified *cgd1_3530* as the

GBP in *C. parvum* and determined the specific sequence necessary for *Cp*GBP binding

(GGTTA) [51]. Using AlphaFold and ChimeraX, we modeled the G-quadruplex structure

formed by this repetitive sequence and visualized how the *Cp*GBP protein interacts with and

stabilizes it. (**Fig 3B**). To identify the location of *C. parvum* telomeres relative to the

centromeres, we engineered transgenic parasites to express a 3HA tag at the gene encoding the

C-terminus of the G-quadruplex binding protein (referred to as GBP-3HA) (**Fig 3C, S3 A-B Fig,**

Table S1). LSCM-A microscopy on infected HCT-8 monolayers (24 hpi) revealed that the

telomeres were located toward the apical side of the nucleus (**Fig 3D**), rather than at the opposite

pole as expected.

The telomere (GBP) and centromere (CENH3) staining patterns overlap in C. parvum.

To confirm that *C. parvum* centromeres and telomeres had overlapping nuclear locations, we

generated an additional line of transgenic parasites that replaced the dispensable thymidine

kinase (*tk*) gene with a second copy of an N-terminally 3HA tagged CENH3 and GBP C-

terminally tagged with 3Ty via CRISPR/Cas9, with an intervening P2A split peptide to facilitate

co-expression of separate proteins (referred to as 3HA-CENH3-GBP-Ty) (**Fig 3E, S1C-D Fig,**

Table S1). HCT-8 cultures were infected with 3HA-CENH3-GBP-Ty parasites, fixed at 6.5 hpi,

and examined by LSCM-A microscopy after staining. We observed that 3HA-CENH3 and

GBP-3Ty signals were both more concentrated in the apical region of the nucleus, as defined by

Hoechst staining. This overlap was evident by fluorescence microscopy and further supported

quantitatively using the Coloc2 plugin in FIJI/ImageJ, which yielded a Pearson correlation

coefficient of 0.96 (**Fig 3F, S2 Fig**).

Detection of mitotically active DNA, microtubules, and the nuclear pore proteins in C. parvum

The phosphorylation of histone H3 (specifically at serine 10 and serine 28) is a highly conserved

mechanism for initiating chromosome condensation in preparation for mitosis [52]. Antibodies

recognizing the phosphorylated version of H3 at serine 10 are commonly used in other systems

to determine which cells are currently undergoing mitosis from late interphase through telophase

[53]. Even though apicomplexan parasites are known not to condense their chromosomes, *C.*

parvum meronts frequently stained positive for the phosphorylated histone H3 (PH3) antibody

(**Fig 4A**), suggesting they are in the process of mitosis. Previous studies have characterized the

distribution of nuclear pores in *Plasmodium falciparum* using a rabbit polyclonal antibody to the

protein NUP116 [54]. We rationalized that the epitope recognized by this antibody might also be

conserved in *C. parvum*, despite their evolutionary distance. We visualized nuclear pores in *C.*

parvum using the rabbit anti- NUP116 antibody and found that it reliably stained the parasite

nucleus in a punctate pattern that was concentrated primarily peripherally in the nucleus (**Fig**

4B). Although we have not verified the identity of this cross-reaction, it presumably represents a

component of nuclear pores that will be used for future studies on nuclear organization in *C.*

parvum.

Among apicomplexan parasites, mitosis has primarily been studied in *T. gondii* and *P.*

falciparum [16, 17, 27, 55]. These prior studies demonstrate that once centrosomes duplicate,

and the nuclear envelope forms a tunnel-like structure, organizing spindle microtubules within it,

spindle microtubules then attach at the centromere and mediate chromosome movement [56]. We

also attempted to visualize this process in *C. parvum* by staining with antibodies that recognize

the centrin-1 nd alpha tubulin (**Fig 4C-D, S4B-C**). We examined cells at 6-7 hpi to capture early

stages of mitosis. We frequently observed two conformations of centrin-1 tubulin structures that

may represent early (**Fig 4C, S4B**) and late mitotic phases (**Fig 4D, S4C**). In Fig 4C, the

duplicated centrin signal shows a single discrete spindle (microtubule signal) overlying

what appears to be a single nucleus, as evident by size and shape. In other examples the

centrin spots have moved further away, and the spindles elongated (Y-shaped) (**Fig 4D,**

S4C). Notably, the nucleus in this image is much larger and diffuse, consistent with it

being at a later stage that is starting to divide.

Composition of centromeres in C. parvum and T. gondii

Centromeres are defined by the deposition of CENH3-containing nucleosomes [27, 57]. Given

the diffuse staining of CENH3, we wanted to further characterize the centromere by identifying

DNA sites bound by this protein. Therefore, we adopted protocols for CUT&RUN to identify

centromeres based on CENH3 binding based on capture with anti-HA antibody in *C. parvum*

[43]. To ensure sufficient HA binding, we developed a CENH3 line tagged with spaghetti

monster (referred to as CENH3-smHA) that contains 10 HA epitope repeats for these

experiments and validated identical staining patterns to CENH3-3HA *C. parvum* (**S3 A-C Fig,**

Table S1). To validate our experimental strategy, we first applied CUT&RUN to *T. gondii*,

whose centromeres have previously been defined by ChIP-seq [27]. Our CUT&RUN data

closely matched earlier findings, with CENH3 enrichment observed at discrete chromosomal loci

corresponding to the previously described centromeres (**Fig 5A, S5A-C Fig**). We also detected

the positive correlation of CENH3 with H3K9me3 and inverse correlation with H3K4me3

reported in earlier ChIP-chip analyses (**Fig 5A, S5A-B Fig**). Based on these signals, we

implemented a stepwise strategy to identify centromeric regions. Valleys in H3K4me3,

consisting of 20 kb windows, sized to match reported centromere lengths (16 kb ± 3.4 kb)[27],

combined with co-enrichment of H3K9me3 with CENH3-smHA were used as criteria for

identifying centromere regions (**S5D Fig**). Applying these rules, we identified a single

centromere in each of the 13 chromosomes of *T. gondii* (**S5E Fig**), with Chr VIIb identified as

part of Chr VIII [58](**S5C Fig**).

We next applied CUT&RUN to *C. parvum*. Initial experiments using a 120-minute MNase

digestion failed to generate usable data, as both sequencing depth and genome coverage were

low compared to negative controls (**S1 Dataset**). Since CENH3 nucleosomes can be

hypersensitive to MNase digestion, as reported previously in *Caenorhabditis elegans* (*C.*

elegans) [57], we tested digestion times of 5, 30, and 60 min. The 30 min digestion yielded the

best coverage and sequencing depth, and these conditions were used for following analyses (**S1**

Dataset, S6A Fig). Notably, sequencing depth of the *C. parvum* samples was more variable and,

in some cases, lower than that of *T. gondii*. This difference primarily reflects technical

challenges associated with generating transgenic *C. parvum* parasites. To ensure data quality and

comparability, we included only samples with high genome coverage (>80%) in downstream

analyses. Unlike *T. gondii*, read mapping profiles for CENH3 in *C. parvum* showed signals

broadly distributed along chromosomes (**Fig 5B**). To locate centromeric regions in *C. parvum*,

we first attempted the *T. gondii* stepwise strategy, but no valid windows were detected. As the

CENH3 distribution resembled that of the holocentric nematode *C. elegans*, we adjusted

thresholds accordingly by restricting the peaks analysis to 400 bp windows (**S6B Fig**). We

identified 423 CENH3-enriched regions distributed across all chromosomes. Heatmaps and

aggregate signal profiles showed that these windows displayed features of CENH3 enrichment

correlated with H3K9me3 deposition and H3K4me3 depletion, suggesting that they represent

centromere-associated chromatin (**S6C-D Fig**). Importantly, these signals were largely absent in

wild type parasites stained with HA antibody as a control, supporting their authenticity (**S6C-D**

Fig). To further refine localization, we excluded windows overlapping with WT-HA signals,

reducing the dataset to 347 high-confidence centromeric windows (**S6B,E Fig**). Taken together,

these results indicate that *C. parvum* possesses many small CENH3 binding sites dispersed

across eight chromosomes, in contrast to the monocentric nature of *T. gondii*.

We further investigated characteristics of *C. parvum* centromeres. In contrast to *T. gondii*,

where centromeres are typically located within large intergenic regions [27], no extended

intergenic regions were observed in *C. parvum* due to the tight compaction of the genome.

Surprisingly, the majority of *C. parvum* CENH3-associated windows were located within gene

coding regions rather than being restricted to intergenic regions (**S2 DataSet2**). To evaluate

evolutionary constraints on CENH3-associated regions, we performed nucleotide diversity (Pi)

analyses on the DNA sequences within these regions and dN/dS analyses on the genes

overlapping these windows. We compared previously generated genomic sequences from

numerous *C. parvum* IIa and IId isolates, which show high genomic similarity. Most CENH3-

associated windows (~82%) showed reduced Pi values compared with genome-wide 400-bp

sliding windows, and only a single gene within the windows exhibited an elevated dN/dS ratio

relative to the genome-wide orthologous gene set (**S6F,G Fig**). These results indicate that

CENH3-associated regions are highly conserved and under strong purifying selection. To

determine whether centromeric regions were also present within intergenic regions of *C. parvum*,

we applied a stepwise identification strategy using a slightly relaxed read depth threshold (**S6F**

Fig). We identified 58 centromeric regions distributed within intergenic regions across all

chromosomes (**Fig. 5C, S3 DataSet**). Heatmaps and aggregate signal profiles for CNEH3 as well

as chromatin marks also supported that they represent centromere-associated chromatin (**Fig.**

5D,E). These results suggest that CENH3 binding in *C. parvum* is broadly distributed throughout

the chromosomes, including both with in coding regions and intergenic regions. To assess

sequence properties of *C. parvum* centromeric windows, motif analysis of *C. parvum*

centromeric windows was performed using MEME, revealing two GA-rich motifs enriched at

CENH3-associated sites (**Fig 5F**). Notable, these GA repeats are also found in other regions of

the genome that were not identified in the CENH3 capture studies.

DISCUSSION

Despite rapid asexual proliferation being required for parasite expansion, the molecular

mechanisms that underlie *C. parvum* mitosis have remained unexplored. To investigate this

process, we examined nuclear division using markers to track multiple rounds of mitosis during

merogony. Nuclear division in *C. parvum* can be visualized by staining with phospho-H3 and

centrin-1, which provide markers for mitotically active chromatin and duplicating centrosomes,

respectively. Surprisingly, both CENH3, a marker for centromeres, and GBP, a marker for

telomeres, were diffusely localized across the apical half of the nucleus. Moreover, these diffuse

apical patterns persisted throughout mitosis, as shown by visualizing nuclear division stages with

centrin-1 and microtubules. To explore the molecular nature of centromeres, we performed

CUT&RUN sequencing of transgenic parasites expressing epitope-tagged CENH3. Although

discrete monocentric peaks were detected in the related parasite *T. gondii*, CENH3 binding sites

were diffusely scattered across ~ >400 sites in the *C. parvum* genome. Overall, these findings

indicate that *C. parvum* has evolved an alternative chromosome structure compared to other

apicomplexan parasites, differences that may have significant functional consequences for

chromosome stability and meiotic recombination.

Following entry of sporozoites (or merozoites) into the host cell, the parasite undergoes

merogony, consisting of 3 rounds of nuclear division prior to cytokinesis to generate infectious

merozoites. By visualizing the number of nuclei based on Hoechst staining, and centrin-1 to

monitor dividing centrosomes, we observed that all nuclei found in a common cytoplasm shared

the same centrin-1 staining pattern (single centrin-1 spot when not dividing and duplicated

centrin-1 spot prior to nuclear division). This pattern suggests that nuclei within a single meront

may divide synchronously, although further studies using synchronized cells and live cell

imaging would be needed to substantiate this hypothesis. This pattern differs from *Plasmodium*

schizogony, where nuclei are observed at different mitotic stages within a single schizont [28]. In

Plasmodium, nuclear pores are important for division as they facilitate the transport of mitotic

molecules through the intact nuclear envelope and associate with the centriolar plaques [31, 59].

Although we did not examine the dynamics of nuclear pores during division, they appear as

peripheral puncta in mature meronts and may serve a similar function in *C. parvum*.

Staining with alpha-tubulin allowed us to visualize *C. parvum* microtubules during mitosis,

particularly at early stages where microtubules form a cluster that may represents the spindle that

will drive chromosome separation during mitosis. The pattern of microtubules described here in

dividing cels is quite different from the stable, elongate microtubules that emanate from the

apical pole of sporozoites and merozoites [60]. Such dramatic re-organization of microtubules

between non-dividing and dividing cells is not unusual for eukaryotic cells, although the

dynamics and regulation of this process have not been examined extensively in *Cryptosporidium*.

Based on qualitative observations of centrosome duplication, centrin-1 positioning, microtubule

organization, and changes in nuclear morphology, our studies suggest an approximate sequence

of events corresponding to sequential rounds of mitosis during early, mid, and late merogony.

Further studies using fluorescently tagged proteins and life cell imaging, combined with

synchronous cell division would be important to validate and extend this model. Regardless of

the exact dynamics of mitosis, these studies provide an important foundation for studies of the

dynamics of centromere staining described below.

Unexpectedly, we found a diffuse nuclear CENH3-3HA staining pattern rather than the

punctate pattern observed in other apicomplexan parasites [27, 29, 34]. We also confirmed that

the diffuse CENH3 staining pattern did not deviate when the method of tagging was altered (e.g.

smHA), when the location of the tag was swapped (N-terminally tagged, e.g. 3HA-CENH3-

GBP-3Ty), or when a tagged second copy of the gene was inserted into a neutral locus of the

genome (3HA-CENH3-GBP-3Ty). We also developed epitope-tagged lines of telomere-binding

protein GBP, similar to studies in *Eimeria* and *Plasmodium* [61, 62]. Unlike in other

apicomplexans, GBP was also localized to the apical half of the nucleus in *C. parvum*,

overlapping closely with CENH3 [18-20, 48]. We further confirmed that the telomeres and

centromeres occupied a similar nuclear location using a parasite strain in which CENH3 and

GBP proteins were simultaneously expressed as tagged second copies. This pattern contrasts

sharply with other apicomplexans where telomers are localized to the opposite pole of the

nucleus from centromeres [48-50]. We sought to determine if centromeres remained diffuse

throughout mitosis by simultaneously visualizing the centrosomes and centromeres by staining

CENH3-3HA with anti-HA and centrin-1 antibodies. We confirmed that in *C. parvum,* the

diffuse centromere staining pattern does not change throughout mitosis, which further contrasts

with the highly focal nature of centromeres in *T. gondii* and *P. falciparum* [26, 27].

Given the highly unusual staining patterns of CENH3 in *C. parvum*, we decided to

investigate CENH3 binding sites on DNA using CUT&RUN sequencing. For comparison, we

also performed CUT&RUN studies on CENH3 in the related parasite *T. gondii* and compared

them to previous studies that used ChIPSeq to define centromeres [27]. In *T. gondii*, CUT&RUN

sequencing identified highly similar CENH3 binding sites to those reported previously by

ChIPSeq [27] with a single predominant peak on each chromosome, including newly defined

peaks on chromosome IV, that had not been identified previously. Centromeres in *T. gondii*

consist of gene-poor regions of ~15 kb that are not defined by any particular sequence or repeats

but instead are positively correlated with H3K9me3 and inversely correlated with H3K4me3,

typical chromatin patterns expected for centromeric regions [18, 27]. Under similar conditions,

CUT&RUN experiments for CENH3 binding sites in *C. parvum* failed to identify similar

singular dominant peaks on each chromosome. Instead, CENH3 binding sites were

hypersensitive to MNase digestion, with the signal being the highest at a 30- min incubation

time, similar to that observed in ChIP-seq studies of the holocentric nematode *Caenorhabditis*

elegans (*C. elegans*) [57]. Under optimal digestive conditions, we found that the CENH3 signal

in *C. parvum* was broadly distributed along chromosomes, with ~ 350 distinct sites in coding

regions and another ~ 60 sites within intergenic regions scattered across 8 chromosomes.

Interestingly, like *T. gondii*, *C. parvum* CENH3 peaks also correlated with H3K9me3 signal and

were found in areas lacking H3K4me3, consistent with a centromere-associated chromatin

pattern. The distribution of CENH3 binding sites in *C. parvum* is similar to that described in *C.*

elegans as polycentric, implying that CENH3 binds to distinct sites rather than diffusely across

the entire chromosome [57].

Collectively, our findings are consistent with *C. parvum* containing holocentric centromeres

based on the following criteria: Firstly, the staining pattern of CENH3 diffusely stains the apical

half of the nucleus under nondividing conditions, in contrast to the focal staining of CENH3 in *T.*

gondii [18, 27] and *P. falciparum* [26]. Secondly, the diffuse pattern of CENH3 staining remains

throughout mitosis, suggesting centromeres do not coalesce into a single regional center. Thirdly,

CENH3 binding sites as defined by CUT&RUN resemble the dispersed pattern of low-intensity

peaks seen in *C. elegans*, which has well-characterized holocentric chromosomes. Although we

interpret this diffuse distribution as consistent with holocentric centromeres, we acknowledge

that an alternative explanation, such as asynchronous chromatid separation or cohesion fatigue,

could also contribute to the observed pattern. However, current imaging resolution does not

allow simultaneous visualization of all microtubules and all CENH3-associated kinetochore

sites, so we cannot fully distinguish between these possibilities. Therefore, we present our

interpretation cautiously while noting that the number of predicted CENH3 loci substantially

exceeds the 8 (1N) regional centromeres expected for a monocentric organization. Finally, the

CENH3 binding sites show evidence of simple GA nucleotide repeats, a feature also seen in *C.*

*elegan*s [57]. In both cases, the GA repeats are also found in other regions of the respective

genomes, so this feature alone is not sufficient to mediate CENH3 binding. One unusual feature

of the CENH3 binding sites in *C. parvum* is that they extend into coding regions, which is not an

expected feature of centromeres. Interestingly, these regions were under strong purifying

selection at the intraspecies level. The dispersed polycentric pattern of CENH3 binding sites in

C. elegans is also unusual in that these regions are concentrated in High Occupancy Targets

(HOT) sites for permissive binding of transcription factors [57]. In *C. elegans*, the CENH3 and

HOT sites appear to serve a dual purpose: in replicating cells they bind CENH3, while they

recruit transcription factors when CENH3 is lost during exit from the cell cycle in differentiated

muscle cells [57]. By analogy, transcriptional silencing during mitosis in *C. parvum* may allow

CENH3 binding in coding regions to facilitate chromatid segregation, with displacement during

the rest of the cell cycle when transcription is active. Alternatively, the CENH3 binding sites in

intergenic regions may function as the primary centromere-localizing signal for formation of

kinetochores.

Holocentric chromosomes have been documented in many species, primarily through direct

cytological observation, yet their functional implications and evolutionary origins remain poorly

understood [37, 57, 63, 64]. Recent comparative and modeling work in fungi shows that

complete transitions between centromere 'types' (point/regional/holocentric) occur progressively

and do not fix by drift alone; rather, selection is needed to drive and stabilize new architectures

while maintaining compatibility with the relatively conserved kinetochore–spindle interface [65].

In this light, *the* many small CENH3-binding sites in *C. parvum* fits a model of progressive

centromere remodeling along an evolutionary continuum, rather than representing an abrupt

departure from other apicomplexan centromere types.

The organization we describe has clear functional implications, as the presence of hundreds

of small CENH3-positive sites suggests that *C. parvum* chromosomes may rely on numerous

'micro-kinetochores' distributed along their length. By spreading spindle attachment across

many small units rather than concentrating mechanical force at a single centromere, the parasite

could effectively diffuse the load generated during chromosome segregation. Such a system

would be particularly advantageous for *C. parvum*, which performs closed mitosis with

uncondensed chromatin and must execute multiple rounds of synchronous division during rapid

merogony. In this framework, the diffuse, apically clustered distribution of CENH3 is not simply

a structural anomaly, but an evolved solution to the mechanical and cell-cycle constraints

imposed by rapid, multi-round division in an intestinal environment. Thus, the centromere

architecture of *C. parvum* likely reflects selective pressures associated with maintaining

chromosome stability, rather than a simple deviation from typical apicomplexan chromosome

biology.

Current evidence suggests that holocentric chromosomes have evolved convergently at least

13 times across distantly related taxa in both the plant and animal kingdoms [36, 37, 64].

Holocentricity has not previously been described in protists, and hence it likely evolved

convergently in *C. parvum*. However, we note that the directionality of centromere evolution

within Apicomplexa remains unresolved. Because centromere organization has not been

characterized in closely related outgroups such as gregarines or chromerids, it is equally

plausible that the common ancestor of Apicomplexa possessed holocentric chromosomes and

that monocentric organization arose secondarily in coccidia and hematozoa. Thus, the

holocentric architecture in *C. parvum* may represent either an independently derived state or the

retention of an ancestral condition. It is also possible that holocentricity is more widespread

among protists, but that it has been overlooked due to their small size and failure to condense

chromosomes during division.

One proposed advantage of holocentricity is its capacity to stabilize chromosomal fragments

during mitosis, thereby mitigating genetic loss. This trait may confer a selective benefit to

organisms frequently exposed to elevated levels of DNA damage due to environmental stressors

or lifestyle factors [36, 63]. Thus, holocentricity may impart genome stability under stress

encountered in the gastrointestinal system where the parasite is exposed to toxic bacterial

metabolites [66] and host immune responses [67]. Despite this potential advantage,

holocentricity also presents a major challenge for coordinating chromatid segregation following

crossover during meiosis I. Well-characterized holocentric species have evolved mechanisms to

circumvent this issue, such as inverted meiosis or, in the case of *C. elegans*, the deployment of

kinetochore proteins that localize to discrete chromosomal regions, effectively mimicking

monocentric behavior [68]. Whether *C. parvum* uses a similar mechanism as *C. elegans* to

localize kinetochore proteins to discrete sites on the chromosome is uncertain, but previous work

from our laboratory examining crossover rates during *C. parvum* meiosis revealed a high

frequency of crossover events and failed to find evidence for inverted meiosis, a process used by

some holocentric organisms [69]. In contrast, interspecies crosses between *C. tyzzeri* and *C.*

parvum revealed that crossovers are strongly suppressed and most chromosomes are inherited

uniparentally without crossovers [70]. Although we have not interrogated the nature of CENH3

binding sites in *C. tyzzeri*, differences in the recognition sequences or pattern of centromere

distribution (i.e. monocentric vs. holocentric) could result in suppression of hybrids due to miotic

or meiotic interference.

Limitations and Future Studies

Examination of the nuclear organization of *C. parvum* throughout the cell cycle revealed a

persistently diffuse centromere staining pattern, suggesting a holocentric chromosome structure.

During division, such multiple centromeric sites dispersed along the chromosome are expected to

bind centrosomes and thereby connect to microtubules for chromosome separation. Although we

visualized events consistent with this process in an artificial timeline, future studies using

fluorescent reporters could potentially provide more precise information on the formation of

centrosomes, coordinated assembly of microtubules, and nuclear division in *C. parvum*.

However, obtaining direct cytological data in support of holocentric centromeres and their

behavior during nuclear division will be challenging in this system because chromosomes do not

condense during division. Alternatively, it may be possible to explore the nature of centromeres

by examining the distribution of other conserved centrosomes and kinetochore binding proteins

during nuclear division. Such studies could shed further light on chromosome tethering in

nuclear division and provide invaluable insights into unique evolutionary adaptations in *C.*

parvum. Additionally, extending these studies to define how centromeres are organized in related

species could provide insight into mechanisms of diversification that drive speciation.

Acknowledgements:

We thank Jennifer Barks for assistance with cell culture, Dr. Wandy L. Beatty, Microbiology

Imaging Facility, Washington University School of Medicine in St. Louis, for assistance with

microscopy, and Dr. Kami Kim, Univ. South Florida, for providing the *T. gondii* CENH3-HA

line used here. We are grateful for helpful comments from Drs. Daniel E Goldberg, Douglas L.

Chalker, and Tim Schedl, Washington University and Dr. Lihua Xiao, South China Agricultural

University. Genome sequencing was performed by the GTAC Genome core at Washington

University in St Louis. Supported in part by the Graduate Research Fellowship Program,

National Science Foundation (to A.K.) and grant AI175150 from the National Institutes of

Health (to L.D.S.).

Author contributions

Conceptualization, AK, LDS; Data curation, AK, WH; Formal analysis, AK, WH; Funding

acquisition, LDS; Investigation, AK, WH, MK, LFJ; Methodology, AK, WH, MK, LFJ;

Resources, WK, WH, MK, LFJ; Validation, AK, WH; Visualization, AK, WH; Supervision,

LDS; Writing – original draft, AK, WH, LDS; Writing – review & editing, AK, WH, LDS.

Data Sources and Availability:

All materials described here are available from the corresponding author based on MTA.

Sequence for the CUT&RUN analyses have been deposited to the NCBI Bioproject #

PRJNA1330822.

Declarations:

The authors declare that they have no conflicts to disclose.

MATERIALS & METHODS

Animal studies and ethical approval.

Animal studies were approved by the Institutional Animal Studies Committee (School of

Medicine, Washington University in St. Louis). *Ifngr1*$^{-/-}$ mice (003288; Jackson Laboratories),

and NOD scid gamma mice (referred to as NSG) (005557; Jackson Laboratories) were bred in-

house in a specific-pathogen-free animal facility on a 12:12 light-dark cycle. Male and female

mice between 8 and 12 weeks of age were used for sex-matched groups used in crossing

experiments. Mice were co-housed with siblings of the same sex throughout the experiments.

Cell culture.

HCT-8 cells (Human ileocecal adenocarcinoma, ATCC CCL-244) derived from a human

ileocecal carcinoma were cultured in RPMI 1640 ATCC Modification medium supplemented

with 10% fetal bovine serum at 37 °C in a 5% CO_2 incubator. After trypsinization, 450,000 cells

were plated on glass coverslips in 24-well plates and used for experiments. Cells were confirmed

to be mycoplasma-free with the e-Myco plus Mycoplasma PCR detection kit.

Immunofluorescence assays & microscopy. Staining protocols have been previously described

and all antibodies are listed in **Table S2** [71]. Briefly, infected HCT-8 cultures were fixed with

4% formaldehyde 8-24 hpi, permeabilized with 0.1% Triton X-100, blocked with 1% bovine

serum albumin (BSA), stained with a primary antibody mixture for 1 h followed by secondary

antibodies for 1 h and Hoechst for 20 min. Primary antibodies were diluted for staining: anti-HA

(rat, 1:500), anti-Aldolase (ALD, rabbit, 1:1000) centrin-1 (mouse, 1:250), anti-Ty (mouse,

1:250), and anti-PH3 (rabbit, 1:500), NUP116 (rabbit, 1:100). All Alexa Fluor secondary

antibodies were diluted at 1:1000 and Hoechst was diluted 1:2000. Coverslips were mounted

onto microscope slides with Prolong Glass Antifade Mountant. Coverslips were visualized using

a Zeiss LSM880 laser scanning confocal microscope equipped with a 63x, 1.4 N.A. Zeiss Plan

Apochromat oil objective lens and Airy scan. Images were acquired with ZEN v2.1, v2.5

software and further analyzed using ImageJ. All staining experiments were performed

independently 2-3 times, and at least 2-5 fields of view were imaged per coverslip.

Expansion Microscopy. Ultrastructure expansion microscopy (U-ExM) was performed as

previously described [72]. HCT-8 cells were infected with the CENH3-3HA or CENH3-smHA

parasites and fixed following the same procedures used for standard immunofluorescence assays

or 16% methanol-free formaldehyde. Fixed samples were embedded overnight in a solution

containing 1% acrylamide and 0.7% formaldehyde to anchor proteins and prevent crosslinking.

The resulting gel was transferred into denaturation buffer and incubated at 95 °C. A second

expansion gel was polymerized on ice using a monomer mixture composed of 19% sodium

acrylate, 10% acrylamide, and 0.1% (1,2-Dihydroxyethylene) bisacrylamide, supplemented with

0.5% ammonium persulfate (APS) and 0.5% TEMED. Polymerized gels were denatured at 95 °C

for 90 min in denaturation buffer (200 mM SDS, 200 mM NaCl, 50 mM Tris-Base, pH 9.0) and

expanded overnight in ultrapure water. Expansion ratios were determined the following day by

measuring gel diameters. Gels were then shrunk in PBS and stained with primary antibodies

(rabbit anti-HA, rat anti-HA, or either rabbit anti-GFP antibodies at 1:200 and rat Pan-Cp,

1:500), followed by secondary antibodies (Alexa Fluor conjugates, 1:500) and Hoechst (1:500)

diluted in freshly prepared 2% PBS/BSA for 3, 12, or 24 h for primary antibodies and 2 h for

secondary antibodies at room temperature. After each staining step, gels were washed three times

in PBS/0.1% Tween for 10 min. Stained gels were re-expanded overnight in ultrapure water

before imaging. Images were acquired using a Zeiss LSM880 laser scanning confocal

microscope equipped with a 63x, 1.4 N.A. Zeiss Plan-Apochromat oil objective and ZEN v2.1 or

v2.5 software.

Immuno-electron Microscopy. Immuno-electron microscopy (EM) was performed as

previously described using CENH3-3HA and CENH3-smHA parasites [73]. HCT-8 monolayers

were grown in 6-well plates and reached confluence before infection with transgenic *C. parvum*

oocysts. After a 20-h incubation, cultures were washed twice in DPBS and fixed for 1 h at 4 °C

in freshly prepared fixative consisting of 4% paraformaldehyde and 0.05% glutaraldehyde in

100 mM PIPES supplemented with 0.5 mM $MgCl_2$ (pH 7.2). Fixed samples were embedded in

10% gelatin and subsequently infiltrated overnight at 4 °C with a cryoprotectant solution of

2.3 M sucrose and 20% polyvinyl pyrrolidone prepared in $PIPES/MgCl_2$. Blocks were trimmed,

rapidly frozen in liquid nitrogen, and sectioned using a Leica Ultracut UCT7

cryo-ultramicrotome. Ultrathin sections (50 nm) were blocked for 30 min in a mixture of 5%

fetal bovine serum and 5% normal goat serum, followed by incubation with rabbit anti-HA, rat

anti-HA, or either of the rabbit anti-GFP antibodies (1:1000) for 1 h at room temperature. After

rinsing in blocking buffer, sections were labeled for 1 h with 18 nm colloidal gold–conjugated

goat anti-rabbit IgG (H+L) diluted 1:20. Labeled sections were then contrasted with 0.3% uranyl

acetate in 2% methyl cellulose and imaged using a JEOL 1200 EX transmission electron

microscope equipped with an AMT 8-megapixel digital camera and AMT Image Capture Engine

v6.0.2. Control sections processed in parallel without primary antibody were included in all

experiments.

CRISPR/Cas9 plasmid construction.

Plasmid construction

Primers used to generate the following plasmids were ordered from IDT and all sequences are

found in **Table S3**. SnapGene (v7.1.1) and NEBuilder Assembly Tool (v2.9.0) were used to

construct plasmid maps and design primers.

The CENH3-3HA tagging plasmid was generated using the previously described ABC-3HA

tagging plasmid [73] by replacing the C-terminal and 3'UTR sequences with equivalent regions

from the *Cp*CENH3 gene amplified from *C. parvum* cDNA to generate the tagging plasmid

CENH3-3HA-Nluc-P2A-Neo^R.

The 3HA-CENH3- GBP-3Ty tagging plasmid was created via a seven-way Gibson assembly

combining a backbone generated from the TK-GFP-Nluc-P2A-Neo^R-TK homology tagging

plasmid previously generated in our lab [74] together with the following fragments: 1) The 5'

homology region of CENH3 (500 bp), the CENH3 CDS, a portion of the 3' homology region of

CENH3 (216 bp) were amplified from the CENH3-3HA tagging plasmid described above. 2)

The 3HA-pLinker region was amplified from a TK-3HA-CENH3-Nluc-P2A-Neo^R-TK plasmid

(data not shown in this study), 3) the GBP CDS was amplified from *C. parvum* genomic DNA,

4) the pLinker-3Ty sequence was amplified from a tagging plasmid previously generated in our

lab [73], and 5) the P2A skip peptide sequence was generated as a gBlock Gene Fragment from

Azenta. The U6:sgTK plasmid previously generated in our lab was utilized for all experiments

with TK locus replacement repair plasmids [74].

The H3.1-3HA and H3.2-3HA tagging plasmids were generated via Gibson assembly using

the 3HA-CENH3-GBP-3Ty plasmid to provide a backbone, the pLinker-3HA region was

amplified from the CENH3-3HA tagging plasmid described above, and the CDS regions for

H3.1 and H3.2 were amplified from *C. parvum* genomic DNA.

The GBP-3HA tagging plasmid was generated using the CENH3-3HA tagging plasmid

described above for the backbone and replacing the C-terminal and 3'UTR sequences with the

GBP CDS sequence amplified from *C. parvum* genomic DNA.

CENH3 and GBP CRISPR/Cas9 plasmids were created via Gibson assembly of the

linearized backbone amplified from the TK CRISPR/Cas9 plasmid (Addgene 122852) and an

oligonucleotide with the sgRNA specific for the CENH3 or GBP gene [74]. The Eukaryotic

Pathogen CRISPR guide RNA/DNA Design tool (http://grna.ctegd.uga.edu) was used to identify

CENH3 and GBP sgRNA sequences.

Amplifying transgenic parasites in mice.

The transfection and amplification of parasites in immunocompromised mice were conducted as

previously described [74-77]. Briefly, 1×10^8 freshly excysted sporozoites were electroporated

with 100 μg of repair plasmid and 66 μg of CRISPR/Cas9 plasmid, then resuspended in 300 μL

of cold DPBS. *Ifngr1$^{-/-}$* mice were first orally gavaged with an 8% (wt/vol) sodium bicarbonate

solution to neutralize stomach acid, followed by the administration of the electroporated

sporozoites. To facilitate the selection of transgenic parasites, paromomycin (16 g/L) was

provided in the drinking water 24 h post-infection. Fecal pellets were collected at intervals post-

infection for luciferase assays and diagnostic PCR analysis to verify parasite growth and proper

gene insertion. Oocysts were isolated from fecal material using bleach treatment, and 20,000

oocysts per mouse were employed for a subsequent round of amplification in NSG mice.

Luciferase assays for mouse experiments. Fecal samples collected from infected mice were

analyzed with the Promega Nano-Glo luciferase assay kit, as previously described [77]. The

luciferase activity was measured using the BioTek Cytation 3 cell imaging multi-mode reader.

The number of relative luminescence units per milligram of feces was obtained by dividing the

average values from two technical replicates by the fecal sample weight.

Quantifying oocyst shedding in mouse experiments. DNA was extracted from fecal pellets

collected from infected mice using the QIAamp PowerFecal DNA kit, following previously

established methods [74, 76, 77]. Quantitative PCR (qPCR) analysis was conducted to determine

the number of oocysts at each time point based on *C. parvum* glyceraldehyde-3-phosphate

dehydrogenase (GAPDH) primers and a standard curve derived from a dilution series (**S2 Table**)

[74]. Genomic DNA (gDNA) amounts were quantified using the QuantStudio 3 real-time PCR

system.

Confirmation of insertions in transgenic parasites by PCR.

To confirm the correct integration of the homology repair template, extracted gDNA from

murine fecal samples and wild type controls were used as the template. PCR amplification was

performed using 1 µL of the extracted gDNA, Q5 Hot Start high-fidelity 2X master mix or

PrimeSTAR GXL premix (TaKaRa), and primers at a final concentration of 500 nM on a Veriti

96-well thermal cycler. All PCR products were resolved on a 1% agarose gel containing GelRed

(1:10,000) and visualized using the ChemiDoc MP imaging system.

Oocyst purification from fecal material and storage.

The *C. parvum* isolate AUCP -1 used for transfections was purified from fecal material collected

from male Holstein calves at the University of Illinois at Urbana-Champaign [78]. All

procedures involving the calves received approval from the Institutional Animal Care and Use

Committee (IACUC). Transgenic oocysts were purified in-house from fecal material collected

from NSG mice, using a saturated sodium chloride solution for oocyst flotation [79]. Post-

purification, oocysts were stored at 4°C in phosphate-buffered saline (PBS) for up to three

months following fecal collection.

Structural predictions

The protein structure of the G-quadruplex binding protein (GBP, *cgd1_3530*) was predicted

using Alphafold 3 (www.alphafoldserver.com). Using ChimeraX

(https://www.cgl.ucsf.edu/chimerax/), we modeled how the G-quadruplex structure formed by

the *Cp*GBP repetitive binding sequence (GGTTA) [51] interacts and is stabilized by the CpGBP

protein.

Channel colocalization quantification

Co-localization analysis was performed using the Coloc2 plugin (FIJI/ImageJ), using default

thresholding parameters. A 2D intensity histogram was generated as part of the Coloc2 output to

visualize the pixel-wise distribution of intensities from both channels and a Pearson correlation

coefficient was quantified.

Parasite Preparation and Lysis for CUT&RUN experiments.

The CENH3-3HA line of *T. gondii* [27] was obtained from the laboratory of Dr. Kami Kim at

University of South Florida. *Tg*CENH3-3HA tachyzoites were propagated in human foreskin

fibroblasts and harvested using established protocols [80]. *C. parvum* sporozoites ($4x10^7$) and *T.*

gondii tachyzoites (1x10^7) were centrifuged at 2,500 RPM for 5 min, supernatants removed, and

parasites were resuspended in 2 mL of cold parasite lysis buffer (10 mM HEPES pH 7.9, 1.5 mM

MgCl2, 10 mM KCl, 0.65% NP40 (supplemented with IGEPAL) and 0.5 mM PMSF

(supplemented with cOmplete EDTA free protease inhibitor cocktail and sterile distilled H$_2$O),

incubated for 10 min on ice, and centrifuged at 3,000 RPM for 10 min at 4°C (previously

described in [81, 82]).

CUT&RUN Processing, Library Preparation, Quality Control & Sequencing.

Following lysis treatment, the lysis buffer was removed, and the nuclei pellet was washed

following the CUT&RUN Kit Version 4 protocol (EpiCypher, Inc, Catalog No. 14-1048).

Isolated nuclei were incubated with either 1.0 μg of rat anti-HA antibody, 0.5 μg of rabbit anti-

H3K4me3 positive control antibody, or 0.5 μg of mouse anti-IgG negative control antibody.

Chromatin digestion and release steps were performed according to the EpiCypher protocol, and

5 ng of CUT&RUN enriched DNA was used for library preparation according to the CUT&RUN

Library Prep Kit User Manual Version 1.5 (EpiCypher, Inc, Catalog No. 14-1001). The

concentrations of CUT&RUN library samples were determined via Qubit and quality control

were assessed with an Agilent 4200 TapeStation.

CUT&RUN Analysis & Visualization.

Multiple CUT&RUN runs were preformed to obtain biological replicates of each sample. DNA

products were sequenced on an Illumina NovaSeq 6000 using 150-bp paired-ends at the GTAC

sequencing core at Washington University in St. Louis. Sequence reads obtained from Illumina

were trimmed for adapter sequences and poor sequence quality (phred score < 25 and trimmed

length < 50) using fastp v0.23.4. The trimmed reads were mapped to the reference *C. parvum*

IOWA-CDC genome using BWA-MEM v0.7.17 [83]. Duplicate reads were marked and

removed in the bam files using the MarkDuplicates algorithm of picard v3.1.1

(https://broadinstitute.github.io/picard/) with REMOVE_DUPLICATES=true,

CREATE_INDEX=true, and ASSUME_SORTED=true. Reads with mapping lengths less than

50 bp were further removed from the bam files using SAMtools v1.7

(http://samtools.sourceforge.net/). Genome coverage and sequence depth were estimated using

the mpileup algorithm of SAMtools. The read depth of each locus was normalized by dividing by

the total number of counts for all loci, then scaled by multiplying by the number of loci in the

genome. Input counts were subtracted from negative control counts. Data was visualized by a R

package 'ggplot2' (https://www.rdocumentation.org/packages/ggplot2). Motifs within

centromeric windows were identified using MEME v5.5.8 (https://meme-

suite.org/meme/tools/meme). Chromosomal distributions of intergenic regions, centromeric

windows, and motifs were visualized with Circos plot generated in TBtools-II v2.311

(https://github.com/CJ-Chen/TBtools-II).

REFERENCES:

1. Tyzzer E. An extracellular coccidium, Cryptosporidium muris (gen. et sp. nov.), of the gastric glands of the common mouse. J Med Res. 1910;23:487-509. PubMed Central PMCID: PMCPMC2098948.
2. Greenberg PD, Koch J, Cello JP. Diagnosis of Cryptosporidium parvum in patients with severe diarrhea and AIDS. Dig Dis Sci. 1996;41(11):2286-90.
3. Striepen B. Parasitic infections: Time to tackle cryptosporidiosis. Nature. 2013;503(7475):189-91. PubMed Central PMCID: PMC24236315.
4. Tzipori S, Widmer G. A hundred-year retrospective on cryptosporidiosis. Trends Parasitol. 2008;24(4):184-9. Epub 20080307. doi: 10.1016/j.pt.2008.01.002. PubMed PMID: 18329342; PubMed Central PMCID: PMCPMC2716703.
5. Sow SO, Muhsen K, Nasrin D, Blackwelder WC, Wu Y, Farag TH, et al. The Burden of Cryptosporidium Diarrheal Disease among Children < 24 Months of Age in Moderate/High Mortality Regions of Sub-Saharan Africa and South Asia, Utilizing Data from the Global Enteric Multicenter Study (GEMS). PLoS Negl Trop Dis. 2016;10(5):e0004729. Epub 2016/05/25. doi: 10.1371/journal.pntd.0004729. PubMed PMID: 27219054; PubMed Central PMCID: PMCPMC4878811.
6. Kotloff KL, Nataro JP, Blackwelder WC, Nasrin D, Farag TH, Panchalingam S, et al. Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. Lancet. 2013;382(9888):209-22.
7. Wanyiri JW, Kanyi H, Maina S, Wang DE, Steen A, Ngugi P, et al. Cryptosporidiosis in HIV/AIDS patients in Kenya: clinical features, epidemiology, molecular characterization and antibody responses. Am J Trop Med Hyg. 2014;91(2):319-28. Epub 2014/05/29. doi: 10.4269/ajtmh.13-0254. PubMed PMID: 24865675; PubMed Central PMCID: PMCPMC4125256.
8. Funkhouser-Jones LJ, Ravindran S, Sibley LD. Defining Stage-Specific Activity of Potent New Inhibitors of Cryptosporidium parvum Growth In Vitro. mBio. 2020;11(2). Epub 2020/03/05. doi: 10.1128/mBio.00052-20. PubMed PMID: 32127445; PubMed Central PMCID: PMCPMC7064746.
9. English ED, Guerin A, Tandel J, Striepen B. Live imaging of the Cryptosporidium parvum life cycle reveals direct development of male and female gametes from type I meronts. PLoS Biol. 2022;20(4):e3001604. Epub 2022/04/19. doi: 10.1371/journal.pbio.3001604. PubMed PMID: 35436284; PubMed Central PMCID: PMCPMC9015140.
10. Beyer TV, Svezhova NV, Sidorenko NV, Khokhlov SE. Cryptosporidium parvum (Coccidia, Apicomplexa): Some new ultrastructural observations on its endogenous development. European Journal of Protistology. 2000;36(2):151-9. doi: 10.1016/s0932-4739(00)80034-6.
11. Pohlenz J, Bemrick WJ, Moon HW, Cheville NF. Bovine cryptosporidiosis: a transmission and scanning electron microscopic study of some stages in the life cycle and of the host-parasite relationship. Vet Pathol. 1978;15(3):417-27. Epub 1978/05/01. doi: 10.1177/030098587801500318. PubMed PMID: 98896.
12. Tzipori S, McCartney E, Lawson GH, Rowland AC, Campbell I. Experimental infection of piglets with cryptosporidium. Res Vet Sci. 1981;31(3):358-68. Epub 1981/11/01. PubMed PMID: 7342229.

13. Current WL, Reese NC. A comparison of endogenous development of three isolates of
Cryptosporidium in suckling mice. J Protozool. 1986;33(1):98-108. Epub 1986/02/01.
PubMed PMID: 3959014.
14. Keithly JS, Langreth SG, Buttle KF, Mannella CA. Electron tomographic and ultrastructural
analysis of the Cryptosporidium parvum relict mitochondrion, its associated membranes,
and organelles. J Eukaryot Microbiol. 2005;52(2):132-40. Epub 2005/04/09. doi:
10.1111/j.1550-7408.2005.04-3317.x. PubMed PMID: 15817118.
15. Wilke G. Development of an in vitro Culture System for Cryptosporidium parvum. Theses
and Dissertations: Washington University in St. Louis; 2020.
16. White MW, Suvorova ES. Apicomplexa Cell Cycles: Something Old, Borrowed, Lost, and
New. Trends Parasitol. 2018;34(9):759-71. Epub 20180802. doi:
10.1016/j.pt.2018.07.006. PubMed PMID: 30078701; PubMed Central PMCID:
PMCPMC6157590.
17. Striepen B, Jordan CN, Reiff S, van Dooren GG. Building the perfect parasite: cell division
in apicomplexa. PLoS Pathog. 2007;3(6):e78. doi: 10.1371/journal.ppat.0030078. PubMed
PMID: 17604449; PubMed Central PMCID: PMCPMC1904476.
18. Gissot M, Walker R, Delhaye S, Huot L, Hot D, Tomavo S. Toxoplasma gondii
chromodomain protein 1 binds to heterochromatin and colocalises with centromeres and
telomeres at the nuclear periphery. PLoS One. 2012;7(3):e32671. Epub 20120309. doi:
10.1371/journal.pone.0032671. PubMed PMID: 22427862; PubMed Central PMCID:
PMCPMC3302879.
19. Duraisingh MT, Voss TS, Marty AJ, Duffy MF, Good RT, Thompson JK, et al.
Heterochromatin silencing and locus repositioning linked to regulation of virulence genes
in Plasmodium falciparum. Cell. 2005;121(1):13-24. doi: 10.1016/j.cell.2005.01.036.
PubMed PMID: 15820675.
20. del Cacho E, Pages M, Gallego M, Barbero JL, Monteagudo L, Sanchez-Acedo C. Meiotic
chromosome pairing and bouquet formation during Eimeria tenella sporulation. Int J
Parasitol. 2010;40(4):453-62. Epub 20091029. doi: 10.1016/j.ijpara.2009.09.008. PubMed
PMID: 19837073.
21. Dey G, Culley S, Curran S, Schmidt U, Henriques R, Kukulski W, et al. Closed mitosis
requires local disassembly of the nuclear envelope. Nature. 2020;585(7823):119-23. Epub
20200826. doi: 10.1038/s41586-020-2648-3. PubMed PMID: 32848252; PubMed Central
PMCID: PMCPMC7610560.
22. Matthews H, Duffy CW, Merrick CJ. Checks and balances? DNA replication and the cell
cycle in Plasmodium. Parasit Vectors. 2018;11(1):216. Epub 20180327. doi:
10.1186/s13071-018-2800-1. PubMed PMID: 29587837; PubMed Central PMCID:
PMCPMC5872521.
23. Rout MP, Obado SO, Schenkman S, Field MC. Specialising the parasite nucleus: Pores,
lamins, chromatin, and diversity. PLoS Pathog. 2017;13(3):e1006170. Epub 20170302.
doi: 10.1371/journal.ppat.1006170. PubMed PMID: 28253370; PubMed Central PMCID:
PMCPMC5333908.
24. Suvorova ES, Francia M, Striepen B, White MW. A novel bipartite centrosome coordinates
the apicomplexan cell cycle. PLoS Biol. 2015;13(3):e1002093. Epub 20150303. doi:
10.1371/journal.pbio.1002093. PubMed PMID: 25734885; PubMed Central PMCID:
PMCPMC4348508.

25. Chen CT, Kelly M, Leon J, Nwagbara B, Ebbert P, Ferguson DJ, et al. Compartmentalized
Toxoplasma EB1 bundles spindle microtubules to secure accurate chromosome
segregation. Mol Biol Cell. 2015;26(25):4562-76. Epub 20151014. doi: 10.1091/mbc.E15-
06-0437. PubMed PMID: 26466679; PubMed Central PMCID: PMCPMC4678015.
26. Hoeijmakers WA, Flueck C, Francoijs KJ, Smits AH, Wetzel J, Volz JC, et al. Plasmodium
falciparum centromeres display a unique epigenetic makeup and cluster prior to and
during schizogony. Cell Microbiol. 2012;14(9):1391-401. Epub 20120515. doi:
10.1111/j.1462-5822.2012.01803.x. PubMed PMID: 22507744.
27. Brooks CF, Francia ME, Gissot M, Croken MM, Kim K, Striepen B. Toxoplasma gondii
sequesters centromeres to a specific nuclear region throughout the cell cycle. Proc Natl
Acad Sci U S A. 2011;108(9):3767-72. Epub 20110214. doi: 10.1073/pnas.1006741108.
PubMed PMID: 21321216; PubMed Central PMCID: PMCPMC3048097.
28. Arnot DE, Ronander E, Bengtsson DC. The progression of the intra-erythrocytic cell cycle of
Plasmodium falciparum and the role of the centriolar plaques in asynchronous mitotic
division during schizogony. Int J Parasitol. 2011;41(1):71-80. Epub 20100917. doi:
10.1016/j.ijpara.2010.07.012. PubMed PMID: 20816844.
29. Iwanaga S, Khan SM, Kaneko I, Christodoulou Z, Newbold C, Yuda M, et al. Functional
identification of the Plasmodium centromere and generation of a Plasmodium artificial
chromosome. Cell Host Microbe. 2010;7(3):245-55. doi: 10.1016/j.chom.2010.02.010.
PubMed PMID: 20227667; PubMed Central PMCID: PMCPMC2996609.
30. Absalon S. A dynamic barrier: remodeling of the nuclear envelope during closed mitosis in
malaria parasites. mSphere. 2025;10(7):e0099924. Epub 20250609. doi:
10.1128/msphere.00999-24. PubMed PMID: 40488521; PubMed Central PMCID:
PMCPMC12306155.
31. Weiner A, Dahan-Pasternak N, Shimoni E, Shinder V, von Huth P, Elbaum M, et al. 3D
nuclear architecture reveals coupled cell cycle dynamics of chromatin and nuclear pores in
the malaria parasite Plasmodium falciparum. Cell Microbiol. 2011;13(7):967-77. Epub
20110428. doi: 10.1111/j.1462-5822.2011.01592.x. PubMed PMID: 21501361.
32. Kursel LE, Malik HS. Centromeres. Curr Biol. 2016;26(12):R487-R90. doi:
10.1016/j.cub.2016.05.031. PubMed PMID: 27326706.
33. Hegemann JH, Fleig UN. The centromere of budding yeast. Bioessays. 1993;15(7):451-60.
doi: 10.1002/bies.950150704. PubMed PMID: 8379948.
34. Kelly JM, McRobert L, Baker DA. Evidence on the chromosomal location of centromeric
DNA in Plasmodium falciparum from etoposide-mediated topoisomerase-II cleavage.
Proc Natl Acad Sci U S A. 2006;103(17):6706-11. Epub 20060414. doi:
10.1073/pnas.0510363103. PubMed PMID: 16617116; PubMed Central PMCID:
PMCPMC1458945.
35. Viera A, Page J, Rufas JS. Inverted meiosis: the true bugs as a model to study. Genome Dyn.
2009;5:137-56. doi: 10.1159/000166639. PubMed PMID: 18948713.
36. Mandrioli M, Manicardi GC. Holocentric chromosomes. PLoS Genet. 2020;16(7):e1008918.
Epub 20200730. doi: 10.1371/journal.pgen.1008918. PubMed PMID: 32730246; PubMed
Central PMCID: PMCPMC7392213.
37. Marques A, Drinnenberg IA. Same but different: Centromere regulations in holocentric
insects and plants. Curr Opin Cell Biol. 2025;93:102484. Epub 20250220. doi:
10.1016/j.ceb.2025.102484. PubMed PMID: 39983583.

38. Bensasson D. Evidence for a high mutation rate at rapidly evolving yeast centromeres. BMC
Evol Biol. 2011;11:211. Epub 20110718. doi: 10.1186/1471-2148-11-211. PubMed
PMID: 21767380; PubMed Central PMCID: PMCPMC3155921.
39. Csink AK, Henikoff S. Something from nothing: the evolution and utility of satellite repeats.
Trends Genet. 1998;14(5):200-4. doi: 10.1016/s0168-9525(98)01444-9. PubMed PMID:
9613205.
40. Talbert PB, Henikoff S. Centromeres drive and take a break. Chromosome Res.
2025;33(1):17. Epub 20250804. doi: 10.1007/s10577-025-09777-z. PubMed PMID:
40759764; PubMed Central PMCID: PMCPMC12321929.
41. Henikoff S, Ahmad K, Malik HS. The centromere paradox: stable inheritance with rapidly
evolving DNA. Science. 2001;293(5532):1098-102. doi: 10.1126/science.1062939.
PubMed PMID: 11498581.
42. Talbert PB, Henikoff S. What makes a centromere? Exp Cell Res. 2020;389(2):111895. Epub
20200206. doi: 10.1016/j.yexcr.2020.111895. PubMed PMID: 32035948.
43. Skene PJ, Henikoff S. An efficient targeted nuclease strategy for high-resolution mapping of
DNA binding sites. Elife. 2017;6. Epub 20170116. doi: 10.7554/eLife.21856. PubMed
PMID: 28079019; PubMed Central PMCID: PMCPMC5310842.
44. Herrera-Solorio AM, Vembar SS, MacPherson CR, Lozano-Amado D, Meza GR,
Xoconostle-Cazares B, et al. Clipped histone H3 is integrated into nucleosomes of DNA
replication genes in the human malaria parasite Plasmodium falciparum. EMBO Rep.
2019;20(4). Epub 20190304. doi: 10.15252/embr.201846331. PubMed PMID: 30833341;
PubMed Central PMCID: PMCPMC6446197.
45. Verma G, Surolia N. Plasmodium falciparum CENH3 is able to functionally complement
Cse4p and its, C-terminus is essential for centromere function. Mol Biochem Parasitol.
2013;192(1-2):21-9. Epub 20131206. doi: 10.1016/j.molbiopara.2013.11.002. PubMed
PMID: 24316361.
46. Naumov A, Kratzer S, Ting LM, Kim K, Suvorova ES, White MW. The Toxoplasma
Centrocone Houses Cell Cycle Regulatory Factors. mBio. 2017;8(4). Epub 20170822. doi:
10.1128/mBio.00579-17. PubMed PMID: 28830940; PubMed Central PMCID:
PMCPMC5565962.
47. Mahajan B, Selvapandiyan A, Gerald NJ, Majam V, Zheng H, Wickramarachchi T, et al.
Centrins, cell cycle regulation proteins in human malaria parasite Plasmodium falciparum.
J Biol Chem. 2008;283(46):31871-83. Epub 20080808. doi: 10.1074/jbc.M800028200.
PubMed PMID: 18693242.
48. Ay F, Bunnik EM, Varoquaux N, Bol SM, Prudhomme J, Vert JP, et al. Three-dimensional
modeling of the P. falciparum genome during the erythrocytic cycle reveals a strong
connection between genome architecture and gene expression. Genome Res.
2014;24(6):974-88. Epub 20140326. doi: 10.1101/gr.169417.113. PubMed PMID:
24671853; PubMed Central PMCID: PMCPMC4032861.
49. Bunnik EM, Cook KB, Varoquaux N, Batugedara G, Prudhomme J, Cort A, et al. Changes in
genome organization of parasite-specific gene families during the Plasmodium
transmission stages. Nat Commun. 2018;9(1):1910. Epub 20180515. doi: 10.1038/s41467-
018-04295-5. PubMed PMID: 29765020; PubMed Central PMCID: PMCPMC5954139.
50. Hollin T, Le Roch KG. From Genes to Transcripts, a Tightly Regulated Journey in
Plasmodium. Front Cell Infect Microbiol. 2020;10:618454. Epub 20201217. doi:

10.3389/fcimb.2020.618454. PubMed PMID: 33425787; PubMed Central PMCID:
PMCPMC7793691.
51. Liu C, Wang L, Lancto CA, Abrahamsen MS. Characterization of a Cryptosporidium parvum
protein that binds single-stranded G-strand telomeric DNA. Mol Biochem Parasitol.
2009;165(2):132-41. Epub 20090204. doi: 10.1016/j.molbiopara.2009.01.013. PubMed
PMID: 19428660.
52. McIntosh JR. Mitosis. Cold Spring Harb Perspect Biol. 2016;8(9). Epub 20160901. doi:
10.1101/cshperspect.a023218. PubMed PMID: 27587616; PubMed Central PMCID:
PMCPMC5008068.
53. Perez-Cadahia B, Drobic B, Davie JR. H3 phosphorylation: dual role in mitosis and
interphase. Biochem Cell Biol. 2009;87(5):695-709. doi: 10.1139/O09-053. PubMed
PMID: 19898522.
54. Guizetti J, Martins RM, Guadagnini S, Claes A, Scherf A. Nuclear pores and perinuclear
expression sites of var and ribosomal DNA genes correspond to physically distinct regions
in Plasmodium falciparum. Eukaryot Cell. 2013;12(5):697-702. Epub 20130308. doi:
10.1128/EC.00023-13. PubMed PMID: 23475702; PubMed Central PMCID:
PMCPMC3647773.
55. Gerald N, Mahajan B, Kumar S. Mitosis in the human malaria parasite Plasmodium
falciparum. Eukaryot Cell. 2011;10(4):474-82. Epub 20110211. doi: 10.1128/EC.00314-
10. PubMed PMID: 21317311; PubMed Central PMCID: PMCPMC3127633.
56. Francia ME, Striepen B. Cell division in apicomplexan parasites. Nat Rev Microbiol.
2014;12(2):125-36. Epub 20140102. doi: 10.1038/nrmicro3184. PubMed PMID:
24384598.
57. Steiner FA, Henikoff S. Holocentromeres are dispersed point centromeres localized at
transcription factor hotspots. Elife. 2014;3:e02025. Epub 20140101. doi:
10.7554/eLife.02025. PubMed PMID: 24714495; PubMed Central PMCID:
PMCPMC3975580.
58. Xia J, Venkat A, Bainbridge RE, Reese ML, Le Roch KG, Ay F, et al. Third-generation
sequencing revises the molecular karyotype for Toxoplasma gondii and identifies
emerging copy number variants in sexual recombinants. Genome Res. 2021;31(5):834-51.
Epub 20210427. doi: 10.1101/gr.262816.120. PubMed PMID: 33906962; PubMed Central
PMCID: PMCPMC8092015.
59. Blauwkamp J, Ambekar SV, Hussain T, Mair GR, Beck JR, Absalon S. Nuclear pore
complexes undergo Nup221 exchange during blood-stage asexual replication of parasites.
Msphere. 2024;9(12). doi: 10.1128/msphere.00750-24. PubMed PMID:
WOS:001352594000001.
60. Haase R, Puthenpurackal A, Maco B, Guerin A, Soldati-Favre D. gamma-tubulin complex
controls the nucleation of tubulin-based structures in Apicomplexa. Mol Biol Cell.
2024;35(9):ar121. Epub 20240724. doi: 10.1091/mbc.E24-03-0100. PubMed PMID:
39046777; PubMed Central PMCID: PMCPMC11449391.
61. Edwards-Smallbone J, Jensen AL, Roberts LE, Totanes FIG, Hart SR, Merrick CJ.
Plasmodium falciparum GBP2 Is a Telomere-Associated Protein That Binds to G-
Quadruplex DNA and RNA. Front Cell Infect Microbiol. 2022;12:782537. Epub
20220222. doi: 10.3389/fcimb.2022.782537. PubMed PMID: 35273922; PubMed Central
PMCID: PMCPMC8902816.

62. Zhao N, Gong P, Li Z, Cheng B, Li J, Yang Z, et al. Identification of a telomeric DNA-
binding protein in Eimeria tenella. Biochem Biophys Res Commun. 2014;451(4):599-602.
Epub 20140813. doi: 10.1016/j.bbrc.2014.08.030. PubMed PMID: 25128826.
63. Melters DP, Paliulis LV, Korf IF, Chan SW. Holocentric chromosomes: convergent
evolution, meiotic adaptations, and genomic analysis. Chromosome Res. 2012;20(5):579-
93. doi: 10.1007/s10577-012-9292-1. PubMed PMID: 22766638.
64. Lucek K, Augustijnen H, Escudero M. A holocentric twist to chromosomal speciation?
Trends Ecol Evol. 2022;37(8):655-62. Epub 20220426. doi: 10.1016/j.tree.2022.04.002.
PubMed PMID: 35484024.
65. Helsen J, Ramachandran K, Sherlock G, Dey G. Progressive coevolution of the yeast
centromere and kinetochore. Nature. 2025. Epub 20251126. doi: 10.1038/s41586-025-
09779-1. PubMed PMID: 41299172; PubMed Central PMCID: PMCPMC12925627.
66. Funkhouser-Jones LJ, Xu R, Wilke G, Fu Y, Schriefer LA, Makimaa H, et al. Microbiota-
produced indole metabolites disrupt mitochondrial function and inhibit Cryptosporidium
parvum growth. Cell Rep. 2023;42(7):112680. Epub 2023/06/29. doi:
10.1016/j.celrep.2023.112680. PubMed PMID: 37384526.
67. Pardy RD, Wallbank BA, Striepen B, Hunter CA. Immunity to Cryptosporidium: insights
into principles of enteric responses to infection. Nat Rev Immunol. 2024;24(2):142-55.
Epub 20230911. doi: 10.1038/s41577-023-00932-3. PubMed PMID: 37697084.
68. Martinez-Perez E, Schvarzstein M, Barroso C, Lightfoot J, Dernburg AF, Villeneuve AM.
Crossovers trigger a remodeling of meiotic chromosome axis composition that is linked to
two-step loss of sister chromatid cohesion. Genes Dev. 2008;22(20):2886-901. doi:
10.1101/gad.1694108. PubMed PMID: 18923085; PubMed Central PMCID:
PMCPMC2569886.
69. Kimball A, Funkhouser-Jones L, Huang W, Xu R, Witola WH, Sibley LD. Mendelian
segregation and high recombination rates facilitate genetic analyses in Cryptosporidium
parvum. PLoS Genet. 2024;20(6):e1011162. Epub 20240617. doi:
10.1371/journal.pgen.1011162. PubMed PMID: 38885280; PubMed Central PMCID:
PMCPMC11213348.
70. Shaw S, Cohn IS, Baptista RP, Xia G, Melillo B, Agyabeng-Dadzie F, et al. Genetic crosses
within and between species of Cryptosporidium. Proc Natl Acad Sci U S A.
2024;121(1):e2313210120. Epub 2023/12/26. doi: 10.1073/pnas.2313210120. PubMed
PMID: 38147547; PubMed Central PMCID: PMCPMC10769859.
71. Wilke G, Ravindran S, Funkhouser-Jones L, Barks J, Wang Q, VanDussen KL, et al.
Monoclonal antibodies to intracellular stages of *Cryptosporidium parvum* define life cycle
progression in vitro. Msphere. 2018;3(3):e00124-18. Epub 2018/06/01. doi:
10.1128/mSphere.00124-18. PubMed PMID: 29848759; PubMed Central PMCID:
PMCPMC5976880.
72. Dos Santos Pacheco N, Soldati-Favre D. Coupling Auxin-Inducible Degron System with
Ultrastructure Expansion Microscopy to Accelerate the Discovery of Gene Function in
Toxoplasma gondii. Methods Mol Biol. 2021;2369:121-37. doi: 10.1007/978-1-0716-
1681-9_8. PubMed PMID: 34313987.
73. Xu R, Beatty WL, Greigert V, Witola WH, Sibley LD. Multiple pathways for glucose
phosphate transport and utilization support growth of Cryptosporidium parvum. Nat
Commun. 2024;15(1):380. Epub 20240109. doi: 10.1038/s41467-024-44696-3. PubMed
PMID: 38191884; PubMed Central PMCID: PMCPMC10774378.

74. Wilke G, Funkhouser-Jones L, Wang Y, Ravindran S, Wang Q, Beatty WL, et al. Forward
genetics in *Cryptosporidium* enabled by complete in vitro development in stem cell-
derived intestinal epithlium. Cell Host Microbe. 2019;doi.org/10.2139/ssrn.3331307.
75. Xu R, Beatty WL, Greigert V, Witola WH, Sibley LD. Multiple pathways for glucose
phosphate transport and utilization support growth of Cryptosporidium parvum. bioRxiv.
2023. Epub 2023/07/10. doi: 10.1101/2023.06.27.546703. PubMed PMID: 37425855;
PubMed Central PMCID: PMCPMC10327089.
76. Akey ME, Xu R, Ravindran S, Funkhouser-Jones L, Sibley LD. Apical Secretory
Glycoprotein Complex Contributes to Cell Attachment and Entry by Cryptosporidium
parvum. mBio. 2023;14(1):e0306422. Epub 2023/02/02. doi: 10.1128/mbio.03064-22.
PubMed PMID: 36722968; PubMed Central PMCID: PMCPMC9973360.
77. Xu R, Feng Y, Xiao L, Sibley LD. Insulinase-like Protease 1 Contributes to Macrogamont
Formation in Cryptosporidium parvum. mBio. 2021;12(2). Epub 2021/03/11. doi:
10.1128/mBio.03405-20. PubMed PMID: 33688009; PubMed Central PMCID:
PMCPMC8092296.
78. Kuhlenschmidt TB, Rutaganira FU, Long S, Tang K, Shokat KM, Kuhlenschmidt MS, et al.
Inhibition of Calcium-Dependent Protein Kinase 1 (CDPK1) In Vitro by
Pyrazolopyrimidine Derivatives Does Not Correlate with Sensitivity of Cryptosporidium
parvum Growth in Cell Culture. Antimicrob Agents Chemother. 2015;60(1):570-9.
79. Current WL. Techniques and laboratory maintenance of *Cryptosporidium*. In: Dubey JP,
Speer CA, Fayer R, editors. Cryptosporidiosis of Man and Animals. Boca Raton, FL: CRC
Press; 1990. p. 44-77.
80. Mehta S, Sibley LD. Actin depolymerizing factor controls actin turnover and gliding motility
in Toxoplasma gondii. Mol Biol Cell. 2011;22(8):1290-9. Epub 20110223. doi:
10.1091/mbc.E10-12-0939. PubMed PMID: 21346192; PubMed Central PMCID:
PMCPMC3078074.
81. Behnke MS, Radke JB, Smith AT, Sullivan WJ, Jr., White MW. The transcription of
bradyzoite genes in Toxoplasma gondii is controlled by autonomous promoter elements.
Mol Microbiol. 2008;68(6):1502-18. Epub 20080421. doi: 10.1111/j.1365-
2958.2008.06249.x. PubMed PMID: 18433450; PubMed Central PMCID:
PMCPMC2440561.
82. Waldman BS, Schwarz D, Wadsworth MH, 2nd, Saeij JP, Shalek AK, Lourido S.
Identification of a Master Regulator of Differentiation in Toxoplasma. Cell.
2020;180(2):359-72 e16. Epub 20200116. doi: 10.1016/j.cell.2019.12.013. PubMed
PMID: 31955846; PubMed Central PMCID: PMCPMC6978799.
83. Huang W, Tang K, Chen C, Arrowood MJ, Chen M, Guo Y, et al. Sequence introgression
from exogenous lineages underlies genomic and biological differences among
Cryptosporidium parvum IOWA lines. Water Res. 2024;254:121333. Epub 20240217.
doi: 10.1016/j.watres.2024.121333. PubMed PMID: 38402753; PubMed Central PMCID:
PMCPMC10994760.
1018
1019

1020

Figure 1: Characterization of centromeric histone H3 (CENH3) and histone H3 candidates

in *C. parvum* and construction of Histone H3-3HA tagged strains. (A) Phylogenetic

relationship of *C. parvum* H3 and centromeric H3 (CENH3) histones and their corresponding top

BLASTP hits in other eukaryotes, apicomplexan parasites, *Saccharomyces cerevisiae*, or *Homo*

sapiens. Cyan stars represent annotated histone H3 genes. (B) Protein sequence alignment of H3

(cyan) and CENH3 (black) in *C. parvum*. Amino acids that are identical between at least two

sequences are highlighted in grey. (C) Diagram of the targeting constructs designed to replace

the endogenous *tk* locus with a second copy of either of the genes annotated as Histone H3-like

proteins in *C. parvum* (*cgd3_2540* or *cgd4_3220*), a 3HA tag, and a Nluc-P2A-NeoR cassette.

(D) Immunofluorescence staining of transgenic H3.1-3HA or (E) H3.2-3HA parasites. HCT-8

cells were infected with transgenic oocysts, fixed at 18 hpi, and stained with rat anti-HA

followed by the secondary antibody Alexa Fluor 488 goat anti-rat IgG. Hoechst was used for

nuclear staining. Images were acquired as Z-stacks using a Zeiss LSM-880 Laser Scanning

Confocal microscope equipped with Airyscan (LSCM-A) and are presented with orthogonal

views. Scale bars, 1 μm. Data represent two biological replicates, each with 3-5 technical

replicates.

1037

Figure 2. The staining pattern of *C. parvum* centromeric histone H3 (CENH3) is distinct from *Toxoplasma* and remains diffuse during mitosis. (A) Diagram of the targeting construct designed to add a 3HA tag and Nluc-P2A-Neo^R cassette to the C-terminus of the *C. parvum* CENH3 (*cgd4_2030*). (B) HCT-8 cells were infected with CENH3-3HA *C. parvum* parasites, fixed at 22 hpi, and stained with rat anti-HA, followed by the secondary antibody Alexa Fluor 488 goat anti-rat IgG followed by Hoechst staining. Images were acquired as Z-stacks using

LSCM-A and are presented with orthogonal views. Scale bars, 1 μm. (C) Cartoon depiction

corresponding to the microscopy image in panel (B). In *Cryptosporidium*, a late-stage meront

contains eight nuclei that have recently completed cytokinesis to form mature merozoites. The

CENH3-3HA signal within each nucleus displays a diffuse staining pattern in *C. parvum*. (D)

HCT-8 cells were infected with CENH3-3HA *T. gondii* parasites, fixed at 24 hpi, and stained

with rat anti-HA and rabbit anti-aldolase (ALD) to visualize the cytosol, followed by secondary

antibodies Alexa Fluor 488 goat anti-rat IgG and Alexa Fluor 568 anti-rabbit IgG. Hoechst was

used for nuclear staining. Images were acquired as Z-stacks using LSCM-A and are presented

with orthogonal views. Scale bars, 1 μm. (E) Cartoon depiction corresponding to the microscopy

image in panel (D). In *T. gondii* tachyzoites, the CENH3-3HA signal within each nucleus

displays a discrete staining pattern. (F) HCT-8 cells were infected with excysted *C. parvum*

sporozoites for 2 h, then washed twice to remove extracellular parasites. Cultures were fixed at

30 min increments between 6-9 hpi to identify the first (trophozoites with one nucleus), second

(early meronts with 2 nuclei), and third (mid-stage meronts with 4 nuclei) mitotic divisions

during the first round of merogony by widefield microscopy. Scale bars, 1 μm. (G) Cultures were

infected with CENH3-3HA *C. parvum* sporozoites for 2 h, washed, and fixed at 6.5 hpi to

capture the first round of mitosis. Cultures were stained with rat anti-HA, rabbit anti-centrin-1,

followed by secondary antibodies Alexa Fluor 488 goat anti-rat IgG, Alexa Fluor 568 goat anti-

rabbit IgG, and lastly Hoechst. Parasites undergoing mitosis were identified by having two

centrin-1 points per nucleus, whereas parasites in interphase had only one centrin-1 point.

Images were acquired as Z-stacks using LSCM-A and are presented with orthogonal views.

Scale bars, 1 μm. Data represent 2-3 biological replicates, each with 3-5 technical replicates.

1067

Figure 3: The telomere (G-strand binding protein, GBP) staining pattern is apical in *C.*

***parvum* and overlaps with the centromere (CENH3) staining pattern.** (A) The 3' overhang

single-stranded DNA at the end of telomeres contains repetitive sequences (GGTTTA)$_n$ in *C.*

parvum that form G-quadruplex structures. (B) The G-quadruplex binding protein (GBP) is

essential for forming and stabilizing these structures. The sequence of *Cp*GBP and its binding

motif has been previously determined and the protein structure is modeled by Alphafold and

ChimeraX [51]. (C) Diagram of the targeting construct designed to add a 3HA tag and Nluc-

P2A-Neo^R cassette to the C-terminus of the G-quadruplex binding proteins gene *(*GBP*,

cgd1_3530)* (D) HCT-8 cells were infected with GBP-3HA parasites, fixed at 6.5 hpi to capture

early meronts (2 nuclei), and stained with rat anti-HA, followed by the secondary antibody Alexa

Fluor 488 goat anti-rat IgG. Hoechst was used for nuclear staining. Images were acquired as Z-

stacks using LSCM-A and are presented with orthogonal views and the scale bar is 1 μm. (E)

Diagram of the targeting construct designed to replace the endogenous *tk* locus with a second

copy of CENH3 N-terminally tagged with 3HA and GBP with a C-terminal 3xTy tag separated

with a P2A split peptide. (F) Immunofluorescence staining of transgenic 3HA-CENH3-GBP-3Ty

parasites. HCT-8 cells were infected with transgenic oocysts, fixed at 6.5 hpi to capture early

meronts (2 nuclei), and stained with rat anti-HA and mouse anti-Ty, followed by a secondary

antibody stain containing Alexa Fluor 488 goat anti-rat IgG and Alexa Fluor 568 goat-anti

mouse. Hoechst was used for nuclear staining. Images were acquired as Z-stacks using LSCM-A

and are presented with orthogonal views. Scale bars, 1 μm. Data represent 2-3 biological

replicates, each with 3-5 technical replicates.

1090

1091

Figure 4: Mitotically active DNA, microtubules, and the nuclear pore proteins as visualized

by antibody staining in *C. parvum*. (A) HCT-8 cells were infected with wild type parasites,

fixed at 18 hpi, and stained with rabbit anti-phosphorylated Histone H3 (PH3) followed by an

Alexa Fluor 568 goat anti-rabbit IgG antibody. DNA was stained with Hoechst and a mid-stage

meront with 4 nuclei is shown. (B) Parasites were prepared identically to panel A, but were

stained with a rabbit polyclonal antibody that recognizes a subunit of the nuclear pore in

Plasmodium falciparum (NUP116) followed by an Alexa Fluor 568 goat anti-rabbit IgG

antibody and Hoechst. A late-stage meronts with 8 nuclei is shown. (C, D) HCT-8 cells were

infected with wild-type parasites, fixed at 6.5 hpi to capture early meronts (2 nuclei), and stained

with rabbit anti-centrin-1 and mouse anti-tubulin, followed by the secondary antibodies Alexa

Fluor 488 goat anti-mouse IgG, Alexa Fluor 568 goat anti-rabbit IgG, and Hoechst. The nucleus

in the first meront (C) may be an example of early mitosis, whereas the second meront (D) may

be an example of a later phase of the mitotic process. Images were acquired as Z-stacks using

LSCM-A and are presented with orthogonal views. Scale bars, 1 μm. Data represent two

biological replicates, each with 2-5 technical replicates.

1107

1108

Figure 5. Genome-wide distribution of CENH3 in *Toxoplasma gondii* and *Cryptosporidium*

parvum. (A) Genome browser view of *T. gondii* chromosome Ia showing CUT&RUN signals for

CENH3-HA (HA antibody), H3K9me3, and H3K4me3. CENH3 signal enrichment positively

correlates with H3K9me3 and inversely with H3K4me3. Peaks of CENH3 enrichment are

indicated (red arrows). (B) Genome browser view of *C. parvum* chromosome I, showing

CUT&RUN signals for CENH3-HA *C. parvum* captured with HA antibody and wild type *C.*

parvum captured with HA antibody (referred to as WT-HA), H3K9me3, and H3K4me3

(captured with antibodies). (C) Chromosomal distribution of the 58 CENH3-enriched windows

in intergenic regions of *C. parvum* chromosomes (Chr#). Numbers below each chromosome

indicate the number of windows. (D) Heatmaps of CUT&RUN signals for the groups shown in

(B) within a 400 bp window around all 58 CENH3-enriched windows . Each row represents an

individual centromeric site, ranked by signal intensity. (E) Average signal profiles for CENH3-

HA (green), WT-HA (gray), H3K9me3 (orange), and H3K4me3 (teal) across the 400 bp window

around CENH3 sites. (F) Circos plot of CENH3-associated motifs. From inner to outer rings:

chromosomes, locations of 347 CENH3-enriched windows found in coding regions (black), and

locations of 58 CENH3-enriched windows found in intergenic regions (black). Representative

sequence logos for motif 1 and motif 2 are shown.

1126

1127

1128

Supporting Information

S1 Fig. Tagged versions of the H3.1, H3.2, and CENH3 and GBP genes were inserted into the tk

locus in *Cryptosporidium parvum*.

S2 Fig. The GBP and CENH3 expression overlaps in *Cryptosporidium parvum.*

S3 Fig. The GBP and CENH3 gene were successfully tagged with HA tag and/or a spaghetti

monster HA tag in C*ryptosporidium parvum*.

S4 Fig. Additional images of centrin and tubulin staining in *Cryptosporidium parvum.*

S5 Fig. Genome-wide distribution of CENH3 in *Toxoplasma gondii*.

S6 Fig. Genome-wide distribution of CENH3 in *Cryptosporidium parvum*.

1138

1139 **S1 Table.** Transgenic *Cryptosporidium* strains used in this study.

1140 **S2 Table.** Key Resources used in the experiments.

1141 **S3 Table.** Oligonucleotides and plasmids used in this study.

1142

1143 **S1 Dataset** Information of sequencing samples from the Cut&Run and summary statistics of the

1144 data.

1145 **S2 Dataset** Genomic location of the CENH3 sites within coding regions.

1146 **S3 Dataset** Genomic location of the CENH3 sites within intergenic regions.

1147