Title: (119/135 characters) Machine Learning Guided Video Analysis Identifies Sound-Evoked Pain Behaviors from Facial Grimace and Body Cues in Mice

Benjamin J. Seicol[1,2], Amelie Valles[2,3], Anna Kohler[2], Elisabeth Glowatzki[1,2,3*], Megan B. Wood[2,4*]

[1]The Center for Hearing and Balance, Johns Hopkins School of Medicine, Baltimore, Maryland, 21205, [2]Department of Otolaryngology Head and Neck Surgery, The Johns Hopkins School of Medicine, Baltimore, Maryland, 21205 [3]Department of Neuroscience, The Johns Hopkins School of Medicine, Baltimore, Maryland, 21205. [4]Department of Hearing & Speech Sciences, Vanderbilt School of Medicine, Nashville, Tennessee, 37232.

*Corresponding authors: Megan B. Wood and Elisabeth Glowatzki

Email: megan.beers.wood@vumc.org and eglowat1@jhmi.edu

Author Contributions: BJS, AK, EG, and MBW designed research; AV, AK, and MBW performed research; AV, AK, BJS, and MBW analyzed data; BJS, EG and MBW wrote and edited the manuscript.

Competing Interest Statement: Authors report no conflict of interest.

Classification: Biological Sciences, Neuroscience, Animal Behavior and Cognition.

Keywords: Auditory pain, Deep learning, Pain behavior, Migraine, TMIE knockout

This PDF file includes:
    Main Text
    Figures 1 to 10
    Supplemental Figures 1-5
    All Figure, Supplemental Figure, Supplemental Video legends

Abstract (224/250 Words)

Humans can experience auditory pain in response to sound, either from extremely loud noise or in cases of pain hyperacusis, where typically tolerable sounds become painful. However, the mechanisms underlying auditory pain remain poorly understood. Developing behavioral methods to measure sound-evoked pain in animal models is critical for elucidating these mechanisms. Here, a deep learning-based approach was developed to measure auditory pain in freely moving mice by analyzing facial grimace and body position from video recordings during sound exposure. Facial

grimace, a validated marker of spontaneous, ongoing pain in mice, was quantified using a deep neural network trained to extract established facial features. Postural changes, additional indicators of pain, were analyzed from the same camera angle. To validate the model, a known painful state, migraine induced by injection of the neuropeptide calcitonin gene-related peptide (CGRP) was used. With this approach the ability to quantify a pain response distinct from baseline behavior was demonstrated, resulting in a defined pain threshold. Sound exposure at high intensities elicited significant changes in facial grimace and body posture, in comparison, surpassing the pain threshold established during migraine validation. These behavioral changes were absent in TMIE-knockout mice, which lack functional cochlear transduction. This automated, high-throughput framework enables objective and sensitive analysis of sound-evoked pain and provides a foundation for future studies investigating the peripheral and central mechanisms of auditory pain.

Significance (119/120 words)

This study introduces a quantitative framework for assessing affective pain using a single-camera setup and machine learning guided analysis to capture and analyze mouse behavior. By integrating two established pain metrics, facial grimace and attenuated movement, this method enables precise, non-invasive quantification of pain-related behaviors. The approach was validated with a well-characterized pain model, migraine, induced by injection of the neuropeptide CGRP, demonstrating the ability to quantify a pain response distinct from baseline behavior. Applying this framework to auditory pain, the data reveal that exposure to intense sound triggers significant nociceptive behavioral responses. These findings provide novel insights into the behavioral manifestations and neural underpinnings of auditory pain, offering a robust tool for studying the mechanisms of pain perception.

Introduction (Word Count: 581)

Humans can experience auditory pain in response to sound. Such pain can either occur as a reaction to loud sounds (> 120 dB SPL; (1-4)) or, in the case of pain hyperacusis, can be a response to typically tolerable noise exposures (5,6,7). When sound exposure becomes painful, the mechanisms that lead to auditory pain remain unknown (8) (9). Processing of that pain information may involve activation of brain regions canonically associated with pain, such as the thalamus, along with paradoxical hypoactivity in the auditory cortex (10, 11, 12). Developing sensitive

behavioral methods to measure sound-evoked pain in laboratory animals will help to determine whether animal models recapitulate human pathology and to further investigate peripheral and central mechanisms responsible for auditory pain (8, 13, 14).

Behavioral reflexes in response to sound onset are widely used to assess animals' auditory sensitivity, for example by acoustic startle response (4, 15, 16). Broadly, pain can be measured in mice either by 1) withdrawal reflex, such as, to thermal or mechanical stimuli (16), 2) operant conditions of forced-choice tasks (17), or 3) behavioral assessment to identify signatures associated with pain (e.g., facial grimace;(18), movement,(19)). During pain perception, mice 'grimace', like humans (21, 22), and therefore pain can be quantified by characteristic changes in their facial features, for example, by eyes squinting (22). Facial grimace has effectively been used to measure spontaneous, ongoing painful states in animal models, for example in the case of migraine (23).

Automated facial grimace detection of spontaneous, ongoing pain in mice, including post-operative and migraine pain, has proved comparable to human scoring, has reduced bias, and has allowed for high-throughput analysis of large datasets (22, 25, 26). Pain evoked by sound exposures for periods of time on the order of minutes is a form of spontaneous, ongoing pain (herein referred to as 'persistent pain'; (24). Persistent pain caused by sounds was analyzed using deep learning in awake, freely moving mice located inside of a sound chamber by measuring behavioral changes in facial grimace and body position (19, 25).

Here, a deep neural network was trained to extract formerly established facial features (22) from video recordings of freely moving mice exposed to various levels of sound intensity presentations. Secondly, body position measurements were added to detect changes in mouse posture and locomotion behaviors associated with pain (26). A known persistent painful state (migraine) provided validation of the model and of thresholds for detecting auditory pain. Sound presentation at high intensities produced changes in behavior surpassing this threshold, and these changes did not occur in TMIE-knockout mice that lack normal cochlear transduction of sound (27–30). Grimace measurements combined with positional information extracted by deep learning from video recordings of mice presented with sound allowed for sensitive, objective, and high-throughput analysis of mouse auditory pain.

Results

Detecting persistent pain in mice has been performed before by using facial grimace in mice (20, 23, 31, 32). Here, facial grimace detection has been adapted from work that has used six defined features of the face from the profile of a mouse's face from videos of freely moving mice (22). It was further possible to expand beyond facial grimace using the same videos and some of these same markers to analyze body position information of the mice. Average values of each facial grimace parameter and separately of four specific body position parameters were calculated within defined periods (epochs) and combined to create an inventory of pain. This inventory was validated, and pain thresholds were determined, based on a known painful stimulus (CGRP-induced migraine). This approach was subsequently used to test the level of sound that elicits pain in mice. Additionally, mice without normal cochlear function were used to determine whether sound transduction in the inner ear is necessary for the generation of auditory pain.

Detection of facial grimace parameters in freely moving mice using a deep neural network

A deep neural network was trained to measure changes in facial grimace by automatic placement of thirteen points on the face of mice with skeletons connecting these points (Figure 1). This approach allowed for high-throughput analysis of thousands of video frames from hours of recordings, greatly reducing the analysis time compared to manual annotation by blinded observers as done before with the mouse grimace scale (20). In this study, facial grimace measurements were taken from mice enclosed in a custom-built clear chamber with scent holes on only one side that encouraged mice to remain mostly in profile with respect to the camera (Figure 1A). Representative images of mice in pain and control illustrate the changes in the face, for example, facial grimace of a mouse in a pain state manifests with clear changes in ear positions and eye squinting, as well as subtle changes in snout and mouth positions (Figure 1B). Changes can also be seen in body position, such as the mouse being hunched over and the head angle pointing down, which will be described in detail below. Based on the video recordings, six facial parameters (22) were calculated from the pose estimated points on the mouse's face (Figure 1C).

DeepLabCut software (DLC; (33, 34)) was used to train a neural network model capable of automatically identifying thirteen markers on the mice with skeletons connecting these points to calculate the six facial parameters (Figure 1D-I). The changes in ear positions were mapped using 3 parameters: 1) 'Ear Ratio' (which goes up as the mouse's ear lies flat against the head thereby increasing the width in relation to the camera in profile view; Figure 1D), 2) 'Ear Position'

(which measures the angle that increases when the mouse's ear rotates further towards its posterior; Figure 1E), and 3) 'Ear Tip Tilt' (which measures the angle of the top half of the ear relative to the bottom half which increases when the ear is flattened against the back of the head; Figure 1F). These three ways of measuring changes in ear position are valuable due to the manner of grimacing in animals that are exaggerated in the ears compared with humans (35–37). The three ear-related measures also each reflect a distinct neuromuscular circuit, arising from the 1) transverse auricular muscle, 2) interscutular muscle, and 3) anterior and posterior auricular levators, respectively (38, 39). Additional face grimace parameters measured were 'Eye Ratio' (which gets smaller with 'eye squinting'; Figure 1G), 'Snout Position' (which gets smaller when the snout is more pointed during pain; Figure 1H) and Mouth Position' (which gets smaller when the bottom lip extends outwards similar to clenching of the jaw; Figure 1I).

Facial grimace scores in mice that experience migraine after CGRP injection indicate pain

Migraine can be elicited by injection of calcitonin gene-related peptide (CGRP; (40) and is a well-known painful stimulus in mice (41). Such migraine-induced pain has been detected by the formerly used mouse grimace scale (23) and was used to test the approach here implemented of detecting persistent pain with the set of facial grimace parameters as described in Figure 1. Mice were injected with CGRP solution (100 µg/kg in saline, i.p.) or injected with saline (vehicle). Earlier work has shown that CGRP-induced migraines in mice peak approximately 30 minutes following administration (23). Video recording began when mice were placed into the chamber for 10 minutes to record baseline behavior ('Baseline'; t = -10 – 0 mins; Figure 2A). Mice then received an injection of either CGRP or saline (t = 0 min) and the video recording was paused. Video acquisition was restarted 20 minutes after injection and recorded for 20 minutes ('All'; t = 20 – 40 mins) to ensure capture of the expected migraine peak (Supplemental Video 1). Each facial grimace parameter measured during Baseline and All can be visualized as a diary plot of all analyzed frames at 30 frames/s, here shown for data from an individual mouse (e.g., Ear Position; Figure 2B). It is notable that there are gaps in the dairy plot illustrating periods where the points could not all be applied to the face, for example when the mouse was turned away from the camera (Supplemental Video 1). Data from all mice (N = 12) were superimposed and are shown in Figure 2C. The expected peak of migraine-like pain from CGRP was anticipated to be around t = 30 minutes. An empirical peak emerged from combined data of facial grimace and body position

parameters (quantified below in Figure 5), showing that during a 5-minute epoch ('Peak'; t = 27 – 32 minutes) surrounding the expected peak mice exhibited significant changes in behavior. This empirical peak was therefore analyzed separately from the remaining 15 minutes of CGRP recording ('Non-Peak').

Median values from Baseline and All epochs were compared using paired, two-tailed, student t-test for each of the six facial grimace measurements, while Baseline, Peak, and Non-Peak epochs were compared using either repeated measures one-way ANOVA for balanced groups or repeated measures mixed effects model for unbalanced groups (Figure 2D-I). Significant differences compared with Baseline were found during Peak for Eye Ratio and Mouth Position, with Eye Ratio showing changes from Baseline also during All and Non-Peak. Specifically, Eye Ratio was significantly reduced in All compared with Baseline (paired t-test, All compared with Baseline median difference, -0.0781 AU, p < 0.005). Eye Ratio further showed significant decreases compared with Baseline in both Peak and Non-Peak epochs (repeated measures mixed effects model, Baseline, Peak, and Non-Peak, p = 0.0025; post hoc comparisons using Tukey's correction, Peak compared with Baseline median difference = -0.0742 AU, p < 0.05, and Non-Peak compared with Baseline median difference = -0.0681 AU, p < 0.05). Mouth Position also showed a significant decrease during Peak compared with Baseline, as well as a significant decrease during Peak compared with Non-Peak (repeated measures mixed effects model, p < 0.0001; post hoc comparisons using Tukey's correction, Peak compared with Baseline median difference = -2.98°, p < 0.005, Peak compared with Non-Peak median difference = -3.49°, p < 0.005). No significant differences from Baseline or between epochs were found in Ear Ratio, Ear Position, Ear Tip Tilt, or Snout Position. Additionally, no significant differences were observed in any measures in saline-injected mice (Supplemental Figure 1).

Figure 2. Changes measured in facial grimace during CGRP-induced migraine

Supplemental Figure 1. Animals injected with saline do not show changes to facial grimace measurements

Supplemental Video 1: CGRP Experiment Example

Body position parameters in mice experiencing migraine after CGRP injection indicate pain

A subset of the pose estimated points used for measuring facial grimace were additionally applied to generate measurements of mouse body position that could indicate pain. Body position parameters were chosen based on the observation of changed locomotive behaviors (42), whereby mice were observed to explore, rear, and generally move less during pain states, as well as maintain a hunched posture more often. To this point, body position measurements were quantified to grossly incorporate changes in these complex behaviors (Figure 3), namely, the 'Relative Nose Tip Position' (Figure 3A; related to exploring), 'Percentage of Nose Tip in Top 1/3 of Chamber' (Figure 3B; captured rearing), 'Vertical Line Crosses of Nose Tip' (Figure 3C; measured turning), and 'Face Inclination as Slope of Nose Tip to Neck' (Figure 3D; showed hunching).

Figure 3. Analysis of Mouse Body Position from Tracking of Nose Tip Coordinates

---

Body position parameters (Figure 4) were plotted in the same way as facial grimace measurements. A representative example of a diary plot showing relative nose tip position during migraine, (Figure 4A). Superimposed plots from all mice for Relative Nose Tip Position (shown in Figure 4B) illustrate a subtle decrease in variability during Peak. Appropriate statistical tests were performed to compare groups, and no significant differences were found for Relative Nose Tip Position (Figure 4C). Peak was significantly different compared to Baseline for Percentage in Top (Figure 4D; repeated measures one-way ANOVA, $p = 0.022$; post hoc comparisons using Tukey's correction, Peak compared to Baseline median difference = -8.25 percentage points, $p < 0.05$), for Vertical Line Crosses (Figure 4E, repeated measures mixed effects model, $p = 0.0004$; post hoc comparisons using Tukey's correction, Peak compared to Baseline median difference = -39 incidence, $p < 0.005$), and for Face Inclination (Figure 4F, repeated measures mixed effects model, $p = 0.03$; post hoc comparisons using Tukey's correction, Peak compared to Baseline median difference = 35.14 percentage points, $p < 0.05$). Difference in All compared with Baseline were found in both Vertical Line Crosses (paired t-test, All compared with Baseline median difference = -24.5 incidences, $p < 0.05$) and Face Inclination (paired t-test, All compared with Baseline median difference = 15.46 percentage points, $p < 0.05$). Additionally, Vertical Line Crosses was significantly decreased during Peak compared with Non-Peak (paired t-test, Peak compared with Non-Peak median difference = -19 incidences, $p < 0.001$).

Figure 4. Movement decreases during CGRP-induced painful migraine as calculated by measures of mouse body position

Combined Facial Grimace and Body Position Scores provide quantification of pain in mice
Individual facial grimace and body position parameters were combined to create two separate composite scores. Each measurement was first converted to a percentage change from Baseline (Supplemental Figure 2) and then added together to calculate the summed values for facial grimace or for body position for both the mice injected with CGRP solution and saline (Figure 5). CGRP-injected mice showed significant changes from Baseline in the summed measurements for both Facial Grimace and Body Positions Scores, while saline-injected mice showed no changes (Figure 5A-B). Specifically, both Facial Grimace Score and Body Position Score were significantly different from Baseline (Facial Grimace Score: paired t-tests, All compared with Baseline median difference = -20.59 percentage points, $p < 0.005$, Peak compared with Baseline median difference = -40.62 percentage points, $p = 0.001$, and Non-Peak compared with Baseline median difference = -19.04 percentage points, $p < 0.01$; Body Position Score: paired t-tests, All compared with Baseline median difference = -162.6 percentage points, $p < 0.001$, Peak compared with Baseline median difference = -179.5 percentage points, $p < 0.001$, and Non-Peak compared with Baseline median difference = -132.4 percentage points, $p = 0.001$). Peak was then compared with Non-Peak in CGRP-injected mice and found to be significantly different for both Facial Grimace Score (paired t-test, Peak compared with Non-Peak median difference = -21.58 percentage points, $p < 0.05$) and Body Position Score (paired t-test, Peak compared with Non-Peak median difference = -47.1 percentage points, $p < 0.05$) indicating mice behavior during the empirical peak is distinct from the non-peak periods. The Non-Peak and Peak median values were used to create thresholds for two levels of pain, Level 1 and Level 2, respectively, as indicated in Figures 5A-B: Level 1 (dotted line; -19.04 percentage points difference from Baseline for facial grimace and -132.4 percentage points for body position) and Level 2 (dashed line; -40.62 percentage points difference from Baseline for facial grimace and -179.5 percentage points for body position).

Selection of the empirical peak which led to the use of Peak and Non-Peak time windows throughout this study (starting with Figure 2) was done based on diary plots binned into one-minute segments across the entire CGRP recording period (All) (Figure 5C-D). Scores for each one-

minute segment are the summed value of all individual measurements as a median percent difference from baseline. It was found that from minute 27 to 32 the binned scores were below the mean value of all CGRP frames in Facial Grimace Score (-27.17 percentage points, Figure 5C, grey bars), and from minute 26 to 33 were below the mean value of all CGRP frames in Body Position Scores (-145.7 percentage points, Figure 5D, grey bars). The empirical peak was, therefore, selected based on the criteria where both scores were below the mean of all frames for both scores (i.e., t = 27 – 32 minutes). It is noteworthy that other one-minute binned values outside of the empirical peak fall below the mean from the entire CGRP recording (Figure 5C-D, grey bars), however, the longest continuous period where both scores meet these criteria around the 30-minute mark was used as the empirical peak. This method confirmed the expected peak of CGRP-induced migraine pain occurs around 30 minutes after injection (23). Due to a higher sampling rate in this study (30 frames/s) the empirical peak time was found to span 5 minutes and was used to define two pain levels.

In summary, automated analysis of video recordings of mouse behavior using both Facial Grimace and Body Position Scores provides a high throughput approach to quantify persistent pain in mice. This method built on established methods for detecting facial grimace (22) and combined this with body position changes using a single camera in the same chamber. Validation of this method using CGRP-induced migraine demonstrated the ability to detect persistent pain and further allowed for the defining of pain thresholds (levels 1 and 2).

Figure 5. Significant differences in Facial Grimace and Body Position scores distinguish two pain levels

Supplemental Figure 2. Percentage Change from Baseline for each measurement used to generate Facial Grimace and Body Position Scores for Saline- and CGRP-injected mice

Behavioral detection of sound-evoked pain

Having validated the above method to measure persistent pain from video recordings of mouse behavior, this approach was next used to detect changes in behavior elicited by prolonged sound exposure. Sensitivity to sound in mice has been detected primarily using behavior measurements

in response to sound onset (4, 14, 15), which is an effective way to monitor changes in sound sensitivity. To measure behavioral changes associated with pain caused by longer duration sound, the following experiments used facial grimace and body position measurements to quantify auditory pain and determine the level of sound that caused pain-related behaviors in mice.

Sound exposure elicits changes in individual measurements of facial grimace and body position

Mouse behavior was next tested during a sound exposure protocol with varying sound intensities using an assessment of facial grimace and body position parameters as described above. Individual facial grimace and body position parameters changed with increasing sound intensities (Figure 6). The experimental design is shown in Figure 6A. Video recordings started 8 minutes after the mice were placed in the acoustic chamber. The recording from 8-10 minutes was used as the baseline recording. Sound exposures of varied sound intensity levels in a set sequence were performed in 2-minute time-periods and were interleaved with 2-minute time-periods of no sound exposure ('no sound'). The highest sound exposure level (120 dB) was placed last in the sequence of exposures, assuming that lasting cochlear damage might occur.

Example diary plots of one mouse show individual measures of Ear Position and Relative Nose Tip Position in response to the sound exposure protocol (Figure 6B-C). Superimposed data from all mice (n = 17) visualize the results from Baseline, 70 dB SPL, and 110 dB SPL periods (Figure 6D-E, raw and labeled video examples shown in Supplemental Video 2 and 3). Median values along with appropriate statistical tests reveal differences compared to Baseline in multiple individual measurements of facial grimace and body position at many intensities of sound. Ear Ratio, Ear Position, Relative Nose Tip Position, Percent in Top, and Vertical Line Crosses were all significantly different at 110 dB SPL compared to Baseline (Figure 6F-G). Specifically, Ear Ratio (repeated measures mixed effects model of all groups, p = 0.0013, post-hoc comparisons using Tukey's correction, 70 dB compared with Baseline median difference = 0.0361 AU, p < 0.01, 90 dB compared with Baseline median difference = 0.0602 AU, p < 0.01, 100 dB compared with Baseline median difference = 0.0927 AU, p < 0.05, 110 dB compared with Baseline median difference = 0.0637 AU, p < 0.01, and 120 dB compared with Baseline median difference = 0.0642 AU, p < 0.01), Ear Position (repeated measures mixed effects model of all groups, p < 0.0001, post hoc comparisons using Tukey's correction, 110 dB compared with Baseline median difference =

8.0°, p < 0.001, 120 dB compared with Baseline mean difference = 10.6°, p < 0.001), Relative Nose Tip Position (repeated measures mixed effects model of all groups, p = 0.0016, post hoc comparisons using Tukey's correction, 90 dB compared with Baseline median difference = -87 pixels, p < 0.05, 100 dB compared with Baseline median difference = -106 pixels, p < 0.01, 110 dB compared with Baseline median difference = -148 pixel, p < 0.01, 120 dB compared with Baseline median difference = -114 pixels, p < 0.05), Percent in Top (repeated measures mixed effects model of all groups, p < 0.0001, post hoc comparisons using Tukey's correction, 80 dB compared with Baseline median difference = -21.4 percentage points, p < 0.05, 90 dB compared with Baseline median difference = -15.0 percentage points, p < 0.01, 100 dB compared with Baseline median difference = -24.1 percentage points, p < 0.01, 110 dB compared with Baseline median difference = -28.9 percentage points, p < 0.001, 120 dB compared with Baseline median difference = -29.4 percentage points, p < 0.01), and Vertical Line Crosses (repeated measures mixed effects model of all groups, p = 0.0078, post hoc comparisons using Tukey's correction, 70 dB compared with Baseline median difference = -23 incidences, p < 0.05, 100 dB compared with Baseline median difference = -28 incidences, p < 0.01, 110 dB compared with Baseline median difference = -22 incidences, p < 0.05, 120 dB compared with Baseline median difference = -27 incidences, p < 0.05) all showed significant differences at multiple intensities. Ear Tip Tilt, Eye Ratio, and Face Inclination were all significant in the mixed effects model but showed no significant post-hoc comparisons. Specifically, Ear Tip Tilt (repeated measures mixed effects model of all groups, p = 0.0148), Eye Ratio (repeated measures mixed effects model of all groups, p = 0.0354), and Face Inclination (repeated measures mixed effects model of all groups, p = 0.0221) showed significant group effects. Finally, Mouth and Snout Position measures had no significant group effects. Taken together, these data suggest that auditory pain may be preferentially detected by individual grimace measurements related to ears, and by changes in body position.

Figure 6. Sound interleaved with silence causes significant changes to facial grimace and position at higher sound levels.

Supplemental Video 2: Interleaved Sound Experiment Example

Supplemental Video 3: Labeled, Interleaved Sound Experiment Example

---

After sound exposures, mouse behavior returned to baseline during 'No Sound' periods except after exposure to sound at 120 dB SPL

The sound exposure protocol for individual mice included a sequence of multiple sound presentations with varying sound pressure levels, and the analysis of the gained data, as presented in Figure 6, relies on the assumption that mouse behavior returns to baseline following each sound presentation. To verify this assumption, 'No Sound' periods following each sound presentation were analyzed. In Figure 7, 'No Sound' periods are arranged in increasing order of the preceding sound exposure level, and not in the order as they were presented in the interleaved sound exposure protocol. Ear ratio showed a significant effect of group (Figure 7A; repeated measures mixed effects model, p = 0.0012), however no significant differences were detected in post hoc comparisons to Baseline in any 'No Sound' period. The remaining facial grimace parameters showed no significant changes following any sound intensity presented compared to Baseline. By contrast, for most body position measurements, the 'No Sound' period after 120 dB SPL sound exposure did show significant differences compared to Baseline (Figure 7B). Specifically, Relative Nose Tip Position (repeated measures mixed effects model, p = 0.0003, post-hoc comparisons using Tukey's correction, post-120 dB vs Baseline median difference = -207 pixels, p < 0.01), Percent in Top (repeated measures mixed effects model, p = 0.0013, post-hoc comparisons using Tukey's correction, post-120 dB compared with Baseline median difference = -32 percentage points, p < 0.001), and Vertical Line Crosses (repeated measures mixed effects model, p = 0.01, post-hoc comparisons using Tukey's correction, post-120 dB vs Baseline median difference = -33 incidences, p < 0.001) all showed significant changes during 'No Sound' following 120 dB SPL, while no significant differences were found in post-hoc comparisons for Face Inclination (repeated measures mixed effects model, p = 0.0161). In this current study, 120 dB SPL was presented last with the consideration that this level of sound might impact subsequent animal behavior, which was confirmed here for multiple individual body position measurements.

In Summary, these findings indicate that for the specific design of the sound protocol used here, after sound exposures mouse behavior returned to baseline during 'No Sound' periods, except after exposure to sound at 120 dB SPL.

Figure 7. Facial grimace and body position measurements return to baseline between all sound intervals except following 120 dB

____

Sound exposure above 100 dB SPL caused changes in behavior consistent with pain in mice

As presented for the migraine model (Figure 5), Facial Grimace and Body Position Scores for both 'Sound' and 'No Sound' were calculated as a sum of the changes from baseline in individual measurements (Supplemental Figure 3) (Figure 8). For comparison, thresholds for two levels of pain, Level 1 and Level 2, as established in the migraine model (Figure 5), were added to Figure 8. Both composite scores for 'Sound' were significantly different from baseline for sound presentation at 100 dB SPL and above. Additionally, median values of both composite scores at 100 dB SPL and above were found to surpass the Level 1 pain threshold of the migraine model (Figure 8A; dotted line). This validates that both stimuli, sound and migraine, elicit significant behavioral changes from Baseline, surpassing a threshold for pain to a similar extent. Specifically, significant differences were found for Grimace Scores (repeated measures mixed effects model, $p < 0.0001$, post-hoc comparisons using Tukey's correction, 100 dB vs Baseline median difference = -22.6 percentage points, $p < 0.01$, 110 dB vs Baseline median difference = -30.9 percentage points, $p < 0.0001$, and 120 dB vs Baseline median difference = -29.6 percentage points, $p < 0.001$) and Position Scores (repeated measures mixed effects model, $p = 0.0012$, post-hoc comparisons using Tukey's correction, 100 dB vs Baseline median difference = -165.1 percentage points, $p < 0.001$, 110 dB vs Baseline median difference = -141.9 percentage points, $p < 0.05$, 120 dB vs Baseline median difference = -98.6 percentage points, $p < 0.001$) at and above 100 dB SPL. These findings indicated that mice exposed to $2 - 20$ kHz sound for two minutes at intensities of 100 dB SPL and greater showed behavioral signatures of pain.

Composite scores for 'No Sound' had no significant effects from the mixed model analyses for facial grimace nor body position (Figure 8B). It is noteworthy that during the period of no sound following 120 dB SPL, the Body Position Score surpassed Level 2 threshold for pain found in the migraine model (Figure 8B; dashed line). This change in behavior following 120 dB SPL is consistent with statistically significant changes observed in individual measurements described above (Figure 7 and Supplemental Figure 3).

____

Figure 8. Sound above 100 dB evokes behavioral changes associated with pain

Supplemental Figure 3. Percentage change from Baseline for each measurement used to generate Facial Grimace and Body Position Scores for Sound- and No-Sound- exposed mice.

Cochlear mechano-transduction is required to generate sound-evoked pain behavior

Pain caused by loud sounds could result from stress forces on auditory structures peripheral to the cochlea, such as sound pressure activating stretch receptors in the tympanic membrane or middle ear (43). To test whether normal cochlear function is necessary for sound-evoked pain, Mice that lack mechano-transduction due to the loss the TMIE protein needed for proper function of the mechano-transduction channel in all cochlear hair cells (TMIE-knockout mice, $Tmie^{-/-}$) and littermate heterozygous or wild-type controls ($Tmie^{CTL}$) were presented with the same interleaved sound protocol described above and tested for pain behavior (27–30). Importantly, both heterozygous and wild-type mice show normal cochlear mechano-transduction and thus can serve as valid controls (44). These mice are maintained on a C57BL/6J background.

Mice lacking Tmie largely showed no changes in individual behavioral measurements in response to sound exposure

Individual measurements of facial grimace and body position in $Tmie^{CTL}$ mice show similar trends as the above C57BL/6J mice (Figure 9). Example diary plots from an individual $Tmie^{CTL}$ mouse demonstrate changes in Ear Position and Relative Nose Tip Position during sound exposures (Figure 9A-B). For data from all mice combined, for facial grimace measurements, Ear Ratio and Ear Position were significantly different from baseline at multiple sound levels (Figure 9C). Specifically, significant increases were found in Ear Ratio (repeated measures mixed effects model, p = 0.0009, post-hoc comparisons using Tukey's correction, 90 dB compared with Baseline median difference = 0.127 AU, p < 0.001, 100 dB compared with Baseline median difference = 0.173 AU, p < 0.01, and 120 dB compared with Baseline median difference = 0.117 AU, p < 0.05) and in Ear Position (repeated measures mixed effects model, p = 0.0245, post-hoc comparisons using Tukey's correction, 110 dB compared with Baseline median difference = 3.1°, p < 0.05, and 120 dB compared with Baseline median difference = 16.5°, p < 0.01). Eye ratio showed a significant group effect but no post-hoc tests were significantly different from baseline (repeated measures mixed effects model, p = 0.0293). For body position measurements, significant reductions were found in Relative Nose Tip Position, Percent in Top, Vertical Line Crosses (Figure

9D). Specifically, Relative Nose Tip Position (repeated measures mixed effects model, $p = 0.014$, post-hoc comparisons using Tukey's correction, 110 dB compared with Baseline median difference = -129 pixels, $p < 0.01$), Percent in Top (repeated measures mixed effects model, $p = 0.0124$, post-hoc comparisons using Tukey's correction, 110 dB compared with Baseline median difference = -15.1%, $p < 0.05$), Vertical Line Crosses (repeated measures mixed effects model, $p = 0.0078$, post-hoc comparisons using Tukey's correction, 90 dB compared with Baseline median difference = -51 incidences, $p < 0.05$, 100 dB compared with Baseline median difference = -53 incidences, $p < 0.01$, 110 dB compared with Baseline median difference = -60 incidences, $p < 0.01$) all showed significant decreases at 110 dB SPL (e.g., see Supplemental Video 4).

Without functional cochlear mechano-transduction, in $Tmie^{-/-}$ mice, most individual facial grimace measurements were not significantly different from baseline in any condition, however, there was one body position measurements that did show significant differences compared with baseline (detailed below). Example diary plots of Ear Position and Relative Nose Tip Position from an individual mouse show no obvious behavioral changes during any sound exposure (Figure 9E-F). For data from all mice combined, individual facial grimace measurements did not show any changes (Figure 9G). However, of the four body position measurements, only Vertical Line Crosses were significantly reduced compared with baseline at 110 dB SPL (Figure 9H; repeated measures mixed effects model, $p = 0.0005$, post-hoc comparisons using Tukey's correction, 110 dB compared with Baseline mean difference = -68 incidences, $p < 0.01$). It is unclear what could cause the decreased turning observed here, however it is noteworthy that Tmie$^{-/-}$ mice have altered vestibular function that causes repeated turning behavior, which caused a two-fold increase in Vertical Line Crosses at baseline compared with $Tmie^{CTL}$ mice (see Figure 9D and H). Relative Nose Tip Position also showed a significant group effect (repeated measures mixed effects model, $p = 0.04$), but the direction of this effect was towards increased position that is not suggestive of pain.

Figure 9. Mice deficient in mechanotransduction do not respond to increased sound levels with significant changes to facial grimace or position associated with pain

Supplemental Figure 4. Percentage change from Baseline for each measurement used to generate Facial Grimace and Body Position Scores for $Tmie^{CTL}$ and $Tmie^{-/-}$ sound-exposed mice

Supplemental Video 4: TMIE control example video

Combined Facial Grimace and Body Position Scores of *Tmie^CTL* and *Tmie^-/-* mice

Individual measurements of both facial grimace and body position were converted into percentage change in baseline (Supplemental Figure 4) and summed to generate composite scores (Figure 10). Facial Grimace and Body Position Scores revealed sound-evoked pain in *Tmie^CTL* mice emerging at 100 dB SPL and greater sound intensities (Figure 10A) replicating the findings from C57BL/6J detailed above. Specifically, Facial Grimace Score and Body Position Score both showed significant changes at 100 and 110 dB SPL consistent with normal hearing C57BL/6J mice presented above (Figure 5) (Facial Grimace Score: repeated measures mixed effects model, p = 0.0004, post-hoc comparisons using Tukey's correction, 90 dB compared with Baseline median difference = - 26.0%, $p < 0.05$, 100 dB compared with Baseline median difference = -53.7%, $p < 0.01$, 120 dB compared with Baseline median difference = -47.9%, $p < 0.001$; Body Position Score: repeated measures mixed effects model, p = 0.0031, post-hoc comparisons using Tukey's correction, 100 dB compared with Baseline median difference = -154.8%, $p < 0.001$, 110 dB compared with Baseline median difference = -140%, $p < 0.0001$).

In *Tmie^-/-* mice, no significant differences were observed for Facial Grimace or Body Positions Scores at any sound intensity (Figure 10B; Supplemental Video 5) suggesting that normal cochlear mechano-transduction is necessary for inducing auditory pain. These results suggest that sound pressure causing the activation of stretch receptors in peripheral auditory structures are not the source of auditory pain as measured and defined here (45–47). However, descending motor fibers into peripheral structures (such as cranial nerve V innervation of the tensor tympani) could become hyperactive due to prior sound exposures that may contribute to sound-evoked pain (45, 48, 49), which would all require cochlear function to generate their subsequent pathophysiological responses, cannot be ruled out.

Figure 10. Mice deficient in mechanotransduction do not show pain-associated behavioral changes when exposed to loud sounds like wildtype mice.
Supplemental Video 5: TMIE knockout example video

In Summary, there study here demonstrates the feasibility of using a single camera angle to measure facial grimace and additional positional information from video recordings of mice to detect behavioral changes associated with persistent pain. By validating this approach with a known painful stimulus (CGRP-induced migraine) and then applying it to an experimental condition (loud sound), this approach allowed for the investigation of auditory pain in mice. Future studies could apply this method to other ongoing, persistent pain conditions or to investigate further the sources and mechanisms responsible for pain caused by sound exposure.

Discussion (word count 1948)

Difficulty measuring when loud sounds become painful in animal models, such as mice, presents a challenge to understanding how auditory pain is generated and processed. To overcome this, sound-evoked pain was quantified using a deep neural network that was trained to place points on the face and body of mice that were video recorded while being presented with different intensities of sound. Using this method, an inventory capable of detecting affective pain in mice was created by analyzing pain-related changes in facial grimace (20–22, 31, 32) and animal position (50–52). Pain thresholds were determined by analyzing mice injected with calcitonin gene-related peptide (CGRP) to induce migraine—a well characterized stimulus that results in a predictable time course during which mice experience spontaneous, ongoing pain (23). Persistent pain from migraine allowed for validation of this new method before applying it to auditory pain, another form of spontaneous, ongoing pain during a stimulus. Sound presented at intensities of 100 dB and above were found to result in pain-related behavioral changes that appeared in aggregated scores of both facial grimace and body position but were not present in mice lacking cochlear mechanotransduction. Taken together, these findings suggest that persistent painful states, such as sound-evoked auditory pain, can be reproducibly measured in mice, and that sound transduction in the inner ear is necessary for sound-evoked pain.

Integrating Facial Grimace and Body Position to Measure Mouse Affective Pain

Mouse facial grimace and position information were combined to create an inventory-like system capable of detecting subtle changes in mouse behavior associated with spontaneous, ongoing pain. This method involves pose estimation by a deep neural network that places key points on the face and body of freely moving mice inside clear chambers (Figure 1). Existing affective pain

measurements focus on the face or movement of the animal and require cameras aimed at the face and overhead, respectively (21, 53, 54). PainFace can capture statistically significant changes in affective pain behavior in mice of many coat colors without a need for retraining the algorithm (21). Moseq captures both movement and complex behavior data in the form of behavioral syllables and can capture behaviors on a subsecond scale (53, 54). The integration of two behavioral measures expands on previous pain assays by automating facial grimace along with measuring multiple parameters of mouse body position using one camera angle. Facial Grimace and Body Position data are processed in a high throughput, non-biased manner from thousands of frames from hours of video recording. By combining facial grimace and body position measurements, two pain levels were discerned in a known pain model (CGRP-induced migraine). Thresholds for pain were determined based on the change in behavior compared to baseline for CGRP-injected mice. CGRP injection resulted in a peak change in behavior from 27-32 minutes after injection (Figures 2-5; Supplemental Figure 2,3). The behavior during the non-peak time of the recording after CGRP-injection was also significantly changed from both baseline and the peak time-period for both behavioral scores. Thus, two thresholds were created, the Level 1 threshold was based on the median of the Non-Peak behavior scores and the Level 2 threshold based on the median of the Peak behavior scores. The increase in number of pain levels differentiated by this method provides an advancement for affective pain measurement.

In this first publication of the combined facial grimace and body position method, all measurements are equally weighted when added together to create the Facial Grimace and Body Position scores in an effort to replicate similar summed scores of affective pain (21). However, the scale of each measurement affects the maximal percentage change from baseline value that contributes to the summed Facial Grimace or Body Position Score. Furthermore, several measurements such as Ear Ratio, and Ear Tip Tilt do not show large changes in units but instead may be best represented by a change in variation at different points during the recording. These will be considered in subsequent iterations of the method. It is important to note that the score method is designed like an inventory that can incorporate additional measurements and even categories of behavior to further refine the ability to detect affective pain, such as *en face* grimace measurements and even a third score encompassing complex behaviors such as grooming, rearing, exploring, circling, defecation, urination, piloerection, and respiration rate, which are known indicators of persistent painful states in animals (26, 55). While recording the animal's behavior

changes to complex behaviors such as grooming, defecation and respiratory rate were observed. The expansion of the combined behavioral method to include new measurements will also be accompanied by a robust analysis of the measurements most indicative of a painful state further refining and simplifying the method for others to use. With the addition of new measurements, it is possible that this approach of scoring multiple behaviors at sub-second resolution will be able to generate a graded scale of affective pain measurement.

Existing methods for detecting auditory-evoked behavior

Several methods are used to measure sound-evoked behavior in rodents. These methods fall into two categories: 1) measurements of reflexive responses to sound 2) measurements of aversion to sound. Acoustic startle responses and sub-second facial responses model loudness perception (12, 56). Conditioned aversion assays measure the preference of a mouse for a well-lit chamber over a dark chamber with noise going against the animal's instinct (56, 57). Thus, animal models looking at the human condition of loudness hyperacusis do exist. The results of reaction time and conditioned aversion assays have been interpreted as aversion to sound, but the conclusion that is due to affective pain is unclear. Thus, the subset of hyperacusis known as pain hyperacusis currently has no animal model that measures affective pain. A mouse model that can measure affective pain to sound is needed to develop a mouse model of pain hyperacusis.

Sound presentation evokes pain behaviors at 100 dB SPL and above

It is known that humans can experience pain to loud sounds at 120 dB and above (58). With that in mind, suprathreshold levels of sound from 70 dB to 120 dB were used to probe whether rodents would display behavioral changes associated with pain—a painful affect—at any of these levels of sound. Periods of silence ('no sound') were interwoven to prevent any adaptation effect by continuous sound presentation or continuous increases to sound level. These 'No Sound' periods also serve as a control that the behavioral changes were restricted to sound presentation that were reversible. Both Facial Grimace and Body Position scores were significantly changed from baseline starting at 100 dB SPL. When comparing facial grimace and body position scores due to sound presentation to those after CGRP injection, 50% of mice display a similar level of affective pain above 100 dB of sound. Observations included that the behavior of the mice changed at 120

dB such that some movements occurred suddenly between bouts of hunching. Thus, additional analysis of this behavior is warranted as it may be due to escape behavior or fear. Furthermore, the silent period after 120 dB was the only silent period where the mice continued to have changes to their movement in both the interleaved and continuous sound presentation experiments (Figure 7, 8, Supplemental Figure 4). This effect persisted for approximately 30 sec into the silent period. It is known that loud noises can cause changes to auditory perception such as short duration tinnitus (59). Further study will elucidate the mechanism causing this persistent effect.

Cochlear mechano-transduction in required for sound-evoked pain behaviors

One question that the authors wanted to address is what peripheral auditory structures were necessary for an animal to experience auditory pain. The middle ear has stretch receptors that could be triggered at high pressure levels to induce pain due to sound or changes in pressure on the tympanic membrane (45–47, 60). The brain is known to be able to perceive pain in the absence of a sensory structure during phantom limb pain, for instance (61). Therefore, a mouse model was chosen with normal development of outer and middle ear, but a dysfunctional inner ear, the $Tmie^{-/-}$ mouse (44). The littermate $Tmie^{CTL}$ animals exhibited behavioral changes associated with sound presentation similar to wildtype animals, suggesting pain perception during loud sound exposure. However, $Tmie^{-/-}$ animals showed no behavioral changes at any sound exposure level that reached the pain threshold for either the facial grimace or body position scores. Thus, cochlear function is needed to detect sound for a normally hearing animal to display a behavioral affect of pain to loud sound. It is still possible that in animals with disordered hearing or changes to the middle ear (such as otitis media), the middle ear could play a role in auditory pain perception. As in tinnitus and phantom limb syndrome, central control of auditory pain may also play a role in disordered auditory pain such as pain hyperacusis (61, 62). However, detection by the peripheral auditory system plays a fundamental role in the generation of auditory pain to loud sound in a normal mouse.

Methods

Animals

Six-week-old C57BL6/J mice of both sexes were used. Both sexes were tested separately. At the time of testing, animals were group-housed in a low noise vivarium. The housing room was maintained at 12:12 h light: dark cycle (7:00 to 19:00). Up to five mice were housed per one filter top shoebox cage ($30 \times 19 \times 13 \text{ cm}^3$). All procedures were approved by the Johns Hopkins University Animal Care and Use Committee and follow the National Institutes of Health ARRIVE Guidelines.

Behavioral testing setup design

Individual polycarbonate containers (testing chamber; interior dimensions: 9 x 5x 5 cm) were fabricated onsite and placed in an acoustic chamber prior to testing. The acoustic chamber was lined with LED lights for filming. At all times, the mice were awake and freely moving. Mice were oriented to maintain a profile position via scent cues from air holes positioned at one end of their polycarbonate containers. When multiple mice were recorded simultaneously, two mice of the same sex were placed individually into two chambers oriented to face opposite each other while maintaining a profile position via scent cues from air holes positioned at opposite ends of each container. The mice were prevented from seeing each other by adding a white opaque barrier between the containers. A GoPro Hero 11 Black™ (San Mateo, CA) was centered to capture mice during the protocol. Overhead speakers were fitted into the sound chamber to allow for sound exposure.

Video Editing

Sound exposure videos were approximately 26 minutes in length and were filmed in 4K resolution at 30 frames per second. Videos were cropped to approximately 1000 x 1500 pixels (height vs width) to analyze one of two mice at a time when multiple mice were recorded simultaneously. The recordings from the CGRP- and saline-injected mice yielded one 10-minute and one 20-minute video filmed in 4K resolution at 30 frames per second. Each video was then cropped using iMovie to focus on a single mouse at a time for the most accurate labeling for each condition. Time segments (epochs) of video were cropped to the desired length to cover a given condition (e.g., CGRP peak epoch, 110 dB SPL, baseline, etc.)

Pose Estimation using DeepLabCut

The deep neural network was trained to place thirteen markers on the face in each video frame where the face is in profile using DeepLabCut (DLC; (33, 34)). The thirteen markers were labeled inner eye, outer eye, eye up, eye down, nose tip, nose bottom, mouth, ear canal, ear tip, ear center, ear medial, ear lateral, and neck (Figure 1; (22)). DLC was installed on a Windows OS computer with an Intel® Core™ i5-7600K CPU @ 3.80 GHz processor and an Nvidia GeForce GTX 1070 graphics card. ResNet-50-based neural network with default parameters was run for up to 800,000 iterations of training on 816 annotated frames from all cohorts of animals. The dataset was split as 95% for training and 5% for testing. After training, annotated images were visually inspected to ensure the markers were placed on the appropriate key points of the body and face. Data for new videos analyzed were filtered based off a greater than 0.95 likelihood (p-cutoff $>= 0.95$). A lower limit of at least 30 analyzed frames per video analyzed was used to consider that epoch valid for statistical comparisons, and all recorded time periods fit this criterion.

Defining Face and Position Parameters

A representative annotated frame showing DLC-generated pose estimated points is shown in Figure 1. Lines between points (skeletons) were created automatically in DLC (Figure 1A). The skeletons used were nose tip to nose bottom, mouth to inner eye, nose bottom to inner eye, inner eye to outer eye, eye up to eye down, inner eye to ear canal, ear canal to ear tip, and medial ear to lateral ear. These skeletons are used to define eye ratio, ear ratio, ear angle, ear position, snout position, and mouth position. Representative images show changes in these parameters between control and pain conditions (Figure 1B). Angles between certain skeletons and ratios of certain skeletons were used to generate measurements of facial parameters, as follows: Ear Ratio was calculated as the ratio of the length of the lines in pixels from Ear Med to Ear Lat and Ear Canal to Ear Tip indicated by white lines (Figure 1C). Ear Position was calculated as the obtuse angle formed by lines connecting Inner Eye, Ear Canal, and Ear Tip with Ear Canal as the vertex as indicated in white (Figure 1D). Ear Tip Tilt was calculated as the angle formed by lines connecting Ear Canal to Ear Center to Ear Tip with Ear Center as the vertex as indicated in white (Figure 1E). Eye Ratio was calculated as the ratio of the length of the lines in pixels from Eye Up to Eye Down and Inner Eye to Outer Eye indicated by white lines (Figure 1F). Snout Position was calculated as

the acute angle formed by lines connecting Nose Tip, Inner Eye, and Nose Bottom with Inner Eye as the vertex as indicated in white (Figure 1G). Mouth Position was calculated as the acute angle formed by lines connecting Nose Bottom, Inner Eye, and Mouth with Inner Eye as the vertex as indicated in white (Figure 1H).

The Relative Nose Tip Position was the distance in pixels that the nose tip point was below the (0,0) coordinate in the top left most corner of the chamber, with increasingly negative values representing the mouse's nose tip at a low place in the chamber (Figure 3A-B). The nose tip pose estimated point was further used to calculate the percentage of analyzed frames where this point is above a horizontal line drawn at -450 pixels (Figure 3C; corresponding to the back of the mouse when all four paws are planted on the chamber floor, 'Percentage of Nose Tip in Top 1/3 of Chamber'), and to count the incidences when the nose tip crosses a vertical line drawn at 900 pixels (Figure 3D; deliberately set off-center to ensure full turns of the mouse were required to count as an incidence of crossing the vertical line, 'Vertical Line Crosses of Nose Tip') . These measurements were calculated as the percentage of frames above horizontal line and total number of vertical line crosses, respectively. The horizontal line in Percentage of Nose Tip in Top 1/3 of Chamber was set at -450 because the y-axis pixel count varied from ~900-1000 based on cropping of the video. The walls of the chamber (top and bottom, including space outside the chamber walls) accounted for 150-300 pixels of that space leaving the interior of the chamber at an average of 900 pixels. Therefore, 1/3 of the chamber is 300, but 150 was added to account for the top chamber wall thus -450. The Percentage of Nose Tip in Top 1/3 of Chamber provides a gross approximation of decreased rearing, exploration of the top portion of the chamber, and climbing behaviors. While Vertical Line Crosses of Nose Tip capture how often the mouse turns inside the chamber, which is influenced by exploration behavior and total locomotion. Finally, Face Inclination was calculated as the percentage of analyzed frames where the nose tip was below the neck (Figure 3E). Mice in pain tend to hunch forward causing the nose tip to be below the neck point, therefore this measure of face inclination would increase the more the mouse was hunched forward.

CGRP or Saline Injections
Mice were placed individually into the testing chamber and allowed ten minutes of acclimation to the container. Ten minutes of video was subsequently recorded to measure baseline behavior for each mouse. Mice were then removed from the container and injected with either calcitonin gene-

related peptide (CGRP; 100 μg/mL in saline, i.p.) or saline (vehicle) and immediately returned to the container. Since this dose of CGRP induces a migraine that peaks 30 minutes after injection (23), video was recorded from 20 – 40 minutes post injection. In total, 30 minutes of video were recorded for each mouse and was subsequently segmented for analysis as described below.

Sound Exposure Protocol

Mice received an acclimation period of 30 minutes to their containers the day before the sound exposure protocol. Calibrations within the acoustic chamber were completed each day before noise exposure took place to ensure exposures were accurate and consistent. The two readings checked were 90dB SPL and 120dB SPL. The calibration consisted of taking readings from the corner of each container and the center of the acoustic chamber using a sound level meter model LxT2 (Larson Davis, Inc., Provo, UT). Mice were placed into the same individual polycarbonate containers (described above) within an acoustic chamber fitted with overhead speakers. The mice were unable to look at each other throughout the exposure to eliminate visual cues from the other mouse. The video was centered in the middle capturing both mice during sound exposures. The sound exposure protocol began with ten minutes without sound played from the speakers, then sound was presented in an interleaved protocol ranging from 70 – 120 dB SPL of broadband (2-20 kHz). The sound was played for two minutes followed by two minutes of no sound after each sound presentation. The two-minute period without sound (from 8-10 minutes in each video) was used as the control for each mouse. Sound presentation was interleaved in the following order: 90, 110, 70, 100, 80, 120 dB SPL.

Data Analysis and Statistics

Filtered CSV files were output once DeepLabCut finished analyzing each video. These files were converted into xlsx files to be quantified. Pairwise comparisons, such as Baseline compared with All – CGRP, were tested using paired, two-tailed, student t-tests. Statistical differences for groups with repeated measures were assessed using repeated measures ANOVA for balanced groups and repeated measures mixed effects model for subgroups tested in longitudinal designs (e.g., CGRP baseline and subgroups within migraine, and sound exposure epochs) and post-hoc Tukey's correction for multiple comparisons.

Acknowledgements

This work was supported by the following funding: a Hearing Health Foundation Emerging Research grant to M.B.W., a Johns Hopkins University internal Blaustein Pain Research Fund Grant to B.J.S. and M.B.W., the John Mitchell Jr. Fund to B.J.S., the Geraldine Dietz Fox Endowed Research Fund to E.G. and a National Institute on Deafness and other Communication Disorders (NIDCD) grant 5R01DC016559 to E.G. The David M. Rubenstein Precision Medicine Center of Excellence in Hearing Fund project 2a to EG; the David M. Rubenstein Professorship to EG. We would also like to thank Kevin Psoter, MPA, PhD for consultations of his biostatistical expertise. We thank Paul Fuchs, PhD, for comments on the manuscript.

Figures

Figure 1: Labels used to train a deep neural network for facial grimace measurements

A) Left, a schematic of the inside of the acoustic booth used for facial grimace video recording. Right, a diagram of the custom-built enclosure from the view of the camera. The ovals on the left wall and top indicate scent holes. Figure not to scale. **B)** Labeled frames from the same video of a mouse in a control condition (left) and a painful condition (right). **C)** A labeled frame of a mouse face showing the 13 markers on the face (numbered, colored dots). The black lines are skeletons generated by the model. Colored dot 1- Nose Tip (upper tip of nose); 2 – Nose Bot (bottom of nose); 3 – mouth; 4 – inner eye (anterior edge of eye); 5 – Outer Eye (caudal edge of eye); 6 – Eye Up (highest point of eye lid on dorsal edge); 7 – Eye Down (lowest point of eyelid on ventral edge); 8 – Ear canal (edge of hair and ear canal); 9 – Ear Tip (edge of pinna furthest from the ear canal); 10 – Ear center (center of ear pinna between points 9, 10, 11 and 12); 11 – Ear Med (halfway up the pinna on the medial side); 12 – Ear Lat (halfway up the pinna on the lateral side of the ear); 13 – Neck (point where the head connects to the body). **D)** Ear Ratio was calculated as the ratio of the length of the lines in pixels from Ear Med to Ear Lat (Point 11 to 12) and Ear Canal to Ear Tip (Point 8 to 9) indicated by white lines. **E)** Ear Position was calculated as the obtuse angle formed by lines connecting Inner Eye, Ear Canal, and Ear Tip with Ear Canal as the vertex as indicated in white. **F)** Ear Tip Tilt was calculated as the angle formed by lines connecting Ear Canal to Ear Center to Ear Tip with Ear Center as the vertex as indicated in white. **G)** Eye Ratio was calculated as the ratio of the length of the lines in pixels from Eye Up to Eye Down (Point 6 to7) and Inner Eye to Outer Eye indicated by white lines (Point 4 to 5). **H)** Snout Position was calculated as the acute angle formed by lines connecting Nose Tip, Inner Eye, and Nose Bottom with Inner Eye as the vertex as indicated in white. **I)** Mouth Position was calculated as the acute angle formed by lines connecting Nose Bottom, Inner Eye, and Mouth with Inner Eye as the vertex as indicated in white.

Figure 2: Changes measured in facial grimace during CGRP-induced migraine

A) Timeline of CGRP injection experiment. Animals were filmed for a 10-minute baseline period. CGRP was injected based on body weight. After a 10-minute period, mice were filmed for 20 minutes surrounding the peak of action CGRP which occurs 30 minutes after injection. **B)** Example of the Ear Position data from the baseline and CGRP time periods of one mouse (N=1). Each dot is the measurement from a single frame of video. **C)** Averaged data for each frame of video for Ear Position from 12 mice (N=12) during Baseline and the entire CGRP Recording. The right graph shows the average Ear Position for 12 mice during the peak time period as indicated in A and the box in the middle graph. The black horizontal line is the average of the baseline time-period as a reference in all graphs. (**D-I**) The following graphs show the averaged data for each facial parameter indicated. Each dot is the averaged data for the indicated time-period for 1 mouse. Baseline is the averaged data for each mouse for all frames from the Baseline video (10 minutes). 'All' is the averaged data for each mouse for all frames after CGRP injection. 'Peak' is the averaged data from minutes 27-31 and after CGRP injection. 'Non-Peak' is the averaged data from time outside of the peak time period (minutes 20-26 and minutes 32 to 40). Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. **D)** Ear Ratio. **E)** Ear Position. **F)** Ear Tip Tilt. **G)** Eye Ratio, Baseline compared to All: paired, two-tailed t-test, **, p=0.0012; repeated measures mixed effects model, **, p=0.0025; post hoc comparisons using Tukey's correction, Baseline compared to Peak, **, p= 0.0215; Baseline compared to Non-Peak, *, p= 0.0111. **H)** Mouth Position, repeated measures mixed effects model, ****, p=<0.0001; post hoc comparisons using Tukey's correction, Baseline compared to Peak, **, p=0.0015; Peak compared to Non-peak, p=0.0026. **I)** Snout Position. If no statistics noted, data was not significantly different.

Figure 3: Analysis of Mouse Body Position from Tracking of Nose Tip Coordinates

A) Example of a full frame of video of a control mouse with x and y axes showing pixel values. The nose tip is indicated with a navy-blue dot. The y-axis pixel value of the nose tip was recorded for each frame and averaged over discrete time periods to create the measurement named Relative Nose Tip Position as a proxy for exploring behavior. **B)** Example of a full frame of video of a control mouse. The white dashed line at -450 pixels on the y-axis indicates the boundary of the top 1/3 of the chamber. The average number of frames in a time-period where the nose tip (navy-blue dot) was above this line based in the y-axis was calculated as the Nose Tip Top % measurement as a proxy for rearing behavior. **C)** Example of a full frame of video of a control mouse. The white dashed line at 900 pixels on the x-axis indicates a boundary which was used to count the number of times the nose tip (navy-blue dot) crossed this line in a time-period. This was used to calculate the Vertical Line Crosses measurement as a proxy for turning behavior. The line is intentionally offset to capture the full turning of the mouse from one side of the cage to the other instead of rearing straight backwards as during grooming. **D)** Example of a mouse face labeled from the DNN training. The y-axis value of the Nose Tip was subtracted from the y-axis value of the Neck point (Neck – Nose Tip; indicated by the white line). The Face Inclination measurement was then calculated as the percentage of frames during a time-period

that the Neck – Nose Tip value was negative meaning the Nose Tip was below the Neck value on the Y-axis as a proxy for hunched body position.

Figure 4: Movement decreases during CGRP-induced painful migraine as calculated by measures of mouse body position

A) Example of the Relative Nose Tip Position data from the baseline and CGRP time periods of one mouse (N=1). Each dot is the measurement of Y-axis Nose Tip Position from a single frame of video. The line at Y-axis -450 was used to calculate the Percent in Top measurement. **B)** Averaged data for each frame of video for Relative Nose Tip Position from 12 mice (N=12) during Baseline and the entire CGRP Recording. The right graph shows the average Relative Nose Tip Position for 12 mice during the peak time period as indicated in Fig 2A and the box in the middle graph. The black horizontal line is the average of the baseline time-period as a reference in all graphs. **C-F)** The following graphs show the averaged data for each body position measurement indicated. Each dot is the averaged data for 1 mouse. Baseline is the averaged data for each mouse for all frames from the Baseline video. 'All' is the averaged data for each mouse for all frames after CGRP injection. 'Peak' is the averaged data from minutes 27-31 and after CGRP injection. 'Non-Peak' is the averaged data from time outside of the peak time period (minutes 20-26 and minutes 32 to 40). Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. **C)** Rel. Nose Tip Position. **D)** Percent in Top: Baseline compared to Peak and Non-

Peak, repeated measures one-way ANOVA, p=0.0221; post hoc comparisons using Tukey's correction, Baseline compared to Non-Peak, *, p= 0.0266. **E**) Vertical Line Crosses: The absolute number of crosses was normalized to a 5-minute time-period to compare baseline (10 min) and All (20 min) to Peak (5 min) and Non-Peak (15 min). Baseline compared to All: paired, two-tailed t-test, *, p=0.0241; Baseline compared to Peak and Non-Peak, repeated measures mixed effects model, **, p=0.0004; post hoc comparisons using Tukey's correction, Baseline compared to Peak, ***, 0.0025; post hoc comparisons using Tukey's correction, Peak compared to Non-Peak, ***, 0.0002. **F**) Face Inclination: Baseline compared to All: paired, two-tailed t-test, *, p=0.0236; Baseline compared to Peak and Non-Peak, repeated measures mixed effects model, *, p=0.03; post hoc comparisons using Tukey's correction, Baseline compared to Peak, *, p=0.0293. If no statistics noted, data was not significantly different.

Figure 5: Significant differences in Facial Grimace and Body Position scores distinguish two pain levels

Facial Grimace Scores were calculated as the added percentage change from baseline for all facial grimace measurements for each mouse. When a facial grimace parameter resulted in a positive change from baseline (Ear Ratio, Ear Position), it was inverted so the percentage change from baseline associated with pain was a negative value. Body Position Scores were calculated as the added percentage change from baseline for all position measurements for each mouse. When a body position measurement resulted in a positive change from baseline (Rel. Nose Tip Position, Face Inclination), it was inverted so the percentage change from baseline associated with pain was a negative value as with the Facial Grimace Score. **A**) Facial Grimace (left) and Body Position (Right) Scores for mice injected with CGRP (N=12). Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. CGRP Facial Grimace Score: Baseline compared to All, paired, two-tailed t-test, **, p=0.0022; Baseline compared to Peak, paired, two-tailed t-test, **, p=0.001; Baseline compared to Non-Peak, paired, two-tailed t-test, *, p=0.0063. Peak compared to Non-Peak, paired, two-tailed, t-test, *., p=0.0263. The dotted line is defined as the score associated with Pain Level 1 and begins at the

median of the Non-Peak data (median = -19.04 percent change from baseline). The dashed line is defined as the score associated with Pain Level 2 and begins at the mean of the Peak data (median = -40.62 percent change from baseline). CGRP Body Position Score (Right); Baseline compared to All, paired, two-tailed t-test,***, p=0.0003; Baseline compared to Peak, paired, two-tailed t-test, ***, p=0.002; Baseline compared to Non-Peak, paired, two-tailed t-test, *, p=0.001. Peak compared to Non-Peak, paired, two-tailed t-test,**, p=0.0369. Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. The dotted line is defined as the score associated with Pain Level 1 and begins at the median of the Non-Peak data (median = -132.4 percent change from baseline). The dashed line is defined as the score associated with Pain Level 2 and begins at the mean of the Peak data (median = -179.5 percent change from baseline). For a mouse to be experiencing pain, the scores must be lower than -19.04 for Facial Grimace and -132.4 for Position. **B**) Facial Grimace (Left) and Body Position (Right) Scores for mice injected with Saline (N=16). Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. Statistical tests were not significant. Saline Facial Grimace Score: Baseline compared to All, paired, two-tailed t-test, p=0.8211. Baseline compared to Peak and Non-Peak, repeated measures mixed model, p=0.5096. Saline Body Position Score: Baseline compared to All, paired, two-tailed t-test, p=0.1673. Baseline compared to Peak and Non-Peak, repeated measures mixed model, p=0.2735. **C**) The percentage change from baseline for each facial grimace measurement was added together and averaged for each minute of the CGRP recording. Each dot corresponds to 1 animal. The dashed line is the mean of all 20 minutes for all mice (-27.17 % change from baseline). Minutes where the median falls below the dashed line are colored gray. Minutes 27-31 show the longest string of median scores below the dashed line. **D**) The percentage change from baseline for each body position measurement was added together and averaged for each minute of the CGRP recording. Each dot corresponds to 1 animal. The dashed line is the median of all 20 minutes for all mice (-145.7 % change from baseline). Minutes where the median falls below the dashed line are colored gray. Minutes 27-34 have the longest string of lowest median scores.

Figure 6: Sound interleaved with silence causes significant changes to facial grimace and position at higher sound levels.

A) Timeline of interleaved sound level experiments. Recording started at 8 minutes after the mice were placed in the acoustic chamber. The recording from 8-10 minutes was used as the baseline recording. Sound exposure was initiated in 2-minute time-periods separated by 2 minutes of silence. Sound was presented in out of decibel-level order as indicated by color and

label. **B**) Example of analyzed frames of Ear Position measurements for one mouse for the entirety of one recording (N=1). The black vertical lines indicate when the sound levels were manually changed on or off. **C**) Example of analyzed frames of Relative Nose Tip Position measurements for one mouse for the entirety of one recording (N=1). The black vertical lines indicate when the sound levels were manually changed on or off. **D**) Averaged Ear Position measurements of analyzed frames from 12 mice (N=12) for Baseline, 70 dB, and 110 dB time periods. The time in minutes is the average on and off time periods for Baseline, 70 dB and 110 dB for each recording. The 70 dB recording occurred after the 110 dB recording, but the graphs have been reordered according to decibel level. Horizontal black lines indicate the average for the baseline graph. **E**) Averaged Relative Nose Tip Position measurements of analyzed frames from 12 mice (N=12) for Baseline, 70 dB, and 110 dB time periods. The time in minutes is the average on and off time periods for Baseline, 70 dB and 110 dB for each recording. The 70 dB recording occurred after the 110 dB recording, but the graphs have been reordered according to decibel level. Horizontal black lines indicate the average for the baseline graph. **F**) The following graphs show the averaged data for each facial parameter indicated. Each dot is the averaged data for the indicated 2-minute time-period for 1 mouse. The sound levels have been reordered from lowest to most intense sound level. Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. Ear Ratio: baseline compared to each sound level, repeated measures mixed model, p=0.0013; post hoc comparison using Tukey's correction, baseline compared to 70 dB,**, p=0.0068, baseline compared to 90 dB, **, p=0.0055, baseline compared to 100 dB, *, p=0.0135, baseline compared to 110 dB, **, p=0.0079, baseline compared to 120 dB, **, p=0.0015. Ear Position: baseline compared to each sound level, repeated measures mixed model, p=<0.0001; post hoc comparison using Tukey's correction, baseline compared to 110 dB, ***, p=0.0001, baseline compared to 120 dB, ***, p=0.0008. Ear Tip Tilt: baseline compared to each sound level, repeated measures mixed model, p=0.0148; Eye Ratio: baseline compared to each sound level, repeated measures mixed model, p=0.0354; Mouth Position: baseline compared to each sound level, repeated measures mixed model, ns, p=0.1308. Snout Position: baseline compared to each sound level, repeated measures mixed model, ns, p=0.5743. **G**) The following graphs show the averaged data for each body position parameter indicated by the title. Each dot is the averaged data for the indicated 2-minute time-period for 1 mouse. The sound levels have been reordered from lowest to most intense sound level. Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. Rel. Nose Tip Position: baseline compared to each sound level, repeated measures mixed model, p=0.0016; post hoc comparison using Tukey's correction, baseline compared to 90 dB, *, p=0.0142, baseline compared to 100 dB, **, p=0.0095, baseline compared to 110 dB, **, p=0.0058, baseline compared to 120 dB, *, p=0.0473. Percent in Top: baseline compared to each sound level, repeated measures mixed model, p=<0.0001; post hoc comparison using Tukey's correction, baseline compared to 80 dB, *, p=0.0186, baseline compared to 90 dB, **, p=0.0022, baseline compared to 100 dB, **, p=0.0026, baseline compared to 110 dB, ***, p=0.0005, baseline compared to 120 dB, **, p=0.0066. Vertical Line Crosses: baseline compared to each sound level, repeated measures mixed model, p=0.0078; post hoc comparison using Tukey's correction, baseline compared to 70 dB, *, p=0.0112, baseline compared to 100 dB, **, p=0.0018, baseline compared to 110 dB, *, p=0.0122, baseline compared to 120 dB, *, p=0.0122. Face Inclination: baseline compared to each sound level, repeated measures mixed model, p=0.0221.

Figure 7: Facial grimace and body position measurements return to baseline between all sound intervals except following 120 dB

A) The graphs show the averaged data for each facial grimace parameter indicated by the title. Each dot is the averaged data for the indicated 2-minute time-period for 1 mouse. The data have been reordered based on the preceding sound level of the no sound period from lowest to most intense sound level. Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. Ear Ratio: baseline compared to each sound level, repeated measures mixed model, p=0.0012. All other facial parameters showed no significant changes. **B)** The graphs show the averaged data for each body position parameter indicated by the title. Each dot is the averaged data for the indicated 2-minute time-period for 1 mouse. The data have been reordered based on the preceding sound level of the silent sound period from lowest to most intense sound level. Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. Rel. Nose Tip Position: baseline compared to each sound level, repeated measures mixed model, p=0.0003; post hoc comparison using Tukey's correction, baseline compared to 120 dB, **, p=0.002. Percent in Top: baseline compared to each sound level, repeated measures mixed model, p=0.0013; post hoc comparison using Tukey's correction, baseline compared to 120 dB, ***, p=0.001. Vertical Line Crosses: baseline compared to each sound level, repeated measures mixed model, p=0.005; post hoc comparison using Tukey's correction, baseline compared to 120 dB, ***, p=0.0001. Face Inclination: baseline compared to each sound level, repeated measures mixed model, p=0.0161.

Figure 8: Sound above 100 dB evokes behavioral changes associated with pain

A) Facial Grimace (left) and Body Position (Right) Scores for mice during exposure to sound (N=17). Facial Grimace Score, Sound: repeated measures mixed effects model, p=<0.0001; post hoc comparisons using Tukey's correction; baseline compared to 100 dB, **, p=0.0025, baseline compared to 110 dB, ****, p=<0.0001, baseline compared to 120 dB, ***, p=0.0003. Body Position Score, Sound: repeated measures mixed effects model, p=0.0012; post hoc comparisons using Tukey's correction; baseline compared to 100 dB, ***, p=0.0003, baseline compared to 110 dB, *, p=0.0174, baseline compared to 120 dB, ***, p=0.0006. The Pain Level 1 threshold is -19.04 for Facial Grimace and -132.4 for Body Position (dotted lines). The Pain Level 2 threshold is -40.62 for Facial Grimace and -179.5 for Body Position (dashed lines). **B**) Facial Grimace (Left) and Body Position (Right) Scores for mice during the no sound periods (N=17). Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. Facial Grimace Score, No Sound: baseline compared to each sound level, repeated measures mixed model, ns, p=0.1157. Body Position Score, No Sound: baseline compared to each sound level, repeated measures mixed model, ns, p=0.2341

Figure 9: Mice deficient in mechanotransduction do not respond to increased sound levels with significant changes to facial grimace or position associated with pain

A) Example of analyzed frames of Ear Position measurements for one *Tmie^CTL* mouse for the entirety of one recording. The black vertical lines indicate when the sound levels were manually

changed on or off. **B**) Example of analyzed frames of Rel. Nose Tip Position measurements for one *Tmie^{CTL}* mouse for the entirety of one recording. The black vertical lines indicate when the sound levels were manually changed on or off. **C**) The following graphs show the averaged data for each facial parameter indicated. Each dot is the averaged data for the indicated 2-minute time-period for 1 *Tmie^{CTL}* mouse (N=10). The sound levels have been reordered from lowest to most intense sound level. Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. Ear Ratio: repeated measures mixed effects model, p=0.0009; post hoc comparisons using Tukey's correction, baseline compared to 90 dB, ***, p=0.0009, baseline compared to 100 dB, **, p=0.0049, baseline compared to 120 dB, *, p=0.0272. Ear Position: repeated measures mixed effects model, p=0.0245; post hoc comparisons using Tukey's correction, baseline compared to 110 dB, *, p=0.027, baseline compared to 120 dB, **, p=0.0033. Eye Ratio: repeated measures mixed effects model, p=0.0293. **D**) The following graphs show the averaged data for each body position parameter indicated by the title. Each dot is the averaged data for the indicated 2-minute time-period for 1 *Tmie^{CTL}* mouse (N=10). The sound levels have been reordered from lowest to most intense sound level. Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. Rel. Nose Tip Position: repeated measures mixed effects model, p=0.014; post hoc comparisons using Tukey's correction, baseline compared to 110 dB, **, p=0.037. Percent in Top: repeated measures mixed effects model, p=0.0124; post hoc comparisons using Tukey's correction, baseline compared to 110 dB, *, p=0.0224. Vertical Line Crosses: repeated measures mixed effects model, p=0.0078; post hoc comparisons using Tukey's correction, baseline compared to 90 dB, *, p=0.0207, baseline compared to 100 dB, **, p=0.0026, baseline compared to 110 dB, **, p=0.0027. Face Inclination: repeated measures mixed effects model, ns, p=0.1787; post hoc comparisons using Tukey's correction, baseline compared to 110 dB, *, p=0.0316. **E**) Example of analyzed frames of Ear Position measurements for one *Tmie^{-/-}* mouse for the entirety of one recording. The black vertical lines indicate when the sound levels were manually changed on or off. **F**) Example of analyzed frames of Rel. Nose Tip Position measurements for one *Tmie^{-/-}* mouse for the entirety of one recording. The black vertical lines indicate when the sound levels were manually changed on or off. **G**) The following graphs show the averaged data for each facial parameter indicated. Each dot is the averaged data for the indicated 2-minute time-period for 1 *Tmie^{-/-}* mouse (N=8). The sound levels have been reordered from lowest to most intense sound level. Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. **H**) The following graphs show the averaged data for each body position parameter indicated by the title. Each dot is the averaged data for the indicated 2-minute time-period for 1 *Tmie^{-/-}* mouse (N=8). The sound levels have been reordered from lowest to most intense sound level. Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. Rel. Nose Tip Position: repeated measures mixed effects model, p=0.0401. Note this relationship shows an increase Rel. Nose Tip Position not associated with pain. Vertical Line Crosses: repeated measures mixed effects model, p=0.0005; post hoc comparisons using Tukey's correction, baseline compared to 110 dB, **, p=0.0028.

Figure 10: Mice deficient in mechanotransduction do not show pain-associated behavioral changes when exposed to loud sounds like wildtype mice.

A) Facial Grimace (Left) and Body Position (Right) Scores for $Tmie^{CTL}$ mice exposed to sound(N=10). Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. Facial Grimace Score, $Tmie^{CTL}$: repeated measures mixed effects model, p=0.0004; post hoc comparisons using Tukey's correction, baseline compared to 90 dB, *, p=0.0417, baseline compared to 100 dB, **, p=0.0011, baseline compared to 120 dB, ***, p=0.0007. Body Position Score, $Tmie^{CTL}$: repeated measures mixed effects model, p=0.0031; post hoc comparisons using Tukey's correction, baseline compared to 100 dB, ***, p=0.0007, baseline compared to 110 dB, ****, p=<0.0001. **B)** Facial Grimace (left) and Body Position (Right) Scores for $Tmie^{-/-}$ mice exposed to sound (N=8). Facial Grimace Score, $Tmie^{-/-}$: repeated measures mixed effects model, ns, p=0.535; Body Position Score, $Tmie^{-/-}$: repeated measures mixed effects model, p=0.0296. For a mouse to be experiencing pain, the scores must be lower than -19.04 for Facial Grimace and -132.4 for Body Position (dotted line). For a mouse to experience Pain Level 2, the scores must be lower than -40.62 for Facial Grimace and -179.5 for Body Position (dashed line).

Supplemental Figure 1: Animals injected with saline do not show changes to facial grimace measurements

A) These graphs show the averaged data for each facial parameter indicated. Each dot is the averaged data for the indicated time-period for 1 mouse. Baseline is the averaged data for each mouse for all frames from the Baseline video (10 minutes). 'All' is the averaged data for each mouse for all frames after CGRP injection. 'Peak' is the averaged data from minutes 28-33 and after CGRP injection. 'Non-Peak' is the averaged data from time outside of the peak time period (minutes 20-26 and minutes 32 to 40). Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. Eye Ratio, repeated measures mixed model, p=0.0465; post hoc comparisons using Tukey's correction, Peak compared to Non-Peak, ***, p=0.0003. **B)** These graphs show the averaged data for each body position measurement indicated. Each dot is the averaged data for 1 mouse. Baseline is the averaged data for each mouse for all frames from the Baseline video. 'All' is the averaged data for each mouse for all frames after CGRP injection. 'Peak' is the averaged data from minutes 27-31 and after CGRP injection. 'Non-Peak' is the averaged data from time outside of the peak time period (minutes 20-26 and minutes 32 to 40). Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. If no statistics noted, data was not significantly different.

Supplemental Figure 2: Percentage Change from Baseline for each measurement used to generate Facial Grimace and Body Position Scores for Saline- and CGRP-injected mice

A) Graphs of Facial Grimace (left) and Body Position (right) parameters for CGRP-injected mice (N=12). Each dot is the average of one mouse for the indicated time-period after being transformed into a percentage compared to the baseline time-period. If a measurement resulted in a positive change from baseline (Ear Ratio, Ear Position, Rel. Nose Tip Position, Face Inclination), it was inverted so the percentage change from baseline associated with pain was a

negative value. Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. Eye Ratio: Baseline compared to All, paired, two-tailed t-test, p=0.0002; Baseline compared to Peak and Non-Peak, repeated measured mixed model, p= 0.0018, ; post hoc comparisons using Tukey's correction, Baseline compared to Peak, *, p=0.0105, Baseline compared to Non-Peak, **, p=0.0024. Mouth Position: Baseline compared to Peak and Non-Peak, repeated measured mixed model, p= 0.008 ; post hoc comparisons using Tukey's correction, Baseline compared to Peak, *, p=0.0355, Peak compared to Non-Peak, *, p=0.0417. Percent in Top: Baseline compared to Peak and Non-Peak, repeated measures Friedman Test, p= 0.0435 ; post hoc comparisons using Dunn's correction, Baseline compared to Peak, *, p=0.0315. Vertical Line Crosses: Baseline compared to Peak and Non-Peak, repeated measured mixed model, ns, p= 0.6286 ; post hoc comparisons using Tukey's correction, Baseline compared to Non-Peak, *, p=0.019. Face Inclination: Baseline compared to All, paired two-tailed t-test, *, p=0.0255. If no statistics noted, data was not significantly different. **B**) Graphs of Facial Grimace (left) and Body Position (right) parameters for saline-injected mice (N=16). Each open circle is the average of one mouse for the indicated time-period after being transformed into a percentage compared to the baseline time-period. If a measurement resulted in a positive change from baseline (Ear Ratio, Ear Position, Rel. Nose Tip Position, Face Inclination), it was inverted so the percentage change from baseline associated with pain was a negative value. Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. Eye Ratio: Baseline compared to Peak and Non-Peak, repeated measures mixed model, p= 0.0381. Vertical Line Crosses: Baseline compared to All, Wilcoxon signed-rank test, p=0.0494; Baseline compared to Peak and Non-Peak, repeated measures one-way ANOVA, p=0.0379. If no statistics noted, data was not significantly different. All groups were tested for normality. In cases where the data from all groups was not normally distributed, a Wilcoxon Ranked sum test was used for comparisons of Baseline to All and a Friedman test was used for the comparison of Baseline to Peak and Non-Peak.

Supplemental Figure 3: Percentage change from Baseline for each measurement used to generate Facial Grimace and Body Position Scores for Sound- and No-Sound- exposed mice.

A) Graphs of Facial Grimace (left) and Body Position (right) parameters for sound intervals of the interleaved sound exposure experiment (N=17). Each dot is the average of one mouse for the indicated time-period after being transformed into a percentage compared to the baseline time-period. Sound time-periods have been reordered according to the increasing sound level. If a measurement resulted in a positive change from baseline (Ear Ratio, Ear Position, Rel. Nose Tip Position, Face Inclination), it was inverted so the percentage change from baseline associated with pain was a negative value. Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. Ear Ratio: repeated measures mixed effects model, p=0.0014; post hoc comparisons using Tukey's correction; baseline compared to 70 dB, **, p=0.0063, baseline compared to 90 dB, *, p=0.0149, baseline compared to 100 dB, *, p=0.0217, baseline compared to 110 dB, *,

p=0.0104, baseline compared to 120 dB, **, p=0.0036. Ear Position: repeated measures mixed effects model, p=<0.0001; baseline compared to 110 dB, ***, p=0.0001, baseline compared to 120 dB, **, p=0.001. Ear Tip Tilt: repeated measures mixed effects model, p=0.0139. Eye Ratio: repeated measures mixed effects model, p=0.0469. Rel. Nose Tip Position: repeated measures mixed effects model, p=0.002; post hoc comparisons using Tukey's correction; baseline compared to 90 dB, *, p=0.0229, baseline compared to 100 dB, *, p=0.0132, baseline compared to 110 dB, *, p=0.0172. Mouth Position, Snout Position, Percent in Top, Vertical Line Crosses, and Face Inclination showed no significant differences between sound levels or from baseline. **B)** Graphs of Facial Grimace (left) and Body Position (right) parameters for no sound intervals of the interleaved sound exposure experiment (N=17). Each dot is the average of one mouse for the indicated time-period after being transformed into a percentage compared to the baseline time-period. No sound time-periods have been reordered according to the preceding sound level. If a measurement resulted in a positive change from baseline (Ear Ratio, Ear Position, Rel. Nose Tip Position, Face Inclination), it was inverted so the percentage change from baseline associated with pain was a negative value. Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. Ear Ratio: repeated measures mixed effects model, p=0.0072; Rel. Nose Tip Position: repeated measures mixed effects model, p=0.0003, post hoc comparisons using Tukey's correction; baseline compared to post 120 dB, **, p=0.005. Ear Position, Ear Tip Tilt, Eye Ratio, Mouth Position, Snout position, Percent in Top, Vertical line Crosses, and Face Inclination showed no significant differences between no sound periods or from baseline.

Supplemental Figure 4: Percentage change from Baseline for each measurement used to generate Facial Grimace and Body Position Scores for $Tmie^{CTL}$ and $Tmie^{-/-}$ sound-exposed mice

A) Graphs of Facial Grimace (left) and Body Position (right) parameters for sound intervals of $Tmie^{CTL}$ mice in the interleaved sound exposure experiment (N=10). Each dot is the average of one mouse for the indicated time-period after being transformed into a percentage compared to the baseline time-period. Sound time-periods have been reordered according to increasing sound level. If a measurement resulted in a positive change from baseline (Ear Ratio, Ear Position, Rel. Nose Tip Position, Face Inclination), it was inverted so the percentage change from baseline associated with pain was a negative value. Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. Ear Ratio, $Tmie^{CTL}$: repeated measures mixed effects model, p=0.0025; post hoc comparisons using Tukey's correction, baseline compared to 90 dB, **, p=0.0017, baseline compared to 100 dB, **, p=0.0092. Ear Position, $Tmie^{CTL}$: repeated measures mixed effects

model, p=0.0349; post hoc comparisons using Tukey's correction, baseline compared to 110 dB, *, p=0.0237, baseline compared to 120 dB, **, p=0.0035. Eye Ratio, $Tmie^{CTL}$: repeated measures mixed effects model, p=0.0321. Rel. Nose Tip Position, $Tmie^{CTL}$: repeated measures mixed effects model, p=0.0192; post hoc comparisons using Tukey's correction, baseline compared to 110 dB, **, p=0.0065. Percent in Top, $Tmie^{CTL}$: repeated measures mixed effects model, p=0.0064; post hoc comparisons using Tukey's correction, baseline compared to 100 dB, **, p=0.0035, baseline compared to 110 dB, ***, p=0.0001. Vertical line crosses, $Tmie^{CTL}$: repeated measures mixed effects model, p=0.0062; post hoc comparisons using Tukey's correction, baseline compared to 100 dB, ***, p=0.0004, baseline compared to 110 dB, ****, p=<0.0001. Face Inclination **B**) Graphs of Facial Grimace (left) and Body Position (right) parameters for sound intervals of $Tmie^{-/-}$ mice in the interleaved sound exposure experiment (N=8). Each dot is the average of one mouse for the indicated time-period after being transformed into a percentage compared to the baseline time-period. Sound time-periods have been reordered according to the increasing sound level. If a measurement resulted in a positive change from baseline (Ear Ratio, Ear Position, Rel. Nose Tip Position, Face Inclination), it was inverted so the percentage change from baseline associated with pain was a negative value. Box and whiskers have a line at the median and the box is the interquartile range with the whiskers extending to the maximum and minimum value of the data. Rel. Nose Tip Position $Tmie^{-/-}$: repeated measures mixed effects model, p=0.0293. Vertical Line Crosses, $Tmie^{-/-}$: repeated measures mixed effects model, p=0.0135; post hoc comparisons using Tukey's correction, baseline compared to 110 dB, ****, p=<0.0001.

Video Figure Legends

<u>Supplemental Video 1:</u> CGRP Experiment Example
10 second video clips from the Baseline recording and 3 segments during the CGRP recording - First 2 Minutes, Peak and Last 2 Minutes. The labels were generated by Deeplabcut and any marker above 0.6 likelihood is shown for visualization. Only markers with a likelihood above 0.95 were used for analysis.

<u>Supplemental Video 2:</u> Interleaved Sound Experiment Example
10 second video clips from the original recording of the Baseline, 110 dB SPL, post 110 dB SPL, and 70 dB SPL of a mouse undergoing the interleaved sound protocol. Sound is included in the video.

<u>Supplemental Video 3:</u> Labeled, Interleaved Sound Experiment Example
10 second video clips from the labeled recording of the Baseline, 110 dB SPL, post 110 dB SPL, and 70 dB SPL time periods of a mouse undergoing the interleaved sound protocol. The labeled video is from the same time periods from the same mouse shown in Supplemental Video 2. The labels were generated by Deeplabcut and any marker above 0.6 likelihood is shown for visualization. Only markers with a likelihood above 0.95 were used for analysis.

<u>Supplemental Video 4:</u> *Tmie* Wildtype Example Video
10 second video clips from an original recording of the Baseline and 110 dB SPL time periods of a *Tmie^{CTL}* mouse undergoing the interleaved sound protocol with sound included. The original clips are followed by the corresponding labeled video. The labels were generated by Deeplabcut and any marker above 0.6 likelihood is shown for visualization. Only markers with a likelihood above 0.95 were used for analysis.

<u>Supplemental Video 5:</u> *Tmie* Knockout Example Video
10 second video clips from an original recording of the Baseline and 110 dB SPL time periods of a *Tmie$^{-/-}$* mouse undergoing the interleaved sound protocol with sound included. The original clips are followed by the corresponding labeled video. The labels were generated by Deeplabcut and any marker above 0.6 likelihood is shown for visualization. Only markers with a likelihood above 0.95 were used for analysis.

References

(Numbered in order of appearance)

1. H. E. von Gierke, D. R. Brown, Wright-Patterson AFB, Ohio. *Benox Report; an Exploratory Study of the Biological Effects of Noise* 47 (1953).

2. G. Frank, M. Kössl, The acoustic two-tone distortions 2f1-f2 and f2-f1 and their possible relation to changes in the operating point of the cochlear amplifier. *Hear Res* **98**, 104–115 (1996).

3. L. P. Sherlock, C. Formby, Estimates of loudness, loudness discomfort, and the auditory dynamic range: normative estimates, comparison of procedures, and test-retest reliability. *J Am Acad Audiol* **16**, 85–100 (2005).

4. S. Manohar, H. J. Adler, K. Radziwon, R. Salvi, Interaction of auditory and pain pathways: Effects of stimulus intensity, hearing loss and opioid signaling. *Hear Res* **393**, 108012 (2020).

5. R. S. Tyler, *et al.*, A review of hyperacusis and future directions: part I. Definitions and manifestations. *Am J Audiol* **23**, 402–419 (2014).

6. M. Pienkowski, *et al.*, A review of hyperacusis and future directions: part II. Measurement, mechanisms, and treatment. *Am J Audiol* **23**, 420–436 (2014).

7. B. Pollard, Clinical advancements for managing hyperacusis with pain. *The Hearing Journal* **72**, 10–12 (2019).

8. R. Salvi, G.-D. Chen, S. Manohar, Hyperacusis: Loudness intolerance, fear, annoyance and pain. *Hear Res* **426**, 108648 (2022).

9. K. N. Jahn, S. T. Kashiwagura, M. S. Yousuf, Clinical phenotype and management of sound-induced pain: Insights from adults with pain hyperacusis. *J Pain* **27**, 104741 (2025).

10. Q. Song, M.-G. Liu, M. Zhuo, Minocycline does not affect long-term potentiation in the anterior cingulate cortex of normal adult mice. *Mol Pain* **11**, 25 (2015).

11. S.-B. Zhou, *et al.*, Enhancement of behavioral nociceptive responses but not itching responses by viewing mirror images in adult mice. *Mol Pain* **18**, 17448069221111158 (2022).

12. K. K. Clayton, *et al.*, Sound elicits stereotyped facial movements that provide a sensitive index of hearing abilities in mice. *Curr Biol* **34**, 1605-1620.e5 (2024).

13. K. Radziwon, R. Salvi, Using auditory reaction time to measure loudness growth in rats. *Hear Res* **395**, 108026 (2020).

14. A. E. Hickox, M. C. Liberman, Is noise-induced cochlear neuropathy key to the generation of hyperacusis or tinnitus? *J Neurophysiol* **111**, 552–564 (2014).

15.   R. H. Salloum, *et al.*, Untangling the effects of tinnitus and hypersensitivity to sound (hyperacusis) in the gap detection test. *Hear Res* **331**, 92–100 (2016).

16.   J. R. Deuis, L. S. Dvorakova, I. Vetter, Methods Used to Evaluate Pain Behaviors in Rodents. *Front Mol Neurosci* **10**, 284 (2017).

17.   K. J. Sufka, Conditioned place preference paradigm: a novel approach for analgesic drug assessment against chronic pain. *Pain* **58**, 355–366 (1994).

18.   Z. Zhang, *et al.*, Automated preclinical detection of mechanical pain hypersensitivity and analgesia. *Pain* **163**, 2326–2336 (2022).

19.   M. Bohic, *et al.*, Mapping the neuroethological signatures of pain, analgesia, and recovery in mice. *Neuron* **111**, 2811-2830.e8 (2023).

20.   D. J. Langford, *et al.*, Coding of facial expressions of pain in the laboratory mouse. *Nat Methods* **7**, 447–449 (2010).

21.   E. S. McCoy, *et al.*, Development of PainFace software to simplify, standardize, and scale up mouse grimace analyses. *Pain* **165**, 1793–1805 (2024).

22.   O. Le Moëne, M. Larsson, A New Tool for Quantifying Mouse Facial Expressions. *eNeuro* **10**, ENEURO.0349-22.2022 (2023).

23.   B. J. Rea, *et al.*, Peripherally administered calcitonin gene-related peptide induces spontaneous pain in mice: implications for migraine. *Pain* **159**, 2306–2317 (2018).

24.   G. J. Bennett, What is spontaneous pain and who has it? *J Pain* **13**, 921–929 (2012).

25.   E. J. Cobos, *et al.*, Inflammation-induced decrease in voluntary wheel running in mice: a nonreflexive test for evaluating inflammatory pain and analgesia. *Pain* **153**, 876–884 (2012).

26.   P. V. Turner, D. S. Pang, J. L. Lofgren, A Review of Pain Assessment Methods in Laboratory Rodents. *Comp Med* **69**, 451–467 (2019).

27.   M. R. Gleason, *et al.*, The transmembrane inner ear (Tmie) protein is essential for normal hearing and balance in the zebrafish. *Proc Natl Acad Sci U S A* **106**, 21347–21352 (2009).

28.   Z. Wu, *et al.*, Mechanosensory hair cells express two molecularly distinct mechanotransduction channels. *Nat Neurosci* **20**, 24–33 (2017).

29.   J. R. Holt, *et al.*, Putting the Pieces Together: the Hair Cell Transduction Complex. *J Assoc Res Otolaryngol* **22**, 601–608 (2021).

30.   R. Fettiplace, D. N. Furness, M. Beurg, The conductance and organization of the TMC1-containing mechanotransducer channel complex in auditory hair cells. *Proc Natl Acad Sci U S A* **119**, e2210849119 (2022).

31. A. H. Tuttle, *et al.*, A deep neural network to assess spontaneous pain from mouse facial expressions. *Mol Pain* **14**, 1744806918763658 (2018).

32. C.-Y. Chiang, *et al.*, Deep Learning-Based Grimace Scoring Is Comparable to Human Scoring in a Mouse Migraine Model. *J Pers Med* **12** (2022).

33. A. Mathis, *et al.*, DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nat Neurosci* **21**, 1281–1289 (2018).

34. T. Nath, *et al.*, Using DeepLabCut for 3D markerless pose estimation across species and behaviors. *Nat Protoc* **14**, 2152–2176 (2019).

35. S. M. Stubsjøen, *et al.*, Exploring non-invasive methods to assess pain in sheep. *Physiol Behav* **98**, 640–648 (2009).

36. J. Stracke, B. Bert, H. Fink, J. Böhner, Assessment of stress in laboratory beagle dogs constrained by a Pavlov sling. *ALTEX* **28**, 317–325 (2011).

37. A. Schroeer, F. I. Corona-Strauss, R. Hannemann, S. A. Hackley, D. J. Strauss, The vestigial pinna-orienting system in humans briefly suppresses superior auricular muscle activity during reflexive orienting toward auditory stimuli. *J Neurophysiol* **132**, 514–526 (2024).

38. E. Friauf, H. Herbert, Topographic organization of facial motoneurons to individual pinna muscles in rat (Rattus rattus) and bat (Rousettus aegyptiacus). *J Comp Neurol* **240**, 161–170 (1985).

39. J. E. Heuser, T. I. Tenkova, Introducing a mammalian nerve-muscle preparation ideal for physiology and microscopy, the transverse auricular muscle in the ear of the mouse. *Neuroscience* **439**, 80–105 (2020).

40. L. H. Lassen, *et al.*, CGRP may play a causative role in migraine. *Cephalalgia* **22**, 54–61 (2002).

41. B. N. Mason, *et al.*, Induction of Migraine-Like Photophobic Behavior in Mice by Both Peripheral and Central CGRP Mechanisms. *J Neurosci* **37**, 204–216 (2017).

42. Hasriadi, P. W. D. Wasana, O. Vajragupta, P. Rojsitthisak, P. Towiwat, Automated home-cage for the evaluation of innate non-reflexive pain behaviors in a mouse model of inflammatory pain. *Sci Rep* **11**, 12240 (2021).

43. M. J. O. Boedts, Tympanic Resonance Hypothesis. *Front Neurol* **11**, 14 (2020).

44. R. T. Bottom, Y. Xu, C. Siebald, J. Jung, U. Müller, Defects in hair cells disrupt the development of auditory peripheral circuitry. *Nat Commun* **15**, 10899 (2024).

45. P. Fournier, *et al.*, Contraction of the stapedius and tensor tympani muscles explored by tympanometry and pressure measurement in the external auditory canal. *Hear Res* **420**, 108509 (2022).

46. V. Rinaldi, M. Cappadona, M. Gaffuri, S. Torretta, L. Pignataro, Chorda tympani nerve, may it have a role in stabilizing middle ear pressure? *Medical hypotheses* **80**, 726–727 (2013).

47. T. J. Rockley, W. M. Hawke, The middle ear as a baroreceptor. *Acta Otolaryngol* **112**, 816–823 (1992).

48. M. Westcott, *et al.*, Tonic tensor tympani syndrome in tinnitus and hyperacusis patients: a multi-clinic prevalence study. *Noise Health* **15**, 117–128 (2013).

49. A. E. Sutton, R. De Jong, G. Kwartowitz, "Tensor Tympani Syndrome." in *StatPearls*, (StatPearls Publishing, 2025).

50. A. S. Filippidis, S. G. Zarogiannis, A. Randich, T. J. Ness, S. Matalon, Assessment of locomotion in chlorine exposed mice by computer vision and neural networks. *J Appl Physiol (1985)* **112**, 1064–1072 (2012).

51. A. M. Kolstad, R. M. Rodriguiz, C. J. Kim, L. P. Hale, Effect of pain management on immunization efficacy in mice. *J Am Assoc Lab Anim Sci* **51**, 448–457 (2012).

52. D. Vuralli, A.-S. Wattiez, A. F. Russo, H. Bolay, Behavioral and cognitive animal models in headache research. *J Headache Pain* **20**, 11 (2019).

53. A. Ashiquzzaman, *et al.*, MoSeq based 3D behavioral profiling uncovers neuropathic behavior changes in diabetic mouse model. *Sci Rep* **15**, 15114 (2025).

54. C. Weinreb, *et al.*, Keypoint-MoSeq: parsing behavior by linking point tracking to pose dynamics. *Nat Methods* **21**, 1329–1339 (2024).

55. N. R. Council, D. on Earth, I. for L. A. Research, C. on Recognition, A. of P. in L. Animals, Recognition and alleviation of pain in laboratory animals. (2010).

56. S. Manohar, G.-D. Chen, L. Li, X. Liu, R. Salvi, Chronic stress induced loudness hyperacusis, sound avoidance and auditory cortex hyperactivity. *Hear Res* **431**, 108726 (2023).

57. S. Manohar, J. Spoth, K. Radziwon, B. D. Auerbach, R. Salvi, Noise-induced hearing loss induces loudness intolerance in a rat Active Sound Avoidance Paradigm (ASAP). *Hear Res* **353**, 197–203 (2017).

58. W. A. Yost, M. C. Killion, Hearing thresholds. *Encyclopedia of acoustics* **3**, 1545–1554 (1997).

59. S. E. Shore, C. Wu, Mechanisms of Noise-Induced Tinnitus: Insights from Cellular Studies. *Neuron* **103**, 8–20 (2019).

60. T. Nagai, T. Tono, Encapsulated nerve corpuscles in the human tympanic membrane. *Arch Otorhinolaryngol* **246**, 169–172 (1989).

61. K. Limakatso, G. J. Bedwell, V. J. Madden, R. Parker, The prevalence and risk factors for phantom limb pain in people with amputations: A systematic review and meta-analysis. *PLoS One* **15**, e0240431 (2020).

62. M. McGill, *et al.*, Neural signatures of auditory hypersensitivity following acoustic trauma. *Elife* **11** (2022).