Title: Seeking smarts: Male chickadees with better spatial cognition sire more extra-pair young

Corresponding author: Carrie L Branch, cbranch6@uwo.ca

Authors: Carrie L Branch*[1,2], Benjamin R Sonnenberg[2,3], Joseph F Welklin[2], Bronwyn G

Butcher[4], Virginia K Heinen[2], Angela M Pitera[2], Lauren M Benedict[2], Eli S Bridge[5], Irby J

Lovette[4], Michael S Webster[4], Vladimir V Pravosudov[2]

Affiliations:

[1] Department of Psychology, The University of Western Ontario, 1151 Richmond St London,

ON, N6A 3K7 Canada

[2] Department of Biology, University of Nevada, Reno, 1664 N Virginia St Reno, NV, 89557 USA

[3] Department of Integrative Biology, Oregon State University, 1500 SW Jefferson Way,

Corvallis, OR, 97331

[4] Cornell Lab of Ornithology, Cornell University, 159 Sapsucker Woods Rd, Ithaca, NY, 14850

USA

[5] Department of Biology, University of Oklahoma, 660 Parrington Oval, Norman, OK, 73019

USA

Abstract

Across the animal kingdom, females commonly mate with more than one male even when they

are monogamous. Extra-pair males rarely contribute resources outside of sperm, therefore, many

hypotheses surrounding the fitness benefits of extra-pair (EP) paternity involve genetic

mechanisms. The 'good genes' hypothesis suggests that socially monogamous females mate

outside their pair bond to increase the fitness of their offspring via indirect genetic benefits. We

tested this hypothesis by quantifying extra-pair paternity in nonmigratory, food-caching

mountain chickadees (*Poecile gambeli*). Chickadees rely on spatial cognition to recover scattered

food caches and variation in spatial cognition is associated with increased survival, longer

lifespan, and is heritable, however, less is known about the relationship between reproductive

success and spatial cognition. In the current study, we had two main objectives; to assess the

relationship between spatial cognition and reproductive output via extra-pair young and to

compare the spatial cognition of EP males to the within pair males they cuckolded. We found

that 1. males with better spatial cognitive abilities have more extra-pair young and produce

heavier offspring in their own nests compared to their poorer performing counterparts, and 2. EP

males have significantly better spatial cognition than the social males they cuckolded. These

results show that males with better spatial cognitive abilities exhibit higher reproductive success

and are consistent with the good genes hypothesis if females gain genetic benefits associated

with better cognitive abilities via EP young.

Key words: extra-pair mating, spatial cognition, indirect fitness, good genes, reproduction

Introduction

Understanding the factors that influence mating decisions in animals and the consequences that follow from these decisions is a central focus of evolutionary biology (1-8). Socially monogamous species provide an interesting case as they form seasonal to long-term pair bonds but produce extra-pair young (EPY) outside the pair bond (9-11). While EPY are common and significantly contribute to lifetime reproductive success, rates of extra-pair paternity (EPP) vary widely within and between species (9, 12). Given that extra-pair males rarely contribute to parental care or provide access to resources outside of sperm (13-15), many hypotheses surrounding the fitness benefits of EPP involve genetic mechanisms, including 'good' or 'compatible' genes, genetic diversity, and insurance against male infertility (13, 15-22). Despite potential fitness benefits, the opportunity to pursue EP matings, particularly for females, may be constrained by mate guarding or breeding density (9,14, 21, 23) resulting in the highly variable EPP rates reported in the literature (9).

What constitutes 'good genes' in a particular system or population can often be esoteric, but we can generally assume that genes passed from one generation to the next that enhance fitness will be passed on via selection, regardless of the mechanism. For example, scatter food-caching species in temperate regions do not migrate seasonally, but instead store food in various locations throughout their home ranges when food is abundant, to use later when food is scare (24, 25). The spatial cognitive abilities of food-caching birds have been well studied, and research shows that these birds rely on spatial learning and memory to store and relocate their food items (25, 26). More recently, research in the wild using 'smart' feeders (Figure 1) showed that selection acts on the spatial cognitive abilities of food-caching mountain chickadees, such that individuals that perform better on a spatial learning and memory task in the wild are more

likely to survive winter and live longer lives compared to those that perform poorly (27, 28).

Furthermore, using genome-wide association and gene network approaches, research shows that

individual variation in spatial cognitive abilities is associated with genetic differences, including

differences in genes involved in the development and function of the hippocampus, a brain

region involved in spatial cognitive function (29, 30). Interestingly, these spatial cognitive

abilities remain stable across birds' natural lifespan (e.g., no detectable senescence (31) and

spatial cognitive abilities do not change with age (27, 28, 32).

While individual variation in spatial learning and memory abilities is associated with

genetic differences and better spatial abilities have been shown to increase overwinter survival

and life span, we know less about how they influence reproductive success. In several systems,

males with better cognition have been shown to produce more young (33-37), however, much of

this work focuses on problem-solving which may not be attributed to cognitive mechanisms

(e.g., 38), leaving scant evidence for a positive relationship between cognition abilities and

reproductive output (37). Previous work in our long-term mountain chickadee system has

provided some evidence for a relationship between the cognitive abilities of the social male and

reproductive success (39). However, like most Passerine birds, mountain chickadees are socially

monogamous; they raise their brood via biparental care (40), while engaging in extra-pair

matings outside the pair that commonly result in EPY (11). In the current study, we quantify and

compare rates of extra-pair paternity across 3 years of breeding and assess the relationship

between male spatial cognitive performance and individual rates of EPP. We predicted that males

with better spatial cognitive abilities, and associated 'good genes', would produce more EPY

than their poorer performing counterparts.

If males with better spatial abilities sire more EPY, several alternative mechanisms could

explain this outcome. For instance, males with better cognitive abilities may be better at

soliciting EPCs, they may be more active and move more widely increasing the likelihood of

encountering females, or they may hold better quality territories frequented by multiple females

(41-44). Alternatively, females may copulate with extra-pair males that have better cognition

compared to that of their social mates, as would be predicted by the good genes hypothesis (1,8).

While the good genes hypothesis has been commonly proposed to explain female EP mating

behavior, there is rather limited support for this hypothesis (18, 21, 23).

Spatial cognition is a fitness-related trait in food-caching chickadees (25, 27), which

provides the opportunity to test whether females produce EPY with males that exhibit better

spatial cognition than the social males they cuckhold. Producing EPY with higher quality males

is expected to benefit females indirectly because EPY should inherit the genes associated with

better cognition. Indeed, females of several species show preferences for males with better

cognition in lab-based, pairwise choice paradigms (45-49). And previous work from our field

system showed that females can discriminate among males based on their cognitive abilities

(whether directly or indirectly via secondary sexual traits is unknown) and adjust their

reproductive investment (39). Considerable research in Passerine birds shows that females are

the primary drivers of extra-pair matings (41, 50, 51), with females often foraying into

neighboring male territories to solicit copulations (specifically in chickadees, tits, and titmice:

15, 52-56). Furthermore, forced copulations are uncommon in most Passerines, and when they

do occur, may be less likely to result in fertilization (57-59). Specifically, extensive work on

breeding behavior in chickadees and their relatives has never reported forced copulations (e.g.,

42, 52, 53, 60-63).

We have shown that social pairs are not formed based on cognitive abilities (64), likely

due to constraints on social pairing. Juvenile chickadees are recruited into flocks following the

breeding season (65, 66), and male-female pairs from the previous breeding season typically stay

together (67, 68), therefore, social mate choice for the newly recruited juveniles is constrained by

the availability of unpaired individuals. Therefore, we predict that females may seek EPCs

opportunistically with males that exhibit better spatial cognitive performance than their social

male, thus increasing their indirect fitness via 'good genes' (1, 8, 14, 19). We could not directly

assess female EPCs, but we used an indirect approach commonly used to address this question:

comparing the phenotype of the extra-pair male to that of the social male (e.g., 14, 42, 52, 53,

69). The good genes hypothesis predicts that females should seek EPCs when their social mate is

of lower quality and with a male that is of better quality than their social male (1, 8, 14, 19). We

tested this hypothesis by comparing the spatial cognition of males siring EPY in a nest to the

social male they cuckolded. If females obtain genetic benefits via EPY, we expect EP males to

have significantly better spatial cognition compared to the within pair males they cuckold.

Figure 1. Photographs of spatial cognitive testing apparatus. A. Side-view of one array suspended in the air. B. A mountain chickadee tagged with a passive integrated transponder (PIT) tag visiting the 'smart' feeder array. Motorized door is open. The blue band on the left leg of the bird is the PIT tag. C. View of the testing apparatus from below, depicting equidistant placement of eight 'smart' feeders. Data presented come from four independent feeder arrays.

Methods

Experimental Design

We studied spatial learning and memory abilities and extra-pair paternity (EPP) rates in mountain chickadees (*Poecile gambeli*) at our long-term field site in the northern Sierra Nevada mountains of North America (Sagehen Experimental Forest in northern California, USA, Sagehen Creek Field Station, University of California Berkeley). Since 2015, we have monitored breeding phenology in uniquely color-banded individuals across two elevations (63, 70) and since 2016 we have measured the spatial cognitive abilities of hundreds of individuals (e.g., 27-32, 39, 64, 71-74). Data for the paternity analyses were available for the 2018, 2020, and 2021 breeding seasons, while analyses on overall patterns of reproductive output are from the entire database (2015 – 2023). Cognitive data from individuals spanning 2016 – 2023 was used in conjunction with the 3 breeding seasons used for paternity analyses. Birds participate in our cognitive tasks opportunistically, therefore, if a bird was not tested in one of the 3 years we had paternity data for, we were still able to include their cognitive data, as performance does not change over time (27, 28, 32).

The breeding season begins in April and continues through July, while the non-breeding season spans from August to March of the following year. Mountain chickadees are secondary

cavity nesting birds that readily use nest boxes at our field site. Every year in late April we begin

monitoring 350 nest boxes throughout our field site for evidence of nest building. Once nesting

material is detected we monitor all stages of breeding through fledging of each nest, recording

information about the date the first egg is laid, start of incubation, and hatching, and record the

clutch size (number of eggs), and the brood size (number of chicks at post-hatch day 16) of each

nest. We process all chicks in the nest box at post-hatch day 16, which includes banding them

with an aluminum United States Geological Survey (USGS) band, weighing them, and obtaining

a small blood sample for genetic and paternity analyses. We trap adult and first-year birds using

mist nets at established feeders during annual autumn trapping and by hand at nest boxes during

the breeding season. Upon capture, birds are fitted with color-bands and a unique Passive

Integrated Transponder (PIT) tag, wing length is measured, and age is estimated using multiple

plumage features (75). When nestlings banded with USGS bands are recruited into our

population from nest boxes, we band them with an additional color band and PIT tag for

identification.

Testing spatial cognition in the wild

Spatial learning and memory testing was conducted using two feeder arrays at each of our

elevation sites (High, elevation range: 2380 m – 2590 m; coordinates: 39.42402, -120.315015

and Low, elevation range: 1965 m – 2070 m; coordinates: 39.443500, -120.243248), following

(27, 31, 32, 71). Within an elevation, the feeder arrays were established ca. 1.5 km apart with

mostly non-overlapping birds visiting each array. Each array consisted of eight equidistant

RFID-based feeders on a square (122 x 122 cm) aluminum frame suspended ca. 4 m above the

ground. Each feeder is equipped with a small, motorized door (Figure 1). Prior to testing, we

kept the feeder doors of the arrays open for approximately one month. Next, the feeder doors

were closed and programmed to open for any bird with a PIT tag, allowing birds to habituate to

the motorized feeder doors for one week. Finally, we measured the spatial learning and memory

abilities of mountain chickadees using a 4-day task. The spatial learning and memory task

involved pseudo-randomly assigning each bird to a single feeder in the 8-feeder array that was

not its most-visited feeder during the previous two habituation stages. During this task, each bird

can only obtain a food reward (black oil sunflower seed) from its single assigned feeder, but all

eight feeders recorded the date, time, and identity of every visit by all PIT tagged individuals. We

measured cognitive performance by estimating the number of location errors (e.g., non-

rewarding feeders visited prior to visiting the correct rewarding feeder, max 7) per trial, over

multiple trials. A trial began when a bird visited any feeder in the array and ended with a visit to

the rewarding feeder. If the bird landed on its rewarding feeder first, it was considered to have

made zero location errors. Chickadees collect a single sunflower seed during each trial and fly

away from the array to nearby trees to either eat or cache the seed before returning to the array

and each trial takes just a few seconds (73).

We used the mean number of location errors per trial over the first 20 trials and the mean

number of location errors per trial over the entire 4-day task as the two metrics of cognitive

performance, following our previous work (27, 31, 32, 71). These measurements capture learning

curves in a single metric (Figure S7); as birds learn the rewarding feeder location, they make

fewer and fewer errors on each consecutive trial. Using the mean number of location errors over

multiple trials captures this improvement – the mean number of location errors per trial over

multiple trials is smaller for individuals that improved faster and larger for individuals that

improved slower (27). Since birds completed different numbers of trials over the entire task, we

used the total number of trials as a covariate in models involving performance on the entire 4-day

task (27, 31, 32, 71). We have previously documented that variation in both metrics is associated

with significant differences in survival (27), longevity (28), in female reproductive decisions

(39), in foraging decisions (73), and in cognitive flexibility (74). For analyses involving spatial

cognitive abilities, we used cognitive performance measured in the first year each individual

completed the task (between 2016 and 2023), as cognitive performance on the spatial learning

and memory task does not change with age (27, 32). All birds in this study completed a

minimum of 20 trials and showed evidence of learning by performing at better than chance (< 3

errors per trial, 27, 28). Males that bred in more than one year were included in the analyses

(repeated measures, controlled for using male ID as a random effect); as such when reporting

results, we report the sample size for unique nests and 'male-years'.

Assessing extra-pair paternity

DNA extraction and ddRAD sequencing

For this study, we aimed to sequence and genotype parents with cognition data and their

nestlings over four consecutive years (2018-2021). Each year, we prioritized sequencing nests of

males and potential EP sires that performed in our cognitive task. Samples from each year were

sequenced independently, and unfortunately, genotypes from 2019 were of low quality and

paternity assignments could not be completed with confidence, therefore we present data from

2018, 2020, and 2021.

Blood samples from adults and nestlings were stored at room temperature in Queens

Lysis buffer until DNA was extracted using Qiagen DNeasy Blood & Tissue kits. The DNA

concentration of each sample was quantified using Qubit Fluorometer 2.0 (Life Technologies)

prior to library preparation and sequencing. We used a double digest restriction-site associated

DNA (ddRAD) sequencing approach from (76) to obtain single nucleotide polymorphisms

(SNPs) for our paternity analyses. Sequencing was completed in batches following field

sampling of each year, therefore, individuals from each breeding year were run together, but

separately from the other years. For each individual, 100-500 ng of extracted DNA was digested

with restriction enzymes SbfI-HF (NEB) and MspI (NEB), and ligated with one of 20 unique P1

adapters and a P2 adapter (P2-MspI). Next, samples were pooled by groups of 20, maintaining

individual identity via unique P1 adapters. Samples were then purified using 1.5x volume of

homemade solid-phase reversible immobilization (SPRI) beads, made with Sera-Mag SpeedBead

Carboxylate-Modified Magnetic Particles (Cytiva) (77). These pooled samples (index groups)

were sent to Cornell University Biotechnology Resource Center (BRC) for size selection

between 450 – 600 bp using BluePippin (Sage Science). Following size selection, adapter ligated

DNA was enriched by 11 cycles of polymerase chain reaction (PCR) with a universal i5 primer

and a unique indexed i7 primer for each index group using Phusion DNA polymerase (NEB).

Reactions were again cleaned using 0.7x volume of homemade SPRI beads (as above) and the

index groups pooled in equimolar ratios to create a single sequencing library run on one lane of

Ilumnia NextSeq500 (150 bp single end, performed at BRC). Sequencing was performed with a

~20% PhiX spike-in to introduce diversity into the library.

SNP Analysis and Paternity assignment

Quality filtering, demultiplexing, and genome alignment

The following were run the same, but separately for each year of samples. First, the quality of all

reads were assessed using FASTQC version 0.11.9

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). All sequences were trimmed using

fastx_trimmer (FASTX-Toolkit) to a length of 147 bp on the 3' end to exclude low-quality SNP

calls. Next, reads containing a single base with a Phred quality score less than 10 were removed

and any sequences that had more than 5% of the bases with a Phred quality score lower than 20

were also removed. We then used *process_radtags* from STACKS version 2.65 to demultiplex

the sequence reads and obtain files with specific sequences for each individual. Next, we aligned

sequences to the chromosome scale reference genome of mountain chickadees (29, 30) using

Bowtie2 version 2.5.1. The resulting .sam files were converted to .bam files and sorted using the

view and *sort* functions in Samtools version 1.18. These aligned reads were used as the input for

the *ref_map.pl* program in Stacks 2.65.

SNP filtering and assigning paternity

SNPs were exported using *populations* in Stacks 2 version 2.65, and each year's samples were

run separately to maximize the number of SNPs included for downstream paternity analyses

(since every individual does not have data at each SNP locus, using a smaller set of individuals-

i.e., individuals from each year- returns more loci for assigning paternity, as we require loci to be

present in most individuals; see below). All male individuals alive during the year of interest

were included as potential sires. Individuals with more than 50% missing SNP data were

removed from the dataset; only 5 individuals from the 2021 dataset did not meet this criterion.

Loci were exported if they were present in 85% of that year's sample population (*–r*). If a RAD

locus had more than one SNP, data were restricted to the first SNP *(–write_single_snp)* to

minimize those in high linkage disequilibrium. We required a minor allele frequency of 0.05 to

process a nucleotide *(–min_maf),* then filtered for loci in Hardy-Weinberg equilibrium ($p<0.05$)

using VCFtools (--*hwe* option) (78, 79). Variant call format (VCF) files were converted to

GENEPOP format using the 'radiator' package in R (80).

Before assigning paternity, we imported the yearly GENEPOP files into CERVUS (81),

then ran maternity tests separately for each elevation in each year to confirm mother-offspring

relationships identified during field observations. We assessed the quality of these assignments

by plotting the distributions of the first and second-ranked mother's scores for each offspring.

First and second-ranked mother pair log likelihood (LOD) scores formed bimodal distributions,

with nearly all first-ranked mother scores above zero and nearly all second-ranked mother scores

below zero. In rare cases when second-ranked mother scores were positive, these were almost

always known sisters or daughters of the mother identified during field observations. Therefore,

first-ranked mother-offspring pairs with positive LOD scores were considered correct

assignments, but pairs with negative assignments were not included in paternity analyses because

we were not confident in the mother assignment. Offspring resulting from a female laying an egg

in a nest that was not her own (i.e., egg dumps) were uncommon and not included in further

statistical analyses.

Paternity analyses were conducted in CERVUS separately for each elevation in each year

by calculating trio LOD scores for each potential father-offspring pair. Trio LOD scores account

for known maternal genotypes when identifying potential fathers. As we did for mother-offspring

pairs, we plotted the distribution of the first and second-ranked male scores and again found a

bimodal distribution centered around zero (Figure S8). Based on these plots, we set cut-off

scores to assign paternity by: all first-ranked male scores at an elevation in a year that were

positive and greater than the top second-ranked male score at that elevation in that year were

assigned as fathers. Offspring sired by the social male at the nest were considered within-pair,

whereas offspring sired by other males were considered extra-pair. Offspring whose social father

was genotyped, but we could not assign a father to, were also considered extra-pair.

Statistical analysis

We tested whether EPP rates differed between the high and low elevations using generalized

mixed effects models with binomial error distributions and 'logit' links, using the R package

'glmmTMB' (82). Our response variable was whether each offspring was sired by an EP father

or not and we only included nests in which more than 50% of the offspring were genotyped and

run through the paternity analysis. Elevation was included as a fixed effect and mother identity

was included as a random effect in each model. A likelihood ratio test showed statistical support

for an interaction between elevation and year ($p<0.05$), so we tested the effect of elevation on

EPP rates separately for each year.

We employed two methods to test whether spatial cognitive ability was associated with

the number of EPY a male sired in a year. Both models included the number of EPY sired as the

response variable, with spatial cognitive performance (refers to separate models run for (1) the

mean number of location errors per trial over the first 20 trials and (2) the mean number of

location errors per trial over the 4-day task), elevation, and year as fixed effects, and male

identity as a random effect to control for repeated measures of males across years. Both models

used the Poisson error distribution and a 'log' link, but each model accounted for zeros in the

dataset differently. The first method included a zero-inflation parameter (ziformula~1) in

'glmmTMB' to account for observations of males with zero EPY, whereas the second method

restricted the dataset to males that sired at least one extra-pair offspring.

It is possible that a relationship between EPP and spatial cognitive ability is driven by

age, as older males often sire more extra-pair offspring than younger males in many systems

(e.g., 69). While chickadees with better spatial cognition in our study system tend to live longer,

cognitive performance does not change with age in the same individuals (27). However, we

tested whether including male age as a fixed effect improved each model using likelihood ratio

tests that compared models with and without male age.

In addition, cognitive ability could come at a cost to breeding performance, so we tested

whether spatial cognitive ability was associated with breeding performance using a generalized

linear mixed model with the generalized Poisson distribution (83). Brood size was included as

the response variable and male and female spatial performance on the first 20 trials, elevation,

and clutch size were included as fixed effects. For these analyses, we used all breeding data

available, therefore, year was included as a random effect in this analysis because it had a

sufficient number of levels (2015 – 2023). Male and female identity were also included as

random effects to control for yearly variation and repeated measures of individual males and

females. In a second analysis, we used a linear model that included mean nestling mass at day 16

as the response variable (in our system, mass of nestlings on day 16 is a significant predictor of

survival and recruitment (84)), and included male and female performance on the first 20 trials of

the spatial cognitive task and elevation as fixed effects, and year (2015 – 2023) and male and

female identity as random effects. Again, year (2015-2023) and male and female identity were

included as random effects to control for yearly variation and repeated measures of individual

males and females.

Siring extra-pair offspring may reduce a male's investment in his own nest, so we tested

whether siring EPY was associated with breeding performance. We used a generalized linear

mixed model with the generalized Poisson distribution to test whether brood size in a male's nest

was affected by the male siring extra-pair offspring (yes or no), and we used a linear mixed

model to test whether the mean mass of the nestlings at day 16 in a male's nest was affected by

whether the male sired any extra-pair offspring (using three years of data on EPPs). Elevation,

clutch size, and year (three years of EPP data) were included as fixed effects and male and

female identity were included as random effects in both models to control for repeated measures

of individual males and females. We tested whether the mass of young sired by extra-pair males

differed from the mass of young sired by social males using a linear mixed model with nestling

mass at day 16 as the response variable and extra-pair status, elevation, and year as fixed effects.

Nest identity was included as a random effect to control for nestlings that came from the same

nests.

If males initiate EPCs, we expect them to be randomly distributed across different nests,

since the fitness benefit of EPY is increased for the male regardless of the females' social mates'

cognitive abilities. On the other hand, if females initiate EPCs, we expect them to select males of

better quality than their social mate – in our case, males with better spatial cognition – as this

could result in indirect benefits to the female via offspring with better cognition and higher

likelihood of survival (27, 28). Therefore, we tested whether the spatial cognitive abilities of

extra-pair males differed from those of the males they cuckolded by comparing spatial cognition

between father types (EP male vs social male within the same nest) using linear mixed models.

Spatial cognitive performance was used as the response variable, then father type (extra-pair

mate versus social mate) and elevation were included as fixed effects. Offspring identity was

included as a random effect to restrict comparisons to within offspring and mother identity was

included as a random effect to control for repeated measures of females. When fitting models to

the total location errors, we log-transformed the response variable to improve residual fit and

included a quadratic term of total trials completed during the 4-day cognitive testing period,

since the number of trials completed influences performance (27, 31, 32, 71). We tested whether

extra-pair males were significantly older than the social males they cuckolded using a chi-square

test.

Both male and female cognitive abilities could affect the likelihood of a nest containing

extra-pair young, so we tested whether cuckoldry was associated with the spatial cognitive

abilities of both parents at a nest using multiple methods. First, we modeled the relationship

between male and female spatial cognitive ability and cuckoldry at the nest and offspring level.

At the nest level, we tested whether male or female spatial cognitive ability predicted whether

any offspring in a nest were sired by extra-pair males. At the offspring level, we tested whether

male or female spatial cognitive ability predicted whether each offspring was sired by an extra-

pair male or not. For both tests, we used generalized linear mixed models with binomial

distributions that included male and female spatial cognitive performance, elevation, and year as

fixed effects. Male and female identity were included as random effects to control for repeated

measures of individual males and females. Second, for nests that had at least one extra-pair

young, we tested if the number of extra-pair young in a nest was associated with male or female

spatial cognitive ability using a generalized linear mixed model with a truncated Poisson

distribution. The number of extra-pair young in the nest was included as the response variable,

while male and female performance on the first 20 trials, year, and elevation were included as

fixed effects. Male and female identity were included as random effects to control for repeated

measures of individual males and females. The effect of male age on cuckoldry rates was tested

by adding male age as a fixed effect to each model and comparing models with and without male

age using likelihood ratio tests.

We also tested whether male spatial cognitive ability differed between nests that had EPY

compared to those that did not. Male spatial cognition was included as the response variable and

elevation and whether the nest had any EPY were included as fixed effects. When modeling the

mean number of location errors per trial on the entire task we used the Gamma distribution and

included a quadratic term for the total number of trials completed as a fixed effect. Including

male identity as a random effect in both models led to convergence errors, so male identity was

not included in the models.

We tested whether the mean number of location errors per trial across the first 20 trials

correlated with the mean number of location errors per trial over the entire 4-day task for males

using a linear model. Mean location errors per trial over the entire task was used as the response

variable while mean location errors over the first 20 trials and a quadratic term of the total

number of trials completed were included as fixed effects. All statistical analyses were performed

in R version 4.1.1 (85) and the residual fit of all models was checked using DHARMa (86).

Statistical significance was calculated using Wald Chi-square tests.

Results

Extra-pair paternity and spatial cognition

Across three years of data collection on EPP (2018, 2020, and 2021), we found that 30.3% of the

total offspring sampled (222 of 732) were sired by extra-pair males and that 70.1% of nests (89

of 127) had at least one EPY. We found no significant differences in the number of EPY sired at

high versus low elevation populations regardless of year sampled (Supplemental Results, Figure

S1, Table S1), therefore, we combined high and low elevations for all later analyses. Data from

breeding birds in this study confirms previous research showing that female chickadees obtain

extra-pair copulations with neighboring males (on average ~100 m from their own nests; Figures

S2 and S3), which are likely familiar males from their nonbreeding flock (52, 65, 66). In

addition, breeding male territories do not appear to cluster based on males' cognitive

performance (Figure S4).

Performance on the spatial learning and memory task predicted the number of EPY a

male sired each year (Figure 2). Males with better spatial learning and memory abilities (e.g.,

fewer mean location errors per trial) sired more EPY, both according to a zero-inflated model

including males that sired no extra-pair young (Mean location errors per trial over first 20 trials:

slope = -0.92, χ^2_1=6.28, P=0.012 (Figure 2, dashed line); Mean location errors per trial over the

entire task: slope = -1.86, χ^2_1=8.12, P=0.004, N=280 (137 individual males; Figure S5)), and to a

model only including males that sired at least one EPY (Mean location errors per trial over first

20 trials: slope = -1.16, χ^2_1=5.96, P=0.015 (Figure 2, solid line); Mean location errors per trial

over the entire task: slope = -2.55, χ^2_1=7.68, P=0.006, N=41 (34 individual males; Figure S5)).

Our model suggests that of the males that sired at least one EPY, the males with the best

cognitive performance may sire between 6 and 7 EPY each year, whereas the males with the

worst cognitive performance will sire only 1 to 2 EPY each year. We have shown that birds with

better spatial cognitive abilities live longer (27) and lack reproductive senescence within their

natural life span (87), therefore, males with better cognition are expected to produce significantly

more offspring over their lifetime. Male age was not associated with the number of EPY sired

(see Supplemental Results).

In addition to testing how cognitive abilities were related to number of EPY, we also

tested whether variation in cognitive abilities was associated with breeding performance. Using 9

years of breeding data (2015-2023), we detected no relationship between the social males' spatial

cognitive abilities and brood size (χ^2_1=2.32, P=0.127, N=205 nests), or the females' spatial

cognitive abilities and brood size (χ^2_1=1.75, P=0.186, N=205 nests). Controlling for the effect of

clutch size on brood size produced similar results (Table S4). However, males with better spatial

cognition raised heavier young (mass at day 16, χ^2_1=4.86, P=0.028, N=202 nests), while female

spatial cognition showed no significant relationship with offspring mass at day 16 (χ^2_1=0.93,

P=0.335, N=202 nests). Based on these findings, we used the 3 years of EPP data to indirectly

assess differences in parental care and overall male condition by comparing the brood size and

mean fledgling mass of young from nests of males that sired EPY versus those that did not sire

EPY. Neither brood size (brood size: χ^2_1=0.30, P=0.584, N=144 nests) nor mean nestling mass

(mean mass: χ^2_1=2.10, P=0.147, N=140 nests) differed significantly among males with and

without EPY. Furthermore, we compared the mass of young sired by extra-pair males to those

sired by the social males raised in the same nests and found no significant difference (χ^2_1=2.83,

P=0.093, N=719 offspring).

Figure 2. Model predictions of the number of extra-pair young a male sired in a year by the mean

number of location errors per trial over the first 20 trials. The dashed line represents a zero-

inflated model for all males, including those with no extra-pair young (N=280; 137 individual

males), and the solid line represents the model prediction for males that sired at least one extra-

pair young (N=41; 34 individual males). Points are slightly *jittered* on the y-axis to improve

readability. Shading represents bootstrapped 95% confidence intervals. Inset depicts 8 nestlings

at 16 days post-hatching. See model predictions for the number of extra-pair young a male sired

in a year by total 4-day testing performance in Figure S5.

Extra-pair males have better spatial cognitive abilities than the social males they cuckold

When there were EPY in a nest, they were sired by males with better spatial learning and

memory abilities (on average) compared to the social male of the female (Mean location errors

per trial over first 20 trials: χ^2_1=14.81, P<0.001, N=71 comparisons of a social male and the

extra-pair male sire (Figure 3); Mean location errors per trial over the total task: χ^2_1=6.08,

P=0.014, N=71 comparisons (Figure S6)). Extra-pair males did not differ significantly in age

from the males they cuckolded (28 offspring were sired by an extra-pair male that was older than

the social male, 19 offspring were sired by an extra-pair male that was the same age as the social

male, and 24 offspring were sired by an extra-pair male that was younger than the social male;

Chi-square test: χ^2_2= 1.72, P=0.424).

Figure 3. Comparing spatial cognitive performance (mean number of location errors per trial

over the first 20 trials) of extra-pair (mean $\pm$ SE = 0.96 $\pm$ 0.07) and social fathers (mean $\pm$ SE =

1.17 $\pm$ 0.07, 71 comparisons). Lines connect males by offspring: blue lines show comparisons

where the extra-pair father performed better than the social father (49 of 71 cases or 69%), and

purple lines show comparisons where the social father performed better than the extra-pair father

(19 of 71 cases, or 26.7%; in 3 cases the extra-pair and social father had the same cognition

score). Shaded distributions represent smoothed histograms of cognitive performance for each

father type. The same analysis is presented for performance over the 4-day testing period in

Figure S6.

The finding that extra-pair males have better spatial cognition than the social male of the

nest (on average) could be explained by the cognitive abilities of the social male or female, as

either sex may have cognitive deficiencies (i.e., females seek EPC to compensate for her or her

social male's poor cognition) or be better at obtaining EPC. Therefore, we assessed whether the

likelihood of EPY in the nest was related to the spatial cognitive abilities of the social male or

the female. The likelihood of being cuckolded was not significantly related to the cognitive

abilities of the social males (Offspring level: Mean location errors per trial over first 20:

χ^2_1=0.25, P=0.620, Mean location errors per trial over the entire task: χ^2_1=0.43, P=0.513, N=348

offspring for 42 males; Nest level: Mean location errors per trial over first 20: χ^2_1=0.05, P=0.808;

Mean location errors per trial over the entire task: χ^2_1=0.22, P=0.642, N=59 nests for 42 males).

In addition, male age did not significantly affect the likelihood of a male being cuckolded (see

Supplemental Results). However, females who performed worse on the first 20 trials of the

spatial cognitive task were more likely to have EPY in their nests (Offspring level: Mean

location errors per trial over first 20: χ^2_1=6.21, P=0.013, N=348 offspring for 42 females);

though this relationship was not significant when we analyzed cognitive performance across the

entire 4-day task (χ^2_1=0.18, P=0.673, N=348 offspring for 42 females), or at the nest level of

analysis (Mean location errors per trial over first 20: χ^2_1=0.21, P=0.645, Mean location errors per

trial over the entire task: χ^2_1=0.13, P=0.723, N=59 nests for 42 females).

495   Similarly, for nests that had at least one EPY, the cognitive abilities of the social male did

not predict the number of EPY in their nests (χ^2_1=1.03, P=0.311, N=41 nests for 35 males),

however, females that performed worse on the task had significantly more EPY in their nests

(χ^2_1=6.79, P=0.009, N=41 nests for 32 females). In other words, females would not have sought

EPCs solely in response to their social mate's poor cognition. Instead, the data suggest that

females may engage in EPCs based on their own cognitive deficiencies, and when doing so, may

seek males with better spatial cognitive abilities than their social mate. As a result, males with

better spatial cognitive abilities exhibit higher fitness; they did not lose paternity in their own

nests (i.e., male spatial cognition does not predict if a male will be cuckolded or how many EPY

are in the nest), they produced similar quality of offspring compared to males with worse

cognitive abilities (e.g., similar brood size and fledgling mass), and gained more EPY overall.

Discussion

Our findings show that spatial cognitive abilities influence extra-pair paternity in free-living,

food-caching mountain chickadees. Males with better spatial learning and memory abilities sire

more EPY than their poorer performing counterparts and, more importantly, EP males have better

spatial cognitive abilities than the social males they cuckold, regardless of age. While we are

unable to rule out effects of sperm competition, the direction of our findings suggest EPP may be

driven by females foraying into male territories, as opposed to male intruders, for several

reasons; 1. Female chickadees (and Passerines more broadly) are known to pursue EPCs with

neighboring males (41, 50, 51, 52, 53, 88, 89), 2. Chickadees have never been reported or

observed engaging in forced copulation (42, 52, 53), and 3. Resident males are highly territorial

and do not tolerate male conspecifics in their territories. In addition, the directional relationship

between the cognitive phenotype of the social and extra-pair male would not be predicted by

male choice but is consistent with good genes hypotheses (1, 8, 14).

If EPCs were driven by male forays, we would expect EPY to be present in nests

regardless of the cognitive abilities of the social male, as there is no clear biological reason for a

male seeking EPCs to selectively target females mated to males with worse cognitive abilities.

That said, it is possible that males with poor spatial cognitive abilities are worse at defending

their territories from other males, which may allow those males with better cognitive abilities to

intrude and copulate with the resident female. We think this explanation is unlikely as males with

worse cognitive abilities tend to be more proactively aggressive (90) and cognitive performance

is not associated with social dominance (91). However, females, would benefit from obtaining

EPCs with males that have better spatial cognition than their social male, as this trait is heritable

(29, 30) and would increase the likelihood of producing offspring with better spatial cognition,

leading to higher overwinter survival (27) and longer lifespan (28). Finally, our finding that the

EP males have better spatial cognitive abilities compared to the social males they cuckhold is

consistent with the good genes hypothesis, that females engage in EPCs with higher quality

males to obtain indirect benefits for their offspring (1, 8, 14). Comparing phenotypic quality of

the EP male to that of the social male they cuckhold has been one of the main methods to test for

the good gene hypothesis. Considering that alleles associated with better spatial cognitive

abilities are not associated with sex (30), producing EPY with better quality males increases

selection pressure on spatial cognitive abilities in both males and females, as both sexes rely on

spatial cognition for survival (92).

Although it remains unknown whether chickadees can evaluate cognitive abilities directly

or indirectly via some correlated trait(s), the life history and breeding ecology of mountain

chickadees provide several possibilities. For example, female chickadees may use secondary

sexual signals, such as song and plumage variation associated with spatial cognitive abilities, to

identify high-performing males (1, 48, 93). Recent work in closely related black-capped

chickadees shows that male cognitive performance is associated with variation in plumage patch

brightness and contrast (93). Alternatively, it is possible that females can directly assess a male's

ability to successfully recover food stores over winter and base mating decisions on this

information in the spring (49, 94). Chickadees are nonmigratory, sedentary birds that aggregate

in flocks of unrelated male-female pairs throughout the nonbreeding season and break apart once

breeding commences (65, 94). Data from this study confirm previous research showing that

female chickadees seek EPCs with neighboring males (15, 52, 53), which would be familiar

males from their winter flock (65, 94). Regardless of how females evaluate male cognition,

mating with males with better cognitive performance should increase female fitness by

increasing the likelihood of offspring inheriting the specific alleles associated with better

cognition, which would be expected to lead to higher offspring survival and longer lifespan (27,

28).

An alternative hypothesis for our findings is that females seek EPCs with other males as

insurance against the social male's infertility (15, 22), again gaining genetic benefits. However,

this hypothesis does not explain the direction of our findings, because male cognitive ability did

not predict the number of offspring a male produced in his own nest (i.e., most cuckolded social

males still fathered chicks in their own nests), suggesting that males with poorer cognition do not

suffer from infertility. That said, we do not have data on the sperm quality of these birds and

therefore cannot directly exclude differences in fertility or sperm competition between males

with better and worse spatial cognition. Related, we have not detected fitness trade-offs or

negative effects associated with better cognitive abilities in our system. We do not see cognitive

or reproductive senescence within birds' naturally occurring life span (31, 87). In fact,

chickadees with better spatial cognition live longer allowing them to produce more offspring

over their lifetime compared to chickadees with worse cognition (28).

Based on our research to date, male chickadees with better spatial cognitive abilities have

higher reproductive success than their poorer performing counterparts: these males produce more

EPY (without losing paternity in their own nests) and raise heavier young, which are more likely

to survive and be recruited into the population (84). Females can potentially gain indirect and

direct benefits from reproducing with males exhibiting better spatial cognition. Producing EPY

with males that have better cognition increases the likelihood that their young inherit good genes

and if their social male has better cognition, they benefit directly by raising heavier chicks. These

fitness benefits would be expected to increase selection on any mechanism allowing

discrimination based on cognition.

Interestingly, a previous study in our system showed that females mated to social males

with better spatial abilities lay larger clutches and fledge larger broods in years of high resource

abundance (39). Together with the present findings, we suggest that female mating decisions,

including EPCs, and reproductive investment may vary opportunistically. When resources are

abundant, females lay larger clutches if their social male has better cognitive abilities (39), and

when males with better spatial cognition than her social mate are available for EPCs, they pursue

those opportunities. Furthermore, we found that females with poorer spatial cognition had more

EPY in their nests, suggesting females' mating decisions could be based on some physiological

signal associated with poorer cognition. At the ultimate level, females with poorer cognition

likely benefit disproportionately by mating with the highest quality male they can find, which in

the case of food-caching birds, means those with better spatial cognitive abilities (27, 28).

An additional alternative explanation for our findings is that birds with better spatial

cognitive abilities have access to better territories and therefore are clustered based on cognitive

abilities. Then, because females have extra-pair copulations with neighboring males (15, 52, 53),

those neighbors would happen to be males with better spatial cognition. However, in our study

system, males do not appear to cluster based on cognitive performance (Figure S4), suggesting

that females are not finding males with better cognitive abilities than their social males by

chance. Furthermore, higher social dominance status is associated with access to better territories

in chickadees (95), but we see no association with dominance status and spatial cognitive

abilities in our system (91) or closely related black-capped chickadees (93). While it is possible

that our results reflect some unmeasured trait(s) linked to spatial cognitive ability, such as

boldness, aggression, or increased exploratory behavior (e.g., 46), previous data from our system

do not support these additional alternative hypotheses. Chickadees with enhanced spatial

cognitive abilities are not more proactively aggressive (90) nor do they show higher rates of

exploratory behavior (e.g., higher space use) (96).

Given the fitness advantages of superior spatial cognitive abilities for food-caching

chickadees, how variation is maintained and why enhanced spatial cognitive abilities have not

evolve to fixation may seem puzzling. While food-caching birds exhibit superior spatial

cognitive abilities compared to their non-caching counterparts (87), considerable variation

persists within species (71). Research from our system demonstrates the highly polygenic nature

of the spatial cognitive phenotype, meaning there are multiple genetic pathways that may

produce the same cognitive phenotype (29, 30, 98). In addition, selection strength varies with

annual variation in winter climate, which in milder years, may allow individuals to survive and

reproduce despite comparatively poor spatial cognitive abilities (27, 63, 70, 71). Finally, we have

documented elevation-related differences in spatial cognition and the hippocampus of mountain

chickadees (71, 72), but there is gene flow among elevations (99) which moves allelic variation

associated with poorer cognition within the population.

The results presented here are an exciting contribution to the field because: 1. We show

that males with better spatial cognition exhibit higher reproductive output, via siring more EPY

and fledging heavier chicks which are more likely to survive and be recruited into the breeding

population, and 2. We show that when there are EPY in the nest, they are sired by males with

better spatial cognition than the social male (on average), suggesting these mating decisions may

be driven by females. However, we cannot completely rule out alternative explanations related to

sperm competition. Regardless of whether these patterns are driven by females or males, males

with better spatial cognitive abilities exhibit higher fitness by producing more offspring, adding

to the evidence for selection on spatial cognitive abilities, including differential survival and

longevity in food-caching birds (27, 28).

Acknowledgements

Thank you to Qwahn Kent, Danielle Schaening-Lopez, Alyssa Nowicki, and Beck Kerdman for

assistance with DNA extraction. We thank two anonymous reviewers for their comments and

suggestions which greatly improved the strength of our findings and clarity of our manuscript.

All procedures were in accordance with University of Nevada Reno IACUC (protocols 20-11-

1103-1, 20-06-1014, 20-08-1062) and California State Fish and Wildlife Scientific Collecting

Permit (S-193630001-20007-001).

Funding

Kessel Fellowship from American Ornithological Society (CLB)

Edward Rose Fellowship from Cornell Lab of Ornithology (CLB)

National Science Foundation Graduate Research Fellowship Program, 2019287870 (BRS)

National Science Foundation Graduate Research Fellowship Program, 2020305313 (LMB)

National Science Foundation IOS1856181 (VVP)

National Science Foundation IOS2119824 (VVP)

Author contributions

Conceptualization: CLB, VVP

Methodology: CLB, BRS, JFW, BGB, VKH, AMP, LMB, ESB, MSW, IJL, VVP

Investigation: CLB, BRS, VKH, AMP, LMB, VVP

Visualization: CLB, JFW, VVP

Funding acquisition: CLB, VVP

Writing – original draft: CLB

Writing – review & editing: CLB, BRS, JFW, BGB, VKH, AMP, LMB, ESB, MSW, IJL, VVP

Declaration of Interests: Authors declare that they have no competing interests.

Data Availability: All data, code, and materials used in the analysis are available at

https://figshare.com/s/ef7f40371b487b9dabca.

REFERENCES

1. Andersson, M & Simmons, LW. Sexual selection and mate choice. *Trends Ecol. Evol*. 21,

296–302 (2006).

2. Rosenthal, G. *Mate Choice: the Evolution of Sexual Decision Making from Microbes to*

Humans. Princeton, NJ, Princeton University Press (2017).

3. Davies, AD, Lewis, Z, & Dougherty LR. A meta-analysis of factors influencing the

strength of mate-choice copying in animals. *Behav. Ecol.* 31, 1279-1290 (2020).

4. Henshaw, JM, Fromhage, L, & Jones, AG. The evolution of mating preferences for

genetic attractiveness and quality in the presence of sensory bias. *PNAS* 119,

e2206262119 (2022).

5. DuVal, EH, Fitzpatrick, CL, Hobson, EA, & Servedio, MR. Inferred attractiveness: A

generalized mechanism for sexual selection that can maintain variation in traits and

preferences over time. *PLoS Biol.* 21, e3002269 (2023).

6. Fisher, RA. *The genetical theory of natural selection.* Oxford, Clarendon Press (1930).

7. Endler, JA. Sensory drive: does sensory biology bias or constrain the direction of

evolution? *Am. Nat.* 139, S1-S153 (1992).

8. Andersson, M. *Sexual Selection*. Princeton, NJ, Princeton University Press (1994).

9. Griffith SC, Owens IPF, & Thuman KA. Extra pair paternity in birds: a review of

interspecific variation and adaptive function. *Mol Ecol*. 11:2195–2212 (2002).

10. Lambert CT, Sabol AC, & Solomon NG. Genetic monogamy in socially monogamous

mammals is primarily predicted by multiple life history factors: a meta-analysis. *Front*

Ecol Evol., 6:139 (2018).

11. Brouwer L, & Griffith SC. Extra-pair paternity in birds. *Molecular Ecology,* 28, 4864-

4882 (2019).

12. Webster, MS, Tarvin, KA, Tuttle, EM, & Pruett-Jones, S. Promiscuity drives sexual

selection in a socially monogamous bird. *Evolution*, *61*, 2205-2211. (2007).

13. Jennions, MD, & Petrie, M. Why do females mate multiply? A review of the genetic

benefits. *Biol. Rev*. 75, 21-64. (2000).

14. Herdtle A, Duncan C, Manser MB, & Clutton-Brock T. Patterns and drivers of female

extra-pair mating in wild Kalahari meerkats. *Behavioral Ecology*, 36: 3 (2025).

15. Lifjeld JT, Kleven O, Amundsen T, & Slagsvold T. Extrapair copulations in coal tits

(*Periparus ater*): Female insurance against male fertility. *Beh. Ecol. Sociobiol.* 79:79.

(2025).

16. Petrie M, Kempenaers B. Extra-pair paternity in birds: explaining variation between

species and populations. *TREE*, 13, 52-58 (1998).

17. Tregenza T, & Wedell N. Genetic compatibility, mate choice, and patterns of parentage:

Invited review. *Molecular Ecology*, 9,1013-1027 (2000).

18. Arnqvist G, & Kirkpatrick M. The evolution of infidelity in socially monogamous

passerines: the strength of direct and indirect selection on extrapair copulation behavior

in females. *American Naturalist*, 165: S26-S37. (2005).

19. Neff BD, & Pitcher TE. Genetic quality and sexual selection: an integrated framework for

good genes and compatible genes. *Mol Ecol.,* 14:19–38. (2005).

20. Akçay E, & Roughgarden J. Extra-pair paternity in birds: review of the genetic benefits.

Evol Ecol Res. 9:855–868. (2007).

21. Forstmeier W, Nakagawa S, Griffith SC, & Kempenaers B. Female extra-pair mating:

adaptation or genetic constraint? *TREE*, 29: 456-464 (2014).

22. Yasui Y, & Yoshimura J. Bet-hedging against male-caused reproductive failures may

explain ubiquitous cuckoldry in female birds. *J Theor Biol*. 437:214–221. (2018).

23. Westneat DF, & Stewart IRK. Extra-pair paternity in birds: causes, correlates, and

conflict. *Annu Rev Ecol Evol Syst*. 34:365–396. (2003).

24. Vander Wall SB. Food hoarding in animals. University of Chicago Press, Chicago.

(1990).

25. Pravosudov VV, & Roth II TC. Cognitive ecology of food hoarding: the evolution of

spatial memory and the hippocampus. *Ann. Rev. Ecol. Evol. System*. 44, 173-193 (2013).

26. Brodin, A. Mechanisms of cache retrieval in long-term hoarding birds. *J Ethol* 23, 77–83

(2005).

27. Sonnenberg, BR, Branch, CL, Pitera, AM, Bridge, E, & Pravosudov, VV. Natural

selection and spatial cognition in wild food-caching mountain chickadees. *Curr. Biol.* 29,

670-676 (2019).

28. Welklin, JF *et al.* Spatial cognitive ability is associated with longevity in a food-caching

bird*. Science*, 385, 1111-1115 (2024).

29. Branch, CL *et al*. The genetic basis of spatial cognitive variation in a food-caching

bird. *Curr. Biol.* 32, 210-219 (2022).

30. Semenov, G *et al.* Genes and gene networks underlying spatial cognition in food-caching

chickadees. *Curr. Biol.* 34, 1930-1939 (2024).

31. Heinen, VK *et al.* Specialized spatial cognition is associated with reduced cognitive

senescence in a food-caching bird. *Proc. R. Soc. B*. 288, 20203180 (2021).

32. Tello-Ramos, MC *et al.* Memory in wild mountain chickadees from different elevations:

comparing first-year birds with older survivors. *Anim. Behav.* 137, 149-160 (2018).

33. Keagy J, Savard JF, & Borgia G. Male satin bowerbird problem-solving ability predicts

mating success. *Anim. Behav.* 78, 809-817 (2009).

34. Medina-Garcia A, & Wright TF. An integrative measure of cognitive performance, but

not individual task performance, is linked to male reproductive output in

budgerigars. *Scientific Reports, 11*, 11775. (2021).

35. Preiszner B, Papp S, Pipoly I, Seress G, Vincze E, Liker A, & Bokony V. Problem-

solving performance and reproductive success of great tits in urban and forest

habitats. *Animal Cognition, 20*, 53-63. (2017).

36. Shaw RC, MacKinlay RD, Clayton NS, & Burns KC. Memory performance influences

male reproductive success in a wild bird. *Curr. Biol.* 29, 1498-1502 (2019).

37. White DJ, Arthur J, Davies HB, Guigueno MF. Cognition and reproductive success in

cowbirds. *Learn Behav* 50, 178–188 (2022).

38. van Horik JO, Madden JR. A problem with problem solving: motivational traits, but not

cognition, predict success on novel operant foraging tasks. *Anim Behav*, 114, 189-198

(2016).

39. Branch CL, Pitera AM, Kozlovsky DY, Bridge ES, & Pravosudov VV. Smart is the new

sexy: female mountain chickadees increase reproductive investment when mated to males

with better spatial cognition. *Ecol. Lett.* 22, 897-903 (2019).

40. Cockburn A. Prevalence of different modes of parental care in birds. *Proceedings of the*

Royal Society B: Biological Sciences, *273*, 1375–1383. (2006).

41. Gray EM. Female red-winged blackbirds accrue material benefits from copulating with

extra-pair males. *Animal Behaviour*, *53*, 625-639. (1997).

42. Mennill DJ, Ramsay SM, Boag PT, & Ratcliffe LM. Patterns of extrapair mating in

relation to male dominance status and female nest placement in black-capped

chickadees. *Behavioral Ecology*, *15*, 757-765. (2004).

43. Ophir AG, Phelps SM, Sorin AB, Wolff JO. Social but not genetic monogamy is

associated with greater breeding success in prairie voles. *Animal Behaviour,* 75, 1143-

1154. (2008).

44. Canal D, Jovani R, & Potti J. Male decisions or female accessibility? Spatiotemporal

patterns of extra pair paternity in a songbird. *Behavioral Ecology*, *23*, 1146-1153. (2012).

45. Spritzer MD, Meikle DB, & Solomon NG. Female choice based on male spatial ability

and aggressiveness among meadow voles. *Anim Behav.* 69, 1121–1130 (2005).

46. Shohet AJ, & Watt PJ. Female guppies *Poecilia reticulata* prefer males that can learn fast.

J Fish Biol. 75, 1123 – 1523 (2009).

47. Snowberg LK, & Benkman CW. Mate choice based on a key ecological performance

trait. *J. Evol. Biol.* 22, 762-769 (2009).

48. Boogert NJ, Fawcett TW, Lefebvre L. Mate choice for cognitive traits: a review of the

evidence in nonhuman vertebrates, *Behav. Ecol.* 22, 447–459 (2011).

49. Chen J, Zou,Y, Sun Y-H, & Ten Cate C. Problem-solving males become more attractive

to female budgerigars. *Science* 363, 166 –167 (2019).

50. Sheldon BC. Male phenotype, fertility, and the pursuit of extra-pair copulations by

female birds. *Proc. R. Soc. B*. 257, 25–30 (1994).

51. Double M, & Cockburn A. Pre-dawn infidelity: females control extra-pair mating in

superb fairy-wrens. *Proc Biol Sci.* 267, 465-70 (2000).

52. Smith SM. Extra-pair copulations in black-capped chickadees: the role of the female.

Behav. 107, 15-23. (1988).

53. Otter K, Ratcliffe L, & Boag P. Extra-pair paternity in the black-capped chickadee.

Condor 96, 218 – 222 (1994).

54. Poesel A, Kunc, HP, Foerster K, JohnsenA, & Kempenaers B. Early birds are sexy: Male

age, dawn song and extrapair paternity in blue tits, *Cyanistes* (formerly *Parus*) *caeruleus*.

Animal Behavior, 72, 531–538. (2006).

55. Kempenaers B, Borgström P, Loës P, Schlicht E, & Valcu M. Artificial night lighting

affects dawn song, extra-pair siring success, and lay date in songbirds. *Current Biology*,

20, 1735–1739. (2010).

56. Bonderud ES, Otter KA, Burg TM, Marini KLD, Reudink MW. Patterns of extra-pair

paternity in mountain chickadees. *Ethology*. 124: 378–386. (2018).

57. Hinde RA. The Conflict between Drives in the Courtship and Copulation of the

Chaffinch. *Behaviour*, *5*, 1–31. (1953).

58. McKinney F, & Evarts S. Sexual Coercion in Waterfowl and Other Birds. *Ornithological*

Monographs, *49*, 163–195. (1998).

59. Hooper, R, Maher, K, Moore, K, McIvor, G, Hosken, D, Thornton, A. Ultimate drivers of

forced extra-pair copulations in birds lacking a penis: jackdaws as a case-study. *R. Soc.*

Open Sci. 11, 231226 (2024).

60. Perrins CM, & Moss D. Reproductive Rates in the Great Tit. *Journal of Animal Ecology*, *44*,

695–706. (1975).

61. Sanz JJ. Effects of geographic location and habitat on breeding parameters of Great

Tits, *The Auk*, 115, 1034-1051 (1998).

62. Kirwan GM, Sly ND, Gosler A, Clement P, Christie, DA, Moura N, & Pyle P. Great

Tit (*Parus major*), version 2.0. In Birds of the World (G. M. Kirwan and N. D. Sly,

Editors). Cornell Lab of Ornithology, Ithaca, NY, USA. (2024).

63. Whitenack LE, *et al.* Complex relationships between climate and reproduction in a

resident montane bird. *R. Soc. Open Sci.* 10, 230554 (2023).

64. Branch CL, *et al.* What's in a mate? Social pairing decisions and cognitive performance

in food-caching mountain chickadees. *Proc. R. Soc. B.* 290, 20231073 (2023).

65. Ekman J. Ecology of non-breeding social systems of Parus. *Wil. Bull.* 101, 263–288

(1989).

66. McCallum D, Grundel R, & Dahlsten DL. Mountain Chickadee (*Poecile gambeli*),

version 1.0. In Birds of the World (A. F. Poole and F. B. Gill, Editors). Cornell Lab of

Ornithology, Ithaca, NY, USA (2020).

67. Pitera AM, Branch CL, Sonnenberg BR, Benedict LM, Kozlovsky DY, & Pravosudov

VV. Reproduction is affected by individual breeding experience but not pair longevity in

a socially monogamous bird. *Behav. Ecol. Sociobiol*. 75, 101 (2021).

68. Welklin JF, Sonnenberg BR, Branch CL, Pitera AM, Whitenack LE, Benedict LM,

Heinen VK, & Pravosudov VV. Social pairing in the absence of reproductive senescence

in a socially monogamous songbird. *Behavioral Ecology*, 36, p.araf083. (2025).

69. Hsu YH, Schroeder J, Winney I, Burke T & Nakagawa S. Are extra-pair males different

from cuckolded males? A case study and a meta-analytic examination. *Mol. Ecol.*, *24*,

1558-1571 (2015).

70. Kozlovsky DY, Branch CL, Pitera AM, Croston R, & Pravosudov VV. Fluctuations in

annual climatic extremes are associated with reproductive variation in resident mountain

chickadees. *R Soc. Open Sci.* 5, 171604 (2018).

71. Croston R *et al.* Individual variation in spatial memory performance in wild mountain

chickadees from different elevations. *Anim. Behav.* 111, 225-234 (2016).

72. Freas CA, LaDage LD, Roth II TC, & Pravosudov VV. Elevation-related differences in

memory and the hippocampus in mountain chickadees, *Poecile gambeli. Anim. Behav.*

84*, 121-127 (2012).

73. Benedict LM *et al.* Information maintenance of food sources is associated with

environment, spatial cognition and age in a food-caching bird. *Anim. Behav.* 182, 153-

172 (2021).

74. Benedict LM, Heinen VK, Sonnenberg BR, Bridge ES, & Pravosudov VV. Learning

predictably alternating food locations across days in a food-caching bird. *Anim. Behav.*

196, 55-81 (2023).

75. Pyle P. Identification Guide to North American Birds, Part I. Second Edition. Sheridan

Books, Ann Arbor, MI, USA. (2022).

76. Thrasher DJ, Butcher BG, Campagna L, Webster MS, & Lovette IJ. Double-digest RAD

sequencing outperforms microsatellite loci at assigning paternity and estimating

relatedness: A proof of concept in a highly promiscuous bird. *Mol. Ecol. Res.* 18, 953-965

(2018).

77. Rohland N, & Reich D. Cost-effective, high-throughput DNA sequencing libraries for

multiplexed target capture. *Genome Res.* 22, 939-46 (2012).

78. Wigginton JE, Cutler DJ, & Abecasis GR. A note on exact tests of Hardy-Weinberg

equilibrium. *Am. J Human Genetics* 76, 887-893 (2005).

79. Danecek P, *et al.* The variant call format and VCFtools. *Bioinformatics* 27, 2156–2158

(2011).

80. Gosselin T. radiator: RADseq Data Exploration, Manipulation and Visualization using R.

R package version 1.1.9 https://thierrygosselin.github.io/radiator/ (2020).

81. Kalinowski S, Taper M, & Marshall T. Revising how the computer program CERVUS

accommodates genotyping error increases success in paternity assignment. *Mol Ecol.* 16,

1099-1106 (2007).

82. Brooks ME, *et al.* glmmTMB balances speed and flexibility among packages for zero-

inflated generalized linear mixed modeling. *R J.* 9, 378–400 (2017).

83. Joe H, & Zhu R. Generalized Poisson distribution: the property of mixture of Poisson and

comparison with negative binomial distribution. *Biometrical Journal: J. Math. Methods*

in Biosci. 47, 219-229 (2005).

84. Whitenack LE, Richmond AA, Sonnenberg BR, Welklin JF, Heinen VK, Pitera AM,

Haley SM, Branch CL, Pravosudov VV. Postnatal dispersal and drivers of successful

recruitment in resident Mountain Chickadees (*Poecile gambeli*). *Ornithology*, ukaf006
(2025).

85. R Core Team. R: A language and environment for statistical computing (2021).

86. Hartig F. DHARMa: Residual diagnostics for hierarchical (multi-level/mixed) regression
models R package Version 0.4.4. (2018).

87. Welklin JF *et al.* No evidence of reproductive senescence within natural life span in
resident mountain chickadees (*Poecile gambeli*). *Behav. Ecol. Sociobiol.* 79, 10 (2025).

88. Hasselquist D, & Sherman PW. Social mating systems and extrapair fertilizations in
passerine birds. *Behav. Ecol.* 12, 457–466. (2001).

89. Kempenaers B, Verheyen GR, Vanderbroeck M, Burke T, Vanbroeckoven C, & Dhondt
AA. Extra-pair paternity results from female preference for high-quality males in the blue
tit. *Nature,* 357, 494–96 (1992).

90. Branch CL, Kozlovsky DY, & Pravosudov VV. Elevation related variation in aggressive
response to mirror image in mountain chickadees. *Behav.* 152, 667-676 (2015).

91. Heinen VK *et al.* Social dominance has limited effects on spatial cognition in a wild
food-caching birds. *Proc. R. Soc. B.* 288, 20211784 (2021).

92. Branch CL, *et al.* Testing the greater male variability phenomenon: male mountain
chickadees exhibit larger variation in reversal learning performance compared to females.
Proc. R. Soc. B. 287, 20200895 (2020).

93. Robayo Noguera L, Stevenson CAL, Wang T, Pasquale MK, Branch, CL. Variation in
plumage reflectance but not song reflects spatial cognitive performance in black-capped
chickadees *(Poecile atricapillus)*. *Anim. Cog.* 28, 14 (2025).

94. Lemmon D, Withiam ML, Barkan CPL. Mate protection and winter pair-bonds in black-

capped chickadees. *Condor* 99, 424–433 (1997).

95. Ratcliffe L, Mennill DJ, & Schubert KA. Social dominance and fitness in black-capped

chickadees in *Ecology and Behaviour of Chickadees and Titmice: An integrative*

approach. Oxford, UK, Oxford University Press (2007).

96. Kozlovsky DY, Branch CL, Freas CA, & Pravosudov VV. Elevation-related

differences in exploration and social dominance in mountain chickadees (*Poecile*

gambeli). *Behav. Ecol. Sociobiol.* 68, 1871-1881 (2014).

97. Brodbeck DR, & Shettleworth SJ. Matching location and color of a compound

stimulus: comparison of a food-storing and nonstoring bird species. *J. Exp.Psychol.* 21,

64-77 (1995).

98. Pritchard JK, Pickrell JK, & Coop G. The genetics of human adaptation: hard

sweeps, soft sweeps, and polygenic adaptation. *Curr. Biol.* 23, R208-R215 (2010).

99. Branch CL, Jahner JP, Kozlovsky DY, Parchman TL, & Pravosudov VV. Absence of

population structure across elevational gradients despite large phenotypic variation in

mountain chickadees *(Poecile gambeli). R. Soc. Open Sci.* 4, 170057 (2017).

100. Sonnenberg BR, *et al.* Natural variation in developmental condition has limited effect

on spatial cognition in a wild food-caching bird. *Proc. R. Soc. B.* 289, 1984 (2022).

101. Sonnenberg, BR *et al.* Long-term winter food supplementation in a montane

environment shows no significant impact on reproductive success in a resident food-

caching bird. *Ornithology* 140, 1-12 (2022).

102. Branch CL *et al.,* Code and data for: Seeking smarts: Male chickadees with better

spatial cognition sire more extra-pair young. Figshare (2024);

https://figshare.com/s/ef7f40371b487b9dabca.

914
915