Title: Male chickadees with better spatial cognition sire more extra-pair young

Corresponding author: Carrie L Branch, cbranch6@uwo.ca

Authors: Carrie L Branch*[1], Benjamin R Sonnenberg[2,3], Joseph F Welklin[2], Bronwyn G

Butcher[4], Virginia K Heinen[2], Angela M Pitera[2], Lauren M Benedict[2], Eli S Bridge[5], Irby J

Lovette[4], Michael S Webster[4], Vladimir V Pravosudov[2]

Affiliations:

[1] Department of Psychology, The University of Western Ontario, 1151 Richmond St London,

ON, N6A 3K7 Canada

[2] Department of Biology, University of Nevada, Reno, 1664 N Virginia St Reno, NV, 89557 USA

[3] Department of Integrative Biology, Oregon State University, 1500 SW Jefferson Way,

Corvallis, OR, 97331

[4] Cornell Lab of Ornithology, Cornell University, 159 Sapsucker Woods Rd, Ithaca, NY, 14850

USA

[5] Department of Biology, University of Oklahoma, 660 Parrington Oval, Norman, OK, 73019

USA

Abstract:

Across animal taxa, females commonly mate with more than one male, even in monogamous

mating systems. These extra-pair (EP) copulations and resulting young may increase the fitness

of the female via a variety of mechanisms, including genetic benefits. North American

chickadees provide an interesting system to study the role of sexual selection via EP paternity,

because they are socially monogamous, nonmigratory birds that rely on spatial cognition to

recover food stores and variation in spatial cognition is associated with increased survival, longer

lifespan, and is heritable. Given spatial cognitive abilities are heritable and associated with direct

survival benefits, these abilities may be under sexual selection if males with better spatial

abilities sire more offspring and females prefer to mate with such males. We aimed to address

these predictions by quantifying extra-pair paternity and comparing spatial abilities of EP males

to those of the social male they cuckold in a wild population of mountain chickadees (*Poecile*

gambeli). We found that 1. males with better spatial cognitive abilities have more EP young and

produce heavier offspring in their own nests compared to their poorer performing counterparts,

and 2. EP males have significantly better spatial cognition than the social males they cuckolded.

These results suggest that sexual selection is involved in the evolution of spatial cognitive

abilities in food-caching chickadees and are consistent with the good genes hypothesis, which

posits that females gain indirect genetic benefits via EP young.

Key words: extra-pair mating, spatial cognition, indirect fitness, genetic benefits, sexual

selection

Introduction

Introduction

Understanding the factors that influence mate choice in animals and the consequences

that follow from these decisions is a central focus of evolutionary biology (1-5). Numerous

hypotheses have been generated to explain patterns of sexual selection, including those

associated with the evolution of seemingly arbitrary attractiveness (e.g., runaway selection (6,7))

or sensory exploitation (e.g., sensory drive (8)), and those at the intersection of sexual and

natural selection, when sexually selected traits have direct fitness consequences (e.g., good genes

(9)). Although it has been recognized for some time that males can be actively involved in mate

choice (10,11), considerable work shows that females tend to be the choosier sex, as females

often invest more in gamete production and parental care (12-15). Multiple enduring hypotheses

argue that females should prefer males of 'high-quality', frequently indicated by secondary

sexual signals including ornaments and courtship displays (9,14,16).

In the literature, 'high-quality' refers to everything from sperm competition (17) and

parasite load (18), to direct benefits such as parental care, nuptial gifts, or territory quality (19-

21), to indirect genetic benefits (e.g., 'good genes') that offspring are likely to inherit

(9,16,22,23). Untangling these potential explanations requires an understanding of the fitness-

related traits of interest, including the direction and type of fitness consequences (e.g., survival or

reproduction), as well as whether variation in the trait is genetically heritable, so that the 'high-

quality' of a parent can be inherited by the offspring (24). Identifying and quantifying fitness-

related traits in the wild that are associated with sexual selection and mate choice can be

challenging especially for complex behaviors, like cognition (25).

Cognitive abilities, including associative learning and memory, allow animals to succeed

across variable environments, providing predictable variation or a buffer against unpredictable

availability of resources (26,27). If variation in cognitive abilities is associated with differential

fitness outcomes (survival, reproduction, or both), and this variation is heritable, natural selection

can shape cognitive variation (24). Furthermore, when heritable cognitive variation directly

confers a fitness advantage, it may be associated with mate preference via secondary sexual

signals or by direct observation of mate behavior (28-32). For example, females of several

species show preferences for males with better cognition in lab-based, pairwise choice paradigms

(30-32), while in the wild, males with better cognition have been shown to produce more young

(33,34). However, there is scant evidence from the wild that females prefer males with better

cognition (28), and little evidence outside of model organisms that shows heritability of

cognitive abilities (35).

In the current study, we investigated the potential role of sexual selection on the evolution

of spatial cognitive abilities by examining extra-pair paternity in a wild population of socially

monogamous mountain chickadees (*Poecile gambeli*) over three breeding seasons. Mountain

chickadees are nonmigratory, food caching birds that rely on specialized spatial cognition to

recover thousands of scattered food stores throughout their territories to survive harsh montane

winters (36,37). Following the breeding season, in the temperate autumn, juvenile chickadees are

recruited into flocks of unrelated male-female pairs (36). Breeding pairs from the recent breeding

season typically stay together, providing little opportunity for social mate choice at this time,

however, being paired within the flock is critical because it increases overwinter survival,

particularly for females (38, 39). Work in our long-term study system using 'smart' feeder arrays

(Figure 1) shows that individual variation in spatial learning and memory abilities confers a

fitness advantage in wild chickadees: individuals with better spatial cognitive abilities are more

likely to survive the winter and live longer compared to those with worse abilities (40, 41) and in

resource abundant years, females lay larger clutches and fledge larger broods when their social

male exhibits better spatial learning and memory abilities (42). Furthermore, using genome-wide

association and gene network approaches, we have shown that individual variation in spatial

cognitive abilities is associated with genetic differences, including genes involved in the

development and function of the hippocampus, a brain region involved in spatial cognitive

function (43, 44). Finally, we have shown that spatial cognitive abilities remain stable across

birds' natural lifespan (e.g., no detectable senescence (45) and does not change with age (40,

41)). Due to their life-history and reliance on specialized spatial cognition, chickadees provide a

great system for studying the role of sexual selection on a cognitive trait in the wild.

Given that individual variation in spatial cognitive abilities in chickadees is genetically

heritable and better cognitive abilities lead to increased overwinter survival and longevity,

specifically (i.e., caches are not used for provisioning young), we predict that spatial cognition

should be an important target of sexual selection via extra-pair mating. We have previously

shown that social pairs are not formed based on cognitive abilities (46), likely due to the

constraints on social mate choice mentioned above. We predicted that males with better spatial

cognitive performance will sire more EPY overall compared to their poorer performing

counterparts and that females opportunistically obtain EP matings with males that perform better

on a spatial cognitive task compared to their social male (9).

Figure 1. Photographs of spatial cognitive testing apparatus. A. Side-view of one array suspended

in the air. B. A mountain chickadee tagged with a passive integrated transponder (PIT) tag

visiting the 'smart' feeder array. Motorized door is open. The blue band on the left leg of the bird

is the PIT tag. C. View of the testing apparatus from below, depicting equidistant placement of

eight 'smart' feeders. Data presented come from four independent feeder arrays.

Methods

Experimental Design

We studied spatial learning and memory abilities and extra-pair paternity rates in

mountain chickadees (*Poecile gambeli*) at our long-term field site in the northern Sierra Nevada

mountains of North America (Sagehen Experimental Forest in northern California, USA,

Sagehen Creek Field Station, University of California Berkeley). Since 2013, we have monitored

breeding phenology in uniquely color-banded individuals across two elevations (47,48) and

measured the spatial cognitive abilities of hundreds of individuals starting in 2015 (e.g., 40-46,

49, 52-54). The breeding season begins in April and continues through July, while the non-

breeding season spans from August to March of the following year. Mountain chickadees are

secondary cavity nesting birds that readily use nest boxes at our field site. Every year in late

April we begin monitoring 350 nest boxes throughout our field site for evidence of nest building.

Once nesting material is detected we monitor all stages of breeding through fledging of each

nest, recording information about the date the first egg is laid, start of incubation, and hatching,

and record the clutch size (number of eggs), and the brood size (number of chicks at post-hatch

day 16) of each nest. We process all chicks in the nest box at post-hatch day 16, which includes

banding them with an aluminum United States Geological Survey (USGS) band, weighing them,

and obtaining a small blood sample for genetic and paternity analyses. We trap adult and first-

year birds using mist nets at established feeders during annual autumn trapping and by hand at

nest boxes during the breeding season. Upon capture, birds are fitted with color-bands and a

unique Passive Integrated Transponder (PIT) tag, wing length is measured, and age is estimated

using multiple plumage features (55). When nestlings banded with USGS bands are recruited

into our population from nest boxes, we band them with an additional color band and PIT tag for

identification.

Testing spatial cognition in the wild

Spatial learning and memory testing was conducted using two feeder arrays at each of our

elevation sites (High, range: 2380 m – 2590 m; coordinates: 39.42402, -120.315015 and Low,

range: 1965 m – 2070 m; coordinates: 39.443500, -120.243248), following 40-46, 49, 52-54.

Within an elevation, the feeder arrays were established ca. 1.5 km apart with mostly non-

overlapping birds visiting each array. Each array consisted of eight equidistant RFID-based

feeders on a square (122 x 122 cm) aluminum frame suspended ca. 4 m above the ground. Each

feeder is equipped with a small, motorized door. Prior to testing, we kept the feeder doors of the

arrays open for approximately one month. Next, the feeder doors were closed and programmed

to open for any bird with a PIT tag, this allowed birds to habituate to the motorized feeder doors

for one week. Finally, we measured the spatial learning and memory abilities of mountain

chickadees using a 4-day task. The spatial learning and memory task involved pseudo-randomly

assigning each bird to a single feeder in the 8-feeder array that was not its most-visited feeder

during the previous two habituation stages. During this task, each bird can only obtain a food

reward (black oil sunflower seed) from its single assigned feeder, but all eight feeders recorded

the date, time, and identity of every visit by all PIT tagged individuals. We measured cognitive

performance by estimating the number of location errors (e.g., non-rewarding feeders visited

prior to visiting the correct rewarding feeder, max 7) per trial, over multiple trials. A trial began

when a bird visited any feeder in the array and ended with a visit to the rewarding feeder. If the

bird landed on its rewarding feeder first, it was considered to have made zero location errors.

Chickadees collect a single sunflower seed during each trial and fly away from the array to

nearby trees to either eat or cache the seed before returning to the array and each trial takes just a

few seconds (54).

We used the mean number of location errors per trial over the first 20 trials and the mean

number of location errors per trial over the entire 4-day task as the two metrics of cognitive

performance, following our previous work (40-46, 49, 52-54). These measurements capture

learning curves in a single metric (Figure S7); as birds learn the rewarding feeder location, they

make fewer and fewer errors on each consecutive trial. Using the mean number of location errors

over multiple trials captures this improvement – the mean number of location errors per trial over

multiple trials is smaller for individuals that improved faster and larger for individuals that

improved slower (40). Since birds completed different numbers of trials over the entire task, we

used the total number of trials as a covariate in models involving performance on the entire 4-day

task (40-46, 49, 52-54). We have previously documented that variation in both metrics is

associated with significant differences in survival (40), longevity (41), in female reproductive

decisions (42), in foraging decisions (53), and in cognitive flexibility (54). For analyses of spatial

cognitive abilities and extra-pair paternity we used cognitive performance measured in the first

year each individual completed the task, as cognitive performance on the spatial learning and

memory task does not change with age (40, 52). In addition, birds were included in analyses if

they performed at least 20 trials and made fewer than mean 3 errors across the 4-day testing

period. Males that bred in more than one year were included in the analyses (controlled for using

male ID as a random effect); as such when reporting results, we report the sample size for unique

nests and 'male-years'.

Assessing extra-pair paternity

188   *DNA extraction and ddRAD sequencing*

For this study, we aimed to sequence and genotype parents with cognition data and their

nestlings over four consecutive years (2018-2021). Each year, we prioritized sequencing nests of

males and potential EP sires that performed in our cognitive task. Samples from each year were

sequenced at the end of each breeding season. Unfortunately, genotypes from 2019 were of low

quality and paternity assignments could not be completed with confidence, therefore we present

data from 2018, 2020, and 2021.

195   Blood samples from adults and nestlings were stored at room temperature in Queens

Lysis buffer until DNA was extracted using Qiagen DNeasy Blood & Tissue kits. The DNA

concentration of each sample was quantified using Qubit Fluorometer 2.0 (Life Technologies)

prior to library preparation and sequencing. We used a double digest restriction-site associated

DNA (ddRAD) sequencing approach from (56) to obtain single nucleotide polymorphisms

(SNPs) for our paternity analyses. Sequencing was completed in batches following field

sampling of each year, therefore, individuals from each breeding year were run together, but

separately from the other years. For each individual, 100-500 ng of extracted DNA was digested

with restriction enzymes SbfI-HF (NEB) and MspI (NEB) and ligated with one of 20 unique P1

adapters and a P2 adapter (P2-MspI). Next, samples were pooled by groups of 20, maintaining

individual identity via unique P1 adapters. Samples were then purified using 1.5x volume of

homemade solid-phase reversible immobilization (SPRI) beads, made with Sera-Mag SpeedBead

Carboxylate-Modified Magnetic Particles (Cytiva) (57). These pooled samples (index groups)

were sent to Cornell University Biotechnology Resource Center (BRC; RRID:SCR_021727) for

size selection between 450 – 600 bp using BluePippin (Sage Science). Following size selection,

adapter ligated DNA was enriched by 11 cycles of polymerase chain reaction (PCR) with a

universal i5 primer and a unique indexed i7 primer for each index group using Phusion DNA

polymerase (NEB). Reactions were again cleaned using 0.7x volume of homemade SPRI beads

(as above) and the index groups pooled in equimolar ratios to create a single sequencing library

run on one lane of Ilumnia NextSeq500 (150 bp single end, performed at BRC). Sequencing was

performed with a ~20% PhiX spike-in to introduce diversity into the library.

SNP Analysis and Paternity assignment

Quality filtering, demultiplexing, and genome alignment

The following were run the same, but separately for each year of samples. First, the quality of all

reads were assessed using FASTQC version 0.11.9

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). All sequences were trimmed using

fastx_trimmer (FASTX-Toolkit) to a length of 147 bp on the 3' end to exclude low-quality SNP

calls. Next, reads containing a single base with a Phred quality score less than 10 were removed

and any sequences that had more than 5% of the bases with a Phred quality score lower than 20

were also removed. We then used *process_radtags* from STACKS version 2.65 to demultiplex

the sequence reads and obtain files with specific sequences for each individual. Next, we aligned

sequences to the chromosome scale reference genome of mountain chickadees (43) using

Bowtie2 version 2.5.1. The resulting .sam files were converted to .bam files and sorted using the

view and *sort* functions in Samtools version 1.18. These aligned reads were used as the input for

the *ref_map.pl* program in Stacks 2.65.

SNP filtering and assigning paternity

SNPs were exported using *populations* in Stacks 2 version 2.65, and each year's samples were

run separately to maximize the number of SNPs and individuals included for downstream

paternity analyses. All male individuals alive during the year of interest were included as

potential sires. Individuals with more than 50% missing SNP data were removed from the

dataset; only 5 individuals from the 2021 dataset did not meet this criterion. Loci were exported

if they were present in 85% of that year's sample population (–*r*). If a RAD locus had more than

one SNP, data were restricted to the first SNP *(–write_single_snp)* to minimize those in high

linkage disequilibrium. We required a minor allele frequency of 0.05 to process a nucleotide *(–*

min_maf), then filtered for loci in Hardy-Weinberg equilibrium (p<0.05) using VCFtools (*--hwe*

option) (58,59). Variant call format (VCF) files were converted to GENEPOP format using the

'radiator' package in R (60).

244   Before assigning paternity, we imported the yearly GENEPOP files into CERVUS (61),

then ran maternity tests separately for each elevation in each year to confirm mother-offspring

relationships identified during field observations. We assessed the quality of these assignments

by plotting the distributions of the first and second-ranked mother's scores for each offspring.

First and second-ranked mother pair LOD scores formed bimodal distributions, with nearly all

first-ranked mother scores above zero and nearly all second-ranked mother scores below zero. In

rare cases when second-ranked mother scores were positive, these were almost always known

sisters or daughters of the mother identified during field observations. Therefore, first-ranked

mother-offspring pairs with positive log likelihood (LOD) scores were considered correct

assignments, but pairs with negative assignments were not included in paternity analyses because

we were not confident in the mother assignment. Offspring resulting from a female laying an egg

in a nest that was not her own (i.e., egg dumps) were uncommon and not included in further

statistical analyses.

257   Paternity analyses were conducted in CERVUS separately for each elevation in each year

by calculating trio LOD scores for each potential father-offspring pair. Trio LOD scores account

for known maternal genotypes when identifying potential fathers. As we did for mother-offspring

pairs, we plotted the distribution of the first and second-ranked male scores and again found a

bimodal distribution centered around zero (Figure S8). Based on these plots, we set cut-off

scores to assign paternity by: all first-ranked male scores at an elevation in a year that were

positive and greater than the top second-ranked male score at that elevation in that year were

assigned as fathers. Offspring sired by the social male at the nest were considered within-pair,

whereas offspring sired by other males were considered extra-pair. Offspring whose social father

was genotyped, but we could not assign a father to, were also considered extra-pair.

268   **Statistical analysis**

We tested whether extra-pair paternity rates differed between the high and low elevations using

generalized mixed effects models with binomial error distributions and 'logit' links, using the R

package 'glmmTMB' (62). Our response variable was whether each offspring was sired by an

extra-pair father or not and we only included nests in which more than 50% of the offspring were

genotyped and run through the paternity analysis. Elevation was included as a fixed effect and

mother identity was included as a random effect in each model. A likelihood ratio test showed

statistical support for an interaction between elevation and year ($p<0.05$), so we tested the effect

of elevation on extra-pair paternity rates separately for each year.

277   We employed two methods to test whether spatial cognitive ability was associated with

the number of extra-pair young a male sired in a year. Both models included the number of extra-

pair young sired as the response variable, with spatial cognitive performance (refers to separate

models run for (1) the mean number of location errors per trial over the first 20 trials and (2) the

mean number of location errors per trial over the 4-day task), elevation, and year as fixed effects,

and male identity as a random effect to control for repeated measures of males across years..

Both models used the Poisson error distribution and a 'log' link, but each model accounted for

zeros in the dataset differently. The first method included a zero-inflation parameter

(ziformula~1) in 'glmmTMB' to account for observations of males with zero EPY, whereas the

second method restricted the dataset to males that sired at least one extra-pair offspring.

It is possible that a relationship between EPP and spatial cognitive ability is driven by

age, as older males often sire more extra-pair offspring than younger males in many systems

(e.g., 63). While chickadees with better spatial cognition in our study system tend to live longer,

cognitive performance does not change with age in the same individuals (40, 41, 52). For each of

the models described, we tested whether including male age as a fixed effect improved each

model using likelihood ratio tests that compared models with and without male age.

In addition, cognitive ability could come at a cost to breeding performance, so we tested

whether spatial cognitive ability was associated with breeding performance using a generalized

linear mixed model with the generalized Poisson distribution (64). Brood size was included as

the response variable and male and female spatial performance on the first 20 trials, elevation,

and clutch size were included as fixed effects. For these analyses, we used all breeding data

available, therefore, year was included as a random effect in this analysis because it had enough

levels (2015 – 2023). Male and female identity were also included as random effects to control

for yearly variation and repeated measures of individual males and females. In a second analysis,

we used a linear model that included mean nestling mass at day 16 as the response variable (in

our system, mass of nestlings on day 16 is a significant predictor of survival and recruitment

(65)), and included male and female performance on the first 20 trials of the spatial cognitive

task and elevation as fixed effects, and year (2015 – 2023) and male and female identity as

random effects. Again, year (2015-2023) and male and female identity were included as random

effects to control for yearly variation and repeated measures of individual males and females.

Siring extra-pair offspring may reduce a male's investment in his own nest, so we tested

whether siring extra-pair young was associated with breeding performance. We used a

generalized linear mixed model with the generalized Poisson distribution to test whether brood

size in a male's nest was affected by the male siring extra-pair offspring (yes or no), and we used

a linear mixed model to test whether the mean mass of the nestlings in a male's nest was affected

by whether the male sired any extra-pair offspring. Elevation, clutch size, and year (3 years of

EPP data) were included as fixed effects and male and female identity were included as random

effects in both models to control for repeated measures of individual males and females. We

tested whether the mass of young sired by extra-pair males differed from young sired by social

males using a linear mixed model with nestling mass at day 16 as the response variable and

extra-pair status, elevation, and year as fixed effects. Nest identity was included as a random

effect to control for nestlings that came from the same nests.

We tested whether the spatial cognitive abilities of extra-pair males differed from the

males they cuckolded by comparing spatial cognition between father types (EP male vs social

male within the same nest) using linear mixed models. Spatial cognitive score was the used as

the response variable, then father type (extra-pair mate versus social mate) and elevation were

included as fixed effects. Offspring identity was included as a random effect to restrict

comparisons to within offspring and mother identity was included as a random effect to control

for repeated measures of females. When fitting models to the total location errors, we log-

transformed the response variable to improve residual fit and included a quadratic term of total

trials completed during the 4-day cognitive testing period, since the number of trials completed

influences performance (40-46, 49, 52-54). Here, we tested whether extra-pair males were

significantly older than the social males they cuckolded using a chi-square test.

The cognitive abilities of either the male or female could affect the likelihood of a nest

containing extra-pair young, so we tested whether cuckoldry was associated with the spatial

cognitive abilities of either parent at the nest using multiple methods. First, we modeled the

relationship between male and female spatial cognitive ability and cuckoldry at the nest and

offspring level. At the nest level, we tested whether male or female spatial cognitive ability

predicted whether any offspring in a nest were sired by extra-pair males. At the offspring level,

we tested whether male or female spatial cognitive ability predicted whether each offspring was

sired by an extra-pair male or not. For both tests, we used generalized linear mixed models with

binomial distributions that included male and female spatial cognition score, elevation, and year

as fixed effects. Male and female identity were included as random effects. Second, for nests that

had at least one extra-pair young, we tested if the number of extra-pair young in a nest was

associated with male or female spatial cognitive ability using a generalized linear mixed model

with a truncated Poisson distribution. The number of extra-pair young in the nest was included as

the response variable, while male and female performance on the first 20 trials, year, and

elevation were included as fixed effects. Male and female identity were included as random

effects to control for repeated measures of individual males and females. The effect of male age

on cuckoldry rates was tested by adding male age as a fixed effect to each model and comparing

models with and without male age using likelihood ratio tests.

We also tested whether male spatial cognitive ability differed between nests that had

extra-pair young compared to those that did not. Male spatial cognition was included as the

response variable and elevation and whether the nest had any extra-pair young were included as

fixed effects. When modeling the mean number of location errors per trial on the entire task we

used the Gamma distribution and included total trials as a fixed effect. Including male identity as

a random effect in both models led to convergence errors, so male identity was not included in

the models.

We tested whether the mean number of location errors per trial on the first 20 trials

correlated with the mean number of location errors per trial over the entire task for males using a

linear model. Mean location errors over the entire task was used as the response variable while

mean location errors over the first 20 trials and the total number of trials completed were

included as fixed effects. All statistical analyses were performed in R version 4.1.1 (66) and the

residual fit of all models was checked using DHARMa (67). Statistical significance was

calculated using Wald Chi-square tests.

Results

Extra-pair paternity and spatial cognition

Extra-pair paternity rates

Across three years of data collection (2018, 2020, and 2021), we found that 30.3% of the total

offspring sampled (222 of 732) were sired by extra-pair males and that 70.1% of nests (89 of

127) had at least one extra-pair young (EPY). We found no significant differences in the number

of extra-pair young sired at high versus low elevation populations regardless of year sampled

(Supplemental Results, Figure S1, Table S1), therefore, we combined high and low elevations for

all later analyses. Data from breeding birds in this study confirmed previous research showing

that female chickadees seek extra-pair copulations with neighboring males (on average ~100 m

from their own nests; Figures S2 and S3), which are likely familiar males from their nonbreeding

flocks (36, 39, 68, 69). In addition, breeding male territories do not appear to cluster based on

cognitive performance (Figure S4).

Male spatial cognitive abilities and extra-pair paternity

Performance on the spatial learning and memory task predicted the number of EPY a male sired

each year (Figure 2). Males with better spatial learning and memory abilities (e.g., fewer mean

location errors per trial) sired more EPY, both according to a zero-inflated model including males

that sired no extra-pair young (Mean location errors per trial over first 20 trials: χ^2_1=6.28,

P=0.012 (Figure 2, dashed line); Mean location errors per trial over the entire task: χ^2_1=8.12,

P=0.004, N=280 (137 individual males; Figure S5)), and to a model only including males that

sired at least one EPY (Mean location errors per trial over first 20 trials: χ^2_1=5.96, P=0.015

(Figure 2, solid line); Mean location errors per trial over the entire task: χ^2_1=7.68, P=0.006, N=41

(34 individual males; Figure S5). This model suggests that of the males that sired at least one

EPY, those with the best cognitive performance may sire between 6 and 7 EPY each year,

whereas males with the worst cognitive performance will sire only 1 to 2 EPY each year. We

have shown that birds with better spatial cognitive abilities live longer (41) and lack reproductive

senescence within their natural life span (70), therefore, males with better cognition are expected

to produce significantly more offspring over their lifetime. Importantly, we show that male age

was not associated with the number of EPY sired (see Supplemental Results).

Do males that sire EPY lose reproductive benefits in their own nests?

Three years of EP paternity data suggest that EP males do not lose fitness in their own nests

compared to males that did not sire any EP offspring, as neither brood size (brood size: χ^2_1=0.30,

P=0.584, N=144 nests) nor mean nestling mass (mean mass: χ^2_1=2.10, P=0.147, N=140 nests)

differed significantly between males that had EPY and males that did not sire any EPY. We also

compared the mass of young sired by extra-pair males to those sired by the social males raised in

the same nests and found no significant difference (χ^2_1=2.83, P=0.093, N=719 offspring).

Social parents' spatial cognitive abilities and reproductive success

In addition to testing how cognitive abilities were related to number of EPY, we also tested

whether variation in cognitive abilities had direct consequences on breeding performance. Across

9 years of breeding data (2015-2023), we detected no relationship between the social males'

spatial cognitive abilities and brood size (χ^2_1=2.32, P=0.127, N=205 nests), or the females'

spatial cognitive abilities and brood size (χ^2_1=1.75, P=0.186, N=205 nests). Controlling for the

effect of clutch size on brood size produced similar results (Table S4). However, males with

better spatial cognition raised heavier young (mass at day 16, χ^2_1=4.86, P=0.028, N=202 nests),

while female spatial cognition showed no significant relationship with offspring mass at day 16

(χ^2_1=0.93, P=0.335, N=202 nests).

Figure 2. Model predictions of the number of extra-pair young a male sired in a year by the mean

number of location errors per trial over the first 20 trials. The dashed line represents a zero-

inflated model for all males, including those with no extra-pair young (N=280; 137 individual

males), and the solid line represents the model prediction for males that sired at least one extra-

pair young (N=41; 34 individual males). Points are slightly *jittered* on the y-axis to improve

readability. Shading represents bootstrapped 95% confidence intervals. Inset depicts 8 nestlings

at 16 days post-hatching. See model predictions for the number of extra-pair young a male sired

in a year by total 4-day testing performance in Figure S5.

Extra-pair males have better spatial cognition than the social males they cuckold

When there were EPY in a nest, they were sired by males with better spatial learning and

memory abilities (on average) compared to those of the social male (Mean location errors per

trial over first 20 trials: χ^2_1=14.81, P<0.001, N=71 comparisons of a social male and the extra-

pair male sire (Figure 3); Mean location errors per trial over the total task: χ^2_1=6.08, P=0.014,

N=71 comparisons (Figure S6)). Furthermore, extra-pair males did not differ significantly in age

from the males they cuckolded (28 offspring were sired by an extra-pair male that was older than

the social male, 19 offspring were sired by an extra-pair male that was the same age as the social

male, and 24 offspring were sired by an extra-pair male that was younger than the social male;

Chi-square test: χ^2_2= 1.72, P=0.424).

Figure 3. Comparing spatial cognitive performance (mean number of location errors per trial

over the first 20 trials) of extra-pair (mean ± SE = 0.96 ± 0.07) and social fathers (mean ± SE =

1.17 ± 0.07, 71 comparisons). Lines connect males by offspring: blue lines show comparisons

where the extra-pair father performed better than the social father (49 of 71 cases or 69%), and

purple lines show comparisons where the social father performed better than the extra-pair father

(19 of 71 cases, or 26.7%; in 3 cases the extra-pair and social father had the same cognition

score). Shaded distributions represent smoothed histograms of cognitive performance for each

father type. The same analysis is presented for performance over the 4-day testing period in

Figure S6.

Does spatial cognition of the social male or female predict presence of EPY?

The finding that extra-pair males have better spatial cognition than the social male of the nest (on

average) could suggest that when females obtain EP copulations (EPC), they may do so

preferentially with males that have better cognition than their social male. But it is also possible

that female cognitive abilities influence their own decisions to seek EPC (i.e., a female may seek

EPC to compensate for her own poor cognition). Therefore, we assessed whether the likelihood

of having EPY in the nest was related to the spatial cognitive abilities of the social male or the

female. The likelihood of being cuckolded was not significantly related to the cognitive abilities

of the social males (Offspring level: Mean location errors per trial over first 20: χ^2_1=0.25,

P=0.620, Mean location errors per trial over the entire task: χ^2_1=0.43, P=0.513, N=348 offspring

for 42 males; Nest level: Mean location errors per trial over first 20: χ^2_1=0.05, P=0.808; Mean

location errors per trial over the entire task: χ^2_1=0.22, P=0.642, N=59 nests for 42 males). In

addition, male age did not significantly affect the likelihood of a male being cuckolded (see

Supplemental Results). However, females who performed worse on the first 20 trials of the

spatial cognitive task were more likely to have EPY in their nests (Offspring level: Mean

location errors per trial over first 20: χ^2_1=6.21, P=0.013, N=348 offspring for 42 females);

though this relationship was not significant when we analyzed cognitive performance across the

entire 4-day task (χ^2_1=0.18, P=0.673, N=348 offspring for 42 females), or at the nest level of

analysis (Mean location errors per trial over first 20: χ^2_1=0.21, P=0.645, Mean location errors per

trial over the entire task: χ^2_1=0.13, P=0.723, N=59 nests for 42 females).

Similarly, for nests that had at least one EPY, the cognitive abilities of the social male did

not predict the number of EPY in their nests (χ^2_1=1.03, P=0.311, N=41 nests for 35 males),

however, females that performed worse on the task had significantly more EPY in their nests

(χ^2_1=6.79, P=0.009, N=41 nests for 32 females). In other words, females may not engage in EPC

solely in response to their social mate's poor cognition. Instead, the data suggest that females

may engage in EPCs based on their own cognitive deficiencies, and when doing so, may seek

males with better spatial cognitive abilities than their social mate (though we are speculating

here, as we did not assess all EPC attempts in the system, only those that resulted in young).

Nevertheless, males with better spatial cognitive abilities have higher fitness; they produce more

EPY overall and we found no evidence of a trade-off for producing more EPY, as cuckoldry rates

and reproductive output (brood size and fledgling mass) were similar among males with better

and worse spatial cognitive performance.

Discussion

Our findings show that sexual selection acts on the spatial cognitive abilities of free-

living, food-caching mountain chickadees via extra-pair paternity. We found that (a) males, but

not females with better spatial cognitive abilities produce higher quality offspring (higher mass,

65) in their social nests, (b) males with better spatial learning and memory abilities sire more

EPY than their poorer performing counterparts and, (c) EP males have better spatial cognitive

abilities than the social males they cuckold, regardless of age. Importantly, our data show that

absolute male cognitive abilities do not predict EPY in the nest, rather that the differences in

cognition between the EP male and the social male are relative to each other –a good quality

male can lose paternity to an even better-quality male.

We specifically quantified EPP in our mountain chickadee system and do not have data

on all EPC attempts by males or females. However, given the extensive research on chickadee

breeding behavior and their closely related Eurasian relatives (tits), we suggest these patterns of

EPP may be driven by females for several reasons; 1. Female parids (chickadees and tits) are

known to pursue EPCs with neighboring males (68, 69, 71-74), 2. Parids have never been

reported or observed engaging in forced copulation (e.g., 36, 74, 75), and 3. Resident males are

highly territorial and do not tolerate male conspecifics in their territories. In addition, the

directional relationship between the cognitive phenotype of the social and extra-pair male would

not be predicted by male choice but is consistent with good genes hypotheses, where females

seek high quality males (in this case, those with better spatial cognitive abilities) for EPC (1, 9,

14). In other words, males would benefit from siring EPY with any female regardless of her

social male's cognitive abilities.

That said, active choice need not be implied, as males with poor spatial cognitive abilities

could be worse at defending their territories from other males, which may allow those males with

better cognitive abilities to intrude and copulate with the resident female. We think this

explanation is unlikely as males with worse cognitive abilities tend to be more aggressive (76,

77) and cognitive performance is not associated with social dominance (78). Females, on the

other hand, would benefit from obtaining EPCs with males that have better spatial cognition than

their social male, as this trait is heritable (43, 44) and would increase the likelihood of producing

offspring with better spatial cognition, leading to higher overwinter survival (40) and longer

lifespan (41). Indeed, females that performed worse on the spatial cognitive task had more EPY

in their nests, while social male cognitive performance did not predict EPY in the nest. This

suggests that females may compensate for their poor cognitive abilities via EPY. Considering

that alleles associated with better spatial cognitive abilities are not sex linked (43), producing

EPY with better quality males increases selection pressure on spatial cognitive abilities in both

males and females, as both sexes rely on spatial cognition for survival (79).

Although it remains unknown whether chickadees can evaluate cognitive abilities directly

or indirectly via some correlated trait(s), the life history and breeding ecology of mountain

chickadees provide several possibilities. For example, female chickadees may use secondary

sexual signals, such as song and plumage variation associated with spatial cognitive abilities, to

identify high-performing males (1, 9, 28, 80). Recent work in closely related black-capped

chickadees shows that male cognitive performance is associated with variation in plumage patch

brightness and contrast (80). Alternatively, it is possible that females can directly assess a male's

ability to successfully recover food stores over winter and base mating decisions on this

information in the spring (30, 38), as chickadees are nonmigratory, sedentary birds that aggregate

in flocks of unrelated male-female pairs throughout the nonbreeding season and break apart once

breeding commences (38, 39). Data from this study confirm previous research showing that

female chickadees seek EPCs with neighboring males (68, 69, 74), which would likely be males

from their winter flock (38, 39). Regardless of if and how females evaluate male cognition,

mating with males with better cognitive performance should increase female fitness by

increasing the likelihood of offspring inheriting the specific alleles associated with better

cognition, which would be expected to lead to higher offspring survival and longer lifespan (40,

41).

An alternative hypothesis for our findings is that females seek EPCs with other males as

insurance against the social male's infertility (81), again gaining indirect genetic benefits. This

hypothesis does not explain the direction of our findings, because male cognitive abilities did not

predict the number of offspring a male produced in his own nest (i.e., cuckolded social males still

fathered chicks in their nests), suggesting that males with poorer cognition do not suffer from

infertility. That said, we do not have data on the sperm quality of these birds and therefore cannot

exclude differences in fertility or sperm competition between males with better and worse spatial

cognition. Related, we have not detected fitness trade-offs or negative effects associated with

better cognitive abilities in our system; we do not see cognitive or reproductive senescence

within birds' naturally occurring life span (45, 70). In fact, chickadees with better spatial

cognition live longer allowing them to produce more offspring over their lifetime compared to

chickadees with worse cognition (41).

Interestingly, a previous study in our system showed that females mated to social males

with better spatial abilities lay larger clutches and fledge larger broods in years of resource

abundance (42). Together with the present findings, we suggest that female mating decisions,

including EPCs and reproductive investment, may vary opportunistically. When resources are

abundant, females lay larger clutches if their social male has better cognitive abilities (42), and

when males with better spatial cognition than her social mate are available for EPCs, they pursue

those opportunities. Indeed, we found that females with poorer spatial cognition had more EPY

in their nests, suggesting female mating decisions could be based on some physiological signal

associated with poorer cognition that serves as a compensatory mechanism. At the ultimate level,

females with poorer cognition likely benefit disproportionately by mating with the highest

quality male they can find, which in the case of food-caching birds, means those with better

spatial cognitive abilities (40, 41).

An additional alternative explanation for our findings is that birds with better spatial

cognitive abilities have access to better territories and therefore are clustered based on cognitive

abilities (82). Then, because females have extra-pair copulations with neighboring males (68, 69,

74), those neighbors would happen to be males with better spatial cognition. However, in our

study system, males do not appear to cluster based on cognitive performance (Figure S4),

suggesting that females are not finding males with better cognitive abilities than their social

males by chance. Furthermore, while higher social dominance status is associated with access to

better territories in chickadees (83), we see no association with dominance status and spatial

cognitive abilities in our system (78) or closely related black-capped chickadees (80). While it is

possible that our results reflect some unmeasured trait(s) linked to spatial cognitive ability, such

as boldness, aggression, or increased exploratory behavior (e.g., 32), previous data from our

system do not support these additional alternative hypotheses. Chickadees with enhanced spatial

cognitive abilities are not more aggressive (76) nor do they show higher rates of exploratory

behavior (e.g., higher space use) (77).

Given the fitness advantages of superior spatial cognitive abilities for food-caching

chickadees, how variation is maintained and why enhanced spatial cognitive abilities have not

evolved to fixation may seem puzzling. While food-caching birds exhibit superior spatial

cognitive abilities compared to their non-caching counterparts (84), considerable variation

persists within species (49). Research from our system demonstrates the highly polygenic nature

of the spatial cognitive phenotype, meaning there are multiple genetic pathways that may

produce the same cognitive phenotype (43, 44, 85). In addition, selection strength varies with

annual variation in winter climate, which in milder years, may allow individuals to survive and

reproduce despite comparatively poor spatial cognitive abilities (47, 48). Finally, we have

documented elevation-related differences in spatial cognition and the hippocampus of mountain

chickadees (49, 50), but there is gene flow among elevations (51) which moves allelic variation

associated with poorer cognition within the population.

Overall, our study provides novel results showing that sexual selection contributes to the

evolution of spatial cognitive abilities in a food-caching species, which rely on such spatial

cognition for survival. We show that males with better spatial cognition exhibit higher

reproductive output, via siring more EPY and fledging heavier chicks, which are more likely to

survive and be recruited into the breeding population (65). We also show that when there are

EPY in the nest, they are sired by males with better spatial cognition than the social male (on

average), which suggests that females may be selectively obtaining EPC with higher quality

males; though we do not have direct experimental evidence of female choice. Whether these

patterns are driven by females or males, males with better spatial cognitive abilities exhibit

higher fitness by producing more offspring, adding to the evidence for selection on spatial

cognitive abilities, including differential survival and longevity in food-caching birds (40, 41).

Acknowledgements

Thank you to Qwahn Kent, Danielle Schaening-Lopez, Alyssa Nowicki, and Beck Kerdman for

assistance with DNA extractions and the Cornell University BRC Bioinformatics Core Facility

(RRID:SCR_021757) for providing computing power for all bioinformatics. All procedures were

in accordance with University of Nevada Reno IACUC (protocols 20-11-1103-1, 20-06-1014,

20-08-1062) and California State Fish and Wildlife Scientific Collecting Permit (S-193630001-

20007-001).

Funding

Kessel Fellowship from American Ornithological Society (CLB)

Edward Rose Fellowship from Cornell Lab of Ornithology (CLB)

National Science Foundation Graduate Research Fellowship Program, 2019287870 (BRS)

National Science Foundation Graduate Research Fellowship Program, 2020305313 (LMB)

Author contributions

Conceptualization: CLB, VVP

Methodology: CLB, BRS, JFW, BGB, VKH, AMP, LMB, ESB, MSW, IJL, VVP

Investigation: CLB, BRS, VKH, AMP, LMB, VVP

Visualization: CLB, JFW, VVP

Funding acquisition: CLB, VVP

Writing – original draft: CLB

Writing – review & editing: CLB, BRS, JFW, BGB, VKH, AMP, LMB, ESB, MSW, IJL, VVP

Competing interests: Authors declare that they have no competing interests.

Data Availability: All data, code, and materials used in the analysis are available at

https://figshare.com/s/ef7f40371b487b9dabca.

References
1. Andersson, M & Simmons, LW. Sexual selection and mate choice. *Trends Ecol. Evol.* **21**,

296–302 (2006).

2. Rosenthal, G. *Mate Choice: the Evolution of Sexual Decision Making from Microbes to*

Humans. Princeton, NJ, Princeton University Press (2017).

3. Davies, AD, Lewis, Z, & Dougherty LR. A meta-analysis of factors influencing the
strength of mate-choice copying in animals. *Behav. Ecol.* **31**, 1279-1290 (2020).

4. Henshaw, JM, Fromhage, L, & Jones, AG. The evolution of mating preferences for
genetic attractiveness and quality in the presence of sensory bias. *PNAS* **119**,
e2206262119 (2022).

5. DuVal, EH, Fitzpatrick, CL, Hobson, EA, & Servedio, MR. Inferred attractiveness: A
generalized mechanism for sexual selection that can maintain variation in traits and
preferences over time. *PLoS Biol.* **21**, e3002269 (2023).

6. Fisher, RA. *The genetical theory of natural selection.* Oxford, Clarendon Press (1930).

7. Prokop, Z.M., Michalczyk, Ł., Drobniak, S.M., Herdegen, M. and Radwan, J. Meta-
analysis suggests choosy females get sexy sons more than "good genes". *Evolution.* **66**,
2665-2673 (2012).

8. Endler, JA. Sensory drive: does sensory biology bias or constrain the direction of
evolution? *Am. Nat.* **139**, S1-S153 (1992).

9. Andersson, M. *Sexual Selection.* Princeton, NJ, Princeton University Press (1994).

10. Bonduriansky, R. The evolution of male mate choice in insects: a synthesis of ideas and
evidence. *Biological Reviews.* **76**, 305-339 (2001).

11. Gowaty, PA, Steinichen, R, & Anderson, WW. Indiscriminate females and choosy males:
within☐and between☐species variation in Drosophila. *Evolution.* **57**, 2037-2045 (2003).

12. Verner, J, Wilson, MF. Mating systems, sexual dimorphism and the role male North
American passerine birds in the nesting cycle. *Oenitol. Monogr.* **9**, 1-76 (1969).

13. Schärer, L, Rowe, L, & Arnquist, G. Anisogamy, chance and the evolution of sex roles.
Trends in Ecology and Evolution. **27**, 260-264 (2012).

14. Rubenstein, DR. *Animal Behavior*. New York, NY, Oxford University Press (2023).

15. Gowaty, PA. Field studies of parental care in birds: new data focus questions on variation among females. *Adv. Study Behav*. **25**, 477-531 (1996).

16. Zahavi, A. Mate selection – a selection for a handicap. *J. Theo. Biol*. **53**, 205-214 (1975).

17. Birkhead, TR, & Møller, A. Sperm Competition in Birds: Evolutionary Causes and Consequences. London, UK, Academic Press (1992).

18. Hamilton, WD, & Zuk, M. Heritable true fitness and bright birds: A role for parasites? *Science*. **218**, 384-387 (1982).

19. Östlund, S & Ahnesjö, I. Female fifteen-spined sticklebacks prefer better fathers. *Animal Behaviour*. **56**, 1177-1183 (1998).

20. Hill, GE. Plumage coloration is a sexually selected indicator of male quality. *Nature*. **350**, 337-339 (1991).

21. Lewis, SM, & Cratsley, CK. Flash signal evolution, mate choice, and predation in fireflies. *Annual Review of Entomology*. **53**, 293-321 (2008).

22. Patricelli, GL, Uy, JAC, & Borgia, G. Multiple males traits interact: Attractive bower decorations facilitate attractive behavioral displays in satin bowerbirds. *Proc. R. Soc. B*. **270**, 2389-2395 (2003).

23. Kodric-Brown, A & Brown, JH. Truth in advertising: The kinds of traits favored by sexual selection. *Am. Nat*. **124**, 309-323 (1984).

24. Endler, JA. *Natural Selection in the Wild*. Princeton, NJ, Princeton University Press (1986).

25. Morand⬜Ferron, J, Cole, EF, & Quinn, JL. Studying the evolutionary ecology of

cognition in the wild: a review of practical and conceptual challenges. *Biological*

Reviews, **91**, 367-389 (2016).

26. Shettleworth, SJ. *Cognition, Evolution, and Behavior*. New York, NY, Oxford University

Press (2010).

27. Johnston, TD. Selective costs and benefits in the evolution of learning. In *Advances in the*

Study of Behavior. Academic Press **12**, 65-106 (1982).

28. Boogert, NJ, Fawcett, TW, Lefebvre, L. Mate choice for cognitive traits: a review of the

evidence in nonhuman vertebrates, *Behav. Ecol.* **22**, 447–459 (2011).

29. Snowberg, LK, & Benkman, CW. Mate choice based on a key ecological performance

trait. *J. Evol. Biol.* **22**, 762-769 (2009).

30. Chen, J, Zou, Y, Sun, Y-H, & Ten Cate, C. Problem-solving males become more attractive

to female budgerigars. *Science* **363**, 166 –167 (2019).

31. Shohet, AJ, & Watt, PJ. Female guppies *Poecilia reticulata* prefer males that can learn

fast. *J Fish Biol.* **75**, 1123 – 1523 (2009).

32. Spritzer, MD, Meikle, DB, & Solomon, NG. Female choice based on male spatial ability

and aggressiveness among meadow voles. *Anim Behav.* **69**, 1121–1130 (2005).

33. Shaw, RC, MacKinlay RD, Clayton, NS, & Burns, KC. Memory performance influences

male reproductive success in a wild bird. *Curr. Biol.* **29**, 1498-1502 (2019).

34. Keagy, J. Savard, JF, & Borgia, G. Male satin bowerbird problem-solving ability predicts

mating success. *Anim. Behav.* **78**, 809-817 (2009).

35. Croston, R, Branch, CL, Kozlovsky, DY, Dukas, R, & Pravosudov, VV. Heritability and

the evolution of cognitive traits. *Behav. Ecol.* **26**, 1447-1459 (2015).

36. McCallum, D, Grundel, R, Dahlsten, DL. Mountain Chickadee (*Poecile gambeli*), version 1.0. In Birds of the World (A. F. Poole and F. B. Gill, Editors). Cornell Lab of Ornithology, Ithaca, NY, USA (2020).

37. Pravosudov, VV, & Roth II, TC. Cognitive ecology of food hoarding: the evolution of spatial memory and the hippocampus. *Ann. Rev. Ecol. Evol. System.* **44**, 173-193 (2013).

38. Lemmon D, Withiam ML, Barkan CPL. Mate protection and winter pair-bonds in black-capped chickadees. *Condor* **99**, 424–433 (1997).

39. Ekman, J. Ecology of non-breeding social systems of Parus. *Wil. Bull.* **101**, 263–288 (1989).

40. Sonnenberg, BR, Branch, CL, Pitera, AM, Bridge, E, & Pravosudov, VV. Natural selection and spatial cognition in wild food-caching mountain chickadees. *Curr. Biol.* **29**, 670-676 (2019).

41. Welklin, JF *et al.* Spatial cognitive ability is associated with longevity in a food-caching bird. *Science*, **385**, 1111-1115 (2024).

42. Branch, CL, Pitera, AM, Kozlovsky, DY, Bridge, ES, & Pravosudov, VV. Smart is the new sexy: female mountain chickadees increase reproductive investment when mated to males with better spatial cognition. *Ecol. Lett.* **22**, 897-903 (2019).

43. Branch, CL *et al.* The genetic basis of spatial cognitive variation in a food-caching bird. *Curr. Biol.* **32**, 210-219 (2022).

44. Semenov, G *et al.* Genes and gene networks underlying spatial cognition in food-caching chickadees. *Curr. Biol.* **34**, 1930-1939 (2024).

45. Heinen, VK *et al.* Specialized spatial cognition is associated with reduced cognitive senescence in a food-caching bird. *Proc. R. Soc. B.* **288**, 20203180 (2021).

46. Branch, CL, *et al.* What's in a mate? Social pairing decisions and cognitive performance

in food-caching mountain chickadees. *Proc. R. Soc. B.* **290**, 20231073 (2023).

47. Kozlovsky, DY, Branch, CL**,** Pitera, AM, Croston, R, & Pravosudov, VV. Fluctuations in

annual climatic extremes are associated with reproductive variation in resident mountain

chickadees. *R Soc. Open Sci.* **5**, 171604 (2018).

48. Whitenack, LE, *et al.* Complex relationships between climate and reproduction in a

resident montane bird. *R. Soc. Open Sci.* **10**, 230554 (2023).

49. Croston, R *et al.* Individual variation in spatial memory performance in wild mountain

chickadees from different elevations. *Anim. Behav.* **111**, 225-234 (2016).

50. Freas, CA, LaDage LD, Roth II, TC, & Pravosudov, VV. Elevation-related differences in

memory and the hippocampus in mountain chickadees, *Poecile gambeli. Anim. Behav.*

84*, 121-127 (2012).

51. Branch, CL, Jahner, JP, Kozlovsky, DY, Parchman, TL, & Pravosudov, VV. Absence of

population structure across elevational gradients despite large phenotypic variation in

mountain chickadees *(Poecile gambeli). R. Soc. Open Sci.* **4**, 170057 (2017).

52. Tello-Ramos, MC *et al.* Memory in wild mountain chickadees from different elevations:

comparing first-year birds with older survivors. *Anim. Behav.* **137**, 149-160 (2018).

53. Benedict, LM *et al.* Information maintenance of food sources is associated with

environment, spatial cognition and age in a food-caching bird. *Anim. Behav.* **182**, 153-

172 (2021).

54. Benedict, LM, Heinen, VK, Sonnenberg, BR, Bridge, ES, & Pravosudov, VV. Learning

predictably alternating food locations across days in a food-caching bird. *Anim. Behav.*

196, 55-81 (2023).

55. Pyle, P. Identification Guide to North American Birds, Part I. Second Edition. Sheridan

Books, Ann Arbor, MI, USA. (2022).

56. Thrasher, DJ, Butcher, BG, Campagna, L, Webster, MS, & Lovette, IJ. Double☐digest

RAD sequencing outperforms microsatellite loci at assigning paternity and estimating

relatedness: A proof of concept in a highly promiscuous bird. *Mol. Ecol. Res.* **18**, 953-965

(2018).

57. Rohland, N, & Reich, D. Cost-effective, high-throughput DNA sequencing libraries for

multiplexed target capture. *Genome Res.* **22**, 939-46 (2012).

58. Wigginton, JE, Cutler, DJ, & Abecasis, GR. A note on exact tests of Hardy Weinberg

equilibrium. *Am. J Human Genetics* **76,** 887-893 (2005).

59. Danecek, P, *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158

(2011).

60. Gosselin, T. radiator: RADseq Data Exploration, Manipulation and Visualization using R.

R package version 1.1.9 https://thierrygosselin.github.io/radiator/ (2020).

61. Kalinowski, S, Taper, M, & Marshall, T. Revising how the computer program CERVUS

accommodates genotyping error increases success in paternity assignment. *Mol Ecol.* **16**,

1099-1106 (2007).

62. Brooks, ME, *et al.* glmmTMB balances speed and flexibility among packages for zero-

inflated generalized linear mixed modeling. *R J.* **9**, 378–400 (2017).

63. Double, M, & Cockburn, A. Pre-dawn infidelity: females control extra-pair mating in

superb fairy-wrens. *Proc Biol Sci.* **267**, 465-70 (2000).

64. Joe, H, & Zhu, R. Generalized Poisson distribution: the property of mixture of Poisson

and comparison with negative binomial distribution. *Biometrical Journal: J. Math.*

Methods in Biosci. **47**, 219-229 (2005).

65. Whitenack, L.E., Richmond, A., Sonnenberg, B.R., Welklin, J.F., Heinen, V.K.,Pitera,

A.M., Branch, C.L., Pravosudov, V.V. Postnatal dispersal and drivers of successful

recruitment in resident Mountain Chickadees, *Ornithology,* ukaf006 (2025).

66. R Core Team. R: A language and environment for statistical computing (2021).

67. Hartig, F. DHARMa: Residual diagnostics for hierarchical (multi-level/mixed) regression

models R package Version 0.4.4. (2018).

68. Smith, SM. Extra-pair copulations in black-capped chickadees: the role of the female.

Behav. **107**, 15-23. (1988).

69. Otter, K, Ratcliffe, L, & Boag P. Extra-pair paternity in the black-capped chickadee.

Condor **96**, 218 – 222 (1994).

70. Welklin, JF *et al.* No evidence of reproductive senescence within natural life span in

resident mountain chickadees (*Poecile gambeli*). *Behav. Ecol. Sociobiol.* **79**, 10 (2025).

71. Hasselquist, D, & Sherman, PW. Social mating systems and extrapair fertilizations in

passerine birds. *Behav. Ecol.* **12**, 457–466. (2001).

72. Kempenaers, B, Verheyen, GR, Vanderbroeck, M, Burke, T, Vanbroeckoven, C, &

Dhondt, AA. Extra-pair paternity results from female preference for high-quality males in

the blue tit. *Nature* **357**, 494–96 (1992).

73. Sheldon, BC. Male phenotype, fertility, and the pursuit of extra-pair copulations by

female birds. *Proc. R. Soc. B*. **257**, 25–30 (1994).

74. Mennill DJ, Ramsay SM, Boag PT, & Ratcliffe LM. Patterns of extrapair mating in

relation to male dominance status and female nest placement in black-capped

chickadees. *Behavioral Ecology*, *15*, 757-765. (2004).

75. Bonderud ES, Otter KA, Burg TM, Marini KLD, Reudink MW. Patterns of extra-pair

paternity in mountain chickadees. *Ethology*. 124: 378–386. (2018).

76. Branch, CL, Kozlovsky, DY, & Pravosudov, VV. Elevation related variation in aggressive

response to mirror image in mountain chickadees. *Behav.* **152**, 667-676 (2015).

77. Kozlovsky, DY, Branch, CL, Freas, CA, & Pravosudov, VV. Elevation-related

differences in exploration and social dominance in mountain chickadees (*Poecile*

gambeli). *Behav. Ecol. Sociobiol.* **68**, 1871-1881 (2014).

78. Heinen, VK *et al.* Social dominance has limited effects on spatial cognition in a wild

food-caching birds. *Proc. R. Soc. B.* **288**, 20211784 (2021).

79. Branch, CL, *et al.* Testing the greater male variability phenomenon: male mountain

chickadees exhibit larger variation in reversal learning performance compared to females.

Proc. R. Soc. B. **287**, 20200895 (2020).

80. Robayo Noguera, L, Stevenson, CAL, Wang, T, Pasquale, MK, Branch, CL. Variation in

plumage reflectance but not song reflects spatial cognitive performance in black-capped

chickadees *(Poecile atricapillus). Anim. Cog.* **28**, 14 (2025).

81. Lifjeld JT, Kleven O, Amundsen T, & Slagsvold T. Extrapair copulations in coal tits

(*Periparus ater*): Female insurance against male fertility. *Beh. Ecol. Sociobiol.* **79**, 79

(2025).

82. Canal D, Jovani R, & Potti J. Male decisions or female accessibility? Spatiotemporal

patterns of extra pair paternity in a songbird. *Behavioral Ecology*, **23**, 1146-1153 (2012).

83. Ratcliffe, L, Mennill, DJ, & Schubert, KA. Social dominance and fitness in black-capped

chickadees in *Ecology and Behaviour of Chickadees and Titmice: An integrative*

approach. Oxford, UK, Oxford University Press (2007).

84. Brodbeck, DR, Shettleworth, SJ. Matching location and color of a compound

stimulus: comparison of a food-storing and nonstoring bird species. *J. Exp.Psychol.* **21**,

64-77 (1995).

85. Pritchard JK, Pickrell JK, & Coop G. The genetics of human adaptation: hard

sweeps, soft sweeps, and polygenic adaptation. *Curr. Biol.* **23**, R208-R215 (2010).

86. Sonnenberg, B.R., Heinen, V.K., Pitera, A.M., Benedict, L.M., Branch, C.L., Bridge,

E.S., Ouyang, J., Pravosudov, V.V. Natural variation in developmental condition has

limited effect on spatial cognition in a wild food-caching bird. *Proceedings of the Royal*

Society B **289**, 1984 (2022).

87. Pitera, AM, Branch, CL, Sonnenberg, BR, Benedict, LM, Kozlovsky, DY, Pravosudov,

VV. Reproduction is affected by individual breeding experience but not pair longevity in

a socially monogamous bird. *Behav. Ecol. Sociobiol.* **75**, 101 (2021).

88. Sonnenberg, BR, *et al.* Long-term winter food supplementation in a montane

environment shows no significant impact on reproductive success in a resident food-

caching bird. *Ornithology* **140**, 1-12 (2022).

89. Branch, CL *et al.,* Code and data for: Seeking smarts: Female chickadees choose extra-

pair males based on spatial cognitive abilities. Figshare (2024);

https://figshare.com/s/ef7f40371b487b9dabca.

Supplemental Materials for: Male chickadees with better spatial cognition sire more extra-pair

young

Authors: Carrie L Branch, Benjamin R Sonnenberg, Joseph F Welklin, Bronwyn G Butcher,

Virginia K Heinen, Angela M Pitera, Lauren M Benedict, Eli S Bridge, Irby J Lovette, Michael S

Webster, Vladimir V Pravosudov

1. Summary of previous work on Sierra Nevada mountain chickadee study system:

The hypothesis presented in the current paper is based on established results in this system, all

collected over multiple years and including different groupings of birds:

1. Survival of juveniles in their first year of life is dependent on their spatial cognitive

abilities, representing directional selection (40).

2. Life span of chickadees depends on their spatial cognitive abilities - birds with better

spatial cognitive abilities live longer (41).

3. Spatial cognitive performance does not change with age (40, 52).

4. Spatial cognitive abilities do not show age-related senescence (45).

5. Spatial cognitive abilities are not associated with social dominance status (78) or with

naturally occurring developmental perturbations (86).

6. Individual variation in spatial cognitive abilities is highly heritable and is associated with

differences in genes related to cognition and brain development (43, 44).

7. Genotypes of birds with the worst cognition disappear from populations before winter

(e.g., homozygous recessive birds), likely due to mortality, yet these genotypes must have

been present at fledging (43, 44), which shows that birds with worse cognition die early

on in their first winter.

8. There is no detectable senescence in reproduction within the naturally occurring lifespan

of chickadees and females increase their reproductive success with age (70).

9. Female cognition is not associated with the number of fledglings (42) or fledgling mass

(current study), however, in a high resource year, male cognition is associated with larger

clutch and brood size in their social nest - a finding driven by female behavior (42).

10. Chickadees are socially monogamous species and form long-term social pairs - divorces

are rare (87). In addition, it is well established that females leave their territories and visit

neighboring male territories for EPCs, which we confirm here, and forced copulation

does not occur in chickadees.

11. Social pairing is not associated with spatial learning and memory (46), likely because

chickadees form social pairs in Fall and new recruits do not have the choice of any mate -

many males are already paired (46, 87), as such the primary way to increase indirect

fitness is via EPC.

12. There are no differences in spatial cognitive abilities between males and females (79), as

both rely on spatial cognition to survive the winter.

13. We have shown that supplementary food does not affect reproductive success in our

system (88).

2. SUPPLEMENTAL RESULTS

Across 3 years of data collection on EPCs and reproductive output, we see that on average, there

are 1.7 EPY per nest, however, if we constrain this analysis to only nests with at least one EPY,

the mean is 2.5 EPY per nest. Nests ranged from 0% EPY to 100% EPY, with the most being 6/6

nestlings sired by extra-pair fathers.

Elevation by year comparisons

Elevation by year comparison of extra-pair rates using binomial generalized mixed effects

modeling revealed a significant interaction between elevation and year, suggesting the direction

of the differences in extra-pair paternity rate between the high and low elevation sites differed

across years (Figure S1; see Table S1 for sample sizes by year and elevation). Therefore, we ran

a separate model for each year. In 2018, the extra-pair paternity rate was higher at the low

elevation site, but this difference was not significant (elevation term: χ^2_1=2.24, P=0.134, N=209

offspring (134 at high, 75 at low)). In 2020, the extra-pair paternity rate was higher at the high

elevation site, but this difference was not statistically significant (elevation term: χ^2_1=2.45,

P=0.118, N=263 offspring (119 at high, 144 at low)). In 2021, the extra-pair paternity rate was

higher at the low elevation site, but this difference was not statistically significant (elevation

term: χ^2_1=1.72, P=0.190, N=260 offspring (141 at high, 119 at low)).

Does male spatial cognition differ between nests that had and did not have extra-pair

young?

Within nests that had spatial cognitive data for the social male, approximately 28% of nests

(28/101) had no extra-pair young and 72% of nests had at least one extra-pair young (73/101).

The spatial cognitive abilities of males that had EPY in their nests did not differ from males that

had no EPY in their nests (Mean location errors per trial over first 20 trials: mean 1.08 errors vs.

1.01 errors; $\chi^2_1=0.42$, P=0.516, Mean location errors per trial over the entire task: mean 0.38

errors vs. 0.34 errors; $\chi^2_1=2.07$, P=0.150, N=101 nests).

Does male age affect extra-pair paternity?

Male age was not associated with the number of EPY a male sired, both according to a zero-

inflated model including males that sired no extra-pair young (Age as fixed effect in the first 20

trials model: $\chi^2_1=0.83$, P=0.362; Age as fixed effect in the entire task model: $\chi^2_1=2.41$, P=0.121,

N=280 (137 individual males)), and using a model only including males that sired at least one

EPY (Age as fixed effect in the first 20 trials model: $\chi^2_1=1.29$, P=0.255; Age as fixed effect in the

entire task model: $\chi^2_1=0.16$, P=0.687, N=41nests; 34 individual males).

Male age did not affect the likelihood of a male being cuckolded (Offspring level: Age as

fixed effect in the first 20 trials model: $\chi^2_1=0.08$, P=0.783, Age as fixed effect in the entire task

model: $\chi^2_1=0.18$, P=0.676, N=348 offspring for 42 males; Nest level: Age as fixed effect in the

first 20 trials model: $\chi^2_1=0.09$, P=0.768; Age as fixed effect in the entire task model: $\chi^2_1=2.02$,

P=0.570, N=59 nests for 42 males). Within nests that had at least one EPY, social male age did

not predict the number of EPY in the nest ($\chi^2_1=0.23$, P=0.635, N=41 nests; 35 individual males).

3. SUPPLEMENTAL FIGURES

Figure S1. Extra-pair (EPY) and within-pair (WPY) paternity rates across 3 years of data

collection and two elevation sites. Number insets represent the number of EPY and WPY for

each year and elevation. EPY rates did not differ significantly by year or elevation.

926
927
928   Figure S2. Histogram plotting the distance of extra-pair male nests from the females' nest. Mean
929   = 167 meters and Median = 136 meters. Range is 56 to 1022 meters between nestboxes.
930

931
Figure S3a. High elevation extra-pair paternity relationships and cognition scores. Maps show
nests with genotyped offspring and the nests of males that sired extra-pair offspring. Red arrows
show extra-pair paternity relationships. The direction of the arrow indicates a social male's nest
at the blunt end connected to the nest he sired extra-pair young in at the sharp end. Arrows that
originate in open space show males that sired extra-pair young but did not breed in our nest
boxes in that year (they likely bred in a natural cavity). Point colors represent the cognition
scores of the social male at each nest. Gray circles represent males without cognition data. Only
nests that included EPY are included on this map, which is a sample of the active nests in each
year (Figure S4).
941
942
943

944
945
Figure S3b. Low elevation extra-pair paternity relationships and cognition scores. Maps show
nests with genotyped offspring and the nests of males that sired extra-pair offspring. Red arrows
show extra-pair paternity relationships. The direction of the arrow indicates a social male's nest
at the blunt end connected to the nest he sired extra-pair young in at the sharp end. Arrows that
originate in open space show males that sired extra-pair young but did not breed in our nest
boxes in that year (they likely bred in a natural cavity). Point colors represent the cognition
scores of the social male at each nest. Gray circles represent males without cognition data. Only
nests that included EPY are included on this map, which is a sample of the active nests in each
year (Figure S4).
955
956
957

Figure S4. Heat maps of male spatial cognitive scores for high and low elevation nestboxes across the 3 years we have extra-pair paternity data from, including those nests that failed before day 16 or were not genotyped. Each point represents a nestbox. Color gradient represents mean location errors of male on the first 20 trials of the spatial cognitive task. Gray circles indicate nestboxes with no cognitive data for the male. Data shows that nestboxes are not clustered by male cognitive performance. Some of the males on the maps did not participate in cognitive testing until after these breeding seasons, but since spatial cognitive ability does not change with age, we present their scores here as well.

969
Figure S5. Model predictions of the number of extra-pair young a male sired in a year by the
mean number of location errors per trial over the 4-day testing period. The dashed line represents
a zero-inflated model for all males, including those with no extra-pair young (N=280; 137
individual males), and the solid line represents the model prediction for males that sired at least
one extra-pair young (N=41; 34 individual males). Shading represents bootstrapped 95%
confidence intervals. Note: data presented in figure do not account for total trials completed,
which was accounted for in statistical analysis.
977

Figure S6. Comparing spatial cognitive performance (mean number of location errors per trial over the 4-day task) of extra-pair and social fathers (total of 71 comparisons). Lines connect males by offspring: blue lines show comparisons where the extra-pair father performed better than the social father (34 of 71 cases or 47.9%), and purple lines show comparisons where the social father performed better than the extra-pair father (37 of 71 cases or 52.1%). Shaded distributions represent smoothed histograms of cognitive performance. Note that the data presented in the figure do not account for the total number of trials completed, which was accounted for in the statistical analysis.

991
992
993 Figure S7a. There is a strong positive correlation between the number of location errors on the
994 entire 4-day task and on the first 20 trials for males completing the task (Coef=0.27, F=121.72,
995 P<0.001).
996
997

998
999 Figure S7b. Representative learning curve of 19 males included in the extra-pair paternity
1000 analysis, from the 2021 breeding season. Dotted line represented chance level performance on
1001 the task at 3.5 mean location errors. Curve includes performance on trials 1 – 20 and over the
1002 entire 4-day task. Errors bars represent standard error of the mean.
1003

1004

1005

1006   8A.
1007

1008
1009   8B.
1010
1011

8C.

Figure S8. Comparisons of Trio LOD scores for 1st and 2nd ranked potential fathers for each offspring by elevation and year. Note the mostly bimodal distribution in which 1st-ranked males (blue) consistently out-performed 2nd-ranked males (red). Offspring with low 1st-ranked male Trio LOD scores are cases where we did not sample the offspring's genetic father.

4. SUPPLEMENTAL TABLES

1040

1041 Table S1. Sample sizes for extra-pair paternity broken down by year and elevation for both the
1042 nest and offspring level.

	Year	Elevation	No EPY	≥1 EPY	Rate
Nest level					
	2018	High	8	17	68.00%
		Low	1	13	92.90%
	2020	High	6	18	75.00%
		Low	8	14	63.60%
	2021	High	9	13	59.10%
		Low	6	14	70.00%
	Total	High	23	48	67.60%
		Low	15	41	73.20%
Offspring level	Year	Elevation	WPY	EPY	Rate
	2018	High	89	45	33.60%
		Low	38	37	49.30%
	2020	High	76	43	36.10%
		Low	110	34	23.60%
	2021	High	113	28	19.90%
		Low	84	35	29.40%
	Total	High	278	116	29.40%
		Low	232	106	31.40%

1059 Table S2. Differences in extra-pair paternity rates between high and low elevations. A likelihood
1060 ratio test indicated that an interaction between elevation and year improved the model, so years
1061 were run separately.
1062
1063 2018

	Estimate	Standard error	Z value
Intercept	-0.85	0.35	-2.42
Elevation (Low)	0.84	0.56	1.50
Random effects	Variance	Std deviation	
Mother ID	1.66	1.29	

1064 N=209 offspring, 39 mothers
1065
1066 2020

	Estimate	Standard error	Z value
Intercept	-0.72	0.39	-1.86
Elevation (Low)	-0.88	0.56	-1.57
Random effects	Variance	Std deviation	
Mother ID	2.15	1.47	

1067 N=263 offspring, 46 mothers
1068
1069 2021

	Estimate	Standard error	Z value
Intercept	-1.79	0.40	-4.53
Elevation (Low)	0.69	0.53	1.31
Random effects	Variance	Std deviation	
Mother ID	1.56	1.25	

1070 N=260 offspring, 42 mothers
1071
1072
1073

1074 Table S3. Relationship between spatial cognitive ability and extra-pair young sired. Two models
1075 are presented for each spatial cognitive measure, a model that included a zero-inflation parameter
1076 for all observations and a model that only included male-years where the male sired at least one
1077 offspring.
1078

1079 **3A. First 20 trials mean location error with zero-inflation parameter**

Conditional model	Estimate	Standard error	Z value
Intercept	0.40	0.54	0.75
First 20 trials mean location error	-0.92	0.37	-2.51
Elevation (Low)	0.61	0.31	1.99
Year 2020	0.67	0.43	1.54
Year 2021	0.74	0.43	1.70
Zero-inflation model	Estimate	Standard error	Z value
Intercept	1.28	0.23	5.55
Random effects	Variance	Std deviation	
Male ID	0.08	0.29	

1080 N=280 male-years (137 unique males over 3 breeding seasons)
1081

1082 **3B. First 20 trials mean location error with positive-only values**

	Estimate	Standard error	Z value
Intercept	0.50	0.63	0.79
First 20 trials mean location error	-1.16	0.48	-2.44
Elevation (Low)	0.74	0.34	2.17
Year 2020	0.73	0.55	1.33
Year 2021	0.76	0.55	1.38
Random effects	Variance	Std deviation	
Male ID	0.11	0.33	

1083 N=41 male-years (34 unique males over 3 breeding seasons)
1084

1085 **3C. Entire task mean location error with zero-inflation parameter**

Conditional model	Estimate	Standard error	Z value
Intercept	0.18	0.46	0.40
Entire task mean location error	-1.86	0.65	-2.85
Elevation (Low)	0.76	0.29	2.65
Year 2020	0.73	0.42	1.75
Year 2021	0.79	0.41	1.91
Zero-inflation model	Estimate	Standard error	Z value
Intercept	1.34	0.22	6.07
Random effects	Variance	Std deviation	
Male ID	0.03	0.17	

1086 N=280 male-years (137 unique males over 3 breeding seasons)

3D. Entire task mean location error with positive-only values

	Estimate	Standard error	Z value
Intercept	0.24	0.56	0.43
Entire task mean location error	-2.55	0.92	-2.77
Elevation (Low)	0.85	0.32	2.65
Year 2020	0.81	0.54	1.50
Year 2021	0.88	0.54	1.64
Random effects	Variance	Std deviation	
Male ID	0.03	0.18	

N=41 male-years (34 unique males over 3 breeding seasons)

1098 Table S4. Relationship between male and female spatial cognitive ability (mean location errors
1099 per trial over the first 20 trials) and breeding performance.
1100
1101
1102 **4A. Brood size**

	Estimate	Standard error	Z value
Intercept	1.75	0.07	26.07
Male First 20 trials mean location error	0.06	0.04	1.52
Female First 20 trials mean location error	-0.05	0.04	-1.32
Elevation (Low)	0.08	0.04	2.10
Random effects	Variance	Std deviation	
Year (2015-2023)	<0.01	0.08	
Male ID	<0.01	0.09	
Female ID	<0.01	0.06	

1103 N=205 nests over 9 years for 108 males and 89 females.
1104
1105
1106 **4B. Brood size relative to clutch size**

	Estimate	Standard error	Z value
Intercept	0.72	0.10	7.39
Male First 20 trials mean location error	0.02	0.02	1.00
Female First 20 trials mean location error	0.02	0.03	0.70
Elevation (Low)	0.02	0.02	0.97
Clutch size	0.14	0.01	12.05
Random effects	Variance	Std deviation	
Year (2015-2023)	<0.01	0.04	
Male ID	<0.01	<0.01	
Female ID	<0.01	<0.01	

1107 N=205 nests over 9 years for 108 males and 89 females.
1108
1109 **4C. Mean nestling mass**

	Estimate	Standard error	Df	T value
Intercept	12.47	0.25	67.48	50.35
Male first 20 trials mean location error	-0.28	0.13	94.91	-2.20
Female first 20 trials mean location error	0.15	0.15	46.89	0.97
Elevation (Low)	-0.38	0.13	56.62	-2.86
Random effects	Variance	Std deviation		
Year (2015-2023)	0.10	0.31		
Male ID	0.04	0.19		
Female ID	0.13	0.35		
Residual	0.41	0.64		

1110 N=202 nests over 9 years for 107 males and 89 females.
1111
1112
1113

1114   Table S5. Test of whether siring extra-pair offspring affected the offspring in a male's own nest.
1115   Whether a male sired extra-pair offspring or not was compared to A) brood size, B) brood size
1116   controlled for clutch size, and C) mean nestling mass.
1117
1118   **5A. Brood size**

	Estimate	Standard error	Z value
Intercept	1.78	0.04	45.66
Sired EPY (Yes)	0.03	0.05	0.55
Elevation (Low)	0.04	0.04	0.95
Year 2020	-0.06	0.04	-1.33
Year 2021	0.06	0.04	1.34
Random effects	Variance	Std deviation	
Male ID	0.01	0.12	
Female ID	<0.01	0.04	

1119   N=144 nests for 99 males and 104 females
1120
1121   **5B. Brood size relative to clutch size**

	Estimate	Standard error	Z value
Intercept	0.71	0.10	7.09
Sired EPY (Yes)	<0.01	0.03	0.03
Elevation (Low)	<-0.01	0.03	-0.17
Clutch size	0.15	0.01	11.41
Year 2020	-0.03	0.03	-0.86
Year 2021	-0.01	0.03	-0.26
Random effects	Variance	Std deviation	
Male ID	<0.01	<0.01	
Female ID	<0.01	<0.01	

1122   N=144 nests for 99 males and 104 females
1123
1124   **5C. Mean nestling mass**

	Estimate	Standard error	Df	T value
Intercept	12.41	0.14	134.89	87.98
Sired EPY (Yes)	0.26	0.18	134.89	1.45
Elevation (Low)	-0.32	0.16	85.75	-1.94
Year 2020	-0.68	0.17	115.82	-3.94
Year 2021	-0.14	0.17	115.13	-0.82
Random effects	Variance	Std deviation		
Male ID	0.12	0.35		
Female ID	0.00	0.00		
Residual	0.61	0.78		

1125   N=140 nests for 98 males and 101 females
1126
1127
1128

1129 Table S6. Test of whether offspring sired by extra-pair fathers differed in mass from offspring
1130 sired by social males.
1131

	Estimate	Standard error	Df	T value
Intercept	12.44	0.15	135.23	82.34
Extra-pair (Yes)	0.14	0.08	694.45	1.68
Elevation (Low)	-0.50	0.15	121.56	-3.25
Year 2020	-0.71	0.19	124.70	-3.80
Year 2021	-0.16	0.19	122.26	-0.83
Random effects	Variance	Std deviation		
Nest ID	0.59	0.77		
Residual	0.72	0.85		

1132 N=719 offspring from 125 nests
1133
1134
1135
1136

1137 Table S7. Comparison of spatial cognition scores between social males and extra-pair males they
1138 were cuckolded by. Log-transformed values are reported in Table 4B – entire task mean location
1139 error model.

1140

1141 7A. First 20 trials mean location error model

	Estimate	Standard error	Df	T value
Intercept	1.00	0.09	37.00	11.21
Father type (social)	0.21	0.05	108.11	3.85
Elevation (Low)	-0.10	0.13	27.36	-0.73
Random effects	Variance	Std deviation		
Offspring ID	0.00	0.00		
Mother ID	0.10	0.32		
Residual	0.11	0.32		

1142 N=71 social male – extra-pair male comparisons (71 offspring) from 32 mothers

1143

1144 7B. Entire task mean location error model

	Estimate	Standard error	Df	T value
Intercept	-1.07	0.06	42.35	-16.78
Father type (social)	0.11	0.05	107.68	2.44
Elevation (Low)	-0.17	0.09	27.30	-1.93
Trials completed	-4.80	0.33	136.81	-14.63
Trials completed^2	1.67	0.33	136.51	5.03
Random effects	Variance	Std deviation		
Offspring ID	0.00	0.00		
Mother ID	0.04	0.19		
Residual	0.08	0.28		

1145 N=71 social male – extra-pair male comparisons (71 offspring) from 32 mothers

1146
1147
1148

1149 Table S8. Test of whether a male's spatial cognition score was associated with whether he was
1150 cuckolded or not.
1151

1152 **8A. First 20 trials mean location error offspring model**

	Estimate	Standard error	Z value
Intercept	-2.05	0.75	-2.74
Male first 20 trials mean location error	-0.20	0.40	-0.50
Female first 20 trials mean location error	1.23	0.49	2.49
Elevation (Low)	0.52	0.41	1.26
Year 2020	-0.83	0.38	-2.15
Year 2021	-0.55	0.39	-1.40
Random effects	Variance	Std deviation	
Male ID	0.23	0.48	
Female ID	0.41	0.64	

1153 N=348 offspring from 42 females and 42 males over 3 breeding seasons
1154

1155 **8B. Entire task mean location error offspring model**

	Estimate	Standard error	Z value
Intercept	-0.79	0.61	-1.31
Male entire task mean location error	-0.67	1.02	-0.65
Female entire task mean location error	0.41	0.97	0.42
Elevation (Low)	0.16	0.44	0.37
Year 2020	-0.81	0.40	-2.02
Year 2021	-0.53	0.40	-1.32
Random effects	Variance	Std deviation	
Male ID	0.05	0.22	
Female ID	0.86	0.93	

1156 N=348 offspring from 42 females and 42 males over 3 breeding seasons
1157

1158 **8C. First 20 trials mean location error nest model**

	Estimate	Standard error	Z value
Intercept	0.35	1.11	0.32
Male first 20 trials mean location error	0.16	0.67	0.24
Female first 20 trials mean location error	0.35	0.76	0.46
Elevation (Low)	0.20	0.64	0.30
Year 2020	0.21	0.73	0.29
Year 2021	-0.73	0.72	-1.01
Random effects	Variance	Std deviation	
Male ID	<0.01	<0.01	
Female ID	<0.01	<0.01	

1159 N=59 nests from 42 females and 42 males over 3 breeding seasons
1160

1161 **8D. Entire task mean location error nest model**

	Estimate	Standard error	Z value
Intercept	0.82	0.91	0.90
Male entire task mean location error	0.80	1.72	0.47
Female entire task mean location error	-0.52	1.48	-0.35
Elevation (Low)	0.07	0.62	0.11
Year 2020	0.32	0.73	0.44
Year 2021	-0.71	0.73	-0.98
Random effects	Variance	Std deviation	
Male ID	<0.01	<0.01	
Female ID	<0.01	<0.01	

1162 N=59 nests from 42 mothers and 42 fathers over 3 breeding seasons

1163

1164

1165 Table S9. Test of whether the number of extra-pair young in nests with at least one extra-pair
1166 young is associated with spatial cognitive ability (performance on the first 20 trials).
1167

	Estimate	Standard error	Z value
Intercept	-0.26	0.55	-0.48
Male First 20 trials mean location error	-0.30	0.30	-1.01
Female First 20 trials mean location error	0.92	0.35	2.61
Elevation (Low)	0.79	0.35	2.23
Year 2020	-0.70	0.34	-2.09
Year 2021	0.09	0.31	0.31
Random effects	Variance	Std deviation	
Father ID	<0.01	<0.01	
Mother ID	<0.01	<0.01	

1168   N=41 nests with extra-pair offspring from 35 males and 32 females.

1169
1170

1171
1172

1173

1174

1175

1176

1177

1178
1179

1180
1181

1182