Effects of chemical modulators on enzyme specificity

Andrew D. Hecht[*,†,‡,¶] and Oleg A. Igoshin[*,†,‡,¶,§,∥,⊥]

†Department of Bioengineering, Rice University, Houston, TX 77005, USA

‡Center for Theoretical Biological Physics, Rice University, Houston, TX 77005, USA

¶Rice Synthetic Biology Institute, Rice University, Houston, TX 77005, USA

§Department of BioSciences, Rice University, Houston, TX 77005, USA

∥Department of Chemistry, Rice University, Houston, TX 77005, USA

⊥To whom correspondence should be addressed

E-mail: ahecht@rice.edu; igoshin@rice.edu

Abstract

Chemical inhibitors bind to enzymes, thereby inhibiting their catalytic activity. While many enzymes catalyze reactions with a single substrate, others, like DNA polymerase, can act on multiple related substrates. Substrate-selective inhibitors (SSIs) target these multi-substrate enzymes to modulate their specificity. Although SSIs hold promise as therapeutics, our theoretical understanding of how different inhibitors influence enzyme specificity remains limited. In this study, we examine enzyme selectivity within kinetic networks corresponding to known inhibition mechanisms. We demonstrate that competitive and uncompetitive inhibitors do not affect substrate specificity, regardless of rate constants. In contrast, noncompetitive and mixed inhibition can alter specificity and can lead to non-monotonic responses to the inhibitor. We show that mixed and non-competitive inhibitors achieve substrate-selective inhibition by altering the effective free-energy barriers of product formation pathways that are enabled by the inhibitor's presence. We then apply this framework to the Sirtuin-family deacylase SIRT2, showing that the suicide inhibitor thiomyristoyl lysine (TM) cannot influence substrate specificity unless there is a direct substrate exchange reaction or biochemical constraints are relaxed. These findings provide insights into engineering systems where cofactor binding modulates metabolic flux ratios.

Introduction

Enzymes are proteins that catalyze chemical reactions involving one or more substrates, thereby playing key roles in cellular function. Given the importance of enzymes in biology, many pharmaceutical drugs target enzymatic processes to produce medically beneficial effects; these drugs are often chemical inhibitors that can slow or halt the activity of a target enzyme. For example, nonsteroidal anti-inflammatory drugs (NSAIDs) act as inhibitors of the cyclooxygenases COX-1 and COX-2 and reduce the production of pro-inflammatory prostaglandins.[1] Tyrosine kinase inhibitors[2,3] and histone deacetylase inhibitors[4] have been

used as cancer therapies to promote apoptosis in cancer cells.

While many enzymes act on only a single substrate, in some cases, enzymes such as DNA polymerase can catalyze reactions involving various chemically similar substrates. For these multi-substrate enzymes, each unique substrate can lead to a corresponding product. Importantly, an enzyme is not necessarily equally efficient with all substrates; depending on the enzyme's ability to discriminate between substrates, some products will be more prevalent than others even with equal substrate concentrations.

Inhibitors that can affect the substrate selectivity of the targeted enzyme, known as substrate-selective enzyme inhibitors (SSIs), are becoming increasingly important and represent a novel class of drugs. In one example, Maianti et al. carried out a high-throughput screen to identify SSIs capable of preferentially inhibiting insulin degradation by insulin-degrading enzyme (IDE) without affecting glucagon degradation.[5] As insulin and glucagon have opposite effects on blood glucose levels, SSIs targeting IDE could serve as a new class of drug for treating type II diabetes. SSIs are also relevant to basic research, as potential confounding factors for reverse chemical genetics screens.[6]

Kinetic modeling can provide insights into how enzyme inhibition affects substrate specificity. Previously, Hendricks et al. developed a kinetic model of the kinase p38 to explain substrate-selective inhibition of substrate phosphorylation.[7] However, to our knowledge, there has been no comprehensive theoretical study on the effects of different enzyme inhibition modalities on substrate specificity.

Prior work by Mallory et al.[8] demonstrated that kinetic barrier perturbations are required to modulate the steady-state ratio of reaction fluxes for alternative biochemical pathway branches. On the other hand, merely changing the stability of states is insufficient to affect the flux ratio.[8] These findings imply that inhibitors that operate by effectively stabilizing certain states will not be substrate-selective.

Here, we introduce kinetic models of common inhibition modalities for multi-substrate enzymes to examine the possible effects of inhibition on substrate selectivity. We demon-

strate that competitive and uncompetitive inhibitors can never affect substrate selectivity, regardless of the underlying kinetic parameters governing the system. In contrast, mixed and partial inhibition modes can not only affect substrate selectivity, but can also yield non-monotonic effects where the substrate selection error can exceed the limiting bounds for intermediate inhibitor concentrations. We examine the thermodynamic basis for substrate-selective inhibition by mixed and partial inhibitors, and show that substrate selectivity can only change when the presence of the inhibitor leads to an alternative product formation pathway with potentially different kinetic barriers. Finally, we construct a kinetic model of substrate selection by the Sirtuin-family de-acylase SIRT2 and show that either direct substrate exchange reactions or a relaxed, unordered binding mechanism are required to observe substrate-selective inhibition.

Theoretical Methods

Kinetic Models of Enzyme Inhibition Modes

Reversible enzyme inhibitors are primarily classified by the inhibition mechanism, which is determined by the structure of both the enzyme and the inhibitor molecule. The most basic form of inhibition is competitive inhibition (Figure 1A), in which the inhibitor directly competes with substrate for access to the enzyme active site.[9] This competition effectively raises the enzyme Michaelis-Menten constant, K_M, i.e., higher substrate concentrations are required to achieve the same reaction velocity. In contrast to competitive inhibitors that bind to the free enzyme, uncompetitive inhibitors bind directly to the enzyme-substrate complex (Fig. 1B) and block the catalytic transformation. Uncompetitive inhibitors generally reduce the maximum enzyme velocity, V_max, but can also affect the observed K_M,.[9]

Mixed inhibitors combine features from competitive and uncompetitive inhibition, and can bind to both the free enzyme and the enzyme-substrate complex (Figure 1). If the inhibitor binds equally well to both enzyme states, the inhibitor is said to be a non-competitive

inhibitor, which is a special case of mixed inhibition. Both mixed and non-competitive inhibitors can affect the enzyme V_{max}, however, only mixed inhibition can affect the apparent K_{M}.[9] Finally, inhibition is occasionally incomplete, leading to partial inhibition (Fig. 1D). In this case, residual catalytic activity can be observed even when fully bound to inhibitor.[10]

Figure 1: Kinetic schemes for competitive (A), uncompetitive (B), mixed (C), and partial inhibition (D). Product formation reactions are shown in blue.

To examine the effects of each inhibition mode on substrate selectivity, we construct kinetic models for each of the schemes in Figure 1; the models are built using techniques previously used to study substrate selection errors in various biological systems,[11–13] and allow us to examine the behavior of each inhibition mode analytically. While in principle, enzymes can have a large number of possible substrates, we limit ourselves to two substrates,

R and W, for simplicity.

The forward chemical master equation allows us to compute the likelihood of observing each state in the model, and is defined by a system of ordinary differential equations for each fixed value of inhibitor concentration, x:

$$\frac{\mathrm{d}\vec{P}(x,t)}{\mathrm{d}t} = \mathbf{K}(x)\vec{P}(t). \tag{1}$$

with the normalization constraint

$$\sum_i P_i(x,t) = 1 \tag{2}$$

The matrix $\mathbf{K}$ encodes the state transitions for each system, with $\mathbf{K}_{ji} = k_{ij}$ for each transition $i \rightarrow j$. We assumed substrate, cofactor, and inhibitor concentrations are kept constant and absorbed these into the rate constants. We only keep the dependence on the inhibitor concentration (x) explicit. A detailed description of each of the inhibition models is provided in the Supplemental Information.

The steady-state probabilities $\vec{P}_{\mathrm{ss}}(x)$ can be computed by enforcing the additional constraint

$$\frac{\mathrm{d}\vec{P}(x,t)}{\mathrm{d}t} = [0, 0, \cdots, 0]^{\mathrm{T}}. \tag{3}$$

The fluxes for each reaction in the system can be readily computed from the system probabilities; for a given reaction $i \rightarrow p$, the flux is simply $J_i = k_{ip}P_i$. The total product formation flux for each substrate can be defined as a sum over all possible product formation reactions; for a substrate with N such reactions, the total flux is simply

$$\hat{J} = \sum_i^N J_i \tag{4}$$

with the ratio of fluxes for two substrates, R and W, being given as

$$\eta = \frac{\hat{J}_W}{\hat{J}_R}. \tag{5}$$

The flux ratio indicates the relative catalytic efficiency for the two substrates; as the flux ratio decreases, the enzyme increasingly favors substrate R over substrate W.

Kinetic schemes that contain thermodynamic (non-dissipative) reaction cycles, such as the mixed and partial inhibition models (Figure 1C,D), are additionally constrained by detailed balance. Detailed balance constrains the rate constants for reactions along the cycle such that the forward and reverse rates for each reaction i are related by

$$\prod_i \frac{k_i}{k_{-i}} = 1. \tag{6}$$

Parameter Sampling

To examine the prevalence of non-monotonic dose-response curves, model parameters were sampled in log-space using the MATLAB function `rand`. Given upper and lower bounds of a and b, for each N parameter model we computed a parameter set $\vec{r}$ as

```
r = 10 .^ (b + (b - a) .* rand(N,1));
```

Generally, the rate constants are sampled from the range $[10^{-4}, 10^4]$ for all models. The model discrimination factors are sampled separately, with the full bounds given in Supplemental Table 1.

For simplicity, we require that the saturating bound (η_∞) be lower than the uninhibited bound (η_0). In the case of mixed inhibition, this is satisfied for all parameter sets with $f_p < f_{off}$; however, we were unable to derive a corresponding constraint for partial inhibition. Therefore, we enforced the constraint by rejecting parameter sets that fail to satisfy $\eta_\infty < \eta_0$. However, we still set the discrimination in catalysis, f_{cat} based on the sampled f'_{off} by scaling f'_{off} by a number sampled from the uniform distribution, i.e. $f_{cat} = U_{[0,1]} f'_{off}$. We also assume $f_{cat} = f'_{cat}$ for partial inhibition.

Detecting Undershoot and Overshoot

For each random parameter set, we first compute the uninhibited and saturation bounds η_0 and η_∞. Next, starting with an initial concentration range $x \in [x_i, x_j]$, we compute the flux ratio $\eta(x)$. In order to ensure that we capture the full range of behavior for each parameter set, we progressively expand the concentration range until the computed flux ratios are sufficiently close to the limits η_0 and η_∞.

Undershoot events occur when the flux ratio $\eta(x)$ becomes smaller than the saturating limit η_∞ for intermediate inhibitor concentrations. For a given parameter set, the total undershoot q_{under} is computed as the difference

$$q_{under} = \min\left(\eta\right) - \eta_\infty. \tag{7}$$

Similarly, we can compute the total overshoot q_{over} for a given parameter set as

$$q_{over} = \max\left(\eta\right) - \eta_0, \tag{8}$$

where we now compare the maximum flux ratio with the uninhibited limit.

In all cases, we enforce a threshold $\epsilon = 0.01$ to avoid parameter sets that only display a slight undershoot or overshoot; when plotting, all parameter sets with undershoot or overshoot under the threshold ϵ are discarded.

Results

Competitive and Uncompetitive Inhibitors Cannot Affect Specificity

First, we determined how the flux ratio η changes as a function of inhibitor concentration x. When the inhibitor is substrate-selective, we expect to see the flux ratio change with

increasing inhibitor concentration, with the derivative $\frac{d\eta(x)}{dx} \neq 0$ for at least some range of concentrations.

For competitive and uncompetitive inhibition (Figure 1A,B) we find that the concentration of inhibitor does not affect the product flux ratio (Figure 2A,B). Specifically, the flux ratio η is independent of the inhibitor concentration x for these inhibition modes and has the same form in both cases:

$$\eta = \frac{f_{\text{on}} f_p \left(k_{\text{off}} + k_p \right)}{f_{\text{off}} k_{\text{off}} + f_p k_p} \tag{9}$$

Thus, regardless of kinetic parameters, the inhibitor cannot affect substrate selectivity for competitive or uncompetitive inhibition. Notably, an uncompetitive inhibitor that only inhibits one of the substrates (i.e., only binds to EW to form EWI but not to ER) is still incapable of affecting selectivity.

In contrast, both mixed inhibition (Figure 1C) and partial inhibition (Figure 1D) can affect substrate selectivity, and are therefore capable of being substrate-selective inhibitors. In both cases, the structure of the reaction network involves reaction cycles (loops), which we assume are non-dissipative, i.e., their rate constants obey the detailed-balance constraint (Eq. 6).

Thermodynamic Basis for Substrate Selective Inhibition

To better understand the connections between our results and those of Mallory et al.,[8] we can consider the coarse-graining, where we cannot explicitly detect enzyme states bound to the inhibitor. In this case, we will be unable to differentiate between states E and EI, ER and ERI, and other pairs of states for more complicated kinetic schemes. In the corresponding coarse-grained model, the pairs of states are lumped together to yield a simplified reaction scheme that can be easily analyzed.

For example, the competitive inhibition scheme (Figure 1A) can be coarse-grained to

Figure 2: Competitive (A) and uncompetitive (B) inhibitors cannot affect substrate selectivity, no matter the underlying kinetic parameters. In contrast, both mixed (C) and partial inhibition (D) can yield substrate-selective inhibition.

₁ yield the effective reaction scheme

$$\text{E}' + \text{P}_{\text{W}} \longleftarrow \text{EW} \rightleftharpoons \text{E}' \rightleftharpoons \text{ER} \longrightarrow \text{E}' + \text{P}_{\text{R}}. \tag{10}$$

₂ Here, we assume that the free enzyme (E) and the inhibited enzyme (EI) are indistinguish-

1   able, and are thus represented by the composite state (E′) so that

$$[E'] = [E] + [EI]. \tag{11}$$

2   At steady-state, we have

$$[E'] = [E] \left(1 + \frac{x}{K_I} \right) \tag{12}$$

3   , where K_I is the association constant for inhibitor binding,

$$K_I = \frac{k_b}{k_u} = \frac{1}{x} \frac{[EI]}{[E]}. \tag{13}$$

4   The effective substrate binding rates for the coarse-grained network are obtained by re-
5   normalizing the original rates. By re-normalizing the rates, the steady-state fluxes for each
6   reaction in the coarse-grained network are identical to those in the original network. The
7   effective substrate association rates for the coarse-grained network are given by

$$\hat{k}_{on}^R = k_{on}^R \frac{[E]}{[E']} = k_{on}^R \left(\frac{1}{1 + \frac{x}{K_I}} \right) \tag{14}$$

$$\hat{k}_{on}^W = k_{on}^W \frac{[E]}{[E']} = k_{on}^W \left(\frac{1}{1 + \frac{x}{K_I}} \right) \tag{15}$$

Here, increasing the inhibitor concentration effectively decreases the substrate association
rate by limiting the concentration of free enzyme. The enzyme-substrate dissociation rates
k_{off}^R and k_{off}^W are unaffected by the presence of inhibitor; thus, competitive inhibitors effectively
perturb the free-energy of the composite state [E′], and therefore the invariance of the flux
ratio holds.

The uncompetitive inhibition scheme (Figure 1B) can be coarse-grained in a similar
manner. Here, we make a similar assumption that the enzyme-substrate complexes (ER
and EW) cannot be distinguished from the enzyme-substrate-inhibitor complexes (ERI and

EWI), which leads to the effective reaction scheme

$$E + P_W \longleftarrow EW' \rightleftharpoons E \rightleftharpoons ER' \longrightarrow E + P_R. \tag{16}$$

The effective substrate dissociation rates for the coarse-grained network are given by

$$\hat{k}_{off}^R = k_{off}^R \frac{[E]}{[E']} = k_{off}^R \left(\frac{1}{1 + \frac{x}{K_I}} \right) \tag{17}$$

$$\hat{k}_{off}^W = k_{off}^W \frac{[E]}{[E']} = k_{off}^W \left(\frac{1}{1 + \frac{x}{K_I}} \right). \tag{18}$$

The effective catalytic rates for the coarse-grained network take a similar form and are given by

$$\hat{k}_{cat}^R = k_{cat}^R \frac{[E]}{[E']} = k_{cat}^R \left(\frac{1}{1 + \frac{x}{K_I}} \right) \tag{19}$$

$$\hat{k}_{cat}^W = k_{cat}^W \frac{[E]}{[E']} = k_{cat}^W \left(\frac{1}{1 + \frac{x}{K_I}} \right). \tag{20}$$

Changes in the inhibitor concentration effectively change the energies of the ER′ and EW′ states, with the catalytic and dissociation rates being scaled by the same factor. The flux ratio is therefore unaffected.

In the case of partial and mixed inhibition (Figure 1C,D), the inhibitor can bind to both the free enzyme (E) and the enzyme-substrate complex (ER and EW). Assuming again that the unbound and inhibitor-bound states cannot be distinguished, we can coarse-grain the network to yield the reduced scheme:

$$E' + P_W \longleftarrow EW' \rightleftharpoons E' \rightleftharpoons ER' \longrightarrow E' + P_R. \tag{21}$$

Below, we demonstrate that for this effective scheme, describing partial and mixed inhibition cases, changes in the inhibitor concentration alter the effective barrier heights in the coarse-

grained networks.

For simplicity, we assume the catalytic rates $k_{\text{cat}}^{\text{R}}$ and $k_{\text{cat}}^{\text{W}}$ are much slower than the other transitions. In that case, we can make a quasi-equilibrium approximation to compute the effective substrate binding and unbinding rates. In that limit, for the coarse-grained network, the effective substrate binding rates are given by

$$\hat{k}_{\text{E}'\rightarrow\text{ER}'} = k_{\text{on}}^{\text{R}} \left(\frac{1}{1 + \frac{x}{K_1}} \right) + k_{\text{on}}'^{\text{R}} \left(\frac{x}{1 + \frac{x}{K_2}} \right) \tag{22}$$

and

$$\hat{k}_{\text{E}'\rightarrow\text{EW}'} = k_{\text{on}}^{\text{W}} \left(\frac{1}{1 + \frac{x}{K_1}} \right) + k_{\text{on}}'^{\text{W}} \left(\frac{x}{1 + \frac{x}{K_3}} \right), \tag{23}$$

where K_1, K_2, and K_3 are the association constants for inhibitor binding to the free enzyme (E) and enzyme-substrate complexes (ER, EW), respectively.

Here, we see that the inhibitor affects the effective rate of both substrate binding and unbinding. By looking at the ratio of the effective binding rates for the R and W substrates

$$\ln\left(\frac{\hat{k}_{\text{E}'\rightarrow\text{EW}'}}{\hat{k}_{\text{E}'\rightarrow\text{ER}'}} \right) = \hat{\epsilon}_W^{\ddagger} - \hat{\epsilon}_R^{\ddagger}, \tag{24}$$

we can see how the inhibitor affects the relative free-energy barriers $\hat{\epsilon}_R^{\ddagger}$ and $\hat{\epsilon}_W^{\ddagger}$ for the R and W substrates, and thus the underlying thermodynamic landscape. Using the coarse-grained network, the ratio is given by

$$\frac{\hat{k}_{\text{E}'\rightarrow\text{EW}'}}{\hat{k}_{\text{E}'\rightarrow\text{ER}'}} = \frac{(K_2 + x)\left(K_3 x \left(K_1 + x\right) f_{\text{on}}' k_{\text{on}}'^{\text{R}} + K_1 f_{\text{on}} k_{\text{on}}^{\text{R}} \left(K_3 + x\right)\right)}{(K_3 + x)\left(K_2 x \left(K_1 + x\right) k_{\text{on}}^{\text{R}} + K_1 k_{\text{on}}^{\text{R}} \left(K_2 + x\right)\right)} \tag{25}$$

where each rate k_i^W is redefined in terms of discrimination factors as $k_i^W = f_i k_i^R$. We see that the ratio is a function of the inhibitor concentration x, which implies, consistent with the results of Mallory et al.,[8] that the inhibitor effectively acts as a perturbation affecting the relative free-energy barriers for the substrate binding reactions. Trivially, we can see that the inhibitor will be unable to affect the ratio of effective rates if both $K_2 = K_3$ and

$f_{\text{on}} = f'_{\text{on}} = 1$; the ratio will deviate from unity for some inhibitor concentration x if either constraint is relaxed. Ultimately, we require either reaction cycles involving substrate binding or the existence of multiple pathways to the product formation reaction(s) for the inhibitor to affect the ratio of effective transition rates for reactions involving different substrates, and consequently, the difference in effective free-energy barriers for said reactions.

Mixed and Partial Inhibition Can Yield Non-Monotonic Responses

Thus far, we have shown that mixed and partial inhibitors can affect enzymatic substrate selectivity. Beyond simply affecting substrate specificity, for some parameter regimes, it is possible to observe non-monotonic responses to changes in the inhibitor concentration. In these cases, the product formation flux ratio η can exceed the limits for the uninhibited enzyme (η_0) and the enzyme fully saturated with inhibitor (η_∞).

Two distinct classes of non-monotonic behaviors are possible depending on the particular model parameters; for simplicity, we constrain the parameters to ensure that the flux ratio decreases as a function of inhibitor concentration. First, for some parameterizations, an undershoot condition is observed, in which the substrate selection error can be improved beyond the saturating ($\eta < \eta_\infty$) limit for intermediate inhibitor concentrations (Figure 3A). Second, other parameterizations can display an overshoot, whereby the substrate selection error is higher than the uninhibited enzyme ($\eta > \eta_0$) for some intermediate inhibitor concentrations (Figure 3B). Beyond undershoot and overshoot, in other cases, the substrate selection error is less sensitive to changes in inhibitor concentration at sub-saturating concentrations, yielding a plateau-like effect (SI Figure 1).

The uninhibited limit η_0 for both mixed and partial inhibition is identical and corresponds to that of a basic Michaelis-Menten enzyme with two possible substrates,[14] which is

$$\lim_{x \to 0} \eta(x) \equiv \eta_0 = \frac{f_{\text{on}} f_{\text{p}}(k_{\text{off}} + k_{\text{p}})}{f_{\text{off}} k_{\text{off}} + f_{\text{p}} k_{\text{p}}}. \tag{26}$$

Figure 3: Substrate-selective enzyme inhibitors can, in principle, have non-monotonic effects on product flux ratios, including overshoot (A) and undershoot (B), where the flux ratio can exceed the limiting bounds. The non-monotonic response to inhibitor arises due to differential effects on the product formation fluxes (C) caused by changes in the effective substrate dissociation constants (D).

The expression for specificity is defined in terms of the rates for reactions involving the

desired substrate, R, as well as discrimination factors f_i. The discrimination factors are

defined for each reaction as $f_{ij} = k_{ij}^{W}/k_{ij}^{R}$; these factors determine the ability of the enzyme

to differentiate between substrates for each reaction.

In contrast, the saturation limit η_∞ varies between mixed and partial inhibition. For

mixed and non-competitive inhibition, the saturating limit is simply

$$\lim_{x \to \infty} \eta(x) \equiv \eta_\infty = f_\mathrm{p} \frac{f_\mathrm{on}}{f_\mathrm{off}}, \tag{27}$$

7   which is the discrimination in catalysis (f_p) scaled by the ratio of discrimination in sub-

8   strate binding and unbinding. The corresponding limit for partial inhibition is somewhat

more complicated; because the enzyme retains residual catalytic activity when bound to the inhibitor, the saturating limit for partial inhibition takes the same structural form as for a typical Michaelis-Menten enzyme (Eq. 26), but with kinetic rates and discrimination factors that in principle may differ from the uninhibited enzyme:

$$\lim_{x \to \infty} \eta(x) \equiv \eta_\infty = \frac{f'_{\mathrm{on}} f'_{\mathrm{p}} (k'_{\mathrm{off}} + k'_{\mathrm{p}})}{f'_{\mathrm{off}} k'_{\mathrm{off}} + f'_{\mathrm{p}} k'_{\mathrm{p}}}. \tag{28}$$

The underlying basis for the non-monotonic responses observed for mixed and partial inhibition can be probed by examining the changes in the product formation fluxes J_{R} and J_{W} and the effective substrate binding and unbinding rates as a function of inhibitor concentration. For one parameterization of the mixed inhibition model (Figure 1C) leading to undershoot (Figure 3A), the J_{R} and J_{W} fluxes become less sensitive to changes in inhibitor at different inhibitor concentration ranges, which causes the flux ratio to change non-monotonically as a function of inhibitor (Figure 3C). Ultimately, the effective substrate dissociation constant ($\hat{K}_D = \hat{k}_{\mathrm{off}}/\hat{k}_{\mathrm{on}}$) provides insights into how changes in inhibitor concentration can differentially affect the R and W product formation fluxes. As the inhibitor concentration increases, the effective dissociation constant for the R substrate saturates at a lower inhibitor concentration than the W substrate, which allows the product formation fluxes to change at different rates (Figure 3D).

In order to verify that these non-monotonic behaviors are general, we randomly sampled 10000 models and checked for overshoot and undershoot numerically. We enforce the inequality

$$f_{\mathrm{p}} < f_{\mathrm{off}}; \tag{29}$$

to ensure that the constraint $\eta_\infty < \eta_0$ is satisfied for the sampled parameters, Thus, as long as the inequality holds, the flux ratio at saturating inhibitor concentrations should be less than the uninhibited limit for mixed inhibitors. The situation for partial inhibition is complicated by the capacity for catalysis when bound to the inhibitor; therefore, when

1 sampling parameters for both models, we reject parameter sets that lead to $\eta_\infty > \eta_0$. Under

2 this sampling regime, we find that parameterizations yielding both undershoot (Figure 4A,B)

3 and overshoot (Figure 4C,D) responses can be readily found for both mixed and partial

4 inhibition modes. Interestingly, in practice, we also find that partial inhibition tends to be

5 less likely to exhibit these responses.

Figure 4: Mixed and partial inhibition can display both undershoot (A,B) and overshoot (C,D), where the flux ratio η can exceed the limiting bounds ($N = 10000$ parameter sets).

1 Suicide Inhibition Cannot in Itself Affect Specificity

In addition to reversible inhibitors, mechanism-based inhibitors (also known as "suicide" inhibitors) are capable of permanently inactivating the enzyme. Such inhibitors compete with *bona fide* substrates for access to the active site in a manner analogous to competitive inhibitors, with a basic reaction scheme given by

$$[E] + [I] \underset{k_{\text{off}}}{\overset{k_{\text{on}}}{\rightleftharpoons}} [EI] \xrightarrow{k_{\text{inact}}} [EI]^\star. \tag{30}$$

However, unlike real substrates, which can proceed through the full enzyme catalytic cycle, the suicide substrate becomes stuck at some covalent intermediate state, leading to enzyme inactivation ($[EI]^\star$).

At steady-state, for example, in cells, the quantity of active enzyme in the system remains constant over time; as enzyme is removed by inactivation, new enzyme is introduced into the system to maintain the steady state. In order to account for this, we modify the governing equation (Eq. 1) to include a production term $\sigma(x)$ that can balance the loss of enzyme by inactivation, i.e.

$$\frac{\mathrm{d}\vec{P}(x,t)}{\mathrm{d}t} = \mathbf{K}(x)\vec{P}(t) + \sigma(x). \tag{31}$$

With this modification, we find that enzyme inactivation by suicide substrates cannot affect substrate specificity at steady state (SI Figure 2). Additionally, time-course simulations of the suicide inhibition model rapidly approach the steady-state limit even in the case that the enzyme is not replaced after inactivation (SI Figure 3).

18 Ordered Binding Prevents TM from Affecting SIRT2 Specificity

The principles formulated for the basic inhibition networks can be applied to more complex schemes. Thus, we next applied our framework to a real system. The Sirtuin family of NAD-dependent de-acylases is particularly interesting, as they are involved in many biolog-

ical processes, including apoptosis and inflammation;[15] in addition, SIRT2 has been impli-

cated in some cancers.[16] Recently, a thiomyristoyl lysine compound (TM) was developed

as a mechanism-based inhibitor of the Sirtuin family member SIRT2;[17] like other suicide

inhibitors, TM proceeds down the SIRT2 catalytic pathway but stalls at an intermediate

step and remains covalently linked to the enzyme. Notably, TM has been shown to inhibit

SIRT2 histone 3 lysine 9 (H3K9) deacetylation while leaving SIRT2 activity towards other

substrates, such as myristoylated peptides, less affected.[6,17]

The precise mechanism by which TM affects SIRT2 substrate selectivity is unclear. It

has been proposed that the effect of TM on SIRT2 substrate specificity can be explained as a

function of relative substrate binding affinities;[6] if the inhibitor binds SIRT2 with a binding

affinity intermediate between some desired "correct" substrate and an undesired "incorrect"

substrate, it may be capable of preferentially inhibiting SIRT2 action against the incorrect

substrate.

Kinetic analyses by Jing et al. provided insights into the TM inhibition mechanism. We

employ these results to develop a model of SIRT2 inhibition. Competition assays under either

(1) NAD saturating conditions (1000 nM NAD) or (2) substrate saturating conditions (100

nM acetylated H3K9) suggest that TM is competitive with substrate and uncompetitive with

NAD.[17] Additionally, SIRT2 has been reported to follow an ordered binding mechanism;[18,19]

with ordered binding, the acyl lysine substrate must bind to SIRT2 to enable NAD binding.

With these constraints, we arrive at the model in Figure 5A. Lineweaver-Burke (LB) plots

generated with the ordered binding model still predict substrate-competitive inhibition by

TM (SI Figure 4A,B), consistent with the results of Jing et al.

Notably, the ordered binding model lacks reaction cycles involving the possible substrates.

Thus, we expect SIRT2 substrate selectivity to be insensitive to TM concentration. Indeed,

we find $\frac{d\eta(x)}{dx} = 0$ for all possible parameters, and that TM cannot affect SIRT2 specificity

while still satisfying the constraints imposed by experimental evidence (Figure 5B).

Figure 5: The NAD-dependent de-acylase SIRT2 is proposed to have an ordered binding mechanism, with acyl-lysine substrates binding prior to NAD (A, scheme adapted from [19]). If TM is competitive with the acyl-lysine substrate and uncompetitive with NAD, SIRT2 deacylation can be modeled with a simple kinetic scheme (B). Under these constraints, we predict that TM cannot affect SIRT2 substrate selectivity (C).

1 Substrate-Selective Inhibition of SIRT2 Requires Direct Substrate

2 Exchanges or Unordered Binding

In order for TM to affect SIRT2 substrate selectivity, we must introduce additional pathways

leading to product formation into the network in Figure 5A. There are at least two ways this

can be achieved without changing our assumptions on the mode of TM inhibition. First,

we can abandon the unordered binding constraint, and allow SIRT2 to bind NAD and acyl

lysine substrates in a random order (Figure 6A); as expected, relaxing the ordered binding

constraint allows TM to affect SIRT2 substrate specificity (Figure 6B). Additionally, LB plots

generated from the unordered binding model still largely agree with a substrate-competitive

inhibition mechanism (SI Figure 4C,D).

Alternatively, we can introduce direct substrate exchange reactions. In such reactions,

either substrate binding or inhibitor binding can displace the currently bound substrate,

which leads to the network in Figure 6C. As with relaxing the ordered binding constraint, introducing direct exchange reactions has the effect of creating cycles of reactions within the network; furthermore, with direct exchange reactions there is no need to relax the ordered binding constraint. Again, we see that allowing for direct exchanges enables TM to affect SIRT2 substrate specificity (Figure 5D).

Similar to the unordered binding model, the direct exchange model again agrees with substrate-competitive inhibition by TM (SI Figure 4E,F). However, at low values of $1/[R]$, the model predicts a slight curving in the LB plot; this curving was also observed in some cases for the unordered binding model. The curving may be an artifact of the precise parameters chosen for the model, which are largely unknown for SIRT2. In real experimental data, the curving in the LB plot could likely be missed due to measurement noise.

Discussion

Here, we have presented an analysis of substrate-selective inhibition for known inhibition modes. Our models suggest that competitive and uncompetitive inhibition cannot affect substrate selectivity (Figure 2). Previous experimental evidence corroborates this finding for competitive inhibition, where enzyme active site inhibitors have generally failed to affect substrate selectivity.[5]

In addition, we have found that the presence of reaction cycles that involve both inhibitor and substrate binding is generally sufficient to observe substrate-selective inhibition; in the basic enzyme inhibition modes we have examined, such cycles are found only in mixed and non-competitive inhibition, as well as in partial inhibition. Cycles that involve only the inhibitor appear to be incapable of affecting substrate selectivity. For example, in a modified version of the SIRT2 ordered binding model where SIRT2 binds to NAD prior to the substrate (a mechanism similar to the related SIRT6[20]), the only reaction cycle involves NAD. Under this alternative reaction mechanism, TM cannot affect specificity (SI Figure

Figure 6: If the TM inhibitor is competitive with substrate and uncompetitive with NAD, either (A) unordered binding or (B) direct substrate exchange reactions are required for TM to affect substrate specificity (C,D).

5).

Prior kinetic models of substrate-selective inhibitors are consistent with our findings. The model of IDE inhibition proposed by Maianti et al. is consistent with our findings. They propose a scheme that is essentially a limiting case of the partial inhibition scheme (SI Figure 6A), whereby the inhibited enzyme is still capable of both binding and cleaving glucagon, while significantly reducing insulin binding and cleavage.[5] While Maianti et al. do not consider the possibility of inhibitor binding to the enzyme-substrate complex, it is more than likely possible, as there are no intrinsic physical barriers preventing these binding reactions. However, even in the absence of reaction cycles substrate-selective inhibition can still be maintained so long as there are multiple catalytically competent states (SI Figure

6). With multiple catalytic states for both substrates, the overall product formation flux ratio is a ratio of sums of fluxes and changes as a function of the inhibitor concentration as inhibitor-bound states become more prevalent.

In our models, we have considered the situation in which the inhibitor can inhibit catalysis for multiple substrates, albeit to possibly different extents. However, in the extreme case in which an inhibitor is competitive with only one substrate out of many, even competitive inhibition (which lacks reaction cycles) can affect substrate selectivity; this is the mechanism that governs the IDE inhibitor identified by Maianti et al., which sterically hinders insulin binding to the active site while leaving glucagon binding unaffected.[5] The kinetic scheme describing IDE inhibition can be considered as a form of partial inhibition, in which catalysis with one substrate (insulin) is completely inhibited while the other (glucagon) is completely unaffected by the presence of inhibitor. Notably, the scheme considered by Maianti et al. lacks reaction cycles,[5] but admits multiple pathways to the glucagon product formation reaction. Overall, this example supports the idea that the existence of multiple pathways to formation is generally a sufficient condition to observe substrate-selective inhibition.

In addition, Robin et al. recently demonstrated that inhibitor binding can paradoxically increase enzyme turnover rates (catalytic events per unit time) at intermediate concentrations, as measured at the single-enzyme level.[21] In their framework, they abandon the typical assumption that system state transitions are Markovian and are exponentially distributed, and instead allow the distribution of transition waiting times to follow an arbitrary distribution. Future work could extend our models in an analogous manner to test whether these effects at the single-enzyme level can also affect substrate selectivity.

Finally, our modeling framework could be applied to complicated biological decision-making processes, such as the DNA repair pathway decision.[22,23] In the context of CRISPR-Cas9-mediated genome editing, the decision between classical non-homologous end joining (cNHEJ) and homology-directed repair (HDR) has a significant impact on the editing outcome. Consequently, there has been substantial work aimed at understanding the repair of

Cas9-mediated DNA breaks.[24–28] Additionally, there have been attempts to bias DNA repair to favor a particular desired editing outcome; this can involve precise timing of Cas9 delivery in conjunction with cell-cycle synchronization[29] as well as co-expression of exogenous factors such as yeast Rad52.[30] While these techniques can work well when editing cells in laboratory settings, they may prove difficult to translate to clinical applications of Cas9.

Conclusions

Overall, we have shown that mixed and partial inhibition are the only reversible inhibition mechanisms capable of affecting substrate selectivity. The physical basis of this selectivity can be explained by analogy to the work of Mallory et al.,[8] which demonstrated that perturbations in the free-energy barriers between states can affect biochemical pathway flux ratios. In mixed and partial inhibition schemes, as well as in systems possessing reaction cycles, the presence of an inhibitor (or other biochemical cofactors) acts as a *de facto* perturbation, affecting the transition barriers. Changes in substrate selectivity, which are nothing more than changes in ratios of biochemical pathway fluxes, are a direct consequence of these perturbations.

We have also demonstrated that the inhibitor TM, targeting the enzyme SIRT2, can affect substrate specificity while satisfying known biochemical constraints. The SIRT2 ordered binding mechanism in which substrate binds prior to the required NAD cofactor,[19] coupled with competitive inhibition of substrate binding, has the effect of eliminating reaction cycles that involve the acylated substrates; as a consequence, the inhibitor is no longer capable of affecting the transition barriers, and thus either a relaxation of constraints or direct substrate exchange reactions is required to restore substrate-selective inhibition.

Acknowledgement

The authors would like to thank Anatoly B. Kolomeisky, Seokjoo Chae, and Zheng Diao for insightful comments on the manuscript.

O.A.I. acknowledges funding support from the Welch Foundation (Grant C-1995).

Supporting Information Available

Model implementation details; Transient suicide inhibition model; Supplemental figures and tables

References

(1) Garavito, R. M.; Mulichak, A. M. The Structure of Mammalian Cyclooxygenases. *Annual Review of Biophysics and Biomolecular Structure* **2003**, *32*, 183–206, DOI: `10.1146/annurev.biophys.32.110601.141906`.

(2) Rosti, G.; Castagnetti, F.; Gugliotta, G.; Baccarani, M. Tyrosine Kinase Inhibitors in Chronic Myeloid Leukaemia: Which, When, for Whom? *Nature Reviews. Clinical Oncology* **2017**, *14*, 141–154, DOI: `10.1038/nrclinonc.2016.139`.

(3) Zhang, Y.; Shen, L.; Peng, Z. Advances in MET Tyrosine Kinase Inhibitors in Gastric Cancer. *Cancer Biology & Medicine* **2024**, *21*, 484–498, DOI: `10.20892/j.issn.2095-3941.2024.0044`.

(4) Carey, N.; La Thangue, N. B. Histone Deacetylase Inhibitors: Gathering Pace. *Current Opinion in Pharmacology* **2006**, *6*, 369–375, DOI: `10.1016/j.coph.2006.03.010`.

(5) Maianti, J. P.; Tan, G. A.; Vetere, A.; Welsh, A. J.; Wagner, B. K.; Seeliger, M. A.; Liu, D. R. Substrate-Selective Inhibitors That Reprogram the Activ-

ity of Insulin-Degrading Enzyme. *Nature Chemical Biology* **2019**, *15*, 565–574, DOI: `10.1038/s41589-019-0271-0`.

(6) Lin, H. Substrate-Selective Small-Molecule Modulators of Enzymes: Mechanisms and Opportunities. *Current Opinion in Chemical Biology* **2023**, *72*, 102231, DOI: `10.1016/j.cbpa.2022.102231`.

(7) Hendriks, B. S.; Seidl, K. M.; Chabot, J. R. Two Additive Mechanisms Impair the Differentiation of 'substrate-Selective' P38 Inhibitors from Classical P38 Inhibitors in Vitro. *BMC Systems Biology* **2010**, *4*, DOI: `10.1186/1752-0509-4-23`.

(8) Mallory, J. D.; Kolomeisky, A. B.; Igoshin, O. A. Kinetic Control of Stationary Flux Ratios for a Wide Range of Biochemical Processes. *Proceedings of the National Academy of Sciences of the United States of America* **2020**, *117*, 8884–8889, DOI: `10.1073/pnas.1920873117`.

(9) Lehninger, A. L.; Nelson, D. L.; Cox, M. M. *Lehninger Principles of Biochemistry*, 5th ed.; W.H. Freeman: New York, 2008.

(10) Masson, P.; Mukhametgalieva, A. R. Partial Reversible Inhibition of Enzymes and Its Metabolic and Pharmaco-Toxicological Implications. *International Journal of Molecular Sciences* **2023**, *24*, 12973, DOI: `10.3390/ijms241612973`.

(11) Mallory, J. D.; Kolomeisky, A. B.; Igoshin, O. A. Trade-Offs between Error, Speed, Noise, and Energy Dissipation in Biological Processes with Proofreading. *The Journal of Physical Chemistry. B* **2019**, *123*, 4718–4725, DOI: `10.1021/acs.jpcb.9b03757`.

(12) Yu, Q.; Mallory, J. D.; Kolomeisky, A. B.; Ling, J.; Igoshin, O. A. Trade-Offs between Speed, Accuracy, and Dissipation in tRNAIle Aminoacylation. *The Journal of Physical Chemistry Letters* **2020**, *11*, 4001–4007, DOI: `10.1021/acs.jpclett.0c01073`.

(13) Midha, T.; Mallory, J. D.; Kolomeisky, A. B.; Igoshin, O. A. Synergy among Pausing, Intrinsic Proofreading, and Accessory Proteins Results in Optimal Transcription Speed and Tolerable Accuracy. *The Journal of Physical Chemistry Letters* **2023**, *14*, 3422–3429, DOI: `10.1021/acs.jpclett.3c00345`.

(14) Banerjee, K.; Kolomeisky, A. B.; Igoshin, O. A. Elucidating Interplay of Speed and Accuracy in Biological Error Correction. *Proceedings of the National Academy of Sciences of the United States of America* **2017**, *114*, 5183–5188, DOI: `10.1073/pnas.1614838114`.

(15) Lu, W.; Ji, H.; Wu, D. SIRT2 Plays Complex Roles in Neuroinflammation Neuroimmunology-Associated Disorders. *Frontiers in Immunology* **2023**, *14*, 1174180, DOI: `10.3389/fimmu.2023.1174180`.

(16) Chen, G.; Huang, P.; Hu, C. The Role of SIRT2 in Cancer: A Novel Therapeutic Target. *International Journal of Cancer* **2020**, *147*, 3297–3304, DOI: `10.1002/ijc.33118`.

(17) Jing, H.; Hu, J.; He, B.; Negrón Abril, Y. L.; Stupinski, J.; Weiser, K.; Carbonaro, M.; Chiang, Y.-L.; Southard, T.; Giannakakou, P. et al. A SIRT2-Selective Inhibitor Promotes c-Myc Oncoprotein Degradation and Exhibits Broad Anticancer Activity. *Cancer Cell* **2016**, *29*, 297–310, DOI: `10.1016/j.ccell.2016.02.007`.

(18) Feldman, J. L.; Dittenhafer-Reed, K. E.; Kudo, N.; Thelen, J. N.; Ito, A.; Yoshida, M.; Denu, J. M. Kinetic and Structural Basis for Acyl-Group Selectivity and NAD(+) Dependence in Sirtuin-Catalyzed Deacylation. *Biochemistry* **2015**, *54*, 3037–3050, DOI: `10.1021/acs.biochem.5b00150`.

(19) Bernhard, S. P.; Ruiz, F. X.; Remiszewski, S.; Todd, M. J.; Shenk, T.; Kulp, J. L.; Chiang, L. W. Structural Basis for Sirtuin 2 Activity and Modulation: Current State and Opportunities. *The Journal of Biological Chemistry* **2025**, *301*, 110274, DOI: `10.1016/j.jbc.2025.110274`.

(20) Klein, M. A.; Liu, C.; Kuznetsov, V. I.; Feltenberger, J. B.; Tang, W.; Denu, J. M. Mechanism of Activation for the Sirtuin 6 Protein Deacylase. *The Journal of Biological Chemistry* **2020**, *295*, 1385–1399, DOI: `10.1074/jbc.RA119.011285`.

(21) Robin, T.; Reuveni, S.; Urbakh, M. Single-Molecule Theory of Enzymatic Inhibition. *Nature Communications* **2018**, *9*, 779, DOI: `10.1038/s41467-018-02995-6`.

(22) Ingram, S. P.; Warmenhoven, J. W.; Henthorn, N. T.; Smith, E. a. K.; Chadwick, A. L.; Burnet, N. G.; Mackay, R. I.; Kirkby, N. F.; Kirkby, K. J.; Merchant, M. J. Mechanistic Modelling Supports Entwined Rather than Exclusively Competitive DNA Double-Strand Break Repair Pathway. *Scientific Reports* **2019**, *9*, 6359, DOI: `10.1038/s41598-019-42901-8`.

(23) Paull, T. T. Reconsidering Pathway Choice: A Sequential Model of Mammalian DNA Double-Strand Break Pathway Decisions. *Current Opinion in Genetics & Development* **2021**, *71*, 55–62, DOI: `10.1016/j.gde.2021.06.011`.

(24) Roidos, P.; Sungalee, S.; Benfatto, S.; Serçin, Ö.; Stütz, A. M.; Abdollahi, A.; Mauer, J.; Zenke, F. T.; Korbel, J. O.; Mardin, B. R. A Scalable CRISPR/Cas9-based Fluorescent Reporter Assay to Study DNA Double-Strand Break Repair Choice. *Nature Communications* **2020**, *11*, 4077, DOI: `10.1038/s41467-020-17962-3`.

(25) Eki, R.; She, J.; Parlak, M.; Benamar, M.; Du, K.-P.; Kumar, P.; Abbas, T. A Robust CRISPR-Cas9-based Fluorescent Reporter Assay for the Detection and Quantification of DNA Double-Strand Break Repair. *Nucleic Acids Research* **2020**, *48*, e126, DOI: `10.1093/nar/gkaa897`.

(26) Song, B.; Yang, S.; Hwang, G.-H.; Yu, J.; Bae, S. Analysis of NHEJ-Based DNA Repair after CRISPR-Mediated DNA Cleavage. *International Journal of Molecular Sciences* **2021**, *22*, 6397, DOI: `10.3390/ijms22126397`.

(27) Miyaoka, Y.; Berman, J. R.; Cooper, S. B.; Mayerl, S. J.; Chan, A. H.; Zhang, B.; Karlin-Neumann, G. A.; Conklin, B. R. Systematic Quantification of HDR and NHEJ Reveals Effects of Locus, Nuclease, and Cell Type on Genome-Editing. *Scientific Reports* **2016**, *6*, 23549, DOI: `10.1038/srep23549`.

(28) Brinkman, E. K.; Chen, T.; de Haas, M.; Holland, H. A.; Akhtar, W.; van Steensel, B. Kinetics and Fidelity of the Repair of Cas9-Induced Double-Strand DNA Breaks. *Molecular Cell* **2018**, *70*, 801–813.e6, DOI: `10.1016/j.molcel.2018.04.016`.

(29) Lin, Y.; Cradick, T. J.; Brown, M. T.; Deshmukh, H.; Ranjan, P.; Sarode, N.; Wile, B. M.; Vertino, P. M.; Stewart, F. J.; Bao, G. CRISPR/Cas9 Systems Have off-Target Activity with Insertions or Deletions between Target DNA and Guide RNA Sequences. *Nucleic Acids Res* **2014**, *42*, 7473–85, DOI: `10.1093/nar/gku402`.

(30) Shao, S.; Ren, C.; Liu, Z.; Bai, Y.; Chen, Z.; Wei, Z.; Wang, X.; Zhang, Z.; Xu, K. Enhancing CRISPR/Cas9-mediated Homology-Directed Repair in Mammalian Cells by Expressing Saccharomyces Cerevisiae Rad52. *The International Journal of Biochemistry & Cell Biology* **2017**, *92*, 43–52, DOI: `10.1016/j.biocel.2017.09.012`.