Tree microbiomes and methane emissions in upland forests

Jonathan Gewirtzman[1†*], Wyatt Arnold[2†], Meghan Taylor[1,3], Hannah Burrows[5], Carter Merenstein[4], David Woodbury[1], Naomi Whitlock[6], Kendall Kraut[6], Leslie Gonzalez[7], Craig R. Brodersen[1,8], Marlyse Duguid[1,8], Peter A. Raymond[1,9], Jordan Peccia[2], Mark A. Bradford[1,8]

[1] Yale School of the Environment, Yale University, New Haven, CT, USA
[2] Department of Chemical and Environmental Engineering, School of Engineering and Applied Science, Yale University, New Haven, CT, USA
[3] WCRP Climate and the Cryosphere, University of Massachusetts, Amherst, Amherst, MA, USA
[4] Harvard College, Harvard University, Boston, MA, USA
[5] Harvard T.H. Chan School of Public Health, Boston, MA, USA
[6] Wesleyan University, Middletown, CT, USA
[7] Yale College, Yale University, New Haven, CT, 06511
[8] The Forest School, Yale School of the Environment, Yale University, New Haven, CT, USA
[9] Yale Center for Natural Carbon Capture, Yale University, New Haven, CT, USA
[†] Contributed equally to this work
* Corresponding author: jonathan.gewirtzman@yale.edu

Summary

- **Rationale**: Trees emit methane and harbor methanogens, but the generality, distribution, and ecosystem significance of these associations remain unclear. We investigated methanogenic colonization across 16 upland forest tree species and evaluated relationships between microbial communities and emissions.
- **Methods**: We measured 1,148 stem fluxes and 276 soil fluxes, quantified methanogens and methanotrophs via droplet digital PCR in 564 samples, characterized communities through 16S rRNA sequencing, and developed models to upscale fluxes ecosystem-wide.
- **Key results**: Methanogens were detected in 97% of heartwood samples (up to 10^7 copies g^{-1}), exceeding mean soil abundances by ~2 orders of magnitude. Wood harbored distinct methanogenic communities (Methanobacteriaceae, Methanomassiliicoccaceae) compared to soils. Species-level methanogen abundance correlated with emissions ($R^2 = 0.40$), though individual-level correlations were weak, reflecting spatial heterogeneity and complex metabolism and transport interactions.
- **Main conclusion**: Methanogens are common in tree microbiomes across diverse species, concentrated in heartwood with substantial individual variation. Species-level abundance patterns partially explain emission variability, establishing internal production as a widespread contributor to upland tree fluxes. Future multi-omics approaches could strengthen abundance-flux relationships, while resolving ecosystem-scale importance requires improved quantification of canopy surface area and flux variability.

Keywords: forest carbon cycle; greenhouse gas; heartwood; methanogen; methanotroph; methane emissions; temperate forest; tree microbiome

Introduction

Forests are key regulators of the global methane (CH_4) cycle, functioning as both sinks and
sources of this potent greenhouse gas. Methane contributes significantly to climate change,
possessing a global warming potential more than 80 times greater than carbon dioxide (CO_2)
over a 20-year period (Wood et al. 2023; Forster et al. 2021). While the role of soil in forest
methane cycling has been extensively studied, emerging evidence reveals that trees themselves
play an active role in methane fluxes (Barba et al., 2019; Gauci, 2025). Tree stems have been
identified as both methane emitters and sinks, suggesting that they may also be important
regulators of methane dynamics in forest ecosystems. However, the biological and ecological
mechanisms underlying these tree-mediated methane fluxes remain poorly understood.

Tree methane emissions may originate from methanogens (methane-producing archaea) and
methanotrophs (methane-oxidizing bacteria) within tree tissues. Methanogens, which generate
methane from H_2/CO_2 or acetate in anaerobic environments (Le Mer & Roger 2001), have been
detected in heartwood of multiple species (Zeikus & Ward 1974; Zeikus & Henning 1975;
Covey et al. 2012; Yip et al. 2019; Flanagan et al. 2021; Feng et al. 2022; Moisan et al. 2025;
Harada et al. 2024; Mochidome et al. 2025). Anaerobic conditions in heartwood (Schink & Ward
1984; Shigo & Hillis 1973; Jensen 1967; Arnold et al. 2025), where oxygen diffusion is
restricted, may create suitable methanogenic niches (Uroz et al. 2016; Mieszkin et al. 2021), but
the extent of colonization, the metabolic pathways, and the overall contribution to forest methane
budgets remain unclear (Putkinen et al. 2021; Barba et al. 2024; Barba, Bradford, et al. 2019).

Methanotrophs oxidize CH_4 to CO_2 and water under aerobic conditions. Detected in sapwood
and bark, they may mitigate emissions before atmospheric release (Leung et al. 2024; Jeffrey et
al. 2021; Carmichael et al. 2024; Gauci et al. 2024). The co-occurrence of methane producers
and consumers suggests complex interactions influencing net fluxes at tree and ecosystem scales,
but quantitative data on their abundance, activity, and spatial distribution remain sparse
(Putkinen et al. 2021). It remains unclear whether methanotroph activity varies with tree species,
physiological traits, or environmental conditions.

Tree-mediated methane emissions vary considerably across species, landscapes, and temporal
scales. In wetland forests, trees primarily act as conduits for soil-derived methane, transporting
CH_4 from anaerobic soils through vascular pathways (Pangala et al. 2013; Pangala et al. 2015;
Pangala et al. 2017; Pitz et al. 2018; Jeffrey et al. 2023; Jeffrey et al. 2019). Conversely, upland
forests, where well-drained soils are traditionally methane sinks, have trees that emit methane
even when adjacent soils consume it (Machacova et al. 2023; Bréchet et al. 2025; Hettwer et al.
2025; Pitz & Megonigal 2017; Warner et al. 2017; Covey & Megonigal 2019), suggesting
internal production rather than soil transport (Pitz & Megonigal 2017). While a number of recent
detections of endophytic methanogens support this hypothesis, factors governing production
including tree species, wood chemistry, and microbial interactions remain poorly understood.

The taxonomic, spatial, and environmental distribution of microbial methane producers and
consumers within trees remains unresolved. Methanogens are thought to primarily associate with

heartwood, but their presence in sapwood or bark lacks systematic investigation (Arnold et al.
2024; Arnold et al. 2025). Methanotroph activity in sapwood and bark likely varies with oxygen
availability and stem water content (Putkinen et al. 2021). Environmental factors including soil
moisture, nutrients, and rhizosphere composition influence methane dynamics, though their
interaction with tree-mediated fluxes is unclear (Epron et al. 2022). Climate and tree-specific
traits such as wood density and anatomical features may regulate microbial colonization and flux
potential (Barba, Bradford, et al. 2019).

The ecological and biogeochemical significance of tree-mediated methane emissions extends
beyond individual trees to entire forest ecosystems and the planet. Given that forests represent a
substantial component of the global methane budget, even small changes in tree methane
emissions could have significant implications for atmospheric CH_4 concentrations. The
recognition that trees in upland forests may contribute to net methane fluxes challenges
traditional assumptions about forest methane cycling and highlights the need for a more
integrative approach to forest methane budgets (Gauci 2025).

Study Objectives

Despite growing evidence of tree methane emissions, the underlying microbial and physiological
mechanisms remain poorly characterized, particularly in upland forests where soils consume
atmospheric methane. We investigated methane fluxes and associated microbial communities
across a moisture gradient from upland to transitional wetland sites at a mixed temperate forest
in Connecticut, USA, addressing four objectives:

(1) Quantify tree methane emissions across environmental gradients. We quantified methane
fluxes from stems and soils across upland, intermediate, and transitional wetland sites,
examining vertical profiles to differentiate soil-derived transport from internal production. We
hypothesized upland trees would show uniform vertical profiles consistent with internal
production, contrasting with exponential height-dependent decline in wetland trees. We assessed
species-specific differences and performed variance partitioning to identify emission drivers.

(2) Characterize methanogenic communities within trees. We determined methanogen
abundance and distribution across heartwood and sapwood using droplet digital PCR and
characterized taxonomic composition via 16S rRNA sequencing. We tested whether tree
methanogens represent distinct communities from soil populations, hypothesizing heartwood
would harbor higher abundances due to lower oxygen. Through co-occurrence analysis and
functional inference, we characterized microbial consortia and metabolic pathways, predicting
fermentative bacteria would co-occur with methanogens to provide substrates.

(3) Establish methanogen abundance-flux relationships. We examined whether methanogen
populations correlate with emission rates at both individual tree and species levels. We tested
correlations at multiple scales to determine whether aggregation across individuals would reduce
noise from spatial heterogeneity and reveal underlying biological relationships between
methanogen abundance and methane flux.

(4) Evaluate ecosystem-scale impacts. Using random forest models incorporating tree
characteristics and environmental drivers, we upscaled measurements to estimate forest-wide
fluxes. We calculated how tree emissions offset soil consumption to determine whether tree
fluxes meaningfully alter upland forest methane sink status.

Here we present 1,148 tree stem flux measurements from 482 individual trees across 16 species;
276 soil flux measurements; and molecular quantification of methanogens and methanotrophs
from 564 tree and soil samples collected across upland to transitional wetland gradients in a
temperate mixed forest. Through integrated flux measurements, molecular quantification, and
community analysis, we aimed to establish whether methanogens are endemic tree microbiome
components and determine their contribution to emissions across species and environmental
gradients. This multi-scale approach addresses fundamental questions about the origin,
regulation, and magnitude of tree-mediated methane emissions, advancing mechanistic
understanding of this emerging component of the global methane cycle.

Materials and Methods

Study site

Field measurements were conducted at Yale-Myers Forest, a 3,213-ha research forest in
northeastern Connecticut, USA (41°56′N, 72°07′W). The forest comprises second-growth
mixed-deciduous vegetation established on abandoned agricultural land since the mid-19th
century, with elevations ranging from 170-300 m a.s.l. The climate is temperate and humid with
mean annual temperature of 9.5°C (summer: 21°C, winter: -2°C) and 123 cm annual
precipitation. Soils consist of glacial till-derived, moderately to well-drained stony loams
overlaying bedrock.

Measurements during 2020-2021 and 2023 were conducted within an 8-ha permanent forest
inventory plot established following standardized protocols, with some additional 2020-2021
measurements in transitional wetland immediately adjacent to the plot. Destructive sampling for
microbiome analysis (2021-2022) occurred in nearby forest stands of similar composition and
developmental history, as destructive sampling was prohibited within the long-term monitoring
plot.

Tree stem methane flux measurements

Chamber designs

Two chamber types were employed for stem flux measurements. Semi-rigid chambers (2020-
2021) were constructed following (Siegenthaler et al. 2016) using transparent polyethylene
terephthalate (PET) sheets . Sheets were framed with 1.5 cm thick × 3 cm wide adhesive-backed
expanded neoprene strips for gas-tight sealing, with two vertical neoprene wedges maintaining
equidistant spacing from the stem. Chamber volume was calculated as: $Vc = (HL/(Dstem + 2T))$
$× [(Dstem + 2T)/2]^2 - [Dstem/2]^2 - Vwedges$, where H = height, L = periphery length, Dstem =
stem diameter, and T = chamber thickness. Chambers were wrapped around stems, secured with
ratchet straps, and gaps sealed with modeling clay tested to confirm no CH_4 or CO_2 off-gassing.

Rigid chambers (2021-2023) were constructed from transparent plastic containers (Rubbermaid)
with arcs cut to fit tree stems. Chamber volumes (0.5-2.0 L) were determined by gas standard
dilution in the laboratory (Jeffrey et al., 12/2020; Siegenthaler et al., 2016). Enclosed surface
area was calculated as the product of the chamber's planar area and arc length. Chambers were
sealed with potting clay (Amaco) and secured with lashing straps. Seal integrity was verified

through visual inspection, blowing on perimeter to check for spikes in CO_2, and confirmed by
monitoring concentration linearity.

Measurement protocol

All chambers were connected via 5-mm internal diameter PVC tubing (Bev-a-Line) to a portable
off-axis integrated cavity output spectroscopy analyzer (GLA131-GGA, Los Gatos Research)
measuring CO_2, CH_4, and H_2O concentrations at 1 Hz with CH_4 precision <0.9 ppb. The
analyzer's 5 L min^{-1} flow rate allowed complete chamber volume turnover within ~30 s.
Measurements lasted 3-10 min per location, with the initial 30 s excluded for equilibration.

During the 2020-2021 temporal survey, measurements were collected from 41 trees (21 upland,
10 intermediate, 10 wetland) and 30 soil collars distributed across 6 plots (2 plots per landscape
position) during 10 sampling campaigns from June 2020 to May 2021. Soil collars were
distributed as follows: upland plots with 12 collars (2 plots × 6 collars), intermediate plots with 6
collars (2 plots × 3 collars), and wetland plots with 12 collar positions (2 plots × 6 collars,
measured under both wetland-dry and wetland-saturated conditions). The 2021 intensive survey
measured fluxes at three heights (50, 125, 200 cm) on the southern aspect of 158 trees across 16
species: *Acer rubrum*, *A. saccharum*, *Betula alleghaniensis*, *B. lenta*, *B. papyrifera*, *Carya ovata*,
Fagus grandifolia, *Fraxinus americana*, *Kalmia latifolia*, *Pinus strobus*, *Prunus serotina*,
Quercus alba, *Q. rubra*, *Q. velutina*, *Sassafras albidum*, and *Tsuga canadensis*. The 2022
vertical profile study measured a single mature *Q. velutina* at seven heights (0.5, 1.25, 2, 4, 6, 8,
10 m), with heights >2 m accessed via arborist climbing equipment. The 2023 survey expanded
to 335 trees with measurements standardized at breast height (125 cm).

Soil methane flux and associated measurements

Soil CH_4 fluxes (2020–2021) were measured using static chambers consisting of PVC collars (25
cm diameter) inserted 5 cm into soil at least 24 h before measurement to minimize disturbance.
A PVC cap chamber was sealed to collars during measurements using the same analyzer system.
Soil moisture was characterized in December 2020 through transects perpendicular to moisture
gradients, with volumetric water content measured at 12 cm depth using a handheld sensor
(HydroSense, Campbell Scientific). Measurements along transects were collected at a spacing of
~30 m between points, ensuring coverage across both fine-scale and broader soil moisture
variability. Soil temperature was recorded at 10 cm depth during flux measurements.

Flux calculations

Methane and CO_2 fluxes were calculated using the goFlux R package v2.0.0 (Rheault et al.
2024), which implements both linear (LM) and non-linear Hutchinson-Mosier (HM) models with
automatic water vapor dilution correction (Hutchinson & Mosier 1981; Hüppi et al. 2018). The
flux equation:

$F = (dC/dt) \times (V_c/A_c) \times (P/RT) \times (1 - X_{H_2O})$

where F = flux (nmol m^{-2} s^{-1} for CH_4; μmol m^{-2} s^{-1} for CO_2), dC/dt = concentration change rate,
Vc = chamber volume (L), Ac = surface area (m^2), P = atmospheric pressure (101.325 kPa), R =
8.314 L kPa K^{-1} mol^{-1}, T = temperature (K), and X_{H_2O} = water vapor mole fraction.

Model selection used the best.flux() function with criteria: goodness-of-fit metrics (MAE, RMSE
< instrument precision; AICc), physical constraints (g-factor < 2.0 where g = HM flux/LM flux;
κ/κmax ≤ 1.0), and quality thresholds (P < 0.05; flux > minimal detectable flux; n ≥ 60
observations). CO_2 flux R^2 served as a chamber seal quality metric, as tree respiration ensures
consistently positive CO_2 fluxes.

Internal gas sampling

Tree internal gas concentrations were measured by drilling holes with 5-mm increment borers
from bark to pith, then sealing with gas-impermeable tape (2021-2022) or butyl rubber stoppers
(2024). After 5 min equilibration, 20 ml gas was extracted using gas-tight syringes (Hamilton)
and stored overpressurized in pre-evacuated 12 ml Exetainer vials (Labco). Analysis at Yale
Analytical and Stable Isotope Center employed gas chromatography with flame ionization
detection for CH_4 and CO_2, and electron capture detection for N_2O and O_2. Calibration used
certified standards (0-60,000 ppm CH_4) with ambient air blanks between runs.

Microbiome sampling and processing

Tree cores were collected at 125 cm height using 5.15-mm increment borers (Haglöf Sweden)
flame-sterilized between trees. Cores were immediately wrapped in sterile aluminum foil and
frozen on dry ice, then stored frozen at -80°C until further analysis. Processing followed methods
we optimized and reported (Arnold et al. 2024), and involved: (i) sectioning into operationally-
defined heartwood (inner 5 cm from pith) and sapwood (outer 5 cm from bark); (ii)
lyophilization for 72 h; (iii) cryogenic grinding (Spex 6775 Freezer/Mill) with 10 min liquid
nitrogen pre-cooling followed by two cycles of 2 min grinding and 2 min cooling at 10 cycles
s^{-1}, with materials flame-sterilized or bleached between samples.

Soil samples were collected at four cardinal directions around each tree at distances equal to tree
circumference or 1 m (whichever greater). Samples were separated by horizon (organic; mineral
to 30 cm depth or refusal), composited by horizon, sieved to 4 mm in the field, and immediately
frozen on dry ice then stored at -80°C until further analysis.

Molecular analyses

DNA was extracted from 100 mg ground wood or 250 mg soil using MagMAX Microbiome
Ultra Nucleic Acid Isolation Kit with KingFisher Apex automated extraction, eluting into 75 μl
(wood) or 100 μl (soil). Samples with potential PCR inhibitors were further processed using
Zymo OneStep-96 PCR Inhibitor Removal Kit. Microbial communities were characterized
through 16S rRNA amplicon sequencing at University of Minnesota Genomics Center following
protocols in Arnold and Gewirtzman et al. (2025). The 16S rRNA V4 region was amplified using
515F/806R primers with PNA blockers reducing plant DNA amplification. Sequencing

employed Illumina MiSeq with 2×300 bp paired-end chemistry and dual indexing. Sequencing
results from this analysis were also reported in (Arnold et al. 2025).

Methanogen (mcrA gene) and methanotroph (pmoA and mmoX genes) abundances were
quantified using droplet digital PCR (Bio-Rad QX200), also using methods reported in (Arnold
et al. 2024). For mcrA quantification (Steinberg & Regan 2009; Kolb et al. 2003), we used our
novel probe-based assay employing FAM-based degenerate primers and probe set (Forward:
ACGACYTRCAGGAYCAGTGY, Probe: WGGWCCWAACTAYCCBAACTACG, Reverse:
TGGTGWCCBACGTTCATYG) which produces a 118 bp amplicon. This was found to reduce
the rate of false positives found in previous methods, and is recommended for environments that
may harbor low abundance methanogen populations. Reaction conditions followed the published
method: 95°C for 10 min, followed by 40 cycles of 94°C for 30 s and 48°C for 1 min 20 s, then
98°C for 10 min. Reactions used ddPCR Supermix for Probes (No dUTP) (Bio-Rad) with primer
concentrations of 900 nM and probe concentration of 250 nM. For pmoA and mmoX genes,
EvaGreen chemistry was employed with previously published primers (Luesken et al. 2011;
McDonald & Murrell 1997; McDonald et al. 1995). All assays included standards and negative
controls. Primers were synthesized at the Keck Oligonucleotide Synthesis facility at Yale
University. Probes were synthesized by IDT using their PrimeTime chemistry.

Bioinformatic and statistical analyses

Sequence data were processed using DADA2 (Callahan et al. 2016) via Nephele v2.23.2 (Weber
et al. 2018), with taxonomic assignment using SILVA v138.1 (16S) database (Quast et al. 2013).
After filtering sequences with <10 reads and removing chloroplast/mitochondrial ASVs, samples
were rarefied to 3,500 reads. Community analyses employed phyloseq and microeco R packages
using weighted/unweighted UniFrac distances (McMurdie & Holmes 2013; Liu et al. 2021).
Functional inference used PICRUSt2 (Douglas et al. 2020) and FAPROTAX (Louca et al. 2016).

Variance partitioning employed linear mixed-effects models in lme4 (Bates et al. 2015) with tree
identity as random effect and species, diameter, and moisture as fixed effects. Random Forest
models using ranger (Wright & Ziegler 2017) were developed to upscale fluxes using
environmental predictors including temperature, moisture, seasonal indices, and taxonomic
information, with separate models for stem and soil fluxes.

Results

1. Tree and Soil Methane Fluxes from Upland to Transitional Wetland Sites

Methane flux measurements from both tree stems and soil revealed distinct patterns across
upland, intermediate, and transitional wetland environments at Yale-Myers Forest during the
2020-2021 survey period. In upland sites, tree stems consistently showed positive CH_4 fluxes
(mean 3.3 µg CH_4 m^{-2} hr^{-1}, range -10.1 to 267), with 69% of measurements indicating methane
emissions. This contrasted sharply with upland soils, which acted as strong methane sinks (mean

-90.9 µg CH₄ m⁻² hr⁻¹, range -247.8 to -25.4), with 100% of measurements showing methane

consumption. At intermediate sites, tree fluxes remained predominantly positive but lower in

magnitude (mean 0.6 µg CH₄ m⁻² hr⁻¹, range -1.2 to 14.7), with 62% of measurements showing

emissions. Soils at these sites maintained strong methane consumption (mean -77.7 µg CH₄ m⁻²

hr⁻¹, range -263.4 to -13.4), comparable to upland sites despite higher soil moisture.

In transitional wetland sites, trees showed the highest emission rates (mean 7.5 µg CH₄ m⁻² hr⁻¹,

range -0.8 to 183.8), with 78% of measurements positive. Soils in these areas exhibited extreme

variability, with a mean flux of 105.9 µg CH₄ m⁻² hr⁻¹ but ranging from -127.2 to 3,642.6 µg CH₄

m⁻² hr⁻¹. The median soil flux in transitional wetlands was -6.0 µg CH₄ m⁻² hr⁻¹, indicating that

despite occasional high emission events, these soils frequently consumed methane (60% of

measurements were negative).

These results highlight a clear distinction between tree-mediated methane emissions and soil

methane fluxes across different landscape positions. While upland soils act as strong methane

sinks, upland tree stems are consistent methane sources, with a one-sample t-test confirming that

upland tree fluxes were significantly greater than zero (p = 0.017). This indicates that methane

produced within tree tissues contributes to net ecosystem methane exchange even in methane-

consuming soils. In contrast, transitional wetland sites show high variability in both tree and soil

fluxes, with episodic conditions that can shift soils from sinks to sources, emphasizing the

importance of landscape context and moisture dynamics in determining net methane fluxes.

Figure 1. Temporal patterns of methane fluxes from tree stems and soils across a hydrological gradient at Yale-Myers Forest, Connecticut, June 2020 to May 2021. Data represent 342 tree flux measurements from 41 trees and 276 soil measurements from 12 soil plots, collected during 10 sampling campaigns. ***Top panel***: Distribution of volumetric water content (%) at 12 cm depth across upland (mean $16.7 \pm 8.2\%$, n=121), intermediate (mean $20.0 \pm 8.7\%$, n=47), and transitional wetland (mean $42.8 \pm 19.4\%$, n=119) sites, showing three distinct hydrological zones. ***Middle panel***: CH₄ flux measurements from tree stems (green) and soils (brown) with individual measurements shown as semi-transparent points and mean values $\pm$ SE for 7-day intervals as solid points connected by lines. Data displayed on pseudo-logarithmic scale with dashed line at zero flux. Upland sites showed tree emissions averaging 0.057 nmol m^{-2} s^{-1} contrasting with soil consumption of -1.57 nmol m^{-2} s^{-1}. Intermediate sites showed tree emissions of 0.011 nmol m^{-2} s^{-1} and soil consumption of -1.35 nmol m^{-2} s^{-1}. Transitional wetland trees averaged 0.129 nmol m^{-2} s^{-1} emissions while soils showed high variability (mean 1.83, median -0.104 nmol m^{-2} s^{-1}). ***Bottom panel***: Seasonal dynamics of soil temperature (red, ranging -0.7 to 19.3°C) and volumetric water content (blue) as loess-smoothed curves.

2. Methane Flux by Height and Analysis of Height Effects

Methane flux measurements at three stem heights (0.5, 1.25, 2.0 m) showed consistent positive fluxes for most upland species. Linear mixed-effects models revealed significant height effects in only 3 of 11 species tested. *Betula alleghaniensis* showed the strongest negative relationship (-0.824 nmol m^{-2} s^{-1} per meter stem height, p = 0.002), with fluxes decreasing from 1.52 ± 0.52 nmol m^{-2} s^{-1} at 0.5 m to 0.29 ± 0.08 nmol m^{-2} s^{-1} at 2.0 m. This robust effect (100% jackknife iterations significant; 11/14 trees with negative slopes) occurred where soil moisture (35.9% VWC) and soil methanogens ($10^{4.8}$ mcrA copies g^{-1}) were highest.

Tsuga canadensis (eastern hemlock) also showed a robust but much weaker negative height effect (-0.019 nmol m^{-2} s^{-1} per meter, p = 0.025; approximately 2% the magnitude of *B. alleghaniensis*), with 82% of jackknife iterations remaining significant and 71% of individual trees showing negative slopes. In contrast, *Acer saccharum* (sugar maple) displayed a positive relationship (+0.200 nmol m^{-2} s^{-1} per meter, p = 0.050) that was not robust, with only 44% of jackknife iterations remaining significant, driven largely by an exceptionally high flux value (1.94 nmol m^{-2} s^{-1}) observed at 2 m on one tree. When this outlier was removed, the height effect became weaker but remained positive (+0.056 nmol m^{-2} s^{-1} per meter, p = 0.022).

Across all species, height effect coefficients showed a strong negative correlation with soil methanogen abundance (r = -0.76, p = 0.007), indicating that species in areas with higher soil methanogens were more likely to show decreasing flux with height—a pattern consistent with soil-to-stem transport. The correlation with soil moisture alone was weaker and non-significant (r = -0.41, p = 0.21), though soil moisture and methanogen abundance were positively correlated (r = 0.50, p < 0.001).

The absence of height-dependent flux patterns in 8/11 species, combined with the strong correlation between estimated height effect and soil methanogen abundance, supports a dual-mechanism model: trees in high soil methanogen environments show height-dependent patterns

typically considered indicative of soil-to-stem transport, while trees in typical upland conditions with low soil methanogens exhibit lower but uniform vertical profiles, at least to 2m height, potentially consistent with internal production.

Figure 2. Methane flux patterns across stem heights and relationships with environmental drivers in upland forest tree species at Yale-Myers Forest, Summer 2021. Data from 434 flux measurements across 146 individual trees. Trees with multiple height measurements enabled mixed-effects modeling to separate within-tree from between-tree variation. **Top panel:** CH_4 flux measurements at three heights (0.5, 1.25, 2.0 m) for 15 tree species. Half-violin boxplots show flux distributions with individual measurements as jittered points (gray). Boxes indicate interquartile range with median; dashed horizontal lines indicate zero flux. Species arranged by height effect coefficient from most negative (left) to most positive (right). Sample sizes indicate number of individual trees per species. **Middle panel:** Height effect coefficients (nmol m^{-2} s^{-1} per cm) from linear mixed-effects models with 95% confidence intervals. Points and error bars colored by significance and direction: significant negative effects shown

in blue (Betula alleghaniensis p = 0.002, Tsuga canadensis p = 0.025), significant positive in red (Acer saccharum p
= 0.050), non-significant negative in light blue, and non-significant positive in light red. Solid circles indicate p <
0.05; open circles indicate p ≥ 0.05. **Bottom panel:** Environmental correlates as z-score normalized heatmap.
Variables include volumetric water content (VWC) and $\log_{10}$-transformed soil methanogen abundance (mcrA gene
copies g^{-1}) in organic, mineral, and depth-weighted soil layers. Blue intensity indicates relative values with darker
blue representing higher z-scores. Species ordered to match middle panel. Height effect coefficients showed strong
negative correlation with soil methanogen abundance (r = -0.76, p = 0.007).

3. Species Trends in Flux and Individual Variance Partitioning

Pooling breast-height (1.25 m) flux measurements from peak growing season surveys in 2021
and 2023 (n = 476 measurements; 141 from 2021, 335 from 2023) revealed substantial variation
both within and among species. These measurements were primarily from upland forest sites
where soils typically consume atmospheric methane. To quantify the relative importance of
different factors driving flux variation, we conducted variance partitioning analysis using linear
models that sequentially included environmental variables (temperature, moisture, tree diameter),
species identity, and their interactions. This analysis revealed that species identity explained only
5.3% of variance, species-environment interactions 8.7%, environmental factors alone <0.01%,
leaving 82.9% unexplained at the individual tree level.

Betula alleghaniensis (yellow birch) showed the highest mean emissions (0.279 ± 0.091 nmol
m^{-2} s^{-1}), followed by *Acer saccharum* (sugar maple, 0.188 ± 0.061 nmol m^{-2} s^{-1}) and *Acer*
rubrum (red maple, 0.170 ± 0.060 nmol m^{-2} s^{-1}). Gymnosperms including *Tsuga canadensis*
(eastern hemlock, 0.026 ± 0.007 nmol m^{-2} s^{-1}) and *Pinus strobus* (white pine, 0.021 ± 0.010
nmol m^{-2} s^{-1}) exhibited consistently lower emissions.

The dominance of unexplained individual variance (82.9%) indicates tree-specific
characteristics—potentially heartwood development, internal decay, microbial colonization
history, wounding, bark features, gas flowpaths, or physiological status—drive emissions more
than species identity or environmental conditions. Species-environment interactions (8.7%)
suggest differential environmental responses among species, though these remain secondary to
individual variability in upland forests.

Figure 3. Variance partitioning and species-level patterns of methane flux from upland forest trees. Data from 476 breast-height (1.25 m) flux measurements across 13 species during peak growing season (summer) surveys in 2021 and 2023 at Yale-Myers Forest upland sites. **Left panel:** Ridge density plots showing distribution of CH$_4$ flux by species, with individual measurements colored by soil volumetric water content (VWC). Black points with error bars show mean ± SE. Species ordered by mean flux. **Top right panel:** Variance components from linear models showing the proportion of flux variance explained by unexplained individual variation (82.9%), species-environment interactions (8.7%), species identity (5.3%), and environmental factors (<0.01%). **Bottom right panel:** Standardized effect sizes from full model showing environmental (green) and species (blue) effects with 95% confidence intervals. Solid points indicate p < 0.05; open points indicate p ≥ 0.05.

4. Quantifying Methanogens Across Tree and Soil Compartments

In a survey of 155 standing trees encompassing 16 species sampled during summer 2021, the vast majority harbored detectable methanogenic archaea within their living wood. Heartwood was particularly enriched in methanogens, with 97% of samples containing detectable mcrA genes. Among positive samples, mean abundance was $10^{3.2}$ copies g^{-1} dry wood (median $10^{2.7}$), with maximum values reaching $10^{6.9}$ copies g^{-1} in *Acer saccharum* (sugar maple). Over half (54%) of heartwood samples exceeded 500 mcrA copies g^{-1}. Sapwood showed significantly lower methanogen abundance (Wilcoxon rank-sum test, p < 0.001), with 69% of samples detectable, averaging $10^{2.3}$ copies g^{-1} (median $10^{2.1}$, max $10^{4.2}$) among positive samples.

Methanogens were substantially less abundant in soils surrounding trees. Only 59% of mineral
soil samples and 53% of organic soil samples contained detectable methanogens. Among
positive samples, mineral soils averaged $10^{1.9}$ copies g^{-1} wet soil (median $10^{1.6}$, max $10^{5.5}$),
while organic horizons averaged $10^{2.0}$ copies g^{-1} (median $10^{1.6}$, max $10^{5.6}$). Given average soil
moisture of 21% VWC, the median difference between heartwood and soil methanogen
abundances exceeded one order of magnitude, with only 6-8% of soil samples exceeding 500
mcrA copies g^{-1}. Notable exceptions occurred in transitional wetland positions, particularly
around *Betula alleghaniensis*, where soil methanogens reached $10^{5.5-5.6}$ copies g^{-1}.

Methanotrophs showed inverse abundance patterns. Combined pmoA and mmoX abundances
were lowest in heartwood ($10^{3.3}$ copies g^{-1}), intermediate in sapwood ($10^{3.5}$ copies g^{-1}), and
highest in soils ($10^{4.9}$ copies g^{-1} in both horizons). This inverse relationship suggests spatial
segregation of methane production and consumption processes.

Based on controlled spiking experiments with wood samples that we performed in related work
(Arnold et al 2024), wood DNA extraction recovers approximately 20% of target sequences,
with losses occurring during freeze-drying, cryo-grinding, and extraction steps. Thus, actual
wood methanogen abundances could potentially exceed our measurements by approximately 5-
fold.

Figure 4. Methanogen and methanotroph abundance across forest compartments from summer 2021 survey.
Left panel: Methanogen abundance (mcrA gene copies g^{-1}) across 16 tree species and four forest compartments at
Yale-Myers Forest, measured by droplet digital PCR with probe-based assays. Bars show mean ± SE on log₁₀ scale.
Species ordered by mean abundance within each compartment; bar colors represent phylogenetic distance.
Horizontal lines indicate compartment-wide geometric means. Sample sizes shown for each species. **Right panel:**
Relationship between methanogen (mcrA) and methanotroph (pmoA + mmoX) abundance across forest
compartments on pseudo-log scales. Points represent individual samples colored by compartment type. Marginal

density plots show distributions for each axis. Centroids (circles with black borders) indicate geometric mean values

for each compartment. n = 147 heartwood, 154 sapwood, 128 mineral soil, 135 organic soil samples.

5. Community and Taxonomic Identity of Methanogens

16S rRNA sequencing corroborated the ddPCR findings, revealing that methanogens constitute a

significant portion of the wood microbiome. In heartwood, methanogens ranged from 0 to 56.4%

of all microbial taxa (mean 3.3±9.3%), while in sapwood they ranged from 0 to 6.37% (mean

0.096±0.61%). Methanogen abundance varied significantly by tree species (ANOVA: 16S rRNA

p<0.01, ddPCR p<0.05), with certain trees serving as preferential hosts. *Acer saccharum*

heartwood contained the highest methanogen concentrations both in absolute terms ($10^{5.13}\pm10^{2.11}$

copies g^{-1}, p<0.1) and as a proportion of the total microbiome (15.7±13.0%, p<0.001). In

contrast, conifer species including *Pinus strobus* and *Tsuga canadensis* harbored the lowest

methanogenic loads (p<0.01), averaging approximately $10^{3.6}$ copies g^{-1} in heartwood and $10^{1.4}$

copies g^{-1} in sapwood.

The wood methanogenic community was dominated by two taxonomic families.

Methanobacteriaceae comprised 2.65±8.12% of heartwood communities on average but reached

up to 56.3% in some samples, while in sapwood they ranged from 0-6.37%.

Methanomassiliicoccaceae showed lower but still substantial abundances, averaging 0.63±2.42%

in heartwood with a maximum of 17.1%, and 0-0.63% in sapwood. Both families were

significantly more abundant in heartwood than sapwood (p<0.001). Other methanogenic groups

occurred only at trace amounts, with the third most abundant family, Methanosarcinaceae,

peaking at just 1.29% of the total community.

Soil methanogenic communities differed markedly from those in wood, being dominated by

Bathyarchaeia rather than wood-associated families. While trace amounts of

Methanobacteriaceae were detected in soils (mineral: 0.025±0.16%, organic: 0.032±0.19%),

Methanomassiliicoccaceae was virtually absent from soil samples. Meanwhile, methanotrophs

(known and potential/facultative) were elevated in soils, with different taxonomic composition

compared to wood (Fig. S1-S2). This compositional divergence between wood and soil

communities argues against passive colonization from soil and suggests selection for specific

methanogenic lineages adapted to the wood environment.

These two dominant wood-associated families utilize distinct metabolic pathways.

Methanobacteriaceae are hydrogenotrophic, primarily reducing CO_2 using H_2 or formate as

electron donors. Methanomassiliicoccaceae are methylotrophic, performing hydrogen-mediated

reduction of methyl groups from compounds such as methanol and potentially lignin-derived

substrates. The co-occurrence of these metabolically distinct groups indicates that multiple

pathways support methanogenesis within living wood, reflecting the diverse chemical

environments and substrate availability within tree tissues.

Figure 5: Methanogenic taxa relative abundance in tree and soil samples. (a) Methanogen relative abundance (percentage of total prokaryotic community) based on 16S rRNA gene sequencing across tree species and forest compartments. Data shown for heartwood, sapwood, mineral soil, and organic soil for 16 tree species at Yale-Myers Forest. (b) Methanogen taxonomic composition showing relative abundances of major methanogenic families across tree species and forest compartments. Stacked bars represent proportion of different methanogenic taxa.

6. Taxonomic Correlations and Inferred Metabolic Pathways

FAPROTAX analysis of 16S rRNA data (Fig. S3) suggested potential metabolic capabilities of detected taxa. Fermentation-related metabolisms were predicted to comprise 16.6% of heartwood and 6.8% of sapwood communities. Putative dark hydrogen oxidation, which requires H_2 availability, appeared enriched in heartwood (3.69%) compared to sapwood (1.15%, p<0.1), with the highest values (11.8%) in *Acer saccharum*. Predicted methylotrophy was also more frequent in heartwood (0.64%) than sapwood (0.02%, p<0.001), peaking at 5.07% in *Acer saccharum* heartwood (p<0.001). These inferred functions are consistent with the availability of hydrogen and methyl substrates that could support methanogenesis, although competition between methylotrophs and methanogens is also possible.

PICRUSt2 functional inference (Fig. S4) indicated differences in predicted metabolic pathways between heartwood and sapwood segments. Heartwood showed greater representation of methanogenesis-associated pathways, including reductive acetyl-CoA, H_2/CO_2 methanogenesis, and acetate methanogenesis. These co-occurred with predicted fermentation pathways (homolactic fermentation, mannan degradation, L-arginine biosynthesis) that may provide hydrogen, acetate, and formate substrates for methanogens. Sapwood, by contrast, was enriched

in predicted aerobic pathways such as purine nucleotide degradation and heme biosynthesis,
along with cell wall synthesis pathways including peptidoglycan and teichoic acid biosynthesis.

Significant correlations with *mcrA* gene copy number, quantified by ddPCR, revealed that
Christensenellaceae, Lachnospiraceae, and Ruminococcaceae were consistently associated with
methanogens (Methanobacteriaceae and Methanomassiliicoccaceae). These associations are
consistent with known syntrophic relationships, in which fermentative bacteria generate
metabolites that are consumed by methanogens. Christensenellaceae are obligate anaerobes that
ferment polysaccharides and proteins, yielding short-chain fatty acids and hydrogen (Morotomi
et al. 2012); Lachnospiraceae are versatile fermenters of plant-derived carbohydrates that
produce acetate, butyrate, lactate, formate, and hydrogen (Kaminsky et al. 2023; Zaplana et al.
2024); and Ruminococcaceae specialize in degrading cellulose and lignocellulose, releasing
acetate, formate, and hydrogen as fermentation products (Ntaikou et al. 2008; Gharechahi et al.
2023; Weimer 2022). Together, these families provide substrates that sustain methanogenic
archaea. Methanobacteriaceae are hydrogenotrophic methanogens that reduce CO_2 with H_2 to
methane, while Methanomassiliicoccaceae are methylotrophic methanogens that use methanol
and methylamines with H_2 as an electron donor (Buan 2018; Boone et al. 2015; Cozannet et al.
2020). Their significant association with *mcrA* copy number therefore supports the interpretation
of complementary metabolic roles that facilitate methane production in anoxic
microenvironments.

The spatial distribution of these pathways reflects a clear redox gradient within tree tissues. Inner
core segments appear to be more reduced and favor anaerobic processes, such as fermentation
and methanogenesis, while outer core segments are more oxidized, favoring aerobic metabolism.
These findings align with redox stratification, where metabolic processes occur along a gradient
from oxygen-rich to oxygen-depleted zones within tree stems, and align with our recent results
from related work finding higher methane and more variable oxygen in heartwood compared to
sapwood in trees at this site (Arnold et al. 2025).

Figure 6: Heatmap showing microbial community composition and methanogen abundance across tree core samples from Yale-Myers Forest. The figure displays bacterial family abundance (log10 16S rRNA gene copies) with samples organized by core type (inner vs. outer) and methanogen abundance (log10 mcrA gene copies). Each column represents an individual tree core sample, with the top color bars indicating 16S rRNA abundance, core type (teal for inner, purple for outer), and mcrA abundance (green gradient). Bacterial families are listed on the y-axis, with red intensity indicating relative abundance within each sample. Families significantly associated with methanogen abundance (p<0.01) highlighted in green on the left margin; non-significant associations are shown in gray. Key methanogenic families (Methanobacteriaceae, Christensenellaceae, Methanomassiliicoccaceae) appear at the bottom, showing higher abundance in samples with elevated mcrA levels, particularly in inner core samples.

7. Individual-Level Methanogen-Flux Relationships and Sampling Limitations

At the individual tree level, correlations between methanogen abundance and methane flux were weak ($R^2 < 0.1$ for both heartwood and sapwood), likely reflecting the heterogeneous distribution of methanogens within tree tissues given increment boring samples only a small fraction of stem volume. Correlations were similarly positive but weak between methanogen abundance and stem internal methane concentration (Fig. S5-S6). The sampling limitations were dramatically illustrated in an intensively sampled black oak (*Quercus velutina*) with measurements at seven heights (0-10 m). Methane fluxes peaked at mid-stem heights (4-6 m) rather than showing uniform or exponentially decreasing patterns, corresponding with the upper limit of visible heart rot. Internal methane concentrations exceeded 1,500 ppm at these heights, yet mcrA abundance from cores varied by over three orders of magnitude, with some samples showing no detectable methanogens despite elevated gas concentrations.

This mismatch between gas concentrations and microbial abundance indicates that methane
production may occur in localized hotspots associated with decay pockets or anaerobic
microsites rather than being uniformly distributed throughout the stem. These hotspots may drive
emissions yet be easily missed by standard increment core sampling, which would not
adequately capture the small-scale patchiness of methanogenic communities. The poor
individual-level correlations reflect regression attenuation: when measurement error in the
independent variable (mcrA abundance) is high due to inadequate spatial sampling, the observed
correlation with the dependent variable (flux) becomes systematically weakened.

Figure 7: Vertical profiles of internal gas concentrations, methanogen abundance, and methane flux in a
felled black oak (Quercus velutina) tree at Yale-Myers Forest. All measurements taken at seven heights along
the stem (0-10 m): (Left) Internal methane concentrations (CH4 ppm) with points colored by oxygen percentage
(17.0-19.0%, yellow to purple gradient). (Center) Methanogen abundance (mcrA gene copies/µL) in heartwood
(brown circles) and sapwood (blue circles) samples, with fitted trend lines. (Right) Methane flux measurements
(CH4 flux units) along the stem height with points colored by internal methane concentration (500-1500 ppm, yellow
to gray gradient). The black curves represent smoothed trend lines for each measurement.

8. Species-Level Methanogen-Flux Relationships

We next examined whether aggregating data to species level would reveal patterns obscured by
within-tree heterogeneity. For this analysis, we calculated area-weighted methanogen
abundances by integrating densities from inner (heartwood) and outer (sapwood) samples,
weighted by their respective cross-sectional areas. Samples were taken approximately 5 cm from
the pith and 5 cm from the bark using increment cores.

Species-level aggregation revealed significant correlations absent at the individual level. While individual trees showed no relationship between area-weighted mcrA and flux ($R^2 < 0.001$, n = 122 trees), species-level medians showed strong positive correlation ($R^2 = 0.40$, p = 0.050 for log-transformed data; n = 10 species with ≥5 observations each). The relationship remained consistent across transformations: untransformed (r = 0.58, p = 0.080) and log-log (r = 0.63, p = 0.049). Mean values produced weaker correlations than medians ($R^2 = 0.30$, p = 0.098), suggesting medians better represent typical species behavior. When normalized to outer stem surface area, methanogen load still correlated with emissions ($R^2 = 0.37$, p = 0.063).

This scale-dependent pattern, where microscale heterogeneity obscures relationships that emerge at broader scales, is not wholly unexpected given the small size of tree cores, the known hotspot-driven spatial variability of methane production, and the limited mobility within wood. The dramatic improvement demonstrates that species-level averaging reduces measurement noise from spatial heterogeneity, revealing an underlying biological relationship between methanogen abundance and methane flux that sampling limitations obscure at finer scales.

Figure 8. Methanogen distribution in tree cross-sections and relationship with methane flux across species at Yale-Myers Forest. (a) Interpolated radial distribution of mcrA gene abundance in tree stem cross-sections at breast height (1.25 m). Each circle represents an individual tree, organized by species (rows, n ≥ 3 trees per species) and individual trees within species (columns). Color intensity shows $\log_{10}$(mcrA copies g^{-1}) interpolated across a 0.5 cm resolution grid, with the plasma color scale showing radial gradients from heartwood (center) to sapwood (periphery). Concentrations were modeled with a linear transition zone between inner wood (≤5 cm from pith) and outer wood (≤5 cm from bark) based on ddPCR quantification of mcrA genes from increment cores. Circle diameter is proportional to tree DBH. (b) Correlation between median area-weighted mcrA abundance and median CH$_4$ flux aggregated by species. Area-weighted mcrA values integrate methanogen concentrations across stem cross-sectional areas, accounting for the relative contributions of heartwood and sapwood zones. Points represent species-level medians (n ≥ 5 trees per species for both measurements) with error bars showing interquartile ranges 25th-75th percentile). Blue line shows linear regression with 95% confidence interval (shaded). Statistical tests shown: Pearson r on $\log_{10}$-transformed data, Spearman ρ, and Kendall τ with p-values. (c) Proportion of trees exceeding methanogen abundance thresholds across species. Percentage of trees with area-weighted mcrA gene copies exceeding 1,000, 5,000, and 10,000 copies g^{-1}.

9. Random Forest Upscaling

Random forest models trained on tree and soil flux measurements predicted seasonal flux patterns across the forest plot (Fig. S7). The tree model achieved out-of-bag $R^2 = 0.15$, with within-species standardized diameter, soil moisture, species-specific moisture interactions (particularly for *Betula alleghaniensis* and *Fraxinus americana*), and taxonomic prior as top predictors. The soil model achieved $R^2 = 0.28$, with moisture-temperature interactions, soil moisture, soil temperature, and seasonal cyclicity as dominant predictors. Models incorporated chamber type as a predictor rather than applying post-hoc calibration. Feature importance was assessed using impurity-based metrics.

On a per-unit-area basis, tree stem emissions averaged 0.07 nmol m^{-2} s^{-1} across the year (monthly means ranged from 0.04-0.12 nmol m^{-2} s^{-1}) while soil uptake averaged 1.76 nmol m^{-2} s^{-1} (monthly means: 1.58-2.05 nmol m^{-2} s^{-1}). Per unit area, stem emissions represented 4.2% of soil uptake, with peak emissions in July (0.12 nmol m^{-2} s^{-1}).

At our study site, based on lateral stem surface area of cylinders to 2 m height (3,555 m^2 across 10.2 ha, representing 3.47% of plot area), tree emissions contributed 0.003 nmol m^{-2} s^{-1} on a plot-area basis, totaling 1.3 mg CH$_4$ m^{-2} yr^{-1} and offsetting 0.14% of soil uptake (890 mg CH$_4$ m^{-2} yr^{-1}). As a scaling exercise, applying our per-unit-area stem flux to the woody area index of 3.07 reported for temperate forests (Gauci et al. 2024)—assuming constant emissions across all woody surfaces—would yield 114 mg CH$_4$ m^{-2} yr^{-1}, offsetting 13% of soil uptake. These estimates are based on measurements restricted to 0-2 m stem height and lack empirical data on vertical flux profiles or site-specific woody area above 2 m. While we observed no evidence of declining fluxes within our measurement range and detected a flux hotspot at ~4 m height in one tree measured to 10 m, insufficient data exist to constrain these uncertainties.

Figure 9: Integrated seasonal and species-level patterns of methane flux across tree and soil compartments at Yale-Myers Forest. Top panel: Monthly random forest-estimated tree stem methane flux for December, April, July, and September, with individual trees represented as points colored by flux magnitude (nmol m^{-2} s^{-1}, plasma color scale) and sized proportionally to basal area (m²). **Middle panel**: Corresponding estimated soil methane flux for the same months, displayed as interpolated raster surfaces. **Bottom panel**: Spatial distribution of annual mean tree CH$_4$ flux for common canopy species. Points colored by annual mean flux (nmol m^{-2} s^{-1}) and sized by basal area.

Discussion

Patterns of Tree Methane Emission from Upland to Wetland

Our study reveals that methane fluxes from trees vary significantly across landscape positions, with upland trees emitting lower levels of methane compared to trees in wetter, lowland areas. This gradient aligns with moisture availability and redox conditions of the surrounding

environment. Upland soils are typically well-drained and oxygen-rich, conditions that promote
methane oxidation by soil methanotrophs. Despite these soils consistently consuming
atmospheric methane (mean -90.9 µg CH_4 m^{-2} hr^{-1}), trees in these areas still emit methane (mean
3.3 µg CH_4 m^{-2} hr^{-1}), suggesting internal methane production within tree tissues (Fig. 1).

Trees in transitional wetland areas exhibited the highest emission rates (mean 7.5 µg CH_4 m^{-2}
hr^{-1}), where soils showed extreme variability—sometimes consuming, sometimes emitting
methane. These higher tree emissions likely result from a combination of soil-derived methane
transport (in areas with high soil methanogens like *Betula alleghaniensis* sites with $10^{4.8}$ mcrA
copies g^{-1} soil), on top of equal or greater internal production by endophytic methanogens as
compared to upland trees. The consistent methane emissions observed across tree species and
landscape positions, even where soils are strong methane sinks, suggest internal production may
be widespread among trees, though emission magnitude varies considerably between individuals.

Methane fluxes measured at multiple heights along tree stems showed no consistent pattern of
decline with height across all but one of the upland tree species sampled, suggesting that the
source of methane in upland trees may differ from the soil-originated methane typically observed
in wetland trees (Fig. 2). In wetland environments, methane produced in anaerobic soils is
commonly transported through tree stems via the xylem or internal gas spaces, resulting in
higher fluxes near the base of the stem and an exponential decrease in methane flux with
increasing height, as methane diffuses outward through stem tissues and into the atmosphere
(Jeffrey et al. 2024; Anttila et al. 2024). This soil-origin transport pattern was not observed in the
primarily upland tree species in this study.

Instead, methane fluxes in upland trees were relatively consistent across stem heights, indicating
that methane production within the tree tissues themselves may be a more important contributor
to methane flux in these environments. This pattern suggests that internal anaerobic niches
within the tree stems—such as in the heartwood or other oxygen-depleted tissues—could serve
as sites for methanogenic activity. The presence of methanogenic archaea in tree cores, as
revealed through microbial community analysis, supports this hypothesis. These microbes are
capable of producing methane in anaerobic microenvironments within tree tissues, independent
of soil methane transport (and were confirmed to do so in anaerobic incubations in related work;
(Arnold et al. 2025).

Nonetheless, it is not possible to rule out some contribution from a soil origin. The observation
of methane emissions from trees growing on methane-consuming soils does not necessarily
indicate exclusive internal production. Multiple non-exclusive mechanisms could contribute to
this apparent paradox (Pitz et al. 2018). Deep soil layers below the zone of active methanotrophy
may harbor methanogenic communities whose products bypass surface oxidation through root
uptake and xylem transport (Bhullar et al. 2013; Watson et al. 1997; Nouchi et al. 1990; Schimel
1995; Bradford et al. 2001). Anoxic microsites within otherwise aerobic soils, associated with
aggregate interiors, root detrituspheres, or temporarily saturated zones, could serve as methane
sources (Lacroix et al. 2023; Zausig et al. 1993; Keiluweit et al. 2018). While our soil sampling

indicated a minimal role of soil methanogens in these communities, further work with RNA and
multi-omics could provide further confirmation to link with rates. Further, it is important to note
that the lack of decrease of flux within the lower portion of the stem could break down with
measurements to higher heights, where a trend could become apparent; or, likelier (as seen in our
high-resolution sampling of a black oak; Fig. 7)), that unpredictable patterns may occur, with
hotspots at different heights aboveground.

Prevalence of Methanogens in Living Trees

Among all samples analyzed, 97.3% of surveyed trees harbored measurable methanogen
populations in their heartwood, compared to 68.8% in sapwood, 58.6% in mineral soil, and
52.6% in organic soil (Fig. 4). With abundances reaching ~10^7copies g^{-1} in some samples, these
are not incidental colonizers but established communities representing more than half of the total
heartwood microbiome in extreme cases. The consistency of this pattern across multiple diverse
species suggests that methanogen colonization of heartwood is a widespread phenomenon in
temperate forests.

The two-order-of-magnitude enrichment of methanogens in wood relative to surrounding soils
indicates that trees provide a unique and favorable habitat for these obligate anaerobes.
Heartwood's low oxygen availability, resulting from limited gas diffusion and consumption by
aerobic microbes, creates anaerobic microsites suitable for methanogenesis; and such patterns
support an emerging understanding of a divergent prokaryotic heartwood microbiome in life and
in death (Cregger et al. 2018; Arnold et al. 2025; Moll et al. 2018). The presence of fermentative
bacteria producing hydrogen and simple organic acids may provide electron donors, while wood
polymers including lignin may supply methylated substrates for methylotrophic methanogens.
While we can only infer function in this work, some co-occurring clades detected here
(*Lachnospiraceae*, *Methanobacteriaceae*; Fig. 5) have been shown to demonstrate such
syntrophic methane production through mechanisms such as interspecies hydrogen transfer in
other environments (Ruaud et al. 2020).

The consistent presence of methanogens across diverse tree species (Yip et al. 2019; Harada et
al. 2024; Feng et al. 2022; Zeikus & Henning 1975) raises intriguing questions about the
evolutionary history and ecological significance of these associations. Tree-methanogen
relationships may represent an ancient symbiosis predating the divergence of major tree lineages,
or convergent colonization of a widespread ecological niche. The species-specific differences in
methanogen abundance, with highest levels in *Acer saccharum* and lowest in conifers, could
reflect variation in wood chemistry, heartwood formation processes, or historical association
with methanogens. Further, the patterns suggest that long-term changes in tree species
composition, forest management practices, or climate-driven shifts in wood properties such as
moisture or substrate availability could substantially alter ecosystem-level fluxes resulting from
these tree-hosted communities.

Distinct Methanogenic Communities in Wood vs Soil

The taxonomic composition of wood methanogens differs markedly from soil communities, with
wood dominated by Methanobacteriaceae and Methanomassiliicoccaceae while soils harbor
primarily Bathyarchaeia (Evans et al. 2015). This compositional divergence argues against
passive colonization from soil and instead suggests selection for specific methanogenic lineages
adapted to the wood environment. The co-occurrence of hydrogenotrophic
(Methanobacteriaceae) and methylotrophic (Methanomassiliicoccaceae) methanogens indicates
multiple metabolic pathways support methanogenesis within trees.

The syntrophic community structure revealed by co-occurrence analysis (Fig. 6) and functional
predictions (Fig. S3-S4). provides mechanistic insight into how methanogenesis is sustained. The
strong correlation between fermentation and methanogenesis pathways, combined with
anticorrelation with aerobic processes, depicts a stratified microbial ecosystem where spatial
segregation enables coexistence of incompatible metabolisms. This microscale heterogeneity
within seemingly homogeneous wood tissue underscores the importance of considering trees as
complex microbial habitats rather than sterile structural material.

Multi-scale Complexity in Linking Microbial Abundance to Methane Flux

The weak correlation between methanogen abundance and flux at the individual tree level
reflects multiple layers of complexity in tree-mediated methane cycling that our current methods
only partially capture. While we detected methanogens through DNA-based approaches targeting
the mcrA gene, this provides only a census of potential methane producers. DNA abundance
does not directly translate to metabolic activity (Rocca et al. 2015); dormant or inactive cells
retain their genetic material, while actively transcribing populations may represent a small
fraction of total community members. RNA-based approaches targeting mcrA transcripts, or
proteomic and metabolomic analyses, would provide more direct measures of active
methanogenesis but present substantial technical challenges in wood tissues due to low biomass
yields and complex matrix effects.

Beyond the limitations of molecular censusing, methane emissions represent the net result of
multiple interacting processes operating across different spatial and temporal scales. Flux
measurements integrate production, consumption, and transport processes that may be spatially
and temporally decoupled. Production occurs in anaerobic microsites where methanogens are
active; consumption happens wherever methanotrophs encounter methane under suitable
conditions; and transport depends on wood anatomy, moisture content, and pressure gradients.
Reactive transport further complicates this picture—methane may be oxidized during diffusion
through aerobic zones, creating steep concentration gradients that influence both flux magnitude
and spatial patterns.

Tree physiology fundamentally shapes these processes through anatomical and physiological
controls. Wood porosity determines gas diffusivity and the potential for advective transport.
Species with ring-porous anatomy may facilitate rapid vertical gas movement through large
earlywood vessels, while diffuse-porous species restrict transport to slower diffusive pathways.
Sapflow dynamics create diurnal and seasonal pressure fluctuations that can enhance or suppress
gas emissions (Barba, Poyatos, et al. 2019; Barba et al. 2021; Bréchet et al. 2025; Anttila et al.
2024). Radial cell architecture, including the presence of ray parenchyma, tyloses, and resin
ducts, may create preferential flow paths that channel gases along specific routes rather than
uniform radial diffusion (Sorz & Hietz 2006; Cǒté 1963). These anatomical features vary not
only among species but also within individual trees based on growth history, injury response, and
position within the stem. Furthermore, these processes operate on different timescales. Soil gas
production and consumption can shift rapidly, on hourly to daily timescales (Parolari et al.
2021), while internal gas production within trees may build up more slowly, persisting for weeks
to months in poorly ventilated or anaerobic stem tissues, and and transport rates may vary from
minutes for advective flow (Jeffrey et al. 2024) to days for diffusive movement across
centimeters of wood (Sorz & Hietz 2006).

Internal methane buildup illustrates this temporal complexity. Elevated concentrations
(approaching 10% in the highest samples; Fig. S1-S2) could result from either high production
rates in active methanogenic zones or low diffusivity creating accumulation even with modest
production. The relationship between concentration and flux is non-linear and depends on
boundary conditions, diffusion coefficients, and the connectivity of gas-filled pore spaces. Hot
moments—periods of elevated flux—may occur when changes in atmospheric pressure,
temperature, or tree water status suddenly enhance transport from zones of accumulated
methane. Such episodic releases could contribute disproportionately to time-integrated emissions
while being easily missed by discrete sampling (Bréchet et al. 2025; Anthony & Silver 2021).

Spatial heterogeneity presents perhaps the greatest challenge for linking microbial abundance to
flux. Our intensive sampling of the black oak dramatically illustrated this issue: even within
zones of elevated methane concentration and emission at 4-6 m height, increment cores showed
only a 50% detection rate for methanogens (Fig. 7). This suggests extremely localized
production sites—decay pockets, wound sites, or other anaerobic niches—that create methane
plumes extending well beyond the source. Standard increment boring, which samples <0.1% of
stem volume, has remarkably low probability of intersecting these hotspots. While we assume
primarily radial diffusion drives emissions, methane originating from a small hotspot may
diffuse three-dimensionally, creating an extended zone of elevated concentration and surface
emission. The measured flux at any point on the bark surface thus integrates contributions from
multiple potential sources at varying distances and depths within the stem.

Addressing these complexities requires methodological advances beyond our current approaches.
High-resolution sampling combining multiple cores at various depths and heights could better
capture the probability distribution of methanogenic zones. Physical reconstruction of wood

anatomy through techniques like X-ray computed tomography would reveal the three-
dimensional network of potential gas conduits and barriers. Combining such structural
information with multi-point gas sampling and isotopic tracers could help distinguish soil-
derived from internally-produced methane and map transport pathways (Megonigal et al. 2020).
Time-series measurements capturing diurnal and seasonal dynamics would reveal the importance
of hot moments and the relationship between environmental drivers and emission patterns.

From Individual Complexity to Species-Level Patterns

Despite this individual-level complexity (Fig. 3), the emergence of strong correlations when data
are aggregated to species level and weighted by cross-sectional area suggests that species-
average methanogen loads better reflect emission potential than point measurements (Fig. 8).
This aligns with recent findings that tree microbiomes show species-level conservation (Arnold
et al. 2025), indicating that wood chemistry, anatomy, and physiology create predictable niches
for microbial colonization. Our observation that area-weighted methanogen abundance correlates
with species-median flux ($R^2 = 0.40$) demonstrates that species-characteristic traits influence
both microbial communities and gas emissions, even amid substantial individual variation.

The dramatic improvement in correlation strength when aggregating from individual to species
level parallels observations in soil systems where microscale heterogeneity obscures
relationships that become apparent at larger scales (Nunan et al. 2020; Waring et al. 2016;
Polussa et al. 2021). This reflects how averaging reduces noise from spatial heterogeneity and
sampling error while preserving the underlying biological signal, illustrating the ecological
principle that patterns obscured by fine-scale variation often emerge at broader scales of
organization. For practical purposes, species-level relationships may prove more useful for
predicting forest-scale methane budgets than attempts to characterize every individual tree's
heterogeneous internal landscape.

Moving forward, reconciling individual and species-level patterns requires embracing this
complexity rather than seeking simple universal relationships. Trees are not homogeneous
chemical reactors but spatially structured habitats where microbial processes, physical transport,
and plant physiology interact across multiple scales. Understanding these interactions demands
interdisciplinary approaches combining microbial ecology, tree physiology, soil science, and
transport physics—a synthesis that promises to reveal how living trees function as dynamic
components of the global methane cycle.

Methodological Considerations

Our study highlights important methodological challenges in quantifying tree-associated
methane cycling. The high spatial heterogeneity of methanogens within trees necessitates
sampling strategies that capture representative volumes rather than point measurements. Future
studies should consider computed tomography or other non-destructive imaging to map
anaerobic zones within trees, combined with targeted sampling of these regions.

The dominance of unexplained variance at the individual tree level indicates that unmeasured
factors strongly influence emissions. Potential drivers include tree health status, presence of heart
rot or other decay, hydraulic properties affecting internal oxygen dynamics, and historical
colonization events. Long-term monitoring of individual trees could help identify whether
emissions are stable traits or vary with tree physiological status.

Implications for Forest Methane Budgets

Our results highlight a fundamental scaling challenge in quantifying tree contributions to forest
methane budgets. At our site, based on lateral stem surface area to 2 m height (3.47% of plot
area), tree emissions offset 0.14% of soil methane consumption (1.3 vs. 890 mg CH_4 m^{-2} yr^{-1};
Fig. 9). However, extrapolating our per-unit-area fluxes to a woody area index of 3.07 (Gauci et
al., 2024), assuming constant emissions across all woody surfaces, yields 114 mg CH_4 m^{-2} yr^{-1}
(13% offset). Whether tree emissions represent a negligible fraction or a substantial offset
depends critically on vertical flux patterns and total woody surface area, both of which remain
largely uncharacterized.

Within our limited measurement range (0-2 m), we observed no evidence of declining fluxes
with height, and detected a flux hotspot at ~4 m in the one tree measured to 10 m. Notably, we
saw no evidence of a recently proposed switch to net uptake with increasing stem height (Gauci
et al. 2024). However, these sparse observations cannot constrain vertical profiles across the
canopy. Without systematic vertical profiling across multiple trees and species, we cannot
determine whether our 0-2 m measurements accurately represent fluxes at canopy height.

Our upland site likely differs mechanistically from wetland systems in the source of tree-stem
methane emissions. In wetlands, exponential decline in flux with height results from diffusive
loss of soil-transported methane (Pangala et al. 2013; Barba, Bradford, et al. 2019). If upland tree
emissions derive primarily from internal stem production rather than basal transport, emissions
would scale with total woody surface area rather than proximity to soil. This mechanistic
difference has critical implications for scaling: internal production suggests fluxes may remain
substantial throughout the canopy, making measurements restricted to lower stems potentially
unrepresentative of whole-tree contributions.

Resolving these uncertainties requires addressing two interconnected measurement challenges.
First, quantifying woody surface area demands direct three-dimensional measurements rather

than allometric estimates. Terrestrial laser scanning (TLS) offers promise for deriving site-
specific woody area component measurements and allometries (Calders et al. 2020). Second,
measuring stem fluxes throughout the canopy requires climbing, canopy access systems, towers,
or walkways—infrastructure rarely available across sites or seasons. Top-down constraints from
tower-based or airborne eddy covariance offer an alternative, allowing deduction of tree
contributions from differences between net ecosystem and soil fluxes (Pangala et al. 2017).
However, towers in upland forests remain sparse, and small tree emissions relative to dominant
soil uptake may fall below detection limits or prove difficult to separate from contributions of
nested wetland areas (Delwiche et al. 2021). The combination of low signal strength, high spatial
variability, and limited infrastructure for canopy-scale measurements represents a major frontier
in understanding tree-stem methane dynamics in upland forests.

Conclusions

This study demonstrates that upland temperate forest trees harbor abundant methanogenic
communities that drive consistent methane emissions independent of soil processes. The
discovery that wood provides a more favorable habitat for methanogens than surrounding soils,
combined with evidence for specialized wood-adapted methanogenic communities,
fundamentally alters our understanding of forest methane cycling. Tree emissions partially offset
soil methane consumption, reducing the net sink strength of upland forests and possibly
representing a climate-relevant flux that may respond differently to environmental change than
traditional soil-atmosphere exchange. Recognition of trees as active players in methane cycling
invites revision of forest greenhouse gas budgets and consideration of how forest management
practices influence this previously overlooked emission source.

Acknowledgements

We thank Josep Barba, Claire Butler, Luke Jeffrey, Andrew Reinmann, David Woodbury, Cade Brown, Ari Gewirtzman, Thomas Harris, Naomi Hegwood, Marsh Hlavka, Fiona Jevon, Ellie Jose, Jonas Karosas, Talia Kolodkin, Camila Ledezma, Laura Logozzo, Taylor Maavara, Naomi Norbraten, Alex Polussa, Adriana Rubenstein, Cyrena Thibodeau, Les Welker, Qespi Wood, Michelle Spicer, and Jackie Matthes for their contributions to this work. We acknowledge our additional collaborators on the forest inventory plot: Liza Comita, Mark Ashton, Simon Queenborough, Stuart Davies and Sean McMahon. We thank Yale School Forests, University of Minnesota Genomics Center, Yale Analytical and Stable Isotope Center, W.M. Keck Biotechnology Resource Laboratory, Yale Center for Genetic Analyses of Biodiversity, and Yale Chemistry Glass Shop for technical support and facilities access.

Funding

This research was supported by the National Science Foundation Graduate Research Fellowship, Yale-Myers Forest Kohlberg-Donohoe Fellowship, Yale Institute for Biospheric Studies Early Grant, Yale School of the Environment-Yale Institute for Biospheric Studies Fellowship, and Yale Planetary Solutions Project Seed Funding.

Competing Interests

The authors declare no competing interests.

Data Availability

The 16S rRNA amplicon sequencing data generated in this study have been deposited in the NCBI Sequence Read Archive (SRA) under BioProject accession number PRJNA1124946. Methane flux measurements, Methanogen and methanotroph abundance data (ddPCR), internal gas concentration data are available at Figshare: https://figshare.com/s/b781d3c5c24f1089ed5d?. Tree and soil characterization data, and associated metadata will be made available upon publication. R code for statistical analyses, Random Forest models, variance partitioning, and figure generation is available on GitHub: https://github.com/jgewirtzman/tree-methanogens and will be made public upon publication.

Author Contributions

J.G. and W.A. contributed equally to this work. J.G. and W.A. conceived and designed the research, curated and analyzed data, acquired funding, conducted fieldwork and laboratory investigations, developed methodology, administered the project, provided resources, developed software and code, supervised research activities, and validated results. M.T. contributed to conceptualization, curated data, conducted investigations, and developed methodology. C.M. performed formal analysis and developed software. J.G., W.A., and C.M. created visualizations. H.B. and D.W. curated data, conducted investigations, and developed methodology. N.W., K.K., and L.G. curated data and conducted investigations. C.R.B. and M.D. acquired funding, developed methodology, and provided resources. P.A.R. contributed to conceptualization, acquired funding, and supervised research. J.P. acquired funding, developed methodology, administered the project, provided resources, and supervised research. M.A.B. acquired funding, developed methodology, administered the project, and supervised research. J.G. wrote the original draft. All authors reviewed and edited the manuscript and approved the final version.

References

Anthony, T.L. & Silver, W.L., 2021. Hot moments drive extreme nitrous oxide and methane emissions from agricultural peatlands. *Global change biology*, 27(20), pp.5141–5153.

Anttila, J. et al., 2024. Model of methane transport in tree stems: Case study of sap flow and radial diffusion. *Plant, cell & environment*, 47(1), pp.140–155.

Arnold, W. et al., 2025. A diverse and distinct microbiome inside living trees. *Nature*, pp.1–10.

Arnold, W. et al., 2024. A method for sampling the living wood microbiome. *Methods in ecology and evolution / British Ecological Society*. Available at: https://besjournals.onlinelibrary.wiley.com/doi/10.1111/2041-210X.14311.

Barba, J., Bradford, M.A., et al., 2019. Methane emissions from tree stems: a new frontier in the global carbon cycle. *The New phytologist*, 222(1), pp.18–28.

Barba, J. et al., 2024. Methane emissions from tree stems - current knowledge and challenges: an introduction to a Virtual Issue. *The new phytologist*, 241(4), pp.1377–1380.

Barba, J. et al., 2021. Spatiotemporal variability and origin of CO2 and CH4 tree stem fluxes in an upland forest. *Global change biology*, 27(19), pp.4879–4893.

Barba, J., Poyatos, R. & Vargas, R., 2019. Automated measurements of greenhouse gases fluxes from tree stems and soils: magnitudes, patterns and drivers. *Scientific reports*, 9(1), p.4005.

Bates, D. et al., 2015. Fitting Linear Mixed-Effects Models Using \textbflme4. *Journal of statistical software*, 67(1). Available at: http://www.jstatsoft.org/v67/i01/ [Accessed September 16, 2021].

Bhullar, G.S., Edwards, P.J. & Olde Venterink, H., 2013. Variation in the plant-mediated methane transport and its importance for methane emission from intact wetland peat mesocosms. *Journal of plant ecology*, 6(4), pp.298–304.

Boone, D.R., Whitman, W.B. & Koga, Y., 2015. Methanobacteriaceae. *Bergey's Manual of Systematics of Archaea and Bacteria*, pp.1–1. Available at: http://dx.doi.org/10.1002/9781118960608.fbm00097.

Bradford, M.A. et al., 2001. Role of CH4 oxidation, production and transport in forest soil CH4 flux. *Soil biology & biochemistry*, 33(12-13), pp.1625–1631.

Bréchet, L.M. et al., 2025. Insights into the subdaily variations in methane, nitrous oxide and carbon dioxide fluxes from upland tropical tree stems. *The new phytologist*, 245(6), pp.2451–2466.

Buan, N.R., 2018. Methanogens: pushing the boundaries of biology. *Emerging topics in life sciences*, 2(4), pp.629–646.

Calders, K. et al., 2020. Terrestrial laser scanning in forest ecology: Expanding the horizon. *Remote sensing of environment*, 251, p.112102.

Callahan, B.J. et al., 2016. DADA2: High-resolution sample inference from Illumina amplicon data. *Nature methods*, 13(7), pp.581–583.

Carmichael, M.J. et al., 2024. Microbial Communities in Standing Dead Trees in Ghost Forests are Largely Aerobic, Saprophytic, and Methanotrophic. *Current microbiology*, 81(8), p.229.

Côté, W.A., Jr, 1963. Structural factors affecting the permeability of wood. *Journal of Polymer*

Science Part C Polymer Symposia, 2(1), pp.231–242.

Covey, K.R. et al., 2012. Elevated methane concentrations in trees of an upland forest.
Geophysical research letters, 39(15). Available at:
https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2012GL052361 [Accessed March
10, 2021].

Covey, K.R. & Megonigal, J.P., 2019. Methane production and emissions in trees and forests.
The New phytologist, 222(1), pp.35–51.

Cozannet, M. et al., 2020. New insights into the ecology and physiology of
Methanomassiliicoccales from terrestrial and aquatic environments. *Microorganisms*, 9(1),
p.30.

Cregger, M.A. et al., 2018. The Populus holobiont: dissecting the effects of plant niches and
genotype on the microbiome. *Microbiome*, 6(1), p.31.

Delwiche, K.B. et al., 2021. FLUXNET-CH_4: a global, multi-ecosystem dataset and analysis of
methane seasonality from freshwater wetlands. *Earth system science data*, 13(7), pp.3607–
3689.

Douglas, G.M. et al., 2020. PICRUSt2 for prediction of metagenome functions. *Nature*
biotechnology, 38(6), pp.685–688.

Epron, D. et al., 2022. Variability in stem methane emissions and wood methane production of
different tree species in a cold temperate mountain forest. *Ecosystems (New York, N.Y.)*,
26(4), pp.784–799.

Evans, P.N. et al., 2015. Methane metabolism in the archaeal phylum Bathyarchaeota revealed
by genome-centric metagenomics. *Science*, 350(6259), pp.434–438.

Feng, H. et al., 2022. Methane emissions may be driven by hydrogenotrophic methanogens
inhabiting the stem tissues of poplar. *The New phytologist*, 233(1), pp.182–193.

Flanagan, L.B. et al., 2021. Multiple processes contribute to methane emission in a riparian
cottonwood forest ecosystem. *The New phytologist*, 229(4), pp.1970–1982.

Forster, P. et al., 2021. The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity.
In V. Masson-Delmotte et al., eds. *Climate Change 2021: The Physical Science Basis.*
Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental
Panel on Climate Change. Cambridge, United Kingdom and New York, NY, USA:
Cambridge University Press, pp. 923–1054.

Gauci, V. et al., 2024. Global atmospheric methane uptake by upland tree woody surfaces.
Nature, 631(8022), pp.796–800.

Gauci, V., 2025. Tree methane exchange in a changing world. *Nature reviews. Earth &*
environment, pp.1–13.

Gharechahi, J. et al., 2023. Lignocellulose degradation by rumen bacterial communities: New
insights from metagenome analyses. *Environmental research*, 229(115925), p.115925.

Harada, M. et al., 2024. Ubiquity of methanogenic archaea in the trunk of coniferous and
broadleaved tree species in a mountain forest. *Antonie van Leeuwenhoek*, 117(1), p.107.

Hettwer, C. et al., 2025. Methane flux from living tree stems in a northern conifer forest.

Biogeochemistry, 168(4), pp.1–15.

Hüppi, R. et al., 2018. Restricting the nonlinearity parameter in soil greenhouse gas flux
calculation for more reliable flux estimates. *PloS one*, 13(7), p.e0200876.

Hutchinson, G.L. & Mosier, A.R., 1981. Improved soil cover method for field measurement of
nitrous oxide fluxes. *Soil Science Society of America journal. Soil Science Society of*
America, 45(2), pp.311–316.

Jeffrey, L.C. et al., 2021. Bark-dwelling methanotrophic bacteria decrease methane emissions
from trees. *Nature communications*, 12(1), p.2127.

Jeffrey, L.C. et al., 2023. Large methane emissions from tree stems complicate the wetland
methane budget. *Journal of geophysical research. Biogeosciences*, 128(12). Available at:
https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2023JG007679.

Jeffrey, L.C. et al., 2024. Rapid bark-mediated tree stem methane transport occurs independently
of the transpiration stream in Melaleuca quinquenervia. *The new phytologist*, 242(1), pp.49–
60.

Jeffrey, L.C. et al., 2019. Rhizosphere to the atmosphere: contrasting methane pathways, fluxes,
and geochemical drivers across the terrestrial–aquatic wetland boundary. *Biogeosciences* ,
16(8), pp.1799–1815.

Jensen, K.F., 1967. Measuring oxygen and carbon dioxide in red oak trees. *Research Note NE-*
74. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest
Experiment Station. 4p., 74. Available at:
https://research.fs.usda.gov/treesearch/download/10474.pdf [Accessed February 13, 2025].

Kaminsky, R.A. et al., 2023. Rumen *Lachnospiraceae* isolate NK3A20 exhibits metabolic
flexibility in response to substrate and coculture with a methanogen. *Applied and*
Environmental Microbiology, 89(10), pp.e00634–23.

Keiluweit, M. et al., 2018. Anoxic microsites in upland soils dominantly controlled by clay
content. *Soil biology & biochemistry*, 118, pp.42–50.

Kolb, S. et al., 2003. Quantitative Detection of Methanotrophs in Soil by Novel pmoA-Targeted
Real-Time PCR Assays. *Applied and environmental microbiology*, 69(5), pp.2423–2429.

Lacroix, E.M. et al., 2023. Consider the anoxic microsite: Acknowledging and appreciating
spatiotemporal redox heterogeneity in soils and sediments. *ACS earth & space chemistry*,
7(9), pp.1592–1609.

Le Mer, J. & Roger, P., 2001. Production, oxidation, emission and consumption of methane by
soils: A review. *European journal of soil biology*, 37(1), pp.25–50.

Leung, P.M. et al., 2024. Wetland tree barks are dynamic hotspots for microbial trace gas
cycling. *bioRxiv*, p.2024.07.02.601631. Available at:
https://www.biorxiv.org/content/10.1101/2024.07.02.601631v1.abstract [Accessed August
13, 2024].

Liu, C. et al., 2021. microeco: an R package for data mining in microbial community ecology.
FEMS microbiology ecology, 97(2). Available at: http://dx.doi.org/10.1093/femsec/fiaa255.

Louca, S., Parfrey, L.W. & Doebeli, M., 2016. Decoupling function and taxonomy in the global

ocean microbiome. *Science*, 353(6305), pp.1272–1277.

Luesken, F.A. et al., 2011. pmoA Primers for detection of anaerobic methanotrophs. *Applied and*
environmental microbiology, 77(11), pp.3877–3880.

Machacova, K. et al., 2023. Methane emission from stems of European beech (Fagus sylvatica)
offsets as much as half of methane oxidation in soil. *The new phytologist*, 238(2), pp.584–
597.

McDonald, I.R., Kenna, E.M. & Murrell, J.C., 1995. Detection of methanotrophic bacteria in
environmental samples with the PCR. *Applied and environmental microbiology*, 61(1),
pp.116–121.

McDonald, I.R. & Murrell, J.C., 1997. The particulate methane monooxygenase gene pmoA and
its use as a functional gene probe for methanotrophs. *FEMS microbiology letters*, 156(2),
pp.205–210.

McMurdie, P.J. & Holmes, S., 2013. phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data M. Watson, ed. *PloS one*, 8(4),
p.e61217.

Megonigal, J.P., Brewer, P.E. & Knee, K.L., 2020. Radon as a natural tracer of gas transport
through trees. *The new phytologist*, 225(4), pp.1470–1475.

Mieszkin, S. et al., 2021. Oak decaying wood harbors taxonomically and functionally different
bacterial communities in sapwood and heartwood. *Soil biology & biochemistry*, 155,
p.108160.

Mochidome, T. et al., 2025. Local methanogenesis drives significant methane emissions from
upper tree trunks in a cool-temperate upland forest. *The new phytologist*. Available at:
http://dx.doi.org/10.1111/nph.70331 [Accessed July 29, 2025].

Moisan, M.-A. et al., 2025. Tissue humidity and pH as important species traits regulating tree
methane emissions in floodplain wetland forests. *The new phytologist*, (nph.70543).
Available at: http://dx.doi.org/10.1111/nph.70543 [Accessed September 9, 2025].

Moll, J. et al., 2018. Bacteria inhabiting deadwood of 13 tree species are heterogeneously
distributed between sapwood and heartwood. *Environmental microbiology*, 20(10),
pp.3744–3756.

Morotomi, M., Nagai, F. & Watanabe, Y., 2012. Description of Christensenella minuta gen.
nov., sp. nov., isolated from human faeces, which forms a distinct branch in the order
Clostridiales, and proposal of Christensenellaceae fam. nov. *International journal of*
systematic and evolutionary microbiology, 62(Pt 1), pp.144–149.

Nouchi, I., Mariko, S. & Aoki, K., 1990. Mechanism of methane transport from the rhizosphere
to the atmosphere through rice plants. *Plant physiology*, 94(1), pp.59–66.

Ntaikou, I. et al., 2008. Hydrogen production from sugars and sweet sorghum biomass using
Ruminococcus albus. *International journal of hydrogen energy*, 33(4), pp.1153–1163.

Nunan, N., Schmidt, H. & Raynaud, X., 2020. The ecology of heterogeneity: soil bacterial
communities and C dynamics. *Philosophical transactions of the Royal Society of London.*
Series B, Biological sciences, 375(1798), p.20190249.

Pangala, S.R. et al., 2017. Large emissions from floodplain trees close the Amazon methane
budget. *Nature*, 552(7684), pp.230–234.
Pangala, S.R. et al., 2015. The contribution of trees to ecosystem methane emissions in a
temperate forested wetland. *Global change biology*, 21(7), pp.2642–2654.
Pangala, S.R. et al., 2013. Trees are major conduits for methane egress from tropical forested
wetlands. *The New phytologist*, 197(2), pp.524–531.
Parolari, A.J., Sizemore, J. & Katul, G.G., 2021. Multiscale legacy responses of soil gas
concentrations to soil moisture and temperature fluctuations. *Journal of geophysical*
research. Biogeosciences, 126(2). Available at: http://dx.doi.org/10.1029/2020jg005865.
Pitz, S.L. et al., 2018. Methane fluxes from tree stems and soils along a habitat gradient.
Biogeochemistry, 137(3), pp.307–320.
Pitz, S. & Megonigal, J.P., 2017. Temperate forest methane sink diminished by tree emissions.
The New phytologist, 214(4), pp.1432–1439.
Polussa, A. et al., 2021. Scale dependence in functional equivalence and difference in the soil
microbiome. *Soil biology & biochemistry*, 163(108451), p.108451.
Putkinen, A. et al., 2021. New insight to the role of microbes in the methane exchange in trees:
evidence from metagenomic sequencing. *The New phytologist*, 231(2), pp.524–536.
Quast, C. et al., 2013. The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. *Nucleic acids research*, 41(Database issue), pp.D590–6.
Rheault, K., Christiansen, J.R. & Larsen, K.S., 2024. goFlux: A user-friendly way to calculate
GHG fluxes yourself, regardless of user experience. *Journal of open source software*, 9(96),
p.6393.
Rocca, J.D. et al., 2015. Relationships between protein-encoding gene abundance and
corresponding process are commonly assumed yet rarely observed. *The ISME journal*, 9(8),
pp.1693–1699.
Ruaud, A. et al., 2020. Syntrophy via interspecies H $_2$ transfer between *Christensenella* and
Methanobrevibacter underlies their global cooccurrence in the human gut. *mBio*, 11(1).
Available at: http://dx.doi.org/10.1128/mbio.03235-19.
Schimel, J.P., 1995. Plant transport and methane production as controls on methane flux from
arctic wet meadow tundra. *Biogeochemistry*, 28(3), pp.183–200.
Schink, B. & Ward, J.C., 1984. Microaerobic and Anaerobic Bacterial Activities Involved in
Formation of Wetwood and Discoloured Wood. *IAWA journal / International Association of*
Wood Anatomists, 5(2), pp.105–109.
Shigo, A.L. & Hillis, W.E., 1973. Heartwood, Discolored Wood, and Microorganisms in Living
Trees. *Annual review of phytopathology*, 11(1), pp.197–222.
Siegenthaler, A. et al., 2016. Technical Note: Semi-rigid chambers for methane gas flux
measurements on tree stems. *Biogeosciences* , 13(4), pp.1197–1207.
Sorz, J. & Hietz, P., 2006. Gas diffusion through wood: implications for oxygen supply. *Trees*,
20(1), pp.34–41.
Steinberg, L.M. & Regan, J.M., 2009. mcrA-targeted real-time quantitative PCR method to

examine methanogen communities. *Applied and environmental microbiology*, 75(13), pp.4435–4442.

Uroz, S. et al., 2016. Ecology of the forest microbiome: Highlights of temperate and boreal ecosystems. *Soil biology & biochemistry*, 103, pp.471–488.

Waring, B.G. et al., 2016. Scale-dependent variation in nitrogen cycling and soil fungal communities along gradients of forest composition and age in regenerating tropical dry forests. *The new phytologist*, 209(2), pp.845–854.

Warner, D.L. et al., 2017. Carbon Dioxide and Methane Fluxes From Tree Stems, Coarse Woody Debris, and Soils in an Upland Temperate Forest. *Ecosystems* , 20(6), pp.1205–1216.

Watson, A. et al., 1997. Oxidation of methane in peat: Kinetics of CH4 and O2 removal and the role of plant roots. *Soil biology & biochemistry*, 29(8), pp.1257–1267.

Weber, N. et al., 2018. Nephele: a cloud platform for simplified, standardized and reproducible microbiome data analysis B. Berger, ed. *Bioinformatics* , 34(8), pp.1411–1413.

Weimer, P.J., 2022. Degradation of cellulose and hemicellulose by ruminal microorganisms. *Microorganisms*, 10(12), p.2345.

Wood, S.A. et al., 2023. Mitigating near-term climate change. *Environmental research letters*, 18(10), p.101002.

Wright, M.N. & Ziegler, A., 2017. ranger: A Fast Implementation of Random Forests for High Dimensional Data in C++ and R. *Journal of Statistical Software*, 77(1), pp.1–17. Available at: http://dx.doi.org/10.18637/jss.v077.i01.

Yip, D.Z. et al., 2019. Methanogenic Archaea dominate mature heartwood habitats of Eastern Cottonwood (Populus deltoides). *The New phytologist*, 222(1), pp.115–121.

Zaplana, T., Miele, S. & Tolonen, A.C., 2024. Lachnospiraceae are emerging industrial biocatalysts and biotherapeutics. *Frontiers in Bioengineering and Biotechnology*, 11 - 2023. Available at: http://dx.doi.org/10.3389/fbioe.2023.1324396.

Zausig, J., Stepniewski, W. & Horn, R., 1993. Oxygen concentration and redox potential gradients in unsaturated model soil aggregates. *Soil Science Society of America journal. Soil Science Society of America*, 57(4), pp.908–916.

Zeikus, J.G. & Henning, D.L., 1975. Methanobacterium arbophilicum sp.nov. An obligate anaerobe isolated from wetwood of living trees. *Antonie van Leeuwenhoek*, 41(4), pp.543–552.

Zeikus, J.G. & Ward, J.C., 1974. Methane Formation in Living Trees: A Microbial Origin. *Science*, 184(4142), pp.1181–1183.

1147   **Supporting Information**

1148   **SI Figures**

1149

1150

Figure S1: Mean relative abundances of potentially methanotrophic groups based on 16S rRNA sequencing
across tree species and forest compartments. Bars show mean relative abundance (log$_{10}$-transformed) ± SE of
bacterial families with putative or facultative methanotrophic capabilities across tree and soil samples at Yale-Myers
Forest. Data shown for inner wood (heartwood-enriched), outer wood (sapwood-enriched), mineral soil, and organic
soil samples. Species codes shown on x-axis (see main text for full species names). Families colored by taxonomy
include taxa with confirmed or potential methane-oxidizing capabilities (Methylococcaceae, Methylocystaceae,
Methylacidiphilaceae) as well as facultative methylotrophs that may metabolize methane under certain conditions
(Methylophilaceae, Methylomonadaceae, and others). These taxa were most abundant in soil samples, particularly
mineral and organic horizons, with lower abundances detected in wood tissues. This pattern contrasts with
methanogens (Figure 5), which showed enrichment in heartwood relative to soils, reflecting the spatial segregation
of methane production (anaerobic wood tissues) and consumption (aerobic soils and potentially outer wood layers)
processes.

Figure S2: Methanogen and methanotroph abundance in a single black oak (Quercus velutina) tree across
tissue types. Heatmap showing the distribution of methanogenic and methanotrophic taxa based on 16S rRNA gene
sequencing across different tissues of a felled black oak at Yale-Myers Forest. Each column represents a different
tissue type: heartwood, sapwood, bark, branches, leaves, roots, leaf litter, and surrounding soil. Rows show bacterial
families grouped by functional role (methanogens vs. methanotrophs). Color intensity indicates $\log_{10}$-transformed
relative abundance, with warmer colors (yellow-orange) showing higher abundance and cooler colors (blue-purple)
showing lower abundance. Methanobacteriaceae (methanogens) were detected exclusively in a pocket of heart rot
within the heartwood, demonstrating the localized nature of methanogenic colonization. Methanotrophic and
methylotrophic taxa (Beijerinckiaceae families including Methylocella, Methyloferula, Methylorosula, and
Methylobacterium-Methylorubrum) were distributed across various intern tissues.

Figure S3: Average relative abundance of different metabolisms. a) Average relative abundance of different metabolisms per core type, as determined via FAPROTAX b) Average relative abundance of different metabolisms per species (heartwood), as determined via FAPROTAX

Figure S4: Functional metabolic pathway analysis comparing inner and outer tree core segments based on PiCRUSt2 predictions. The heatmap shows the relative abundance of predicted metabolic pathways across tree core samples, organized by core type (inner in teal, outer in red) and methanogen abundance (log10 mcrA qPCR copies, green gradient at top). Each row represents a specific metabolic pathway, with pathway names listed on the y-axis. Color intensity indicates mean pathway abundance counts, with red showing higher abundance and blue showing lower abundance. The right color bar shows statistical significance (-logFDR), with green indicating significantly enriched pathways. Inner core samples show enrichment of anaerobic pathways including methanogenesis from H_2 and CO_2, methanogenesis from acetate, and various fermentation pathways. Outer core samples are enriched in aerobic pathways such as purine nucleotide degradation (aerobic), heme biosynthesis (aerobic), and sulfur metabolism pathways, reflecting the redox gradient from anaerobic inner cores to more oxygenated outer cores.

1191

Figure S5. Species-level variation in internal methane and oxygen concentrations from upland forest tree stems at Yale-Myers Forest. Internal CH₄ concentration measurements at breast height (1.25 m) across 16 tree species, with individual trees shown as points distributed using a beeswarm layout to visualize sample density. Points colored by internal O₂ concentration, with cooler colors (blue) indicating lower oxygen and warmer colors (red) indicating higher oxygen availability. Black points with error bars show species mean ± standard error. Species ordered by mean CH₄ concentration from lowest (bottom) to highest (top). Y-axis displayed on pseudo-logarithmic scale to accommodate wide concentration ranges.

1199

Figure S6. Internal gas composition and methanogen abundance correlate with methane flux from upland forest tree stems. (A) CH_4 flux versus internal CO_2 concentration. **(B)** CH_4 flux versus internal O_2 concentration. **(C)** CH_4 flux versus heartwood methanogen abundance (mcrA gene copies g^{-1}). **(D)** CH_4 flux versus internal CH_4 concentration at three stem heights (50, 125, 200 cm), with separate panels for each height. Points colored by tree species (legend, phylogenetic order). Axes displayed on pseudo-logarithmic scales (log_{10} for mcrA in panel C) to accommodate wide concentration ranges. Black lines show linear regression fits with 95% confidence intervals (shaded gray). Relationships between internal gas composition, methanogen colonization, and surface emissions suggest that methane production within tree tissues drives observed flux patterns.

1208

1209

Figure S7: Random forest model performance, feature importance, and response functions for tree stem and soil methane fluxes. (a-b) Observed versus predicted CH₄ fluxes showing hexagonal binning to visualize point density. Model performance is evaluated using Lin's concordance correlation coefficient (CCC), which accounts for both precision and accuracy, and coefficient of determination (R^2). Red line indicates 1:1 agreement. Tree model: CCC = 0.329, R^2 = 0.163, n = 453; Soil model: CCC = 0.500, R^2 = 0.292, n = 262. (c-d) Feature importance rankings from random forest models showing the relative contribution of predictor variables to model performance. Tree model includes species identity, species×moisture interactions, and within-species standardized DBH (deconfounded for species identity). Soil model includes environmental drivers and their interactions. (e-f) Partial dependence plots showing the marginal effect of four key predictors on predicted CH₄ flux while averaging over all other variables. Shaded regions represent ±1 standard deviation across the prediction ensemble. Tree predictors: air temperature, soil moisture, soil temperature, and within-species DBH (z-score removes between-species size differences). Soil predictors: air temperature, soil moisture, soil temperature, and seasonal index (residual temporal variation). (g-h) Monthly predictions of mean CH₄ flux from tree stems and soil across the annual cycle, displayed on pseudo-log scale to accommodate the wide range of fluxes and negative values (soil uptake). Predictions integrate species composition, environmental conditions, and seasonal patterns from the random forest models.

1225

1226

1227   **Figure S8:** Photo of representative semi-rigid chamber used 2020-2021.

1228

1229

1230   **Figure S9**: Photo of representative rigid chamber used 2021-2023.

1231

Tissue-Specific Microbiomes of a Single Black Oak

Microbial community analysis of a single felled black oak (*Quercus velutina*) revealed distinct
patterns of methanogen and methanotroph-related taxa across various tree tissues, including
heartwood, sapwood, bark, branches, leaves, roots, and leaf litter. No methanogens were detected
in soil samples from this tree, further supporting the hypothesis that methane emissions in upland
trees are driven by internal methanogenic activity within tree tissues, rather than by soil-to-stem
methane transport.

The only methanogen genus detected in the tree was *Methanobacterium* (family
Methanobacteriaceae), which was found exclusively in heartwood and sapwood. This taxon was
most abundant in heartwood, suggesting that internal anaerobic environments within tree tissues
provide suitable conditions for methanogenesis. No methanogens were detected in oxygen-rich
tissues such as bark, branches, or foliage, indicating that oxygen availability is a key limiting
factor for methanogenic activity within trees.

In contrast, several methanotroph-related taxa were detected across various tree tissues,
particularly in bark, sapwood, and leaf litter. These taxa were classified into three categories
based on their methane-oxidation capabilities: known methanotrophs, potential methanotrophs,
and methanotroph associates.

Known Methanotrophs: *Methylocella* (family Beijerinckiaceae) was detected in heartwood,
sapwood, bark, and foliage. This genus is a facultative methanotroph capable of oxidizing both
methane and multicarbon compounds, indicating that methane oxidation may occur within tree
tissues. The widespread presence of Methylocella across different tissues suggests its potential
role in reducing internal methane concentrations before methane is emitted from tree stems.

Potential Methanotrophs: *Methyloferula* (Beijerinckiaceae) was found in sapwood, bark, and
leaf litter. This genus is considered an obligate methanotroph, utilizing soluble methane
monooxygenase (sMMO) to oxidize methane. Its presence in tree-associated tissues further
supports the possibility of methane oxidation occurring within the tree.

Methylorosula (Beijerinckiaceae) was detected in heartwood, sapwood, and leaf litter. While
closely related to known methanotrophs, its capacity for methane oxidation has not been
conclusively demonstrated. It is thus categorized as a potential methanotroph.

Methanotroph Associates: *Methylobacterium-Methylorubrum* and *Roseiarcus* (family
Beijerinckiaceae) were detected across multiple tissues, including bark, sapwood, roots, and leaf
litter. These genera are pink-pigmented facultative methylotrophs (PPFMs) that are known to
utilize methanol and other single-carbon compounds, but they do not directly oxidize methane.
Their presence in methane-rich environments suggests they may play a role in supporting
methanotrophic activity or participating in co-metabolism with methanotrophs.