1

Resilin, the gene for the molecular spring: its roles in flight and

jumping of *Drosophila*

5   Miyuna Hagiwara[1], Yoshiki Eto[2], Yoko Keira[2], Hiroyuki O. Ishikawa[2],

6   Takaomi Sakai[1], Toshiro Aigaki[1], Tsunaki Asano[1,]*

8   [1] Department of Biological Sciences, Tokyo Metropolitan University
9   [2] Department of Biology, Chiba University
10
*Corresponding Author: asano-tsunaki@tmu.ac.jp, Minami Osawa 1-1, Hachioji city,
Tokyo, Japan
Email: asano-tsunaki@tmu.ac.jp
Tell: +81-426-771-111
Fax: +81-426-772-559

Abstract

Resilin is a member of the chitin-binding protein family that was originally found as a

major component of an elastic matrix present in insects. The knockdown of the gene

for this protein in *Drosophila melanogaster* causes the characteristic downturned wing

posture, and the knockdown flies cannot move their wings. The jump distance of the

knockdown flies is around 50% shorter than that of the controls. Null mutant flies

show the same phenotypes, which can be rescued by introducing a 4.7 kb genomic

fragment harboring the whole coding region of *Resilin*. *In vitro* experiments have

shown that the elasticity of the matrix made of Resilin is due to the

dityrosine-mediated polymerization of Resilin molecules. Dual oxidase (Duox) is the

most likely candidate for the *in vivo* polymerization of Resilin molecules. *Duox*

knockdown induces phenotypes similar to those observed in wing posture and jump

performance of the *Resilin* knockdown and knockout flies, which can be rescued by the

overexpression of *Duox* gene from a beetle. These findings suggest that Duox is an

essential factor for the proper function of Resilin as the major component of the resilin

matrix.

Introduction

Insects are often regarded as the most successful animals on Earth with their extreme

diversity, enormous biomass, and great impact on the global ecosystem. There have

been intensive debates as to why insects have been so successful, and their ability to

fly must have had great impacts on their evolutionary process and the expansion of

their habitat to a wide range of terrestrial environments. It is considered that the ability

to fly is advantageous in the competition for survival and the creation of new niches

not easily accessible to other animals, which is supported by the fact that 99% of the

described insect species are winged.

During powered flight, insects flap their wings by moving their flight muscles in

the thorax. Repetitive and rapid contractions of flight muscles are transformed into

high-frequency wing motion, generating a force to lift the insect's body[1,2]. The hinges

connecting the wings and the body have thick tendons, which may be involved in

energy transformation from the flight muscles to the wings[3,4]. Accumulation of

mechanical loads derived from the high-frequency vibration during flight could

eventually destroy structures around the wing hinges. However, the thick tendon of the

wing hinges is composed of a rubber-like matrix, which may presumably

absorb/relieve such stresses, and prevent the hinges from breaking down[3,4,5]. In

addition to their powered flight, insects may also use the properties of such rubber-like

matrices for jumping. Similarly to flight, jumping is a high-velocity locomotion

through the air, which may be another advantage of insects that is not found in other

invertebrates, with only a few exceptions of some arthropods, such as sand hoppers or

jumping spiders[6,7,8]. It has been reported that the rubber-like matrices are found in the

joints of the hind legs of the locust or in the pleural arch at the top of the hind legs of

the flea. Both of which are positioned in places that can be easily imagined to be used

as a spring providing highly efficient energy kinetics for achieving long-distance and

high-velocity locomotion[7,9,10].

The first description of the rubber-like matrices in insects was the discovery of

tendons in the wing hinges of dragonflies and locusts[4]. At that time, the rubber-like

matrices were named "resilin" after the Latin word "resilire," which means resilience,

and were defined as materials that can be stained with methylene blue or toluidine blue.

After that, rubber-like matrices (hereafter, referred to as the "resilin matrix") were

found in the salivary pump and feeding pump of bags, or the tymbal of cicadas for

sound production[11,12]. The connection between jumping mechanisms and the

localization of the resilin matrix at leg joints has also been discussed in insects such as

locusts that have the thick resilin matrix at the joints of their hind legs[9,13]. Relatively

recent reports on the functions of the resilin matrix include examples from bed bugs

and midges. Male common bedbugs (*Cimex lectularius* and *Cimex hemipterus*)

puncture the intersegmental membrane of the female abdomen with a needle to inject

their sperm. After the male detaches from the female, the pore is likely closed,

probably owing to the elasticity of the resilin matrix[14,15]. Larvae of the phantom midge

(*Chaoborus americanus*) can regulate their buoyancy by altering the volume of their

air sacs. This is achieved through a pH-dependent transition in the shape of the resilin

matrix that composes the air sacs. During this process, membrane ATPase controls the

proton content of the resilin matrix[16].

The physical properties of the resilin matrix are not affected considerably by

deep freezing, high temperature, or treatments with alcohols or fixatives. In contrast,

the resilin matrix is rapidly dissolved by proteolytic digestion, indicating that proteins

are its major constituents[4]. Subsequently, a study showed that proteins in the resilin

matrix are crosslinked together through the formation of di- or tri-tyrosine, in which

tyrosine residues are covalently crosslinked to each other[17]. It was also shown that di-

and tri-tyrosine emit the characteristic blue fluorescence under UV irradiation, and this

property has been used as a useful tool to observe the localization of the resilin matrix

noninvasively (hereafter, di- and tri-tyrosine are collectively referred to as dityrosine).

In 2001, Ardell and Andersen reported a candidate gene encoding a protein composing

the resilin matrix. Partial amino acid sequences of peptides isolated from the protease

digest of the resilin matrix taken from the desert locust *Schistocerca gregaria* were

determined by Edman degradation. The obtained sequences were used for BLAST

search in the genomic database of the fruit fly *Drosophila melanogaster* to find

candidate genes encoding *Drosophila* proteins homologous to the locust proteins[18].

Two proteins (CG15920 and CG9036) were found to have the sequences that matched

well with those obtained from the resilin matrix of the locust. On the basis of similarity

in amino acid compositions (high amounts of proline, glycine, and tyrosine), Ardell &

Andersen concluded that CG15920 was the most likely candidate, and tentatively

named the *CG15920 "Resilin"* as the gene for a major protein composing the resilin

matrix. Since that study, *CG15920* has been called *Resilin* (hereafter, *Rsl*). Until

recently, however, there had been no reports on the gene function analysis to directly

demonstrate that the gene encoding resilin is actually involved in insect flight or

jumping. Furthermore, it was also not known how resilin molecules are polymerized *in*

vivo, although *in vitro* experiments using a recombinant Rsl protein of *D. melanogaster*

have shown that Rsl polymerization is necessary for the high elasticity of synthetic

resilin matrix[5,19-21].

Here, using the newly obtained *Rsl* mutant of *D. melanogaster*, we characterized

the phenotypes and showed that *Rsl* has roles in preventing the breakdown of the wing

hinge structure over time and in efficient jumping. We also tried to show which genes

were involved in the polymerization of the Rsl molecule. Dual oxidase (Duox) is a

membrane protein with the extracellular peroxidase domain and the transmembrane

NADPH-oxidase domain. By tissue-specific *Duox* knockdown and its rescue

experiment, we showed that Duox was indeed the factor for Rsl polymerization. We

also showed that the Duox-mediated polymerization of Rsl was indispensable for the

proper function of the Rsl/resilin matrix both in flight and jumping, which may be the

critical factors for the evolution and success of insects as terrestrial organisms.

Results

Rsl encodes a protein with "elastic repeats"

Rsl is composed of three exons, and two isoforms *Rsl-RA* and *Rsl-RB* (with and

without exon 2, respectively) are expressed (Figure 1A)[22,23]. Rsl is the secreted protein

with a signal sequence (Figure 1B)[18]. The polypeptide encoded by *Rsl-RA* is divided

into three parts: 1) repeats of N-terminally located "elastic modulus", 2) chitin-binding

motif, and 3) repeats of C-terminally located "elastic modulus". The polypeptide

encoded by *Rsl-RB* is devoid of the large portion of the chitin-binding motif (Figure

1B)[18,24].

The elastic moduli[24] (hereinafter referred to as elastic repeats) that are indicated

by light-blue or light-green boxes (Figure 1B) are characterized by the consensus

sequences containing proline, glycine, and tyrosine. The consensus sequences are also

found in other insect proteins that are annotated as resilin[23,24]. The tyrosine residue is

considered important for the formation of intermolecular crosslinks[18]. As first

described by Ardell and Andersen[18], the consensus sequences of the elastic repeats in

the N- and C-terminal regions are different from each other (Figure 1B). In *D.*

melanogaster, the N-terminal repeats (-**P**--Y̲G-**P**G---) have two prolines with an

intervening tyrosine, whereas the C-terminal repeats (GY̲S-GR**P**GG) have only one

tyrosine and one proline. In the multiple sequence alignment[23,24], the N-terminal

repeats are conserved among species, but there are at least two subtypes in the

consensus sequences of the C-terminal repeats. In many insects, the consensus is

GY̲**P**SGG**P**GG with two prolines (type-A repeat). In contrast, many fly species and a

portion of hemipteran insects have different consensus sequences characterized by the

presence of only one proline (Figure 1B). In a comprehensive database search of

arthropod genomes, we found that in amino acid sequences of resilins from the

palaeopteran insects (Odonata and Ephemeroptera: the orders of the most primitive

winged insects) or those from the collembolan (the primitive non-insect hexapods), no

consensus sequences of the C-terminal repeats are found (Figure S1). Considering the

branching pattern of the phylogenetic tree illustrating the evolutionary process of

insects (see Figure S1), it can be hypothesized that the C-terminal repeats are a trait

that was obtained in accordance with the emergence of a new type of winged insects

(Neoptera). In the three-dimensional model of *Drosophila* Rsl (Q9V7U0 in the uniplot

database), which was predicted using AlphaFold Server (https://alphafoldserver.com)

powered by AlphaFold 3[25] (Figure 1C), both the N- and C-terminal regions have a

spring-like "coiled" shape (panels b and c, respectively), with each elastic repeat

corresponding to one cycle of the spring (panels d and e). The presence of the beta turn

at the short PGG sequence motif in the C-terminal repeats (arrowheads in panel c) is

consistent with the previous prediction[18]. Our predictions using AlphaFold 3 showed a

spring-like shape similar to the Rsl model in the resilin of other insects, except for

species such as *Bombyx mori* (Lepidoptera) and *Folsomia candida* (Collembola)

(Figure S2). The spring-like "coiled" shape in the models is intriguing, considering the

function of the resilin as a component of spring-rubber-like biomaterials.

Rsl is expressed in the wing hinge and the leg joint

In previous studies of the two Drosophilidae species[22,26], it was shown that *Rsl* was

mainly expressed in late pupal stages. In our RT-PCR analysis (Figure 2A), the

apparent signals of the isoforms *Rsl-RA* and *Rsl-RB* were detected from the pupal to

early adult stages. From 2 to 4 days after puparium formation (APF), the signals for

Rsl expressions are strong, and the emergence of the signal for the expression of

Rsl-RB precedes that of *Rsl-RA*. In western blotting with the specific antibody, two

signals corresponding to the molecular masses of 57k and 52k can be seen (Figure 2B).

The upper and lower signals correspond to mature Rsl-PA and Rsl-PB without a signal

sequence, respectively (Figure S3). The emergence of the Rsl-PB signal precedes that

of Rsl-PA, which is consistent with the temporal patterns obtained by RT-PCR analysis

(Figure 2A). Rsl signals almost disappear after eclosion, which is consistent with the

observation that resilin molecules are polymerized to become insoluble materials[17]. Rsl

polymer cannot penetrate into the SDS-PAGE gel and are not detectable by western

blot analyses.

To see the spatial patterns of *Rsl* expression, we prepared the *Rsl* promoter-Gal4

driver that covers the genomic region around 2.3 kbp upstream of the first ATG (Figure

1A). The expression was visualized using the *UAS-Rsl-RA::mCherry* reporter. The

photos in Figure 2C show pupae on day 3 APF. The mCherry signal was visible at the

hinges of the wings, spiracles, compound eyes, and labrum (Figure 2C, panels a, b, and

c). In the observation of the adult on day 3 after eclosion, the mCherry signal can be

seen on the ventral side of the leg joints between the trochanter and the femur, and the

roots of the mechanosensory hairs of the leg and wing (Figures 2D and 2E). These

expression patterns were obtained using flies carrying the *Rsl*-promoter Gal4 sequence

integrated into the 3rd chromosome. The mCherry signal pattern obtained with the

Rsl-promoter Gal4 sequence on the second chromosome was nearly identical to the

data obtained with *Rsl*-promoter Gal4 on the third chromosome, except that the signal

in the compound eyes was very weak (Figure S4). In western blot analysis of the

compound eye (Figure 2F), the signals of Rsl can be seen in the wild-type fly, and this

signal disappeared in the sample from Rsl^{KO} that we generated (Figures 1A and S5).

The results of the western blot analysis indicate that the signal in the wild-type fly

corresponds to Rsl, and that the compound eye surely expresses *Rsl*.

To observe the details of the internal structure at the wing hinge, we removed the

outer cuticle. The mCherry signal was clearly visible just beneath the wing hinge

(Figure 2G, panel i), and the structure exhibited blue fluorescence under UV irradiation

(Figure 2H). The intensity of this blue fluorescence signal decreased markedly and

increased under acidic and alkaline conditions, respectively (see Figure S6), consistent

with the characteristics of dityrosine previously reported[17]. This structure, which is

positive for both the mCherry signal and blue fluorescence, is the tergopleural tendon.

Previous studies have shown that this tendon is the structure connecting the direct

flight muscle and the root of the wing at the wing hinge[27-29].

Since Rsl-PA has the chitin-binding domain (Figure 1B), we consider that this

domain may function in chitin binding. To visualize the distribution of the chitin fiber

inside the tergopleural tendon, we expressed ChtVis-tdTomato with *Rsl*-promoter-Gal4.

A strong signal can be seen along the horizontal midline, indicating the presence of

chitin along the midline (Figure 2I). This is similar to the spatial pattern of

Rsl-RA::mCherry (Figure 2H). Next, we compared the distributions of

Rsl-RA::mCherry and Rsl-RB::mCherry (with and without the sequence of the

chitin-binding domain, respectively) that were separately expressed (Figure 2J). The

spatial pattern of Rsl-RA::mCherry, which is denser along the horizontal central line

(panel p), contrasts with the more uniform distribution of Rsl-RB::mCherry (panel q).

In the calculation for the biased distribution of mCherry signals (see Materials and

Methods), the value was higher in flies expressing Rsl-RA::mCherry than in those

expressing Rsl-RB::mCherry. This indicates a more uniform distribution of

Rsl-RB::mCherry, which may be caused by the absence of a chitin-binding domain.

Rsl may work to prevent the breakdown of a tendon in the wing hinge that is

required for sustainable flight

To examine the effects of the suppression of _Rsl_ expression, we knocked down _Rsl_[30]

with the _Rsl_-promoter Gal4 driver, and the expression was suppressed to around the

baseline levels (Figure S7). In the _Rsl_ knockdown flies, no visible phenotypes were

observed in the very early stages of the adult, but after one day, all the knockdown

individuals were unable to move their wings and started to display the characteristic

downturned wing posture (Figure 3A). The same wing posture has been observed in

multiple studies including the analysis of _pink1_, which functions in muscle

development[31,32]. The phenotypes observed in our _Rsl_ knockdown are the same as

those previously reported for the _Rsl_ knockdown flies[21]. To further characterize the

function of _Rsl_, we used the null mutant (Rsl^{KO}) that was prepared in this study (Figure

S5). The homozygous Rsl^{KO} flies showed the characteristic downturned wing posture,

and they could not move their wings, which are the same phenotypes as those observed

in *Rsl* knockdown flies (Figure 3C, see Extended data Video 2). To rescue these

phenotypes, we generated flies with the transgene of the 4.7 kbp genomic fragment

harboring the *Rsl* coding region on the 3rd chromosome (*Rsl*-Genomic rescue: *Rsl-GR*)

(Figure 1A). The *Rsl*^{KO} flies with *Rsl-GR* did not show the downturned wing posture

(Figures 3C and 3D), and they could move their wings and fly (see Supplementary

mov. S3), indicating that *Rsl-GR* rescued the phenotypes of *Rsl*^{KO}. Then, we also tried

to rescue the phenotypes of *Rsl*^{KO} using the Gal4/UAS system (Figure 3D). Although

flies expressing each isoform only gradually showed the wing posture phenotype, the

phenotype was almost rescued. Combining the two isoforms did not show markedly

different results with those obtained from either *Rsl-RA* or *Rsl-RB* alone.

It was shown in *D. melanogaster* that the tergopleural tendon was stained with

the anti-Rsl antibody[28], which is consistent with the presence of the signal for the

Rsl::mCherry reporter at the tergopleural tendon (Figures 2G and 2H). In Figure 3E,

the structure around the tergopleural tendon of the wild-type fly was illustrated

according to the previous descriptions [28,29]. The proximal side of the tergopleural

tendon is attached to the inner side of the cuticle located at the wing hinge (indicated

by the yellow arrowhead), and the direct flight muscles (tergopleural muscles 1 and 2:

TP-1 and TP-2, respectively) are attached to both the tergopleural tendon and the

cuticle of the body wall. In observation of the inner structure of wing hinges,

dumpy-YFP (*dpy-YFP*) was used to visualize the boundary between the tendon and the

muscle (Figure 3F, panel a). The green signal of Dpy-YFP surrounds the tergopleural

tendon emitting the blue fluorescence upon UV irradiation (Figure 3F, panel a). Like

the schematic figure (Figure 3E), the direct flight muscles are attached to the

tergopleural tendon that is visualized with Dpy-YFP. As shown in panel b of Figure 3F,

in the newly emerged Rsl^{KO} fly, no blue fluorescence signal of dityrosine was seen.

The loss of dityrosine signals can be explained by the absence of Rsl, which is

supposed to provide the tyrosine residues as the source of dityrosine. The tergopleural

tendon of the newly emerged Rsl^{KO} fly remained linked to the root of the wing (white

arrowheads), which may correlate with the absence of wing posture or movement

abnormalities (Figure 3D). In contrast, in Rsl^{KO} flies exhibiting the downturned wing

posture (panel c), the tergopleural tendon was ruptured into two pieces (white

arrowheads), confirming that the connection between the muscle and the root of the

wing was almost lost. In contrast, in Rsl^{KO} flies with Rsl-GR (panel d), the blue

fluorescence that is surrounded by the Dpy-YFP signal is clearly visible. The structures

of TP-1 and TP-2 are the same as those observed in wild-type flies, and no damage

was observed. This is consistent with the Rsl^{KO} flies with Rsl-GR showing no

phenotypes of wing posture and movement abnormalities (Figures 3C and 3D;

Extended data Video 3).

In the fields of anatomy and biomechanics, many reports define "resilin" as an

extracellular structure that exhibits blue fluorescence, which is characteristic of

dityrosine[33]. The blue fluorescence signal has been found in various body parts,

including the intersegmental membrane, compound eyes, and wings[34,35]. In this study,

to characterize the nature of the blue fluorescence in detail, we focused on the wings,

since the presence of the blue fluorescence signal in wings has been reported in

multiple insect species[34,35]. In the fly with Rsl-RA::$mCherry$ expression that was driven

by Rsl-$Gal4$, (Figure S8, panels, a and b), strong mCherry signal can be seen in several

places including the tergopleural tendon. In contrast, there is almost no signal at the

wing, although with UV, blue fluorescence can be seen at the proximal part of the

wings (panel c). This finding suggests that *Rsl* is not expressed even in the area

positive for blue fluorescence. In the magnified photo of the wild type fly's wing

(Figure S8, panel e), the arrows indicate distinct blue fluorescence signals at the

proximal part of the wing, roughly corresponding to the area with landmarks adopted

in a previous study (panel d)[36]. These signals became weaker drastically under acidic

conditions (panel f), suggesting that they corresponded to dityrosine. In the wing of the

Rsl[KO] fly (panel g), however, the signals were still seen, indicating that not Rsl but

other tyrosine-rich substances are the cause of the blue fluorescence signal. These

findings support the idea that dityrosine-rich structures exist without *Rsl* in *D.*

melanogaster.

Rsl is polymerized by the enzyme for ROS production

It has been shown that Duox is involved in dityrosine formation in ecdysozoans such

as nematodes and insects[37-39]. The ROSs produced in the peroxidase domain may

oxidize a tyrosine residue to form a tyrosine phenoxyl radical, followed by the reaction

with the neighboring tyrosine residue to produce dityrosine or trityrosine[37,38]. Duox

may be the most plausible candidate factor responsible for Rsl polymerization, because

Rsl molecules are polymerized through dityrosine formation[17]. To demonstrate this, we

examined the effect of *Duox* knockdown on Rsl polymerization or other phenotypic

characteristics. In this study, we performed tissue-specific *Duox* knockdown with

Rsl-Gal4, because whole body knockdown or knockout of *Duox* leads to lethality[37,40].

We used two UAS-lines, NIG stock #3131R-3 (*NIG3131R-3*) and the previously

reported *UAS-Duox-RNAi$_{976-1145}$*[40], for *Duox* knockdown. In both cases, immediately

after eclosion, *Duox* knockdown flies were able to fold and move their wings normally,

but within a day, all of them exhibited the downturned wing posture, similar to *Rsl*

knockdown and knockout (Figures 4A and 4B). After emergence of this wing posture,

both the *Duox* knockdown flies could not move their wings, similarly to *Rsl*

knockdown and knockout flies (hereafter, we mainly performed phenotypic

characterization using *NIG3131R-3*).

To examine the cause of the downturned wing posture in *Duox* knockdown flies,

we performed western blot analysis to determine whether Rsl molecules were

polymerized. As shown in Figure 4C, in the *Duox* knockdown flies, the signal for Rsl

can be seen at the same electrophoretic mobility of Rsl monomers (Figure 4C, right

panel), suggesting that a portion of Rsl molecules remained unpolymerized. This is in

contrast to the patterns of the control (Figure 4C, left panel) and wild-type flies (Figure

2B), in which the Rsl signals disappeared after eclosion. Another explanation for the

detection of *Rsl* in the *Duox* knockdown flies could be the enhancement of *Rsl*

expression after becoming adults. However, there were no marked differences in the

patterns of *Rsl* expression between the control and the *Duox* knockdown flies (Figure

S9), which does not negate the model of the Duox-mediated Rsl polymerization

(Figure 4D).

We also observed the structure around the tergopleural tendon of the *Duox*

knockdown flies (Figure 4E). In the control flies (panels a and b), the structures were

almost the same as those of the wild type (Figure 3F), as can be judged from the

presence of the blue fluorescence signal at the tergopleural tendon. In the *Duox*

knockdown flies, the signal for dityrosine in the tergopleural tendon (indicated by the

dotted line in panel c) was very weak. In the *Duox* knockdown flies, the TP-1 and TP-2

were completely detached from the tergopleural tendon with the signal of Dpy (panel

d), as illustrated in panel e.

To rescue the phenotypes caused by *Duox* knockdown, we prepared a UAS-line

for expressing of *T. castaneum Duox* (*TcDuox*) as a transgene with a base sequence that

is not similar to that of *D. melanogaster Duox*. As shown in Figure 4F, *Duox*

knockdown flies with additional *TcDuox* expression did not exhibit the downturned

wing posture, even one week after eclosion (Figures 4F and 4G). The flies could move

their wings normally and fly (our observation). In western blot analysis, the Rsl signal

disappeared within two days after eclosion (Figure 4H, right panel), in contrast to the

control flies in which the Rsl signal remained detectable even after two days (left

panel). The anatomical structure around the tergopleural tendon was almost the same

as that of the wild type (Figure 4I, right panel), which can be judged from the presence

of the blue fluorescence signal in the tergopleural tendon, and also the connection

between the tergopleural tendon and TP muscles. These findings indicate that the

expression of *TcDuox* rescued the phenotypes of the *Duox* knockdown flies, and that

Duox is the actual factor responsible for Rsl polymerization and the proper functioning

of Rsl/resilin matrix in the maintenance of the structure around the wing hinges. We

then examined biochemically the Duox-mediated Rsl polymerization through

dityrosine formation. Here, commercially available peroxidase was used as the catalyst,

instead of the peroxidase domain of Duox, because we were unable to synthesize the

functional peroxidase domain of Duox successfully. In the western blot analysis of the

truncated version of the recombinant Rsl (reRsl), which was synthesized in an *E. coli*

system, reRsl migrated at an electrophoretic mobility corresponding to the predicted

molecular mass of 35 k (Figure 4J, left panel, lane a). After treatment with either

peroxidase or hydrogen peroxide (lanes b and c), reRsl can be seen at its original

mobility. In contrast, in the presence of both peroxidase and hydrogen peroxide, the

reRsl signal was shifted to an electrophoretic mobility corresponding to a much higher

molecular mass (lane d). As indicated by closed arrowheads, the strong smear signal

was seen from the entry site of the concentration gel. The shift of the Rsl signal to an

electrophoretic mobility corresponding to high-molecular masses indicates that the Rsl

molecules formed a high-molecular mass complex. In the western blot analysis with

the antibody against dityrosine (right panel), signals are visible only in the sample

treated with both peroxidase and hydrogen peroxide (lane h). The signal is strongest at

the entry site of the concentration gel (closed arrowhead), indicating the presence of

reRsl, which is highly polymerized by the formation of a large number of dityrosines.

Both the expressions of *Rsl* and *Duox* are required for high jump performance

It has been proposed that the resilin matrix may enable high performance in jumping in

several insects, such as the flea and locust[10, 21,41,42]. In this study, as a method to

quantify the jump performance in *D. melanogaster*, we measured the jump distance

using de-winged flies (Figure 5A). In the control flies, their jump distance was around

2 cm, but in the *Rsl* knockdown flies, their jump distance was shorter by around half of

that of the control (Figure 5B). Also, in Rsl^{KO} flies, the jump distance was almost half

of that of the wild-type flies (Figure 5C, Extended data Video 4 and 5). The jump

distance of the Rsl^{KO} flies with *Rsl-GR* was almost the same as that of the wild-type

flies (Figure 5C), indicating that the phenotype of shorter jump distance was rescued

by *Rsl-GR*. In the rescue experiment using the Gal4/UAS system, we were unable to

obtain data clearly showing the successful rescue with either *UAS-Rsl-RA* or

UAS-Rsl-RB. Therefore, presently, we do not have information on the functional

differences in jumping between Rsl-PA and Rsl-PB (not shown).

Here, in addition to *Rsl*, we examined the contribution of *Duox* to fly jumping.

Duox was knocked down with either of *NIG3131R-3* or *UAS-Duox-RNAi$_{976-1145}$*, and in

both cases, the jump distances of the knockdown flies were around half of that of the

controls (Figure 5D). To rescue this phenotype, *TcDuox* was additionally expressed in

the *Duox* knockdown flies (*Rsl>NIG3131R-3*). With the expression of *TcDuox*, the

jump distance was increased to around twofold that of the *Duox* knockdown flies

(Figure 5E), indicating that *TcDuox* expression compensated for the reduction in the

level of the endogenous *Duox* expression.

Discussion

Duox is the oxidase that is widely conserved in metazoans, and is known as the thyroid

oxidase in mammals[43,44]. There have been many studies on the diverse roles of Duox,

including defense responses against intestinal wounds, immune responses against

pathogens, the regulation of stem cell development/differentiation, and extracellular

matrix formation[37-40,45-48]. In particular, in ecdysozoans such as insects and nematodes,

it is considered that the Duox-mediated crosslinking through dityrosine formation is

very important for the strengthening and stabilization of the extracellular matrix[37-39,49].

In this study, we showed that Rsl was polymerized by Duox, in which inter-Rsl

molecule dityrosine formation may occur (Figures 4C, 4D, 4H, and 4J). In insects, this

is the first study to identify the target protein of the Duox-mediated polymerization.

The polymerization of Rsl may be used as a model to study the formation of the

dityrosine matrix, which is distributed widely in metazoan taxa, including annelids and

molluscs[50,51].

Until this study, two reports on the function of the resilin-encoding gene in

Drosophilae species had been published, both by Moussian's group. Here, we were

able to specify the internal structures related to the emergence of this phenotype

(Figure 3F). As shown in Figure 3F, the tergopleural tendon was ruptured in Rsl^{KO} flies.

The direct flight muscles (TP-1 and TP-2) that are linked to the tergopleural tendon are

required for the movement of the wings that are upturned, as has been demonstrated in

a neurogenetic experiment[52]. Probably, in the Rsl knockdown and Rsl^{KO} flies, the

tergopleural tendon may have been damaged owing to the accumulation of stress from

the wing movement, since the occurrence of this phenotype is time-dependent (Figures

3B and 3D). After the rupture of the tergopleural tendon, energy transfer from the

muscle to the wings may no longer occur.

In the insect database of the full-length CDS or long-read RNA-seq results, we

found two splicing isoforms of resilins with or without the chitin-binding domain (our

observation), which indicates the possibility that the presence of these two isoforms is

important. The localization of Rsl-PA::mCherry (with the chitin-binding domain) in the

tergopleural tendon (Figure 2H) is actually similar to that observed in the distribution

of chitin fibers that were visualized using the ChtVis-tdTomato chitin reporter (Figure

2I). This suggests that the chitin-binding domain of Rsl-PA is functional *in vivo*, which

is consistent with the finding of an *in vitro* experiment using recombinant Rsl, showing

the binding of the chitin-binding domain to chitin[23]. The binding of the Rsl molecule to

chitin fibers seems to have a certain significance for the formation and proper

functions of the tergopleural tendon. However, when we compared the extent of

phenotypic rescue by overexpressions of *Rsl-RA* and *Rsl-RB* in *Rsl^{KO}* flies, the

downturned wing phenotypes were rescued, in almost the same proportion (Figure 3D).

This suggests that the chitin-binding domain is not necessarily required for Rsl

functions at the wing hinge. In the wing hinge, it is possible that other cuticle proteins

with both tyrosine residues and the chitin-binding domain may compensate for the

absence of the chitin-binding ability of Rsl-PB by mediating the linkage between chitin

fibers and Rsl-PB.

The polymerization of elastic protein moieties often occurs to give the matrix

high elasticity, as seen in other elastic materials, such as elastin and abductin[53-55]. *In*

vitro experiments designed to polymerize recombinant Rsls support this theory[5,20,53]. In

these studies, either peroxidase, photocatalysts, or chemical modifiers were used to

artificially polymerize Rsl, and the resulting materials treated with these reagents

exhibited high elasticity. The representative molecule forming the elastic matrix in

mammals is elastin, which is referred to as the "molecular spring" because of its roles

in restoring the structure and providing elasticity to tissues[53-55]. The tyrosine residues

of Rsl may form intermolecular crosslinks, which may be responsible for the formation

of the resilin matrix and its high resilience. According to the UniProt website

(https://www.uniprot.org/uniprotkb/Q9V7U0/entry), the chitin-binding domain of the

resilin model has a pLDDT score greater than 90, indicating high precision in the 3D

model prediction. However, the scores for the N- and C-terminal repeats are estimated

to be low (pLDDT score <70), which indicates that these regions could be intrinsically

disordered proteins. This is consistent with previous predictions that the Rsl molecule

has a random structure, lacking alpha helices or beta sheets[23,56]. However, low scores

in AlphaFold's predictions may simply reflect insufficient information about proteins

possessing similar sequences that could serve as references for known protein

structures during 3D modeling. At present, there is insufficient evidence to rule out the

possibility that Rsl possesses a structure similar to that of the predicted model. To

accurately understand the thermodynamic mechanism by which polymerized Rsl

molecules exhibit high elasticity, it would be beneficial to confirm whether Rsl

actually possesses a coil-like structure.

As shown in Figure 4A, the knockdown of *Duox* induced the same downturned

wing posture as that of the *Rsl* mutant. This result can be explained by the failure of

Rsl polymerization, as shown in the western blot in Figure 4C. Even in the adult stage,

Rsl polypeptides remained detectable at an electrophoretic mobility corresponding to

those of the Rsl monomers. *Duox* knockdown also induced the reduction in the

intensity of dityrosine signal in the tergopleural tendon (Figure 4I), strongly supporting

the idea that Duox is the factor for dityrosine-mediated Rsl polymerization. The

reduction or loss of the elasticity of the resilin matrix at the tergopleural tendon may

have occurred, finally leading to the rupture of the tergopleural tendon. It is also

possible that the link between the tergopleural tendon and the muscle is lost, probably

because of the weak bonding between them, as can be assumed from the detachment of

the muscle from the tergopleural tendon (Figure 4E).

The model that Duox is the factor for Rsl polymerization was based on the

studies of the nematode *Caenorhabditis elegans*, showing that the gene for Duox is

involved in the dityrosine-mediated polymerization of collagens in the cuticle

matrix[38,49]. It is possible that this process may be controlled by humoral factors,

because it is stage-dependent (Figure 2B). It has been shown that the functions of

Duox are regulated by the EF-hand motif present at the intracellular part of this

protein[40,45]. We observed an increase in the cytosolic calcium level in the cells

surrounding the tergopleural tendon in the late pupal stages (our observation). The next

step is to clarify whether the increase in the cytoplasmic calcium level actually

contributes to the activation of Duox and the subsequent dityrosine formation,

followed by elucidation of the molecular mechanisms underlying the stage-dependent

regulation of Rsl polymerization during the process of adult development.

Regarding jumping, both *Rsl^{KO}* and *Duox* knockdown flies showed shortened

jump distances, suggesting the loss of the polymerized Rsl may have been the cause of

this phenotype (Figure 5). In a recent study on the resilin gene of the locust *S. gregaria*,

the authors showed the lower jump performance of RNAi individuals (reduced jump

velocity and irreversible reduction in the jump distance during repetitive jumps)[57]. It

was also shown that the joints of the hind legs (between the femur and the tibia) were

more easily broken in the RNAi individuals. In *D. melanogaster*, the resilin matrix is

present on the ventral side of the leg joint (Figure 2); from the finding, we proposed a

simple model on how the resilin matrix works in fly jumping (Figure S10). Before

jumping, the resilin matrix becomes tense to store mechanical energy, and when the fly

jumps, the stored energy may be liberated from the resilin matrix during transition to

its relaxed state.

As shown in the observation of the Rsl^{KO} flies, we found the wing structures still

showing blue fluorescence (Figure S8). The lack of Rsl is also supported by the

previous observation that there are no reporter signals at the proximal part of the

wing[21]. In many reports, the term 'resilin' has been used to refer to the cuticular or

other extracellular structures with blue fluorescence. However, the data from Rsl^{KO}

flies show the existence of the non-resilin matrix with dityrosine, indicating that the

matrices positive for blue fluorescence do not necessarily contain Rsl molecules. It is

possible that such the non-resilin matrix with blue fluorescence may have mechanical

properties quite different from those of the resilin matrix, because blue fluorescence

can be found in very hard parts such as the tip of the mandible of copepods or

ragworm[58,59]. Future studies will reveal the mechanical characteristics or composition

of the non-resilin matrix with blue fluorescence and the biological significance of their

existence.

Materials and Methods

Fly stocks and maintenance

Fly stocks were reared at 25°C under a 12h light and dark cycle. They were fed the

standard cornmeal‑agar‑glucose medium containing propionic acid and n-butyl

p-hydroxybenzoate (7.21% cornmeal, 3.20% glucose, 8.01% yeast, and 0.64% agar

supplemented with 9.9 mL/L butyl *p*-hydroxybenzoate and 3.0 mL/L propionic acid) .

w^1118 (our laboratory stock, Canton-S background) was used as the wild-type strain.

UAS-eGFP (BL#1521), *UAS-GFP-RNAi* (BL#9331), *vas-int* (BL#40161), AttP2

(BL#36303), AttP5 (BL#34766), and *vas-Cas9* (BL#51323) were obtained from the

Bloomington Drosophila Stock Center. *UAS-resilin-RNAi* (*KK106773*) in VIE260b

(#101068) was obtained from the Vienna Drosophila Resource Center. *UAS-Duox-RNAi*

(#3131R-3) and *y²cho²v¹; Sco/CyO* (#TBX-0007) were obtained from Fly Stocks of the

National Institute of Genetics. *UAS-Duox-RNAi$_{976-1145}$*[40] was kindly provided by Dr.

Won-Jae Lee (Seoul National University, Korea). All mutants and lines that were used

for knockdown were crossed into the *w^1118* control background for at least five

generations.

RNA isolation and reverse-transcription PCR

Total RNA was extracted from individuals at each growth stage using RNAiso Plus

(D9108A, Takara, Japan). cDNA was synthesized from 1 μg of total RNA using Prime

Script RT Reagent Kit with gDNA Eraser (DRR047A, Takara). cDNAs were amplified

using Prime star Max (R045A, Takara). To obtain patterns of stage-specific gene

expressions, a three-step thermal cycling profile was used as follows: 95 °C for 1 min,

followed by 25 cycles at 98 °C for 10 s, 55 °C for 5 s, and 72 °C for 5 s. The primers

used in the RT-PCR are listed in Supplementary Table 1.

Generation of fly strains

Rsl^{KO} was obtained by the CRSPR/Cas9-mediated deletion of the whole coding region

with two gRNAs[60]. The gRNAs were expressed from a transgene with a ubiquitous *pol*

III promoter derived from *Drosophila snRNA:U6:96Ab*[61]. For construction of DNA to

generate flies expressing gRNAs, two primer sets for *Rsl-gRNA1* and *-gRNA2* (listed in

Supplementary Table 2) were annealed, and the resulting DNA fragments were

separately subcloned into *pBFv-U6.2* and *pBFv-U6.2B*, respectively, that were

predigested with *Bbs*I. *pBFv-U6.2* with the sequence of *Rsl-gRNA1* was digested with

*EcoR*I and *Not*I, and one of the resulting DNA fragments (with the sequence of

Rsl-gRNA1) was ligated into one of the *EcoR*I and *Not*I fragment of *pBFv-U6.2B* (with

the sequence of *Rsl-gRNA2*). The resulting vector with both the sequences for

expressing *Rsl-gRNA1* and *-gRNA2* was introduced into AttP2 flies using a phiC31

integration system[60,61]. The flies with the transgene were crossed with *vas-Cas9* flies

and the offspring were crossed with $y^2cho^2v^1$; *Sco/CyO* virgins to establish lines with

deletions. The genotypes of the established lines were checked by genomic PCR

analysis using the primers listed in Supplementary Table 2.

For the construction of the plasmid vector for generating transgenic flies

(*Rsl-Gal4, Rsl* genomic rescue (*Rsl*-GR)*, UAS-Rsl-RA*, and *UAS-Rsl-RB*), genomic

DNA and cDNA obtained from w^{1118} were used as templates of PCR. For the subcloning

of the PCR fragments, an In-Fusion HD cloning kit (Z9648N, Takara) was used. To

construct the vector of *Rsl-Gal4*, the 2.3 kbp PCR fragment for the *Rsl* promoter region

amplified using the primer set (Supplementary Table 2) was subcloned into *pBPGw*

(Addgene; https://www.addgene.org/) that was predigested with *Kpn*I and *Bgl*II. To

generate flies with the transgene for the genomic rescue of Rsl^{KO}, we prepared a 4.7 kb

genomic fragment containing the *Rsl* CDS (from 1.0 kb upstream to 0.5 kb downstream

of the transcription start and stop sites of *Rsl*) which was amplified by PCR using the

primer set (Supplementary Table 2). The DNA fragment was subcloned into *pBFv-U6.2*

predigested with *EcoR*I and *Not*I. For the construction of the plasmid vector to generate

UAS-Rsl-RA and *UAS-Rsl-RB*, PCR fragments corresponding to the whole coding

regions of *Rsl-RA* and *Rsl -RB,* respectively, were amplified using the primer set

(Supplementary Table 2) with the cDNA. They were subcloned into *pUAST-attB*

(laboratory stock) predigested with *Not*I and *Xba*I. For the construction of the vector for

UAS-TcDuox, the DNA fragment corresponding to the whole coding region of the duox

gene of the red flour beetle *Tribolium castaneum* was amplified by PCR using the *T.*

castaneum cDNA and the primer set (Supplementary Table 2) (*T. castaneum* was a kind

gift from Dr. Toru Togawa, Nihon University, Japan). For the construction of

UAS-Rsl-RA::mCherry and *UAS-Rsl-RB::mCherry*, the whole coding region of each

isoform was amplified by PCR using the primer set (Supplementary Table 2) with

cDNA from iso-1 (BL2057) as the template. A fragment including the sequence of

mCherry with an N-terminal flexible 4 × GlyGlySer linker was PCR-amplified from

UASp-Elyssim-mCherry[62] using the primer set (Supplementary Table 2). The two DNA

fragments were subcloned into *EcoR*I fragment of *pUAST-attB* by the SLiCE method

(Motohashi, 2015). Each vector was introduced into flies with AttP5 (*UAS-Rsl-RA,*

UAS-Rsl -RB, and *UAS-Tcduox*) or AttP2 (*Rsl-Gal4*, *Rsl-GR, UAS-Rsl-RA::mCherry*,

and *UAS-Rsl-RB::mCherry*) using a phiC31-based integration system[63,64].

Generation of antibody

A part of Rsl (Lys478-Gly590) was chosen as the antigen to raise the antibody against

Rsl. In this study, pET-32 (Novagen, USA) was used to express the Rsl antigen as a

fusion protein with the N-terminally located thioredoxin. The DNA fragment

corresponding to the antigen was amplified with the primer set (Supplementary Table 1)

using the cDNA of *w*[1118]. The PCR product was digested with *Xho*I and *Bgl*II, and

ligated into pET-32 predigested with the same enzyme set. The vector constructed was

used to transform the *E. coli* BL21 strain. After the IPTG induction of the recombinant

protein expression (1 mM, 4 h of culture at 37°C）, the *E. coli* cells were harvested by

centrifugation, and ruptured by sonication in 50 mM Tris-HCl (pH 7.5) containing 150

mM sodium chloride. From the extract, the recombinant protein was purified by

Ni-NTA affinity chromatography using Ni-NTA Agarose (30210, Qiagen, Germany).

The N-terminally located thioredoxin was removed by thrombin treatment () to break

the peptide bond between the thioredoxin and the Rsl antigen, and the Rsl antigen was

used for the immunization of a rabbit. The serum obtained from the rabbit was used as

the anti-Rsl antibody.

Western blot analysis of fly samples

Five individuals at different stages (white pupae, pupae, and adults) were homogenized

in 400 µL of 2×SDS-PAGE sample buffer (12.5 mM Tris at pH 6.8, 20% glycerol, 4%

SDS, 2% 2-mercaptoethanol, and 0.001% bromophenol blue) and boiled for 10 min at

100°C. The proteins in a 5 µL aliquot were separated by SDS-PAGE using 12.5%

polyacrylamide gel and transferred to PVDF membranes (IPFL00010, Merck KGaA,

Germany). After blocking using 5% skim milk with TBST (0.1% Tween20), the

membranes were incubated with the rabbit anti-Rsl antibody (1: 3000 dilution) and

mouse anti-α-Tubulin antibody [DM1A] (ab7291, Abcam, UK) (1: 5000 dilution) in

Can Get signal solution 1 (NKB-201, Toyobo, Japan) at 4°C overnight. For the

secondary antibody, IRDye-conjugated anti-rabbit and anti-mouse IgGs (925-68071 and

925-32210, respectively, LI-COR, USA) were used with Can Get signal solution 2

(NKB-301, TOYOBO, Japan) (1: 20000 dilution, for 2 h at 25°C）. For luminescence

imaging, an Odyssey DLx imaging system (LI-COR, USA) was used.

Measurement of jump distance

One day before the experiments, the flies were de-winged under mild CO_2 anesthesia.

Jump distance was measured using the equipment, which was set in a room kept as

25°C. At the bottom of the glass beaker, 2% agarose was placed. After hardened, the

agarose solution, a mesh for scaling was put on the agarose gel, and the surface was

covered with additional 2% agarose. Five flies per group were housed in the equipment

and their jumping were tracked for 5 minutes at 480 fps with Tough TG-5

(4545350-051112, OLYMPUS, Japan) and the jump distances were measured on the

basis of intervals of the mesh. The longest jump distance in each group was used as the

representative value. The experiments were performed at least eight times for each

genotype and sex.

613   Crosslinking of Rsl

Horseradish peroxidase was used for the crosslinking reaction of recombinant Rsl[21,65].

100 µg of the recombinant Rsl (CSB-EP892693DLU, WUHAN HUAMEI BIOTECH,

USA) was dissolved with 100 µl of 50 mM potassium phosphate buffer (pH7.0).

Horseradish peroxidase (P-8375, Sigma, USA) was dissolved with 0.1 M potassium

phosphate buffer (pH 6.0) to a final concentration of 20 µg/µL. 1 µL of the recombinant

Rsl solution was mixed with 5 µL of 50 mM potassium phosphate buffer (pH 7.0). Then,

1 µL of horseradish peroxidase solution and 3 µL of 0.3% hydrogen peroxide (18084-00,

KANTO CHEMICAL, Japan) were added, and the mixture was incubated at 30°C for 1

h. After incubation, the reaction mixture was mixed with 10 µL 2×SDS-sample buffer

and boiled for 10 min at 100°C. Western blot analyses of Rsl and dityrosine were

performed using the rabbit anti-Rsl antibody (this study) and the mouse monoclonal

anti-dityrosine antibody (MDT-020P, JaICA, Japan). The experimental procedures of

western blot analysis were the same as those described in section 2.5.

Histology and microscopy

To observe the internal structure of the wing hinge, the adult thorax was divided into

two parts (left and right) along the midline. Then the body parts around the wing hinge

were obtained by trimming with microscissors. The tissues were fixed using 4%

paraformaldehyde with PBS (pH 7.0). After blocking using 5% BSA (A7906, Merck

KGaA) with PBST (0.05% Tween20), these samples were incubated in

phaloidin-iFluor488 (CAY20549, Cayman Chemical, USA) with PBST (1:500) at 25°C

overnight. An OLYMPUS BX60 (OLYMPUS, Japan) equipped with an

WRAYCAM-VEX830 (WRAYMER, Japan) was used for whole-body or dityrosine,

mCherry, and YFP signal detection. An OLYMPUS SZX7 (OLYMPUS, Japan)

equipped with an OLYMPUS DP71 (OLYMPUS, Japan) was used for observing the

phenotypes of wing posture.

Quantification of biased Rsl::mCherry distribution

The standard deviation (std) and the mean intensity of the mCherry signal in the resilin

matrix were quantified using ImageJ. The std/mean was used as the index of the biased

Rsl::mCherry distribution[66].

Statistical analyses

Statistical analyses were performed using R version 3.6.0 (https://www.r-project.org/).

The Wilcoxon rank sum test was used to compare the jump distances. The student t-test

was used to compare the intensities of mCherry signals in the resilin matrix in the wing

hinges. The Steel‑Dwass test was used to compare the expression levels of *Rsl*, *Duox,*

and *TcDuox*.

Acknowledgements

UAS-Duox-RNAi$_{976–1145}$ was a kind gift of Dr. Won-Jae Lee (Seoul National University,
Korea). *T. castaneum* was a kind gift from Dr. Toru Togawa (Nihon University, Japan).
We are grateful to Mr. Kensuke Aizawa for his contributions to our preliminary
experiments in the early period. We thank the National Institute of Genetics, the
University of Indiana. Bloomington *Drosophila* Stock Center and the Vienna
Drosophila Resource Center for providing fly lines. This work is supported by JSPS
KAKENHI Grant Number JP23H02225 and 21J21904 and the Sasakawa Scientific

661 Research Grant from The Japan Science Society.

663 **Author contributions**

670 **Competing interests**

References

1.   Ellington, C. P. The aerodynamics of hovering insect flight. VI. Lift and power requirements. *Philos. Trans. R. Soc. London. B, Biol. Sci.* **305**, 145–181 (1984).

2.   Ellington, C. P. Power and efficiency of insect flight muscle. *J. Exp. Biol.* **VOL. 115**, (1985).

3.   Andersen, S. O. & Weis-Fogh, T. Resilin. A Rubberlike Protein in Arthropod Cuticle. *Adv. In Insect Phys.* **2**, 1–65 (1964).

4.   Weis-Fogh, T. A Rubber-Like Protein in Insect Cuticle. *J. Exp. Biol.* **37**, (1960).

5.   Elvin, C. M. *et al.* Synthesis and properties of crosslinked recombinant pro-resilin. *Nature* **437**, 999–1002 (2005).

6.   Parry, D. A. & Brown, R. H. J. The Hydraulic Mechanism of the Spider Leg. *J. Exp. Biol.* **36**, 423–433 (1959).

7.   Wan, C., Hao, Z. & Feng, X. Structures, properties, and energy-storage mechanisms of the semi-lunar process cuticles in locusts. *Sci. Reports 2016 61* **6**, 1–13 (2016).

8.   Wan, C. & Gorb, S. N. Body-catapult mechanism of the sandhopper jump and its biomimetic implications. *Acta Biomater.* **124**, 282–290 (2021).

9.   Burrows, M. Development and deposition of resilin in energy stores for locust jumping. *J. Exp. Biol.* **219**, 2449–2457 (2016).

10.   Sutton, G. P. & Burrows, M. Biomechanics of jumping in the flea. *J. Exp. Biol.* **214**, 836–847 (2011).

11.   Bennet-Clark, H. C. Negative Pressures Produced in the Pharyngeal Pump of the Blood-Sucking Bug, Rhodnius Prolixus. *J. Exp. Biol.* **40**, 223–229 (1963).

12.   Young, D. & Bennet-Clark, H. C. The Role of the Tymbal in Cicada Sound Production. *J. Exp. Biol.* **198**, 1001–1020 (1995).

13.   Lyons, R. E. *et al.* Molecular and functional characterisation of resilin across three insect orders. *Insect Biochem. Mol. Biol.* **41**, 881–890 (2011).

14.   Stutt, A. D. & Siva-Jothy, M. T. Traumatic insemination and sexual conflict in the bed bug Cimex lectularius. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 5683–5687 (2001).

15.   Michels, J., Gorb, S. N. & Reinhardt, K. Reduction of female copulatory damage by resilin represents evidence for tolerance in sexual conflict. *J. R. Soc. Interface* **12**, (2015).

16.   McKenzie, E. K. G., Kwan, G. T., Tresguerres, M. & Matthews, P. G. D. A pH-powered mechanochemical engine regulates the buoyancy of Chaoborus midge larvae. *Curr. Biol.* **32**, 927-933.e5 (2022).

17. Andersen, S. O. The cross-links in resilin identified as dityrosine and trityrosine. *BBA - Gen. Subj.*
93, (1964).

18. Ardell, D. H. & Andersen, S. O. Tentative identification of a resilin gene in Drosophila
melanogaster. *Insect Biochem. Mol. Biol.* **31**, 965–970 (2001).

19. Balu, R., Dutta, N. K., Dutta, A. K. & Choudhury, N. R. Resilin-mimetics as a smart biomaterial
platform for biomedical applications. *Nat. Commun. 2021 121* **12**, 1–15 (2021).

20. Li, L. & Kiick, K. L. Resilin-based materials for biomedical applications. *ACS Macro Lett.* **2**,
635–640 (2013).

21. Qin, G. *et al.* Recombinant exon-encoded resilins for elastomeric biomaterials. *Biomaterials* **32**,
9231–9243 (2011).

22. Lerch, S. *et al.* Resilin matrix distribution, variability and function in Drosophila. *BMC Biol.* **18**,
1–15 (2020).

23. Ohkubo, S., Shintaku, T., Mine, S., Yamamoto, D. S. & Togawa, T. Mosquitoes Possess
Specialized Cuticular Proteins That Are Evolutionarily Related to the Elastic Protein Resilin.
Insects **14**, 941 (2023).

24. Andersen, S. O. Studies on resilin-like gene products in insects. *Insect Biochem. Mol. Biol.* **40**,
541–551 (2010).

25. Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with AlphaFold 3.
Nature **630**, 493–500 (2024).

26. Lerch, S., Yang, Y., Flaven-Pouchon, J., Gehring, N. & Moussian, B. Resilin is needed for wing
posture in Drosophila suzukii. *Arch. Insect Biochem. Physiol.* **111**, e21913 (2022).

27. Neff, D. Bursicon, Metamorphosis and Development of Resilin in the Fruit Fly. *Theses, Diss.*
Capstones (2012).

28. Wong, D. C. C., Pearson, R. D., Elvin, C. M. & Merritt, D. J. Expression of the rubber-like
protein, resilin, in developing and functional insect cuticle determined using a Drosophila anti-rec
1 resilin antibody. *Dev. Dyn.* **241**, 333–339 (2012).

29. Fabian, B., Schneeberg, K. & Beutel, R. G. Comparative thoracic anatomy of the wild type and
wingless (wg1cn1) mutant of Drosophila melanogaster (Diptera). *Arthropod Struct. Dev.* **45**,
611–636 (2016).

30. Dietzl, G. *et al.* A genome-wide transgenic RNAi library for conditional gene inactivation in
Drosophila. *Nat. 2007 4487150* **448**, 151–156 (2007).

31. Greene, J. C. *et al.* Mitochondrial pathology and apoptotic muscle degeneration in Drosophila
parkin mutants. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 4078–4083 (2003).

32.   Park, J. *et al.* Mitochondrial dysfunction in Drosophila PINK1 mutants is complemented by

739   parkin. *Nature* **441**, 1157–1161 (2006).

740   33.   Michels, J., Appel, E. & Gorb, S. N. Functional diversity of resilin in Arthropoda. *Beilstein*

741   *Journal of Nanotechnology* (2016) doi:10.3762/bjnano.7.115.

742   34.   Appel, E., Heepe, L., Lin, C. P. & Gorb, S. N. Ultrastructure of dragonfly wing veins: Composite

743   structure of fibrous material supplemented by resilin. *J. Anat.* **227**, 561–582 (2015).

744   35.   Michels, J. & Gorb, S. N. Detailed three-dimensional visualization of resilin in the exoskeleton of

745   arthropods using confocal laser scanning microscopy. *J. Microsc.* **245**, 1–16 (2012).

746   36.   Gidaszewski, N. A., Baylac, M. & Klingenberg, C. P. Evolution of sexual dimorphism of wing

747   shape in the Drosophila melanogaster subgroup. *BMC Evol. Biol.* **9**, 1–11 (2009).

748   37.   Anh, N. T. T., Nishitani, M., Harada, S., Yamaguchi, M. & Kamei, K. Essential role of duox in

749   stabilization of Drosophila wing. *J. Biol. Chem.* (2011) doi:10.1074/jbc.M111.263178.

750   38.   Edens, W. A. *et al.* Tyrosine cross-linking of extracellular matrix is catalyzed by Duox, a

751   multidomain oxidase/peroxidase with homology to the phagocyte oxidase subunit gp91phox. *J.*

752   *Cell Biol.* (2001) doi:10.1083/jcb.200103132.

753   39.   Hurd, T. R., Liang, F. X. & Lehmann, R. Curly Encodes Dual Oxidase, Which Acts with Heme

754   Peroxidase Curly Su to Shape the Adult Drosophila Wing. *PLOS Genet.* **11**, e1005625 (2015).

755   40.   Ha, E. M., Oh, C. T., Bae, Y. S. & Lee, W. J. A direct role for dual oxidase in Drosophila gut

756   immunity. *Science (80-.).* **310**, 847–850 (2005).

757   41.   Burrows, M., Shaw, S. R. & Sutton, G. P. Resilin and chitinous cuticle form a composite

758   structure for energy storage in jumping by froghopper insects. *BMC Biol.* **6**, 1–16 (2008).

759   42.   Burrows, M. & Sutton, G. P. Locusts use a composite of resilin and hard cuticle as an energy

760   store for jumping and kicking. *J. Exp. Biol.* **215**, 3501–3512 (2012).

761   43.   De Deken, X. *et al.* Cloning of Two Human Thyroid cDNAs Encoding New Members of the

762   NADPH Oxidase Family. *J. Biol. Chem.* **275**, 23227–23233 (2000).

763   44.   Dupuy, C. *et al.* Purification of a novel flavoprotein involved in the thyroid NADPH oxidase.

764   Cloning of the porcine and human cDNAs. *J. Biol. Chem.* (1999) doi:10.1074/jbc.274.52.37265.

765   45.   Ha, E. M. *et al.* Coordination of multiple dual oxidase–regulatory pathways in responses to

766   commensal and infectious microbes in drosophila gut. *Nat. Immunol. 2009 109* **10**, 949–957

767   (2009).

768   46.   Lee, K. A. *et al.* Bacterial-derived uracil as a modulator of mucosal immunity and gut-microbe

769   homeostasis in drosophila. *Cell* **153**, 797–811 (2013).

47.   Moreira, S., Stramer, B., Evans, I., Wood, W. & Martin, P. Prioritization of Competing Damage

and Developmental Signals by Migrating Macrophages in the Drosophila Embryo. *Curr. Biol.* **20**,

464–470 (2010).

48.   Niethammer, P., Grabher, C., Look, A. T. & Mitchison, T. J. A tissue-scale gradient of hydrogen

peroxide mediates rapid wound detection in zebrafish. *Nat. 2009 4597249* **459**, 996–999 (2009).

49.   Thein, M. C. *et al.* Combined Extracellular Matrix Cross-linking Activity of the Peroxidase

MLT-7 and the Dual Oxidase BLI-3 Is Critical for Post-embryonic Viability in Caenorhabditis

elegans. *J. Biol. Chem.* **284**, 17549–17563 (2009).

50.   Suderman, R. J., Dittmer, N. T., Kramer, K. J. & Kanost, M. R. Model reactions for insect cuticle

sclerotization: Participation of amino groups in the cross-linking of Manduca sexta cuticle protein

MsCP36. *Insect Biochem. Mol. Biol.* **40**, 252–258 (2010).

51.   Kumar, S., Molina-Cruz, A., Gupta, L., Rodrigues, J. & Barillas-Mury, C. A peroxidase/dual

oxidase system modulates midgut epithelial immunity in anopheles gambiae. *Science (80-.).* **327**,

1644–1648 (2010).

52.   O'Sullivan, A. *et al.* Multifunctional Wing Motor Control of Song and Flight. *Curr. Biol.* **28**,

2705-2717.e4 (2018).

53.   Carvajal, M. F. C. A. *et al.* Dynamics in natural and designed elastins and their relation to elastic

fiber structure and recoil. *Biophys. J.* **120**, 4623–4634 (2021).

54.   Jamhawi, N. M., Koder, R. L. & Wittebort, R. J. Elastin recoil is driven by the hydrophobic effect.

Proc. Natl. Acad. Sci. U. S. A. **121**, e2304009121 (2024).

55.   Trębacz, H. & Barzycka, A. Mechanical Properties and Functions of Elastin: An Overview.

Biomol. 2023, Vol. 13, Page 574 **13**, 574 (2023).

56.   Qin, G., Hu, X., Cebe, P. & Kaplan, D. L. Mechanism of resilin elasticity. *Nat. Commun.* **3**,

(2012).

57.   Rogers, S. M. *et al.* RNAi of the elastomeric protein resilin reduces jump velocity and resilience

to damage in locusts. *Proc. Natl. Acad. Sci. U. S. A.* **122**, e2415625121 (2025).

58.   Michels, J., Vogt, J. & Gorb, S. N. Tools for crushing diatoms - Opal teeth in copepods feature a

rubber-like bearing composed of resilin. *Sci. Rep.* **2**, 1–6 (2012).

59.   Birkedal, H. *et al.* Halogenated veneers: Protein cross-linking and halogenation in the jaws of

Nereis, a marine polychaete worm. *ChemBioChem* **7**, 1392–1399 (2006).

60.   Kondo, S. & Ueda, R. Highly Improved gene targeting by germline-specific Cas9 expression in

Drosophila. *Genetics* **195**, 715–721 (2013).

61.   Wakiyama, M., Matsumoto, T. & Yokoyama, S. Drosophila U6 promoter-driven short hairpin

RNAs effectively induce RNA interference in Schneider 2 cells. *Biochem. Biophys. Res. Commun.*

331, 1163–1170 (2005).

62. Hirai, K. *et al.* The Drosophila nucleoporin ELYS is required for parental chromosome arrangement at fertilization. *G3 Genes|Genomes|Genetics* **15**, 104 (2025).

63. Bischof, J., Maeda, R. K., Hediger, M., Karch, F. & Basler, K. An optimized transgenesis system for Drosophila using germ-line-specific φC31 integrases. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 3312–3317 (2007).

64. Groth, A. C., Fish, M., Nusse, R. & Calos, M. P. Construction of transgenic Drosophila by using the site-specific integrase from phage phiC31. *Genetics* **166**, 1775–1782 (2004).

65. Qin, G. *et al.* Expression, cross-linking, and characterization of recombinant chitin binding resilin. *Biomacromolecules* **10**, 3227–3234 (2009).

66. Zhang, X. *et al.* Mechanoresponsive regulation of myogenesis by the force-sensing transcriptional regulator Tono. *Curr. Biol.* **34**, 4143-4159.e6 (2024).

67. Misof B. *et al*. Phylogenetics resolves the timing and pattern ofinsect evolution. Science, 346, No. 6210 (2014).

 Figure legends

Figure 1. Exon−intron organization in the *Rsl* locus and protein structures of Rsl. (A)

The top is a schematic of the *Rsl* locus in the wild-type fly. Black and white boxes are

CDSs and UTRs, respectively. The exon compositions of the two isoforms are shown

below. The schematics below show the *Rsl* locus in *Rsl^{KO}* flies; the regions chosen to

generate the transgenes for the rescue experiments (*Rsl-GR*) and for the Gal4 driver

(*Rsl-Gal4*) are shown. The short bars indicate two gRNAs designed for the deletion of

CDS to generate *Rsl^{KO}* flies. (B) Domain-motif organization of Rsl isoforms are shown.

Each of the functional domain or consensus is shown below. Rsl-PB has a small portion

of the chitin-binding R&R consensus sequence (red box) encoded by the 5' part of exon

3. (C) The 3D model of Rsl-PA predicted in AlphaFold Server is shown. Panel a is the

whole Rsl-PA model. Blue and green parts are the N-terminal and C-terminal repeats,

respectively. In panels b and c, the N-terminal and C-terminal repeats are enlarged with

their proline and tyrosine residues illustrated by blue and purple colors, respectively. In

panel c, beta turns are indicated by closed arrowheads. Panels d and e show the views in

which two repeats are extracted from panels b and c, respectively, with schematics for

two cycles of their polypeptide backbones.

Figure 2. Temporal patterns of *Rsl* expression and spatial patterns of the reporter. (A)

RT-PCRs of *Rsl-RA* and *Rsl-RB* from embryo to day 5 adult are shown. (B) Western

blots of the Rsl protein from white pupa to day 5 adult are shown. In both A and B, FL,

WL, WP, P1-4, and A0-5 are the abbreviations of feeding larva, wandering larva, white

pupa, pupa days 1−4, and adult day 0−5, respectively. (C−H and J) Spatial distribution

of the Rsl-RA::mCherry reporter expressed with *Rsl-Gal4* is shown. (C) Panels a – c

show the views from dorsal, lateral, and ventral directions, respectively. Photos on the

left side were taken under white light. In each panel, white arrowheads indicate the

mCherry signals at the wing hinges, spiracles, or labrum. The dotted circle indicates the

outline of the compound eye. (D and E) The midleg of the flies (D) and the edge of the

wing (E) are shown. Photos of the mCherry signal and bright field were merged. Scale

bar: 50 μm. (F) The western blots of Rsl in the compound eyes dissected from day-3

pupae of the wild-type and Rsl^{KO} flies are shown. The electrophoretic mobilities of

Rsl-PA and Rsl-PB are indicated with arrowheads. (G) The Rsl-RA::mCherry

852 localization at the wing hinge is shown. The outer cuticle was removed together with

853 the wing (arrowhead) to observe the internal structure (panel h). In panel i, the area

854 enclosed in the square in panel h is enlarged. Scale bar: 500 μm. (H) Panel j is the

855 magnified view of the wing hinge of the fly expressing *Rsl-RA::mCherry*. Panel k is the

856 photo taken under UV radiation, and panel l is the merged photo of panels j and k. Scale

857 bar: 50 μm. (I) The distribution pattern of ChtVis-tdTomato that was expressed with

858 *Rsl-Gal4* is shown. Scale bar: 50 μm. In both H and I, the wavelengths of the filters

859 used for excitation and the detection of dityrosine are 330−385 and >420 nm,

860 respectively. (J) The biased distributions of Rsl::mCherry signals are compared. Panels

861 p and q are the photos of the resilin matrix visualized with Rsl-RA::mCherry and

862 Rsl-RB::mCherry, respectively. In panel r, the deviations of the biased distribution were

863 calculated as described in Materials and Methods.

865 Figure 3. Phenotypes of *Rsl* knockdown and mutant flies. (A) The wing postures of the

866 control fly (left: *Rsl>GFP-RNAi*) and *resilin* knockdown fly (right: *Rsl>Rsl-RNAi*) are

867 compared. Scale bar: 0.5 mm. (B) The proportion of flies with normal wing posture on

each day against the total number of flies is indicated (closed circles: *Rsl>GFP-RNAi*,

grey circles: *Rsl>Rsl-RNAi*). (C) The wing postures of wild-type (w^{1118}), Rsl^{KO} (w^{1118};

Rsl^{KO}), and rescued (w^{1118}; *resilin*^{KO}; *resilinGR*) flies are compared. Scale bar: 0.5 mm.

(D) The temporal changes in the proportion of flies with normal wing posture after

eclosion are shown. The fly genotypes are w^{1118} (closed circles), *resilin*^{KO} (gray circles),

resilin^{KO}; *resilinGR* (open circles), *resilin*^{KO}; *resilin-Gal4* (closed triangles), *resilin*^{KO},

UAS-resilin-RA (gray triangles), *resilin*^{KO}, *UAS-resilin-RB* (open triangles), *resilin*^{KO},

UAS-resilin-RA; resilin-Gal4 (closed squares), *resilin*^{KO},*UAS-resilin-RB; resilin-Gal4*

(gray squares), and *Rsl*^{KO},*UAS-Rsl-RA/Rsl*^{KO},*UAS-Rsl-RB; Rsl-Gal4* (open squares). (E)

The middle (Direction A) is a schematic diagram of the wing hinge that is viewed from

the midline toward the wing (blue arrow in the dorsal view of the fly). The right

(Direction B) shows the wing hinge viewed from posterior to anterior direction (red

arrow in the dorsal view of the fly). Blue shows the tergopleural tendon, which connects

to the cuticle where the wing attaches (yellow arrowheads). Red indicates direct flight

muscles TP1 and TP2[52]. (F) The inner structure of the wing hinge in *Dpy-YFP* was

observed (panel a, left part), and its schematic is shown (right part). The red signals

indicate muscles stained with phalloidin. The blue signal corresponds to blue

fluorescence emitted upon UV irradiation. Green signals indicate Dpy-YFP. White

arrow heads indicate the cuticle structure beneath the wing hinge attached to the

proximal end of the tergopleural tendon. Panels b and c show the inner structures and

the illustrations of the wing hinge of Rsl^{KO} flies (with and without the downturned wing

phenotype, respectively). Panel d shows those of the Rsl^{KO} fly with Rsl-GR. In Figure

3F, the adults 72 h after eclosion were used for taking the photos, except for panel b for

which adults 0 h after eclosion were used. Yellow arrowheads indicate the internal

cuticle structure corresponding to the root of the wing hinge. White arrowheads indicate

the tergopleural tendon devoid of any blue fluorescence signal in Rsl^{KO} flies. Asterisks

indicate Dpy-YFP signals attached to flight muscles other than TP-1 and TP-2. Scale

bar: 50 μm.

Figure 4. Analyses of *Duox* functions in Rsl polymerization. (A) The wing postures of

the control (*Rsl>GFP-IR*) and *Duox* knockdown flies (*Rsl>NIG 3131R-3*, and

Rsl>Duox-RNAi$_{976–1145}$) are compared. Scale bar: 0.5 mm. (B) The temporal changes in

the proportions of flies with normal wing posture after eclosion are shown. The flies

used were the same as those in (A) (closed circles: *Rsl>GFP-RNAi*, grey closed circles:

Rsl>NIG3131R-3, and gray open circles: *Rsl>Duox-RNAi$_{976–1145}$*). (C) Western blot

analyses of Rsl protein levels were performed using samples from day-4-APF pupae to

day-2-after-eclosion adults. The left and right are flies of the control (*Rsl>GFP-RNAi*)

and *Duox* knockdown (*Rsl>NIG3131R-3*), respectively. (D) The possible model of

Duox-mediated Rsl polymerization (after Edens et. al., 2001[38]) (E) The inner structures

of the wing hinge in the two control flies, *Rsl-Gal4/+* (panel a) and *NIG3131R-3/+*

(panel b), are shown. The changes in the structure in the *Duox*-knockdown flies

(*Rsl>NIG3131R-3*) are shown (panels c and d). Panel e on the right is the schematic of

panels c and d. TP-1 and TP-2 muscles were stained with phalloidin. In

Duox-knockdown flies, the dotted line (panels c and d) indicates the outline of the

dityrosine signal corresponding to the weak blue fluorescence signal indicated by

triangles in the photo taken with a longer exposure time. Scale bar: 50 μm. (F) Wing

posture of the *Duox* knockdown fly (*Rsl>NIG3131R-3, eGFP*) and that of the *Duox*

knockdown fly with the additional *TcDuox* expression (*Rsl> NIG3131R-3, TcDuox*).

Scale bar: 0.5 mm. (G) Difference in the proportion changes of flies with normal wing

posture was compared between *Rsl> NIG3131R-3, eGFP* (gray circle) and

Rsl>NIG3131R-3, TcDuox (closed circle) flies. (H) Western blot analyses were

performed to examine the levels of the Rsl protein from day-4-APF pupae to

day-2-after-eclosion adults are shown. The left shows the *Duox*-knockdown individuals

(*Rsl>NIG3131R-3, eGFP*) and the right shows the rescued individuals

(*Rsl>NIG3131R-3, TcDuox*). (I) The photos and illustrations of the inner structure of the

wing hinge in *Duox*-knockdown (left panels) and rescued flies (right panels) are shown.

Scale bar: 50 μm (J) Western blot analyses of reRsl protein and dityrosine levels. reRsl

was incubated in the presence (+) or absence (-) of 20 μg/μL horseradish peroxidase

(HRP) and/or 0.3% hydrogen peroxide (H_2O_2) in 50 mM sodium phosphate buffer (pH

7.0) for 1 h at 30°C. Closed arrowheads show the signals stacking at the entry site of the

concentration gels. The open arrowhead shows the signal stacking at the border between

the concentration and separation gels. Curly bracket shows smear signals in the stacking

gel.

Figure 5. Contributions of *Rsl* and *Duox* to fly's jumping. (A) Experimental scheme to

measure jump distance of de-winged flies. (B) The jump distances between the control

flies (*Rsl>GFP-RNAi*, white box) and *Rsl* knockdown flies (*Rsl>Rsl-RNAi*, gray box)

are compared. Cross marks indicate the average. (Wilcoxon rank sum test, **: $P < 0.01$,

Rsl>GFP-RNAi: n=8, *Rsl>Rsl-RNAi*: n=12). (C) The Rsl^{KO} phenotype of jump distance

was rescued with *Rsl-GR*. The flies used were the wild-type (w^{1118}, white box), Rsl^{KO}

(w^{1118}; Rsl^{KO}, gray box), and Rsl^{KO} with *Rsl-GR* (w^{1118}; Rsl^{KO}; *Rsl-GR*, dark gray box).

Cross marks indicate the average (Wilcoxon rank sum test, Bonferroni correction, w^{1118}:

n=10, Rsl^{KO}: n=10, Rsl^{KO} with *Rsl-GR*: n=11). (D) Effect of *Duox* knockdown on jump

performance was examined. The graph shows the jump distances of the control flies

(*Rsl>GFP-RNAi,* white box), the *Duox* knockdown flies (*Rsl> NIG 3131R-3*, indicated

by the light gray box, and *Rsl>Duox-RNAi$_{976-1145}$* indicated by the gray box) (Wilcoxon

rank sum test, Bonferroni correction, *Rsl>GFP-RNAi*: n=9, *Rsl> NIG 3131R-3*: n=12,

Rsl>Duox-RNAi$_{976-1145}$: n=8). (E) The phenotype of *Duox* knockdown flies was rescued

by the additional expression of *TcDuox*. The graph shows the jump distances of the

Duox knockdown flies (*Rsl> NIG3131R-3, eGFP,* white box) and the flies with *TcDuox*

expression (*Rsl> NIG3131R-3, TcDuox*, gray box) (Wilcoxon rank sum test, **: *P <*

0.01, *Rsl> NIG3131R-3, eGFP*: n=9, *Rsl> NIG3131R-3, TcDuox*: n=7).

Figure 1

953
954

955   Figure 2

956
957

958   Figure 3

960   Figure 4-1

A Rsl >GFP-RNAi   Rsl >NIG3131R-3   Rsl >Duox-RNAi_{976–1145}

B Proportion of flies with normal wings (%) vs Days after eclosion (Day)

C Rsl >GFP-RNAi   Rsl >NIG3131R-3
   P4 A0 A1 A2   P4 A0 A1 A2
   Rsl   ◄ Rsl-PA
                                 ◄ Rsl-PB
   α-Tubulin

D Rsl molecule ROS
   Peroxidase homology domain
   NADPH oxidase domain   Outside
   EF hand motifs   Inside

E Rsl-Gal4 / +   NIG3131R-3/ +   Rsl>NIG3131R-3
   a   b   c UV+phalloidin d Dpy+phalloidin e

962   Figure 4-2

964 Figure 5