# Intravital calcium imaging of meningeal macrophages reveals niche-specific dynamics and aberrant responses to brain hyperexcitability

Simone Carneiro-Nascimento[1], Chao Wei[1], Anna Gutterman[1], Dan Levy[1]*

*Correspondence:

Dan Levy

dlevy1@bidmc.harvard.edu

[1]Department of Anesthesia, Critical Care and Pain Medicine, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, United States

# Abstract

The meninges, which envelop and protect the brain, host a dense network of resident macrophages with diverse roles in regulating homeostasis and neuroinflammation. Despite their importance, we have a limited understanding of their behavior in vivo. Many dynamic cellular functions of macrophages involve intracellular $Ca^{2+}$ signaling. However, to our knowledge, virtually nothing is known about the spatiotemporal $Ca^{2+}$ dynamics of meningeal macrophages. We developed a chronic intravital two-photon imaging approach and related computational analysis tools to interrogate meningeal macrophage $Ca^{2+}$ dynamics, at a subcellular resolution, in a novel Pf4$^{Cre}$:TIGRE2.0$^{GCaMP6s/wt}$ reporter mouse model. Using imaging in awake mice, we characterized the $Ca^{2+}$ activity of meningeal macrophages at steady state and in response to cortical spreading depolarization (CSD), an aberrant pro-inflammatory brain hyperexcitability event, implicated in migraine, traumatic brain injury, and stroke. In homeostatic meninges, macrophages in the dural perivascular niche exhibited several $Ca^{2+}$ dynamic features, including event duration and signal frequency spectrum, distinct from those of in the interstitial, non-perivascular niche. Simultaneous tracking of meningeal macrophage $Ca^{2+}$ dynamics and local vasomotion revealed a subset of dural perivascular macrophages whose activity was coupled to behaviorally-driven diameter fluctuations of their associated vessels. Most perivascular and non-perivascular meningeal macrophages displayed propagating intracellular $Ca^{2+}$ activity as well as synchronized intercellular $Ca^{2+}$ elevations, likely driven by extrinsic factors. In response to CSD, the majority of perivascular and non-perivascular meningeal macrophages showed a persistent decrease in $Ca^{2+}$ activity, while a smaller subset displayed $Ca^{2+}$ elevations.

Mechanistically, CGRP/RAMP1 signaling mediated the increase but not the decrease in CSD-mediated $Ca^{2+}$ signaling. Collectively, our results highlight a previously unknown diversity of meningeal macrophage $Ca^{2+}$ dynamics at steady state and in response to an aberrant brain hyperexcitability event linked to neuroinflammation.

**Keywords:** Meningeal macrophages, calcium imaging, dural vasomotion, cortical spreading depolarization, migraine

# Introduction

Resident macrophages are key myeloid immunocytes that play an important role in innate immune surveillance and defense across various peripheral tissues and organs [1-3]. The central nervous system also harbors a large subset of parenchymal macrophages, known as microglia, and several distinct subsets of macrophages localized to the brain's border tissues, including the choroid plexus, perivascular spaces, and the meningeal compartments that cover, protect, and support the brain [4-8]. Macrophages are the predominant immune cell type within the brain meninges, and recent studies have demonstrated the diverse ontogeny, transcriptomic profiles, and immune functions of meningeal macrophages at steady state [5, 8-12] and in several neuropathological conditions [8, 13-16].

Cytoplasmic calcium ($Ca^{2+}$) signaling underlies a wide variety of cellular homeostatic and inflammatory processes in macrophages [17-23]. In addition to intracellular $Ca^{2+}$ elevation, distinct spatiotemporal dynamics - including oscillation patterns, intracellular propagations, and intercellular synchronization of $Ca^{2+}$ signals - may regulate different macrophage functions during steady state and pathophysiology [20-25]. Despite our increased understanding of the diverse molecular signatures and contributions of meningeal macrophages to homeostasis and neuroinflammation, virtually nothing is known about their $Ca^{2+}$ signaling heterogeneity in both healthy and diseased states.

To comprehensively characterize the $Ca^{2+}$ dynamics of individual macrophages localized

to the brain meninges, we combined intravital two-photon $Ca^{2+}$ imaging in a novel reporter

mouse line, in which the $Ca^{2+}$ reporter GCaMP6s is expressed in platelet factor 4 ($Pf4^+$)

meningeal macrophages, with an event-based signaling analysis pipeline. Our data

reveal several distinct spatiotemporal $Ca^{2+}$ dynamic features in perivascular versus non-

perivascular meningeal macrophages, including a unique coupling between the $Ca^{2+}$

signals of dural perivascular macrophages and behaviorally-driven vasomotion of their

associated dural vessels at steady state. Furthermore, our data uncover both increases

and decreases in $Ca^{2+}$ activity in distinct subsets of meningeal macrophages in response

to cortical spreading depolarization (CSD), a pathophysiological brain hyperexcitability

event linked to neuroinflammation [26] in migraine, traumatic brain injury, and stroke [27].

Mechanistically, our data suggest the calcitonin gene-related peptide/receptor activity-

modifying protein 1 (CGRP/RAMP1) axis uniquely mediates CSD-evoked macrophage

$Ca^{2+}$ elevation, indicating a brain-to-meninges neuroimmune signaling pathway involving

the activation of meningeal peptidergic sensory neurons.

# 39    Results

## 40    Characterizing macrophage $Ca^{2+}$ signaling features in homeostatic

## 41    brain meninges

Previous intravital imaging studies exploring the spatiotemporal dynamics of tissue-

resident macrophages have primarily used CX3C motif chemokine receptor 1 (CX3CR1)-

based mouse reporter strains [13, 22, 28, 29]. However, other resident monocyte-derived

cells are labeled in these reporter mice [30, 31]. Moreover, brain microglia also express

CX3CR1 [13, 31-33], limiting the use of these reporter mice for resolving the

spatiotemporal subcellular $Ca^{2+}$ dynamics of macrophages localized to the relatively thin

meningeal layers covering the brain parenchyma. To systematically characterize

meningeal macrophage $Ca^{2+}$ dynamics *in vivo* and avoid contamination from $Ca^{2+}$ signals

arising from other resident monocyte-derived meningeal cells and superficial

parenchymal microglia, we leveraged recent findings showing the expression of Pf4 in

meningeal macrophages but not in parenchymal microglia, [10, 16, 34], and generated

transgenic reporter mice expressing the highly sensitive $Ca^{2+}$ indicator GCaMP6s (using

TIGRE2.0-based Ai162D mice [35]) in Pf4$^{+}$ macrophages (using Pf4$^{Cre}$ mice [36]).

Notably, nearly 100% of meningeal macrophages are labelled in Pf4$^{Cre}$-based reporter

mice [34]. Although Pf4 is also expressed by megakaryocytes [36], these cells are absent

in the brain meninges [5].

Anesthetic agents impact intracellular $Ca^{2+}$ signaling, including in brain macrophages and

other non-excitable glial cells [37-39]. We therefore investigated subcellular $Ca^{2+}$ activity

of meningeal macrophages in awake, behaving mice. We imaged meningeal

macrophages via a chronic cranial window implanted together with a restraining headpost

over the intact dura mater overlying the posterior neocortex. This chronic window

approach produces minimal inflammatory responses in the cortex and meninges below

the window [40, 41]. After at least 7 days of recovery, mice were gradually habituated

across multiple days to head restraint while free to run on a wheel (**Fig 1b**). We tracked

$Ca^{2+}$ transients of GCaMP6s-labeled meningeal macrophages using high-speed two-

photon microscopy in 37 fields of view (FOVs) from 7 mice. We first corrected the imaging

movies for locomotion-evoked meningeal translational shifts using rigid registration [41].

Movies were then processed using the AQuA2 data analysis platform [42], which

implements an unbiased event-based approach to capture spatiotemporal $Ca^{2+}$ event

dynamics (**Fig 1C**). Based on spatial analysis, we assigned events (n=1361), with their

corresponding $Ca^{2+}$ features, to a given macrophage, with data gathered from a total of

503 cells.

**Figure 1. Imaging meningeal macrophage $Ca^{2+}$ dynamics in awake behaving mice. (A)**
Pf4$^{Cre}$:GCaMP6s$^{fl/wt}$ reporter mouse construct for imaging meningeal macrophages $Ca^{2+}$ activity. **(B)**
Experimental procedure for two-photon imaging of meningeal macrophage $Ca^{2+}$ activity. Following the
implantation of a headpost and a cranial window, mice were habituated to head restraint and subjected to
two-photon microscopy while head-fixed on a running wheel to study meningeal macrophage $Ca^{2+}$ activity.
**(C)** Macrophage $Ca^{2+}$ imaging processing pipeline.
83

We characterized $Ca^{2+}$ dynamics in two anatomically distinct macrophage populations:

perivascular and interstitial non-perivascular (n=122; n=381, respectively, **Fig 2A and Video S1**).

The vast majority of perivascular macrophages (93.4%, n=114) displaying ongoing $Ca^{2+}$ activity

were associated with vessels in the dura mater. The two meningeal macrophage subpopulations

exhibited several distinct $Ca^{2+}$ activity features. While the total area of $Ca^{2+}$ activity in the peri- and

non-perivascular macrophages was similar (**Fig 2B**), the signal perimeter of perivascular

macrophages was significantly higher (**Fig 2C**) and exhibited a more elongated shape (**Fig 2A,**

**2D**). The $Ca^{2+}$ event duration in the perivascular macrophages was also longer compared to the

non-perivascular subpopulation (**Fig 2F**). The peak $Ca^{2+}$ activity level (Max $DF/F_0$, **Fig 2E**) and

event rate (**Fig 2G**), nonetheless, were similar in the two meningeal macrophage subpopulations.

**Figure 2. $Ca^{2+}$ dynamic features of meningeal macrophages at steady state. (A)** Left: Mean projection
of an example FOV depicting perivascular (P, red; 1, 2) and non-perivascular (NP, blue; 3, 4) meningeal
macrophages (white squares). Scale bar 50 μm. Right: Corresponding macrophages with representative
900-second $Ca^{2+}$ activity traces (top) and their fluorescence signal at selected time points (bottom). **(B-G)**
AQuA2-based morphological and $Ca^{2+}$ event functional features of perivascular (P=122 cells) and non-
perivascular (NP, n=381 cells) meningeal macrophage. (B) Event area, (C) Event perimeter, (D) Event
circularity, (E) Event max $DF/F_0$, (F) Event duration, (G) Event rate. Data (B-G) represent median ± IQR.
***p<0.001, ****p<0.0001, Mann-Whitney U-test.  n=37 fields of view (FOVs) from 7 mice.
104

Distinct $Ca^{2+}$ signal frequency spectra may underlie different biological functions of

macrophages [22]. We therefore analyzed meningeal macrophage $Ca^{2+}$ signal waveforms

by performing a multi-step signal processing and clustering analysis. We clustered cells

based on two distinct patterns of $Ca^{2+}$ activity. Cells in Cluster 1 (n=40) exhibited a more

noisy-like activity pattern characterized by multiple frequencies, while cells in Cluster 2

(n=463) displayed a single dominant frequency at 0.01 Hz (**Fig 3A**). To further explore

these cellular signaling differences, we combined the clustering data with AQuA2-derived

features and observed that macrophages in Cluster 1 showed a larger event perimeter

and lower circularity and peak magnitude compared to those in Cluster 2 (**Fig 3D-F**).

Cluster 1 cells also had a lower signal-to-noise ratio compared to Cluster 2 cells (**Fig 3I**),

but had differences in terms of event area, duration, or rate (Figures 3C, G, H). There

was a significant association between cell cluster (1 vs. 2) and cell type, with Cluster 1

predominantly comprised of perivascular macrophages, and Cluster 2 including primarily

non-perivascular macrophages (**Fig 3J**), further indicating that these two meningeal

macrophage subpopulations have distinct $Ca^{2+}$ signaling properties.

**Figure 3. Intracellular $Ca^{2+}$ signal frequency spectra of macrophage subsets in the steady-state meninges (A)** Left: Clustering of meningeal macrophage $Ca^{2+}$ activity based on frequency-domain features and peak detection. Cluster 1 (purple, n=40) and Cluster 2 (green, n=463). Right: Power spectrum density (PSD) of $Ca^{2+}$ signals for each cluster. **(B)** Example $DF/F_0$ heatmaps of Cluster 1 and Cluster 2 macrophages. **(C-H)** Morphological and $Ca^{2+}$ functional features of Cluster 1 and Cluster 2 macrophages. (C) Event area. (D) Event perimeter (E) Event circularity. (F) Event max $DF/F_0$. (G) Event duration. (H) Event rate. (I) Signal-to-noise ratio from Clusters 1 and 2. **(J)** Distribution of cell types across clusters. Data (C-I) represents median ± IQR. ***$p<0.001$, ****$p < 0.0001$, Mann-Whitney U-test Data (J) represents the cell proportion. ****$p<0.0001$, Chi-square test.

Intracellular $Ca^{2+}$ signal propagation underlies diverse cellular functions and has been recently identified in macrophages *in vitro* [21] and in skin-resident macrophages *in vivo* [43]. By assessing the $Ca^{2+}$ signal propagation maps of each event within a determined cell, we identified two distinct patterns of activity: propagating events in which $Ca^{2+}$ signals

traveled throughout the entire cell, and stationary events (**Fig 4A**). Propagating events

had varied signal source regions and directionality. Most macrophages (perivascular,

94.3%, n=115; non-perivascular, 86.9%, n=331) exhibited only propagating events, while

a small minority of cells displayed a mix of propagating and stationary events or

exclusively stationary activity events (**Fig 4B**).

Macrophage intercellular communication, including synchronized activity, potentially

across connected macrophage networks, has been implicated in maintaining tissue

homeostasis and immune function [18, 21, 44, 45]. To investigate meningeal macrophage

intercellular interactions, we characterized spatiotemporal relationships between $Ca^{2+}$

events in distinct cells within each FOV. We compared temporal factors, including the

relative onset latency ($t0_A - t0_B$) between different macrophages and the duration of the

first occurring event ($t1_A - t0_A$) (**Fig 4C**). We also compared the distances between

macrophage pairs exhibiting concurrent events and the number of synchronous events

(**Fig 4D**). Finally, we calculated the proportion of perivascular and non-perivascular

macrophages exhibiting synchronous $Ca^{2+}$ events. Across all FOVs, nearly half of the

macrophages (49.3%) exhibited synchronous $Ca^{2+}$ activity, involving cells at a wide range

of distances (**Fig 4E**), suggesting that spatial proximity does not influence event

synchronicity. Interestingly, there was poor correlation between the distance of cell pairs

exhibiting synchronous events and their timing (**Fig 4F**), suggesting that an external

factor, rather than intercellular coordinated networked activity, mediates this intercellular

signaling. When comparing synchronous activity among peri- and non-perivascular

macrophages, most cells exhibited mixed interactions between the same and different

subtypes (**Fig 4G**).

**Figure 4. Intracellular propagation and intercellular synchronization of meningeal macrophage Ca$^{2+}$ activity**. **(A)** Spatial maps of two distinct Ca$^{2+}$ events. Left: Propagating Ca$^{2+}$ activity. Right: Stationary Ca$^{2+}$ activity. **(B)** Distribution of event propagation profiles in perivascular and non-perivascular meningeal macrophages. **(C)** Schematic analysis paradigm for detecting synchronous Ca$^{2+}$ activity in meningeal macrophages. **(D)** Synchronous Ca$^{2+}$ events among meningeal macrophages within a FOV. Left: Mean projection of an example FOV showing Ca$^{2+}$ activity in distinct macrophages (colored/numbered). Scale bar 50 µm. Right: Spatial map of macrophages exhibiting synchronous Ca$^{2+}$ activity. Lines connect macrophages with synchronized Ca$^{2+}$ activity, and colors indicate the extent of Ca$^{2+}$ event synchronization. **(E)** Distribution of distances across macrophage pairs showing different numbers of synchronous Ca$^{2+}$ events. **(F)** Linear regression showing poor correlation between distances of macrophage with synchronized Ca$^{2+}$ activity and event delay. **(G)** Proportion of macrophages exhibiting a specific synchronized interaction (cells interacting only with the same subtype: P-P/NP-NP; cells interacting only with a different subtype: P-NP; mixed (cells interacting with the same and different subtypes).

**Dural perivascular macrophage Ca$^{2+}$ activity is tuned to behaviorally-**

**driven dural vasomotion**

Brain border macrophages in the leptomeninges and related parenchymal perivascular spaces regulate pial arterial vasomotion indirectly by affecting vessel stiffness [6]. Yet, interaction between vascular-associated macrophages residing in the dura mater, the outermost meningeal layer, and dural vasomotion remains unknown. We employed a Generalized Linear Model (GLM) approach to investigate functional interaction between dural perivascular macrophage Ca$^{2+}$ signals and behaviorally-driven dural vessel dynamics (**Fig 5A**). Dural arteries constrict during locomotion, while pial arteries dilate [46]. We analyzed the locomotion-associated responses of 86 meningeal vessels (32 FOVs from 5 mice) and identified a subset (22%, n=19), whereby the diameter changes were well fit by a GLM using the locomotion state as a predictor. Of these, we identified 74% (n=14) as dural vessels based on their GLM's negative coefficients consistent with constriction (**Fig 5B and 5C**). Next, we fitted the Ca$^{2+}$ signal observed in perivascular macrophages associated with these dural vessels (n=35) to a GLM using the diameter changes as a predictor variable. Overall, the Ca$^{2+}$ activity of 83% (n=29) of these dural macrophages was well predicted by the model (average deviance explained across all well-fit macrophages: 0.43±0.15, mean±SD). Analysis of the macrophage-vascular models' beta coefficients revealed two distinct interactions. About half of the macrophages (55%, n=16) exhibited negative coefficients (i.e., increase and decrease in Ca$^{2+}$ activity associated with dural vasoconstriction and recovery, respectively; **Fig 5B and 5D**). The remaining macrophages (45%, n=13) exhibited positive coefficients (i.e., decrease and increase in Ca$^{2+}$ activity in response to dural constriction and recovery,

respectively; **Fig 5B and 5 E**). The coefficients for increased and decreased macrophage

Ca$^{2+}$ activity peaked near zero delay relative to the vasoconstriction and were not

statistically different (**Fig 5D-E and 5H**). These data provide evidence that dural

perivascular macrophages are functionally coupled to dural vasomotion, either

responding to or mediating such behaviorally driven vascular response.

**Figure 5. Ca$^{2+}$ signals of dural perivascular macrophages are functionally coupled to**
**behaviorally driven dural vasomotion. (A)** Experimental paradigm: Locomotion data were
acquired during imaging in awake-behaving mice. Behaviorally-evoked changes in meningeal
vessel diameter were obtained using segmentation of vessels labeled with a tracer during

macrophage $Ca^{2+}$ and further tested for coupling with $Ca^{2+}$ signals. **(B)** Example data of
macrophages with $Ca^{2+}$ activity tuned to locomotion-related dural vessel vasomotion. Locomotion
bouts (top trace) and dural vessel diameter (black trace) that was well fit by a GLM (red line) using
locomotion state as a predictor. Note vasoconstriction during locomotion, indicating a dural
vessel. The two bottom traces depict the $Ca^{2+}$ signal (black traces) of distinct meningeal
macrophages that were well-fit by a GLM using dural vessel diameter as a predictor, showing
either a negative coefficient (purple) or a positive coefficient (green). **(C)** Temporal profile of dura
vessels GLM coefficients averaged across all well-fitted vessels (n=19). Traces represent the
median across all well-fit ROIs, with shaded regions indicating IQR. **(D, E)** Temporal profiles of
GLM coefficient values for macrophages' $Ca^{2+}$ activity averaged across all well-fitted cells (purple,
negative coefficient, n=16; green, positive coefficient, n=13). Traces represent the median across
all well-fit macrophages, with shaded regions indicating IQR. **(F)** The deviance explained
(goodness-of-fit estimates) of the vessel diameter data included in the GLM used to predict
macrophage $Ca^{2+}$ activity were not statistically different for macrophages showing negative (n=16)
and positive (n=13) coefficients, suggesting similar interaction levels. **(G)** Comparison of peak
positive and negative coefficients of well-fit macrophage $Ca^{2+}$ activity/vasomotion GLMs. **(H)**
Center of mass of GLM coefficients indicating that dural vessel diameter changes drive
bidirectional changes in fluorescence signal at zero delay. Data (F-H) are median ± IQR.
****$p<0.0001$ (Mann-Whitney U-test).
231

## An acute aberrant pro-inflammatory brain hyperexcitability event

## drives diverse $Ca^{2+}$ dynamics in meningeal macrophages

CSD is an aberrant brain hyperexcitability event linked to parenchymal inflammation and

pain in migraine, traumatic brain injury, and stroke [27, 47, 48], and could also affect

meningeal macrophages [49]. In anesthetized mice subjected to an acute CSD episode,

a small subset of meningeal macrophages undergoes morphological changes resembling

an inflammatory state [26]. Given the direct anatomical and functional connection

between the brain and meninges [11, 50] and the involvement of intracellular $Ca^{2+}$ influx

in macrophage inflammatory activation [51], we asked whether brain-to-meninges

signaling drives intracellular $Ca^{2+}$ elevations in meningeal macrophages in the wake of

CSD. We therefore triggered a single CSD episode in the frontal cortex [52, 53] and

characterized the related changes in meningeal macrophage $Ca^{2+}$ dynamics. In each

experiment, we verified CSD induction based on the associated acute meningeal

deformation and/or pial vasoconstriction observed in awake animals ([53] and **Video S2**).

We studied CSD-related $Ca^{2+}$ dynamics in 249 macrophages (perivascular, n=64; non-

perivascular, n=185; 13 FOVs from 10 mice). For each cell, we compared $Ca^{2+}$ event

rates during the passage of the CSD wave (1 min) and the PostCSD period (30 min) to

baseline (PreCSD, 30 min) to assess acute and persistent $Ca^{2+}$ responses, respectively.

We characterized cells as exhibiting persistently increased (event rate > 2× PreCSD),

decreased (event rate < 0.5× PreCSD), or unchanged responses. Given the low $Ca^{2+}$

activity rate observed under steady state and the likelihood that no spontaneous $Ca^{2+}$

elevations occur during the brief period of the CSD event, we considered cells to be either

acutely activated or exhibiting an unchanged $Ca^{2+}$ response. CSD evoked both acute and

persistent $Ca^{2+}$ activity changes in (**Fig 6 A-E and Video S2**). While smaller subsets of

meningeal macrophages exhibited acute (21.3%, n=53) and/or persistent increase

(22.1%, n=55) in their $Ca^{2+}$ activity, the majority of cells (58.6%, n=146) exhibited a

persistent decrease. An acute increase was observed more often in peri- vs non-

perivascular macrophages (perivascular, 32.8%, n=21; non-perivascular, 17.3%, n=32,

**Fig 6F**). Persistent changes in $Ca^{2+}$ activity were observed similarly in peri- and non-

perivascular macrophages (increases; perivascular, 28.2%, n=18; non-perivascular,

18.4%, n=37; decreases; perivascular, 50.0%, n=32; non-perivascular, 61.6%, n=114,

**Fig 6G**). The macrophages' propensity to develop a persistent $Ca^{2+}$ increase was

unrelated to their acute response (**Fig 6H**), suggesting that the mechanisms underlying

these two temporal responses are distinct. However, cells showing no acute activation

were more likely to display decreased $Ca^{2+}$ activity post-CSD (**Fig 6H**). Finally, we

observed that macrophages exhibiting a persistent increase in $Ca^{2+}$ activity had a lower

baseline activity when compared to cells showing a persistent (**Fig 6I**), suggesting that

this post-CSD response is influenced by the macrophages' basal response activity.

**Figure 6. Diverse meningeal macrophage Ca$^{2+}$ dynamics following cortical spreading depression (CSD). (A)** Experimental setup and example data: (top) A small burr hole was drilled above the frontal cortex 7 days before Ca$^{2+}$ imaging. Mice were pretreated with saline 30 min before imaging baseline macrophage Ca$^{2+}$ activity (30 min, PreCSD). CSD was then induced with a pin prick, and macrophage Ca$^{2+}$ activity was assessed during CSD (1 min during CSD), and

post-CSD (30 min, post-CSD). (Bottom) A raster plot of macrophage $Ca^{2+}$ activity showing the
acute and persistent increases and persistent decrease in response to CSD. **(B)** Example of
macrophage $Ca^{2+}$ fluorescence changes following CSD. Images depict the mean projection over
the specific experimental timeline. Arrows indicate an increase or a decrease in $Ca^{2+}$ activity.
Scale bar 50 μm. **(C)** Individual responses of perivascular (P, n=21) and non-perivascular (NP,
n=32) macrophages showing an acute increase in $Ca^{2+}$ activity. **(D)** Individual response of P
(n=18) and NP (n=37) macrophages exhibiting a persistent increase in $Ca^{2+}$ activity. **(E)** Individual
responses of P (n=32) and NP (n=114) macrophages showing a persistent decrease in $Ca^{2+}$
activity. **(F)** Proportion of P and NP macrophages showing an acute increase in $Ca^{2+}$ activity or
no acute change. **(G)** Proportion of P and NP macrophages showing a persistent increase,
decrease, or no change in $Ca^{2+}$ activity. **(H)** Proportion of macrophages displaying distinct
persistent responses stratified based on their acute response. **(I)** Baseline (PreCSD) $Ca^{2+}$ activity
in macrophages exhibiting persistent increased (n=55) or decreased activity (n=146).  Data (C-E)
are median ± IQR. ****$p<0.0001$ (Wilcoxon signed rank test). Data (F-H) represents the proportion
of cells, **$p<0.01$; ****$p<0.0001$ (Chi-square test). Data (I) are median ± IQR. ****$p<0.0001$ (Mann-
Whitney U-test).
293

## CGRP/RAMP1 signaling mediates CSD-evoked persistent increase in meningeal macrophage $Ca^{2+}$ activity

Many meningeal macrophages are localized near peptidergic sensory axons [16]. In the

wake of CSD, cortex-to-meninges signaling enhances the responsiveness of meningeal

sensory neurons and can drive the release of the neuropeptide CGRP from their

peripheral nerve endings [52-54]. CGRP-expressing sensory neurons regulate tissue

immunity, including meningeal macrophage function, via the CGRP/RAMP1

neuroimmune axis [16, 55]. We therefore asked whether the CSD-related changes in

meningeal macrophage $Ca^{2+}$ dynamics we observed involve CGRP/RAMP1 signaling.

We pretreated mice with the RAMP1 antagonist BIBN4096 and then imaged meningeal

macrophage $Ca^{2+}$ activity (42 cells; perivascular, n=14; non-perivascular, n=28; 3 FOVs

from 3 mice) before and after CSD. As expected, RAMP1 blockade did not affect CSD

triggering [56]. When compared to control saline treatment, RAMP1 blockade also did not

reduce basal macrophage $Ca^{2+}$ activity (**Fig 7B**). Blocking CGRP/RAMP1 signaling

neither affected the incidence of acute increases in $Ca^{2+}$ activity (**Fig 7A and 7C**) nor the

magnitude of that response (**Fig 7D**). RAMP1 antagonism, however, inhibited the CSD-

evoked persistent increase in the macrophage's $Ca^{2+}$ activity, without affecting the

incidence of the persistent decrease (**Fig 7E**). The data suggest that in the wake of CSD,

the CGRP/RAMP1 axis, likely involving a peripheral sensory neuroimmune signaling, is

responsible for the prolonged enhancement of $Ca^{2+}$ signaling in a subset of meningeal

macrophages, which could potentially mediate their pro-inflammatory response.

**Figure 7. CGRP/RAMP1 signaling mediates CSD-related persistent increase in meningeal**
**macrophage $Ca^{2+}$ activity. (A)** Experimental setup and example data: (top) A small burr hole
was drilled above the frontal cortex 7 days before $Ca^{2+}$ imaging. Mice were pretreated with the
RAMP1 antagonist BIBN4096 (BIBN) 30 min before imaging baseline macrophage $Ca^{2+}$ activity
(30 min, PreCSD). CSD was then induced with a pin prick, and macrophage $Ca^{2+}$ activity was
assessed during CSD (1 min during CSD), and post-CSD (30 min, post-CSD). (Bottom) A raster
plot of macrophage $Ca^{2+}$ activity showing the acute and persistent increases and persistent

decrease in response to CSD. **(B)** RAMP1 inhibition does not affect the baseline (PreCSD) event
rate. Data compared between macrophages imaged in saline-treated mice (n=249 cells) and
BIBN-treated mice (n=42 cells). **(C)** RAMP1 antagonism does not affect the CSD-evoked acute
increase in macrophage $Ca^{2+}$. Proportion of macrophages showing an acute response (increase
vs. no change) in saline-treated mice (n=249 cells) and BIBN-treated mice (n=42 cells). **(D)**
RAMP1 antagonism does not affect the magnitude of the acute macrophage $Ca^{2+}$ signal. Event
max $DF/F_0$ in macrophages showing an acute $Ca^{2+}$ increase in saline-treated mice (n=53 cells)
and BIBN-treated mice (n=8 cells). **(E)** RAMP1 antagonism distinctly inhibits the persistent
increase in macrophage $Ca^{2+}$ activity post CSD. Proportion of macrophages showing persistent
increase, persistent decrease, or no persistent change in saline-treated mice (n=249 cells) and
BIBN-treated mice (n=42 cells). **(F)** Data (B, C, E) represents the proportion of cells. *p<0.05,
Fisher's exact test. Data (D) are median ± IQR.

# Discussion

Resident macrophages in the brain meninges are essential for maintaining brain

homeostasis, regulating central nervous system immune surveillance, and mediating

neuroimmune responses under pathological conditions [12, 14, 16, 57-59]. Macrophages

rely on $Ca^{2+}$ signaling to mediate many of their functions [17-23]. Here, using two-photon

microscopy in awake, behaving Pf4[Cre]:GCaMP6s reporter mice, we describe the

heterogeneity of meningeal macrophage $Ca^{2+}$ dynamics at steady state and in response

to CSD, an aberrant cortical hyperexcitability event associated with migraine, traumatic

brain injury, and stroke. Our data suggest the presence of two subsets of macrophages

in discrete meningeal niches, perivascular and non-perivascular, that exhibit several

distinct $Ca^{2+}$ signal properties at steady state. We further demonstrate $Ca^{2+}$ activity in

dural perivascular macrophages tuned to behaviorally-driven dural vasomotion. Finally,

we describe opposing $Ca^{2+}$ responses of meningeal macrophages following CSD, and

demonstrate the contribution of the CGRP/RAMP1 axis in mediating CSD-evoked

persistent $Ca^{2+}$ elevations.

The exact link between the distinct $Ca^{2+}$ signal properties of meningeal macrophage

subsets observed herein and their homeostatic function remains to be established. The

lower event magnitude and noisier signal observed in dural perivascular macrophages

may reflect functional interactions with the pulsation dynamics of dural vessels [46].

Indeed, by combining vascular and macrophage $Ca^{2+}$ imaging, we demonstrate a tight

temporal association between the diameter fluctuations of the dural vessels and the $Ca^{2+}$

signal of their associated macrophages in awake, locomoting mice. Direct vascular-

macrophage coupling may underlie this interaction, involving macrophages sensing

vascular-related mechanical changes via Piezo1 signaling [60]. Functional vascular-

macrophage interaction may also involve mural cells as inter-mediators [58, 61]. Studying

whether macrophage $Ca^{2+}$ signaling regulates dural vasomotion will require an

experimental approach that has yet to be developed, allowing for their selective

manipulation. The paucity of ongoing $Ca^{2+}$ activity in perivascular macrophages situated

in the leptomeninges we observed supports recent findings that subdural perivascular

macrophages indirectly affect pial and parenchymal vasomotion via extracellular matrix

remodeling [6].

Intracellular $Ca^{2+}$ signal propagation has been observed in various non-excitable cells,

such as astrocytes [62]. We show that the majority of meningeal macrophages, including

peri- and non-perivascular, exhibit intracellular $Ca^{2+}$ signals that propagate throughout the

entire cell, suggesting microdomain intracellular $Ca^{2+}$ signal elevation, following release

from internal stores. By characterizing the spatiotemporal relationships between $Ca^{2+}$

signals in distinct cells, we also demonstrate synchronous events that are independent of

spatial proximity, suggesting that synchronous $Ca^{2+}$ activity is not driven by intercellular

communication but rather by an external signal. Interestingly, our data indicate that

synchronized events involve both peri- and non-perivascular macrophages, despite

having distinct $Ca^{2+}$ elevation features, suggesting that these meningeal macrophage

subtypes similarly sense external signals.

Cortex-to-meninges signaling involves a relatively slow flow of soluble molecules in the

cerebrospinal fluid reaching the subarachnoid space and advancing via arachnoid cuff

exit points into the dura mater [11, 50]. Our finding of an acute $Ca^{2+}$ elevation in a subset

of extrasinusoidal perivascular dural macrophages coinciding with the CSD event

suggests a rapid transfer of soluble signaling factors from the cortex across all meningeal

layers [47]. Nevertheless, we cannot exclude a mechanically driven macrophage

response to the acute meningeal deformation produced by the neuronal and glial swelling

and shrinkage of the cortical extracellular space during CSD [53, 63, 64]. Our data also

indicate a delayed and prolonged increase in $Ca^{2+}$ signal in a relatively small subset of

macrophages post-CSD, which could underlie their proinflammatory-like morphological

change [49, 51]. Our findings also support the view that meningeal neuroimmune

CGRP/RAMP1 axis serves as a mechanism responsible for this macrophage $Ca^{2+}$

response, likely via the activation of their CGRP receptors [16]. Whether the relatively

small subset of meningeal macrophages featuring increased $Ca^{2+}$ signaling serves a

protective role [65, 66] or a proinflammatory, destructive function [67] remains to be

elucidated. Intriguingly, our data points to a persistent decrease in macrophage $Ca^{2+}$

activity post-CSD, not involving CGRP/RAMP1 signaling, as the most prevalent response,

raising the possibility that a hyperexcitable cortex dampens meningeal immunity.

# Conclusions

We provide a detailed characterization of macrophage $Ca^{2+}$ dynamics in homeostatic

meninges, thereby expanding our understanding of their biological diversity. The coupling

of dural perivascular macrophage $Ca^{2+}$ signals and dural vasomotion may represent a

unique homeostatic functional dural macrophage-vascular unit that controls dural

perfusion. The diversity of meningeal macrophage $Ca^{2+}$ responses to CSD further

highlights the complexity of brain-to-meninges neuroimmune signaling and meningeal

macrophage function in neurological disorders such as migraine, traumatic brain injury,

and stroke. Our study also provides essential genetic and data analysis tools to further

understand the molecular signaling underlying macrophage function at steady state and

neuropathological conditions.

# Methods

## Animals

All experimental procedures were approved by the Beth Israel Deaconess Medical

Center Institutional Animal Care and Use Committee. Experiments were conducted on

adult Pf4$^{Cre/wt}$:TIGRE2.0$^{GCaMP6s/wt}$ $Ca^{2+}$ reporter mice (8-17 weeks) (9 males, 5 females).

Mice were generated by crossing Pf4$^{Cre}$ (C57BL/6-Tg (Pf4-icre) Q3Rsko/J, Jackson

laboratory, Strain #008535) mice with GCaMP6s$^{fl/fl}$ (Ai162D; B6.Cg-Igs7$^{tm162.1(tetO-}$

GCaMP6s,CAG-tTA2)Hze/J, Jackson laboratory, Strain #031562) mice. Animals were

genotyped by Transnetyx Inc.

## Surgical procedures

Animals were anesthetized using isoflurane in 100% $O_2$ (induction: 3%; maintenance:

1.5-2%) and placed on a heating pad with a rectal probe attached to a stereotaxic frame

to monitor animal body temperature during surgery. Animals received dexamethasone (8

mg/kg, i.p.) and Meloxicam SR (4 mg/kg, s.c.) to reduce inflammation and improve

surgical outcomes. An eye ointment was used to prevent ocular drying. Mice were

implanted with a titanium headpost and a 3 mm glass cranial window (1.5 mm lateral and

2 mm posterior to Bregma) over an intact dura covering the left posterior neocortex [53].

Immediately after surgery, the mouse cage was placed on a water-circulating heating pad

for faster recovery. Animals were then single-housed with access to a running wheel and

a hut and allowed to recover for at least one week.

## Wheel running acclimation

436 After the cranial window surgery, mice were allowed to recover for at least a week. To

437 reduce stress associated with head-fixation during imaging and habituate to wheel

438 running, the mice received multiple training sessions (10 min to 1 h over 3–4 days). In

439 each session, the mouse was placed on a 3D printed running wheel, with its headpost

440 attached to two clamps, and allowed to locomote freely.

## Two-photon $Ca^{2+}$ imaging

Awake-behaving mice were head-fixed to the running wheel by its headpost (**Figure 1**).

We used a two-photon microscope (Neurolabware) with a Nikon 16X, 0.8 NA objective to

acquire images at 15.5 Hz with digital zoom set at 4X (312 x 212 $\mu m^2$ FOV). A MaiTai

laser set to 920 nm with 25–40 mW power was used to excite fluorescence. The Scanbox

package for MATLAB (Neurolabware) was used to control the microscope and acquire

images and wheel running data. To image the meningeal vasculature, mice were

administered 70kDa TRITC-Dextran tracer (50 mg/kg, i.v.; Sigma-Aldrich).

**Behavioral tracking during two-photon imaging**

We recorded running speed in MATLAB using a custom-made encoder (Arduino) coupled

to the 3D-printed running wheel. See below for details of analyses of behavioral variables.

**Induction of CSD and pharmacological treatment**

For CSD induction, a 1 mm burr hole was drilled at the frontal bone (1.5 mm anterior to

the cranial window) to allow access to the brain cortical surface. A small amount of

silicone elastomer (Kwik-Cast, WPI) was placed to cover the burr hole opening, and the

animal was left to recover for at least one week. CSD was induced using a brief 2-second

cortical pinprick [53]. CSD induction was confirmed by the identification of a short-lasting

meningeal deformation and/or transient pial constriction [53]. On the experimental day,

30 minutes before baseline recording, mice were pretreated with the CGRP/RAMP1

antagonist BIBN4096 (0.3 ml, 1 mg/kg, i.p., Tocris) [54] or 0.3 ml of saline (Vehicle

control).

## Quantification and statistical analysis

**Two-photon imaging movie processing**

We used a discrete Fourier transform to perform rigid registration to correct translation changes due to brain motion during locomotion. Movies were then downsampled to 1.03 Hz. Locomotion signals were detected as described [41]. All image processing and locomotion signal extraction were performed in MATLAB 2021b (Mathworks).

**$Ca^{2+}$ signal detection pipeline**

For detecting macrophage $Ca^{2+}$ signals, we used the Activity Quantification and Analysis (AQuA2) platform that implements an event-based approach with advanced machine learning techniques for temporal and spatial segmentation of $Ca^{2+}$ fluorescence events [38]. Importantly, this computational platform captures event dynamics beyond traditional ROI-based approaches. The following user-defined parameters were input: 0.49 µm/pixel spatial resolution, 1.03 Hz temporal resolution, 1-second minimal event duration detection; window of event size between 25% (125 µm$^2$/ 523 pixels) and 150% (750 µm$^2$/ 3140 pixels). Every event and cell identified was followed by a manual visual check. Subsequently, each cell was labelled as perivascular or non-perivascular according to its location relative to vessels. Morphological features (area, perimeter, circularity) and spatiotemporal aspects of the $Ca^{2+}$ signals (i.e., $\Delta F/F_0$ dynamics, frequency, amplitude, duration) were used to analyze cell-specific characteristics and $Ca^{2+}$ activity profiles. We employed the AQuA2 automatically-generated function [42] to characterize intracellular $Ca^{2+}$ propagation. Intercellular $Ca^{2+}$ activity was evaluated by analyzing temporally co-

occurring events (synchronized event pairs), as well as their corresponding spatial

localization in the FOV. All post-processing of AQuA2-generated data was performed

using MATLAB 2021b (Mathworks).

**Clustering of Ca$^{2+}$ dynamics**

We used a Savitzky-Golay filter to detrend and smooth Ca$^{2+}$ activity traces. Polynomial

order and frame size were optimized for each cell by selecting the combination of

parameters that yielded a better signal-to-noise ratio. From the filtered signals, we

extracted the dominant frequency using Fast Fourier Transform, and peak counts using

minimum peak prominence (threshold of 10% signal amplitude range). The optimal

number of clusters was determined using the Elbow method, after which k-means

clustering was applied to group cells based on their signal characteristics.

**Analysis of CSD-related changes in macrophage Ca$^{2+}$ dynamics**

For analyzing the effects of CSD on macrophages' Ca$^{2+}$ dynamics, we divided the activity

of each cell into three phases: 'PreCSD' (minutes 0 – 30), 'DuringCSD' (minutes 30 – 31),

and 'PostCSD' (minutes 31 – 61). Ca$^{2+}$ event rate during CSD and PostCSD was

compared against that of the PreCSD baseline for evaluation of acute and persistent Ca$^{2+}$

responses, respectively. Ca$^{2+}$ responses were categorized as increased (event rate > 2x

PreCSD), decreased (event rate < 0.5x PreCSD), or unchanged.

**Analysis of locomotion**

We extracted running speed (cm/sec) from the wheel encoder. To infer the locomotion state, we first concatenated all velocity signals obtained from a given mouse across all experiments and trained a two-state Hidden Markov Model using the MATLAB function 'hmmtrain'. Then, the locomotion state was inferred for each individual imaging run by applying the MATLAB function 'hmmviterbi' with the model trained on the concatenated data. Locomotion bouts were defined as periods when the locomotion state was sustained for at least two seconds.

**Vascular signals**

We calculated changes in vascular diameter by first generating a maximum intensity projection of the red channel and drawing polygons (ROIs) around each vessel. Subsequently, for each frame, pixels inside each ROI were extracted, and a Radon transform was applied to get a 1D vessel profile. Radon transform of the first frame was used as a reference to normalize the results of all subsequent frames, resulting in a time series of normalized diameter traces.

**General linear models (GLM)**

We investigate the functional interaction between dural perivascular macrophage $Ca^{2+}$ activity and dural vasomotion by fitting a Gaussian GLMs using the GLMnet package (MATLAB 2021b), with elastic net regularization ($\alpha=0.01$) and ten-fold cross-validation. We employed a two-step modeling. The first model was used to identify dural vessels by evaluating the correlation of vessel diameter changes to locomotion state, classifying dural or pial vessels according to their vasoconstriction and vasodilation

dynamics, respectively [46]. The second modeling step evaluated the relative contribution

of perivascular dural $Ca^{2+}$ activity signals to dural vessel diameter changes. All signals

were downsampled to 1.03 Hz to match the frame rate used when extracting fluorescence

$Ca^{2+}$ signals in AQuA2. To allow potential anticipatory or delayed responses of diameter-

to-locomotion or $Ca^{2+}$ fluorescence-to-diameter, the time interval used to analyze the

response to the predictor was set from -60 s to +60 s. The GLM was trained on 75% of

the data, and all predictions and model performances reported are from the remaining

25% testing set. A threshold of 0.1 goodness-of-fit deviance explained was set. For each

predictor temporal shift, a response coefficient was generated. Centroid delay between

predictor and response was calculated, weighted by the absolute value of the coefficient

(i.e., center-of-mass, COM).

**Statistical analysis**

All statistical analysis was performed using GraphPad Prism 10.4 and MATLAB 2021b.

Data were analyzed using a Wilcoxon matched-pairs signed rank sum test or a Mann-

Whitney U-test. Distribution of categorical data was analyzed using the Chi-square or

Fisher's exact test. P-values are indicated as follows: $*p<0.05$, $**p<0.01$, $***p<0.001$,

$****p<0.0001$.

**Data availability.**

All data needed to evaluate the conclusions in the paper are present in the paper.

The code used for analyzing the data in this study was deposited in the Levy Lab

GitHub account.

Code for movie processing is available at:

https://github.com/levylabheadache/MovieProcessing/tree/SCN;

Code for locomotion analysis is available at:

https://github.com/levylabheadache/Locomotion/tree/SCN;

Code for post Aqua2-processing is available at:

https://github.com/levylabheadache/Aqua2Processing;

Code for vascular segmentation is available at:

https://github.com/levylabheadache/Vasculature/tree/SCN;

Code for GLM is available at:

https://github.com/levylabheadache/GeneralLinearModel_Macrophages

Any additional information is available from the corresponding author upon request.

**Abbreviations**

AQuA2 – Activity Quantification and Analysis

$Ca^{2+}$ – Cytoplasmic calcium

CGRP – Calcitonin gene-related peptide

CSD – Cortical spreading depolarization (depression)

CX3CR1 – CX3C Motif chemokine receptor 1

GLM – General Linear Model

NP – Non-perivascular

P – Perivascular

Pf4 – Platelet factor 4

RAMP1 – Receptor activity modifying protein 1

TRITC – Tetramethylrhodamine isothiocyanate

# References

584    1.    Okabe Y, Medzhitov R. Tissue biology perspective on macrophages. Nat Immunol.
585    2016;17(1):9-17. doi: 10.1038/ni.3320. PubMed PMID: 26681457.
586
587    2.    Guilliams M, Thierry GR, Bonnardel J, Bajenoff M. Establishment and Maintenance
588    of   the   Macrophage   Niche.   Immunity.   2020;52(3):434-51.   doi:
589    10.1016/j.immuni.2020.02.015. PubMed PMID: 32187515.
590
591    3.    Mass E, Nimmerjahn F, Kierdorf K, Schlitzer A. Tissue-specific macrophages: how
592    they develop and choreograph tissue biology. Nature reviews. 2023;23(9):563-79. Epub
593    20230315.   doi:   10.1038/s41577-023-00848-y.   PubMed   PMID:   36922638;   PubMed
594    Central PMCID: PMCPMC10017071.
595
596    4.    Kierdorf K, Masuda T, Jordao MJC, Prinz M. Macrophages at CNS interfaces:
597    ontogeny and function in health and disease. Nat Rev Neurosci. 2019;20(9):547-62. Epub
598    20190729. doi: 10.1038/s41583-019-0201-x. PubMed PMID: 31358892.
599
600    5.    Rustenhoven J, Drieu A, Mamuladze T, de Lima KA, Dykstra T, Wall M, et al.
601    Functional characterization of the dural sinuses as a neuroimmune interface. Cell.
602    2021;184(4):1000-16 e27. Epub 2021/01/29. doi: 10.1016/j.cell.2020.12.040. PubMed
603    PMID: 33508229; PubMed Central PMCID: PMCPMC8487654.
604
605    6.    Drieu A, Du S, Storck SE, Rustenhoven J, Papadopoulos Z, Dykstra T, et al.
606    Parenchymal border macrophages regulate the flow dynamics of the cerebrospinal fluid.
607    Nature.   2022;611(7936):585-93.   Epub   20221109.   doi:   10.1038/s41586-022-05397-3.
608    PubMed PMID: 36352225; PubMed Central PMCID: PMCPMC9899827.
609
610    7.    Masuda T, Amann L, Monaco G, Sankowski R, Staszewski O, Krueger M, et al.
611    Specification of CNS macrophage subsets occurs postnatally in defined niches. Nature.
612    2022;604(7907):740-8.   Epub   20220420.   doi:   10.1038/s41586-022-04596-2.   PubMed
613    PMID: 35444273.
614
615    8.    Amann L, Fell A, Monaco G, Sankowski R, Wu HZQ, Jordao MJC, et al.
616    Extrasinusoidal macrophages are a distinct subset of immunologically active dural
617    macrophages.   Sci   Immunol.   2024;9(102):eadh1129.   Epub   20241220.   doi:
618    10.1126/sciimmunol.adh1129. PubMed PMID: 39705337.
619
620    9.    Mrdjen D, Pavlovic A, Hartmann FJ, Schreiner B, Utz SG, Leung BP, et al. High-
621    Dimensional Single-Cell Mapping of Central Nervous System Immune Cells Reveals

Distinct Myeloid Subsets in Health, Aging, and Disease. Immunity. 2018;48(2):380-95 e6. Epub 20180206. doi: 10.1016/j.immuni.2018.01.011. PubMed PMID: 29426702.

10.    Van Hove H, Martens L, Scheyltjens I, De Vlaminck K, Pombo Antunes AR, De Prijck S, et al. A single-cell atlas of mouse brain macrophages reveals unique transcriptional identities shaped by ontogeny and tissue environment. Nat Neurosci. 2019;22(6):1021-35. Epub 2019/05/08. doi: 10.1038/s41593-019-0393-4. PubMed PMID: 31061494.

11.    Smyth LCD, Xu D, Okar SV, Dykstra T, Rustenhoven J, Papadopoulos Z, et al. Identification of direct connections between the dura and the brain. Nature. 2024;627(8002):165-73. Epub 20240207. doi: 10.1038/s41586-023-06993-7. PubMed PMID: 38326613.

12.    Vara-Perez M, Movahedi K. Border-associated macrophages as gatekeepers of brain homeostasis and immunity. Immunity. 2025;58(5):1085-100. Epub 20250504. doi: 10.1016/j.immuni.2025.04.005. PubMed PMID: 40324381; PubMed Central PMCID: PMCPMC12094687.

13.    Rua R, Lee JY, Silva AB, Swafford IS, Maric D, Johnson KR, et al. Infection drives meningeal engraftment by inflammatory monocytes that impairs CNS immunity. Nat Immunol. 2019;20(4):407-19. Epub 20190318. doi: 10.1038/s41590-019-0344-y. PubMed PMID: 30886419; PubMed Central PMCID: PMCPMC6481670.

14.    Rebejac J, Eme-Scolan E, Arnaud Paroutaud L, Kharbouche S, Teleman M, Spinelli L, et al. Meningeal macrophages protect against viral neuroinfection. Immunity. 2022;55(11):2103-17 e10. Epub 20221101. doi: 10.1016/j.immuni.2022.10.005. PubMed PMID: 36323311.

15.    De Vlaminck K, Van Hove H, Kancheva D, Scheyltjens I, Pombo Antunes AR, Bastos J, et al. Differential plasticity and fate of brain-resident and recruited macrophages during the onset and resolution of neuroinflammation. Immunity. 2022;55(11):2085-102 e9. Epub 20221012. doi: 10.1016/j.immuni.2022.09.005. PubMed PMID: 36228615.

16.    Pinho-Ribeiro FA, Deng L, Neel DV, Erdogan O, Basu H, Yang D, et al. Bacteria hijack a meningeal neuroimmune axis to facilitate brain invasion. Nature. 2023;615(7952):472-81. Epub 20230301. doi: 10.1038/s41586-023-05753-x. PubMed PMID: 36859544; PubMed Central PMCID: PMCPMC10593113.

17.    Desai BN, Leitinger N. Purinergic and calcium signaling in macrophage function and plasticity. Front Immunol. 2014;5:580. Epub 20141127. doi: 10.3389/fimmu.2014.00580. PubMed PMID: 25505897; PubMed Central PMCID: PMCPMC4245916.

18.    Zumerle S, Cali B, Munari F, Angioni R, Di Virgilio F, Molon B, et al. Intercellular Calcium Signaling Induced by ATP Potentiates Macrophage Phagocytosis. Cell Rep.

668  2019;27(1):1-10 e4. doi: 10.1016/j.celrep.2019.03.011. PubMed PMID: 30943393;
669  PubMed Central PMCID: PMCPMC6449513.
670
671  19.    Seegren PV, Downs TK, Stremska ME, Harper LR, Cao R, Olson RJ, et al.
672  Mitochondrial Ca(2+) Signaling Is an Electrometabolic Switch to Fuel Phagosome Killing.
673  Cell Rep. 2020;33(8):108411. doi: 10.1016/j.celrep.2020.108411. PubMed PMID:
674  33238121; PubMed Central PMCID: PMCPMC7793167.
675
676  20.    Nascimento Da Conceicao V, Sun Y, Ramachandran K, Chauhan A, Raveendran
677  A, Venkatesan M, et al. Resolving macrophage polarization through distinct Ca(2+) entry
678  channel that maintains intracellular signaling and mitochondrial bioenergetics. iScience.
679  2021;24(11):103339. Epub 20211023. doi: 10.1016/j.isci.2021.103339. PubMed PMID:
680  34816101; PubMed Central PMCID: PMCPMC8591423.
681
682  21.    Taghdiri N, Calcagno DM, Fu Z, Huang K, Kohler RH, Weissleder R, et al.
683  Macrophage calcium reporter mice reveal immune cell communication in vitro and in vivo.
684  Cell Rep Methods. 2021;1(8). Epub 20211213. doi: 10.1016/j.crmeth.2021.100132.
685  PubMed PMID: 35079727; PubMed Central PMCID: PMCPMC8786215.
686
687  22.    Mehari FT, Miller M, Pick R, Bader A, Pekayvaz K, Napoli M, et al. Intravital calcium
688  imaging in myeloid leukocytes identifies calcium frequency spectra as indicators of
689  functional states. Sci Signal. 2022;15(744):eabe6909. Epub 20220726. doi:
690  10.1126/scisignal.abe6909. PubMed PMID: 35881691.
691
692  23.    Seegren PV, Harper LR, Downs TK, Zhao XY, Viswanathan SB, Stremska ME, et
693  al. Reduced mitochondrial calcium uptake in macrophages is a major driver of
694  inflammaging. Nat Aging. 2023;3(7):796-812. Epub 20230605. doi: 10.1038/s43587-023-
695  00436-8. PubMed PMID: 37277641; PubMed Central PMCID: PMCPMC10353943.
696
697  24.    Vaeth M, Zee I, Concepcion AR, Maus M, Shaw P, Portal-Celhay C, et al. Ca2+
698  Signaling but Not Store-Operated Ca2+ Entry Is Required for the Function of
699  Macrophages and Dendritic Cells. J Immunol. 2015;195(3):1202-17. Epub 20150624. doi:
700  10.4049/jimmunol.1403013. PubMed PMID: 26109647; PubMed Central PMCID:
701  PMCPMC4506881.
702
703  25.    Schappe MS, Stremska ME, Busey GW, Downs TK, Seegren PV, Mendu SK, et
704  al. Efferocytosis requires periphagosomal Ca(2+)-signaling and TRPM7-mediated
705  electrical activity. Nat Commun. 2022;13(1):3230. Epub 20220609. doi: 10.1038/s41467-
706  022-30959-4. PubMed PMID: 35680919; PubMed Central PMCID: PMCPMC9184625.
707
26.    Kaya Z, Belder N, Sever-Bahcekapili M, Erdener SE, Donmez-Demir B, Bagci C,
et al. Spreading depolarization triggers pro- and anti-inflammatory signalling: a potential
link to headache. Brain. 2025;148(7):2522-36. doi: 10.1093/brain/awaf015. PubMed
PMID: 39823578; PubMed Central PMCID: PMCPMC12378585.

27.    Harriott AM, Ayata C. Spreading depolarization as a therapeutic target in migraine.
Nat Rev Neurol. 2025;21(10):529-43. Epub 20250814. doi: 10.1038/s41582-025-01128-
0. PubMed PMID: 40813918.

28.    Ren X, Manzanares LD, Piccolo EB, Urbanczyk JM, Sullivan DP, Yalom LK, et al.
Macrophage-endothelial cell crosstalk orchestrates neutrophil recruitment in inflamed
mucosa. J Clin Invest. 2023;133(15). Epub 20230801. doi: 10.1172/JCI170733. PubMed
PMID: 37261911; PubMed Central PMCID: PMCPMC10378177.

29.    Ishikawa-Ankerhold H, Busch B, Bader A, Maier-Begandt D, Dionisio F, Namineni
S, et al. Novel multiphoton intravital imaging enables real-time study of Helicobacter pylori
interaction with neutrophils and macrophages in the mouse stomach. PLoS Pathog.
2024;20(9):e1012580. Epub 20240930. doi: 10.1371/journal.ppat.1012580. PubMed
PMID: 39348445; PubMed Central PMCID: PMCPMC11478878.

30.    Chinnery HR, Ruitenberg MJ, McMenamin PG. Novel characterization of
monocyte-derived cell populations in the meninges and choroid plexus and their rates of
replenishment in bone marrow chimeric mice. J Neuropathol Exp Neurol. 2010;69(9):896-
909. doi: 10.1097/NEN.0b013e3181edbc1a. PubMed PMID: 20720507.

31.    Jung S, Aliberti J, Graemmel P, Sunshine MJ, Kreutzberg GW, Sher A, et al.
Analysis of fractalkine receptor CX(3)CR1 function by targeted deletion and green
fluorescent protein reporter gene insertion. Mol Cell Biol. 2000;20(11):4106-14. doi:
10.1128/MCB.20.11.4106-4114.2000. PubMed PMID: 10805752; PubMed Central
PMCID: PMCPMC85780.

32.    Roth TL, Nayak D, Atanasijevic T, Koretsky AP, Latour LL, McGavern DB.
Transcranial amelioration of inflammation and cell death after brain injury. Nature.
2014;505(7482):223-8. Epub 2013/12/10. doi: 10.1038/nature12808. PubMed PMID:
24317693; PubMed Central PMCID: PMCPMC3930079.

33.    Liu YU, Ying Y, Li Y, Eyo UB, Chen T, Zheng J, et al. Neuronal network activity
controls microglial process surveillance in awake mice via norepinephrine signaling. Nat
Neurosci. 2019;22(11):1771-81. Epub 20191021. doi: 10.1038/s41593-019-0511-3.
PubMed PMID: 31636449; PubMed Central PMCID: PMCPMC6858573.

34.    McKinsey GL, Lizama CO, Keown-Lang AE, Niu A, Santander N, Larpthaveesarp
A, et al. A new genetic strategy for targeting microglia in development and disease. Elife.
2020;9. Epub 20200623. doi: 10.7554/eLife.54590. PubMed PMID: 32573436; PubMed
Central PMCID: PMCPMC7375817.

35.    Daigle TL, Madisen L, Hage TA, Valley MT, Knoblich U, Larsen RS, et al. A Suite
of Transgenic Driver and Reporter Mouse Lines with Enhanced Brain-Cell-Type Targeting
and Functionality. Cell. 2018;174(2):465-80 e22. doi: 10.1016/j.cell.2018.06.035.
PubMed PMID: 30007418; PubMed Central PMCID: PMCPMC6086366.

36.     Tiedt R, Schomber T, Hao-Shen H, Skoda RC. Pf4-Cre transgenic mice allow the generation of lineage-restricted gene knockouts for studying megakaryocyte and platelet function in vivo. Blood. 2007;109(4):1503-6. Epub 20061010. doi: 10.1182/blood-2006-04-020362. PubMed PMID: 17032923.

37.     Umpierre AD, Bystrom LL, Ying Y, Liu YU, Worrell G, Wu LJ. Microglial calcium signaling is attuned to neuronal activity in awake mice. Elife. 2020;9. Epub 20200727. doi: 10.7554/eLife.56502. PubMed PMID: 32716294; PubMed Central PMCID: PMCPMC7402678.

38.     Thrane AS, Rangroo Thrane V, Zeppenfeld D, Lou N, Xu Q, Nagelhus EA, et al. General anesthesia selectively disrupts astrocyte calcium signaling in the awake mouse cortex. Proc Natl Acad Sci U S A. 2012;109(46):18974-9. Epub 2012/11/01. doi: 10.1073/pnas.1209448109. PubMed PMID: 23112168; PubMed Central PMCID: PMCPMC3503159.

39.     Zhou H, Neudecker V, Perez-Zoghbi JF, Brambrink AM, Yang G. Age-dependent cerebral vasodilation induced by volatile anesthetics is mediated by NG2(+) vascular mural cells. Commun Biol. 2024;7(1):1519. Epub 20241115. doi: 10.1038/s42003-024-07200-7. PubMed PMID: 39548262; PubMed Central PMCID: PMCPMC11568297.

40.     Goldey GJ, Roumis DK, Glickfeld LL, Kerlin AM, Reid RC, Bonin V, et al. Removable cranial windows for long-term imaging in awake mice. Nature protocols. 2014;9(11):2515-38. Epub 2014/10/03. doi: 10.1038/nprot.2014.165. PubMed PMID: 25275789; PubMed Central PMCID: PMCPMC4442707.

41.     Blaeser AS, Sugden AU, Zhao J, Carneiro-Nascimento S, Shipley FB, Carrie H, et al. Trigeminal afferents sense locomotion-related meningeal deformations. Cell Rep. 2022;41(7):111648. doi: 10.1016/j.celrep.2022.111648. PubMed PMID: 36384109.

42.     Mi X, Chen AB, Duarte D, Carey E, Taylor CR, Braaker PN, et al. Fast, accurate, and versatile data analysis platform for the quantification of molecular spatiotemporal signals. Cell. 2025;188(10):2794-809 e21. Epub 20250408. doi: 10.1016/j.cell.2025.03.012. PubMed PMID: 40203826.

43.     Leon Guerrero PA, Rasmussen JP, Peterman E. Calcium dynamics of skin-resident macrophages during homeostasis and tissue injury. Mol Biol Cell. 2024;35(12):br26. Epub 20241113. doi: 10.1091/mbc.E24-09-0420. PubMed PMID: 39535893; PubMed Central PMCID: PMCPMC11656469.

44.     Paterson N, Lammermann T. Macrophage network dynamics depend on haptokinesis for optimal local surveillance. Elife. 2022;11. Epub 20220328. doi: 10.7554/eLife.75354. PubMed PMID: 35343899; PubMed Central PMCID: PMCPMC8963880.

45.     Behmoaras J, Mulder K, Ginhoux F, Petretto E. The spatial and temporal activation of macrophages during fibrosis. Nature reviews. 2025. Epub 20250604. doi: 10.1038/s41577-025-01186-x. PubMed PMID: 40467841.

46.     Gao YR, Drew PJ. Effects of Voluntary Locomotion and Calcitonin Gene-Related Peptide on the Dynamics of Single Dural Vessels in Awake Mice. J Neurosci. 2016;36(8):2503-16. Epub 2016/02/26. doi: 10.1523/JNEUROSCI.3665-15.2016. PubMed PMID: 26911696; PubMed Central PMCID: PMCPMC4764666.

47.     Levy D, Moskowitz MA. Meningeal Mechanisms and the Migraine Connection. Annual review of neuroscience. 2023. Epub 20230313. doi: 10.1146/annurev-neuro-080422-105509. PubMed PMID: 36913712.

48.     Kursun O, Yemisci M, van den Maagdenberg A, Karatas H. Migraine and neuroinflammation: the inflammasome perspective. J Headache Pain. 2021;22(1):55. Epub 20210610. doi: 10.1186/s10194-021-01271-1. PubMed PMID: 34112082; PubMed Central PMCID: PMCPMC8192049.

49.     Schain AJ, Melo-Carrillo A, Borsook D, Grutzendler J, Strassman AM, Burstein R. Activation of pial and dural macrophages and dendritic cells by cortical spreading depression. Ann Neurol. 2018;83(3):508-21. Epub 2018/02/03. doi: 10.1002/ana.25169. PubMed PMID: 29394508; PubMed Central PMCID: PMCPMC5965700.

50.     Kipnis J. The anatomy of brainwashing. Science. 2024;385(6707):368-70. Epub 20240725. doi: 10.1126/science.adp1705. PubMed PMID: 39052816.

51.     Chauhan A, Sun Y, Sukumaran P, Quenum Zangbede FO, Jondle CN, Sharma A, et al. M1 Macrophage Polarization Is Dependent on TRPC1-Mediated Calcium Entry. iScience. 2018;8:85-102. Epub 20180920. doi: 10.1016/j.isci.2018.09.014. PubMed PMID: 30293012; PubMed Central PMCID: PMCPMC6174824.

52.     Zhao J, Levy D. Cortical Spreading Depression Promotes Persistent Mechanical Sensitization of Intracranial Meningeal Afferents: Implications for the Intracranial Mechanosensitivity of Migraine. eNeuro. 2016;3(6):ENEURO.0287-16.2016. Epub 2017/01/28. doi: 10.1523/ENEURO.0287-16.2016. PubMed PMID: 28127585; PubMed Central PMCID: PMCPMC5242377.

53.     Blaeser AS, Zhao J, Sugden AU, Carneiro-Nascimento S, Andermann ML, Levy D. Sensitization of meningeal afferents to locomotion-related meningeal deformations in a migraine model. Elife. 2024;12. Epub 20240208. doi: 10.7554/eLife.91871. PubMed PMID: 38329894; PubMed Central PMCID: PMCPMC10942541.

54.     Zhao J, Levy D. The CGRP receptor antagonist BIBN4096 inhibits prolonged meningeal afferent activation evoked by brief local K(+) stimulation but not cortical spreading depression-induced afferent sensitization. Pain Rep. 2018;3(1):e632. Epub

2018/02/13. doi: 10.1097/PR9.0000000000000632. PubMed PMID: 29430561; PubMed
Central PMCID: PMCPMC5802320.

55.    Deng L, Gillis JE, Chiu IM, Kaplan DH. Sensory neurons: An integrated component
of innate immunity. Immunity. 2024;57(4):815-31. doi: 10.1016/j.immuni.2024.03.008.
PubMed PMID: 38599172.

56.    Jin X, Morais A, Sasaki Y, Zhai Q, Banerjee P, Harriott A, et al. Small-molecule
CGRP antagonist atogepant does not affect cortical spreading depression susceptibility
in rats. J Headache Pain. 2025;26(1):177. Epub 20250805. doi: 10.1186/s10194-025-
02127-8. PubMed PMID: 40764901; PubMed Central PMCID: PMCPMC12326636.

57.    Rustenhoven J, Kipnis J. Brain borders at the central stage of neuroimmunology.
Nature. 2022;612(7940):417-29. Epub 20221214. doi: 10.1038/s41586-022-05474-7.
PubMed PMID: 36517712; PubMed Central PMCID: PMCPMC10205171.

58.    Min H, O'Neil SM, Xu L, Moseman EA, Kurtzberg J, Filiano AJ. Mural cells interact
with macrophages in the dura mater to regulate CNS immune surveillance. J Exp Med.
2024;221(2). Epub 20240109. doi: 10.1084/jem.20230326. PubMed PMID: 38193859;
PubMed Central PMCID: PMCPMC10783178.

59.    Dalmau Gasull A, Glavan M, Samawar SKR, Kapupara K, Kelk J, Rubio M, et al.
The niche matters: origin, function and fate of CNS-associated macrophages during
health and disease. Acta Neuropathol. 2024;147(1):37. Epub 20240212. doi:
10.1007/s00401-023-02676-9. PubMed PMID: 38347231; PubMed Central PMCID:
PMCPMC10861620.

60.    Atcha H, Jairaman A, Holt JR, Meli VS, Nagalla RR, Veerasubramanian PK, et al.
Mechanically activated ion channel Piezo1 modulates macrophage polarization and
stiffness sensing. Nat Commun. 2021;12(1):3256. Epub 20210531. doi: 10.1038/s41467-
021-23482-5. PubMed PMID: 34059671; PubMed Central PMCID: PMCPMC8167181.

61.    Tong L, Hill RA, Damisah EC, Murray KN, Yuan P, Bordey A, et al. Imaging and
optogenetic modulation of vascular mural cells in the live brain. Nature protocols.
2021;16(1):472-96. Epub 20201209. doi: 10.1038/s41596-020-00425-w. PubMed PMID:
33299155.

62.    Semyanov A, Verkhratsky A. Astrocytic processes: from tripartite synapses to the
active milieu. Trends Neurosci. 2021;44(10):781-92. Epub 20210831. doi:
10.1016/j.tins.2021.07.006. PubMed PMID: 34479758.

63.    Takano T, Tian GF, Peng W, Lou N, Lovatt D, Hansen AJ, et al. Cortical spreading
depression causes and coincides with tissue hypoxia. Nat Neurosci. 2007;10(6):754-62.
Epub 2007/05/01. doi: 10.1038/nn1902. PubMed PMID: 17468748.

64. Rosic B, Dukefoss DB, Abjorsbraten KS, Tang W, Jensen V, Ottersen OP, et al. Aquaporin-4-independent volume dynamics of astroglial endfeet during cortical spreading depression. Glia. 2019;67(6):1113-21. Epub 2019/02/23. doi: 10.1002/glia.23604. PubMed PMID: 30791140; PubMed Central PMCID: PMCPMC6594042.

65. Lu YZ, Nayer B, Singh SK, Alshoubaki YK, Yuan E, Park AJ, et al. CGRP sensory neurons promote tissue healing via neutrophils and macrophages. Nature. 2024;628(8008):604-11. Epub 20240327. doi: 10.1038/s41586-024-07237-y. PubMed PMID: 38538784; PubMed Central PMCID: PMCPMC11023938.

66. Hasegawa T, Lee CYC, Hotchen AJ, Fleming A, Singh R, Suzuki K, et al. Macrophages and nociceptor neurons form a sentinel unit around fenestrated capillaries to defend the synovium from circulating immune challenge. Nat Immunol. 2024;25(12):2270-83. Epub 20241125. doi: 10.1038/s41590-024-02011-8. PubMed PMID: 39587345; PubMed Central PMCID: PMCPMC11588661.

67. Fattori V, Zaninelli TH, Rasquel-Oliveira FS, Heintz OK, Jain A, Sun L, et al. Nociceptor-to-macrophage communication through CGRP/RAMP1 signaling drives endometriosis-associated pain and lesion growth in mice. Sci Transl Med. 2024;16(772):eadk8230. Epub 20241106. doi: 10.1126/scitranslmed.adk8230. PubMed PMID: 39504351.

## Funding

The study was supported by NIH grants: R21NS130561; R01NS115972 and R01NS133625 to D.L.

## Authors information

D.L. conceived the project. S.C-N and D.L. wrote the manuscript. S.C-N performed two-photon imaging and data analysis with help from D.L. S.C-N, C.W., and A.G. performed surgeries.

**Authors and affiliations**

[1]Department of Anesthesia, Critical Care and Pain Medicine, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, United States

## Corresponding author

927     Correspondence to: dlevy1@bidmc.harvard.edu (Dan Levy)

## Competing interests

931     The authors declare that they have no competing interests.

**Supplementary video legends:**

**Video S1. Macrophage Ca$^{2+}$ dynamics in homeostatic meninges of awake mice**

Example two-photon imaging showing Ca$^{2+}$ activity in perivascular and non-perivascular meningeal macrophages.

**Video S2. Acute and persistent changes in meningeal macrophage Ca$^{2+}$ activity in response to CSD.**

Example two-photon imaging of meningeal macrophage Ca$^{2+}$ activity at baseline, during, and following CSD. The arrow indicates a macrophage showing an acute Ca$^{2+}$ elevation, the arrowhead depicts a delayed and persistent Ca$^{2+}$ elevation, and the asterisk, a macrophage showing a persistent decreased Ca$^{2+}$ activity.